1 **Laminar endocytic machinery revealed by quantitative multiplexed super-**
2 **resolution imaging**
3

4 **Tai Kiuchi[1]*, Ryouhei Kobayashi[2], Shuichiro Ogawa[1]†, Louis L. H. Elverston[2] and**
5 **Naoki Watanabe[1, 2]**
6
7
8 [1] Department of Pharmacology, Kyoto University Faculty of Medicine, Sakyo-ku, Kyoto
9 606-8501, Japan
10
11 [2] Laboratory of Single-Molecule Cell Biology, Kyoto University Graduate School of
12 Biostudies, Sakyo-ku, Kyoto 606-8501, Japan
13
14 * Correspondence should be addressed to T.K. (kiuchi.tai.6z@kyoto-u.ac.jp)
15
16 Tel: +81-75-753-4397, Fax: +81-75-753-4394
17
18 † Present address: Nagahama Institute of Bio-Science and Technology, 1266 Tamura-cho,
19 Nagahama, Shiga 526-0829, Japan
20

Abstract

Few algorithms exist to elucidate colocalization between multiple molecules. Here we present a method to map molecular complexes using multiplexed super-resolution microscopy IRIS. We developed antiserum-derived Fab IRIS probes that enabled nanoscale quantification of endocytic proteins in a clathrin-coated structure (CCS). Our Protein Cluster coloring (PC-coloring), which employs pixel-based principal component analysis and clustering, maps protein clusters of distinct ratios. Subsequent correlation analysis between two proteins detects their complex formation in a specific PC-colored region. Upon EGF stimulation, EGFR-, EGFR-Grb2- and Grb2-dominant clusters lined up from the CCS rim. Along the interior of Grb2-clusters, Eps15, FCHo1/2 and intersectin-1 formed a complex with Grb2. The interior of Eps15-FCHo1/2-clusters was lined with Eps15-intersectin-1-clusters. The results reveal laminar organization of complexes among EGFR, Grb2 and CCS components.

In clathrin-coated structures (CCSs), a variety of proteins colocalize to from molecular complexes for receptor endocytosis (*1*). These localizations have been elucidated by immunoelectron microscopy and super-resolution microscopy (*2–7*). Super-resolution microscopy with a resolution of a few nanometers has been reported (*8*, *9*), allowing visualization of molecular localization within molecular complexes. However, multiple labeling of endogenous proteins with antibodies is impossible in a region smaller than the antibody size (> 10 nm) (*10*, *11*). Especially in multiplexed super-resolution microscopy, physical interference between antibodies may reduce the accessibility to the antigens as previously reported (*12*). Due to the incomplete labeling, accuracy in colocalization analysis cannot be improved solely by increasing the resolution. Although an interaction network of proteins in CCSs has been proposed based on *in vitro* protein-protein binding (*13*, *14*), the spatial relationship between their colocalization sites remains to be determined.

A multiplexed super-resolution microscopy, named image reconstruction by integrating exchangeable single-molecule localization (IRIS) was developed to overcome the labeling problem (*15*). IRIS employs fluorescent probes (IRIS probes) that rapidly associate with and dissociate from the endogenous target (*15*). The number of labels can surpass that of target molecules without causing physical interference between the probes (*12*). By sequentially incubating the samples with multiple probes, a large number of targets can be stained in a single specimen. The IRIS probes have ever been produced

from protein fragments, Fab fragments of mouse monoclonal antibodies, and engineered

antibody and nanobody fragments (*12*, *15*, *16*). Here we introduce a new IRIS probe from

antiserum. The antiserum contains polyclonal antibodies against multiple epitopes in the

target protein. Therefore, antiserum-derived Fab probes are expected to stain targets,

some epitopes of which are not exposed in a complex.

Spatial relationship among CCS-associated proteins

Using antiserum-derived Fab probes, we visualized the distribution of eight CCS-

associated proteins, which were clathrin light chain (CLC), adaptor protein 2 complex

beta subunit (AP2β2), EGFR, Grb2, transferrin receptors (TfR), FCHo1 and 2 (FCHo1/2),

Eps15 and intersectin-1 (ITSN1) (Fig. 1A). These Fab probes were verified to label the

endogenous targets in HeLa cells (figs. S1–8, Table S1, and Methods). We improved the

localization accuracy using normalized Gaussian with mean localization and standard

error (SE) of a probe bound over multiple frames (*9*) (Methods). In the localization

distribution of a long-binding probe, the normalized Gaussian represents the probability

distribution of the true target location, according to the central limit theorem (Fig. 1B,

bottom graph). The SE decreased with the number of the localizations, reaching around

1 nm (fig. S9A). The mode of the SEs was reduced to 10.7 nm compared to 24.2 nm in

the standard deviations (SDs) of the localization distribution (fig. S9B, C). The resolution

of the Gaussian rendering IRIS image with mean localization is 2.26 times better than

that of the IRIS image reconstructed by integrating the localizations (Fig. 1C and fig. S10).

The labeling densities in the target-localized area were 1.5 – 6.6 or higher per 10 nm pixel

(fig. S11E). These densities exceed the maximum occupied density (0.9 per 10 nm pixel)

of an antibody (12 nm in size) or an EGFR dimer (11.9 nm in size) (*17*).

The IRIS images with improved resolution and high-density labeling revealed clusters

of the eight proteins in a single CCS (Fig. 1D). Cells were stimulated with a saturated

concentration (100 ng/ml) of EGF at room temperature for delayed endocytosis (*18*) (fig.

S2A, B). Under this condition, EGFR and Grb2 clusters were localized at the outer rim

of the CCS, the width of which ranged from around 50 nm to 100 nm (Fig. 1E, arrows).

FCHo1/2, Eps15 and ITSN1 clusters were found at the interior of Grb2 clusters (Fig. 1G–

I, arrows). Interestingly, EGFR clusters were more extended outward at the CCS rim than

Grb2 clusters (Fig. 1F, arrow), suggesting either stepwise formation of EGFR-Grb2

complex or the recruitment of Grb2 by a molecule(s) other than EGFR.

Upon EGF stimulation at 37°C, EGFR and Grb2 clusters were localized at CCS rims

at 3 min (fig. S12B, yellow arrowheads). However, the number of EGFR clusters

decreased at 5 min (fig. S12C, cyan arrowheads), suggesting internalization of EGFR.

Interestingly many Grb2 clusters without EGFR clusters remained at the CCS rims (fig.
S12C, green arrowheads). These clusters also colocalized with mCherry (mCh)-SOS1
(Ras GEF) and cCbl-mCh (E3 ligase of EGFR) (*19*) in IRIS images with anti-mCh
nanobody (*20*) (fig. S13C, D). These results suggest that EGFR and Grb2 clusters may
contain molecular complexes of different compositions.
98
PC-coloring for multiple protein clusters
To map the molecular composition at the colocalization site of the targets, we
developed an image analysis, Protein Cluster coloring (PC-coloring). PC-coloring
classifies pixels in the merged IRIS image by principal component analysis (PCA) and
clustering based on the ratio of the standardized (z-score) intensities of multiple targets
(Fig. 2A, top panels). A PCA score image was created by plotting the raw intensity of the
targets (Fig. 2A, top left panel) in the coordinates of the first and second principal
components (PCs) (Fig. 2a, top center panel). In the PCA score image, EGFR (red)-
dominant, Grb2 (green)-dominant, Eps15 (cyan)-dominant and AP2β2 (blue)-dominant
pixels were located away from the center of the image. The pixels were then clustered
using k-means method, a commonly used clustering method (Fig. 2A, top right panel).
The clustered pixels are shown by eight-colored circles on a PCA cluster image (Fig. 2A,
bottom left panel). This coloring was performed based on the similarity of the target
intensity ratio in the dendrogram, the mean z-score intensities of the target (Fig. 2A,
middle panels) and the position of the clustered pixels in the PCA cluster image (Fig. 2A,
bottom left panel and Methods). The colors of the clustered pixels were then assigned on
the original IRIS image, generating a super-resolved (SR) cluster image (Fig. 2A, bottom
center panel). In the SR cluster image, EGFR-dominant (red), EGFR- and Grb2-dominant
(yellow), Grb2-dominant (green) pixels and orange pixels were localized in this order
from the outside of the CCS rim (Fig. 2B, C).
Colocalization was further characterized by the Pearson correlation coefficient
between the targets in each colored pixel (Fig. 2F–H and fig. S14). The correlation
coefficients between EGFR and Grb2 were 0.79, 0.93 and 0.76 in red, yellow, and green
pixels, respectively (Fig. 2F and fig. S14B, black arrows). These high correlation
coefficients at a 5 nm pixel size, which is close to the size of the protein, strongly indicate
the formation of the complex of EGFR and Grb2. In other colors, the correlation
coefficient was low. These results indicate complex formation of EGFR and Grb2 in
specific regions near the CCS rim (Fig. 2B). We also noted that the correlation coefficient
between Grb2 and Eps15 was 0.8 in orange pixels, indicating the complex formation of
Grb2 and Eps15 in orange pixels (Fig. 2C, G, arrow and fig. S14B, open arrow), which

is not apparent in the original IRIS image (Fig. 1D). The slopes of the regression lines in

green, yellow and red pixels were larger in this order (Fig. 2H). This increase in the

composition ratio of Grb2 to EGFR toward the center of CCSs suggests gradual complex

formation of EGFR with Grb2. PC-coloring in combination with the correlation analysis

allows us to identify the sites of complex formation between specific combinations of

targets in multiplexed super-resolution imaging.

Laminar distribution of the protein clusters at CCS rims

To further examine colocalization sites of CCS components (Eps15, FCHo1/2 and

ITSN1) with EGFR and Grb2, we created SR cluster images (Fig. 3 and figs. S15–17). In

the SR cluster image of EGFR, Grb2 and Eps15, EGFR-dominant (orange), EGFR-Grb2-

dominant (yellow) and Grb2-dominant (green) pixels lined up from the outside the CCS

rim (Fig. 3A). Grb2-Eps15-dominant (magenta) pixels were localized along the edges of

Grb2-dominant (green) pixels. On the central side of the CCS, EGFR-Eps15-dominant

(cyan) and Eps15-dominant (blue) pixels were localized. Among these color pixels, the

yellow, magenta and cyan pixels had a high correlation coefficient of 0.82 between EGFR

and Grb2, 0.8 between Grb2 and Eps15, and 0.78 between EGFR and Eps15, respectively,

indicating their complex formation (Fig. 3A, red arrows). Eps15 are known to be bound

to EGFR upon EGF stimulation (*21*). The distribution of cyan pixels indicates that EGFR

may form a complex with Eps15 primarily on the central side of the CCS, not in the EGFR

clusters at the outer rim of CCSs. Notably, colocalization of Grb2 and Eps15 (magenta

pixels) was confined to the boundaries of the Grb2 clusters and the CCS rim. In other SR

cluster images, magenta pixels were also confined to the boundaries (Fig. 3B, C). In the

scatter plots of Grb2 and Eps15, Eps15 intensity increased with Grb2 intensity at a certain

intensity ratio in the magenta pixels (Fig. 3E, arrow and fig. S18C). This indicates

complex formation of Grb2 and Eps15 with a certain composition ratio in most magenta

pixels. On the other hand, in the scatter plots of EGFR and Eps15, the distribution of

magenta pixels showed only a partial correlation compared that of the cyan pixels (Fig.

3F, arrows, and fig. S18C). This suggests that the complex of EGFR and Eps15 is not

formed in a large fraction of magenta pixels. This was also seen in the scatter plots of

other CCS components (fig. S19C, S20C, right graph, arrows). These results indicate that

CCS rims are layered from the outside with EGFR-dominant sites, the sites of EGFR-

Grb2 complex formation, and Grb2-dominant sites, and the sites of Grb2-CCS component

complex formation.

Next, we examined a spatial relationship among FCHo1/2, Eps15 and ITSN1 (Fig. 4A

and fig. S21A). In the SR cluster image, FCHo1/2-dominant (red), FCHo1/2-Eps15-

dominant (yellow) and Eps15-dominant (green) pixels were localized at the CCS rim (Fig.

4B and fig. S21B). At the interior of these colored pixels, Eps15-ITSN1-dominant

(purple) and ITSN1-dominant (blue) pixels were localized. Among these color pixels,

yellow and purple pixels had a high correlation coefficient of 0.73 between FCHo1/2 and

Eps15, and 0.78 between Eps15 and ITSN1, respectively (Fig. 4C, arrows). FCHo1/2-

ITSN1-dominant (orange) and FCHo1/2-Eps15-ITSN1-dominant (pale pink) pixels also

had a high correlation coefficient, but these clustered pixels were small and dispersed

(Fig. 4B, C). Overexpression of mCh-FCHo2 highlighted its localization to the CCS rim

(fig. S22A). The SR image again shows that the interior of Eps15-FCHo1/2 dominant

pixels is lined with Eps15-ITNS1 dominant pixels (fig. S22A). These results indicate that

Eps15 colocalizes with different partners from the rim to the center of the CCS in a

stepwise manner, which is not apparent in the original IRIS image (Fig. 4A and fig. S21A).

In the scatter plots, FCHo1/2 and ITSN1 intensity increased with Eps15 intensity at a

certain ratio in yellow and purple pixels, respectively (fig. S23C, arrows). FCHo1/2,

Eps15 and ITSN1 are known to bind to each other (*5*, *22–24*). However, our results

suggest that Eps15 forms a complex with FCHo1/2 and ITSN1 in distinct regions. These

results reveal the laminar distribution of distinct clusters composed of FCHo1/2, Eps15

and ITSN1 from the rim to the center of CCSs.

Endocytosis sites of membrane receptors

We investigated the sites of EGFR and TfR clusters in a CCS. In the merged IRIS image,

TfR clusters were localized in spots (fig. S24A, cyan arrowheads). The SR cluster image

showed that TfR-dominant pixels (cyan) were separated from EGFR-dominant (red) and

Grb2-dominant (green) pixels (fig. S24B). In the scatter plots of TfR and EGFR or Grb2,

the TfR-dominant pixels (cyan) were largely separated from both EGFR-dominant (red)

and Grb2-dominant (green) pixels (fig. S24E, center and right, arrows), compared to the

position of the red and green pixels in the scatter plot of EGFR and Grb2 (fig. S24E, left,

arrows). Since EGFR and TfR are expected to accumulate at the endocytic sites, the

separation of these dominant pixels suggest that their endocytic sites are different.

Discussion

Colocalization analyses using single-molecule localization-based super-resolution

microscopy have been reported for Clus-Doc (*25*), ClusterViSu (*26*) and Coloc-Tesseler

(*27*), but these analyses are performed on only two targets. In this study, we developed

PC-coloring for colocalization analysis of three or more targets. In the first step, PC-

coloring creates the color map of colocalization sites characterized by the intensity ratio

and correlation coefficient of the targets. Hence, in PC-coloring, PCA extracts features
about molecular complexes from the multiplexed intensity data. By assigning the features
on the original image, their spatial relationship is visualized. In the second step, the
detailed correlation between individual target pairs is further examined in the scatter plots.
A high correlation coefficient may reflect the formation of the molecular complexes
because correlation is analyzed with highly accurate quantitative data in protein-size
pixels. PC-coloring was enabled by the advantages of IRIS. The high-density labeling
with the antiserum-derived Fab probes allowed faithful visualization of distributions of
multiple endogenous proteins. This high density also allowed improved resolution using
the large number of probes bound over multiple frames. These resulted in greater
separation of the pixels by PCA.
Immunoelectron microscopy studies reported that EGFR is localized inside of CCSs
upon EGF stimulation (3, 4). In the present study, the clusters of EGFR and Grb2 were
found at the CCS rim. The discrepancy may arise from the temperature or time of EGF
simulation (4°C vs room temperature or 37°C for 3 min). At the CCS rim, the localizations
of various endocytic proteins have been reported (2, 5, 6). In addition, many clathrin-
coated microcages of 50 – 60 nm diameter are reported to emerge from CCS rims of cells
treated with a proton ionophore at pH 6.3 (28), similar to pH inside endocytic vesicles
containing EGF (29). Therefore, the CCS rim has been considered an endocytic site of
receptor (6, 30). EGF-induced colocalization of EGFR, Grb2 and CCS components
shown in this study provides direct evidence for endocytosis at the CCS rim. In addition,
the separation of EGFR-dominant and Grb2-dominant regions from TfR-dominant
regions suggests EGFR-specific endocytosis sites at the CCS rim. This is consistent with
the facts that Grb2 is required for endocytosis of EGFR, but not TfR (31), and that the
sorting of EGFR and TfR occurs early in the endocytosis process on the CCSs (32).
Furthermore, the Grb2 clusters between the EGFR and CCS component clusters suggests
that Grb2 anchors EGFR to the CCS components. This is supported from the domain
structure of Grb2, in which its SH2 domain binds to phosphorylated EGFR, and its two
SH3 domain bind to the proline-rich domains of Eps15 and Dynamin (33, 34). The half-
life of Grb2 binding to EGFR is about five sec (35), which is shorter than the endocytosis
process of 1 – 2 min (36). Thus, Grb2 may translocate from EGFR to CCS components
to form complexes. Our results indicate six layers in CCSs from the outside rims (Fig.
4D), as follows: (i) EGFR-dominant sites, (ii) the sites of EGFR-Grb2 complex formation,
(iii) Grb2-dominant sites, (iv) the sites of Grb2-CCS component complex formation, (v)
the sites of Eps15-FCHo1/2 complex formation, (vi) the sites of Eps15-ITSN1 complex
formation. After EGF endocytosis, the remaining Grb2 clusters could trap another EGFR

cluster. PC-coloring can update the *in vitro* network of protein interactions (*13*, *14*) to the

in vivo map of protein complexes with their composition ratio.

References

M. Mettlen, P. H. Chen, S. Srinivasan, G. Danuser, S. L. Schmid, Regulation of Clathrin-Mediated Endocytosis. *Annual review of biochemistry* **87**, 871-896 (2018) doi:10.1146/annurev-biochem-062917-012644

F. Tebar, T. Sorkina, A. Sorkin, M. Ericsson, T. Kirchhausen, Eps15 is a component of clathrin-coated pits and vesicles and is located at the rim of coated pits. *The Journal of biological chemistry* **271**, 28727-28730 (1996) doi:10.1074/jbc.271.46.28727

L. E. Johannessen, N. M. Pedersen, K. W. Pedersen, I. H. Madshus, E. Stang, Activation of the epidermal growth factor (EGF) receptor induces formation of EGF receptor- and Grb2-containing clathrin-coated pits. *Molecular and cellular biology* **26**, 389-401 (2006). doi:10.1128/MCB.26.2.389-401.2006

M. Kazazic, V. Bertelsen, K. W. Pedersen, T. T. Vuong, M. V. Grandal, M. S. Rødland, L. M. Traub, E. Stang, I. He. Madshus, Epsin 1 is involved in recruitment of ubiquitinated EGF receptors into clathrin-coated pits. *Traffic* **10**, 235-245 (2009) doi:10.1111/j.1600-0854.2008.00858.x

L. Ma, P. K. Umasankar, A. G. Wrobel, A. Lymar, A. J. McCoy, S. S. Holkar, A. Jha, T. Pradhan-Sundd, S. C. Watkins, D. J. Owen, L. M. Traub, Transient Fcho1/2·Eps15/R·AP-2 Nanoclusters Prime the AP-2 Clathrin Adaptor for Cargo Binding. *Developmental Cell* **37**, 428-443 (2016). doi:10.1016/j.devcel.2016.05.003

K. A. Sochacki, A. M. Dickey, M. P. Strub, J. W. Taraska, Endocytic proteins are partitioned at the edge of the clathrin lattice in mammalian cells. *Nature cell biology* **19**, 352-361 (2017). doi:10.1038/ncb3498

M. Mund, J. A. van der Beek, J. Deschamps, S. Dmitrieff, P. Hoess, J. L. Monster, A. Picco, F. Nédélec, M. Kaksonen, J. Ries, Systematic Nanoscale Analysis of Endocytosis Links Efficient Vesicle Formation to Patterned Actin Nucleation. *Cell* **174**, 884-896 (2018). doi: 10.1016/j.cell.2018.06.032

F. Balzarotti, Y. Eilers, K. C. Gwosch, A. H. Gynnå, V. Westphal, F. D. Stefani, J. Elf, S. W. Hell, Nanometer resolution imaging and tracking of fluorescent molecules with minimal photon fluxes. *Science* **355**, 606-612 (2017). doi: 10.1126/science.aak9913

S. C. M. Reinhardt, L. A. Masullo, I. Baudrexel, P. R. Steen, R. Kowalewski, A. S. Eklund, S. Strauss, E. M. Unterauer, T. Schlichthaerle, M. T. Strauss, C. Klein, R. Jungmann, Ångström-resolution fluorescence microscopy. *Nature* **617**, 711-716 (2023). doi: 10.1038/s41586-023-05925-9

10   M. Sauer, Localization microscopy coming of age: from concepts to biological
impact. *Journal of cell science* **126**, 3505-3513 (2013). doi: 10.1242/jcs.123612

11   P. Kanchanawong, C. M. Waterman, Localization-based super-resolution
imaging of cellular structures. *Methods Mol. Biol* **1046**, 59-84 (2013). doi:
10.1007/978-1-62703-538-5_4

12   Q. Zhang, A. Miyamoto, S. Watanabe, T. Arimori, M. Sakai, M. Tomisaki, T.
Kiuchi, J. Takagi, N. Watanabe, Engineered fast-dissociating antibody fragments
for multiplexed super-resolution microscopy. *Cell reports Methods* **2**, 100301
(2022). doi: 10.1016/j.crmeth.2022.100301

13   E. M. Schmid, H. T. McMahon, Integrating molecular and network biology to
decode endocytosis. *Nature* **448**, 883-888 (2007). doi: 10.1038/nature06031

14   H. T. McMahon, E. Boucrot, Molecular mechanism and physiological functions
of clathrin-mediated endocytosis. *Nature reviews molecular cell biology* **12**, 517-
533 (2011). doi: 10.1038/nrm3151

15   T. Kiuchi, M. Higuchi, A. Takamura, M. Maruoka, N. Watanabe, Multitarget
super-resolution microscopy with high-density labeling by exchangeable probes.
Nature methods **12**, 743-746 (2015). doi:10.1038/nmeth.3466

16   T. Miyoshi, Q. Zhang, T. Miyake, S. Watanabe, H. Ohnishi, J. Chen, H. D
Vishwasrao, O. Chakraborty, I. A. Belyantseva, B. J. Perrin, H. Shroff, T. B.
Friedman, K. Omori, N. Watanabe, Semi-automated single-molecule microscopy
screening of fast-dissociating specific antibodies directly from hybridoma
cultures. *Cell reports* **34**, 108708 (2021). doi:10.1016/j.celrep.2021.108708

17   S. R. Needham, S. K. Roberts, A. Arkhipov, V. P. Mysore, C. J. Tynan, L. C.
Zanetti-Domingues, E. T. Kim, V. Losasso, D. Korovesis, M. Hirsch, D. J. Rolfe,
D. T. Clarke, M. D. Winn, A. Lajevardipour, A. H. A. Clayton, L. J. Pike, M.
Perani, P. J. Parker, Y. Shan, D. E. Shaw, M. L. Martin-Fernandez, EGFR
oligomerization organizes kinase-active dimers into competent signalling
platforms. *Nature Communications* **7**, 13307 (2016). doi: 10.1038/ncomms13307

18   G. M. Hillman, J. Schlessinger, Lateral diffusion of epidermal growth factor
complexed to its surface receptors does not account for the thermal sensitivity of
patch formation and endocytosis. *Biochemistry* **21**, 1667-1672 (1982).
doi:10.1021/bi00536a030

19   G. Levkowitz, H. Waterman, S. A. Ettenberg, M. Katz, A. Y. Tsygankov, I. Alroy,
S. Lavi, K. Iwai, Y. Reiss, A. Ciechanover, S. Lipkowitz, Y. Yarden, Ubiquitin
ligase activity and tyrosine phosphorylation underlie suppression of growth factor
signaling by c-Cbl/Sli-1. *Molecular Cell* **4**, 1029-1040 (1999).

doi:10.1016/s1097-2765(00)80231-2

20   P. C. Fridy, Y. Li, S. Keegan, M. K. Thompson, I. Nudelman, J. F. Scheid, M.
Oeffinger, M. C. Nussenzweig, D. Fenyö, B. T. Chait, M. P. Rout, A robust
pipeline for rapid production of versatile nanobody repertoires. *Nature methods*
11, 1253-1260 (2014). doi:10.1038/nmeth.3170

21   S. van Delft, C. Schumacher, W. Hage, A. J. Verkleij, P. M. van B. en
Henegouwen, Association and colocalization of Eps15 with adaptor protein-2 and
clathrin. *The Journal of cell biology* **136**, 811-821 (1997). doi:
10.1083/jcb.136.4.811

22   A. S. Sengar, W. Wang, J. Bishay, S. Cohen, S. E. Egan, The EH and SH3 domain
Ese proteins regulate endocytosis by linking to dynamin and Eps15. *The EMBO*
Journal **18**, 1159-1171 (1999). doi:10.1093/emboj/18.5.1159

23   T. Koh, V. I. Korolchuk, Y. P. Wairkar, W. Jiao, E. Evergren, H. Pan, Y. Zhou,
K. J. T. Venken, O. Shupliakov, I. M. Robinson, C. J. O'Kane, H. J. Bellen, Eps15
and Dap160 control synaptic vesicle membrane retrieval and synapse
development. *The Journal of cell biology* **178**, 309-322 (2007).
doi:10.1083/jcb.200701030

24   W. M. Henne, E. Boucrot, M. Meinecke, E. Evergren, Y. Vallis, R Mittal, H. T.
McMahon, FCHo proteins are nucleators of clathrin-mediated endocytosis.
Science **328**, 1281-1284 (2010). doi: 10.1126/science.1188462

25   S. V. Pageon, P. R. Nicovich, M. Mollazade, T. Tabarin, K. Gaus, Clus-DoC: a
combined cluster detection and colocalization analysis for single-molecule
localization microscopy data. *Molecular biology of the cell* **27**, 3627-3636 (2016).
doi: 10.1091/mbc.E16-07-0478

26   L. Andronov, I. Orlov, Y. Lutz, J. Vonesch, B. P. Klaholz, ClusterViSu, a method
for clustering of protein complexes by Voronoi tessellation in super-resolution
microscopy. *Scientific reports* **6**, 24084 (2016). doi: 10.1038/srep24084

27   F. Levet, G. Julien, R. Galland, C. Butler, A. Beghin, A. Chazeau, P. Hoess, J.
Ries, G. Giannone, J. B. Sibarita, A tessellation-based colocalization analysis
approach for single-molecule localization microscopy. *Nature Communications*
10, 2379 (2019). doi: 10.1038/s41467-019-10007-4

28   J. Heuser, Effects of cytoplasmic acidification on clathrin lattice morphology. *The*
Journal of cell biology **108**, 401-411 (1989). doi:10.1083/jcb.108.2.401

29   C. H. Heo, M. K. Cho, S. Shin, T. H. Yoo, H. M. Kim, Real-time monitoring of
vesicle pH in an endocytic pathway using an EGF-conjugated two-photon probe.
Chemical Communications **52**, 14007-14010 (2016). doi:10.1039/c6cc08036g

30 M. Lampe, S. Vassilopoulos, C. Merrifield, Clathrin coated pits, plaques and
adhesion. *Journal of structural biology* **196**, 48-56 (2016).
doi:10.1016/j.jsb.2016.07.009

31 F. Huang, A. Khvorova, W. Marshall, A. Sorkin, Analysis of clathrin-mediated
endocytosis of epidermal growth factor receptor by RNA interference. *The*
Journal of biological chemistry **279**, 16657-16661 (2004).
doi:10.1074/jbc.C400046200

32 D. Leonard, A. Hayakawa, D. Lawe, D. Lambright, K. D. Bellve, C. Standley, L.
M. Lifshitz, K. E. Fogarty, S. Corvera, Sorting of EGF and transferrin at the
plasma membrane and by cargo-specific signaling to EEA1-enriched endosomes.
Journal of cell science **121**, 3445-3458 (2008). doi:10.1242/jcs.031484

33 Z. Wang, M. F. Moran, Requirement for the adapter protein GRB2 in EGF
receptor endocytosis. *Science* **272**, 1935-1939 (1996).
doi:10.1126/science.272.5270.1935

34 C. Parachoniak, M. Park, Distinct recruitment of Eps15 via Its coiled-coil domain
is required for efficient down-regulation of the met receptor tyrosine kinase. *The*
Journal of biological chemistry **284**, 8382-8394 (2009). doi:
10.1074/jbc.M807607200

35 M. Morimatsu, H. Takagi, K. G. Ota, R. Iwamoto, T. Yanagida, Y. Sako,
Multiple-state reactions between the epidermal growth factor receptor and Grb2
as observed by using single-molecule analysis. *Proceedings of the National*
Academy of Sciences of the United States of America **104**, 18013-18018 (2007).
doi:10.1073/pnas.0701330104

36 D. Loerke, M. Mettlen, D. Yarar, K. Jaqaman, H. Jaqaman, G. Danuser, S. L.
Schmid, Cargo and dynamin regulate clathrin-coated pit maturation. *PloS Biology*
7, e1000057 (2009). doi: 10.1371/journal.pbio.1000057

37 T. Kiuchi, E. Ortiz-Zapater, J. Monypenny, D. R. Matthews, L. K. Nguyen, J.
Barbeau, O. Coban, K. Lawler, B. Burford, D. J. Rolfe, E. de Rinaldis, D. Dafou,
M. A. Simpson, N. Woodman, S. Pinder, C. E. Gillett, V. Devauges, S. P. Poland,
G. Fruhwirth, P. Marra, Y. L. Boersma, A. Plückthun, W. J. Gullick, Y. Yarden,
G. Santis, M. Winn, B. N. Kholodenko, M. L. Martin-Fernandez, P. Parker, A.
Tutt, S. M. Ameer-Beg, T. Ng, The ErbB4 CYT2 variant protects EGFR from
ligand-induced degradation to enhance cancer cell motility. *Science signaling* **7**,
ra78 (2014). doi:10.1126/scisignal.2005157

38 M. Higuchi, K. Ishiyama, M. Maruoka, R. Kanamori, A. Takaori-Kondo, N.
Watanabe, Paradoxical activation of c-Src as a drug-resistant mechanism. *Cell*

reports **34**, 108876 (2021). doi: 10.1016/j.celrep.2021.108876

39   M Ovesný, P. Křížek, J. Borkovec, Z. Svindrych, G. M. Hagen, ThunderSTORM:
a comprehensive ImageJ plug-in for PALM and STORM data analysis and super-
resolution   imaging.   *Bioinformatics*   **30**,   2389-2390   (2014).   doi:
10.1093/bioinformatics/btu202

40   G. Gut, D. M. Herrmann, L. Pelkmans, Multiplexed protein maps link subcellular
organization   to   cellular   states.   *Science*   **361**,   eaar7042   (2018).   doi:
10.1126/science.aar7042

41   J. Heuser, Three-dimensional visualization of coated vesicle formation in
fibroblasts.   *The   Journal   of   cell   biology*   **84**,   560-583   (1980).
doi:10.1083/jcb.84.3.560

42   J. M. Larkin, W. C. Donzell, R. G. Anderson, Potassium-dependent assembly of
coated pits: new coated pits form as planar clathrin lattices. *The Journal of cell*
biology **103**, 2619-2627 (1986). doi:10.1083/jcb.103.6.2619

METHODS

Plasmids and reagents

Expression plasmids for human CLC (#57452), human TfR (#61505), human Eps15 (#27696), human ITSN1 Long (#47395), mouse FCHo2 (#27686), human Epsin2 (#22242), mouse Hip1r (#), human Dynamin2 (#27689) and human AP2μ2 (#27672) were purchased from Addgene. cDNA encoding human AP2β2 (Clone ID: 3532472) and human FCHo1 (Clone ID: 5218319) were purchased from Horizon (Dharmacon). cDNA encoding nanobodies for anti-EGFP and anti-mCh were synthesized by gBlocks Gene Fragments (Integrated DNA Technolodies) according to reported sequences (Clone ID: LaG-3 and LaM-1) (20). Expression plasmids for human EGFR, human cCbl and human Grb2 were previously described (37, 38). These cDNAs and their fragments were inserted into pGEX-6P-1 vector using PCR for protein purification from E. coli. EGF was purchased from Pepro Tech. DyLight 488 NHS Ester was purchased from Thermo Fisher Scientific.

Cell culture

HeLa cells were cultured in DMEM supplemented with 10% fetal bovine serum. For stimulation with 100 ng/ml EGF, the cells were serum-starved overnight. For plasmid transfection, FuGENE HD (Promega) was used, and the cells were cultured for 24 hours before IRIS imaging.

Purification of antigen proteins and nanobodies

In this study, 4 to 5 mg of a recombinant protein per a Fab probe was required for rabbit antiserum preparation, the affinity purification of polyclonal antibodies and Fab probe purification. To increase the yield of the recombinant protein, a general purification protocol for glutathione S-transferase (GST)-fused protein from E. coli was improved using an excess of glutathione resins. BL21(DE3) competent cells were transformed with pGEX-6P-1 vector encoding GST-fused protein. After 35 μM IPTG induction, 6L cultures of the cells were incubated for 16 to 20 hours at 20°C with shaking. The collected cells were lysed in 50 ml lysis buffer (20 mM Tris pH7.3, 150 mM NaCl, 0.25% Triton X-100, 1 mM DTT) containing protease inhibitor cocktail (Nacalai Tesque) and sonicated. Glutathione resins (GST-Accept, Nacalai Tesque) in 30 to 60 ml bed volume was added to the supernatant of the cell lysate, and then was rotated for 90 min at 4 °C. The resins were transferred to two or three columns constructed from a 50-ml plastic syringe. The resins were washed with the lysis buffer and a high Na buffer (20 mM Tris pH7.3, 500 mM NaCl, 0.25% Triton X-100, 1 mM DTT), and three times for 5 min with a buffer A

(20 mM Tris pH7.3, 500 mM NaCl, 5 mM MgCl, 0.25% Triton X-100, 1 mM DTT, 1mM

ATP). The resins were further washed five times with a PreScission reaction buffer (50

mM Tris pH7.3, 150 mM NaCl, 1 mM EDTA, 1 mM DTT) for antiserum preparation.

For affinity column preparation, the resins were washed five times with a $NaHCO_3$ buffer

(200 mM $NaHCO_3$, 500 mM NaCl) to remove primary amines from the buffer. Two ml

of 50% slurry of the resins were aliquoted into thirty to sixty 2-ml tubes. The resins were

subject to PreScission protease and rotated overnight at 4°C to elute the target proteins.

The use of the 2-ml tubes is critical in this elution step. For unknown reasons, some

proteins such as FCHo1, Dyn2 and cCbl were reproducibly lost in this step when 50-ml

or 15-ml tubes were used, even if low-protein adsorption tubes (PROTEOSAVE,

Sumitomo Bakelite Co.) were used. The eluted protein solution was in turn transferred to

an ultrafiltration column (Amicon Ultra-15, Merck Millipore) and concentrated by

centrifugation. Eighty to ninety percentage of the eluted protein was recovered from the

column, if the water level of the solution was not lowered below the height of the

column's filter during centrifugation. The resins were again filled with the buffer to obtain

the second and third eluates, which were added to the first eluate and all concentrated to

2 ml. Since the predicted molecular weights of TfR1-60, TfR93-179, Grb2 109-159 and

Grb2 160-217 are small (6–10 kDa), for antiserum preparation, the GST-fused protein

fragments were eluted with 20 mM reduced glutathione. The reduced glutathione was

removed by dialysis. By this protocol using an excess of glutathione resins, the yield of

EGFR 985-1210, which had a low yield, increased about 11-fold. Seventeen antigens,

some with molecular weights around 100 kD, were obtained in milligram quantities in

one or two experiments, for FCHo2 in three or four experiments (Table S2). Nanobodies

for EGFP and mCh were purified using the usual amount of the glutathione resins from

BL21 cells because of their high yield.

The used resins were collected and reused after being cleaned according to the cleaning

protocol for a glutathione column (GSTrap™ FF, Cytiva). The resins were transferred to

two or three columns constructed from a 50-ml plastic syringe. GST remained on the

resins was removed by adding 20 mM reduced glutathione three times. The resins were

washed five times with PBS. Precipitated substances were removed with 6 M guanidine

hydrochloride (denaturing agent). The resins were washed fifteen times with PBS.

Hydrophobically bound substances were removed by adding 70% ethanol three times.

The resins were stored in 20% ethanol at 4°C. No protein bands were detected when the

cleaned resins were subjected to SDS–polyacrylamide gel electrophoresis.

Production of antiserum-derived Fab probes

IgG antibodies in a rabbit antiserum were purified by a Protein A column (HiTrap
rProteinA FF 5 ml, Cytiva). After dialysis with PBS, the IgG antibodies was applied to
an affinity column (HiTrap NHS-activated HP 1 ml, Cytiva) coupled to 1.5–3 mg of its
antigen protein. For IgG antibodies from antiserum against GST-tagged TfR 1-60, TfR
93-179, Grb2 109-159 or Grb2 160-217, the affinity column was coupled to GST-cleaved
TfR 1-60, TfR 93-179 or Grb2 full length, respectively. In addition, the column was
linked to a GST-coupled affinity column to remove anti-GST antibodies. The polyclonal
antibodies were eluted with 100 mM glycine buffer at pH 3.5 (or 3.0) and pH 2.0. The
pH of the eluate was immediately neutralized with 1 M Tris buffer at pH 9.0, and then
dialyzed in PBS. To find the eluted fraction that can stain for endogenous targets, HeLa
cells were immunostained. For fluorescent labeling, the fraction was added to a 200 μl
bed volume of antigen-conjugated resins (NHS-activated Sepharose 4 Fast Flow, Cytiva)
to protect their antigen binding sites. The antigen bound to the resins was more than twice
the molar amount of the antibodies in the fraction. After overnight rotation at 4°C, the
antibodies were labeled with twice the molar amount of Dylight 488 NHS Ester as the
total amount of the antigens and antibodies on the resins. After rotation at room
temperature for 1 hour, the resins were transferred to a low-protein adsorption 1.5-ml
tubes (PROTEOSAVE, Sumitomo Bakelite Co.), whose bottom was perforated with a
fine needle. The tube was placed on top of a 2-ml tube to construct a homemade microspin
column. The resins were washed five times with PBS. The antibodies were eluted three
times with 200 μl of 100 mM glycine buffer at pH 2.0. The pH of the eluate was
immediately neutralized with 1 M Tris buffer at pH 9.0. The eluted antibodies were mixed
with the first and second eluates, and then digested with papain to generate Fab fragments.
Their Fc domains were removed using Protein A Sepharose CL-4B (Cytiva). To reduce
the non-specific signals of the Fab, the Fab was further bound to antigen-conjugated resin.
After overnight rotation at 4°C, the resins were transferred to the handmade microspin
column constructed from a 0.5-ml tube and a 1.5-ml tube. The resins were washed five
times each with 100 μl of PBS and 100 mM glycine buffer at pH 5.0. The Fabs were
eluted ten and five times with 100 μl of 100 mM glycine buffer at pH 3.0 and pH 2.0,
respectively. The pH of the eluate was immediately neutralized with 1 M Tris buffer at
pH 9.0. The labeling ratio of the Fabs was estimated to be about 0.6 by enzyme-linked
immunosorbent assay (ELISA). The fraction eluted at pH 3.0, which contains the highest
amount of the Fabs, was used for IRIS super-resolution imaging. In each of these
purification steps, the polyclonal antibodies and Fabs were placed in the low-protein
adsorption tubes to protect them from being lost. This protocol was established using anti-
CLC antiserum (fig. S1).

Multiplexed super-resolution imaging

Multiplexed super-resolution imaging

Multiplexed IRIS super-resolution imaging was performed as previously described (*15*).

HeLa cells on a 24 mm round coverslip were fixed with 3.7% PFA in a cytoskeleton

buffer (10 mM MES pH 6.1, 90 mM KCl, 3 mM $MgCl_2$, 2 mM EGTA) for 20 min, and

then permeabilized with 0.2% Triton-X100 in a HEPES buffered solution (10 mM pH

7.3, 90 mM KCL, 3 mM $MgCl_2$, 100 μM DTT) for 5 min. After blocking with 4% bovine

serum albumin in the HEPES buffered solution for 30 min, the coverslip was mounted in

an open chamber (A7816, Thermo Fisher Scientific). On the coverslip, a handmade

chamber with a volume of approximately 50 μl was constructed with a vacuum grease

and a 1 cm diameter plastic ring. As a retractable lid, a 15 mm round coverslip was placed

on top of the ring and another 24 mm round coverslip was placed on top of the open

chamber to double the lid. The cells were subjected to the IRIS probe in an imaging

solution (50 mM Tris pH 8.0, 90 mM KCl, 3 mM $MgCl_2$) with an oxygen scavenging mix

(5 U/ml pyranose oxidase, 10 mM cysteamine hydrochloride, 60 μg/ml catalase, 10%

glucose) (*12*). The probe concentrations in the imaging solution, measured as the

concentration of the conjugated fluorescent dye (DyLight 488), were 80 nM anti-EGFR

Fab, 40 nM anti-CLC Fab, 100 nM anti-Eps15 Fab, 40 nM anti-TfR Fab, 80 nM anti-

Grb2 Fab, 30 nM anti-AP2β2 Fab, 60 nM anti-ITSN1 Fab and 60 nM anti-FCHo1 Fab.

The single molecule speckle (SiMS) images were acquired by TIRF microscopy using an

inverted microscope (Olympus IX83-ZDC) equipped with an IX3-ZDC2 Z-drift

compensator (Olympus), an UPlanApo 60×/1.50 NA oil objective lens (Olympus),

ORCA-Flash4.0 V3 Digital CMOS camera (Hamamatus) and a 488 nm laser line (150

mW, Coherent). A set of a bright field image (exposure time, 500 ms) and 500 SiMS

images (exposure time, 100ms) was acquired repeatedly. The bright field images were

used to correct for stage drift (see below). The total number of SiMS images acquired was

as follows: 250,000 (anti-EGFR Fab), 100,000 (anti-CLC Fab), 300,000 (anti-Eps15 Fab),

200,000 (anti-TfR Fab), 400,000 (anti-Grb2 Fab), 200,000 (anti-AP2β2 Fab), 400,000

(anti-ITSN1 Fab) and 400,000 (anti-FCHo1 Fab). After IRIS imaging of each probe, the

probe was washed out 15 times with 0.2% Triton-X100 in the HEPES buffered solution

and 10 times with the HEPES buffered solution. The remaining fluorescence of the probe

was completely photobleached in the HEPES-buffered solution with the oxygen

scavenging mix.

Image reconstruction

Image reconstruction

In hundreds of thousands of the SiMS images, the 10^6 to 10^8 localizations of fluorescent

speckles were determined using ThunderSTORM plugin (*39*) in ImageJ
(http://rsb.info.nih.gov/ij/). Stage drift during long-time image acquisition for each probe
was corrected by an autocorrelation function using the bright field image of fixed cells
and non-fluorescent beads attached to the coverslip as previously described (*15*). After
the drift correction for each probe, stage drift between two different probes was corrected
by calculating cross-correlation functions between 30 bright field images per probe. The
30×30 cross-correlation functions were computed using the 10-fold magnified images
with a bicubic method (pixel size, 10.83 nm). The drift distances between the two probes
were calculated, and its standard error was around 1 nm.

To improve the localization accuracy, the mean localization and standard error of a
probe bound over two frames were used. The localizations of a long-binding probe were
distributed within a radius of 54 nm. The duration of the quenching state due to blinking
was one frame (exposure time: 100 ms). Thus, localizations that were continuous or one
frame apart within a radius of 54 nm were extracted as the localizations of the same probe
bound over two frames. To reduce the contribution of the first localization as a reference,
the mean of the extracted localizations was calculated. Using the mean localization as a
new reference, the localizations were re-extracted using the same filter. The mean and
standard error of the re-extracted localizations were calculated. These calculations were
computed using our plug-in with multi-threaded processing in ImageJ. In addition, the
accuracy of the drift correction using the bright field image was estimated from
differences between the drift-corrected localizations of the same probe in two SiMS
images acquired before and after the bright field image. The mean of the difference was
about 3 nm. Together with the accuracy of the drift correction between the two different
probes, the standard error of the localizations less than 5 nm was converted to 5 nm. Using
the mean localization and the standard error, an IRIS image was reconstructed at 5 nm
pixel size by using a Normalized Gaussian rendering in the ThunderSTORM. When
Gaussian rendering was not used, the IRIS image was reconstructed by integrating the
localizations as previously described (*15*).

PC-coloring
PC-coloring was computed using a set of our plug-ins in ImageJ and scripts in Matlab
(MathWorks). Each process using the ImageJ plug-ins or Matlab scripts is indicated at
the end of the following sentences as (ImageJ) or (Matlab), respectively.

A data set of the intensities of each target and the coordinates of its pixel in the IRIS
images was output (ImageJ). In the multiplexed intensity data, a pixel with target
intensities of all 0's was excluded. To match the intensity distribution among targets, the

intensity data was standardized (z-score) by each target (Matlab). To calculate the
intensity ratio at each pixel, if this z-score intensity data had a value less than 0, an
appropriate value was added to all the intensities to make them greater than or equal to 0
(Matlab). Using the intensity ratios, PCA was performed, and PCA scores were output
(Matlab). A binarized image was created from the summed IRIS image of each target to
exclude pixels with low intensity (ImageJ). Using the binarized image, the pixels with
low intensity were also removed from the PCA score data (ImageJ). To interpret the PCA
result, a PCA score image was created by plotting the raw intensity of each IRIS image
on the coordinates of the first and second PCs (ImageJ). Then the PCA score images of
each target were merged (ImageJ). The PCA score data were clustered by k-means
method (Matlab). The number of clusters set by the k-means method was 25 in this study.
The similarity of the intensity ratio between the clustered pixels was calculated as the
distance by group average method (ImageJ). The two clustered pixels with the closest
similarity were combined, and then the group average distance between the new clustered
pixels and the other clustered pixels was calculated for their linkage distances (ImageJ).
Using the linkage distances, a dendrogram was created (Matlab). The number of the
clustered pixels was renumbered in order from left to right of the leaf nodes in the
dendrogram (Matlab). The mean z-score intensities of each target in the cluster pixels
were output (ImageJ).
A SR cluster image was created by assigning the number of the clustered pixels to the
pixel in the original IRIS image (ImageJ). To examine the adjacent clustered pixels on
the SR cluster image, 8-connectivity analysis was performed (*40*). One pixel in the
clustered pixels in the SR cluster image was selected, and the cluster numbers of the
surrounding eight pixels were obtained (ImageJ). If any of the cluster numbers of the
eight pixels is different from that of the center pixel, the cluster number was recorded as
the pixel located at the border of the adjacent cluster (ImageJ). By performing this process
on all pixels in the clustered pixels, the cluster numbers of the boundary pixels were
tallied, and their percentages were output as proximity between the clustered pixels
(ImageJ). A boundary pixel without any cluster number were excluded from this
percentage.
A PCA cluster image was created by drawing circles on the coordinates of the first and
second PCs (ImageJ). The center of the circle was that of the clustered pixels. The area
of the circle was proportional to the number of pixels in the clustered pixels. The two
circles were connected by a line with a width proportional to the proximity between the
clustered pixels on the SR cluster image, generating a PCA cluster image with proximity
(ImageJ). In the PCA cluster image with proximity, the top three lines in their proximity

were shown. The sum of the three proximity, on average, exceeds 70% of all proximity. In the SR cluster image, adjacent clustered pixels were also adjacent in the PCA cluster image. This indicates that the adjacent relationship of the clustered pixels in the SR cluster image is also preserved in the PCA cluster image.

The clustered pixels in the SR cluster image and the PCA cluster image with proximity were colored (ImageJ). The clustered pixels were colored based on the dendrogram, the mean z-score intensity of target, and the position of the clustered pixels in the PCA cluster image (ImageJ). The clustered pixels with high similarity in intensity ratio at adjacent positions in the PCA cluster image were colored the same. The coloring was adjusted to reflect the distribution of the target intensities on the PCA score image (ImageJ). In each color pixels, the z-score intensity of the target was plotted on a scatter plot. The Pearson correlation coefficient was calculated (ImageJ). In PC-coloring, there is no principle limit to the number of targets. However, pixel separation by PCA decreases with the number of targets. In this study, three to four targets were practical.

ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant Number 24H01282, 21K06168, 19H04961, 17K07384, 15H01635 (T.K.) and 22H00456 (N.W.), by the Asahi Glass Foundation, Takeda Science Foundation (T.K.), Uehara Memorial Foundation (N.W.) and by CREST grant number JPMJCR15G5 (N.W.).

AUTHOR CONTRIBUTIONS

T.K. improved a method for protein purification from E. coli and developed a method for Fab probe production from antiserum, image reconstruction and PC-coloring. T.K. designed and conducted experiments including protein purification, Fab probe production and microscopic imaging. T.K developed all image analysis programs and analyzed the data. T.K. supervised the project and wrote the manuscript with support from N.W. R.K. contributed to the production of Fab probes for CLC and Eps15, and nanobodies for EGFP and mCherry, constructed expression plasmids and purified proteins. S.O. and L.E. purified proteins.

Fig. 1. Multiplexed IRIS super-resolution image of eight targets. (**A**) The merged IRIS image of eight targets reconstructed by integrating localizations at 20 nm sized pixel. HeLa cells were stimulated with 100 ng/ml EGF for 20 min at room temperature. (**B**) Improved localization accuracy using normalized Gaussian with mean localization. A fluorescent speckle of a long-binding probe (upper left). IRIS image reconstructed by integrating localizations of the long-binding probe at 5 nm sized pixel (upper right). In the bottom graph, the cross-sectional profile of the reconstructed image (blue) was plotted

to show the distribution of the localizations, whose standard deviation indicates the
localization accuracy. A normalized two-dimensional Gaussian with mean and SE of the
localizations (red) were plotted to show the accuracy of the mean localization. (**C**)
Merged IRIS images reconstructed by integrating localizations (upper) and by Gaussian
rendering with mean localizations (lower) in the part of the boxed area shown in **A**. (**D**)
Spatial relationship of the eight targets in the CCS in the box area shown in **A**. The IRIS
image was reconstructed by Gaussian rendering with mean localizations at 5 nm sized
pixel. (**E**) The enlarged image of the box area shown in **D**. (**F**–**I**) The merged IRIS image
of three targets in **E**. The target names are indicated in the panel.
665

Fig. 2. Colocalization analysis by PC-coloring. (A) Outline of PC-coloring (also see

Methods). The merged IRIS image of EGFR (red), Grb2 (green), Eps15 (cyan) and

AP2β2 (blue) shown in **Fig. 1D** (top left). The pixels in the IRIS image are classified by
principal component analysis (PCA) based on the ratios of the standardized (z-score)
intensities. A PCA score image is created by plotting the raw intensity of the target of the
IRIS image (top left) in the coordinates of the first and second principal components (PC1,
PC2) (top center). These pixels are clustered by k-means method (top right). A
dendrogram is created based on the similarity of the intensity ratio in the clustered pixels
(middle left). The mean z-score intensity of each target is calculated in the clustered pixels
(middle right). The number of the clustered pixels in the top right panel is the same as the
number shown in the dendrogram. The clustered pixels are shown by eight-colored circles
in the coordinates of the first and second PCs as a PCA cluster image (bottom left). The
center of the circle indicates that of the clustered pixels. Its area is proportional to the
number of pixels in the clustered pixels. These circles are connected by lines indicating
the proximity between the clustered pixels (Methods). The top three lines in the proximity
are shown. The circle is colored based on the dendrogram, the mean z-score intensities
and the position of the clustered pixels in the PCA cluster image (Methods). The colors
of the clustered pixels are assigned on the original IRIS image as a SR cluster image
(bottom center). The mean z-score intensity of each target is calculated in each color
(bottom right). (**B–E**) Extraction of the red, yellow and green pixels (**B**), the yellow, green
and orange pixels (**C**), the red, pale pink and cyan pixels (**D**), and the orange, pale pink,
cyan, slate blue, and blue (**E**) in the SR cluster image. (**F**, **G**) Scatter plots of z-score
intensities between EGFR and Grb2 (**F**), and between Grb2 and Eps15 (**G**). (**H**) Scatter
plots of red, yellow and green pixels. Their regression lines are drawn to estimate the
composition ratio of Grb2 to EGFR. The scatter plots of other target combination are
shown in **fig. S14**.
693

Fig. 3. Colocalization analysis of EGFR, Grb2 and Eps15 (A), FCHo1/2 (B) or ITSN1 (C). (**A–C**) From left to right: the merged IRIS images reconstructed by Gaussian rendering with mean localizations at 5 nm sized pixel, the SR cluster images, the PCA cluster image, the mean z-score intensities of the targets in the clustered pixels and the correlation coefficients between the targets. These PC-coloring with PCA and clustering were performed on the merged IRIS images shown in **figs. S15–17**. HeLa cells were stimulated with 100 ng/ml EGF for 20 min at room temperature. (**D–F**) Scatter plots of z-score intensities of the targets at the yellow, magenta and cyan pixels shown in **A**. These pixels were plotted between Grb2 and EGFR (**D**), between Grb2 and Eps15 (**E**), and between EGFR and Eps15 (**F**). The scatter plots of other target combination are shown in **fig. S18**.

Fig. 4. Colocalization analysis of FCHo1/2, Eps15 and ITSN1. (**A**) The merged IRIS
image of FCHo1/2 (red), Eps15 (green), ITSN1 (cyan) and AP2β2 (blue) was
reconstructed by Gaussian rendering with mean localizations at 5 nm sized pixel. (**B**) The
SR cluster image (left), the PCA score image (right upper) and the PCA cluster image
(right lower) of FCHo1, Eps15 and ITSN1. (**C**) The mean z-score intensities of the targets
in the clustered pixels (upper) and the correlation coefficients between the targets (lower).
These PC-coloring with PCA and clustering were performed on the merged IRIS images
shown in **fig. S21**. (**D**) laminar organization of EGFR, Grb2 and CCS components at CCS
rims in EGFR endocytosis.

A Immunostaining with purified antibody

B IRIS imaging with labeled Fab before Fab purification

C IRIS imaging with labeled Fab after Fab purification

Fig. S1. Development of a method for producing Fab probe from antiserum using
anti-clathrin light chain (CLC) antiserum (see Methods). (**A**) Immunostaining with
anti-CLC antibodies purified by affinity chromatography. The merged image of TagRFP-
CLC (red) and anti-CLC antibodies (green). HeLa cells were stained with the antibodies
to find the eluted fraction capable of staining endogenous CLC. Endogenous CLC was
localized in spots in non-expressing cell similar to expressed CLC (green arrow). (**B**) IRIS
imaging of CLC using fluorescent dye (Dylight 488)-conjugated Fab probe. The merged

image of the bright field image (gray) and the fluorescent image of TagRFP-CLC (red)
(leftmost). The merged image of the fluorescent image of TagRFP-CLC (red) and the
IRIS image using the Fab probe (green) in the boxed area in the leftmost image (three
images on the right). The purified anti-CLC antibodies were labeled while bound to CLC-
conjugated resin to protect its antigen-binding site, which stained endogenous CLC with
less non-specific signals by immunostaining. Their Fab fragments were generated by
papain digestion. In the IRIS image using the Fab probe, CLC was visualized in the CLC-
expressing cell (red arrow). But endogenous CLC was not visualized in the non-
expressing cell, where there were non-specific signals (green arrow). (**C**) IRIS imaging
of CLC using the Fab probe after Fab purification. The merged image of the bright field
image (gray) and the fluorescent image of TagRFP-CLC (red) (leftmost). The merged
image of the fluorescent image of TagRFP-CLC (red) and the IRIS image using the
purified Fab probe (green) in the boxed area in the leftmost image (three images on the
right). To reduce the non-specific signals of the Fab probe, the Fab probe was further
purified with CLC-conjugated resin. The IRIS image using the purified Fab probe showed
endogenous CLC in the non-expressing cell (double arrows).
742

Fig. S2. Preparation of anti-EGFR Fab probe. (**A**) Time-lapse imaging of EGFR-
tagRFP and EGFP-CLC. HeLa cells before and after stimulation with 100 ng/ml EGF for
20 min at room temperature for the delayed endocytosis of EGFR. (**B**) Enlarged images

in the boxed area shown in **A**. Even 20 min after the stimulation, EGFR was accumulating

in CCSs. (**C**) Immunostaining with anti-EGFR antibodies purified by affinity

chromatography. The merged image of EGFR-tagRFP (red) and anti-EGFR antibodies

(green). After EGF stimulation, HeLa cells were stained with the antibodies to find the

eluted fraction capable of staining endogenous EGFR. Endogenous EGFR was localized

in spots in non-expressing cell similar to expressed EGFR (arrow). (**D**) IRIS imaging

using anti-EGFR Fab probe in EGFR-tagRFP-expressing cells. The merged image of the

fluorescent image of EGFR-tagRFP (red) and the IRIS image (green). (**E**) IRIS imaging

of using anti-EGFR Fab probe in non-expressing cells. EGFR was localized in cluster

along the rim of some region with a size of several hundred nanometers (presumably

CCSs) (**D**, **E**, yellow arrowheads).

Fig. S3. Preparation of anti-Grb2 Fab probe. (**A**, **B**) Immunostaining with anti-Grb2
antibodies purified by affinity chromatography. The merged image of anti-Grb2
antibodies (green) and mCherry (mCh)-Grb2 (red) (**A**) or EGFR-tagRFP (red) (**B**). HeLa
cells were stimulated with 100 ng/ml EGF for 20 min at room temperature. The cells were
stained with the antibodies to find the eluted fraction capable of staining endogenous Grb2.

Endogenous Grb2 was localized in spots in non-expressing cell (**A**, arrow). The spots of
Grb2 were colocalized with EGFR-tagRFP (**B**, arrow). As a supplement, this Grb2
antibody could not stain Grb2 with mCh fused to its C-terminus. (**C–E**) IRIS imaging
using anti-Grb2 Fab probe. The merged image of the fluorescent image of mCh-Grb2
(red) and the IRIS image (green). Grb2 were localized along the rim of some region with
a size of several hundred nanometers (presumably CCSs) in mCh-Grb2-expressing cell
(**D**) and in non-expressing cell (**E**) shown in **C** (yellow arrowheads).
772

Fig. S4. Preparation of anti-transferrin receptor (TfR) Fab probe. (**A**, **B**)
Immunostaining with anti-TfR antibodies purified by affinity chromatography. The
merged image of anti-TfR antibodies (green) and TfR-pHuji (red) (**A**) or TagRFP-CLC
(red) (**B**). HeLa cells were stained with the antibodies to find the eluted fraction capable
of staining endogenous TfR. Endogenous TfR was distributed in spots in non-expressing

cell similar to expressed TfR (**A**, arrow). The spots of TfR were colocalized with TagRFP-

CLC (**B**, arrow). (**C**) IRIS imaging using anti-TfR Fab probe. The merged image of the

fluorescent image of TagRFP-CLC (red) and the IRIS image (green). The high intensity

spots of TfR were localized in CCSs shown by TagRFP-CLC (yellow arrowheads).

Fig. S5. Preparation of anti-adaptor protein 2 complex beta subunit (AP2β2) Fab

probe. (**A**) Immunostaining with anti-AP2β2 antibodies purified by affinity

chromatography. The merged image of AP2μ2-mCh (red) and anti-AP2β2 antibodies

(green). AP2β2 forms a heterotetrameric complex of α, β2, μ2 and σ2 subunits and binds

to clathrin heavy chain. HeLa cells expressing AP2μ2-mCh were stained with the

antibodies to find the eluted fraction capable of staining endogenous AP2β2. (**B**) IRIS

imaging using anti-AP2β2 Fab probe. The merged image of the fluorescent image of

AP2µ2-mCh (red) and the IRIS image (green). Endogenous AP2β2 was localized like

CCSs in the non-expressing cell (green arrow) and colocalized with the accumulation of

AP2µ2-mCh (yellow arrow).

Fig. S6. Preparation of anti-FCHo1 Fab probe. (**A–C**) Immunostaining with anti-
FCHo1 antibodies purified by affinity chromatography. The merged image of anti-
FCHo1 antibodies (green) and mCh-FCHo1 (red) (**A**), TagRFP-CLC (red) (**B**) or mCh-
FCHo2 (red) (**C**). HeLa cells were stained with the antibodies to find the eluted fraction
capable of staining endogenous FCHo1. Endogenous FCHo1 was distributed in spots in
non-expressing cell similar to expressed FCHo1 (**A**, arrow). The spots of FCHo1 were
colocalized with TagRFP-CLC (**B**, arrow). In addition, this anti-FCHo1 antibody stained
mCh-FCHo2 (**C**, arrow), which indicates that this antibody can stain both FCHo1 and 2
(FCHo1/2). (**D**) IRIS imaging using anti-FCHo1 Fab probe. The merged image of the
fluorescent image of mCh-FCHo2 (red) and the IRIS image (green). FCHo1/2 was
localized along the rim of some region (presumably CCSs) with a size of several hundred
nanometers in mCh-FCHo2-expressing cell (right upper image) and in non-expressing
cell (right lower image) (cyan arrowheads). In addition, the multi-compartmental
localizations were also observed (yellow arrowheads), which appears to be the rims of
several clathrin plaques in close proximity observed by electron microscopy (*41*, *42*).
811

Fig. S7. Preparation of anti-Eps15 Fab probe. (**A**) Immunostaining with anti-Eps15 antibodies purified by affinity chromatography. The merged image of Eps15-mCh (red) and anti-Eps15 antibodies (green). HeLa cells were stained with the antibodies to find the eluted fraction capable of staining endogenous Eps15. Endogenous Eps15 was distributed in spots in non-expressing cell similar to expressed Eps15 (arrow). (**B**, **C**) IRIS imaging using anti-Eps15 Fab probe. The merged image of the fluorescent image of Eps15-mCh (red) and the IRIS image (green). The expressed (**B**) and endogenous (**C**) Eps15 were visualized. These images show that Eps15 were not only localized along the rim of some region (presumably CCSs) with a size of several hundred nanometers (cyan arrowheads), but also as multi-compartmental localizations (yellow arrowheads).

Fig. S8. Preparation of anti-intersectin-1 (ITSN1) Fab probe. (**A**, **B**) Immunostaining with anti-ITSN1 antibodies purified by affinity chromatography. The merged image of anti-ITSN1 antibodies (red) and EGFP-ITSN1 (green) (**A**). The merged image of anti-ITSN1 antibodies (green) and TagRFP-CLC (red) (**B**). HeLa cells were stained with the

antibodies to find the eluted fraction capable of staining endogenous ITSN1. Endogenous
ITSN1 was distributed in spots in non-expressing cell similar to expressed ITSN1 (**A**,
arrow). The spots of ITSN1 were colocalized with TagRFP-CLC (**B**, arrow). (**C**, **D**) IRIS
imaging using anti-ITSN1 Fab probe. The expressed (**C**) and endogenous (**D**) ITSN1
were visualized. These images showed that ITSN1 were not only localized along the rim
of some region (presumably CCSs) with a size of several hundred nanometers (cyan
arrowheads), but also as multi-compartmental localizations (yellow arrowheads).
836

Fig. S9. Localization accuracy of Fab probes. (A) Scatter plot of standard errors (SEs)

of localizations of a probe bound over multiple frames. The localizations of a probe bound

over two or more frames were extracted, and the mean localizations and SE were

calculated for each probe (Methods). The SE indicates the accuracy of mean localizations

and estimates the resolution of the Gaussian rendering image. The SE decreased with the

number of the localizations and reached around 1 nm. (**B**) Histogram of the SE and

standard deviations (SD) of the localizations. The SD was calculated from the localization

distribution of a probe bound over multiple frames. The SD indicates the spread of the

localizations of the same probe and estimates the resolution of the IRIS image by

integrating the localizations. The localization data in **A**, **B** is from anti-Grb2 probe in **Fig.**

1A. (**C**) The mode of the SEs and SDs from the eight probes. On average, the mode of

the SE, 10.7 nm, was smaller than that of the SD, 24.2 nm, indicating 2.26-fold

improvement in the resolution. (**D**) Mean number of observed frames of a probe. The

localization data in **C**, **E** is from the probes in **Fig. 1A**.

853
Fig. S10. Improvement of the resolution. (**A**) Merged IRIS image of EGFR (red), Grb2
(green), TfR (cyan) and AP2β2 (blue) in the boxed area shown in **Fig. 1A**. The IRIS
image was reconstructed by integrating localizations at 20 nm sized pixel. (**B**) The IRIS
image was reconstructed by integrating localizations at 5 nm sized pixel in the boxed area
in **A**. (**C**) The IRIS image was reconstructed by Gaussian rendering with each localization
and SE at 5 nm sized pixel in the boxed area in **A**. Each localization and SE were
calculated by Gaussian fitting of a fluorescent speckle. (**D**) The IRIS image was
reconstructed by Gaussian rendering with mean and SE of localizations at 5 nm sized
pixel in the boxed area in **A**. The localizations of a probe bound over two or more frames
were extracted, and their mean localizations and SE were calculated (Methods). (**E–G**)
Cross-sectional profiles of EGFR and Grb2 in the boxed area in **B** (**E**), in **C** (**F**) and in **D**
(**G**). The IRIS image by Gaussian rendering using mean localizations reveals that the
location of EGFR and Grb2 clusters were shifted at the CCS rim (**G**, arrow).

Fig. S11. Labeling densities of Fab probes. (**A–C**) Histogram of the raw intensities per

10 nm pixel in the IRIS image reconstructed by integrating localizations (**A**) and

overcount-corrected localizations (**B**) and by Gaussian rendering with mean localizations

(**C**). The number of localizations is overcounted more than the number of probe binding
events due to probes bound over multiple frames. Therefore, the probes bound over
multiple frames were extracted, and their mean localization was calculated as one
localization (Methods). The overcounted localizations was corrected by combining the
mean localization data with the localization data of once-bound probe. Therefore, in IRIS
image reconstructed by integrating the overcount-corrected localizations, the intensity per
pixel represents the true density of the bound probes. To measure the intensities per 10
nm pixel in their target-localized regions, a binarized image was created by thresholding
an intensity in the IRIS image reconstructed by integrating localizations at 10 nm pixel
size. Using the binarized image, the regions with low target intensity were masked in the
IRIS image. The intensity per 10 nm pixel in the masked IRIS image was plotted on a
histogram. (**D**) Mean intensity per 10 nm pixel of each target in the masked IRIS image.
The mean density of the overcount-corrected localizations (i.e. the true density of bound
probes) was 3.9 labels per 10 nm pixel, which is 4.4 times the maximum density at which
12 nm sized antibodies can physically bind. In the Gaussian rendering IRIS image, only
probes bound over two frames were used, so the intensity per 10 nm pixel was about one-
quarter of that in the overcount-corrected IRIS image. (**E**) The masked IRIS images
reconstructed by Gaussian rendering with mean localizations in 8 pseudo color. In the
target-localized regions, the intensity per 10 nm pixel was 1.5 – 6.6 or higher. These
localization data is from the probes in **Fig. 1A**.

891
Fig. S12. Localizations of EGFR and Grb2 before and after EGFR stimulation at
37°C. (**A**) Before the stimulation, EGFR was spread over the plasma membrane. Grb2

was not localized in the CCSs. (**B**) After 3 min of the stimulation, EGFR and Grb2 were

colocalized in clusters at the CCS rims (yellow arrowheads). (**C**) After 5 min of the

stimulation, EGFR clusters decreased at CCS rims (cyan arrowheads). They accumulated

in a round shape about 200 nm in size (red arrowheads), where Grb2 were partially

localized. Many Grb2 clusters remained at the CCS rim (green arrowheads). These IRIS

images were reconstructed by Gaussian rendering with mean localizations at 5 nm sized

pixel.

A

EGFP-CLC,
anti-EGFP Nb
IRIS image

EGFP-CLC

anti-EGFP Nb
IRIS image

400 nm

B

mCh-FCHo2,
anti-mCh Nb
IRIS image

mCh-FCHo2

anti-mCh Nb
IRIS image

400 nm

C

EGFR, Grb2,
mCh-SOS, AP2β2

EGFR

Grb2

mCh-SOS

200 nm

D

EGFR, Grb2,
Cbl-mCh, AP2β2

EGFR

Grb2

Cbl-mCh

200 nm

Fig. S13. Localizations of EGFR, Grb2 and SOS1 or cCbl at CCS rims. (**A**) The
fluorescence image of EGFP-CLC and IRIS image using anti-EGFP nanobody (Nb)-
tagRFP. (**B**) The fluorescence image of mCh-FCHo2 and IRIS image using anti-mCh Nb-
EGFP. These IRIS images were reconstructed by integrating localizations at 20 nm sized
pixel. (**C**, **D**) Colocalization of mCh-SOS1 (**D**) and cCbl-mCh (**D**) with EGFR and Grb2
clusters at CCS rims. HeLa cells expressing mCh-SOS1 or cCbl-mCh were stimulated
with 100 ng/ml EGF for 3 min at 37°C. These IRIS imaging were performed with anti-
mCh Nb-EGFP, anti-EGFR Fab probe, anti-Grb2 Fab probe and anti-AP2β2 Fab probe.
In these IRIS images, the colocalizations of EGFR and Grb2 with SOS1 or cCbl (white
arrowheads) and the colocalizations of Grb2 with SOS1 or cCbl (magenta arrowheads)
were observed. These IRIS images were reconstructed by Gaussian rendering with mean
localizations at 5 nm sized pixel.

Fig. S14. Correlation analysis of the colored pixels shown in Fig. 2. (**A**) The same
panels as in **Fig. 2A** were shown to present the colored pixels. (**B**) Pearson correlation
coefficients between their targets. (**C–H**) Scatter plots of z-score intensities of the targets
at the pixels in the IRIS images shown in **Fig. 2A**. The z-score intensities were plotted
between EGFR and Grb2 (**C**), Eps15 (**D**) or AP2β2 (**E**), between Grb2 and Eps15 (**F**) or
AP2β2 (**G**), and between Eps15 and AP2β2 (**H**). The scatter plots of **C** and **F** are the same
graphs shown in **Fig. 2F**, **G**.

A EGFR, Grb2, Eps15

B SR cluster image

300 nm

923
Fig. S15. Merged IRIS image and SR cluster image of CCSs around the CCS shown
in Fig. 3A. (**A**, **B**) Merged IRIS image of EGFR (red), Grb2 (green) and Eps15 (cyan)
(**A**) and their SR cluster image (**B**). The colors of the pixels in the SR cluster image are
the same as the colors used in **Fig. 3A**. The CCS in the boxed area is shown in **Fig. 3A**.
The IRIS image was reconstructed by Gaussian rendering with mean localizations at 5
nm sized pixel.
930

Fig. S16. Merged IRIS image and SR cluster image of CCSs around the CCS shown

in Fig. 3B. (**A**, **B**) Merged IRIS image of EGFR (red), Grb2 (green) and FCHo1/2 (cyan)

(**A**) and their SR cluster image (**B**). The colors of the pixels in the SR cluster image are

the same as the colors used in **Fig. 3B**. The CCS in the boxed area is shown in **Fig. 3B**.

The IRIS image was reconstructed by Gaussian rendering with mean localizations at 5

nm sized pixel.

Fig. S17. Merged IRIS image and SR cluster image of CCSs around the CCS shown
in Fig. 3C. (**A**, **B**) Merged IRIS image of EGFR (red), Grb2 (green) and ITSN1 (cyan)
(**A**) and their SR cluster image (**B**). The colors of the pixels in the SR cluster image are
the same as the colors used in **Fig. 3C**. The CCS in the boxed area is shown in **Fig. 3C**.
The IRIS image was reconstructed by Gaussian rendering with mean localizations at 5
nm sized pixel.

Fig. S18. Correlation analysis of the colored pixels shown in Fig. 3A and fig. S15B.
(**A**) Scatter plots of z-score intensities of the targets at the pixels in the merged IRIS image
of EGFR, Grb2 and Eps15. The mean z-score intensities of the targets in the colored
pixels and the correlation coefficients between the targets are shown in **Fig. 3A**. The z-
score intensities were plotted between Grb2 and EGFR (left) or Eps15 (center), and
between EGFR and Eps15 (right). (**B**) Scatter plots for comparing EGFR-dominant
(orange), Grb2-dominant (green) and Eps15-dominant (blue) pixels. (**C**) Scatter plots for
comparing Grb2-EGFR dominant (yellow), Grb2-Eps15 dominant (magenta) and EGFR-
Eps15 dominant (cyan) pixels. The scatter plots are the same graphs shown in **Fig. 3D–
F**.

Fig. S19. Correlation analysis of the colored pixels shown in Fig. 3B and fig. S16B.
(**A**) Scatter plots of z-score intensities of the targets at the pixels in the merged IRIS image
of EGFR, Grb2 and FCHo1/2. The mean z-score intensities of the targets in the colored
pixels and the correlation coefficients between the targets are shown in **Fig. 3B**. The z-
score intensities were plotted between Grb2 and EGFR (left) or FCHo1/2 (center), and
between EGFR and FCHo1/2 (right). (**B**) Scatter plots for comparing EGFR-dominant
(orange), Grb2-dominant (green) and FCHo1/2-dominant (blue) pixels (**C**) Scatter plots
for comparing Grb2-EGFR dominant (yellow), Grb2-FCHo1/2 dominant (magenta) and
EGFR-FCHo1/2 dominant (cyan) pixels.

Fig. S20. Correlation analysis of the colored pixels shown in Fig. 3C and fig. S17B.
(**A**) Scatter plots of z-score intensities of the targets at the pixels in the merged IRIS image
of EGFR, Grb2 and ITSN1. The mean z-score intensities of the targets in the colored
pixels and the correlation coefficients between the targets are shown in **Fig. 3C**. The z-
score intensities were plotted between Grb2 and EGFR (left) or ITSN1 (center), and
between EGFR and ITSN1 (right). (**B**) Scatter plots for comparing EGFR-dominant
(orange), Grb2-dominant (green) and ITSN1-dominant (blue) pixels (**C**) Scatter plots for
comparing Grb2-EGFR dominant (yellow), Grb2-ITSN1 dominant (magenta) and EGFR-
ITSN1 dominant (cyan) pixels.

Fig. S21. Merged IRIS image of FCHo1/2 (red), Eps15 (green), ITSN1 (cyan) and

AP2β2 (blue) (A) and the SR cluster image (B). The colors of the pixels in the SR
cluster image are the same as the colors used in **Fig. 4B**. The CCS in the boxed area is
shown in **Fig. 4**. The IRIS image was reconstructed by Gaussian rendering with mean
localizations at 5 nm sized pixel.

Fig. S22. Colocalization analysis of FCHo1/2, Eps15 and ITSN1 in HeLa cells
overexpressing mCh-FCHo2. (A) Merged IRIS images of FCHo1/2 (red), Eps15 (green),

ITSN1 (cyan) and AP2β2 (blue) and the SR cluster image. The IRIS images were
reconstructed by Gaussian rendering with mean localizations at 5 nm sized pixel. (**B**)
PCA cluster image of FCHo1, Eps15 and ITSN1. (**C**) The mean z-score intensities of the
targets in the colored pixels (left) and the correlation coefficients between the targets
(right). In the cells overexpressing mCh-FCHo2, the localization of FCHo1/2 along the
CCS rim was enhanced.
996

Fig. S23. Correlation analysis of the colored pixels shown in Fig. 4 and fig. S21B. (**A**)
Scatter plots of z-score intensities of the targets at the pixels in the merged IRIS image of
FCHo1/2, Eps15 and ITSN1. The z-score intensities were plotted between Eps15 and
FCHo1/2 (left) or ITSN1 (center), between FCHo1/2 and ITSN1 (right). (**B**) Scatter plots
for comparing FCHo1/2-dominant (red), Eps15-dominant (green) and ITSN1-dominant

(blue) pixels (**C**) Scatter plots for comparing FCHo1/2-Eps15 dominant (yellow),

FCHo1/2-ITSN1 dominant (orange), FCHo1/2-Eps15-ITSN1 dominant (pale pink) and

Eps15-ITSN1 dominant (purple) pixels.

1006

Fig. S24. Colocalization analysis of EGFR, Grb2, TfR and AP2β2. (**A**) Merged IRIS

image of EGFR (red), Grb2 (green), TfR (cyan) and AP2β2 (dark blue) was reconstructed

by Gaussian rendering with mean localizations at 5 nm sized pixel. (**B**, **C**) The SR cluster

image (**B**) and the PCA cluster image with proximity (**C**) of EGFR, Grb2, TfR and AP2β2.

(**D**) The mean z-score intensities of the targets in the colored pixels (left) and the

correlation coefficients between the targets (right). The IRIS images including 112 CCSs

were subjected to PCA and cluster analysis with the same eigenvectors and clustering.

The cells were stimulated with 100 ng/ml EGF for 20 min at room temperature or for 3

min at 37°C in three independent experiments. (**E**) Scatter plots of z-score intensities of

the targets at the pixels in the IRIS images of the 112 CCSs. The z-score intensities were

plotted between EGFR and Grb2 (left) or TfR (center), and between Grb2 and TfR (right).

1019

1020

 Table S1. List of antibodies purified from antisera.

Antigens with amino acid numbers	Affinity purification pH-eluted fraction	Antibody-staining		Super-resolution imaging by Fab probe		
		expressed antigen	endogenous antigen	expressed antigen	endogenous antigen	low non-specific signals
CLC	pH3.5	P	N			
	pH2.0	P	P	P	P	P
AP2β2 1-591	pH3.5		N			
	pH2.0		P		P	N
AP2β2 616-951	pH3.5		P			
	pH2.0		P		P	P
EGFR 985-1210	pH3.5	P	P			
	pH2.0	P	P	P	P	P
Grb2	pH3.5	P	N			
	pH2.0	P	N			
Grb2 109-159	pH3.5	P	N			
	pH2.0	P	N			
Grb2 160-217	pH3.5	P	P			
	pH2.0	P	P	P	P	P

TfR 1-60	pH3.5	P	N			
	pH2.0	P	N			
TfR 93-179	pH3.5	P	P			
	pH2.0	P	P	P	P	P
Eps15	pH3.0	P	P	P	P	P
	pH2.0	P	N			
ITSN1	pH3.5	P	P			
	pH2.0	P	P	P	P	P
FCHo1	pH3.5	P	N			
	pH2.0	P	P	P	P	P
FCHo2	pH3.5	P	N			
	pH2.0	P	P	P	P	N
Epsin2	pH3.5	P	N			
	pH2.0	P	P	P	P	N
Hip1r	pH3.5	P	P			
	pH2.0	P	P	P	P	N
Dynamin2	pH3.5	P	N			
	pH2.0	P	N			

cCbl	pH3.5	P	N	/	/	/
	pH2.0	P	P	P	P	N

1022 P: Positive, N: Negative, Shaded row: Fab probes used in this study

1023

1024 **Table S2. List of antigens purified from a 6L E. coli culture in one or two experiments, for**
1025 **FCHo2 in three or four experiments.**

Antigen	Species	Amino acid numbers	Molecular weight	Concentration	Volume	Total amount
CLC	Human	Full length	25 kD	2.3 mg/ml	4 ml	9.2 mg
AP2β2	Human	1-591	66 kD	1.2 mg/ml	2.1 ml	2.5 mg
		616-951	36 kD	1.5 mg/ml	2.2 ml	3.3 mg
EGFR	Human	985-1210	25 kD	0.9 mg/ml	2 ml	1.8 mg
Grb2	Human	Full length	25 kD	2.6 mg/ml	2.6 ml	6.7 mg
GST-tagged Grb2	Human	109-159	32 kD	2.7 mg/ml	1.9 ml	5.1 mg
		160-217	32 kD	3.7 mg/ml	5 ml	18.5 mg
GST-tagged TfR	Human	1-60	32 kD	8.9 mg/ml	1.3 ml	11.5 mg
		93-179	36 kD	4 mg/ml	2 ml	8 mg
Eps15	Human	Full length	138 kD	1 mg/ml	4 ml	4 mg
ITSN1	Human	1-1220 (Short isoform full length)	137 kD	1.1 mg/ml	2.4 ml	2.6 mg
FCHo1	Human	Full length	97 kD	3.2 mg/ml	3 ml	9.6 mg

FCHo2	Mouse	Full length	85 kD	0.8 mg/ml	2.8 ml	2.2 mg
Epsin2	Human	1-456	48 kD	2.6 mg/ml	4 ml	10.4 mg
Hip1r	Mouse	Full length	138 kD	1 mg/ml	3 ml	3 mg
Dynamin2	Human	Full length	98 kD	0.86 mg/ml	2 ml	1.7 mg
cCbl	Human	Full length	120 kD	1.6 mg/ml	2.2 ml	3.5 mg

1026

1027

1028

1029

1030