Nitrogen Availability and Utilisation of Oligopeptides by Yeast in Industrial Scotch Grain Whisky Fermentation

Hidde Yaël Berg[1,2#], Georg Arju[3], Ildar Nisamedtinov[1,4]

[1] Department of Chemistry and Biotechnology, School of Science, Tallinn University of Technology, Tallinn, Estonia
[2] AS TFTAK, Tallinn, Estonia
[3] Institute of Chemistry, University of Tartu, Tartu, Estonia
[4] Lallemand Inc., Montreal, Canada

[#] **Correspondence:**
Hidde Yaël Berg
hiberg@taltech.ee

Abstract

Scotch grain whisky is produced with a substantial proportion of unmalted grains, which can result in nitrogen deficiency for yeast in the fermentable grain mash. This study examined nitrogen source availability and utilisation by three commercial whisky strains (*Saccharomyces cerevisiae*) during Scotch grain whisky fermentation, focusing on oligopeptides. Peptide uptake kinetics in synthetic whisky mash with defined peptides showed that oligopeptides of up to nine amino acids were taken up by the strains, albeit with some variability between the strains. The study found that peptides with appropriate molecular weights could replace free amino acids without negatively affecting fermentation kinetics. Moreover, fermentation performance improved when additional nitrogen was provided via peptides rather than diammonium phosphate. Analysis of industrial grain mash indicated that despite low initial yeast assimilable nitrogen, residual proteolytic activity from malt increased nitrogen availability during fermentation. Approximately 30% of the nitrogen consumed by yeast during grain mash fermentation was derived from peptides. LC-HRMS peptide analysis revealed complex dynamics of peptide formation, degradation, and utilisation. This study highlights the importance of oligopeptides in ensuring optimal fermentation efficiency in Scotch grain mash and similar substrates.

Keywords: Scotch grain whisky, *Saccharomyces cerevisiae*, Yeast Assimilable Nitrogen (YAN), Oligopeptides

Introduction

A readily available supply of easily assimilable nitrogen compounds, including amino acids and low-molecular-weight oligopeptides, is required for *Saccharomyces cerevisiae* to attain optimal fermentation efficiency during whisky fermentation (7,11,39). Assimilated amino acids and peptides can play both anabolic and catabolic roles in yeast, for example, through the biosynthesis of structural and functional proteins and nucleic acids (20,25) or the formation of higher alcohols and their associated esters that impart flavour properties of the final product (15,18). A prevalent belief is that most of the assimilable nitrogen for yeast in wort is produced during the malting process (1,9). Unlike the production process of malt whisky, which the whole mash bill is composed of malted grains, Scotch grain whisky worts are primarily derived from unmalted grains (38). Consequently, it is crucial to consider a potential limitation of nitrogenous compounds in this process.

Scotch grain whisky is usually produced from mash containing approximately 90% (w/w) unmalted grains, most commonly wheat (1,38). The unmalted grains are mechanically milled to release starch from the grain's protein matrix and then mixed with process liquor, which is a combination of process water, backset, distillation condensate, and weak worts (9). The resulting grain slurry is subsequently

cooked to gelatinise the starch. The cooking temperature varies based on the specific gelatinisation
temperature of the starch. Nevertheless, most Scotch grain distilleries typically use much higher
temperatures (95-145°C) for starch gelatinisation (9). The cooked grain slurry is then combined with
milled malt slurry in a conversion tank and maintained at mashing temperatures (63-65°C) for up to 30
min (9). The primary objective of this process, termed conversion, is to transform gelatinised starch into
fermentable sugars (mostly maltose, glucose, and maltotriose) (38). Traditionally, wort has been
separated from the grains to provide a clear fermentable liquid. However, the wort separation step has
been largely eliminated in modern grain whisky distilleries; instead, the entire mash is transferred to the
fermenter after cooling and pitching with yeast (9).

As mentioned above, it is widely considered that a significant proportion of the yeast-assimilable
nitrogen present in the final wort is derived from the malt. During the malting process, proteolytic
enzymes are released into the endosperm to degrade grain storage proteins (i.e. hordeins, glutelins,
albumins, and globulins) into peptides and free amino acids (21). Many of these enzymes, including
aminopeptidases and some endoproteinases, are sensitive to heat and are largely inactivated at
temperatures above 55°C, used during the conversion step (8,23). However, studies have described the
formation of soluble nitrogenous compounds during conversion by heat-stable endoproteinases and
carboxypeptidases (23,29). These proteases are inactivated only at temperatures above 70°C (23).
Unlike brewing, the malt stream in the Scotch grain whisky production process does not reach this
temperature, nor does it contain a separation and boiling step (9,31,38). Consequently, in addition to the
conversion step, the proteolytic activity can also be present during fermentation. In the context of Scotch
grain whisky production, where the malted substrate constitutes on average 10% of the mash bill, this
may be crucial for supplying sufficient nitrogen for yeast to ensure good fermentation kinetics,
especially as the addition of exogenous proteolytic enzymes or nitrogen supplements is prohibited (38).

The utilisation of oligopeptides as a yeast nitrogen source has received limited attention in the field of
Scotch grain whisky production. Our previous study on wine yeast showed that oligopeptides can
supply yeast with sufficient nitrogen to complete fermentation without the need for additional amino
acids (6). Thus, oligopeptides may also be an important secondary source of nitrogen for yeast during
Scotch grain whisky fermentation.

In this study, we aim to investigate the availability, production, and utilisation of different yeast-
assimilable nitrogen sources during Scotch grain whisky fermentation. Three commercial strains of
Saccharomyces cerevisiae, commonly used in whisky production, were selected for analysis. The focus
was on the role of oligopeptides as a secondary source of nitrogen for yeast. Our findings provide
insights into nitrogen availability in Scotch grain whisky fermentation and highlight the importance of
oligopeptides for ensuring optimal fermentation efficiency.

Materials and Methods

Strains
Three commercial *S. cerevisiae* whisky yeast strains (DistilaMax® GW, NT, and XP) were provided by
Lallemand Inc. (Montreal, QC, Canada). The strains were stored as cryocultures in 25% glycerol at -80
˚C. The relevant strain information is presented in Table 1.

Each yeast inoculum was prepared by streaking the cryoculture on a yeast peptone dextrose (YPD) agar
plate (1% yeast extract, 2% bacteriological peptone, and 2% glucose) and pitching a single colony into
a shaker flask containing YPD broth. The flasks were incubated overnight at 30°C, with shaking at 150
rpm. Cells were washed twice with equal volumes of sterile 0.9% NaCl prior to inoculation into
fermenters to deliver 5×10^6 cells/mL, based on OD600 measurements.

Fermentations

Synthetic whisky mash fermentations

Fermentation experiments were performed on synthetic whisky mash (SWM) to confirm the importance of peptides as a nitrogen source to whisky yeast. In addition, the potential benefits of nitrogen supplementation, either through organic or inorganic means, were investigated.

Preparation of the bovine serum albumin hydrolysate

A bovine serum albumin (BSA) enzymatic hydrolysate was used as a source of peptides in a synthetic whisky mash (SWM). The hydrolysate was prepared according to a previously reported method (4,6).

The hydrolysate was produced from BSA (40 g/L) in a 1 L benchtop fermenter (Applikon, Delft, The Netherlands) using the industrial protease COROLASE®7089 (AB Enzymes, Darmstadt, Germany). The protease dose rate was 0.5% w/w BSA, and hydrolysis was performed at pH 7 (maintained by titration using 2 M NaOH) for 20 h at 50 °C. After completion of BSA hydrolysis, the hydrolysate was filtered with a 10 kDa Vivaflow® 200 10,000 MWCO Hydrosart crossflow cassette (Sartorius, Göttingen, Germany) to remove the protease and larger peptides. To further concentrate the smaller MW peptide fraction ($\leq$ 2 kDa), another filtration step was performed using a Vivaflow® 200 2,000 MWCO Hydrosart crossflow cassette (Sartorius, Göttingen, Germany). The hydrolysate was then freeze-dried and stored at −20 °C until further use.

Synthetic whisky mash (SWM) composition

The composition of SWM is an adaptation of the synthetic grape must described by Salmon and Barre (32). Citric and malic acids were excluded from the medium, and the sugar concentration, free amino acid ratios (Table S1), and pH were adjusted to mimic whisky mash. These values were based on measurements taken from a commercial whisky production process.

The base SWM contained 150 g/L maltose and glucose (9:1 w/w) and was modified by adding different nitrogen sources (Table 2). Thus, in the first experiment (AP230), 130 mg/L of nitrogen was added with amino acids, and 100 mg/L was added in the form of small molecular weight peptides (smaller than 2 kDa) from the BSA hydrolysate. In the second experiment, all nitrogen (200 mg/L) was supplied in the form of small-MW peptides from the BSA hydrolysate (P230). To study the effect of additional nitrogen supplementation with (in)organic sources, 70 mg/L of nitrogen was added to the P230 medium from either the BSA hydrolysate (PP300) or diammonium phosphate (PD300). The pH of all the media was adjusted to 5.2. The media were then filter-sterilised using a 0.22 μm Steritop® vacuum-driven disposable filtration system (Merck-Millipore, Burlington, MA, USA) prior to yeast inoculation.

Fermentations

All fermentations on synthetic media (100 mL) were performed in triplicate at 24 °C in 100 mL Pyrex™ bottles equipped with a GL45 open-top PBT screw cap and PYREX™ Media Bottle Septum (Corning Inc., Corning, NY, USA). A gas outlet was installed to prevent overpressure by piercing the septum with a Sterican® Ø 0.8 x 40 mm single-use hypodermic needle (B. Braun, Melsungen, Germany) attached to a Millex-FG 0.2 μm hydrophobic PTFE filter (Merck KGaA, Darmstadt, Germany).

Samples (2 mL) were collected every 6 h for the first 48 h, every 24 h until 120 h, and every 48 h until 192 h. Monitoring of CO_2 production was continued until the endpoint (288 h), after which a final sample was collected. The biomass density of the samples was assessed by measuring the optical density at 600 nm using an Ultrospec® 10 Cell Density Meter (Biochrom, Ltd., Cambridge, UK). The specific production rate of CO_2, monitored gravimetrically, was used as the main indicator of the fermentation progress. After centrifugation at 9600 × g for 10 min at 4 °C in a MicroCL 21R Microcentrifuge (Thermo Fisher Scientific, MA, USA), the supernatant was filtered through Minisart RC 0.2 μm syringe filters (Sartorius, Göttingen, Germany) and stored at -20 °C until further analysis.

Grain mash fermentations

Unfiltered grain mash, produced from approximately 10% malted barley and 90% unmalted wheat, was obtained from a commercial Scotch grain whisky distillery. Fermentation was performed in 500 mL of grain mash in 1 L Biobundle benchtop fermenters (Applikon, Delft, the Netherlands) using the cultivation control software BIOEXPERT V2 v2.96 (Applikon, Delft, the Netherlands). The fermenters were equipped with pH and temperature sensors. The production of CO_2 was measured with an HPR-20 R&D Benchtop Gas Analysis System (Hiden Analytical Ltd., Warrington, UK) using MASsoft 10 Professional software (Hiden Analytical Ltd., Warrington, UK). To allow for stable measurement of CO_2 production, the headspace was flushed at a rate of 100 ml/min (5% CO_2, 95% N_2).

The fermentation temperature was raised from 25 °C to 34 °C in the first 24 h to mimic the heat production that occurs during the factory process due to yeast activity, and then maintained at 34 °C for 60 h. All fermentations were performed in triplicate. Samples (10 mL) were collected every 12 h from the bottom of the fermenter using a sample port and a 10 ml sterile syringe (B. Braun, Melsungen, Germany).

The formation of free amino acids and peptides due to residual proteolytic activity in the grain mash was analysed in an uninoculated grain mash by simulating the changes in temperature, pH, and ethanol concentration during fermentation. The change gradients of the latter parameters were based on the actual measurements of grain mash fermentation with DistilaMax® NT, and are shown in

Table **3**. The uninoculated mash contained 200 μg/mL G418 (InvivoGen, Toulouse, France) and 2 ppm Lactoside 247™ (Lallemand, Montreal, QC, Canada) to prevent microbial growth.

Analysis

Preparation of grain mash samples for amino acids and peptide analysis

Grain mash samples (10 mL) were centrifuged at 13000 × g for 15 min at 20°C in a Hettich® ROTANTA 460/460R centrifuge (Andreas Hettich GmbH & Co. KG, Tuttlingen, Germany) to separate the solids. The liquid phase was then filtered using Minisart RC 0.2μm syringe filters (Sartorius, Göttingen, Germany) to remove any residual biomass.

Sugars were removed from the grain mash by fermentation with a peptide transporter knockout (PepKO) strain (5,6). The latter was necessary to avoid the loss of amino acids due to Maillard reactions when total amino acid analysis was carried out under high temperature and acidic conditions. First, the samples were heated at 75 °C for 20 min to remove any residual proteolytic activity. In addition, the uninoculated grain mash samples were freeze-dried in a Heto PowerDry PL3000 Freeze Dryer (Thermo Fisher Scientific, MA, USA) and reconstituted in MilliQ water to remove ethanol, which would otherwise prevent fermentation. To ensure proper fermentation with the PepKO stain, 100 μL of a nutrient stock (Table S2) containing minerals, vitamins, and 100 mg/L nitrogen from NH_4^+ was added to the samples. Because the PepKO strain used was a descendant of a wine stain, the nutrient stock was also supplied with 0.1% glucoamylase (MEGA PACIFIC TECHNOLOGY INC, Arcadia CA, USA) to break down maltose and maltotriose into more easily assimilable glucose (3). In addition, because the PepKO strain contained KanMX as a marker for the deletion of *FOT* peptide transporter genes (5), 200 μg/mL G418 (InvivoGen, Toulouse, France) was added to prevent microbial contamination. Fermentation with the PepKO strain was performed in 15 mL conical tubes (Eppendorf AG, Hamburg, Germany) at 32 ºC and at 150 rpm. The lids of the tubes were slightly opened to prevent overpressure. The inoculum (20 μL) was prepared as previously described. The end of the fermentation process was indicated by the cells dropping from the culture medium. As the samples contained variable concentrations of residual sugars, the fermentation time was sample dependent. Cells were separated from the samples by centrifugation at 9600 × g for 10 min at 4 °C in a MicroCL 21R Microcentrifuge (Thermo Fisher Scientific, MA, USA) and filtered using Minisart RC 0.2 μm syringe filters (Sartorius, Göttingen, Germany).

180 The resulting samples were filtered to obtain a ≤ 2 kDa fraction using Vivaspin 2 centrifugal
181 concentrators (Sartorius AG, Göttingen, Germany) according to the manufacturer's instructions. The
182 retentate was washed twice with 2 mL of MilliQ water. The permeate was freeze-dried and reconstituted
183 in Milli-Q water to obtain the final sample.

Analysis of free and bound amino acids

Concentrations of free and peptide-bound amino acids were analysed on a Waters ACQUITY UPLC®
system (Waters Corporation, Milford, MA, USA) coupled to Waters ACQUITY UPLC tuneable
ultraviolet (TUV) detector (Waters Corporation, Milford, MA, USA) after derivatisation using Waters
AccQ-Tag chemistry, as described by Fiechter and Mayer (17). For total amino acids, the samples were
first hydrolysed with 6M HCl + 1% (v/v) phenol at 105 °C for 22 h in vacuum using an Eldex H/D
workstation (Eldex Laboratories Inc., Napa, CA, USA). The peptide-bound amino acids were calculated
by subtracting the measured free amino acids from the total amino acids.

Peptide analysis

Sample preparation

The SWM samples were first mixed (1:1) with methanol in BRANDplates™ pureGrade™ 96-Well
Microplates (BRAND GMBH + CO KG, Wertheim, Germany). The plates were then centrifuged at 560
× g using a BioSan LMC-3000 centrifuge (Biosan, City, Company) to remove the precipitate. Then, 20
μL from each well was transferred to a Waters™ round well polypropylene 350 μL 96-well sample
collection plate (Waters Corporation, Milford, MA, USA) and diluted with Milli-Q water (160 μL) to a
final volume of 180 μL. All samples were spiked with 0.5 ppm caffeine (20 μL) to a final volume of
200 μL. Caffeine was used as an internal standard (reference housekeeping ion) during subsequent
analysis.

Ice-cold grain mash samples (100 μL) were first mixed with 900 μL of ice-cold methanol in 1.5 mL
Eppendorf® tubes (Eppendorf, Hamburg, Germany) and centrifuged at 11,200 × g at 4 °C for 20 min in
a MicroCL 21R Microcentrifuge (Thermo Fisher Scientific, MA, USA) to remove the precipitate. The
samples were then freeze-dried in a Heto PowerDry PL3000 Freeze Dryer (Thermo Fisher Scientific,
MA, USA) and reconstituted in 100 μL Milli-Q water. All samples were spiked with 0.5 ppm caffein to
a final volume of 110 μL.

Liquid chromatography-mass spectrometry (ultra-high-pressure liquid chromatography ion mobility separation-enabled high-resolution mass spectrometry (UHPLC-IMS-HRMS))

The peptides were analysed using the methodology described by Arju et al. (2022). Briefly, the Waters
I-Class Plus (SM-FL) UPLC system (Waters Corporation, Milford, MA, USA) was coupled with a
Waters Vion IMS-QTof Mass Spectrometer equipped with a LockSpray II Exact Mass Source Enclosure
and MKII tool-free ESI probe assembly directly connected to the column outlet. Nitrogen was used as
collision gas. The instrument was controlled using Waters UNIFI 1.9.4 (3.1.0, Waters Corporation,
Milford, MA, USA).

The instrument was operated in positive polarity, sensitivity mode (31,000 FWHM at 556.2766 m/z),
and labile ion mobility tuning mode. The analysis type was set as Peptide Map (IMS), and the
experiment type was set to MSe. Data were acquired in the HDMSe mode with a scan time of 0.165 s
for synthetic and 0.250 s for grain mash. The following manual quadrupole profile was used for SWM:
mass 150/250/450 (m/z), dwell time 60/20 (% scan time), ramp time 10/10 (% scan time); and for grain
mash: mass 162/300/500 (m/z), dwell time 35/35 (% scan time), ramp time 15/15 (% scan time).

The injection volume was 5 μL. Full loop (5 μL) injection mode was used with a 3-volume overfill.
The samples were analysed using an Acquity UPLC HSS T3 Column (1.8 μm, 1 × 150 mm, Waters
Corporation, Milford, MA, USA) maintained at 45 °C. The initial flow rate was 0.15 mL/min. The
gradient for synthetic mash was as follows: a 0–0.5 min hold at 3% B; 0.5–5.5 min linear gradient, 3%–
28% B; 5.5–7 min linear gradient, 28%–95% B; 7–7.5 min hold at 95% B accompanied by a linear flow

rate increase of 0.15–0.3 mL/min; 7.5–7.6 min linear gradient, 99%–3% B accompanied by a linear
flow rate decrease of 0.3–0.15 mL/min; and 7.6–10 min hold at 3% B. The gradient for grain mash was
as follows: a 0-1 min gold at 1% B; 1-16 min linear gradient, 1-30% B; 16-17 min linear gradient, 30-
95% B accompanied by a linear flow rate increase of 0.15-0.2 mL/min; a 17-17.5 min hold at 95% B
accompanied with a linear flow rate increase of 0.2-0.3 mL/min; 17.5-17.6 min gradient, 95-1% B
accompanied with linear flow rate decrease of 0.3-0.15 mL/min; a 17.6-20 min hold at 1% B.

Data Treatment and Statistical Analysis

Data was treated and analysed using R v4.3.2 (R Core Team 2023) and RStudio (RStudio Team 2023).
The growth rate and maximum cell population in SWM were estimated by fitting the optical density
curves with the "summarizegrowth" R-function from the GrowthCurver package (33). The significance
of ANOVA between the mean values was assessed using Tukey's HSD method.

Peptide mapping for peptide analysis was conducted according to the method described by Arju et al.
(2022) (4). The time window for peptide detection was extended to 14 min to accommodate the
extended gradient used to analyse the grain mash samples. Peptide mapping in the grain mash samples
of peptide masses was performed against five barley storage proteins and nine wheat storage proteins
(Table S3).

Data processing of amino acids and peptides from the strain characterisation in SWM was performed,
as described by Berg et al. (6). A list of the mass spectrometric data of di- to nonapeptides identified in
the BSA hydrolysate in this study is shown in Table S6.

All data were visualised using ggplot2 (40).

Data Availability

248   Data files and the R script for data treatment and analysis can be found at:
249   https://zenodo.org/records/12154681

250   All data are contained in the manuscript and the supplemental material.

Results

Effect of oligopeptides on the fermentation kinetics in synthetic whisky mash

Fermentation experiments were performed on synthetic whisky mash (SWM) to study the uptake of
peptides as a nitrogen source by the whisky yeast strains used in this study (Table 1) under different
conditions of nitrogen supply (Table 2). In the first experiment, the fermentation medium (AP230)
contained 130 mg N/L from free amino acids (FAA) and 100 mg N/L from peptides with a molecular
weight (MW) ≤ 2 kDa. In the second medium (P230) the FAA were deficient (30 mg N/L), and peptides
were the only substantial source of nitrogen (200 mg N/L). Finally, the effect of additional nitrogen
supplementation (70 mg N/L) with either diammonium phosphate (DAP) or peptides was studied
(media PP300 and PD300, respectively). The effect on the fermentation kinetics is shown in Figure 1.

In the experiments with a total N content of 230 mg N/L, containing either free amino acids (56.5% of
N) and peptides (43.5% of N) (AP230) or only peptides (P230), the three strains displayed no significant
differences in maximum cell population (Figure 1A), maximum CO_2 production rate (Vmax, Figure
1B), and fermentation time (Figure 1C). DistilaMax® GW was the only strain that exhibited a
significantly higher Vmax value in the P230 medium, resulting in a slight, yet statistically insignificant
reduction in overall fermentation time (80% attenuation, determined based on the maximum theoretical
CO_2 production from 150 g/L maltose and glucose (9:1)). These findings corroborate previous results
from the experiments we conducted with a wine yeast and suggest that peptides individually can also
provide the necessary nitrogen to these whisky strains (6).

Among the three strains, DistilaMax® XP exhibited the lowest fermentation performance (Vmax and fermentation time) at 230 mg N/L (Figure 1), which could be attributed to its higher nitrogen demand than that of the other strains. Interestingly, with additional nitrogen supplementation (70 mg N/L) with diammonium phosphate (DAP, medium PD300), the fermentation kinetics did not improve, whereas a positive impact was observed with an increased peptide concentration (medium PP300). Thus, in the PP300 medium, the maximum cell population and Vmax increased by 52% and 62%, respectively, while the fermentation time decreased by 42%, compared to the P230 medium. Similarly, DistilaMax® GW in the PP300 medium showed a 28% increase in Vmax and a 36% decrease in fermentation time, although the biomass density remained similar to that in the P230 medium. In the case of DistilaMax® NT, the maximum cell population and Vmax increased by 17% and 11%, respectively, resulting in a 23% decrease in fermentation time, albeit not statistically different from the fermentation time in the lower nitrogen-containing P230 medium. Interestingly, DAP supplementation (PD300 medium) had a negative effect on the maximum cell population and Vmax by DistilaMax® GW and NT. Nevertheless, despite variations in the maximum cell population and Vmax, the fermentation time remained unaffected.

A principal component analysis plot was constructed to support these observations (Figure 1D). The experiments in the AP230 and P230 media clustered together by strain, indicating that peptides can provide the necessary nitrogen to these whisky strains when FAA are limited, without affecting fermentation efficiency. For DistilaMax® XP, the experiments conducted on the PD300 medium cluster with the AP230 and P230 lower nitrogen-containing media, showing no or minimal effect of ammonium supplementation on this strain. The negative effect of DAP supplementation on the other strains can be observed by their respective PD300 medium clustering away from their respective AP230 and P230 clusters. The positive effect of increased nitrogen content from peptide supplementation (PP300) on DistilaMax® XP can be seen by its clustering with DistilaMax® GW in the AP230 and P230 media. DistilaMax® NT in the PP300 medium also moved closer to this cluster, whereas DistilaMax® GW in the PP300 medium formed its own group.

Oligopeptide uptake by whisky yeast strains in synthetic whisky mash

In addition to the fermentation kinetics, peptide consumption by the three whisky yeast strains was monitored during the first 72 hours of fermentation (Figures 2, S1-4). Intensive uptake of peptides with different chain lengths (up to nine amino acids) simultaneously with FAA was observed in both AP230 and P230 media for all strains (Figure S4). The conserved ability of the three strains to take up peptides from the environment was also evident from their similar peptide consumption profiles in these two media.

In the medium where additional nitrogen was added in the form of DAP (PD300), ammonia was consumed first, which resulted in delayed uptake of di- and tripeptides and limited the uptake of peptides with longer chain lengths in all strains (Figures 2, S1-4). In addition, the strains revealed distinct patterns of peptide uptake in the medium with increased peptide content (PP300), which may be attributed to their individual nitrogen requirements. For example, DistilaMax® GW showed similar consumption levels for the di- and tripeptides in the PP300 medium as in the P230 medium (Figure 2), while peptides with a longer chain length were taken up more slowly. In DistilaMax® NT, this effect was less pronounced, whereas in DistilaMax® XP, all peptides were taken up simultaneously. As the methodology utilised in this study does not enable the exact quantification of individual peptides, it was impossible to determine the exact quantity of nitrogen contributed by each peptide length group to the overall nitrogen intake.

Together with the observed effects of peptides on fermentation performance (Figure 1), these results support our hypothesis that peptides can serve as a substantial alternative source of nitrogen for yeast.

Nitrogen availability and the role of peptides in Scotch grain whisky fermentation

To study the role of peptides as a yeast nitrogen source in Scotch grain whisky fermentation, a grain mash sample from an industrial Scotch grain whisky production process was obtained, and fermentation was conducted in one-litre benchtop fermenters using the three strains previously characterised for their peptide uptake capability in SWM.

All strains completed fermentation in the grain mash, albeit with distinct fermentation kinetics (Figure 3). DistilaMax® GW showed the fastest fermentation, ceasing active fermentation by 24 hours. After this point, the CO_2 production rate significantly decreased, and fermentation was complete after approximately 48 hours. In contrast, with DistilaMax® NT and XP the CO_2 production rate started to cease at 36 hours with fermentation completion at 48 hours. The ethanol content of the fermentation with DistilaMax® NT was the highest at 8.8 ± 0.2 %(v/v), followed by DistilaMax® XP at 8.7 ± 0.2 %(v/v), and DistilaMax® GW at 8.5 ± 0.4 %(v/v), which were not significantly different from each other (α=0.05). The ethanol yield was, on average, 97% of the target of the sampled distillery, which aims at a final ethanol concentration of 9% (v/v).

The initial grain mash contained 90.2 ± 6.73 mg N/L from FAA (Figure 4). According to previously published studies, this amount does not meet the minimum requirement for satisfactory fermentation efficiency (150-250 mg N/L) for the given fermentable sugar content (19,20). However, the grain mash also contained oligopeptides, some of which could be a potential source of N for yeast. In total, 70.4 ± 9.9 mg N/L was determined to be derived from peptides with a MW ≤ 2 kDa. Thus, the sampled grain mash initially contained up to 160.6 ± 12.0 mg N/L of potentially yeast-assimilable nitrogen.

The production of grain mash for Scotch grain whisky fermentation does not involve boiling, which could result in residual proteolytic activity during fermentation originating from the malt. To determine the amount of additional potentially yeast-assimilable nitrogen liberation, an experiment was conducted using an uninoculated grain mash, where the actual grain mash fermentation conditions were simulated by gradually changing the pH and ethanol concentration (Table 3, Figure 4). The results showed an increase in nitrogen content in both FAA and, to a lesser extent, peptides ≤ 2 kDa. The increase of potentially available nitrogen content amounted to a total of 58.4 ± 6.28 mg N/L in 36 hours, with 45.8 ± 4.1 mg N/L from FAA and 12.6 ± 4.7 mg N/L from peptides. Therefore, a total of 219.0 ± 13.5 mg N/L was available for fermentation, with 38% of it derived from peptides with a MW ≤ 2 kDa. Corroborating the observed fermentation kinetics and literature data, this quantity was sufficient to complete fermentation (20).

Although the strains displayed varying fermentation and nitrogen uptake kinetics, the amounts of nitrogen consumed from the grain mash were similar (Figure 4). Thus, DistilaMax® GW consumed 186.8 ± 14.0 mg N/L, with 128.2 ± 7.9 mg N/L derived from FAA and 58.6 ± 11.5 mg N/L from peptides. Similarly, DistilaMax® XP consumed 193.6 ± 14.5 mg N/L, with 134.5 ± 7.9 mg N/L derived from FAA and 59.1 ± 12.2 mg N/L from peptides. DistilaMax® NT consumed 175.3 ± 13.7 mg N/L, with 126.0 ± 7.9 mg N/L derived from FAA and 49.4 ± 11.2 mg N/L from peptides. Thus, peptide-derived nitrogen in these fermentations comprised ~30% of the total nitrogen assimilated by yeast.

Peptide production, degradation and consumption (based on the change in their MS peak intensities) at different sampling points were compared (Figure 5). In total, 153 peptides ranging from 2 to 23 AA in length could be followed. Peptide annotations were obtained by mapping the peptide masses to the six most abundant barley storage proteins (hordeins) and nine wheat storage proteins (glutenins) (Table S3) (24,34). However, this approach is subject to a degree of identification inaccuracy due to the numerous proteins that were included in peptide screening, as well as those that were not included. Consequently, we decided to attribute only the peptide length to the observed peptide candidate masses, as suggested in another study (4).

Analysis of peptides in the uninoculated mash revealed complex dynamics of peptide production and
degradation in the grain mash due to malt proteases (Figure 5A). During the first 12 hours of incubation,
degradation of peptides occurred, as shown by the decreased relative concentration of most peptides, as
well as the increased concentration of free amino acids (Figure 4). The hydrolysis continued between
12 and 36 hours, first with increasing relative concentrations of higher MW peptides (12 to 24 hours),
which were then further degraded into smaller ones between 24 and 36 hours, suggesting an interplay
between various proteases and peptidases. The simultaneous production and degradation of peptides
also explains the limited change in the contribution of peptides to the overall increase in the potentially
yeast-assimilable nitrogen content of the grain mash, as shown in Figure 4.

The complex kinetics of peptide formation and degradation due to proteolysis by malt-derived enzymes
makes it difficult to precisely interpret the extent of their consumption by yeast (Figure 5B-D). Thus,
the changes in the relative concentration of small MW peptides (two to five AA in chain length) during
fermentation in the first 12 hours were very similar to those observed in the uninoculated mash
incubation experiment. In contrast, a higher number of peptides with a negative fold change in the
relative concentration during later stages of fermentation when free amino acids were depleted suggests
their consumption by yeast. In all three strains, the net change between the start and end of fermentation
suggested the consumption of di-to-pentapeptides (average fold change of 0.6 compared to 0.9 in the
uninoculated mash).

The only major difference among the three yeast strains was observed in the DistilaMax® XP
fermentation experiment (Figure 5D), in which the apparent increase in peptides was much more
pronounced during the first stage of fermentation (0 to 12 hours). The exact reasons for this are unknown
but may be related to the slower fermentation kinetics of this strain compared to the other two strains.
Bound amino acid analysis of the 2 kDa fraction in DistilaMax® XP did not show a substantial increase
in the nitrogen content of peptides between 0 and 12 hours (Figure 4). Therefore, it was not possible to
determine the significance of this observation without quantitative data on specific peptides at more
frequent sampling points.

Overall, these findings indicate that the production, degradation, and consumption of peptides during
Scotch grain whisky fermentation is a highly dynamic and complex process.

Discussion

Around 90% of the raw material used in the production of Scotch grain whisky consists of unmalted
grains, which may lead to nitrogen deficiency for yeast in the fermentable grain mash. This study
investigated the availability and utilisation of nitrogen sources by yeast during the fermentation of
Scotch grain whisky, with a particular focus on the role of oligopeptides. First, three commercial
Saccharomyces cerevisiae whisky strains were assessed for their ability to take up peptides from
peptide-supplemented synthetic whisky mash, using various nitrogen sources. All three strains were
able to complete fermentation with peptides as the sole nitrogen source, demonstrating that peptides
with a suitable chain length can satisfy the nitrogen requirements of yeast. We previously characterised
a *Saccharomyces cerevisiae* wine yeast (59A, haploid variant of the commercial wine strain EC1118
(28)) for its ability to take up peptides with different chain lengths in a similar analysis and found that
it could utilise peptides up to seven amino acids in length, which were the highest chain length peptides
determined by the analyses used (6). By further optimising the peptide mapping analysis, in the current
study we suggest that the three whisky strains can take up peptides of up to at least nine amino acids in
length. Like the wine strain 59A, these whisky strains took up smaller peptides first, with the only
exception being that the whisky strains preferred di- and tripeptides, whereas in 59A, tetrapeptides were
also preferred (6). This can be explained by the activity of the Fot family of peptide transporters, which
is specific to wine yeasts (5,6,14,28). Ptr2 (previously characterised as a di- and tripeptide transporter
(30)) is presumably the active di- and tripeptide transporter in whisky strains, explaining why
tetrapeptides were taken up later with the other larger peptides.

Measurements taken from the Scotch grain mash obtained from an industrial distillery indicate that when a significant proportion of the mash bill is unmalted (approximately 90% in this case), the initial YAN content of the grain mash (~160 mg N/L) is relatively low, considering that the minimum nitrogen requirement for satisfactory fermentation efficiency for the mash with the respective fermentable sugar concentration remains between 150 and 250 mg N/L (20) and that not all peptides in the ≤2 kDa molecular weight (MW) fraction might be fully assimilable by yeast. However, our study showed that the residual proteolytic activity that remained in the mash after conversion provided an additional 36% of potentially assimilable nitrogen to yeast during fermentation, of which the majority (80%) was in the form of free amino acids and 20% was contributed by peptides with MW ≤ 2 kDa. The fermentation experiments with three commercial whisky strains indicated that the final nitrogen content in the mash remained sufficient to support complete fermentation, with the strains taking up ~ 185 mg N/L on average, of which ~30% can be attributed to peptides. Thus, the remaining proteolytic activity in the mash seems essential for the fermentation process, and peptides form a considerable source of nitrogen for yeast during Scotch grain whisky fermentation.

Several protease families are involved in the germination process of barley, and are therefore present in mash (35). Heat-stable endoproteinases and carboxypeptidases may retain their activity in the mash during subsequent fermentation because the temperature of the conversion process does not exceed 70 °C (23,29,35), which was observed in our experiments. The LC-HRMS analysis of relative concentrations of individual peptides in uninoculated mash, in which the temperature, pH, and ethanol concentration were changed to simulate the fermentation process, suggested that these conditions favoured the activity of both proteolytic enzyme classes of malt, resulting in the production of peptides and FAA. However, this process is dynamic and depends on the fermentation stage. The formation of free amino acids and small peptides occurred mostly at the beginning of fermentation (0 to 24 hours, T= 24 – 34 °C, pH= 5.2 – 4.3, EtOH= 0 – 50 g/L), whereas higher MW peptides started to accumulate in the later stages (12 hours and further). Because of this highly dynamic process, involving both production and degradation of peptides, limited variation in the nitrogen concentration in the MW ≤ 2 kDa peptide fraction of the uninoculated mash was observed. The extent of peptide uptake in the fermenting mash during the first phases of fermentation (0 to 12 hours) was difficult to estimate because of the co-occurring activity of different malt proteases; however, the consumption of di- to pentapeptides was more apparent between 12 and 36 hours.

The current MS-based methodology for studying the composition and concentration of peptides in unelucidated matrices, such as Scotch grain mash, has limitations. First, not all protein sequences in barley and wheat have been fully sequenced (13). Thus, in the present study, the obtained peptide masses were mapped only to the major (most abundant) known storage proteins in barley (hordeins) and wheat (glutenins) for annotation (24,34). However, this approach is subject to a degree of identification inaccuracy due to the numerous protein sequences (total of 15) that were included for peptide mapping, as well as potentially unknown proteins that were not included but were also degraded into peptides. This limits the annotation of peptides to their size (number of AA residues in the chain length) instead of the exact AA sequence. As the MS peak intensity of peptides can be highly dependent on the amino acid composition and sequence (26), only the relative quantification of peptides by fold-change analysis of individual detected peptides was performed in this study. In combination with fractionation and subsequent AA analysis of the obtained molecular fractions, it was possible to explain these trends more quantitatively. However, some inaccuracies may have occurred in this case, for example due to retention of some smaller peptides on the cut-off filter. Ideally, the qualitative and quantitative analyses of peptides should be integrated into a single analysis. One method that currently exists makes use of the UV absorbance of peptides by coupling a photodiode array detector between LC and MS. The UV peak areas can be converted to absolute concentrations based on the law of Lambert-Beer, which requires the molar extinction coefficient of the peptide (10). However, this analysis would require extended chromatographical times to resolve most UV peaks in complex media, in addition to being hindered by the highly variable ionisation efficiencies of short peptides (26), which are required for the

quantification of different peptides that co-elute under one UV peak (37). Further research to develop qualitative and quantitative analyses of peptides by MS in complex fermentation samples is therefore required.

Although the addition of nitrogen nutrients to grain whisky fermentation is not allowed by regulation (38), the effects of additional nitrogen supplementation with either peptides or by the often industrially applied ammonium salt DAP on the fermentation performance in synthetic whisky mash were studied. In these experiments, supplementation with peptides resulted in higher biomass accumulation and CO_2 production rates, and consequently, shorter fermentation times, than with the medium with a lower nitrogen content. This is in stark contrast to the effect of ammonium supplementation on fermentation performance, which was neutral or even negative. These results seem counterintuitive considering the popularity of DAP or urea in industry; however, they are in line with what has been observed in other studies. In one study, nitrogen supplementation with a peptone induced higher biomass and ethanol production than ammonium sulphate (12). Another study revealed slower fermentation rates with ammonium compared to free amino acids, delivering equivalent amounts of nitrogen (2). Gene expression has been shown to be differentially reprogrammed based on the nitrogen source added (22): when ammonia was added, there was a higher expression of genes involved in amino acid biosynthesis, whereas the addition of amino acids resulted in a higher expression of genes related to protein biosynthesis. Thus, a possible explanation for the observed differences in the effect of nitrogen supplementation is that (peptide-derived) amino acids can be directly used in protein biosynthetic processes, while ammonium must first be incorporated into amino acids through *de novo* biosynthesis, requiring cellular resources including carbon and energy (27). Under the conditions used in this study, ammonia delayed peptide uptake, possibly through nitrogen catabolite repression (36). We hypothesised that this negative effect of ammonium supplementation on fermentation kinetics is due to the interplay between two factors: on the one hand, the cells need to *de novo* synthesise all the required amino acids, and on the other hand, the presence of ammonia hinders the uptake of peptides which could otherwise supply readily usable amino acids.

In conclusion, these results show that peptides can play a considerable role as nitrogen sources in Scotch grain whisky fermentation. The whisky yeast strains were able to take up a wide variety of peptides and use them effectively to attain complete fermentation. In industrial Scotch grain mash, the concentration of yeast assimilable nitrogen, including that produced during fermentation owing to residual proteolytic activity proved to be sufficient, with peptides being a considerable source of nitrogen.

Acknowledgements

The authors want to acknowledge Martin Johannes Talu for his assistance with the experiments, Struan Reid for his advice throughout the project, and Mairéad Walsh for her help in editing this document.

Disclosure statement

All authors have read and agreed to the published version of the manuscript and declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

This work was funded by the Estonian Research Council project RESTA13.

References

1. Agu, R. C., Bringhurst, T. A., and Brosnan, J. M. Production of grain whisky and ethanol from wheat, maize and other cereals. 112 (4) :314–323 2006.

2. Albers, E., Larsson, C., Lidén, G., Niklasson, C., and Gustafsson, L. Influence of the nitrogen source

on Saccharomyces cerevisiae anaerobic growth and product formation. 62 (9) :3187–3195

1996.

3. Arju, G. Development and implementation of high throughput peptidomics for microbial studies.

2022.

4. Arju, G., Berg, H. Y., Lints, T., and Nisamedtinov, I. Methodology for analysis of peptide

consumption by yeast during fermentation of enzymatic protein hydrolysate supplemented

synthetic medium using UPLC-IMS-HRMS. 8 (4) :145 2022.

5. Becerra-Rodríguez, C., Taghouti, G., Portier, P., Dequin, S., Casal, M., Paiva, S., et al. Yeast plasma

membrane fungal Oligopeptide Transporters display distinct substrate preferences despite their

high sequence identity. 7 (11) :963 2021.

6. Berg, H. Y., Arju, G., Becerra-Rodríguez, C., Galeote, V., and Nisamedtinov, I. Unlocking the secrets

of peptide transport in wine yeast: insights into oligopeptide transporter functions and nitrogen

source preferences. *Appl. Environ. Microbiol.* (online). 10.1128/aem.01141-23, 2023.

7. Brice, C., Sanchez, I., Tesnière, C., and Blondin, B. Assessing the mechanisms responsible for

differences between nitrogen requirements of saccharomyces cerevisiae wine yeasts in

alcoholic fermentation. 80 (4) :1330–1339 2014.

8. Briggs, D. E., Hough, J. S., Young, T. W., and Stevens, R. *Malting and brewing science: Malt and

sweet wort, volume 1*. Chapman and Hall, London, England. 1981.

9. Bringhurst, T. A., Harrison, B. M., and Brosnan, J. Scotch whisky: Raw material selection and

processing. In: *Whisky and Other Spirits*. Elsevier, pp.137–203. 2022.

10. Butré, C. I., Sforza, S., Gruppen, H., and Wierenga, P. A. Introducing enzyme selectivity: a

quantitative parameter to describe enzymatic protein hydrolysis. 406 (24) :5827–5841 2014.

11. Crépin, L., Nidelet, T., Sanchez, I., Dequin, S., and Camarasa, C. Sequential use of nitrogen

compounds by Saccharomyces cerevisiae during wine fermentation: a model based on kinetic

and regulation characteristics of nitrogen permeases. 78 (22) :8102–8111 2012.

12. Cruz, S. H., Cilli, E. M., and Ernandes, J. R. Structural complexity of the nitrogen source and

influence on yeast growth and fermentation. 108 (1) :54–61 2002.

13. Daly, M., Bromilow, S. N., Nitride, C., Shewry, P. R., Gethings, L. A., and Mills, E. N. C. Mapping

coeliac toxic motifs in the prolamin seed storage proteins of barley, rye, and oats using a curated

sequence database. *Front. Nutr.* (online). 10.3389/fnut.2020.00087, 2020.

14. Damon, C., Vallon, L., Zimmermann, S., Haider, M. Z., Galeote, V., Dequin, S., et al. A novel fungal

family of oligopeptide transporters identified by functional metatranscriptomics of soil

eukaryotes. 5 (12) :1871–1880 2011.

15. Dickinson, J. R., Salgado, L. E. J., and Hewlins, M. J. E. The catabolism of amino acids to long

chain and complex alcohols in Saccharomyces cerevisiae. 278 (10) :8028–8034 2003.

16. Duteurtre, B., Bourgeois, C., and Chollot, B. Study of the assimilation of proline by brewing yeasts.

77 (1) :28–35 1971.

17. Fiechter, G., Pavelescu, D., and Mayer, H. K. UPLC analysis of free amino acids in wines: profiling

of on-lees aged wines. 879 (17–18) :1361–1366 2011.

18. Hazelwood, L. A., Daran, J.-M., van Maris, A. J. A., Pronk, J. T., and Dickinson, J. R. The Ehrlich

pathway for fusel alcohol production: A century of research on Saccharomyces cerevisiae

metabolism. 74 (12) :3920–3920 2008.

19. Hill. Extraction of FAN from malting barley during malting and mashing. *J. Am. Soc. Brew. Chem.*

(online). 10.1094/asbcj-2014-0113-01, 2014.

20. Hill, A., and Stewart, G. Free Amino Nitrogen in brewing. 5 (1) :22 2019.

21. Jaeger, A., Zannini, E., Sahin, A. W., and Arendt, E. K. Barley protein properties, extraction and

applications, with a focus on brewers' spent grain protein. 10 (6) :1389 2021.

22. Jiménez-Martí, E., and del Olmo, M. L. Addition of ammonia or amino acids to a nitrogen-depleted

medium affects gene expression patterns in yeast cells during alcoholic fermentation. 8 (2)

:245–256 2008.

23. Jones, B. L., and Marinac, L. The effect of mashing on malt endoproteolytic activities. 50 (4) :858–

864 2002.

24. Khalid, A., Hameed, A., and Tahir, M. F. Wheat quality: A review on chemical composition,

nutritional attributes, grain anatomy, types, classification, and function of seed storage proteins

in bread making quality. 10 :1053196 2023.

25. Lekkas, C., Stewart, G. G., Hill, A. E., Taidi, B., and Hodgson, J. Elucidation of the role of
560   nitrogenous wort components in yeast fermentation. 113 (1) :3–8 2007.

561   26. Liigand, P., Kaupmees, K., and Kruve, A. Influence of the amino acid composition on the ionization
562   efficiencies of small peptides. 54 (6) :481–487 2019.

563   27. Ljungdahl, P. O., and Daignan-Fornier, B. Regulation of amino acid, nucleotide, and phosphate
564   metabolism in Saccharomyces cerevisiae. 190 (3) :885–929 2012.

565   28. Marsit, S., Mena, A., Bigey, F., Sauvage, F.-X., Couloux, A., Guy, J., et al. Evolutionary advantage
566   conferred by an eukaryote-to-eukaryote gene transfer event in wine yeasts. 32 (7) :1695–1707
567   2015.

568   29. Osman, A. M., Coverdale, S. M., Cole, N., Hamilton, S. E., Jersey, J., and Inkerman, P. A.
569   Characterisation and assessment of the role of barley malt endoproteases during malting and
570   Mashing1. 108 (1) :62–67 2002.

571   30. Perry, J. R., Basrai, M. A., Steiner, H. Y., Naider, F., and Becker, J. M. Isolation and characterization
572   of a Saccharomyces cerevisiae peptide transport gene. 14 (1) :104–115 1994.

573   31. Pyke, M. The manufacture of scotch grain whisky. 71 (3) :209–218 1965.

574   32. Salmon, J. M., and Barre, P. Improvement of nitrogen assimilation and fermentation kinetics under
575   enological conditions by derepression of alternative nitrogen-assimilatory pathways in an
576   industrial Saccharomyces cerevisiae strain. 64 (10) :3831–3837 1998.

577   33. Sprouffske, K., and Wagner, A. Growthcurver: an R package for obtaining interpretable metrics
578   from microbial growth curves. 17 (1) :172 2016.

579   34. Steiner, E., Gastl, M., and Becker, T. Protein changes during malting and brewing with focus on
580   haze and foam formation: a review. 232 (2) :191–204 2011.

581   35. Stewart, G., Kellershohn, J., and Russell, I., eds. *Whisky and other spirits*. Academic Press, San
582   Diego, CA. 2021.

583   36. Ter Schure, E. G., Silljé, H. H. W., Vermeulen, E. E., Kalhorn, J.-W., Verkleij, A. J., Boonstra, J., et
584   al. Repression of nitrogen catabolic genes by ammonia and glutamine in nitrogen-limited
585   continuous cultures of Saccharomyces cerevisiae. 144 (Pt 5) (5) :1451–1462 1998.

37. Vreeke, G. J. C., Lubbers, W., Vincken, J.-P., and Wierenga, P. A. A method to identify and quantify

the complete peptide composition in protein hydrolysates. 1201 (339616) :339616 2022.

38. Walker, G., and Hill, A. Saccharomyces cerevisiae in the Production of Whisk(e)y. 2 (4) :38 2016.

39. Walker, M. E., Nguyen, T. D., Liccioli, T., Schmid, F., Kalatzis, N., Sundstrom, J. F., et al. Genome-

wide identification of the Fermentome; genes required for successful and timely completion of

wine-like fermentation by Saccharomyces cerevisiae. 15 (1) :552 2014.

40. Wickham, H. *Ggplot2*. Springer International Publishing, Cham, Switzerland. 2016.

Table 1 – Whisky strains used in this study. The information displayed in the table has been summarised from
the provided technical data sheets of the respective strains (Lallemand Inc., Montreal, QC, Canada).

Strain	Region	Type of whisky recommended for	Alcohol tolerance (v/v)	Temperature range	Diastatic phenotype
DistilaMax® GW	North America	Grain Whisky	15%	20-34 °C	No
DistilaMax® NT	Scotland	Grain and malt whisky	12%	20-36 °C	No
DistilaMax® XP	Scotland	Malted grain whisky	15%	20-35 °C	Yes

597 **Table 2 – Nitrogen composition of the synthetic media used in this study.** All media contained 150 g/L maltose
598 and glucose (9:1, w/w). All values are displayed as milligrams of nitrogen per litre. Free proline was not taken
599 into account as it is not or poorly assimilated by yeasts during fermentation (16,25).

	AP230 (Amino acids + Peptides)	P230 (Peptides)	PP300 (Additional peptides)	PD300 (Additional ammonia)
Amino acids	130	30*	40*	30*
Peptides	100	200	260	200
Ammonia	-	-	-	70
Total	**230**	**230**	**300**	**300**

600 *Nitrogen deriving from free amino acids in the BSA hydrolysate

601 **Table 3 – Temperature, pH, and ethanol concentration gradients used in uninoculated grain mash to study**
602 **free amino acid and peptide production by native malt proteases.** The ethanol concentration was increased
603 stepwise by injecting 96% ethanol into the bioreactor.

Time (h)	Temperature (°C) (Gradient)	Ethanol (g/L) (Manual every 12h)	pH (Gradient)
0	25	0	5.2
12	29.5	20	4.75
24	34	50	4.3
36-60	34	70	4.3

Figure 1 – Cell growth and fermentation kinetics in synthetic whisky mash with different nitrogen conditions. Letters denote the statistical groups from Tukey's tests ($P < 0.05$), with lower-case letters indicating different groups of strains per medium and upper-case letters indicating the groups of media for each strain. The "time to 80% attenuation" was chosen as not all fermentations reached 100% of attenuation during the set experimental time. Principal component analysis (PCA) (D) was performed using the three fermentation parameters depicted in A, B, and C.

Figure 2 – **Consumption of peptides in synthetic whisky mash by the three whisky strains.** Peptides were grouped according to the number of amino acid residues: 2, dipeptides; 3, tripeptides; 4, tetrapeptides; 5, pentapeptides; 6, hexapeptides; 7, heptapeptides; 8, octapeptides; and 9, nonapeptides. The area under the curve (AUC) was calculated from the relative abundance curve of each peptide. The AUC values were compared to those of a virtual negative control (100% abundance over 72 hours). The consumption of a particular peptide was level 0 when its abundance AUC was equal to or higher than 80% of the negative control AUC; level 1, 60–80%; level 2, 40–60%; level 3, 20–40%; and level 4, when the peptide abundance AUC was lower than 20% of the control AUC.

621
Figure 3 – CO₂ production rate and ethanol concentration by volume (ABV) during laboratory-scale simulated Scotch grain whisky fermentation with three commercial whisky strains.

Figure 4 – Changes in FAA and oligopeptide (≤ 2 kDa) content during fermentation of Scotch grain mash and uninoculated grain mash. One experiment with uninoculated grain mash, simulating the fermentation process with changing pH and EtOH concentration, was conducted to determine the background production of potentially available nitrogen sources for yeast (Table 3). Free proline was not taken into account as it is poorly assimilated by yeast during fermentation (16,25).

631
632 **Figure 5 – Fold changes in peptide abundance between samples of the uninoculated Scotch grain mash**
633 **and the fermentations of the same mash with DistilaMax® GW, NT and XP.** The "total change" depicts the
634 fold change between peptides candidates at the 0-hour sample and the 36-hour sample.