A VgrG2b fragment cleaved by caspase-11/4 promotes *Pseudomonas aeruginosa*

infection through suppressing the NLRP3 inflammasome

Yan Qian[1,2,3,5], Qiannv Liu[1,2,3,5], Weitao Li[1,2,3], Chunlei Wang[1,2,3], Chun Kong[1,2,3],

Mengqian Li[1,2,3], Shuo Wang[4], Pengyan Xia[1,2,3*]

[1]Department of Immunology, School of Basic Medical Sciences, Peking University,

Beijing, 100191, China

[2]NHC Key Laboratory of Medical Immunology, Peking University, Beijing, 100191,

China

[3]Key Laboratory of Molecular Immunology, Chinese Academy of Medical Sciences,

Beijing, 100191, China

[4]CAS Key Laboratory of Pathogen Microbiology and Immunology, Institute of

Microbiology, Chinese Academy of Sciences, 100101 Beijing, China

[*]Address correspondence and reprint requests to Dr. Pengyan Xia, Department of

Immunology, School of Basic Medical Sciences, Peking University, Beijing, 100191,

China, Tel: 86-10-82805759; E-mail: xiap@pku.edu.cn

[5]These authors contributed equally to this work.

Running title: Caspase-11 cleaves VgrG2b to inhibit the NLRP3 inflammasome.

Abstract

The T6SS of *Pseudomonas aeruginosa* plays an essential role in the establishment

of chronic infections. Inflammatory cytokines mediated by inflammasomes are crucial for

the body to resist bacterial infections. Here we found that during the infection of *P.*

aeruginosa, non-canonical inflammasome was activated in macrophages, but the

activation of downstream NLRP3 inflammasome was inhibited. The VgrG2b of *P.*

aeruginosa is recognized and cleaved by caspase-11, generating a free C-terminal

fragment. The VgrG2b C-terminus can bind to NLRP3, inhibiting the activation of the

NLRP3 inflammasome by rejecting NEK7 binding to NLRP3. Administrating a specific

peptide that inhibits the cleavage of VgrG2b by caspase-11 to mice can significantly

improve their survival rate during infection. Our discovery elucidates a mechanism by

which *P. aeruginosa* inhibits host immune response, providing a new approach for the

future clinical treatment of *P. aeruginosa* infections.

Key words: *P. aeruginosa*; non-canonical inflammasome; caspase-11/4; VgrG2b; NLRP3.

Introduction

Non-canonical inflammasomes are composed of caspase-11 from mice or caspase-4/5 from humans, which can recognize bacteria lipopolysaccharides (LPS) in the cytoplasm[1-3]. Once activated, non-canonical inflammasomes induce the maturation of gasdermin D (GSDMD) to form pores on cell and mitochondrial membranes[4-6]. It also activates the NOD-, LRR- and pyrin domain-containing protein 3 (NLRP3) inflammasome[5,7]. NLRP3 is an important intracellular receptor that can respond to various endogenous and exogenous stimulus signals[8,9]. NLRP3 recruits NIMA-related kinase 7 (NEK7) and undergoes polymerization, inducing the activation of caspase-1 and maturation of interleukin-1β (IL-1β)[8,10]. Non-canonical inflammasomes promote the assembly and final activation of the NLRP3 inflammasome through activating the nuclear orphan receptor NUR77 with the help of LPS and mitochondrial dsDNA[11].

Pseudomonas aeruginosa is an opportunistic pathogenic bacterium widely distributed in the environment, which can cause acute and chronic infections in humans[12,13]. *P. aeruginosa* has multiple secretion systems that are used to secrete virulence factors to the external environment or host cells to promote its survival and infection[13]. Among them, the type III secretion system (T3SS) and the type VI secretion system (T6SS) are two important secretion systems that play important roles in acute and chronic infections of *P. aeruginosa*, respectively[14,15]. T3SS can directly inject effector proteins into target cells to alter their function or cause target cell death[16,17]. *P. aeruginosa* encodes three different T6SSs, namely Hcp secretion island-I (H1)-T6SS, H2-T6SS, and H3-T6SS[18,19]. These T6SSs play a crucial role in biofilm formation, competition between bacteria, and toxicity to host cells[20,21]. H1-T6SS only targets bacteria and is used for competition between bacteria[22]. H2-T6SS and H3-T6SS are involved in interactions between bacteria and host cells[23,24].

The effector proteins of H2-T6SS can target both bacteria and eukaryotic cells[21,23]. T6SS is composed of structural proteins and virulence factors[21,25]. Structural proteins such as hemolysin-coregulated protein (Hcp), valine-glycine repeat protein G (VgrG), and proline-alanine-alanine-arginine repeats (PAAR) form a stable spike-like structure[26,27]. Toxic factors are transported to the extracellular space through T6SS, thereby exerting

toxic effects on other bacteria or host cells[17,21]. VgrG2b is an important component of

T6SS, which usually binds tightly to other components of T6SS such as the Hcp protein,

together forming the core component of the secretion system[28]. The VgrG2b protein has a

special structure for recognizing and binding effector proteins, which are transported to

the tip of the secretion system and injected into the target cells[25,28]. VgrG2b entering

eukaryotic cells can also regulate the biological functions of host cells through binding to

cytoskeletal proteins[28,29].

Here we found that *P. aeruginosa* infection activates caspase-11 but inhibits the

activation of NLRP3 inflammasomes in macrophages. The VgrG2b entering the host cell

is recognized and cleaved by caspase-11 to generate a free C-terminal fragment. This

fragment can bind to NLRP3 and inhibit its binding to NEK7, thereby preventing the

activation of the NLRP3 inflammasome. The inhibitory effect of *P. aeruginosa* effectively

avoids further activation of the host immune system and plays a crucial role in establishing

bacterial infection. Our findings may provide a new perspective for the treatment of *P.*

aeruginosa infections in the future.

Results

***P. aeruginosa* causes abnormal signaling of non-canonical inflammasome**

Non-canonical inflammasomes are activated during the infection of Gram-negative

bacteria such as *P. aeruginosa in vivo*[2,7,30]. Obstruction of the inflammasome pathway can

lead to chronic infections[31,32]. To investigate whether T6SS of *P. aeruginosa* modulated

the non-canonical inflammasome pathway to establish chronic infections, we used an

RetS mutant strain that specifically expresses T6SS[14,33,34]. As an extracellular bacterial, *P.*

aeruginosa synthesizes outer membrane vesicles (OMV) to induce non-canonical

inflammasome activation[35]. We therefore added *P. aeruginosa* and OMVs to

macrophages *in vitro* to simulate the real situation of infection *in vivo* as much as

possible[36]. We found that activation of caspase-11 and cleavage of GSDMD in BMDM

cells were not affected after incubation with *P. aeruginosa* and OMVs (Fig. 1A, B).

However, activation of caspase-1 was significantly inhibited in cells infected with *P.*

aeruginosa (Fig. 1A). Consistent with the results of immunoblotting, the secretion of IL-1β

was diminished after infection with *P. aeruginosa* (Fig. 1C). Although the mortality rate of *P.*

aeruginosa-infected cells was increased (Fig. 1D) and the survival rate was decreased

(Fig. 1E), there was not much difference compared to cells stimulated with intracellular *E.*

coli. These results indicate that the activation of NLRP3 inflammasomes post

non-canonical inflammasome stimulation is inhibited in the presence of *P. aeruginosa*.

Similar results were observed in human macrophage experiments, with normal caspase-4

activation and GSDMD cleavage after *P. aeruginosa* infection (Fig. S1A, B). However, the

activation of caspase-1 was suppressed and IL-1β could not be secreted normally in

these cells (Fig. S1A, C). *P. aeruginosa* infection, the cell mortality rate was increased (Fig.

S1D) and the survival rate was decreased (Fig. S1E), showing no much difference with

cells treated only with intracellular *E. coli*.

There might be two mechanisms by which NLRP3 inflammasomes were inhibited

during this process. One was that the NLRP3 inflammasome lacked necessary stimuli for

activation, and the other was that the NLRP3 inflammasome was inhibited by a bacterial

substance. The release of mitochondrial DNA into the cytoplasm is a prerequisite for

NLRP3 activation in the non-canonical inflammasome pathway. We first detected the

localization of GSDMD N-terminus on the mitochondrial membrane during *P. aeruginosa*

infection. The results indicate that GSDMD-N can be normally localized on the

mitochondrial membrane (Fig. 1F). In addition, the release of mitochondrial DNA into the

cytoplasm was not affected by *P. aeruginosa* infection (Fig. S1F). The binding of NUR77

and NLRP3 was not affected during the infection of *P. aeruginosa* (Fig. S1G). However,

the binding of NLRP3 to NEK7 was significantly inhibited (Fig. S1H). These data suggest

that the inhibition of NLRP3 is potentially influenced by a certain inhibitory factor from *P.*

aeruginosa.

To verify this, we introduced LPS and mitochondrial DNA into macrophages to

induce NLRP3 inflammasome activation, bypassing the caspase-11-GSDMD

prerequisite[11]. To avoid the activation of the AIM2 inflammasome by mitochondrial DNA,

we conducted the following experiments under an *Aim2* knockout background. We found

that after introducing LPS and mitochondrial DNA, NLRP3 inflammasomes were

activated (Fig. 1G). However, NLRP3 activation was inhibited (Fig. 1G) and the secretion

of IL-1β was tremendously inhibited (Fig. 1H) after incubation with *P. aeruginosa*. The

percentage of cells undergoing pyroptosis (Fig. 1I) and the cell viability (Fig. 1J) did not

change due to infection with *P. aeruginosa*. This was consistent with previous reports

suggesting that GSDMD cleavage by caspase-11, rather than subsequent GSDMD

cleavage by NLRP3, is the main factor causing cell death[37,38]. Next, we used *Gsdmd* and

Aim2 double knockout cells to detect the effect of *P. aeruginosa* on NLRP3

inflammasome activation. We found that after *Gsdmd* knockout, the NLRP3

inflammasome became active when LPS and mitochondrial DNA were introduced into

macrophages. However, NLRP3 was still inhibited after incubation with *P. aeruginosa*

(Fig. S1I). The secretion of IL-1β was also inhibited due to *Gsdmd* deficiency (Fig. S1J).

The cell death rate was low due to the absence of GSDMD (Fig. S1K), while the

percentage of viable cells was relatively high (Fig. S1L). These results suggest that

potent inhibitory factors of *P. aeruginosa* may act upstream of GSDMD. Therefore, we

conducted infection experiments using *Casp11* and *Aim2* double knockout cells. After

incubation with *P. aeruginosa*, the activation of NLRP3 inflammasome was not affected

(Fig. 1K). The secretion of IL-1β was no longer inhibited (Fig. 1L). And the cell death rate

was high due to the processing of GSDMD by activated caspase-1 (Fig. 1M), while the

survival rate was low (Fig. 1N). These data indicate that the inhibitory factors of *P.*

aeruginosa requires caspase-11 to take effect.

Caspase-11 cleaves VgrG2b of *P. aeruginosa*

In order to search for potential substrates of caspase-11 in *P. aeruginosa*, we

designed a systematic screening system (Fig. 2A). Simply, we constructed two

fluorescent plasmids, one containing green fluorescent protein (GFP), polyclonal sites,

and mitochondrial outer membrane localization signals, and the other containing red

fluorescent protein (RFP) and mitochondrial outer membrane localization signals. We

extracted the mRNA of *P. aeruginosa* and prepared it into a library for insertion into GFP

containing plasmids. We co-transfected it with the RFP plasmid into HEK293T cells and

sorted the monoclonal cells containing both RFP and GFP signals into a 96-well plate

using flow cytometry. Subsequently, the plasmid encoding caspase-11 was transfected

into cells, and cells with disappearance of green signal on mitochondria were selected.

The target proteins in these cells were cleaved by caspase-11, causing the detachment of

GFP from mitochondria. After sequencing, we found that VgrG1a and VgrG2b repeatedly

appeared in cells that met the screening criteria.

We cloned the genes of VgrG1a and VgrG2b and cleaved them using mouse

caspase-11. We found that they can both be cleaved (Fig. S2A and Fig. 2B). We also

found that they can be cleaved by human caspase-4 (Fig. S2B and Fig. 2C), but not by

caspase-11/4 mutants lacking enzymatic activities. These results indicate that our

screening system is reliable. The number of target genes screened was very limited,

which was consistent with the particularly narrow substrate spectrum of caspae-11[39]. We

also tested other members of the VgrG family and found that neither mouse caspase-11

nor human caspase-4 could cleave these members (Fig. S2C–L). VgrG1a and VgrG2b

belong to the secretion systems of H1-T6SS and H2-T6SS, and H1-T6SS does not have

an effect on eukaryotic cells[22,28]. Therefore, our subsequent research focused on VgrG2b.

Our next concern was whether VgrG2b entered macrophages. We found that after

co-culturing of *P. aeruginosa* with BMDM cells, the VgrG2b protein appeared in

macrophages (Fig. 2D). Immunofluorescence experiments also showed the presence of

bacterial proteins in macrophages after co-incubation with *P. aeruginosa* (Fig. 2E). We

then incubated *P. aeruginosa* with wild-type BMDM cells and found that VgrG2b was

cleaved in macrophages, while the cleavage band of VgrG2b disappeared in *Casp11$^{-/-}$*

BMDM cells (Fig. 2F). We restored caspase-11 expression in *Casp11$^{-/-}$* BMDM cells and

found that the presence of caspase-11 induced VgrG2b cleavage (Fig. 2G). We also

rescued *Casp11$^{-/-}$* BMDM cells with wild-type and enzymatic mutant caspase-11 and

found that mutant caspase-11 could not induce VgrG2b cleavage (Fig. 2H). In addition,

we incubated human macrophages with *P. aeruginosa*. As expected, VgrG2b was cleaved

in human cells (Fig. 2I). Finally, to verify that *P. aeruginosa* affected inflammasome

activation through H2-T6SS, we generated a *P. aeruginosa* strain with VgrG2b deficiency.

We found that *P. aeruginosa* lacking VgrG2b no longer inhibited the NLRP3

inflammasome (Fig. 2J and Fig. S2M) and IL-1β could be secreted normally in the mutant

bacteria-treated cells (Fig. 2K). However, the percentage of cell death was not affected

(Fig. 2L) and the level of cell viability remained unchanged among mutants and wild-type

controls (Fig. 2M). These data indicate that H2-T6SS plays an important role in inhibiting

host immune response through VgrG2b during *P. aeruginosa* infection, primarily targeting

NLRP3 inflammasome activation rather than caspase-11 activation and GSDMD

cleavage.

VgrG2b cleavage at its C-terminus is crucial for NLRP3 inhibition

Caspase-11 recognizes its substrate through a tertiary structure formed by its

p22/p10 fragments[40]. To further verify that VgrG2b was a substrate of caspase-11, we

co-expressed caspase-11 p22/p10 and VgrG2b in HEK293T cells.

Co-immunoprecipitation results showed that VgrG2b bound to the p22/p10 complex (Fig.

3A). VgrG2b also bound to the p22/p10 complex of human caspase-4 (Fig. S3A). These

results suggest that VgrG2b can indeed be recognized by caspase-11/4. Based on the

size of the cleaved VgrG2b fragment, we speculated that the cleavage site should be at its

C-terminus. We constructed mutant plasmids carrying aspartic acid substitutions for

VgrG2b and found that only the D883A mutant was no longer cleaved by caspase-11 (Fig.

3B). We also verified that this VgrG2b mutant could not be cleaved by human caspase-4

(Fig. 3C). Meanwhile, we constructed a chimeric protein containing VgrG2a N-terminus, a

linker sequence around D883 and VgrG2b C-terminus (Fig. S3B). This chimeric protein

could be cleaved by caspase-11 (Fig. 3D). Another chimeric protein containing VgrG2b

N-terminus, a linker sequence around D883 and GFP cannot be cleaved by caspase-11

(Fig. S3C). Based on the substrate cleavage characteristic of caspase-11 that it needs to

bind the target first to mediate cleavage[40], we speculated that the VgrG2b C-terminus was

the site where VgrG2b bound to caspase-11. Therefore, we tested the binding of VgrG2b

C-terminus with caspase-11. Indeed, the C-terminus of VgrG2b bound to the p22/p10

complex (Fig. 3E). Similarly, VgrG2b C-terminus also bound to p22/p10 of human

caspase-4 (Fig. S3D). VgrG2b was recombinantly expressed and incubated with the

p22/p10 complex. We found that caspase-11 induced the cleavage of the recombinant

VgrG2b (Fig. 3F).

We generated a *P. aeruginosa* strain with a D883A mutation in the VgrG2b gene.

After incubating this bacterium with mouse BMDM, mutant VgrG2b did not undergo

cleavage (Fig. 3G). Consistently, the NLRP3 inflammasome was no longer inhibited (Fig.

3H, I) and the secretion of IL-1β was restored post incubation with this *P. aeruginosa*

mutant (Fig. 3J), indicating that VgrG2b cleavage by caspase-11 is crucial for NLRP3

inhibition. The cell death and survival rates were not affected (Fig. 3K, L). Similarly, after

incubation of the *P. aeruginosa* mutant with human macrophages, the NLRP3

inflammasome was not suppressed (Fig. S3E, F) and IL-1β was secreted normally (Fig.

S3G). The cell death and survival rates were not affected (Fig. S3H, I). These results

indicate that VgrG2b cleavage at its C-terminus is crucial for NLRP3 inhibition.

VgrG2b C-terminus interacts with NLRP3

In order to investigate how VgrG2b C-terminus inhibited the NLRP3 inflammasome,

we performed a yeast two-hybrid screening using the C-terminus of VgrG2b as a bait and

screened a mouse bone marrow cDNA library. In the positive clones obtained from the

screening, the NLRP3 fragments repeatedly appeared (Fig. 4A). We validated their

interaction in HEK293T cells and found that VgrG2b C-terminus indeed bound to NLRP3

(Fig. 4B). Full-length VgrG2b also bound to NLRP3 (Fig. 4C), but the N-terminus of

VgrG2b did not bind to NLRP3 (Fig. S4A). VgrG2b C-terminus was introduced into BMDM

cells through cell-penetrating peptides, and it was found that the C-terminus interacted

with endogenous NLRP3 (Fig. 4D). Similarly, VgrG2b C-terminus also interacted with the

human NLRP3 (Fig. S4B). In addition, we incubated *P. aeruginosa* with mouse BMDM

cells and found that NLRP3 interacted with the C-terminus of VgrG2b rather than the

full-length one (Fig. 4E). The same result was observed after incubation of *P. aeruginosa*

with human macrophages, where NLRP3 bound to the C-terminus of VgrG2b (Fig. S4C).

The colocalization between VgrG2b C-terminus and NLRP3 was observed inside BMDM

cells (Fig. 4F). The VgrG2b C-terminus did not bind to mouse GSDMD (Fig. S4D), ASC

(Fig. S4E), nor caspase-1 (Fig. S4F), indicating that the binding targets of VgrG2b in host

cells is very specific. We further searched for the region where VgrG2b C-terminus

interacted with NLRP3 (Fig. 4G), and found that their interaction area was at the

NACHT-LRR junction of NLRP3 (Fig. 4H, I).

In *Casp11* knockout BMDM cells, VgrG2b could not be cleaved and did not bind to

NLRP3 (Fig. 4J). However, in *Gsdmd* knockout cells, VgrG2b still underwent cleavage

and bound to NLRP3 (Fig. S4G). In the absence of GSDMD, the NLRP3 inflammasome

was not activated[5,37]. This result indicates that the binding of VgrG2b C-terminus to

NLRP3 is independent of NLRP3 activation, but requires a cleaved segment of VgrG2b

processed by caspase-11. In all, these results suggest that the cleaved VgrG2b

C-terminus binds NLRP3.

VgrG2b C-terminus suppresses the NLRP3 inflammasome

Whether VgrG2b C-terminus directly inhibited the activation of NLRP3

inflammasomes was the focus of our next exploration. We validated this hypothesis in the

HEK293T recombination system. The VgrG2b C-terminus significantly inhibited the

activation of NLRP3 inflammasome originated from mouse (Fig. 5A) or human (Fig. S5A).

In addition, overexpression of VgrG2b C-terminus suppressed the activation of classical

NLRP3 inflammasomes in both mouse (Fig. 5B) and human macrophages (Fig. S5B).

These data indicate that VgrG2b C-terminus exhibits a broad-spectrum inhibition of the

NLRP3 inflammasome. Introducing VgrG2b C-terminus into BMDM cells through

cell-penetrating peptides can significantly inhibit the activation of NLRP3 post intracellular

LPS stimulation, without affecting the activation of caspase-11 and GSDMD (Fig. 5C, D).

The secretion of IL-1β was significantly decreased (Fig. 5E). The cell death (Fig. 5F) and

survival rates (Fig. 5G) were not affected. The activation of NLRP3 could also be inhibited

by introducing VgrG2b C-terminus through cell-penetrating peptides when classical

NLRP3 inflammasomes stimuli were present (Fig. S5C), accompanied by a reduction of

IL-1β secretion (Fig. S5D), a decrease in cell death (Fig. S5E) and an increase in cell

viability (Fig. S5F).

We next searched for the specific mechanism by which VgrG2b C-terminus

inhibited the NLRP3 inflammasome. We first tested the ability of NLRP3 to catalyze ATP

hydrolysis after the addition of VgrG2b C-terminus and found that the hydrolysis potential

of NLRP3 was not affected (Fig. S5G). Meanwhile, VgrG2b C-terminus did not influence

the binding of NLRP3 to mitochondrial DNA (Fig. S5H), as well as to Nur77 (Fig. 5H). The

binding of NLRP3 to NEK7 was immensely inhibited by the C-terminus of VgrG2b (Fig. 5I).

VgrG2b C-terminus did not bind to NEK7 (Fig. S5I). Domain mapping showed that the

VgrG2b C-terminus bound to the same NLRP3 segment as NEK7 (Fig. 4G–I). We

therefore detected the competitive binding between VgrG2b C-terminus and NEK7 to

NLRP3. VgrG2b C-terminus competed against NEK7 (Fig. 5J), while NEK7 couldn't

compete with VgrG2b C-terminus (Fig. S5J), indicating that VgrG2b C-terminus has a

stronger binding ability with NLRP3 compared to NEK7.

VgrG2b cleavage is essential for *P. aeruginosa* infection

Finally, we infected mice with a *P. aeruginosa* strain carrying a D883A mutation in

the VgrG2b gene. The survival rate of mice was greatly improved when infected with the

mutant strain (Fig. 6A). the level of IL-1β in the serum was also increased (Fig. 6B), while

other inflammatory cytokines such as TNF-α and IL-6 showed little difference among

wild-type and mutant bacteria (Fig. 6C, D). The bacterial load in the lungs of mice infected

with the mutant strain was also decreased (Fig. 6E). We also detected the activation of

inflammasomes in macrophages from lung lavage fluid and found that the activation of the

NLRP3 inflammasome was activated in cells infected with the mutant strain (Fig. 6F), but

the cell viability was not affected (Fig. 6G). We detected the protein levels of cytokines in

the lavage fluid and found that the level of IL-1β was increased from mice infected with *P.*

aeruginosa mutant strain (Fig. 6H). Other inflammatory cytokines such as TNF-α and IL-6

showed little difference (Fig. 6I, J).

We designed a short peptide of VgrG2b C-terminus containing the VgrG2b linker

region with a D883A substitution and a transmembrane TAT sequence. Peptides were

then myristoylated[41], aiming to target this peptide to inner membrane and suppress

bacterial VgrG2b cleavage through competing with wild-type VgrG2b for caspase-11 (Fig.

S6A). We first conducted tests at the cellular level and found that adding this peptide to

BMDM cells after incubation with *P. aeruginosa* relieved the inhibition of the NLRP3

inflammasome (Fig. S6B) and the secretion of IL-1β was significantly increased (Fig.

S6C). The cell death rate did not change significantly and the viability rate was not

affected by this peptide (Fig. S6D, E). The same effect was observed in human

macrophages, where this designed peptide could relieve the inhibition of *P. aeruginosa* on

the NLRP3 inflammasome (Fig. S6F) and the secretion of cytokine IL-1β was also

increased (Fig. S6G). Although it had no effect on cell death and viability (Fig. S6H, I),

Finally, we delivered this peptide into the lungs of mice infected with *P. aeruginosa*

through nebulization, and found that the survival rate of mice was increased (Fig. 6K). The

level of IL-1β in the serum was increased (Fig. 6L), while other inflammatory factors such

as TNF-α and IL-6 remained unchanged (Fig. 6M, N). The bacterial load was also

decreased (Fig. 6O), indicating the potential medicinal value of this designed peptide in

the future clinical treatment of *P. aeruginosa* infections.

Discussion

In this study, we found that the structural protein VgrG2b from *P. aeruginosa* can be

cleaved by caspase-11 in host cells. The C-terminal fragment of VgrG2b bound to NLRP3,

inhibiting the activation of the NLRP3 inflammatory and the secretion of IL-1β. The

delivery of VgrC2b C-terminus mimics significantly improved the infection symptoms

caused by *P. aeruginosa*, providing an optional treatment approach for chronic infections

of *P. aeruginosa*.

The infection of *P. aeruginosa*, especially those in chronic infections, is an important

factor causing pulmonary fibrosis in clinicals[13,17]. LPS from Gram-negative bacterium can

be transported to host cells and activate non-canonical inflammasomes within them[2,3].

There are many ways for LPS from bacteria to be transported into host cells, including

binding with HMGB1 originated from the liver and entering the interior of the cells through

the surface of macrophages, as well as enveloping and delivering LPS into the host cells

through extracellular vesicles[42,43]. We observed the activation of non-canonical

inflammasomes in macrophages after infection with *P. aeruginosa*, indicating that LPS

was also transported into the cells. *P. aeruginosa* is not a typical intracellular bacterium,

and its LPS entry into target cells is likely through the out membrane vesicles *in vivo*.

P. aeruginosa mainly utilizes the T6SS secretion system to promote the

establishment of chronic infections[17]. During this process, H2-T6SS plays many important

roles, responsible for delivering known and yet to be explored virulence factors into host

cells[23,28]. Here we have found that VgrG2b is not only responsible for the delivery of

bacterial virulence proteins, but it can also serve as a virulence factor on its own. Its

uniqueness lies in the fact that it requires cutting to be effective. Whether there are other

proteins secreted through the H2-T6SS in *P. aeruginosa* that need to be cleaved by host

proteases before they exert their function is a very worthwhile research topic. Of course,

this processing from the host is not limited to cleavage, but may also be post-translational

modifications of proteins such as phosphorylation or ubiquitination[44,45]. Systematically

identifying these possible modifications and their substrates will be of great significance

for understanding the pathogenic mechanism of *P. aeruginosa*.

The non-canonical inflammasome receptor caspase-11 has a very narrow substrate

spectrum[39]. The process of substrate cleavage by inflammatory caspases can be divided

into two parts. Firstly, inflammatory caspases bind to a part of the substrate, which is often

not their cleavage site, and the other is the recognition of the true site of their cleavage.

This mechanism ensures that limited proteins are cleaved by host cells when

inflammasomes are activated, thereby confining the severity of the inflammatory response.

Here, we found that caspase-11/4 specifically cleaves the structural proteins of the

secretion system of *P. aeruginosa*, and the generated new fragments can play an

inhibitory role. This is a new way for pathogenic microorganisms to hijack the host's

immune system. Because traditional hijacking usually directly affects the host immune

system without cutting. It is very effective for *P. aeruginosa* using VgrG2b fragment to

inhibit NLRP3. On the one hand, VgrG2b transports virulence factors into the host cell,

and on the other hand, the cleaved substrate plays a role in inhibiting the NLRP3

inflammasome.

In summary, we have discovered a new pattern of recognition between bacterial

proteins and the host immune system during *P. aeruginosa* infection, which involves

hijacking the host's protease to produce a new virulence factor that can inhibit the host's

immune response.

Experimental procedures

Antibodies and reagents

Antibodies against NLRP3 (15101), IL-1β (12703, 12426), cleaved IL-1β (83186,

63124), GSDMD (93709), caspase-1 (3866, 24232, 2225) and ASC (13833, 67824) were

purchased from Cell Signaling Technology. Antibodies against caspase-1 (ab207802) and

GSDMD (ab215203) were purchased from Abcam. Antibodies against caspase-11 were:

ab180673 from Abcam, 14340 from Cell Signaling Technology, 14-9935-82 from

Invitrogen, M029-3 from MBL Beijing Biotech. Anti-c-Myc magnetic beads (88843) were

purchased from Pierce. Antibody against 6xHis tag (MA121315) was purchased from

Invitrogen. Antibodies against FLAG tag (F3165), β-actin (A1978) and anti-FLAG M2 magnetic beads (M8823) were purchased from Sigma-Aldrich. Antibodies against c-Myc (sc-40) and GST (sc-138) were purchased from Santa Cruz Biotechnology. Anti-HA tag antibody (HX1820) was purchased from Huaxingbio (Beijing). HRP-conjugated goat anti-mouse IgG (SA00001-1) and HRP-conjugated goat anti-rabbit IgG (SA00001-2) were purchased from Proteintech Group. Alexa Fluor Plus 488-conjugated goat anti-mouse IgG (A32723) and Alexa Fluor 594-conjugated goat anti-rabbit IgG (A32740) were purchased from Invitrogen. Alexa Fluor Plus 488-conjugated donkey anti-goat IgG (bs-0294D-AF488) and Alexa Fluor 594-conjugated donkey anti-rabbit IgG (bs-0295D-AF594) were purchased from Bioss (Beijing).

Glutathione sepharose 4B resin (17075601) was purchased from Cytiva. Protein A/G PLUS-Agarose beads (sc-2003) were purchased from Santa Cruz Biotechnology. DAPI (2879038) was purchased from PeproTech (BioGems). SMART MMLV reverse transcriptase (639524) was purchased from Clontech, Takara Bio. Random primer (Hexadeoxyribonucleotide mixture, pd(N)6) (3801) was purchased from Takara Bio. Protease inhibitor cocktail (11697498001) was purchased from Roche. Total RNA extraction kit (8034111) was purchased from Dakewe (Beijing). Ni-NTA agarose beads (R90115) and AminoLink coupling resin (20381) were purchased from Invitrogen. LPS (B46894) was purchased from Innochem (Beijing). DOTAP was: D10530 from Psaitong (Beijing) and 11202375001 from Roche. Nigericin (481990), Gramicidin (368020), ATP (A6559) were purchased from Merck Millipore. Cell-penetrating peptides: ab142343 from Abcam and sc-396807 from Santa Cruz. Human M-CSF (300-25) and murine M-CSF (315-02) were purchased from PeproTech. CellTiter-Glo luminescent cell viability assay kit (G7570) and CytoTox 96 non-radioactive cytotoxicity assay kit (G1780) were purchased from Promega.

Bacteria and CFU determination

E. coli (CGMCC1.2389, ATCC11775) was from China General Microbiological Culture Collection Center. *Pseudomonas aeruginosa* (PAO1) was a gift from Dr. Lvyan Ma (Institute of Microbiology, Chinese Academy of Sciences). *E. coli* and *P. aeruginosa* were preserved in LB plates and inoculated to liquid LB medium, followed by shaking at 37 °C

until they grew to the exponential phase (optical density=0.6–0.8). Bacteria were diluted

and spread into LB plates to determine the CFU numbers. When the bacteria were

cultured to the mid-exponential stage, they were washed three times with PBS and

resuspended in PBS or DMEM.

P. aeruginosa OMV purification

When *P. aeruginosa* was cultured to the mid exponential stage, cultures were

centrifugated to obtain the supernatant. After filtering through a 0.22 μm filter, the

supernatant was placed in a discontinuous sucrose gradient tube and centrifuged at a

speed of 100, 000 g for 16 h. The precipitate was resuspended by PBS and centrifuged

again to remove residual culture medium components. The precipitate was resuspended

with PBS again, resulting in the OMV sample. The sample was immediately used or

frozen and stored in a −80 °C freezer. 20 μg/ml OMV was used in each *in vitro*

experiment.

P. aeruginosa knockout and knockin strategy

The knockout plasmid pK18mobsacBGm was digested with BamHI and HindIII, and

then recovered by agarose gel electrophoresis. Primers were designed to clone

homologous arms of approximately 900 bp upstream and downstream of the target gene.

This knockout strategy preserved the starting 99bp and ending 99bp of the target gene.

The upstream and downstream homologous arms were then connected with the

linearized pk18mobsacBGm vector using homologous recombination, and the knockout

vector pk18mobsacBGm-target were obtained through sequencing. Primers used were:

RetS-up-F: 5'-ctcggtacccgggggatccccgccgtgcgcgacatgctcgccggcaa-3',   RetS-up-R:

5'-agcacgtcgctgc   ccggcgaagtcccttcg-3',   RetS-down-F:

5'-ccttcgaagggacttcgccgggcagcgacgtgctccgg-3',   RetS-down-R:

5'-cgttgtaaaacgacggccagtgccaagcttcgagggtcaggcaggcgag-3'   for   RetS   knockout;

KO-VgrG2b-up-F: 5'-attcgagctcggtacccgggggatcccttgatggaaaagagtttcaagacc-3',

KO-VgrG2b-up-R: 5'-tctcgaggaaataatctcgaacgataggctcgcagagcgcttcttccagt-3',

KO-VgrG2b-down-F: 5'-actggaagaagcgctctgcgagcctatcgttcgagattatttcctcga-3',

KO-VgrG2b-down-R: 5'-taaaacgacggccagtgccaagcttgacgacgtcgggggttctctgcctt-3'   for

VgrG2b   knockout;   KI-Myc-VgrG2b-up-F:

5'-tcgagctcggtacccgggggatccgagcacatcaccctgatgtgcggcggcgcct-3', KI-Myc-VgrG2b-up-R1:

5'-ttcgaaccgcgggccctctagactcgagcggtatcccgttgggaagttttcagt-3',   KI-Myc-VgrG2b-up-R2:

5'-atcctcttctgagatgagtttttgttcgaaccgcgggccctctagactc-3',   KI-Myc-VgrG2b-down-F:

5'-aaaaactcatctcagaagaggatctgtgaccaatgaaatgcaagaccttgctca-3',

KI-Myc-VgrG2b-down-R: 5'-taaaacgacggccagtgccaagcttcggccccgccggcgccactggcgaa-3'

for   VgrG2b-Myc   knockin;   VgrG2b-D883A-up-F:

5'-aattcgagctcggtacccgggggatccgagcaccagg gcgtggggcacgacg-3', VgrG2b-D883A-up-R:

5'-aggtcttgcatttcattggtcacagatcctcttctgagat   gag-3',   VgrG2b-D883A-down-F:

5'-actcatctcagaagaggatctgtgaccaatgaaatgcaagaccttgctc a-3', VgrG2b-D883A-down-R:

5'-taaaacgacggccagtgccaagctttttcgccggccaggcagaattcgacg-3' for VgrG2b D883A knockin.

The recipient bacterium *P. aeruginosa*, co-plasmid pRK2013, and donor bacterium

(pK18mobsacBGm-target) were inoculated in LB liquid medium (with corresponding

antibiotics added), and incubated overnight in a shaking table at 37 °C and 200 rpm. 1 ml

of co-plasmid and donor bacterial culture medium were collected separately and

centrifuged at 5, 000 rpm for 2 m. Pellets were collected and resuspended in 200 μl LB

medium. Co-plasmids and donor bacteria were mixed in a 1:1 ratio, and inoculated on LB

solid culture medium at 37 °C for 2 hours. Simultaneously, the recipient strain *P.*

aeruginosa was placed in 42 °C for 2 hours of cultivation. 1mL of receptor bacteria was

collected and centrifuged at 5, 000 rpm for 2 min and resuspended in 200 μl LB liquid

medium. The bacteria were spotted onto a mixed plaque of co-plasmids and donor

bacteria and cultured at 37 °C for 4 hours. The plaque was picked up and suspended in

the 500–800 μl LB liquid medium and plated onto LB plates with Irg and Gm resistance,

followed by culture overnight at 37 °C. A single colony was selected and transferred again

to LB plates with Irg and Gm resistance. The next day, bacteria were drawn a line on a

20–22% sucrose agar plate and cultured at 37 °C for 24–48 hours. Single colonies were

selected again according to the above method. Colonies that could grow on LB plates but

not on Gm resistant plates were chosen and subjected to PCR validation.

Cell component fractionation and mitochondrial DNA quantification

Cells were fractionated according to the guidelines of a cell fractionation kit from

Abcam (ab109719). Mitochondria were isolated from macrophages using a mitochondrial

isolation kit from Abcam (ab110170) or from Invitrogen (89874), followed by mitochondrial

DNA purification through a mitochondrial DNA isolation kit from Abcam (ab65321). Total

cytosolic DNA was extracted using a DNeasy Blood & Tissue Kit (QIAGEN) following

manufacturer's instructions. For mitochondrial DNA quantification, samples containing

mitochondrial DNA were subjected to RT-PCR analyses using the following primers:

5'-GAGATGACCAAATTTACAATG-3' (forward) and 5'-TCCTGTTCCTGCTCCTGCTTC-3'

(reverse) for mouse mitochondrial gene COX1.

Mice and infection

$Casp4^{-/-}$ (S-KO-01332), $Gsdmd^{-/-}$ (S-KO-12963) and $Aim2^{-/-}$ (S-KO-09889) mice

were purchased from Cyagen Biosciences (Jiangsu, China). Mice were intraperitoneally

challenged with poly(I:C) at a dose of 2 mg/kg body weight for 6 h and then intranasally

infected with 1×10^9 cfu of *P. aeruginosa* strain PAO1, followed by survival calculation at

the indicated days. Sera were collected and subjected to ELISA assays. For

bronchoalveolar lavage fluid (BALF) collection, mice were anaesthetized, followed by

thoracic cavity opened and a flush of circulating blood cells by injecting about 10 ml

ice-cold PBS into the right ventricle. Mouse neck was then dissected to expose trachea. A

puncture needle was inserted into the upper end of the trachea, followed by repeated

rinsing with 0.8 ml PBS for five times. BALF collections were centrifuged at 300 g at 4 °C

for 5 min, followed by ELISA assays of the supernatants or macrophage isolation in the

cell pellets.

Primary cell separation and culture

To generate BMDM cells, bone marrow cells were flushed from mouse femurs and

tibias, followed by red blood cell removal using red blood cell lysis buffer. Cells were

resuspended in alpha-MEM containing 10% (v/v) fetal bovine serum and 10 ng/ml M-CSF

and grown at 37 °C with a 5% CO_2 humidified atmosphere for 7 days. To produce human

macrophages, human bloods collected from healthy donors were mixed with an equal

volume of saline and then added to lymphocyte separation medium, followed by

centrifugation and PBMC collection. Human PBMCs were incubated with APC-conjugated

anti-CD14 antibody on ice for 30 min, followed by incubation with anti-APC-conjugated

magnetic beads. Monocytes were enriched on a magnetic separator. To test the purity of

sorted monocytes, cells were stained with APC conjugated anti-CD14 antibody, followed

by FACS examination. Cells with purity above 95% were used for subsequent

experiments. Cells were resuspended in RPMI-1640 medium containing 10% (v/v) fetal

bovine serum and 50 ng/ml human M-CSF, followed by growing at 37 °C with a 5% CO_2

humidified atmosphere for 7 days. Healthy blood samples were provided by the

researchers volunteering. Informed consents were obtained from all subjects and

experiments conformed to related principles. Study was licensed by the Ethics Committee

of Institute of Microbiology, Chinese Academy of Sciences.

Inflammasome induction

For canonical NLRP3 inflammasome activation, cells were incubated with 1 μg/ml

LPS for 3 h, followed by stimulation with 10 μM nigericin for 30 min, 0.5 μM gramicidin for

1 h and 2 mM ATP for 30 min. Cells were grown at 37 °C with a 5% CO_2 humidified

atmosphere. For non-canonical inflammasome stimulation, macrophages were primed

overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS using DOTAP

for 16 h. Otherwise, macrophages were primed overnight with 1 μg/ml Pam3CSK4 and

incubated with *P. aeruginosa* strain PAO1 at an MOI of 30 and OMVs for 2 h. Cells were

then supplemented with fresh medium containing 100 μg/ml Gentamycin and cultured for

16 h.

Plasmid transfection and lentiviral production

Transfection reagent JetPRIME (Polyplus Transfection) was used to conduct

transfection in HEK293T cells. HEK293T cells were plated 14 h earlier to achieve a

density of about 70% when transfection was performed. Lentiviral plasmids containing

exogenous genes of interest and packaging plasmids pSPAX2 and pMD2.G were

transfected into HEK293T cells at a ratio of 4:3:1. 48 h later, supernatants were collected,

followed by centrifugation at 1,000 rpm for 5 min and filtration using 0.45 μm sterile

syringe filters (Merck Millipore). The filtrate was then concentrated with a 100 kD Amicon

ultra centrifugal filter unit (Merck Millipore). Macrophage cells were incubated with

lentiviruses for 6 h, followed by culture medium replacement with fresh medium. 2 μg/ml

puromycin was added to the medium 48 h post infection for 3 days.

Cell viability and cytotoxicity assay

Cell pellets from cells stimulated with inflammasome activators were used to

measure cell viability through checking cellular ATP levels with a CellTiter-Glo

luminescent cell viability assay kit (Promega). The supernatants from cells stimulated with

inflammasome activators were collected and subjected to lactate dehydrogenase (LDH)

assay to assess cell death using a CytoTox 96 non-radioactive cytotoxicity assay kit

(Promega). Both the supernatants and cell pellets were collected for immunoblotting

analyses.

ELISA

ELISA kits detecting human IL-1β (D711068), mouse IL-1β (D721017), mouse IL-6

(D721022) and mouse TNF-α (D721217) were purchased from BBI life sciences

(Shanghai). Cell culture supernatants were collected and subjected to ELISA assays

following manufacturer's instructions.

Recombinant protein expression and purification

For recombinant expression of GST-tagged proteins, cDNA of interest was cloned

into pGEX-6P-1 vector. Plasmids were transformed into *E. coli* strain BL21 and the

bacteria grew in LB medium supplemented with 100 μg/ml ampicillin at 37 °C for 8 h,

followed by addition of IPTG to a final concentration of 0.5 mM and shaking at 150 rpm at

37 °C for 3 h. Cells were harvested and resuspended with PBS containing 1%

Triton-X-100 and lysed by sonication. Cell lysates were centrifuged at 13,000 rpm at 4 °C

for 30 min and filtered through sterile syringe filters. For crude purification of recombinant

proteins, the filtered lysate was loaded on column containing glutathione sepharose 4B

resin, followed by washing with PBS containing 1% Triton-X-100. The protein was

eventually eluted by 1 ml 10 mM L-Glutathione solution. For removal of LPS

contaminations, crudely purified proteins were further eluted through Pierce high-capacity

endotoxin removal spin columns and heparin columns to remove LPS and bacterial

genomic DNA, respectively. GST tags were removed by a PreScission protease.

Immunoprecipitation

HEK293T cells were seeded on 6-well culture plates and were transfected with

plasmids 16 h later. 24 h post transfection, cells were washed twice with PBS and lysed in

a pre-chilled buffer containing 150 mM NaCl, 50 mM Tris (pH 7.5), 1 mM EDTA, 0.5%

digitonin, 1% protease inhibitor cocktail and 10% glycerol on ice for 30 min. For

immunoprecipitation using primary cells, cells post stimulation or treatments were lysed in

a pre-chilled PBS buffer containing 0.5% digitonin, 1% protease inhibitor cocktail on ice

for 30 min. Cell lysates were centrifuged at 13,000 rpm at 4 °C for 10 min. Primary

antibodies and their isotype control IgG were immobilized to AminoLink Coupling Resin

(ThermoFisher Scientific) following the manufacturer's instructions. Supernatants were

incubated with immobilized antibodies at 4 °C for 2 h. Resins were washed three times

with PBS containing 0.5% digitonin, followed by immunoprecipitant detachment in 0.1 M

Glycine-HCl, pH 2.7. Supernatants were neutralized by adding 1/10 the volume of 2 M

Tris-HCl, pH 8.0, followed by immunoblotting with the indicated antibodies.

Immunofluorescence

Cells adhered to covers pre-coated with poly-L-lysine were stimulated with

inflammasome activators, followed by fixation with 4% paraformaldehyde for 10 min at

room temperature. Cells were then washed twice with PBS and permeabilized with PBS

containing 0.1% digitonin for 10 min, followed by washing with PBS for twice and blockage

in 10% normal goat serum at 37 °C for 30 min. Cells were incubated with primary antibody

for 2 h, followed by washing with PBS for three times and further incubation with

fluorescence-conjugated secondary antibody for 1 h. Nuclei were stained with DAPI. Cells

were visualized through an UltraVIEW VoX imaging system (PerkinElmer).

Statistical analysis

No statistical methods were used to predetermine sample size. Experiments were

independently repeated at least three times to achieve statistical significance. No

randomization or blinding procedures was used in this study. No samples were

excluded from the analysis. Data were shown as means ± SD of three technical

replicates. Data with normal distribution determined by Shapiro–Wilk normality test

were statistically analyzed by two tailed Student's t tests if not specified. The

Gehan-Breslow-Wilcoxon test was used for the analysis of survival data. Data were

analyzed by GraphPad Prism 9.0. P-values ≤ 0.05 were termed as significant; NS

means non-significant where $P>0.05$.

Acknowledgements

We thank Yingyu Chen (Peking University) for technical help. This work was supported by the National Natural Science Foundation of China (92369104, 82271790, 92169113), Beijing Natural Science Foundation (JQ23028, 7212067), the National Key R&D Program of China (2021YFA1300202, 2022YFC2302900), Strategic Priority Research Programs of the Chinese Academy of Sciences (XDB29020000), Key Research Program of Frontier Sciences of Chinese Academy of Sciences (ZDBS-LY-SM025), CAS Project for Young Scientists in Basic Research (YSBR-010), Fok Ying Tung Education Foundation to P.X., Youth Innovation Promotion Association of CAS to S.W.

Author contributions

Y.Q. and Q.L. performed experiments and analyzed data; W.L., C.W., C.K., M.L. and S.W. performed experiments; P.X. initiated the study, designed and performed experiments, analyzed data, and wrote the paper.

References

1. Hagar, J.A., Powell, D.A., Aachoui, Y., Ernst, R.K., and Miao, E.A. (2013). Cytoplasmic LPS activates caspase-11: implications in TLR4-independent endotoxic shock. Science (New York, N.Y.) *341*, 1250-1253. 10.1126/science.1240988.

2. Kayagaki, N., Wong, M.T., Stowe, I.B., Ramani, S.R., Gonzalez, L.C., Akashi-Takamura, S., Miyake, K., Zhang, J., Lee, W.P., Muszyński, A., et al. (2013). Noncanonical inflammasome activation by intracellular LPS independent of TLR4. Science (New York, N.Y.) *341*, 1246-1249. 10.1126/science.1240248.

3. Shi, J., Zhao, Y., Wang, Y., Gao, W., Ding, J., Li, P., Hu, L., and Shao, F. (2014). Inflammatory caspases are innate immune receptors for intracellular LPS. Nature *514*, 187-192. 10.1038/nature13683.

4. Huang, L.S., Hong, Z., Wu, W., Xiong, S., Zhong, M., Gao, X., Rehman, J., and Malik, A.B. (2020). mtDNA Activates cGAS Signaling and Suppresses the YAP-Mediated Endothelial Cell Proliferation Program to Promote Inflammatory Injury. Immunity *52*, 475-486.e475. 10.1016/j.immuni.2020.02.002.

5. Kayagaki, N., Stowe, I.B., Lee, B.L., O'Rourke, K., Anderson, K., Warming, S., Cuellar, T., Haley, B., Roose-Girma, M., Phung, Q.T., et al. (2015). Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. Nature *526*, 666-671. 10.1038/nature15541.

6. Shi, J., Zhao, Y., Wang, K., Shi, X., Wang, Y., Huang, H., Zhuang, Y., Cai, T., Wang, F., and Shao, F. (2015). Cleavage of GSDMD by inflammatory caspases determines

pyroptotic cell death. Nature *526*, 660-665. 10.1038/nature15514.

7. Rathinam, V.A.K., Zhao, Y., and Shao, F. (2019). Innate immunity to intracellular LPS.
Nature immunology *20*, 527-533. 10.1038/s41590-019-0368-3.

8. Barnett, K.C., Li, S., Liang, K., and Ting, J.P. (2023). A 360° view of the inflammasome:
Mechanisms of activation, cell death, and diseases. Cell *186*, 2288-2312.
10.1016/j.cell.2023.04.025.

9. Christgen, S., Place, D.E., and Kanneganti, T.D. (2020). Toward targeting inflammasomes:
insights into their regulation and activation. Cell research *30*, 315-327.
10.1038/s41422-020-0295-8.

10. He, Y., Zeng, M.Y., Yang, D., Motro, B., and Núñez, G. (2016). NEK7 is an essential
mediator of NLRP3 activation downstream of potassium efflux. Nature *530*, 354-357.
10.1038/nature16959.

11. Zhu, F., Ma, J., Li, W., Liu, Q., Qin, X., Qian, Y., Wang, C., Zhang, Y., Li, Y., Jiang, D., et al.
(2023). The orphan receptor Nur77 binds cytoplasmic LPS to activate the non-canonical
NLRP3 inflammasome. Immunity *56*, 753-767.e758. 10.1016/j.immuni.2023.03.003.

12. Cendra, M.D.M., and Torrents, E. (2021). Pseudomonas aeruginosa biofilms and their
partners in crime. Biotechnology advances *49*, 107734.
10.1016/j.biotechadv.2021.107734.

13. Rossi, E., La Rosa, R., Bartell, J.A., Marvig, R.L., Haagensen, J.A.J., Sommer, L.M.,
Molin, S., and Johansen, H.K. (2021). Pseudomonas aeruginosa adaptation and
evolution in patients with cystic fibrosis. Nature reviews. Microbiology *19*, 331-342.
10.1038/s41579-020-00477-5.

14. Goodman, A.L., Kulasekara, B., Rietsch, A., Boyd, D., Smith, R.S., and Lory, S. (2004). A
signaling network reciprocally regulates genes associated with acute infection and
chronic persistence in Pseudomonas aeruginosa. Dev Cell *7*, 745-754.
10.1016/j.devcel.2004.08.020.

15. Faure, E., Mear, J.B., Faure, K., Normand, S., Couturier-Maillard, A., Grandjean, T.,
Balloy, V., Ryffel, B., Dessein, R., Chignard, M., et al. (2014). Pseudomonas aeruginosa
type-3 secretion system dampens host defense by exploiting the NLRC4-coupled
inflammasome. American journal of respiratory and critical care medicine *189*, 799-811.
10.1164/rccm.201307-1358OC.

16. Hauser, A.R. (2009). The type III secretion system of Pseudomonas aeruginosa: infection
by injection. Nature reviews. Microbiology *7*, 654-665. 10.1038/nrmicro2199.

17. Qin, S., Xiao, W., Zhou, C., Pu, Q., Deng, X., Lan, L., Liang, H., Song, X., and Wu, M.
(2022). Pseudomonas aeruginosa: pathogenesis, virulence factors, antibiotic resistance,
interaction with host, technology advances and emerging therapeutics. Signal
transduction and targeted therapy *7*, 199. 10.1038/s41392-022-01056-1.

18. Hachani, A., Lossi, N.S., Hamilton, A., Jones, C., Bleves, S., Albesa-Jové, D., and Filloux,
A. (2011). Type VI secretion system in Pseudomonas aeruginosa: secretion and
multimerization of VgrG proteins. The Journal of biological chemistry *286*, 12317-12327.
10.1074/jbc.M110.193045.

19. Mougous, J.D., Cuff, M.E., Raunser, S., Shen, A., Zhou, M., Gifford, C.A., Goodman, A.L.,
Joachimiak, G., Ordoñez, C.L., Lory, S., et al. (2006). A virulence locus of Pseudomonas
aeruginosa encodes a protein secretion apparatus. Science (New York, N.Y.) *312*,

1526-1530. 10.1126/science.1128393.

20.   Chen, L., Zou, Y., She, P., and Wu, Y. (2015). Composition, function, and regulation of
T6SS in Pseudomonas aeruginosa. Microbiological research *172*, 19-25.
10.1016/j.micres.2015.01.004.

21.   Ho, B.T., Dong, T.G., and Mekalanos, J.J. (2014). A view to a kill: the bacterial type VI
secretion system. Cell host & microbe *15*, 9-21. 10.1016/j.chom.2013.11.008.

22.   Hood, R.D., Singh, P., Hsu, F., Güvener, T., Carl, M.A., Trinidad, R.R., Silverman, J.M.,
Ohlson, B.B., Hicks, K.G., Plemel, R.L., et al. (2010). A type VI secretion system of
Pseudomonas aeruginosa targets a toxin to bacteria. Cell host & microbe *7*, 25-37.
10.1016/j.chom.2009.12.007.

23.   Jiang, F., Waterfield, N.R., Yang, J., Yang, G., and Jin, Q. (2014). A Pseudomonas
aeruginosa type VI secretion phospholipase D effector targets both prokaryotic and
eukaryotic cells. Cell host & microbe *15*, 600-610. 10.1016/j.chom.2014.04.010.

24.   Russell, A.B., LeRoux, M., Hathazi, K., Agnello, D.M., Ishikawa, T., Wiggins, P.A., Wai,
S.N., and Mougous, J.D. (2013). Diverse type VI secretion phospholipases are
functionally plastic antibacterial effectors. Nature *496*, 508-512. 10.1038/nature12074.

25.   Durand, E., Cambillau, C., Cascales, E., and Journet, L. (2014). VgrG, Tae, Tle, and
beyond: the versatile arsenal of Type VI secretion effectors. Trends in microbiology *22*,
498-507. 10.1016/j.tim.2014.06.004.

26.   Leiman, P.G., Basler, M., Ramagopal, U.A., Bonanno, J.B., Sauder, J.M., Pukatzki, S.,
Burley, S.K., Almo, S.C., and Mekalanos, J.J. (2009). Type VI secretion apparatus and
phage tail-associated protein complexes share a common evolutionary origin.
Proceedings of the National Academy of Sciences of the United States of America *106*,
4154-4159. 10.1073/pnas.0813360106.

27.   Shneider, M.M., Buth, S.A., Ho, B.T., Basler, M., Mekalanos, J.J., and Leiman, P.G.
(2013). PAAR-repeat proteins sharpen and diversify the type VI secretion system spike.
Nature *500*, 350-353. 10.1038/nature12453.

28.   Sana, T.G., Baumann, C., Merdes, A., Soscia, C., Rattei, T., Hachani, A., Jones, C.,
Bennett, K.L., Filloux, A., Superti-Furga, G., et al. (2015). Internalization of Pseudomonas
aeruginosa Strain PAO1 into Epithelial Cells Is Promoted by Interaction of a T6SS
Effector with the Microtubule Network. mBio *6*, e00712. 10.1128/mBio.00712-15.

29.   Wood, T.E., Howard, S.A., Förster, A., Nolan, L.M., Manoli, E., Bullen, N.P., Yau, H.C.L.,
Hachani, A., Hayward, R.D., Whitney, J.C., et al. (2019). The Pseudomonas aeruginosa
T6SS Delivers a Periplasmic Toxin that Disrupts Bacterial Cell Morphology. Cell reports
29, 187-201.e187. 10.1016/j.celrep.2019.08.094.

30.   Balakrishnan, A., Karki, R., Berwin, B., Yamamoto, M., and Kanneganti, T.D. (2018).
Guanylate binding proteins facilitate caspase-11-dependent pyroptosis in response to
type 3 secretion system-negative Pseudomonas aeruginosa. Cell Death Discov *4*, 3.
10.1038/s41420-018-0068-z.

31.   Huus, K.E., Joseph, J., Zhang, L., Wong, A., Aaron, S.D., Mah, T.F., and Sad, S. (2016).
Clinical Isolates of Pseudomonas aeruginosa from Chronically Infected Cystic Fibrosis
Patients Fail To Activate the Inflammasome during Both Stable Infection and Pulmonary
Exacerbation. J Immunol *196*, 3097-3108. 10.4049/jimmunol.1501642.

32.   Phuong, M.S., Hernandez, R.E., Wolter, D.J., Hoffman, L.R., and Sad, S. (2021).

Impairment in inflammasome signaling by the chronic Pseudomonas aeruginosa isolates from cystic fibrosis patients results in an increase in inflammatory response. Cell death & disease *12*, 241. 10.1038/s41419-021-03526-w.

33. Allsopp, L.P., Wood, T.E., Howard, S.A., Maggiorelli, F., Nolan, L.M., Wettstadt, S., and Filloux, A. (2017). RsmA and AmrZ orchestrate the assembly of all three type VI secretion systems in Pseudomonas aeruginosa. Proceedings of the National Academy of Sciences of the United States of America *114*, 7707-7712. 10.1073/pnas.1700286114.

34. Han, Y., Wang, T., Chen, G., Pu, Q., Liu, Q., Zhang, Y., Xu, L., Wu, M., and Liang, H. (2019). A Pseudomonas aeruginosa type VI secretion system regulated by CueR facilitates copper acquisition. PLoS pathogens *15*, e1008198. 10.1371/journal.ppat.1008198.

35. Deo, P., Chow, S.H., Han, M.L., Speir, M., Huang, C., Schittenhelm, R.B., Dhital, S., Emery, J., Li, J., Kile, B.T., et al. (2020). Mitochondrial dysfunction caused by outer membrane vesicles from Gram-negative bacteria activates intrinsic apoptosis and inflammation. Nat Microbiol *5*, 1418-1427. 10.1038/s41564-020-0773-2.

36. Esoda, C.N., and Kuehn, M.J. (2019). Pseudomonas aeruginosa Leucine Aminopeptidase Influences Early Biofilm Composition and Structure via Vesicle-Associated Antibiofilm Activity. mBio *10*. 10.1128/mBio.02548-19.

37. Broz, P., Pelegrín, P., and Shao, F. (2020). The gasdermins, a protein family executing cell death and inflammation. Nature reviews. Immunology *20*, 143-157. 10.1038/s41577-019-0228-2.

38. Kayagaki, N., Warming, S., Lamkanfi, M., Vande Walle, L., Louie, S., Dong, J., Newton, K., Qu, Y., Liu, J., Heldens, S., et al. (2011). Non-canonical inflammasome activation targets caspase-11. Nature *479*, 117-121. 10.1038/nature10558.

39. Shi, X., Sun, Q., Hou, Y., Zeng, H., Cao, Y., Dong, M., Ding, J., and Shao, F. (2023). Recognition and maturation of IL-18 by caspase-4 noncanonical inflammasome. Nature *624*, 442-450. 10.1038/s41586-023-06742-w.

40. Wang, K., Sun, Q., Zhong, X., Zeng, M., Zeng, H., Shi, X., Li, Z., Wang, Y., Zhao, Q., Shao, F., and Ding, J. (2020). Structural Mechanism for GSDMD Targeting by Autoprocessed Caspases in Pyroptosis. Cell *180*, 941-955.e920. 10.1016/j.cell.2020.02.002.

41. Zuber, M.X., Strittmatter, S.M., and Fishman, M.C. (1989). A membrane-targeting signal in the amino terminus of the neuronal protein GAP-43. Nature *341*, 345-348. 10.1038/341345a0.

42. Deng, M., Tang, Y., Li, W., Wang, X., Zhang, R., Zhang, X., Zhao, X., Liu, J., Tang, C., Liu, Z., et al. (2018). The Endotoxin Delivery Protein HMGB1 Mediates Caspase-11-Dependent Lethality in Sepsis. Immunity *49*, 740-753.e747. 10.1016/j.immuni.2018.08.016.

43. Vanaja, S.K., Russo, A.J., Behl, B., Banerjee, I., Yankova, M., Deshmukh, S.D., and Rathinam, V.A.K. (2016). Bacterial Outer Membrane Vesicles Mediate Cytosolic Localization of LPS and Caspase-11 Activation. Cell *165*, 1106-1119. 10.1016/j.cell.2016.04.015.

44. Ashida, H., Kim, M., and Sasakawa, C. (2014). Exploitation of the host ubiquitin system by human bacterial pathogens. Nature reviews. Microbiology *12*, 399-413.

744   10.1038/nrmicro3259.

745 45. Li, Z., Liu, W., Fu, J., Cheng, S., Xu, Y., Wang, Z., Liu, X., Shi, X., Liu, Y., Qi, X., et al.
746   (2021). Shigella evades pyroptosis by arginine ADP-riboxanation of caspase-11. Nature
747   *599*, 290-295. 10.1038/s41586-021-04020-1.

 Figure legends

Figure 1. *P. aeruginosa* infection suppresses NLRP3 inflammasome. (A–E) Wild-type

BMDM cells were primed overnight with 1 μg/ml Pam3CSK4 and incubated with *E. coli* or

P. aeruginosa (ΔRetS PAO1) at an MOI of 30 and 20 μg/ml OMVs for 2 h. Cells were then

supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells were lysed and

immunoblotted as indicated 16 h post infection (A). Band intensities of cleaved

caspase-11 (top) and GSDMD (bottom) were quantified and compared to that of β-actin

(B). Cell culture supernatants were collected for an ELISA assay to determine the

secreted IL-1β protein levels 16 h post infection (C). Cytotoxicity was determined by LDH

release assay in cell culture supernatants 16 h post infection (D). Cell viability was

determined by an ATP quantification assay in cell pellets 16 h post infection (E). (F)

Wild-type BMDM cells were primed overnight with 1 μg/ml Pam3CSK4 and incubated with

E. coli or ΔRetS PAO1 at an MOI of 30 and 20 μg/ml OMVs for 2 h. Cells were then

supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells were subjected

to cellular component fractionation and immunoblotted as indicated 16 h post infection.

(G–J) *Aim2$^{-/-}$* BMDM cells were primed overnight with 1 μg/ml Pam3CSK4, followed by

transfection of 2 μg/ml LPS and 2 μg/ml extracted BMDM mitochondrial DNA using

DOTAP with or without the incubation of ΔRetS PAO1 at an MOI of 30 for 2 h. Cells were

then supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells were

lysed and immunoblotted 16 h post infection (G). Cell culture supernatants were collected

for an ELISA assay to determine the secreted IL-1β protein levels 16 h post infection (H).

Cytotoxicity was determined by LDH release assay in cell culture supernatants 16 h post

infection (I). Cell viability was determined by an ATP quantification assay in cell pellets 16

h post infection (J). (K–N) *Casp11$^{-/-}$;Aim2$^{-/-}$* BMDM cells were primed overnight with 1

μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS and 2 μg/ml extracted BMDM

mitochondrial DNA using DOTAP with or without the incubation of ΔRetS PAO1 at an MOI

of 30 for 2 h. Cells were then supplemented with fresh medium containing 100 μg/ml

Gentamycin. Cells were lysed and immunoblotted 16 h post infection (K). Cell culture

supernatants were collected for an ELISA assay to determine the secreted IL-1β protein

levels 16 h post infection (L). Cytotoxicity was determined by LDH release assay in cell

culture supernatants 16 h post infection (M). Cell viability was determined by an ATP

quantification assay in cell pellets 16 h post infection (N). Data were shown as means±SD.

Experiments were repeated three times with similar results.

Figure 2. *P. aeruginosa* VgrG2b is cleaved by caspase-11. (A) Screening strategy for

identifying caspase-11 substrates in *P. aeruginosa*. Briefly, *P. aeruginosa* cDNAs were

cloned into GFP vectors containing a mitochondrial localization signal. RFP vectors with a

mitochondrial localization signal were used as transfection controls. HEK293T cells were

co-transfected with these vectors and caspase-11. Cells losing GFP signals on

mitochondria were sequenced. (B, C) Plasmids encoding VgrG2b and mouse caspase-11

p22/p10 (B) or human caspase-4 p22/p10 (C) were co-transfected into HEK293T cells for

24 h, followed by immunoblotting with antibodies against the indicated proteins. (D)

Wild-type BMDM cells were primed overnight with 1 μg/ml Pam3CSK4 and incubated with

ΔRetS PAO1 or VgrG2b-Myc knockin ΔRetS PAO1 at an MOI of 30 and 20 μg/ml OMVs

for 2 h. Cells were then supplemented with fresh medium containing 100 μg/ml

Gentamycin. Cells were lysed and immunoblotted as indicated 16 h post infection. (E)

Wild-type BMDM cells were primed overnight with 1 μg/ml Pam3CSK4 and incubated with

VgrG2b-Myc knockin ΔRetS PAO1 at an MOI of 30 and 20 μg/ml OMVs for 2 h. Cells were

then supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells were fixed

and stained with antibodies against Myc or caspase-11 16 h post infection. Nucleus was

stained with DAPI. Scale bar, 5 μm. (F) *Casp11*$^{+/+}$ and *Casp11*$^{-/-}$ BMDM cells were primed

overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS using DOTAP

with or without the incubation of VgrG2b-Myc knockin ΔRetS PAO1 at an MOI of 30 for 2 h.

Cells were then supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells

were lysed and immunoblotted 16 h post infection. (G) *Casp11*$^{-/-}$ BMDM cells were

infected with lentiviruses encoding a control vector of caspase-11. Cells were primed

overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS using DOTAP

with or without the incubation of VgrG2b-Myc knockin ΔRetS PAO1 at an MOI of 30 for 2 h.

Cells were then supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells

were lysed and immunoblotted as indicated 16 h post infection. (H) *Casp11*$^{-/-}$ BMDM cells

were infected with lentiviruses encoding wild-type or mutant caspase-11. Cells were

primed overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS using

DOTAP with or without the incubation of VgrG2b-Myc knockin ΔRetS PAO1 at an MOI of

30 for 2 h. Cells were then supplemented with fresh medium containing 100 μg/ml

Gentamycin. Cells were lysed and immunoblotted as indicated 16 h post infection. (I)

Human macrophages were primed overnight with 1 μg/ml Pam3CSK4, followed by

transfection of 2 μg/ml LPS using DOTAP with or without the incubation of VgrG2b-Myc

knockin ΔRetS PAO1 at an MOI of 30 for 2 h. Cells were then supplemented with fresh

medium containing 100 μg/ml Gentamycin. Cells were lysed and immunoblotted as

indicated 16 h post infection. (J–M) Wild-type BMDM cells were primed overnight with 1

μg/ml Pam3CSK4, followed by incubation of VgrG2b-Myc knockin (KI) or knockout (KO)

ΔRetS PAO1 at an MOI of 30 and 20 μg/ml OMVs for 2 h. Cells were then supplemented

with fresh medium containing 100 μg/ml Gentamycin. Cells were lysed and

immunoblotted as indicated 16 h post infection (J). Cell culture supernatants were

collected for an ELISA assay to determine the secreted IL-1β protein levels 16 h post

infection (K). Cytotoxicity was determined by LDH release assay in cell culture

supernatants 16 h post infection (L). Cell viability was determined by an ATP quantification

assay in cell pellets 16 h post infection (M). Data were shown as means±SD. Experiments

were repeated three times with similar results.

Figure 3. Caspase-11 cleaves VgrG2b at D883. (A) Plasmids encoding VgrG2b and

mutant caspase-11 p22/p10 were co-transfected into HEK293T cells for 24 h, followed by

immunoprecipitation with a control IgG or antibody against Myc. Precipitates were

immunoblotted as indicated. (B) Plasmids encoding caspase-11 p22/p10 and VgrG2b

mutants were co-transfected into HEK293T cells for 24 h, followed by immunoblotting with

antibodies against the indicated proteins. (C) Plasmids encoding caspase-4 p22/p10 and

VgrG2b variants were co-transfected into HEK293T cells for 24 h, followed by

immunoblotting with antibodies against the indicated proteins. (D) Plasmids encoding

VgrG2a-N-VgrG2b-C chimera and caspase-11 p22/p10 were co-transfected into

HEK293T cells for 24 h, followed by immunoblotting with antibodies against the indicated

proteins. (E) Plasmids encoding VgrG2b C-terminus and mutant caspase-11 p22/p10

were co-transfected into HEK293T cells for 24 h, followed by immunoprecipitation with a

control IgG or antibody against Myc. Precipitates were immunoblotted as indicated. (F)

Recombinant VgrG2b were incubated with caspase-11 p22/p10 for 4 h, followed by

immunoblotting with antibodies against the indicated proteins. (G) Wild-type BMDM cells

were primed overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS

using DOTAP with or without the incubation of VgrG2b wild-type (KI) or mutant (D883A)

knockin ΔRetS PAO1 at an MOI of 30 for 2 h. Cells were then supplemented with fresh

medium containing 100 μg/ml Gentamycin. Cells were lysed and immunoblotted 16 h post

infection. (H–L) Wild-type BMDM cells were primed overnight with 1 μg/ml Pam3CSK4,

followed by transfection of 2 μg/ml LPS using DOTAP with or without the incubation of

VgrG2b wild-type (KI) or mutant (D883A) knockin ΔRetS PAO1 at an MOI of 30 for 2 h.

Cells were then supplemented with fresh medium containing 100 μg/ml Gentamycin. Cells

were lysed and immunoblotted as indicated 16 h post infection (H). Band intensities of

cleaved caspase-11 (top) and GSDMD (bottom) were quantified and compared to that of

β-actin (I). Cell culture supernatants were collected for an ELISA assay to determine the

secreted IL-1β protein levels 16 h post infection (J). Cytotoxicity was determined by LDH

release assay in cell culture supernatants 16 h post infection (K). Cell viability was

determined by an ATP quantification assay in cell pellets 16 h post infection (L). Data were

shown as means±SD. Experiments were repeated three times with similar results.

Figure 4. VgrG2b C-terminus interacts with NLRP3. (A) Yeast two-hybrid screening

was performed using VgrG2b C-terminus as bait and a mouse bone marrow cDNA library

was screened. The interaction strength between bait and preys were visualized by

X-α-gal assays. (B, C) Plasmids encoding VgrG2b C-terminus (B) or full-length (C) and

NLRP3 were co-transfected into HEK293T cells for 24 h, followed by immunoprecipitation

with a control IgG or antibody against Myc. Precipitates were immunoblotted as indicated.

(D) VgrG2b C-terminus proteins were introduced into BMDM cells with the help of

cell-penetrating peptides (CPP) for 6 h. Cells were lysed and immunoprecipitated with a

control IgG or antibody against Myc. Precipitates were immunoblotted as indicated. (E)

BMDM cells were primed overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2

μg/ml LPS using DOTAP with the incubation of VgrG2b knockin (KI) ΔRetS PAO1 at an

MOI of 30 for 2 h. Cells were then supplemented with fresh medium containing 100 μg/ml

Gentamycin. Cells were lysed and immunoprecipitated with a control IgG or antibody

against NLRP3 16 h post infection. Precipitates were immunoblotted as indicated. (F)

BMDM cells were primed overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2

μg/ml LPS using DOTAP with the incubation of VgrG2b knockin (KI) ΔRetS PAO1 at an

MOI of 30 for 2 h. Cells were then supplemented with fresh medium containing 100 μg/ml

Gentamycin. Cells were fixed and stained with antibodies against Myc and NLRP3 16 h

post infection. Nucleus was stained with DAPI. Scale bar, 5 μm. (G) Scheme for NLRP3

truncations. (H, I) Plasmids encoding VgrG2b C-terminus and NLRP3 truncations were

co-transfected into HEK293T cells for 24 h, followed by immunoprecipitation with a control

IgG or antibody against FLAG. Precipitates were immunoblotted as indicated. (J)

Casp11[+/+] and *Casp11*[−/−] BMDM cells were primed overnight with 1 μg/ml Pam3CSK4,

followed by transfection of 2 μg/ml LPS using DOTAP with the incubation of VgrG2b-Myc

knockin (KI) ΔRetS PAO1 at an MOI of 30 for 2 h. Cells were then supplemented with

fresh   medium   containing   100 μg/ml   Gentamycin.   Cells   were   lysed   and

immunoprecipitated with a control IgG or antibody against NLRP3 16 h post infection.

Precipitates were immunoblotted as indicated. Data were shown as means±SD.

Experiments were repeated three times with similar results.

Figure 5. VgrG2b C-terminus suppresses the NLRP3 inflammasome. (A) Plasmids

encoding VgrG2b C-terminus and the indicated murine proteins were co-transfected into

HEK293T cells for 24 h, followed by immunoblotting with antibodies against the indicated

proteins. (B) Wild-type BMDM cells were infected with lentiviruses encoding a control

vector or VgrG2b C-terminus and primed with 1 μg/ml LPS for 3 h, followed by stimulation

of 10 μM nigericin for 30 min, 0.5 μM gramicidin for 1 h and 2 mM ATP for 30 min. Cells

were lysed and immunoblotted as indicated. (C–G) Wild-type BMDM cells were primed

overnight with 1 μg/ml Pam3CSK4, followed by transfection of 2 μg/ml LPS using DOTAP

with or without the transfection of VgrG2b C-terminus proteins using cell-penetrating

peptides (CPP) for 2 h. Cells were then supplemented with fresh medium. Cells were

lysed and immunoblotted 16 h post transfection (C). Band intensities of cleaved

caspase-11 (top) and GSDMD (bottom) were quantified and compared to that of β-actin

(D). Cell culture supernatants were collected for an ELISA assay to determine the

secreted IL-1β protein levels 16 h post infection (E). Cytotoxicity was determined by LDH

release assay in cell culture supernatants 16 h post infection (F). Cell viability was

determined by an ATP quantification assay in cell pellets 16 h post infection (G). (H, I)

Wild-type BMDM cells were primed overnight with 1 μg/ml Pam3CSK4, followed by

transfection of 2 μg/ml LPS using DOTAP with or without the transfection of VgrG2b

C-terminus proteins using cell-penetrating peptides (CPP) for 2 h. Cells were then

supplemented with fresh medium. Cells were lysed and immunoprecipitated with a control

IgG or antibody against NLRP3 16 h post transfection. Precipitates were immunoblotted

with antibody against Nur77 (H) or NEK7 (I). (J) Plasmids encoding VgrG2b C-terminus,

NEK7 and NLRP3 were co-transfected into HEK293T cells for 24 h, followed by

immunoprecipitation with a control IgG or antibody against NLRP3. Precipitates were

immunoblotted as indicated. Data were shown as means±SD. Experiments were repeated

three times with similar results.

Figure 6. *P. aeruginosa* suppresses NLRP3 inflammasome *in vivo*. (A–E) Wild-type

mice were intraperitoneally challenged with poly(I:C) at a dose of 2 mg/kg body weight for

6 h and then intranasally infected with 1×10^9 cfu of VgrG2b (KI) or mutant (D883A)

knockin ΔRetS PAO1, followed by survival calculation at the indicated days (A). Serum

protein levels of IL-1β (B), TNF-α (C) and IL-6 (D) were examined through ELISA assays

24 h post infection. Bacterial load was examined in mouse lung tissues 4 days post

infection (E). (F–J) Wild-type mice were intraperitoneally challenged with poly(I:C) at a

dose of 2 mg/kg body weight for 6 h and then intranasally infected with 1×10^9 cfu of

VgrG2b (KI) or mutant (D883A) knockin ΔRetS PAO1. Bronchoalveolar lavage fluids

(BALF) were collected 16 h post infection. Cells were immunoblotted as indicated (F). Cell

viability was also determined by an ATP quantification assay in cell pellets (G). Protein

levels of IL-1β (H), TNF-α (I) and IL-6 (J) in BALF supernatants were examined through

ELISA assays. (K–O) Wild-type mice were intraperitoneally challenged with poly(I:C) at a

dose of 2 mg/kg body weight for 6 h and then intranasally infected with 1×10^9 cfu of ΔRetS

PAO1 and administrated with 20 μg TAT sequence containing VgrG2b C-terminus peptide,

followed by survival calculation at the indicated days (K). Serum protein levels of IL-1β (L),

TNF-α (M) and IL-6 (N) were examined through ELISA assays 24 h post infection.

Bacterial load was examined in mouse lung tissues 4 days post infection (O). Data were

shown as means±SD. Experiments were repeated three times with similar results.

Figure 1