Identifying Genetic Variations in *emm*89 *Streptococcus pyogenes* Associated with Severe Invasive Infections

Masayuki Ono[1,2], Masaya Yamaguchi[1,2,3,4,5,20]*, Daisuke Motooka[3,6,7], Yujiro Hirose[1],

Kotaro Higashi[1], Tomoko Sumitomo[1,8], Tohru Miyoshi-Akiyama[9], Rumi Okuno[10],

Takahiro Yamaguchi[11], Ryuji Kawahara[11], Hitoshi Otsuka[12], Noriko Nakanishi[13], Yu

Kazawa[14], Chikara Nakagawa[15], Ryo Yamaguchi[16], Hiroo Sakai[17], Yuko Matsumoto[18],

Tadayoshi Ikebe[19], and Shigetada Kawabata[1,4]

1. Department of Microbiology, Graduate School of Dentistry, The University of Osaka,

Osaka, Japan

2. Bioinformatics Research Unit, Graduate School of Dentistry, The University of

Osaka, Osaka, Japan

3. Bioinformatics Center, Research Institute for Microbial Diseases, The University of

Osaka, Osaka, Japan

4. Center for Infectious Diseases Education and Research, The University of Osaka,

Osaka, Japan

5. Laboratory of Microbial Informatics, Microbial Research Center for Health and

Medicine, National Institutes of Biomedical Innovation, Health and Nutrition, Osaka,

Japan

6.   NGS Core Facility, Research Institute for Microbial Diseases, The University of

Osaka, Osaka, Japan

7.   Integrated Frontier Research for Medical Science Division, Institute for Open and

Transdisciplinary Research Initiatives (OTRI), The University of Osaka, Osaka, Japan

8.   Department of Oral Microbiology, Graduate School of Biomedical Sciences,

Tokushima University, Tokushima, Japan

9.   Pathogenic Microbe Laboratory, Department of Infectious Diseases, National Center

for Global Health and Medicine, Tokyo, Japan

10.   Department of Microbiology, Tokyo Metropolitan Institute of Public Health, Tokyo,

Japan

11.   Department of Bacteriology, Osaka Institute of Public Health, Osaka, Japan

12.   Department of Public Health Sciences, Yamaguchi Prefectural Institute of Public

Health and Environment, Yamaguchi, Japan

13.   Department of Infectious Diseases, Kobe Institute of Health, Hyogo, Japan

14.   Department of Microbiology, Fukushima Prefectural Institute for Public Health,

Fukushima, Japan

15. Division of Microbiology, Kyoto City Institute of Health and Environmental

Sciences, Kyoto, Japan

16. Sapporo Public Health Office, Hokkaido, Japan

17. Niigata City Institute of Public Health and the Environment, Niigata, Japan

18. Microbiological Testing and Research Division, Yokohama City Institute of Public

Health, Kanagawa, Japan

19. Department of Bacteriology I, National Institute of Infectious Diseases, Tokyo,

Japan

20. Lead contact.

*Correspondence: yamaguchi.masaya@nibn.go.jp

Abstract

Streptococcus pyogenes causes mild human infections as well as life-threatening invasive

diseases. Since the mutations known to enhance virulence to date account for only half of

the severe invasive infections, additional mechanisms/mutations need to be identified.

Here, we conducted a genome-wide association study of *emm*89 *S. pyogenes* strains to

comprehensively identify pathology-related bacterial genetic factors (SNPs, indels, genes,

or k-mers). Japanese (n=311) and global (n=666) cohort studies of strains isolated from

invasive or non-invasive infections revealed 17 and 1,075 SNPs/indels and 2 and 169

genes, respectively, that displayed associations with invasiveness. We validated one of

them, a non-invasiveness-related point mutation, *fhuB* T218C, by structure predictions

and introducing it into a severe invasive strain and confirmed that the mutant showed

slower growth in human blood. Thus, we report novel mechanisms that convert *emm*89

S. pyogenes to an invasive phenotype and a platform for establishing novel treatments

and prevention strategies.

Introduction

Streptococcus pyogenes is a human-restricted gram-positive pathogen associated with a

wide spectrum of diseases. While *S. pyogenes* often causes non-invasive diseases,

including pharyngitis and impetigo in children, it is also known as a "flesh-eating

bacterium" owing to its involvement in life-threatening invasive diseases, such as

necrotizing fasciitis and streptococcal toxic shock syndrome (STSS)[1,2]. In 2005, more

than 0.6 million people were estimated to have invasive *S. pyogenes* infections[3], and the

reported incidence of invasive *S. pyogenes* infections continues to increase globally[1]. In

cases of severe infection, rapid bacterial growth and profound metabolic acidosis

necessitate urgent surgical inspection and extended debridement with empiric

antibacterial chemotherapy[4,5]. However, even with proper treatment, the mortality rate of

patients with STSS remains high, ranging from 23–81%[6]. Moreover, although several

protective vaccine candidates against *S. pyogenes* exist, no safe and effective commercial

vaccine has yet been licensed for human use[2,7].

S. pyogenes has been classified into at least 240 *emm* types based on a

hypervariable region sequence of the *emm* gene, which encodes the M protein. This

hypervariable region of the M protein is responsible for type-specific antigenicity and

binds with high affinity to C4b-binding protein, a major fluid phase inhibitor of the

classical and lectin pathways of the complement system that confers resistance to

opsonophagocytosis[8]. Since the mid to late 2000s, *emm*89 strains have been increasingly

isolated from samples obtained from patients with invasive diseases, becoming one of the

most frequently identified lineages in developed countries[9,10]. For example, in the United

Kingdom, no more than ten cases of invasive diseases caused by *emm*89 *S. pyogenes* were

reported annually in the 1990s. In the 2010s, however, approximately 150 cases were

reported out of the 1,000–1,500 invasive cases caused by all *emm* types of *S. pyogenes* in

each year[11]. Similarly, in Japan, in the 2010s, there was an increase in the number of

patients with *emm*89 *S. pyogenes*-induced severe invasive infections. In 2018, *emm*89

strains were isolated from 36% of patients diagnosed with *S. pyogenes*-induced severe

invasive infections, following *emm*1 strains, which were isolated from 57% of patients[9].

*S. pyogenes emm*89 strains have been genetically sub-clustered into three clades

according to the *nga* promoter region patterns and the presence/absence of the *hasABC*

locus, which is responsible for hyaluronan capsule synthesis. Clade 3 is distinct from

clades 1 and 2 in terms of two features: overexpression of virulence factors NAD

glycohydrolase (NADase) and streptolysin O (SLO), owing to mutations in the promoter

region of the *nga-ifs-slo* operon, and the lack of a hyaluronan capsule[11-13]. Although clade

3 strains have frequently been isolated from invasive diseases, their numbers from non-

invasive infections have also increased. We previously reported no difference in the

isolation frequencies of clade 3 strains between invasive and non-invasive diseases, at

least in Japan, and concluded that the mutations in clade 3 are not responsible for the gain

of invasiveness[14]. Therefore, there must be other genetic features within the *emm*89

strains that determine their phenotypes.

During infections caused by *emm*1 *S. pyogenes*, mutations in the two-component

system, CovR/S, promote high virulence[15]. These mutations cause the upregulation of

DNase, hyaluronan capsule, IL-8 protease, C5a peptidase, streptokinase, NADase, SLO,

and superantigen SpeA, as well as the downregulation of cysteine protease SpeB and

streptolysin S[16,17]. The resulting mutants can prevent neutrophil death and subsequently

promote tissue destruction and systemic infections. Epidemiologically, Ikebe *et al.*

reported that nonsense mutations in *covR* and/or *covS* are present in 46.3% of *S. pyogenes*

strains isolated from severe invasive infections in Japan, but only 1.69% of isolates from

non-severe ones[18]. Moreover, these studies indicated that the *covR*/*S* mutation is not

responsible for all invasive clinical strains. Thus, we hypothesize that other mechanisms

are involved in the development of invasiveness.

In the present study, we aimed to explore novel hypervirulent mechanisms of *S.*

pyogenes by performing a pan-genome-wide association study (pan-GWAS) on *S.*

pyogenes. Our pan-GWAS focused on the *emm*89 lineage of *S. pyogenes* to detect lineage-

specific factors and minimize false positives due to lineage differentiation. For a

comprehensive analysis, we constructed the core genome and pan-genome of *emm*89 *S.*

pyogenes strains and evaluated the effect of single-nucleotide polymorphisms (SNPs) on

core gene alignment and accessory clusters of orthologous genes (COGs). In addition, we

performed a k-mers-based GWAS to detect SNPs in the intergenic regions and multiple

mutations. We collected and sequenced *emm*89 clinical strains isolated in Japan during

2016–2021, in addition to public *emm*89 genome sequences. Using these sequences, we

investigated the bacterial factors associated with severe invasive infections in Japan and

globally, using pan-GWAS. Based on the bacterial protein structural predictions, we then

selected candidates with high potential relevance to the phenotype. Finally, we introduced

an SNP related to non-invasiveness into a clinical strain isolated from a severe invasive

infection and examined the alteration of the bacterial phenotype through an *ex vivo*

infection assay.

Results

Collection of *emm*89 *S. pyogenes* clinical isolates in Japan and construction of

cohorts

We collected clinical *S. pyogenes* strains isolated between 2016 and 2021 from patients

with non-invasive and severe invasive infections in Japan. The Ministry of Health, Labour,

and Welfare of Japan has defined the clinical criteria of severe invasive β-hemolytic

streptococcal infections as STSS, based on the STSS 2010 Case Definition of the Centers

for Disease Control and Prevention in the US, with minor modifications, including the

addition of encompassing symptoms in the central nervous system (Table S1)[19,20].

For the *emm*89 clinical isolates, we collected T serotype TB3264 and untypable

strains, in addition to *emm* genotype-identified strains. T-typing is a serologically-based

approach that is often used as an alternative or supplement to *emm* typing. T-antigens are

trypsin-resistant surface antigens exhibiting extensive antigenic diversity[21]. Isolates of a

given *emm* type frequently share the same T serotype pattern[22,23]. The T serotype TB3264

corresponds to the genotype *emm*89 or *emm*94[21,24]. A total of 207 clinical isolates were

collected with the cooperation of the National Institute of Infectious Diseases and ten

public health institutes nationwide (Tables S2 and S3). We performed draft genome

sequencing of the strains and identified their *emm* types. In total, 150 of these were

determined as *emm*89, followed by 24 and 19 strains as *emm*4 and *emm*12, respectively.

To focus on the pathogenic mechanisms underlying severe invasive infections in the

*emm*89 cohort, we used 150 *emm*89 strains for subsequent analyses (Figure 1A, Tables

S2, and S3). We previously determined the draft genome sequences of 161 *emm*89 strains

isolated in Japan between 2011 and 2019 and determined their phenotypes using the same

criteria (Table S2)[14]. We combined these two sets and finally considered a total of 311

*emm*89 strains, including 135 severe invasive and 176 non-invasive isolates, as the

Japanese cohort.

We also collected public genome sequences of *emm*89 *S. pyogenes* strains isolated

from nine countries to further characterize the genetic properties of the Japanese cohort

(Table S2)[25-27]. In this study, the phenotypes of these strains were considered invasive if

the diagnoses included severe infections, STSS, invasive infections, necrotizing fasciitis,

bacteremia, or sepsis, and isolation sites were described as normally sterile sites, such as

the blood, brain, kidney, muscular tissue, or brain. Consequently, we identified 666 strains

in the global cohort, including 420 isolates from invasive cases and 246 from non-

invasive ones (Table S2).

Pan-genome and phylogenetic analyses reveal both shared and distinct features in

the Japanese and global cohorts

To determine the core genes and gene distribution in both cohorts, we performed pan-

genome analyses. In the Japanese cohort, 1,417 core genes common to more than 99% of

all isolates were determined out of the 3,334 different genes detected within the 311

strains. In contrast, the global cohort was more diverse, with 4,743 different genes, of

which 1,327 were core genes (Figure 1B).

Next, we calculated the phylogenetic relationships based on the maximum

likelihood of the core gene sequences (Figure 1C and S2). The tree for the global cohort

branched into four clusters, with clusters A, B1, B2, and B3. Cluster B3 included 640

genetically similar strains isolated mainly from Europe, North America, and Japan,

whereas cluster A comprised 19 strains isolated from Oceanian countries, Kenya,

Lebanon, the US, and Japan (Figure 1C). The phylogenetic tree for the Japanese cohort

could also be clustered as in the case of the global cohort, with no significant difference

in the proportions of strains classified into each sub-cluster (chi-square test, p=0.13;

Figure S2 and Table S4). Thus, we concluded that the overall phylogenetic features of

*emm*89 strains were distributed similarly in Japan and other areas, especially Europe and

North America. Within cluster B3, we identified a non-invasive strain from Japan that had

no identical pattern to the reported *nga* promoter variations (Figure 1C)[28]. This pattern is

likely a subtype of clade 3 as it shares the haplotype $A_{-27}G_{-22}T_{-18}$, which is distinctive of

clade 3, but has a mutation in the –10 box (Figure S2)[28]. Thus, we named this novel *nga*

promoter variation type 3.4[28]. Multilocus sequence typing (MLST) analysis revealed that

in cluster B3, 522 strains (80.9%) were ST101, and 96 strains (14.8%) were ST646

(Figure 1C). Notably, ST646 was the second most dominant type and a Japan-specific

lineage. Moreover, they only differed in the 295[th] nucleotide in the *murI* locus, one of the

seven loci that determine MLST, suggesting that both lineages have a genetically close

relationship. Eight novel MLSTs were determined (ST1450, 1451, 1452, 1454, 1455,

1456, 1461, and 1463) and 15 novel MLST strains were detected in the Japanese cohort

(Figure S1). Taken together, using phylogenetic approaches, we found that most strains

from Japan and countries in Europe and North America share genetically close

relationships, with only one unique lineage in Japan, ST646.

Pan-GWASes detect SNPs/indels associated with invasiveness that are both common

and specific to Japan and other countries

To discover all types of genetic variants in whole genes within *emm*89 *S. pyogenes*

associated with (severe) invasiveness, we applied pan-genome analysis and performed

three types of independent analyses, including pan-GWASes targeting SNPs in core genes

and the presence or absence of all genes, and a GWAS targeting other variants located in

intergenic regions spanning several nucleotides. The present pan-GWAS of bacteria

within a single *emm* type minimized lineage effects, thus reducing false positives.

We extracted SNPs and single-nucleotide indels from core gene alignments and

detected 24,627 and 47,060 SNPs/indels in the Japanese and global cohorts, respectively.

Subsequent pan-GWASes identified SNPs/indels associated with severe invasiveness in

a Japanese cohort and invasiveness in a global cohort. To control for population bias, we

calculated pairwise distance matrices and selected seven and three dimensions for the

analyses of the Japanese and global cohorts, respectively (Figure S3A and S3B). For each

cohort, we performed a permutation test by conducting 1,000 iterations of calculations

with randomly permuted phenotypes, with the significance level set at the 5[th] percentile

of the 1,000 minimal *p*-values ($p=5.75\times10^{-4}$ and $p=5.75\times10^{-4}$ for the Japanese and global

cohorts, respectively). The pan-GWAS of the Japanese cohort detected 17 SNPs/indels in

13 core genes (Figure 2A and Table S5). Of the 17 significant variants, there were 7

single-nucleotide deletions (SNDs), 7 SNPs causing non-synonymous amino acid

substitutions, and 3 SNPs causing synonymous substitutions. The *covS* gene (also known

as *csrS*), encoding a sensor kinase of the two-component system CovR/S, contains four

SNDs with the lowest *p*-values ($p=1.16\times10^{-7}$ for the 39[th], 40[th], and 46[th] nucleotides, and

$p=1.15\times10^{-6}$ for the 125[th] nucleotide). These four deletions were associated with severe

invasive infections.

We also performed a pan-GWAS for the global cohort and detected 1,075

SNPs/indels significantly related to invasive infections among the 360 core genes (Figure

2B, 2C and Table S6). Among the significant SNPs/indels, 725 caused synonymous

substitutions and 319 caused non-synonymous substitutions or frameshift mutations.

Moreover, 19 SNPs induced nonsense mutations, whereas the effects of 12 SNPs/indels

were unpredictable because of a lack of reference sequences (Table S3). Notably, 96

SNPs/indels accumulated in a single gene, *murJ*, which is involved in peptidoglycan

biosynthesis, whereas 53 and 51 SNPs/indels were detected in *murE* and *group_1008*,

respectively (Figure S3C). The SNP with the lowest *p*-value ($p=1.35\times10^{-14}$) was *lacE,*

which encodes the EIICB component of the lactose-specific phosphotransferase system

(Figure 2C). This mutation was found to be associated with an invasive phenotype and

mainly observed in strains isolated in the US. Compared with the significant 17

SNPs/indels in the Japanese cohort, 10 SNPs/indels were also detected in the global

cohort, including 4 SNDs in *covS* and 1 SNP each in 6 loci (Figure 2C). Deletions at the

covS locus were common among strains from several countries, including Japan. In

contrast, SNPs in six loci, *gatA, group_1102, group_647, iscS_1, recU*, and *fhuB*, were

present exclusively in Japan (Figure 2C and Table S6). These results suggest that several

bacterial mechanisms cause severe invasive *S. pyogenes* infections, and some prevail

worldwide, such as *covS* mutations, whereas others are specific to Japan.

Pan-GWAS on COGs reveals 2 and 109 genes associated with severe invasiveness in

the Japanese cohort and invasiveness in the global cohort, respectively

Next, we examined the associations of accessory COGs with severe invasiveness

and global invasiveness in Japanese patients. A permutation test determined significance

levels as p-values of 1.09×10^{-4} and 7.72×10^{-5} for the Japanese and global cohorts,

respectively. Two significant genes were detected in the pan-GWAS for the Japanese

cohort: *group_184*, which encodes a hypothetical protein, and *divIC*, which encodes a

septum formation initiator protein ($p=8.81\times10^{-6}$ and $p=6.72\times10^{-6}$, respectively; Figure

3A and Table S7). Although analysis of the global cohort revealed the presence of 169

genes that were significantly related to invasiveness, no genes were identical or

homologous to the two genes detected in the Japanese cohort (Figure 3B, 3C, and Table

S8). Approximately 90% (152 of 169) of the significant COGs were found to be

associated with non-invasiveness. Among the 169 genes, 37 encoded phage-related genes

and 11 encoded mobile genetic elements (MGEs) such as transposase, integrase, and

recombinase. The gene with the lowest p-value was *group_829*, which encodes a

transposase and is related to the invasive phenotype (Figure 3C and Table S8). In addition,

group_2689, which encodes a multidrug efflux transporter permease, *rhaR_3*, which

encodes a transcriptional regulator involved in rhamnose metabolism, and *group_1829*,

which encodes a vitamin B12 import transporter permease, were also substantially

associated with the invasive phenotype. We identified several gene distribution patterns

associated with invasiveness, suggesting that multiple independent genetic factors cause

invasive infections (Figure 3C). On the other hand, the present pan-GWAS found no genes

encoding known virulence factors significantly associated with invasiveness, thus further

analysis of the relationships of detected distribution patterns with prophages and MGEs

was performed. For calculating the pairwise correlation of the presence of significant

COGs in the 666 strains, the COGs were clustered into eight coexisting groups, seven of

which contained phage- and/or MGEs-related genes (Figure S4). The largest group

comprised 65 genes including phage proteins, while the second largest with 42 genes was

found to be associated with non-invasive infections, and included *group_2689*,

group_1833, and *ermA1*, encoding TetR/AcrR family transcriptional regulator, multidrug

efflux system permease protein, and rRNA adenine N-6-methyltransferase, respectively.

Taken together, genes associated with invasiveness were found to encode mobile genetic

factors and transporters, whereas major virulence factors were not significantly associated

with invasiveness.

K-mers-based GWAS detects both distinctive and identical variants compared to the

SNPs- and COGs-based pan-GWASes

To detect SNPs/indels and multiple mutations in the entire genome, we extracted 31-nt-

length k-mers from whole genomes and performed a GWAS. The k-mers-based GWAS

can handle polymorphisms spanning more than one base, such as indels, inversions, and

translocations, in both the coding and non-coding regions.

In the Japanese cohort, the k-mers-based GWAS detected two regions containing

causative variants associated with severe invasiveness (Table S9). As shown in the de-

Bruijn graphs, overlapping k-mers were concatenated into a single node, and edges

represent variability among the sequences (Figure 4A–F). The set of connected nodes and

edges comprising each de-Bruijn graph is called a complex. Nodes were determined to

be significant if their q-value was less than 0.05. Significant nodes are shown in red and

blue, indicating an association with severe invasive or non-invasive infections,

respectively. The complex comprising the nodes with the lowest q-value ($q=1.49\times10^{-2}$)

was Comp_11 in the *covS* locus (Table S9). The causative mutations in *covS* were an SND

and 10 nucleotide polymorphisms (Figure 4A). This deletion resulted in a frameshift

mutation that shortened the length of CovS from 500 to 35 amino acids, leading to

increased invasiveness, as previously reported in other *emm* types[17]. Another complex

significantly associated with severe invasiveness is Comp_2 ($q=4.22\times10^{-2}$; Table S9).

Comp_2 is a highly variable region containing eight hypothetical protein-coding genes,

with high similarity within the first 75 bp. Significant k-mers were also mapped to the

first 26 bp of *group_184* and 20 bp upstream (Figure 4B). These findings suggest that

group_184 possibly contributes to severe invasiveness through not only its presence but

also by that of the upstream region.

Next, we analyzed the global cohort and identified mutations that were

significantly associated with invasiveness in five regions (Table S10). The mutation with

the lowest *q*-value ($q=1.90\times10^{-2}$) existed in Comp_7 and was identical to the SND in *covS*

present in the k-mers-based GWAS of the Japanese cohort. Thus, this SND is global, as

are the four SNDs detected in the SNPs/indels-based pan-GWAS.

Two significant k-mers were present in Comp_6, which were found to be an

intergenic region of 270 bp ($q=1.90\times10^{-2}$; Table S10 and Figure 4C). We speculated that

this region is related to regulation of gene expression. However, no promoter sequences

were identified by utilizing MLDSPP, BacPP, and BLAST, thus the significance of this

region remains to be clarified[29,30]. In Comp_24, with a high sequence variation containing

group_141, *group_142*, and *group_143*, which encode transposases, the presence of a

281-bp sequence consisting of several k-mers was significantly associated with the

invasive phenotype ($q=1.90\times10^{-2}$; Table S10 and Figure 4D). Furthermore, *group_141*

was also identified in the COGs-based pan-GWAS as a non-invasiveness-related gene

along with *group_139* and *group_467*, which encode transposase and uncharacterized

protein, respectively (Table S8 and Figure S4). Taken together, the absence of an MGE

containing *group_141*, and the presence of another MGE harboring *group_142* and

group_143 may result in an invasive phenotype. In Comp_10, n27458, the *sagG* locus

encoding the ATP-binding protein of the efflux transporter of SLS, was significantly

correlated with the non-invasive phenotype (q=2.40×10^{-2}; Table S10 and Figure 4E).

However, a significant SNP in n27458, *sagG* (A882G), was found to cause a synonymous

mutation. The other significant mutation was present in the *fhuB* locus, encoding a

putative ferrichrome transport system permease (q=2.40×10^{-2}; Table S10). This mutation

was identical to SNP T218C detected in the SNP/indels-based pan-GWAS (Figure 4F).

Moreover, this mutation changes the 73rd residue from valine to alanine in FhuB, which

is a putative ferrichrome transport system permease.

Therefore, the k-mer approach identified multiple variants, including the mutation

identified in the SNPs/indels-based pan-GWAS, *fhu*B SNP T218C. In addition, while the

mutation detected in *covS* differed from that detected in the SNPs/indels-based pan-

GWAS, both caused frameshift mutations.

AlphaFold-based prediction of the impact of the identified mutations on function

To assess the impact of mutations on protein function, we predicted the protein structure

using AlphaFold[31]. Here, we present structural predictions for three representative

proteins: LacE, whose mutation was observed mainly in invasive strains from the US;

CovS, whose invasion-related deletions prevailed worldwide; and FhuB, which carries a

prominent mutation in the Japanese cohort and is associated with non-invasiveness.

The invasive-related SNP in *lacE* substitutes the 554th glycine in LacE with valine.

LacE was predicted to be a membrane protein with nine transmembrane regions (Figure

S5A). LacE is the EIICB component of the lactose-specific phosphotransferase system,

and the EIIC and EIIB domains correspond to the transmembrane and intracellular

regions, respectively. As shown in magenta in the model, the 554th residue is predicted to

be in the intracellular EIIB domain (Figure S5B).

Next, we predicted a homodimerized CovS model using AlphaFold, as the CovS

of *S. pyogenes* forms homodimers[32]. SOSUI predicted that CovS has two transmembrane

regions (Figure 5A). Mutations detected in the SNP/indels- and k-mers-based GWASes

were predicted to shorten the CovS protein to 35 and 45 amino acids, respectively. As the

intracellular domain of CovS is in the C-terminal region and is involved in the

phosphorylation of the transcriptional regulator CovR, frameshift mutations leading to

CovS truncation would inactivate the protein, and thus, CovR function (Figure 5B).

As described above, the SNP T218C in the *fhuB* locus substitutes the 73rd valine of

FhuB with alanine. FhuB is a component of an ATP-binding cassette transporter system

that utilizes ferrichrome, which is one of the carriers of Fe^{3+}. FhuB is predicted to localize

to the cell membrane and form a channel with FhuG (Figure 5C). SOSUI suggested that

FhuB and FhuG are 9-transmembrane proteins. The FhuBG complex can bind to one

molecule of the extracellular ferrichrome-binding lipoprotein, FhuD, and two molecules

of the intracellular ATP-binding protein, FhuC. Therefore, we constructed a structural

model of the FhuBCCDG complex (Figure 5D), which implied that the 73rd residue of

FhuB exists in a region adjacent to FhuD. The mutation was predicted to induce formation

of a small indentation on the molecular surface, thus increasing the surface area accessible

to the solvent, and is considered to potentially affect the stability of the hydrophobic bond

between FhuB and FhuD, and thus ferrichrome transport (Figure 5E). The SNP G538A

in *fhuD* was Japan-specific, significantly related to severe invasiveness, and caused the

V180I mutation in FhuD (Figure S3C). The prediction suggested that the 180th residue is

in an α-helix distant from the active site or the interactive sites with FhuB. Both valine

and isoleucine are branched-chain amino acids, and the amino acid residue is located on

a rigid structure, the α-helix (Figure 5D). Thus, we believe that this mutation is less likely

to cause structural changes than the other non-synonymous mutations.

The *fhuB* T218C mutation inhibits the growth of a severe invasive strain in human

blood

Based on the pan-GWAS and GWAS results and predicted protein structures, we focused

on the SNP *fhuB* T218C. We constructed a mutant strain, in which the SNP *fhuB* T218C

was introduced, to further investigate its potential virulence. We selected the strain TK02

which carries the wild-type (WT) allele T218 in *fhuB* and was originally isolated from a

sample obtained from a patient with severe invasive infection in Japan. We used a several

times-passaged TK02 strain, TK02', as a WT strain and introduced the SNP *fhuB* T218C

into it *via* allelic exchange mutagenesis with a thermo-sensitive shuttle vector. We then

confirmed that there were no differences between the WT and *fhuB* T218C strains using

whole-genome resequencing.

To reveal the effects of the SNP on invasiveness, we performed a transcriptomic

analysis of the WT and *fhuB* T218C strains in THY broth and human blood. Principal

component analysis revealed that the differences in the overall transcriptional profiles

between the strains were more remarkable in blood than in THY (Figure 6A, 6B, and

Tables S11–14). Among the overlapping genes, we found that the expression of CovR-

regulating genes, including *speB*, *nga-ifs-slo* operon, and *sag* operon, was upregulated in

the blood, compared to that in THY, in both the strains (Figure 6C, Tables S12, and S13).

CovR has also been reported to regulate *sda1*, which plays a key role in invasive disease

progression in *emm*1 strains; however, *sda1* does not exist in the *emm*89 reference strain

MGAS27061 or WT strain[15]. In human blood, the mutant strain resulted in the

downregulation of *mga* and *emm* expression and upregulation of the expression of genes

encoding virulence factors, such as *speC*, *scpA/B*, *endoS*, *fba*, *ska*, and *sfbX*, compared to

those observed in the WT strain (Table S14). Although Mga regulates the expression of

surface and secreted molecules important for colonization and immune evasion[33], no

strong expression changes were observed in the Mga regulon, except in the *emm* gene.

Notably, *fhuB*, *fhuC*, and *fhuD* were upregulated in the *fhuB* mutant in both human blood

and THY (Tables S11 and S14). Moreover, the transcriptomic analysis revealed that both

WT and *fhuB* T218C strains displayed upregulation of the *speB, nga-ifs-slo* operon, and

sag operons in human blood. In addition, the *fhuB* T218C mutation caused the

upregulation of *fhuBCD* expression. To determine whether *fhuB* T218C mutation affects

iron transport, we measured the intracellular free ferric ion concentration in each strain

and observed no differences among the strains in either environment (Figure 7A).

Next, to investigate the effects of SNP on bacterial survival in human blood, we

performed a bactericidal assay using human blood. At 2 and 3 h after incubation, the *fhuB*

T218C mutant strain exhibited a significantly decreased survival rate than that of the WT

strain (Figure 7B). To further determine the blood components that the attenuated survival

of the mutant can be attributed to, we compared bacterial survival rates in erythrocytes,

polymorphonuclear cells, plasma, heat-inactivated plasma, and brain heart infusion broth

(Figure 7C–7G). Notably, after incubation with erythrocytes, polymorphonuclear cells,

or plasma, the *fhuB* T218C mutant strain exhibited a significantly lower survival index

than that of the WT strain, at 2 and 3 h after incubation, as observed in whole blood

(Figure 7C–7E and 7G). However, there were no significant differences between the

survival rates of the WT and mutant strains in heat-inactivated plasma, suggesting that

the mutant strain is susceptible to complement (Figure 7F). Taken together, the

polymorphism T218C in *fhuB* impaired the survival of severe invasive strains in human

blood through interactions with erythrocytes.

Discussion

The present study focused on *emm*89 *S. pyogenes*, known to cause increasing rates of

invasive infections worldwide, and also assessed differences between *emm*89 strains

causing invasive and non-invasive infections. By focusing on bacteria within closed

phylogenies, false positives caused by confounding lineage effects were minimized, thus

contributing to a higher level of accuracy of the pan-GWAS. We constructed a workflow

for bacterial pan-GWAS and GWAS and explored the bacterial genetic factors related to

severe or invasive infections to reveal the plausible mechanisms of pathogenesis. We

independently performed pan-GWASes and GWASes for Japanese and global cohorts to

investigate the associations of variants with strictly defined phenotypes in severe invasive

S. pyogenes infections and with phenotypes in a broader context of invasive infections.

Several GWASes have been conducted on *S. pyogenes* to date. Davies *et al.* extracted k-

mers from the genomes of 1,944 strains of *S. pyogenes* with broad *emm* genotypes and

assessed their associations with invasiveness, and Kachroo *et al.* performed GWAS on k-

mers using 442 *emm*28 strains isolated from either invasive or non-invasive infections[27,34].

However, no studies have described the combined analysis of SNPs, genes, and k-mers

exclusively using *emm*89 strains that share genetically close genomes and, thus, a large

number of core genes. The investigations we performed led to the discovery of a

tremendous number of causative variants, including not only SNPs in core genes but also

accessory genes and insertions spanning intergenic regions.

Spontaneous mutations in *covR/S* genes potentiate the transition from localized to

invasive infection by M1T1 *S. pyogenes* [17]. We demonstrated that some SNDs in the *covS*

locus were significantly related to the phenotypes in both the SNPs/indels- and k-mers-

based GWASes. CovS forms a two-component system with CovR that regulates the

transcription of operons encoding virulence factors when phosphorylated by the histidine

sensor kinase CovS[35]. Sumby *et al.* suggested that systemic invasive infections are caused

by the overexpression of virulence factors through the inactivation of CovS, as a result of

the introduction of a depletive mutation in the *covS* locus after infection[15,16]. Several

mutation sites causing such overexpression, such as the 7 nt insertion in *covS* of *emm*1 *S.*

pyogenes and the insertion of adenine at the 877[th] nucleotide, have been reported[15,16].

Ikebe *et al.* demonstrated variations in the lengths of mutant CovS proteins by analyzing

the gene sequences of 164 STSS strains in Japan[18]. In line with their study, we found that

the shortening of CovS that potentially occurred owing to frameshifts caused by SNDs is

significantly associated with invasive infections, potentially contributing to the

upregulation of virulence factors and invasiveness.

In contrast, we observed several factors associated with invasiveness to be

independent of *covR*/*S* mutations. Based on conformational predictions, we selected SNPs

with non-synonymous substitutions that were likely to affect protein function.

Transcriptomic analysis findings suggested that the Japan-specific *fhuB* mutation

associated with non-severe invasive infections contributes to the growth rate of *S.*

pyogenes in human blood by adapting to the environment. In addition, *emm*89 clade 3

carries the identical promoter region pattern of the *nga* operon as *emm*1 strains, and the

pattern conferred similarly high expression of *nga* and *slo*[11,12,28]. Our RNA-seq data

demonstrated that a severe invasive Japanese strain without *covS* mutations increased the

expression of *speB*, *nga*, *slo*, and genes in the *sag* operon in the blood. Although *covR/S*

mutations downregulate the expression of SpeB and SLS, SpeB and SLS act as virulence

factors, allowing *S. pyogenes* to invade host tissues[36-39]. These data suggest that severe

invasive infections have multiple gene expression profiles in addition to *covR/S* mutation-

induced profiles within a single lineage, *emm*89, and that the synergy between optimizing

bacterial survival in human blood and upregulating multiple virulence factors contributes

to the development of severe invasive infections.

Two possible roles of the *FhuB* mutation in the pathogenesis of severe invasive

infections are thus proposed. First, the mutation in *FhuB* could increase bacterial

susceptibility to free radicals generated in the presence of ferric ions in the blood.

Generally*,* ferric ions are essential for the survival of almost all living organisms.

Catalyzed by ferric ions, the Fenton and Haber–Weiss reactions generate hydroxyl

radicals in erythrocytes[40]. Previously, we provided evidence that the iron in erythrocytes

partially inhibits pneumococcal growth *via* a free radical-based mechanism[7]. Although no

significant difference was observed in the intracellular ferric ion concentration in the

present study, the structural changes in FhuB may have caused an increased generation of

free radicals in bacterial cells and the prevention of bacterial survival in human blood.

Second, the *FhuB* mutation could affect bacterial transcriptional profiles and result in

reduced fitness for survival in human blood. We also observed a lower survival rate of the

mutant strain with polymorphonuclear cells (Figure 7D). Lower *emm* transcription in the

mutant may lead to attenuated immune evasion. The M protein can bind C4-binding

protein and factor H and inactivate the deposited C4b and C3b, leading to limited surface

opsonization[41]. Furthermore, complements could have inhibited the proliferation of the

mutant strain, as suggested by the activation of the membrane attack complex (Figure 7D

and 7E). Although the mechanisms of the *FhuB* V73A mutation in the interaction with

each blood component must be further elucidated, our results indicated that the 73rd

residue of FhuB is a key factor for bacterial survival in human blood during systemic

infection.

Recently, He *et al.* reported that antibodies against the ferrichrome-binding protein

FtsB, also known as FhuD, decrease blood bacterial burden and skin abscess formation

in murine models infected with *emm*1 *S. pyogenes*[42]. FhuD is a lipoprotein that has been

studied as a potential vaccine candidate against *Staphylococcus aureus*[43]. The detailed

function and effect of the FhuBCCDG complex in *S. pyogenes* remain unknown; however,

at least in *emm*1 strains, the complex can be expressed and localized abundantly on the

cell membrane. Hence, this complex may be a promising target for vaccines against *S.*

pyogenes.

Our gene-based pan-GWAS revealed no significant correlation with the

distribution of genes encoding virulence factors. This finding minimizes the possibility

that invasive infections are the result of the acquisition of virulence factors by non-

invasive strains and indirectly supports the hypothesis that changes in gene expression

profiles cause invasive infections. Although no vaccine has been commercialized against

S. pyogenes, given the possibility that the pathogen has multiple gene expression patterns,

even within a single lineage, it may be difficult to develop a universal vaccine with a

single antigen, and a vaccine containing multiple antigens may be effective.

In the global cohort, 169 genes were related to the phenotype, including 37 phage-

and 11 MGE-related genes. The genome of *S. pyogenes* is rich in prophages, phage-like

chromosomal islands, and MGEs[44]. Prophages and MGEs sometimes function as vectors

of virulence factors and antibiotic-resistance genes through interspecies and intraspecific

transmission. Common superantigen-coding genes, such as *speA*, *speC*, *speG*, *speH*, *speI*,

speJ, *speK*, and *speL*, and DNase-coding genes, such as *spd1*, are exogenous genes

derived from prophages[45]. In addition, in the genome of *S. pyogenes,* there are broad

MGEs where *erm* and *mefA* related to macrolide resistance and *tetM* involved in

resistance to tetracyclines are located[46]. In the present study, we observed no direct

association between invasiveness and virulence of genes located in the prophages and

MGEs. On the other hand, a cluster comprising 49 non-invasiveness-associated genes

including antibiotic-resistance genes was identified. Furthermore, among the genes

showing a significant correlation with the infectious phenotype, approximately 90% (152

of 169) were associated with non-invasiveness. One possible explanation is that

significantly related genes reflect the process of not only gain of factors but also loss of

those affecting fitness cost. Given that we also identified hypothetical genes associated

with this phenotype, we assumed that there is an unknown mechanism contributing to the

pathogenesis of invasive infections.

In the present study, MLST analysis revealed that 522 of the 666 strains belonged

to ST101. Following ST101, ST646 was the second most prevalent lineage in the

Japanese cohort and was not detected in any other country. In addition, the difference

between ST101 and ST646 was found only at the 295th nucleotide in *mulI*, among the

seven genes subjected to MLST typing. Our phylogenetic trees also suggested that ST101

and ST646 have a close phylogenetic relationship. Five previous studies have described

ST646 in *S. pyogenes*, and all of them reported *emm*89 ST646 strains isolated in Japan[47-

51]. Furthermore, all ST646 strains were isolated from the 4,249 *emm*89 *S. pyogenes* strains

in the MLST database PubMLST[52]. Therefore, ST646 appears to be specific to *emm*89

strains and a unique lineage in Japan. Notably, Ubukata *et al.* also reported that ST646

strains began increasing after 2012, whereas the ST101 lineage was dominant until 2008[51].

Taken together, we conclude that ST646 is possibly a relatively new and Japan-specific

lineage of *emm*89 *S. pyogenes.*

Our study had three limitations. First, because we sequenced bacterial genomes

using short-read sequencing, we could not detect large bacterial genome rearrangements

by comparing complete genome sequences. Thus, we were unable to investigate the

effects of long genomic structural dynamics, such as inversions, on pathogenesis. Second,

the clinical information associated with the isolates was limited; therefore, our analyses

did not reflect host information, such as age, sex, and underlying health conditions. Third,

we only used bacterial genetic distances based on core genome sequences as covariates

in the pan-GWAS. A combined GWAS of the host and pathogen in *S. pyogenes* infection

would highlight the relationship between host risk factors and bacterial genetic variants.

In this study, we revealed the genotype-phenotype associations found in not only

the known factors represented by *covS* but also factors that are related to invasiveness,

including *fhuB*. Moreover, we experimentally validated the contribution of the *fhuB*

mutation to bacterial survival in human blood. This study demonstrates the potential of

our genomic statistical approach for elucidating the pathogenesis of invasive infections.

Further analyses of the invasiveness-related factors identified in this study could provide

a platform for establishing novel treatments and preventive strategies against invasive

infections.

Materials and methods

Clinical isolates in Japan

Clinical isolates were collected from public health institutions in Tokyo, Osaka,

Yamaguchi, Fukushima, Kobe, Kyoto, Amagasaki, Sapporo, and Niigata, Japan. We

defined the strains collected as STSS according to the Infectious Diseases Control Law

in Japan. Non-invasive strains were defined based on diagnostic names, including

asymptomatic, pharyngitis, tonsillitis, or non-invasive infections. Strains with no

diagnostic names for the non-STSS strains were defined as non-invasive based on the

isolate sites. Information on all the strains included in this study is presented in Table S3.

The collected *S. pyogenes* strains were cultured at 37ºC in an atmosphere

containing 5% CO_2, in Todd Hewitt broth supplemented with 0.2% yeast extract (THY;

both from BD Biosciences, Franklin Lakes, NJ, USA) and stored in THY broth with 30%

glycerol (Nacalai Tesque, Kyoto, Japan), at –80°C.

Genomic DNA sequencing of the clinical isolates

The *S. pyogenes* strains were cultured until the exponential growth phase (OD_{600}=0.3–

0.4). Bacterial cells were lysed with T10E1N100 buffer (10 mM Tris-HCl buffer, 100

mM sodium chloride, and 1 mM EDTA), 10 units/mL mutanolysin (Sigma-Aldrich, St.

Louis, MO, USA), 10 mg/mL lysozyme (Fujifilm Wako Pure Chemical Co., Osaka,

Japan), 0.5 mg/mL achromopeptidase (Fujifilm Wako Pure Chemical Co.), and 0.3

mg/mL RNase A (Promega, Madison, WI, USA). Next, genomic DNA was extracted

from each lysate using a Maxwell® RSC instrument (Promega), according to the

manufacturer's instructions, and 250 bp paired-end libraries were then generated from the

extracted DNA using a Nextera XT DNA Kit (Illumina, San Diego, CA, USA). Libraries

were sequenced using a NovaSeq 6000 system (Illumina) at the Genome Information

Research Center, Research Institute for Microbial Diseases, The University of Osaka,

Osaka, Japan. On average, the number of reads was 5,433,301 (range 3,437,124–

9,117,301).

Collection of published genome sequences

We previously sequenced the draft genomes of 161 *emm*89 clinical isolates collected in

Japan between 2011 and 2019[14]. We defined strains derived from STSS as "severe

invasive," and those obtained from pharyngitis, tonsillitis, and superficial skin lesions as

"non-invasive" phenotypes.

To obtain public genome sequences of *emm*89 strains isolated from other countries,

we downloaded draft genome sequences in FASTA format from the National Center for

Biotechnology Information (NCBI) database, using Fasterq-dump v.2.9.6. The phenotype

of each strain, whether invasive or non-invasive, was defined according to the definitions

in the respective references that reported the strains[25-27].

Genomic data processing and pan-genome analysis

All processes and analyses were performed using the National Institute of Genetics (NIG)

supercomputer and SQUID at the D3 center of The University of Osaka (Osaka, Japan).

We constructed a workflow for bacterial pan-GWAS and other bioinformatic processes

(Figure S6). All collected sequences were subjected to quality checks using Fastp v.0.20.1,

with a default cutoff value of Q>15[53]. For newly isolated strains in Japan, *emm* typing

was performed using the emm-typing-tool v.0.0.1, and only sequences of strains

determined as *emm*89 were used for the following analyses[54]. All *emm*89 sequence data

were then subjected to *de novo* assembly using SKESA v.2.4.0, with default parameters[55].

Next, the MLST of each sequence was performed using MLST v.2.19.0[56,57]. Clade typing

was performed using BLAST v.2.13.0, with reference to the three *nga* promoter region

sequences[13]. After the genes were annotated with Prokka v.1.14.5, the pan-genome of all

sequences was calculated using Roary v.3.12.0, with the parameters "-e -mafft -r -qc -cd

99"[58,59]. Roary generated a core gene alignment and the distribution of all genes among

the strains. To extract SNPs/indels from core genes, including single-nucleotide indels,

snp-sites v.2.5.1 with the option "-v" was used. The output files were further processed

using BCFtools v.1.9, with the parameter "norm -m –", enabling analysis of multiple

alleles in the pan-GWAS. In parallel, k-mers were extracted using DBGWAS v.0.5.3, and

the length of k-mers was set as 31 nt using the "-k 31" parameter of DBGWAS[60].

Phylogenetic analysis

Phylogenetic relationships were calculated from the core gene alignment, using IQ-tree

v.1.16.12[61], based on maximum likelihood. The substitution model was automatically

selected considering the Akaike's and Bayesian information criteria by setting the "-m

MFP" parameter of IQ-tree[62]. Phylogenetic trees were constructed using iTOL v.6.6[63].

The similarity of clustering in the two phylogenetic trees was statistically examined using

Pearson's chi-square test with R v.4.0.3[64], followed by *post-hoc* analysis using residual

analysis adjusted with the Holm's method, if $p<0.05$.

GWAS

To investigate the associations between phenotypes and genotypes, including SNPs/indels

and genes, we performed a pan-GWAS using pyseer v.1.3.4[65]. The VCF file of

SNPs/indels or the gene distribution matrix was designated as the genotype. To remove

biases derived from lineages, we added information on the genetic distances between all

pairs of strains as covariates using mash v.2.3[66]. Briefly, *de novo* assemblies were

compressed through conversion into minimum hash values using the command "mash

sketch -s 10000." Subsequently, the commands "mash dist" and "square_mash" were

utilized to generate a genetic distance matrix expressed with Jaccard coefficients[66]. The

obtained matrix underwent eigenvalue decomposition, and the number of eigenvalues

used for multidimensional scaling was visually determined using the plot of the

relationships between the eigenvalues and contribution ratios. The number of eigenvalues

and the distance matrix were then added as pyseer parameters. The pyseer calculation was

iterated 1,000 times with randomized phenotypes, and the 5-percentile value of the

minimal p-value in each calculation was set at the significance level. Using R and the

package ggplot2, the results were visualized as a Manhattan plot for the SNPs/indels-

based pan-GWAS and a volcano plot for the genes-based pan-GWAS, respectively[67].

Heatmaps of the strains possessing significant variants were generated using Excel

(v.16.66.1; Microsoft, Redmond, WA, USA). The correlation of the presence of

significant COGs was calculated and visualized using the R program.

The k-mers-based GWAS was carried out using DBGWAS[60]. K-mers were

considered significant at a false discovery rate (q-value) of <0.05. DBGWAS-calculated

complexes, which are regions encompassing the k-mers, were significantly related to

pathology, and de-Bruijn graphs were generated based on these complexes. The

sequences of the k-mers were outputted and mapped to a reference sequence using

Geneious Prime v.2022.0.1 (Biomatters, Auckland, New Zealand) to identify the

mutations indicated by the k-mers. For the reference strain, we adopted MGAS27061,

which was isolated from an invasive case in the USA and whose complete chromosomal

sequence has been used as the reference sequence of *emm*89 clade 3[68]. Promoter

sequences in intergenic regions were predicted using web-based tools, MLDSPP and

BacPP [29,30]. Additionally, BLAST was employed to search the promoter sequences of *S.*

pyogenes strain SF370 registered in the CDBProm database

(https://aw.iimas.unam.mx/cdbprom/)[69].

Protein structure prediction

Significant variants found in the pan-GWAS, resulting in non-synonymous substitutions

in proteins, were searched by converting nucleotide sequences into amino acid sequences

using EMBOSS Transeq v.6.6.0.0[70]. To assess whether these mutations affected the

protein function, protein structure prediction models were constructed using AlphaFold

v.2.2.2[31]. The calculations were performed five times for each model, by setting the option

multimer_predictions_per_model=5. We predicted multimer models using the option --

model_preset="multimer" if a protein is reported or anticipated to form a multimer. For

each monomer, we selected the model with the best predicted local difference distance

test (an indicator of local structural accuracy) score[71]. For each multimer, AlphaFold was

calculated and expressed as a weighted combination of the interface-predicted TM and

predicted TM scores (ipTM + pTM). pTM is a metric for overall topological accuracy

and ipTM is used to measure the structural accuracy of the protein-protein interface[72].

The transmembrane regions of the proteins were predicted using SOSUI

(https://harrier.nagahama-i-bio.ac.jp/sosui/mobile/)[73]. The structures of the obtained

model were visualized using PyMOL v.2.5 (Schrödinger, LLC., New York, NY, USA).

Construction of the *fhuB* T218C mutant strains

We used the several times-passaged *S. pyogenes* TK02 strain, TK02', as the WT strain.

The TK02 strain was originally isolated from a patient with severe invasive infection[14].

The whole genome of TK02' was sequenced, and the generated fasta file is available in

Document S1. A point mutation, *fhuB* T218C, was introduced using the temperature-

sensitive shuttle vector, pSET4s, as reported previously[74]. Sanger sequencing confirmed

the presence of the point mutation. In addition, we resequenced the draft genome to

confirm that there were no differences apart from the point mutation, as described above.

The bacterial strains, primers, and plasmids used in this study are listed in Tables

S15 and S16. *Escherichia coli* strain DH5α (Takara Bio, Shiga, Japan) was used as a host

for the plasmid derivatives. All *E. coli* strains were cultured in Luria Bertani broth, at

37°C, with agitation. For selection and maintenance of mutants, antibiotics were added

to the media at the following concentrations: carbenicillin (Nacalai Tesque), 100 μg/mL

for *E. coli*; and spectinomycin (Fujifilm Wako Pure Chemical Corporation), 100 μg/mL

for *E. coli* and *S. pyogenes.*

Transcriptomic analysis

The *fhuB* WT and mutant strains were harvested in 30 mL of THY broth, of which 1 mL

was dispensed into 10 mL of THY and the remainder was centrifuged and resuspended in

2 mL of heparinized human blood. Bacterial mixtures with THY or blood were dispensed

into three aliquots (n=3 for each condition) and incubated at 37ºC for 3 h. THY samples

were centrifuged and resuspended in RNA Shield (Zymo Research, Irvine, CA, USA).

For each blood sample, 2 volumes (1 mL) of RNA protection bacteria reagent (Qiagen,

Hilden, Germany) were added. L5, included in the PureLink™ Total RNA Blood

Purification Kit (Thermo Fisher Scientific, Waltham, MA, USA), was added to remove

erythrocytes. The bacterial cell wall was mechanically lysed in Lysing Matrix B using a

MagNA Lyser (Roche, Basel, Switzerland). After centrifugation, the total bacterial RNA

was extracted using a Quick-RNA™ Miniprep Kit (Zymo Research), according to the

manufacturer's instructions. Full-length cDNA was generated using the SMART-Seq®

HT Kit (Takara Bio), according to the manufacturer's instructions. Pair-end libraries were

generated using a Nextera XT DNA Kit and sequenced using a NovaSeq 6000 system

(both from Illumina, San Diego, CA, USA). Sequenced data were preprocessed using

Trimmomatic v.0.33 and FastQC v.0.12.1. The reads were mapped to the complete

MGAS27061 genome (NCBI reference sequence: NZ_CP013840.1) using STAR

v.2.7.0a. After a second quality check using FastQC, read counting was performed using

featureCounts v.1.5.2[75]. Differentially expressed genes were identified using iDEP v.0.96

and gene annotations from NCBI and Prokka were combined[76]. Plots were created using

iDEP and the R package ggplot2.

Intracellular ferric ion assay

Human plasma was obtained through centrifugation of heparinized human blood, after 30

min of incubation at 37ºC. The WT and *fhuB* mutant strains were harvested at the

exponential phase, resuspended into 1 mL of THY or serum, and then incubated at 37ºC

for 3 h. Viable bacterial cells were counted as colony-forming units (CFUs) by plating the

diluted samples onto THY agar plates. Intracellular ferric ions were measured using a

QuantiChrom™ Iron Assay Kit (BioAssay Systems, Hayward, CA, USA), according to

the manufacturer's instructions. Briefly, 50 µL of standards or samples in 96-well plates

were mixed with 200 µL QuantiChrom™ Working Reagent and incubated at 20–24˚C for

an hour. The optical density at the wavelength of 600 nm was measured using an Infinite®

200 Pro F Plex Instrument (TECAN, Männedorf, Switzerland). The assay was repeated

three times, and the results of the respective experiments were combined. Statistical

analyses were performed using the Mann–Whitney U test.

Bactericidal assay

The bactericidal assay was performed as described previously, with minor

modifications[36,77-79]. Briefly, whole blood was collected from healthy adults. Human

neutrophils and erythrocytes were prepared using PolymorphPrep™ (Serumwerk

Bernburg, Bernburg, Germany), according to the manufacturer's instructions.

Heparinized human blood was centrifuged at $500 \times g$ for 30 min, to isolate erythrocytes

and polymorphonuclear cells, which were then suspended in Roswell Park Memorial

Institute (RPMI)-1640 medium containing L-glutamine and Phenol Red (Fujifilm Wako

Pure Chemical Corporation). Heat-inactivated plasma was prepared at 56ºC for 30 min.

Subsequently, 195 µL of heparinized human whole blood, erythrocytes in RPMI-1640,

polymorphonuclear leukocytes in RPMI-1640, plasma, heat-inactivated plasma, or brain

heart infusion broth (BD Biosciences), and 5 µL of early exponential phase bacteria with

$0.9–2.0 \times 10^4$ CFUs/well were mixed in 96-well plates and incubated at 37°C, in an

atmosphere containing 5% CO_2, for 1, 2, and 3 h. Viable bacterial cells were counted as

CFUs by plating the diluted samples onto THY agar plates. The growth index was

calculated as the number of CFUs at a specified time point divided by the number of

CFUs in the initial inoculum. The assay was repeated three times, and the results of the

respective experiments were combined. Statistical analyses were performed using the

Mann–Whitney U test. Differences were considered statistically significant at $p<0.05$,

using Prism v.7.0c (GraphPad, La Jolla, CA, USA).

Ethical approval

Studies involving human participants were reviewed and approved by the Institutional

Review Board of Osaka University Graduate School of Dentistry (approval nos. H26-E43

and H29-E16-2). The donors provided written informed consent to participate in the

human blood bactericidal assay. For the *S. pyogenes* collection, as we retrospectively

obtained clinical isolates of *S. pyogenes*, we utilized an opt-out consent procedure instead

of obtaining written informed consent from the patients.

Data availability

759   Data for the 207 sequenced *S. pyogenes* genomes were deposited in the DDBJ sequence

760   read archive, under BioProject PRJDB16457. The DRR run number is DRR511668-

761   DRR511874.

Acknowledgments

We would like to thank the NGS core facility of the Genome Information Research Center

at the Research Institute for Microbial Diseases of The University of Osaka for their

support in the DNA sequencing and data analysis and the Bioinformatic Research Unit of

Graduate School of Dentistry, The University of Osaka for their support in the

bioinformatics analysis. This study was partially performed on the National Institute of

Genetics (NIG) supercomputer at the Research Organization of Information and Systems

National Institute of Genetics. This study was partly completed using SQUID at the D3

center, The University of Osaka, Japan, under the "Joint Usage/Research Center for

Interdisciplinary Large-scale Information Infrastructures (JHPCN)" (Project ID:

EX22701, jh230035, jh240003, and jh250016). Masayuki Ono and Kotaro Higashi were

recipients of the Iwadare Scholarship from the Iwadare Scholarship Foundation. We wish

to express our gratitude to Mami Tateshita (Sapporo City Institute of Public Health) as

well as the medical institutions that participated in the collection of clinical strains.

This study was partly supported by AMED (JP17fk0108044, JP20fk0108130,

JP20wm0325001, and JP243fa727001h), the Japan Society for the Promotion of Science

KAKENHI (grant numbers 20KK0210, 22H03262, 22K19618, 22K19619, 23H03073,

23K19687, and 24K19854), the Takeda Science Foundation, Naito Foundation, and Joint

Research Program of the Research Center for GLOBAL and LOCAL Infectious Diseases,

Oita University (2022B05, 2025B01, and 2025B13). This study was conducted as part of

"The Nippon Foundation - Osaka University Project for Infectious Disease Prevention."

This study was supported by JST SPRING (grant number JPMJSP2138). The funders had

no role in the study design, data collection or analysis, decision to publish, or preparation

of the manuscript.

Author contributions

M.Y., M.O., and S.K. designed the study. M.O., M.Y., and K.H. designed and performed

bioinformatics analysis. D.M. performed the next-generation sequencing. M.O.

performed the in vitro and in vivo experiments. M.Y., Y.H., T.S., and S.K. contributed to

the setup of the experiments. R.O., T.Y., R.K., H.O., N.N., Y.K., C.N., R.Y., H.S., and

Y.M. collected clinical strains. M.O. and M.Y. wrote the original manuscript. All authors

reviewed and edited the manuscript.

Declaration of interests

797   The authors declare no competing interests.

References

1.   Craik, N., Hla, T., Cannon, J., Moore, H., Carapetis, J.R., and Sanyahumbi, A.

(2022). Global Disease Burden of *Streptococcus pyogenes*. In *Streptococcus*

pyogenes: Basic Biology to Clinical Manifestations [Internet] J.J. Ferretti, D.L.

Stevens, and V.A. Fischetti, eds. (University of Oklahoma Health Sciences

Center), Chapter 21.

2.   Brouwer, S., Rivera-Hernandez, T., Curren, B.F., Harbison-Price, N., De Oliveira,

D.M.P., Jespersen, M.G., Davies, M.R., and Walker, M.J. (2023). Pathogenesis,

epidemiology and control of Group A *Streptococcus* infection. Nat. Rev.

Microbiol. 21, 431–447. 10.1038/s41579-023-00865-7.

3.   Carapetis, J.R., Steer, A.C., Mulholland, E.K., and Weber, M. (2005). The global

burden of group A streptococcal diseases. Lancet Infect. Dis. 5, 685-694.

10.1016/S1473-3099(05)70267-X.

4.   Stevens, D.L., Bisno, A.L., Chambers, H.F., Dellinger, E.P., Goldstein, E.J.,

Gorbach, S.L., Hirschmann, J.V., Kaplan, S.L., Montoya, J.G., Wade, J.C., et al.

(2014). Practice guidelines for the diagnosis and management of skin and soft

tissue infections: 2014 update by the Infectious Diseases Society of America. Clin.

Infect. Dis. 59, e10–e52. 10.1093/cid/ciu444.

5.   Stevens, D.L., and Bryant, A.E. (2016). Severe Group A Streptococcal Infections.

In *Streptococcus pyogenes*: Basic Biology to Clinical Manifestations [internet],

J.J. Ferretti, D.L. Stevens, and V.A. Fischetti, eds. (University of Oklahoma

Health Sciences Center).

6.   Walker, M.J., Barnett, T.C., McArthur, J.D., Cole, J.N., Gillen, C.M., Henningham,

A., Sriprakash, K.S., Sanderson-Smith, M.L., and Nizet, V. (2014). Disease

manifestations and pathogenic mechanisms of Group A Streptococcus. Clin.

Microbiol. Rev. 27, 264–301. 10.1128/CMR.00101-13.

7.   Yamaguchi, M., Terao, Y., Mori-Yamaguchi, Y., Domon, H., Sakaue, Y., Yagi, T.,

Nishino, K., Yamaguchi, A., Nizet, V., and Kawabata, S. (2013). *Streptococcus*

pneumoniae invades erythrocytes and utilizes them to evade human innate

immunity. PLOS ONE 8, e77282. 10.1371/JOURNAL.PONE.0077282.

8.   Bessen, D.E., Smeesters, P.R., and Beall, B.W. (2018). Molecular Epidemiology,

Ecology, and Evolution of Group A Streptococci. Microbiol. Spectr. 6.

10.1128/microbiolspec.CPP3-0009-2018.

9.   Ikebe, T., Okuno, R., Kanda, Y., Sasaki, M., Yamaguchi, T., Otsuka, H., Kazawa,

Y., Suzuki, M., Ohya, H., Uchida, K., and Ohnishi, M. (2021). Molecular

characterization and antimicrobial resistance of group A streptococcus isolates in

streptococcal toxic shock syndrome cases in Japan from 2013 to 2018. Int. J. Med.

Microbiol. 311, 151496. 10.1016/j.ijmm.2021.151496.

10.   Fay, K., Onukwube, J., Chochua, S., Schaffner, W., Cieslak, P., Lynfield, R., Muse,

A., Smelser, C., Harrison, L.H., Farley, M., et al. (2021). Patterns of Antibiotic

Nonsusceptibility Among Invasive Group A *Streptococcus* Infections-United

States, 2006–2017. Clin. Infect. Dis. 73, 1957–1964. 10.1093/cid/ciab575.

11.   Turner, C.E., Abbott, J., Lamagni, T., Holden, M.T.G., David, S., Jones, M.D.,

Game, L., Efstratiou, A., and Sriskandan, S. (2015). Emergence of a new highly

successful acapsular group a Streptococcus clade of genotype *emm*89 in the

United Kingdom. mBio 6, e00622. 10.1128/mBio.00622-15.

12.   Zhu, L., Olsen, R.J., Nasser, W., Beres, S.B., Vuopio, J., Kristinsson, K.G.,

Gottfredsson, M., Porter, A.R., DeLeo, F.R., and Musser, J.M. (2015). A molecular

trigger for intercontinental epidemics of group A Streptococcus. J. Clin. Invest.

125, 3545–3559. 10.1172/JCI82478.

13.   Zhu, L., Olsen, R.J., Nasser, W., De La Riva Morales, I., and Musser, J.M. (2015).

Trading Capsule for Increased Cytotoxin Production: Contribution to Virulence

of a Newly Emerged Clade of *emm*89 *Streptococcus pyogenes*. mBio 6, e01378–

e01315. 10.1128/mbio.01378-15.

14.   Hirose, Y., Yamaguchi, M., Takemoto, N., Miyoshi-Akiyama, T., Sumitomo, T.,

Nakata, M., Ikebe, T., Hanada, T., Yamaguchi, T., Kawahara, R., et al. (2020).

Genetic Characterization of *Streptococcus pyogenes emm*89 Strains Isolated in

Japan From 2011 to 2019. Infect. Microbes Dis. 2, 160–166.

10.1097/IM9.0000000000000038.

15.   Walker, M.J., Hollands, A., Sanderson-Smith, M.L., Cole, J.N., Kirk, J.K.,

Henningham, A., McArthur, J.D., Dinkla, K., Aziz, R.K., Kansal, R.G., et al.

(2007). DNase Sda1 provides selection pressure for a switch to invasive group A

streptococcal infection. Nat. Med. 13, 981–985. 10.1038/nm1612.

16.   Sumby, P., Whitney, A.R., Graviss, E.A., DeLeo, F.R., and Musser, J.M. (2006).

Genome-wide analysis of group a streptococci reveals a mutation that modulates

global phenotype and disease specificity. PLoS Pathog. 2, e5.

10.1371/journal.ppat.0020005.

17.   Cole, J.N., Barnett, T.C., Nizet, V., and Walker, M.J. (2011). Molecular insight

into invasive group A streptococcal disease. Nat. Rev. Microbiol. 9, 724–736.

10.1038/nrmicro2648.

18.   Ikebe, T., Ato, M., Matsumura, T., Hasegawa, H., and Sata, T. (2010). Highly

Frequent Mutations in Negative Regulators of Multiple Virulence Genes in Group

A Streptococcal Toxic Shock Syndrome Isolates. PLOS Pathog. 6, e1000832.

10.1371/journal.ppat.1000832.

19.   Ministry of Health, Labour and Welfare. Severe Invasive Group A Streptococcal

Infections. https://www.mhlw.go.jp/bunya/kenkou/kekkaku-kansenshou11/01-

05-06.html.

20.   Centers for Disease Control and Prevention, C.f.D.C.a. Streptococcal Toxic Shock

Syndrome (STSS) (*Streptococcus pyogenes*) 2010 Case Definition.

https://ndc.services.cdc.gov/case-definitions/streptococcal-toxic-shock-

syndrome-2010/.

21.   Ikebe, T., Murai, N., Endo, M., Okuno, R., Murayama, S., Saitoh, K., Yamai, S.,

Suzuki, R., Isobe, J., Tanaka, D., et al. (2003). Changing prevalent T serotypes

and *emm* genotypes of *Streptococcus pyogenes* isolates from streptococcal toxic

shock-like syndrome (TSLS) patients in Japan. Epidemiol. Infect. 130, 569–572.

22.   Beall, B., Facklam, R.R., Elliott, J.A., Franklin, A.R., Hoenes, T., Jackson, D.,

Laclaire, L., Thompson, T., and Viswanathan, R. (1998). Streptococcal *emm* types

associated with T-agglutination types and the use of conserved *emm* gene

restriction fragment patterns for subtyping group A streptococci. J. Med.

Microbiol. *47,* 893–898. 10.1099/00222615-47-10-893.

23.   Johnson, D.R., and Kaplan, E.L. (1993). A review of the correlation of T-

agglutination patterns and M-protein typing and opacity factor production in the

identification of group A streptococci. J. Med. Microbiol. 38, 311–315.

10.1099/00222615-38-5-311.

24.   Katsukawa, C., Tamaru, A., Morikawa, Y., and Oda, K. (2002). M protein gene

(*emm*) typing of *Streptococcus pyogenes*. Kansenshogaku Zasshi 76, 238–245.

10.11150/kansenshogakuzasshi1970.76.238.

25.   Beres, S.B., Olsen, R.J., Saavedra, M.O., Ure, R., Reynolds, A., Lindsay, D.S.J.,

Smith, A.J., and Musser, J.M. (2017). Genome sequence analysis of *emm*89

Streptococcus pyogenes strains causing infections in Scotland, 2010–2016. J. Med.

Microbiol. 66, 1765–1773. 10.1099/jmm.0.000622.

26.   Chochua, S., Metcalf, B.J., Li, Z., Rivers, J., Mathis, S., Jackson, D., Gertz, R.E.,

Jr., Srinivasan, V., Lynfield, R., Van Beneden, C., et al. (2017). Population and

Whole Genome Sequence Based Characterization of Invasive Group A

Streptococci Recovered in the United States during 2015. mBio 8, e01422-17.

10.1128/mBio.01422-17.

27.   Davies, M.R., McIntyre, L., Mutreja, A., Lacey, J.A., Lees, J.A., Towers, R.J.,

Smeesters, P.R., Frost, H.R., Price, D.J., G Holden, M.T., et al. (2019). Atlas of

group A streptococcal vaccine candidates compiled using large-scale comparative

genomics. Nat. Genet. 51, 1035–1043. 10.1038/s41588-019-0417-8.

28.   Turner, C.E., Holden, M.T.G., Blane, B., Horner, C., Peacock, S.J., and

Sriskandan, S. (2019). The Emergence of Successful *Streptococcus pyogenes*

Lineages through Convergent Pathways of Capsule Loss and Recombination

Directing High Toxin Expression. mBio 10, 1–20. 10.1128/mBio.02521-19.

29. Paul, S., Olymon, K., Martinez, G.S., Sarkar, S., Yella. V.R., and Kumar, A. (2024).

MLDSPP: Bacterial Promoter Prediction Tool Using DNA Structural Properties

with Machine Learning and Explainable AI. J Chem Inf Model. 64, 7, 2705–2719.

10.1021/acs.jcim.3c02017

30. de Avila e Silva, S., Echeverrigaray, S., and Gerhardt G.J. (2011). BacPP:

Bacterial promoter prediction - A tool for accurate sigma-factor specific

assignment in enterobacteria. J Theor Biol. 287, 92–99.

31. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly

accurate protein structure prediction with AlphaFold. Nature, 583–589.

10.1038/s41586-021-03819-2.

32. Jain, I., Danger, J.L., Burgess, C., Uppal, T., and Sumby, P. (2020). The group A

Streptococcus accessory protein RocA: regulatory activity, interacting partners

and influence on disease potential. Mol. Microbiol. 113, 190–207.

10.1111/mmi.14410.

33. Ribardo, D.A., and McIver, K.S. (2006). Defining the Mga regulon: Comparative

transcriptome analysis reveals both direct and indirect regulation by Mga in the

group A streptococcus. Mol. Microbiol. 62, 491–508. 10.1111/j.1365-

2958.2006.05381.x.

34.   Kachroo, P., Eraso, J.M., Beres, S.B., Olsen, R.J., Zhu, L., Nasser, W., Bernard,

P.E., Cantu, C.C., Saavedra, M.O., Arredondo, M.J., et al. (2019). Integrated

analysis of population genomics, transcriptomics and virulence provides novel

insights into *Streptococcus pyogenes* pathogenesis. Nat. Genet. 51, 548–559.

10.1038/S41588-018-0343-1.

35.   Vega, L.A., Malke, H., and McIver, K.S. (2016). Virulence-Related

Transcriptional Regulators of *Streptococcus pyogenes*. In *Streptococcus*

pyogenes : Basic Biology to Clinical Manifestations [Internet], J.J. Ferretti, D.L.

Stevens, and V.A. Fischetti, eds (University of Oklahoma Health Sciences Center).

36.   Terao, Y., Mori, Y., Yamaguchi, M., Shimizu, Y., Ooe, K., Hamada, S., and

Kawabata, S. (2008). Group A streptococcal cysteine protease degrades C3 (C3b)

and contributes to evasion of innate immunity. J. Biol. Chem. 283, 6253–6260.

10.1074/jbc.M704821200.

37.   Sumitomo, T., Nakata, M., Higashino, M., Jin, Y., Terao, Y., Fujinaga, Y., and

Kawabata, S. (2011). Streptolysin S contributes to group A streptococcal

translocation across an epithelial barrier. J. Biol. Chem. 286, 2750–2761.

10.1074/jbc.M110.171504.

38. Honda-Ogawa, M., Ogawa, T., Terao, Y., Sumitomo, T., Nakata, M., Ikebe, K.,

Maeda, Y., and Kawabata, S. (2013). Cysteine proteinase from Streptococcus

pyogenes enables evasion of innate immunity via degradation of complement

factors. J. Biol. Chem. 288, 15854–15864. 10.1074/jbc.M113.469106.

39. Lin, A., Loughman, J.A., Zinselmeyer, B.H., Miller, M.J., and Caparon, M.G.

(2009). Streptolysin S inhibits neutrophil recruitment during the early stages of

Streptococcus pyogenes infection. Infect. Immun. 77, 5190–5201.

10.1128/IAI.00420-09.

40. Cimen, M.Y. (2008). Free radical metabolism in human erythrocytes. Clin. Chim.

Acta. 390, 1–11. 10.1016/j.cca.2007.12.025.

41. Laabei, M., and Ermert, D. (2019). Catch Me if You Can: *Streptococcus pyogenes*

Complement Evasion Strategies. J. Innate. Immun. 11, 3–12. 10.1159/000492944.

42. He, L.Y., Yu, Y.B., Liu, Y., Le, Y.J., Li, S., and Yang, X.Y. (2022). Immunization

with the lipoprotein FtsB stimulates protective immunity against *Streptococcus*

pyogenes infection in mice. Front. Microbiol. 13, 969490.

10.3389/fmicb.2022.969490.

43.   Nguyen, M.T., and Gotz, F. (2016). Lipoproteins of Gram-Positive Bacteria: Key

Players in the Immune Response and Virulence. Microbiol. Mol. Biol. Rev. 80,

891–903. 10.1128/MMBR.00028-16.

44.   Bessen, D.E., McShan, W.M., Nguyen, S.V., Shetty, A., Agrawal, S., and Tettelin,

H. (2015). Molecular epidemiology and genomics of group A Streptococcus.

Infect. Genet. Evol. 33, 393–418. 10.1016/j.meegid.2014.10.011.

45.   McShan, W.M., McCullor, K.A., and Nguyen, S.V. (2019). The Bacteriophages of

Streptococcus pyogenes. Microbiol. Spectr. 7. 10.1128/microbiolspec.GPP3-

0059-2018.

46.   Berbel, D., Camara, J., Gonzalez-Diaz, A., Cubero, M., Lopez de Egea, G., Marti,

S., Tubau, F., Dominguez, M.A., and Ardanuy, C. (2021). Deciphering mobile

genetic elements disseminating macrolide resistance in *Streptococcus pyogenes*

over a 21 year period in Barcelona, Spain. J. Antimicrob. Chemother. 76, 1991–

2003. 10.1093/jac/dkab130.

47.   Wajima, T., Chiba, N., Morozumi, M., Shouji, M., Sunaoshi, K., Sugita, K.,

Tajima, T., Ubukata, K., and Group A Streptococcus Surveillance Study Group.

(2014). Prevalence of macrolide resistance among group A streptococci isolated

from pharyngotonsillitis. Microb. Drug. Resist. 20, 431–435.

10.1089/mdr.2013.0213.

48.   Hasegawa, T., Hata, N., Matsui, H., Isaka, M., and Tatsuno, I. (2017).

Characterisation of clinically isolated *Streptococcus pyogenes* from

balanoposthitis patients, with special emphasis on emm89 isolates. J. Med.

Microbiol. 66, 511–516. 10.1099/jmm.0.000460.

49.   Tatara, K., Gotoh, K., Okumiya, K., Teramachi, M., Ishimoto, K., Tanaka, Y.,

Iwahashi, J., Shindou, S., Yamashita, Y., and Watanabe, H. (2020). Molecular

epidemiology, antimicrobial susceptibility, and characterization of

fluoroquinolone non-susceptible *Streptococcus pyogenes* in Japan. J. Infect.

Chemother. 26, 280–284. 10.1016/j.jiac.2019.10.004.

50.   Tanaka, Y., Gotoh, K., Teramachi, M., Ishimoto, K., Tsumura, N., Shindou, S., and

Yamashita, Y. (2016). Molecular epidemiology, antimicrobial susceptibility, and

characterization of macrolide-resistant *Streptococcus pyogenes* in Japan. J. Infect.

Chemother. 22, 727–732. 10.1016/j.jiac.2016.06.013.

51.   Ubukata, K., Wajima, T., Morozumi, M., Sakuma, M., Tajima, T., Matsubara, K.,

Itahashi, K., and Iwata, S. (2020). Changes in epidemiologic characteristics and

antimicrobial resistance of *Streptococcus pyogenes* isolated over 10 years from

Japanese children with pharyngotonsillitis. J. Med. Microbiol. 69, 443–450.

10.1099/jmm.0.001158.

52.   Jolley, K.A., Bray, J.E., and Maiden, M.C.J. (2018). Open-access bacterial

population genomics: BIGSdb software, the PubMLST.org website and their

applications. Wellcome Open Res. 3, 124. 10.12688/wellcomeopenres.14826.1.

53.   Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one

FASTQ   preprocessor.   Bioinformatics   34,   i884–i890.

10.1093/bioinformatics/bty560.

54.   Kapatai, G., Coelho, J., Platt, S., and Chalker, V.J. (2017). Whole genome

sequencing of group A Streptococcus: development and evaluation of an

automated pipeline for *emm* gene typing. PeerJ 5, e3226. 10.7717/peerj.3226.

55.   Souvorov, A., Agarwala, R., and Lipman, D.J. (2018). SKESA: strategic k-mer

extension for scrupulous assemblies. Genome Biol. 19, 153. 10.1186/s13059-018-

1540-z.

56.   Jolley, K.A., and Maiden, M.C. (2010). BIGSdb: scalable analysis of bacterial

genome variation at the population level. BMC Bioinformatics 11, 595.

10.1186/1471-2105-11-595.

57.   Maiden, M.C.J., Jansen Van Rensburg, M.J., Bray, J.E., Earle, S.G., Ford, S.A.,

Jolley, K.A., and McCarthy, N.D. (2013). MLST revisited: the gene-by-gene

1019 approach to bacterial genomics Europe PMC Funders Group. Nat. Rev. Microbiol.

1020 11, 728–736. 10.1038/nrmicro3093.

1021 58. Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation.

1022 Bioinformatics 30, 2068–2069. 10.1093/bioinformatics/btu153.

1023 59. Page, A.J., Cummins, C.A., Hunt, M., Wong, V.K., Reuter, S., Holden, M.T.,

1024 Fookes, M., Falush, D., Keane, J.A., and Parkhill, J. (2015). Roary: rapid large-

1025 scale prokaryote pan genome analysis. Bioinformatics 31, 3691–3693.

1026 10.1093/bioinformatics/btv421.

1027 60. Jaillard, M., Lima, L., Tournoud, M., Mahé, P., van Belkum, A., Lacroix, V., and

1028 Jacob, L. (2018). A fast and agnostic method for bacterial genome-wide

1029 association studies: Bridging the gap between k-mers and genetic events. PLOS

1030 Genet. 14, e1007758. 10.1371/journal.pgen.1007758.

1031 61. Nguyen, L.T., Schmidt, H.A., von Haeseler, A., and Minh, B.Q. (2015). IQ-TREE:

1032 a fast and effective stochastic algorithm for estimating maximum-likelihood

1033 phylogenies. Mol. Biol. Evol. 32, 268–274. 10.1093/molbev/msu300.

1034 62. Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., and Jermiin,

1035 L.S. (2017). ModelFinder: fast model selection for accurate phylogenetic

1036 estimates. Nat. Methods 14, 587–589. 10.1038/nmeth.4285.

63.   Letunic, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool

for phylogenetic tree display and annotation. Nucleic Acids Res. 49, W293–W296.

10.1093/nar/gkab301.

64.   R Core Team. (2022). A language and environment for statistical computing.

https://www.R-project.org/.

65.   Lees, J.A., Galardini, M., Bentley, S.D., Weiser, J.N., and Corander, J. (2018).

pyseer: A comprehensive tool for microbial pangenome-wide association studies.

Bioinformatics 34, 4310–4312. 10.1093/bioinformatics/bty539.

66.   Ondov, B.D., Treangen, T.J., Melsted, P., Mallonee, A.B., Bergman, N.H., Koren,

S., and Phillippy, A.M. (2016). Mash: fast genome and metagenome distance

estimation using MinHash. Genome Biol. 17, 132. 10.1186/s13059-016-0997-x.

67.   Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis, 2 Edition

(Springer-Verlag New York). 10.1007/978-3-319-24277-4.

68.   Beres, S.B., Kachroo, P., Nasser, W., Olsen, R.J., Zhu, L., Flores, A.R., de la Riva,

I., Paez-Mayorga, J., Jimenez, F.E., Cantu, C., et al. (2016). Transcriptome

remodeling contributes to epidemic disease caused by the human pathogen

Streptococcus pyogenes. mBio 7, 1–14. 10.1128/mBio.00403-16.

69.   Martinez, G.S., Perez-Rueda, E., Kumar, A., Dutt, M., Maya, C.R., Ledesma-

Dominguez, L., Casa, P.L., Kumar, A., de Avila e Silva, S., and Kelvin, D.J.

(2024). CDBProm: the Comprehensive Directory of Bacterial Promoters. NAR

Genom Bioinform. 6, 1, lqae018. 10.1093/nargab/lqae018.

70.   Rice, P., Longden, I., and Bleasby, A. (2000). EMBOSS: the European Molecular

Biology Open Software Suite. Trends Genet. 16, 276–277. 10.1016/s0168-

9525(00)02024-2.

71.   Mariani, V., Biasini, M., Barbato, A., and Schwede, T. (2013). lDDT: a local

superposition-free score for comparing protein structures and models using

distance difference tests. Bioinformatics 29, 2722–2728.

10.1093/bioinformatics/btt473.

72.   Yin, R., Feng, B.Y., Varshney, A., and Pierce, B.G. (2022). Benchmarking

AlphaFold for protein complex modeling reveals accuracy determinants. Protein

Sci. 31, e4379. 10.1002/pro.4379.

73.   Hirokawa, T., Boon-Chieng, S., and Mitaku, S. (1998). SOSUI: classification and

secondary structure prediction system for membrane proteins. Bioinformatics 14,

378–379. 10.1093/bioinformatics/14.4.378.

74.   Takamatsu, D., Osaki, M., and Sekizaki, T. (2001). Thermosensitive suicide

vectors for gene replacement in *Streptococcus suis*. Plasmid 46, 140–148.

10.1006/PLAS.2001.1532.

75.   Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general

purpose program for assigning sequence reads to genomic features.

Bioinformatics 30, 923–930. 10.1093/bioinformatics/btt656.

76.   Ge, S.X., Son, E.W., and Yao, R. (2018). iDEP: an integrated web application for

differential expression and pathway analysis of RNA-Seq data. BMC

Bioinformatics 19, 534. 10.1186/s12859-018-2486-6.

77.   Lancefield, R.C. (1957). Differentiation of group A streptococci with a common

R antigen into three serological types, with special reference to the bactericidal

test. J. Exp. Med. 106, 525–544. 10.1084/jem.106.4.525.

78.   Yamaguchi, M., Goto, K., Hirose, Y., Yamaguchi, Y., Sumitomo, T., Nakata, M.,

Nakano, K., and Kawabata, S. (2019). Identification of evolutionarily conserved

virulence factor by selective pressure analysis of *Streptococcus pneumoniae*.

Commun. Biol. 2, 96. 10.1038/s42003-019-0340-7.

79.   Takemura, M., Yamaguchi, M., Kobayashi, M., Sumitomo, T., Hirose, Y., Okuzaki,

D., Ono, M., Motooka, D., Goto, K., Nakata, M., et al. (2022). Pneumococcal

BgaA Promotes Host Organ Bleeding and Coagulation in a Mouse Sepsis Model.

Front. Cell. Infect. Microbiol. 12, 844000. 10.3389/fcimb.2022.844000

Figures

Figure 1. Pan-genomic and phylogenetic characterization of the collected *emm*89 *S.*

***pyogenes* isolates.**

(A) *emm* genotyping of the 207 clinical isolates collected in Japan. (B) Pan-genome

analysis of the Japanese and global cohorts. All genes detected in each cohort were

classified into four groups, according to prevalence: core, soft-core, shell, and cloud genes.

(C) Phylogenetic tree for the global cohort, based on the sequences of the core genes.

From the inside, the color bars show clusters, phenotypes, MLSTs, clades, countries

where strains were isolated, and global regions, in the order mentioned. The root of the

tree was set as the mid-point. The scale located upper left indicates 0.001 times

substitution of the bases on average. Arrowhead indicates strain belonging to the novel

clade.

MLST, multilocus sequence typing.

1105

Figure 2. Pan-genome-wide association study on SNPs/indels.

(A–B) Manhattan plots for the Japanese (A) and global (B) cohorts. The X-axis shows

the location of each SNP/indel on the core gene alignment, while the Y-axis indicates the

p-value. For each cohort, a permutation test was performed by iterating the calculations

1,000 times with randomly permuted phenotypes, with the significance level set at the 5th

percentile of the 1,000 minimal p-values (p=5.75×10^{-4} and p=1.05×10^{-4} for the Japanese

and global cohorts, respectively). Plots with lower p-values than genome-wide significant

levels are colored magenta and blue, based on the direction of their effect size (positive

and negative, respectively). (C) Distribution heatmap for the global cohort, with the

strains possessing the significant SNPs/indels colored orange. Only the 20 SNPs with the

lowest p-values are shown in this heatmap. Colored bars above indicate countries and

phenotypes, and magenta bars represent invasive phenotypes. Using the Roary program,

gene names starting with "Group_" were automatically assigned. Position indicates the

location of each SNP/indel on the core gene alignment. The full results are shown in Table

S6.

SNP, single-nucleotide polymorphism; indel, insertion/deletion.

1122

Figure 3. Pan-genome-wide association study on gene presence.

(A–B) Volcano plots for the Japanese (A) and global (B) cohorts. The X-axis shows the

effect size, while the Y-axis indicates the *p*-value. Plots with a lower *p*-value than the

genome-wide significant levels ($p=1.09\times10^{-4}$ and $p=7.72\times10^{-5}$ for the Japanese and

global cohorts, respectively) have been colored magenta and blue, based on the direction

of their effect size (positive and negative, respectively). (C) Distribution heatmap for the

global cohort, with the strains possessing the significant genes colored orange. Only the

20 genes with the lowest p-values are shown in this heatmap. Colored bars above indicate

countries and phenotypes, and magenta bars represent invasive phenotypes. Using the

Roary program, gene names starting with "Group_" were automatically assigned. The full

results are shown in Table S8.

1134

Figure 4. K-mers related to pathology.

Detailed results of the GWAS on k-mers for the Japanese cohort in two genomic regions: *covS* (A, top left) and *group_184* (B); and the global cohort in five regions: *covS* (A, top right), an intergenic region (C), *group_141-143* (D), *sagG* (E), and *fhuB* (F). (A–F top)

de-Bruijn graphs generated using DBGWAS. The respective nodes in the graphs indicate

k-mers, with the significant nodes indicated using green arrowheads. The size of each

node corresponds to the allele frequency. (A–F, bottom) Alignments of k-mers or maps

of k-mers on the genome sequence of MGAS27061 around regions including significant

mutations. Each base is colored according to the interpreted amino acids. (A) The

arrowheads indicate significant k-mers that cause a frameshift mutation, truncating CovS

protein to 35 amino acids. (B) The significant k-mers indicate that the presence of a

sequence mapped on the first 26 bp of group_184 and its upstream 20 bp is related to the

pathology. (C) Two significant k-mers indicate the same intergenic region of 270 bps. (D)

The significant k-mers are indicated using arrowheads.

1149

Figure 5. Predicted protein structure models.

(A) Snake-like plot of transmembrane regions of CovS estimated using SOSUI.

Frameshift mutations, detected by the SNPs/indel- and k-mer-based GWASes, cause

truncation of CovS at the indicated 35[th] and 45[th] residues, respectively. (B) Structural

model of the CovS homodimer (ipTM + pTM=0.614). The putative transmembrane

regions are colored orange. The upper part is the sensor domain, while the lower is the C-

terminal kinase domain involved in the phosphorylation of the transcriptional regulator

CovR. (C) Snake-like plot depicting the transmembrane regions of FhuB and FhuG. The

73rd residue of FhuB is indicated by an arrowhead and in magenta. (D) Structural model

of the FhuBDCCG complex (ipTM + pTM=0.791). The putative transmembrane regions

of FhuB and FhuG are colored green-yellow and peach, respectively. The upper part of

the model is located in the extracellular region, while the lower part is in the cytoplasm.

(E) The 73rd valine in FhuB, shown in magenta, was substituted with alanine. The

molecular surface is illustrated with a wireframe and that of the predicted indentation is

shown with an arrowhead.

ipTM + pTM: Weighted combination of interface-predicted TM and predicted TM scores.

ipTM is used to measure structural accuracy in the protein-protein interface, while ipTM

is a metric for overall topological accuracy.

1168

Figure 6. Transcriptome analysis of the *fhuB* T218C mutant strain in THY and

human blood.

(A) Principal component analysis plot of RNA-seq data. (B) Differentially expressed

genes in four comparisons. (C) Plot of gene expression in the WT strain *vs*. *fhuB* T218C

mutant in human blood. Significantly upregulated and downregulated genes in the WT

are colored red and blue, respectively. The shapes of the plots indicate relative

transcriptional changes between THY and blood. Genes depicted with upward triangles

are either significantly upregulated in the WT or downregulated in the mutant strain, in

1177   blood *vs.* THY. The downward triangle plots indicate genes that are either downregulated

1178   in the WT or upregulated in the mutant strain, in blood *vs.* THY. WT, wild-type; THY,

1179   Todd Hewitt broth supplemented with 0.2% yeast extract.

1180

Figure 7. Effects of the *fhuB* T218C mutant strain on ferric ion uptake and bacterial

survival in human blood.

(A) Intracellular ferric ion assay. The WT and *fhuB* T218C mutant strains were incubated

in healthy human blood for 3 h, following which the intracellular ferric ion concentrations

were measured. (B) Bactericidal assay in human blood. The WT and *fhuB* T218C mutant

were mixed with healthy human blood to measure the ratio of bacterial counts at 1, 2, and

3 h to those at 0 h after infection. (C–F) Bactericidal assay in human erythrocyte-rich

medium (C), polymorphonuclear cell-rich medium (D), plasma (E), and plasma

inactivated by heating at 56ºC for 30 min (F). (G) Bacterial growth in brain heart infusion

broth. The data were pooled from three independent experiments, each performed in

sextuplicate. Thick bars and error bars indicate means and quartiles, respectively.

Statistical significance was determined using the Mann–Whitney test with Benjamini–

Hochberg's correction.

1194

Table 1. Comparison of clustering on the phylogenetic trees for the Japanese and global cohorts.

Clusters	A	B1	B2	B3	
Japanese		2	2	3	416
Global		19	2	5	639
Standard residuals		-4.33	-1.85	-2	5.14
Chi-squared test			$p = 0.13$		

Table 2. Complexes including k-mers significantly related to severe invasiveness detected in the Japanese cohort.

Complexes	p-value	q-value	Genes in complexes
Comp_11	4.41E-05	1.49E-02	*covS*
Comp_2	3.91E-05	4.22E-02	*group_184, group_185, group_186, group_1278, group1443, group_1519, group_1520, group_2169*

Table 3. Complexes including k-mers significantly related to invasiveness detected in the global cohort.

Complexes	p-value	q-value	Genes in complexes
Comp_7	2.35E-06	1.90E-02	*group_130, covS*
Comp_6	1.05E-05	1.90E-02	(intergenic region)
Comp_24	1.96E-06	1.90E-02	*group_141, group_142, group_142*
Comp_10	2.20E-05	2.40E-02	*sagG, sagI*
Comp_15	2.05E-05	2.40E-02	*fhuB, fhuD*

MLSTs
Phenotypes
Clusters

Tree scale: 0.001 ├──────┤

MLSTs
☐ 28 ■ 101 ■ 1450 ■ 1451 ■ 1452 ■ 1463 ■ 646 ■ 1454 ■ 1456 ☐ 1285 ■ 1461

Phenotypes
☐ Non-invasive ■ Invasive

Clusters
■ A ■ B1 ■ B2 ■ B3

Figure S1. Phylogenetic tree for the Japanese cohort.
From the inside, color bars show clusters, phenotypes, and MLSTs, in the order mentioned. The root of the tree was set as the mid-point.
The scale located in the upper left indicates 0.001 times substitution of the bases on average.

Figure S2.   Variations in the *nga* promoter region.
Sequences of the *nga* promoter are classified into four major genotypes, according to three key residues (shown as -27, -22, and -18 in the figure) affecting promotor activity. Turner *et al.* further determined subtypes using mutations other than the key residues. FS24 strain has the same allele $A_{-27}G_{-22}T_{-18}$ as the type 3 or clade 3 but has an SNP highlighted in yellow in the -10 box.

Figure S3. Pan-genome-wide association study on SNPs/indels.
(A–B) Scree plot of eigenvalues generated by decomposition of the genetic distance matrix for the Japanese (A) and global (B) cohort. To determine the number of principal components used as covariates for pan-GWAS, the plot in which the subsequent principal components seem to have relatively small eigenvalues was sought. Consequently, we used seven and three principal components as covariates for the Japanese and global cohorts, respectively. (C) Number of significant SNPs accumulated in single genes found in the pan-GWAS for the global cohort. Only the top 30 genes are shown in this graph.

Figure S4. Pairwise correlation of significant COGs.
Each cell indicates the correlation coefficient of each significant COG pair on its presence in the global cohort. The COGs are sorted based on hierarchical clustering. Annotations and phenotypes are represented by circles and cells located on the right and bottom of the heatmap. The dotted line on the tree divides the COGs into eight clusters.

Figure S5. Predicted protein structure model of LacE.

(A) Snake-like plot of transmembrane regions of LacE estimated using SOSUI. The 554th residue has been indicated with an arrowhead and in magenta. (B) Structure model of LacE generated with AlphaFold (pLDDT=0.902). The putative transmembrane regions are colored in orange. LacE has two domains: the transmembrane domain EIIC and intracellular domain EIIB. The 554th glycine substituted with valine is colored in magenta.

pLDDT: The best predicted local difference distance test indicating local structural accuracy.

Figure S6.   Pipeline for bacterial pan-genome analysis, pan-GWAS, and GWAS constructed in this study.
We developed a pipeline for whole genome analyses working on the National Institute for Genetics (NIG) Supercomputer and SQUID at the Cybermedia Center, Osaka University.