Female fruit flies use social cues to make egg clustering decisions

Emily R. Churchill[1], Emily K. Fowler[2], Lucy A. Friend[2], Marco Archetti[3], Douglas W. Yu[2], Andrew

F. G. Bourke[2], Tracey Chapman[2*] and Amanda Bretman[1*]

* Corresponding authors

1. School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, UK

2. School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

NR4 7TJ, UK

3. Department of Biology, Pennsylvania State University, University Park, PA 16802, USA

ORCID information:

Emily R. Churchill – 0000-0003-0910-8600

Emily K. Fowler – 0000-0001-5290-7630

Lucy A. Friend – 0000-0001-5728-9384

Marco Archetti – 0000-0002-8303-1632

Douglas W. Yu – 0000-0001-8551-5609

Andrew F. G. Bourke – 0000-0001-5891-8816

Tracey Chapman – 0000-0002-2401-8120

Amanda Bretman – 0000-0002-4421-3337

Keywords

Cooperative oviposition; Laying rate; Public goods; Density; Fitness; *Drosophila melanogaster*

Abstract

The ability to respond plastically to environmental cues is a key determinant of fitness. In

changing environments, females plastically adjust the number and placement of eggs they

lay, allowing them to optimise the level of resources available for offspring by minimising

over-exploitation whilst also accruing potential cooperative benefits such as through

communal larval feeding. This suggests that the pattern of female egg laying should vary

across differing social environments, to balance potential for cooperative benefits versus

competitive costs of egg clustering. Here we test this prediction in the fruit fly *Drosophila*

melanogaster. We developed a mathematical null model, which verified that females lay eggs

on substrates in significantly non-random patterns. We then tested the effects of

manipulating social density and egg density on the egg laying patterns of mated females. As

the density of adult females was increased, females laid more eggs, laid them faster, and laid

more eggs in clusters. Females also preferred to place eggs within existing egg clusters -and

most egg clusters were of mixed maternity. These results show that females in groups, or

females that encounter already laid eggs, adjust their laying to place more eggs in mixed

maternity clusters. Overall, the results are consistent with our predictions and reveal that

females exhibit active choice in egg investment and egg deposition patterns in response to

variation in social environments.

Background

In natural contexts, social environments and availability of resources are highly variable, exposing populations to fluctuating competition. Phenotypic plasticity can play a key role in ensuring individuals maximise their lifetime reproductive success (1, 2). Because optimal strategies may vary with fluctuating levels of competition, individuals that can respond and make appropriate behavioural and physiological adjustments to the prevailing environment should show increased fitness. This should be particularly true in females, who generally exert greater control over reproductive investment in progeny. For example, females can control paternity (3-5), the number and sex ratio of offspring via ejaculate manipulation (6-8), and choice of nesting/oviposition site locations. For females that deposit eggs upon a substrate, egg site location can be a vital determinant of hatching success. Consistent with this, oviposition patterns appear to be plastic in response to environmental variation (9-12).

Competition, cooperation, and cheating amongst females

At increasing population densities, there is expected to be a greater risk of competition but also increased potential for cooperation, and the relative importance of the two is expected to differ as individuals progress through life stages. For example, when food is limiting and competition high, eggs and larvae are typically more vulnerable to cannibalism than are adults, due to their more restricted mobility (13-15). On the other hand, in some species, there is a benefit to being with conspecifics when foraging or hunting. For example, Eurasian Coots (*Fulica atra*) copy each other's behaviours to reduce foraging costs (16), and larvae in metamorphosing species often benefit from increased efficiencies resulting from communal feeding (17, 18). Females can use the presence of conspecifics as a cue to modify reproductive investment in response to changing social conditions. For example, increasing adult female density reduces offspring production and body size in Western Mosquitofish (*Gambusia affinis*) (19). In contrast, in the Willow Leaf beetle (*Plagiodera versicolora*) fitness effects of increasing adult female density can vary with development, resulting in a decrease in egg hatching, but an increase in egg clutch number and duration of laying period (20).

Among females of species that deposit their eggs upon substrates, it is also often found that eggs are laid in conspecific clusters, e.g. in fish (21), birds (22), reptiles, amphibians (23), and invertebrates (24). Egg clustering may increase the local population density of

offspring and thus the potential not only for competition but also cooperation at early life-

history stages. Benefits from cooperation could include reduced energy expenditure for

ovipositing females by minimising search times, reduced egg predation risk (the dilution

effect' hypothesis (25)), increased protection from abiotic challenges, or increased larval

communal feeding.

However, cooperative interactions can be exploited by cheater individuals. In some

egg-laying species mothers also provide their eggs with diffusible defensive compounds. For

example, medfly (*Ceratitis capitata*) females provision the surface of their eggs with

antimicrobial peptides, which spread into the surrounding environment, creating a zone of

microbial growth inhibition (26). Other examples include in the Green lacewing moth

(*Ceraeochrysa cubana*), in which eggs are endowed with an anti-predator alkaloid (27), and

in fruit flies (*Drosophila melanogaster*), in which eggs are coated with anticannibalism

compounds (28). These defensive substances represent potential public goods (29, 30) as all

eggs nearby could benefit, even eggs that do not themselves provide the defensive compound

itself. This raises the intriguing possibility that in egg clusters of mixed maternity, there could

be 'co-operator' eggs coated with defensive compounds, and non-defended 'cheater' eggs

that benefit from their placement within the defensive diffusion radius. In this scenario,

cheaters benefit by avoiding the energetic expense of provisioning eggs with defensive

compounds. Egg clustering might also provide benefits as well as cheating opportunities

through communal feeding. For example, if eggs are laid on harder substrates there could be

benefits for all via the collective processing of food, while simultaneously allowing the

possibility for cheaters to benefit by utilising resources liberated by the processing of food by

their cluster mates. Such co-operators and cheats could stably coexist within a population if

the benefits of public goods have a non-linear relationship with fitness (30, 31).

In summary, we expect that natural selection can favour either egg dispersion

(competition) or egg-clustering (cooperation and cheating) behaviour. Which of these is

selected for will depend on the balance of abiotic and selective forces. The potential fitness

benefits of joining an existing egg cluster could depend on key factors such as number, quality,

fertility, and age of eggs already present. Based on existing evidence that females can vary

the number of eggs they lay according to social density (9, 10, 32) we predict, and test here,

the idea that females should use number of eggs and/or number of adult conspecifics as cues

to direct their egg-clustering patterns.

Drosophila melanogaster as a model for understanding variation in oviposition patterns

In wild populations, *D. melanogaster* larvae and adults periodically occur at high densities

around fallen, fermenting fruit. For females this fruit is both a source of nutrition and

potential oviposition sites. However, these key resources are patchy and ephemeral, and they

vary in their nutrient availability (density, ripeness, and state of decomposition). Access to

such resources is also shaped by the immediate social environments (number of conspecific

or heterospecific males and females utilising the same resource) (33). To oviposit in a way

that maximises fitness, female *Drosophila* should assess the nutritional quality, degree of

interspecific competition, risk of pathogens and parasites present at each oviposition site, as

well as the ongoing search costs, and potential benefits and costs (e.g. from overexploitation

leading to dispersal) for developing offspring (34).

Consistent with this reasoning, female *Drosophila* have a social attraction to

conspecifics on oviposition resources (35, 36) which they can detect directly or via

pheromones, markings, or the presence of eggs or larvae (37-41). This attraction could arise

because it allows females to copy the site-selection choices of others (37, 38, 40) to enable

cooperative feeding among larvae (42), or to gain other public goods benefits of egg

clustering, as outlined above. Existing evidence suggests that female *Drosophila* can adjust

their fecundity responses according to their previous or current social environments, with

females laying fewer eggs after pre-mating exposure to conspecifics (9, 43) but laying eggs

more quickly within larger current post-mating social groups (32).

Drosophila eggs can often be found in extremely close proximity, in large groups or

clusters (Fig. S1). Hence, after deciding to lay on substrates with existing conspecific eggs,

females must also decide whether to lay eggs that join existing clusters. Females generally lay

one egg at a time (44), explore oviposition sites between laying, and can retain eggs until an

optimal laying site is found (45). Therefore, clustering of eggs suggests that females make

repeated egg-laying decisions and that females lay their eggs in non-random distributions.

However, whether the temporal or spatial distribution of egg laying is significantly non-

random has not previously been formally tested. In addition, social density is reported to

affect the number and location of eggs laid by *D. melanogaster* (9, 10, 32), but nothing is yet

known about how it might affect clustering rates or patterns. Here we address these key gaps

in our knowledge of the social influences on oviposition patterns.

We first test, by using a mathematical model, a key hypothesis:

• (H1) that female *D. melanogaster* do indeed place their eggs non-randomly within a food
patch

We then conducted experiments to determine key drivers of the significantly non-
random egg deposition observed. We did this by testing whether the decisions to join existing
clusters responded to experimentally manipulated adult female group size and egg presence
or density.

Due to the potential for resource competition, cooperation, or public goods benefits,
we predicted that, as adult female social density increased, females should:

• (H2) decrease laying latency (if copying other's laying decisions (38))

• (H3) invest more in egg production (either through laying more eggs ((46)), or fewer eggs
of higher quality)

• (H4) alter egg-clustering pattern (if clustering balances cooperative larval feeding, whilst
avoiding competition)

• (H5) incur fitness benefits via increased egg clustering (increased egg-adult viability of
highly clustered eggs)

• (H6) lay eggs in clusters of mixed -maternity to potentially gain public goods-related
benefits

In addition to adult presence (H2-6), we also predicted that the presence of existing
egg clusters should elicit comparable responses in egg laying (as females should use similar
density cues). Due to the expected advantages of clustering (17, 18, 25-28, 42), and assuming

benefits increase with cluster size, we hypothesised that (H7) females should be more willing

to lay their eggs within existing clusters, and more eggs should be laid in larger clusters as the

number and size of existing clusters present increases.

Methods

Our experimental strategy was first to construct a model, parameterised with data, to test

the hypothesis that females exhibit non-random egg placement (H1). Having confirmed that

egg laying patterns were significantly non-random, we then conducted experiments to test

the effect of social group size on the rate and likelihood of egg laying, egg clustering patterns

(H2-H4) and fitness (egg-adult viability as a function of egg clustering (H5)). An additional

experiment was then conducted using females that produced dyed eggs, to measure the

frequency of egg clusters with mixed maternity (H6). Finally, we tested the effect of exposing

standard sized groups of females to environments containing variable numbers of egg

clusters, and recording the rate and pattern of egg laying (H7).

Fly husbandry and rearing of experimental flies

Fly rearing and all experiments were conducted at 25°C on a 12h light:dark cycle (09:00-21:00

GMT). Wild type flies originating from a Dahomey laboratory population maintained in cages

were used throughout. We collected eggs from cages using Petri dishes (*Sarstedt*

#82.1473.001) filled with grape juice agar-based medium (50g agar (*Fisher Scientific*

#10048991), 600ml red grape juice (*Young's Brew red wine enhancer*), 42ml Nipagin solution

(methylparaben, 10% w/v solution, dissolved in 95% Ethanol) per 1.1l H_2O). Once eggs had

hatched, we transferred first instar larvae to a 40ml plastic vial (*Sarstedt #58.490*) containing

7ml of a standard sugar-yeast-agar (SYA) medium (100g brewer's yeast (*Buy Wholefoods*

Online), 50g sugar (*Tate & Lyle*), 15g agar, 30ml Nipagin solution, and 3ml propionic acid

(*Fisher Scientific #10193190*) per litre of medium), at a standard density of 100 larvae per vial

(i.e., 'standard density vials'). Experimental flies were then collected under light ice

anaesthesia, within six hours of eclosion to ensure virginity.

Egg-clustering model to test for non-random egg deposition (H1)

To ascertain whether the proportion of observed egg clustering differed from a null

expectation (H1), we produced a model to calculate random distributions of eggs in a

standard vial. We defined a 'cluster' as any group of two or more eggs in which the main

bodies of the eggs were in direct contact (Fig. S1). In the model, given that the observed egg

to dish area is 1/5770, we assume that there are 5770 positions available for egg laying; there

is a total number (parameter Ɛ) of eggs per vial. Assuming that all eggs are laid sequentially

and at random, we calculated the expected size of clusters (defined as the number of eggs

per cluster) for any number of total eggs laid (Ɛ).

The preference for clustering eggs next to an existing egg (hereafter 'clustering

preference') is defined by the probability of a clustering decision. Let us call К the probability

that a fly will lay an egg next to an already existing egg. After the first egg is placed at random,

where Ɛ > 1, the next egg is either laid in a random position (probability of probability of 1 -

К) or next to an existing egg (probability of К); in the latter case, the position of the egg is

chosen at random amongst only positions with existing eggs.

The expected number of eggs per cluster can be predicted based on К and Ɛ. When К

= 0 (no preference for egg clustering), very few clusters with 2 eggs occur and virtually no

clusters with more than 2 eggs (2.5% with Ɛ = 233 (the maximum number of eggs observed in

our experiment); 1% with Ɛ = 100; no clusters for Ɛ below 70). Therefore, if Ɛ << 5770 (as is

the case in our experiment), randomly laid eggs form almost exclusively singly laid eggs. When

К = 1 (absolute preference for egg clustering), all eggs are clustered in one large cluster in a

single position.

At intermediate values of К, some variation in cluster size emerges. We simulated the

expected number and size of egg clusters for different values of К from 0 to 1 in intervals of

0.1 (using the average of 10 different simulations per К value). We then compared these

expected distributions to the empirically observed distributions of eggs laid by females from

the differing social density treatments (groups of 1, 2, 4, or 8) described below, to estimate

the most likely preference К leading to that distribution. For each fly, we performed a

Kolmogorov-Smirnov test (implemented in Mathematica 13. 1 (49)) between the expected

and observed distribution, for all К values. We considered the К value yielding the highest

(least significant) P values to be the most likely К value for that fly.

Effects of adult social density on egg laying rate, egg number, egg-clustering decisions and fitness

(egg-adult viability) (H2-H5)

We investigated how variation in adult social group size affected egg laying decisions:

specifically, the speed of egg laying (H2), the number of eggs laid (H3), egg clustering patterns

(H4), and fitness (egg-adult viability of clustered versus non-clustered eggs) (H5). The data

from this experiment were also used to parameterise the model, in order to test for non-

random egg laying patterns (H1). Experimental flies were collected and housed in same-sex

groups of 10 in standard vials. At four days post-eclosion, females were randomly assigned to

one of the following social group treatments: 1, 2, 4, or 8 females and kept in these conditions

for 48h (N = 30 vials each treatment; Fig. S1). Males were held for six days until they were

used for matings.

At 6 days post-eclosion, males were introduced to all the female social group

treatment vials for 2h at a density of 3:2 males:females (or 2:1 for isolated females) to enable

choice (Fig. S1). To ensure that this was a set up that would result in all females successfully

mating within the 2h period allowed, we separately recorded female mating frequency in an

identical experimental set up. In this we observed that 97.3% of females mated, with no

significant difference in mating frequency between treatments (X^2 = 1.79, d.f. = 3, p = 0.617).

After the 2h mating period in the main experiment, we transferred all mated females

to fresh standard vials to observe egg laying decisions. Every 2 – 3h, we counted the number

of eggs laid, number of egg clusters, and size of egg clusters. Observations were made at

14:00, 16:00, 19:00, and 22:00 on day 1 of laying, and 10:00 and 12:00 on day 2 (i.e., 2, 4, 7,

10, 22, and 24h after the end of the mating period, respectively). Differences in counts

(number of eggs, number of clusters, and size of clusters) between timepoints were calculated

to give overall egg laying latencies, numbers and clustering rates (H2-H4). We calculated the

proportion of eggs in clusters at each timepoint by summing the total number of eggs in each

cluster and dividing by the overall total number of eggs laid in the vial. All females were

removed after 12:00 on day 2 and vials were then kept for 14 days, to count the total number

of eclosed adults from each of the social exposure treatments (H5).

Testing for mixed maternity of egg clusters (H6)

To determine whether egg clusters were of single or mixed maternity when more than one female was present (H6), we conducted an additional experiment, to test the egg clustering patterns of groups of 4 females. In each treatment group there was one wild type (Dahomey) focal and 3 non-focal females from the *scarlet* strain (recessive *scarlet* eye colour mutation backcrossed into the Dahomey wild type > 4 times). These non-focal *scarlet* females were fed Sudan Black B dye that stained laid eggs, allowing the eggs of focal versus non-focal females to be identified. First instar larvae of Dahomey wild type and *scarlet* strains were raised in cultures of 100 larvae per vial as described above. Focal Dahomey females were collected from cultures reared and maintained on standard medium, whereas *scarlet* females were derived from cultures containing Sudan Black B dye (100g brewer's yeast, 50g sugar, 15g agar, 30ml Nipagin solution, 3ml propionic acid, and 1.4g Sudan Black B powder (*Sigma-Aldrich,* cat#199664) dissolved in 14ml corn oil (*Mazola* 100% pure corn oil) per litre of medium). Focal Dahomey virgin females were collected from the standard medium vials, and non-focal *scarlet* virgin females were collected from Sudan Black B vials and maintained on that same medium until use. Upon collection, all females were housed in groups of 10 and then Dahomey focal females were housed in groups of 4 from 5 days post-eclosion. At 4 days post-eclosion, *scarlet* females were wing clipped for identification, and also housed 4 per vial. Dahomey and *scarlet* males for matings were collected from standard medium vials and housed in groups of 10 until use.

Seven days post-eclosion, males were transferred into the female vials for mating: 8 Dahomey males were added to each group of 4 Dahomey females, and 6 *scarlet* males for each group of 4 *scarlet* females. The flies were given 4 hours to mate.

One female from each of the Dahomey vials was then transferred to fresh standard media alongside three *scarlet* dye-fed females at 12:00 GMT (N = 29). We then counted the number of eggs laid at 10.5h post-mating. Eggs were categorised as clustered or not clustered (Fig. S1) and according to whether they were laid by a focal or non-focal female. This allowed us to calculate the proportion of eggs that were clustered, and whether the cluster contained eggs from both focal and non-focal females.

Effects of existing egg cluster sizes on female egg-clustering decisions (H7)

To assess whether the size of egg clusters already present in an environment alters

subsequent female laying decisions (H7), we conducted a final experiment in which we

presented focal females with a laying environment that already contained varying numbers

of eggs and egg clusters. To create the egg clusters, singly laid eggs were collected from SYA

medium-filled Petri dishes and transferred into vials to create four different egg cluster sizes:

1, 4, 7, and 10 eggs per cluster. Females have a strong tendency to lay eggs at vial edges (Table

S2; (9, 47, 48)), hence we mimicked this tendency by placing all egg clusters at the edge of the

egg treatment vials.

To obtain the Dahomey females and males for the experiment, we collected virgin flies

from standard density vials. Prior to the tests, females were maintained alone, males in

groups of four per vial. At seven days post-eclosion, males and females were paired and

observed to ensure they had all successfully mated. After mating, groups of 4 females were

then transferred to each of the 4 types of egg cluster treatment vials (1 egg: N = 20 vials; 4

eggs: N = 17 vials; 7 eggs: N = 20 vials; 10 eggs: N = 21 vials). After the introduction of the focal

females, vials were checked at 30 min intervals from 13:00 until 21:30, at which point if

females had not laid eggs they were removed. If eggs were laid by the introduced focal

females, we counted the number of eggs laid and categorised them in relation to the existing

egg cluster present (clustered or not clustered). All vials were then frozen at -20°C for imaging.

Vials were imaged to record inter egg distances, using a video camera (Sony Handycam

HDR-CX405). Using ImageJ's multiple point selector tool (50) we captured x-y coordinates of

each egg and converted these into Euclidean pairwise distances. Selected coordinates were

taken from the edge points of eggs that were in closest proximity, resulting in clustered eggs

having a distance value of zero.

Statistical analysis

All statistical analyses were conducted using R v 4. 2. 2 (51), and mean values and standard

deviations are reported within the text. Graphs were produced using packages 'ggplot2',

'cowplot' and 'grid' (52, 53), with plotted standard errors calculated using 'Rmisc' (54).

We analysed the effect of egg and adult social density on the likelihood that females
laid eggs and produced adult offspring, by using generalised linear models (GLMs) with
binomial errors. We used the package 'AER' (55) to test for overdispersion, and where that
was found, we used GLMs with a quasi-Poisson distribution to account for this over-dispersion
and analyse the effect of adult density on the number of offspring produced. Linear models
were used to investigate the effect of adult density on egg-clustering patterns (egg laying and
clustering rates), and total eggs laid was included as a fixed effect to account for increased
likelihood of clustering when more eggs are present in the environment.

When analysing variables throughout multiple timepoints, we used linear mixed
effects models; the timepoint was included as a fixed effect in the model to account for
differences in rates and the test vial was included as a random effect – using functions in
packages 'lme4' (56) and 'lmerTest' (57). We tested for Pearson's product-moment
correlations between the proportion of eggs laid in clusters and subsequent egg-adult
viability. In all cases, post-hoc tests (models including only pairwise comparisons) were used
to identify significant differences in the six pairwise comparisons available.

In the first experiment investigating the effects of social density, there was variation
in sampling effort between treatments, due to the differences in female group sizes (and lack
of identification of focal females). To account for this, all significant results from the initial
analyses were reanalysed with a randomised subset of data so that the number of individual
flies (rather than number of vials) per treatment were similar. In this reduced subset, all vials
from solitary females were included (N = 30 females), half of those in pairs (N = 30 females),
eight of those from groups of 4 (N = 32 females), and four of those from groups of 8 (N = 32
females). These analyses were congruent with those of the initial analyses, suggesting the
initial analyses were robust.

Further details on the model syntax for all analyses and all data are available from the Dryad
Digital Repository (10.5061/dryad.mgqnk9971).

Results

Egg laying patterns of females were significantly non-random (H1)

Across the four social treatments, clustering preferences (values of K that, based on the

Kolmogorov-Smirnov test, yielded no significant difference between the simulated and

observed distributions) were between 0.3 and 0.5 (Table S1).

For all social treatments with two or more females, patterns of egg clustering in the

empirical data were significantly more clustered than those predicted by the null model (H1;

Wilcoxon Signed Rank test comparisons with simulations using K = 0; paired: $z(6) = -2.29$,

$p = 0.0220$, $r = 0.866$, N = 7 vials; groups of four: $z(25) = -4.47$, $p = 7.95 \times 10^{-6}$, $r = 0.876$, N = 26

vials; groups of eight: $z(27) = 4.63$, $p = 3.63 \times 10^{-6}$, $r = 0.875$, N = 28 vials). Therefore, we

conclude that the egg-clustering patterns of all grouped females followed a non-random

distribution (Table S1 and Fig. 1). However, solitary females did not cluster their eggs

significantly more than the null model predicted ($z(2) = -1.36$, $p = 0.174$, $r = 0.786$, N = 3 vials),

suggesting that their laying patterns are random – although it is worth noting that this effect

could be due to low power as clustering preference values are similar to the grouped

treatments. Clustering preferences also varied among social treatments: solitary and paired

females had a significantly lower clustering preference in comparison to females housed in

groups of 4 or 8 (Table S1 and Fig. 1). The findings support H1, that eggs are laid in non-

random patterns. In addition, the results suggest that egg clustering patterns are plastic and

responsive to the female's social environment.

Figure 1. **Egg clustering is non-random and increases with increasing adult social density.** (a) Comparison of the clustering proportion model outputs and empirical egg laying patterns observed in *D. melanogaster,* for four social group sizes (1 (pink), 2 (green), 4 (light blue), and 8 (dark blue) adult females per group). The average Kolmogorov-Smirnov P values at K (clustering preference) values ranging from 0 to 1 in 0.1 increments – where the null distribution is K = 0 (no eggs clustered), and K = 1 (all eggs laid in one large cluster). The best-fit values of K are those that produced the least significant (highest P value) difference between simulated and observed data. (b-e) The Kolmogorov-Smirnov P values for individual replicates in the four social group sizes (1, 2, 4, and 8 adult females), with the values for each vial shown in the pale, thin lines and the average value indicated with a bold line. The vertical grey dotted lines show clustering preference values.

Females in larger social groups laid eggs at a faster rate (H2), laid more eggs (H3), and clustered their eggs more (H4), but showed no differences in fitness (egg-adult survival, H5)

In line with H2 and H3, females in larger social groups were significantly more likely to lay eggs (X^2 = 20.2, d.f. = 116, p = 1.51 x 10^{-4}). Eggs were present in 100% of vials housing eight females, a significantly higher proportion than for vials containing a solitary female (66.7%, X^2 = 15.9, d.f. = 58, p = 6.76 x 10^{-5}) or those with two females (83.3%, X^2 = 7.39, d.f. = 58, p = 0.00657). There were also significantly more vials containing eggs from the four female treatment (96.7%) in comparison to the solitary female treatment (X^2 = 10.2, d.f. = 58, p = 0.00140).

Females in larger social groups also laid more eggs per female than did the females from the other treatments (solitary females: 7 ± 10 eggs per female (N = 30 vials); pairs: 9 ± 9 eggs per female (N = 30 vials); groups of four: 18 ± 8 eggs per female (N = 30 vials); groups of eight: 19 ± 5 eggs per female (N = 30 vials); $F_{3, 116}$ = 265, p = 2.93 x 10^{-8}; Fig. 2a). Consistent

with this, post-hoc tests showed that females in solitary and paired social densities laid fewer

eggs than females in groups of four and eight (all $p \ll 0.001$). To control for the increased

sampling effort in treatments with larger groups of females, the analysis was repeated with a

randomised subset of data to make number of flies (rather than the number of vials) in each

treatment similar. The results of these analyses were comparable with the main analysis

(larger social groups showed significantly higher likelihood of egg laying: $X^2 = 50.0$, d.f. = 54,

$p = 0.0271$ and significantly higher numbers of eggs laid: $F_{3, 54} = 137$, $p = 0.00427$).

An analysis of the data at the final timepoint (24h post-mating) supported H4 and

showed that females housed in larger groups laid a higher proportion of their eggs in clusters

(solitary females: 22.6% ± 26.9% (N = 20 vials); pairs: 38.1% ± 28.9% (N = 25 vials); groups of

four: 62.7% ± 19.2% (N = 29 vials); groups of eight: 73.7% ± 13.3% (N = 30 vials); total number

of eggs laid included in model: $F_{3, 99} = 28.4$, $p = 2.53 \times 10^{-13}$; all pairwise comparisons $p < 0.05$;

Fig. 2b). We included the total number of eggs laid in the model as we found that there was

a positive relationship between the number of eggs and the proportion of eggs that were

clustered ($F_{1, 99} = 6.66$, $p = 0.0113$), although this was not the key driver of social density

effects on clustering proportions. In all grouped female treatments, the proportion of

clustering also increased over time (individual vial included in model as a random effect: pairs:

$F_{1, 58} = 19.1$, $p = 5.26 \times 10^{-5}$; groups of four: $F_{1, 95} = 75.8$, $p = 9.99 \times 10^{-14}$; groups of eight:

$F_{1, 111} = 74.7$, $p = 4.45 \times 10^{-14}$). There was no significant effect of time on the proportion of

clusters in eggs laid by solitary females ($F_{1, 40} = 1.15$, $p = 0.289$).

Analyses of the number of clusters and size of the largest cluster followed a similar

pattern and both increased with increasing social density (number: $F_{3, 376} = 463$,

$p = 2.20 \times 10^{16}$; size: $F_{3, 299} = 58.3$, $p = 2.20 \times 10^{-16}$; Figs. S2-3), and over time (number:

$F_{1, 376} = 631$, $p = 2.20 \times 10^{-16}$; size: $F_{1, 299} = 67.1$, $p = 7.62 \times 10^{-15}$). These results were again

robust to differences in sampling effort. When the analysis was repeated using the same

number of individuals (rather than vials) at the final timepoint, it was still evident that females

in larger groups laid a higher proportion of their eggs in clusters ($F_{3, 41} = 6.40$, $p = 0.00117$) and

that the number and size of clusters still increased with increasing social density (number: $F_{3,}$

$_{134} = 199$, $p = 2.20 \times 10^{16}$; size: $F_{3, 102} = 25.1$, $p = 2.96 \times 10^{-12}$).

The only hypothesis not supported was of a fitness benefit for eggs laid in clusters in

terms of increased egg-adult survival (H5; Fig 2c). There was no effect of social density (1, 2,

4, or 8 females) on the proportion of eggs that survived to adult stage ($F_{3, 99}$ = 0.292, p = 0.831;

Fig. S4) and we found no correlation between the proportion of eggs laid in clusters and egg-

adult offspring viability in these densities (t = -0.485, d.f. = 101, p = 0.629; Fig. 2c).

Figure 2. **Females held in larger social groups laid more eggs, more quickly, and in a higher proportion of**

clusters, although this did not impact egg-adult survival. (a) Females housed in groups of four and eight were

quicker to lay, and laid more eggs per female, than those kept in solitude or in pairs. Eggs laid per individual

were calculated by dividing total number of observed eggs by the number of females in the vial. Social treatment

means and standard errors are shown for the four housing densities at each observation time point. The grey

box indicates period of dark (21:00 – 09:00 GMT). (b) Females housed in groups of four or eight laid a higher

proportion of their eggs in clusters compared to those kept in solitude or in pairs. Proportion of eggs clustered

was calculated by summing all cluster sizes and dividing by the total number of eggs counted. (c) There was no

effect of the proportion of egg clustering on egg-adult viability (linear regressions with 95% confidence intervals).

Overall, the results supported hypotheses 2, 3 and 4 and show that females respond

plastically to their social environment and lay eggs more quickly, lay more eggs and lay more

eggs in clusters, as the social group size increases. However, there was no indication that eggs

laid in clusters benefitted via increased egg-adult viability (H5).

Egg clusters are usually of mixed-maternity (H6)

In the groups of 4 females comprising 1 standard-fed focal wild type and 3 Sudan Black B dye-

fed non-focals, we observed that of the focal eggs that had been laid in clusters, a mean of

79.5% of them were in mixed-maternity clusters. This supports the hypothesis (H6) that eggs

in clusters are usually of mixed maternity.

Females prefer to lay their eggs within existing clusters rather than with singly laid eggs (H7)

In the final experiment in which we manipulated egg cluster sizes, we found that females

were more likely to add at least one of their eggs to existing clusters of 4, 7, or 10 eggs, than

they were to lay with a singly placed existing egg (X^2 = 12.5, d.f. = 156, p = 4.18 x 10^{-4}; Fig. 3a).

The total proportion of eggs laid that joined existing eggs was also higher when existing eggs

were clustered ($F_{3, 154}$ = 4.75, p = 0.00340; Fig. 3b). Proportions were calculated from the

number of eggs females were able to lay within the 30 min observation window (between 1

– 19 eggs per vial). Post-hoc tests showed that egg cluster sizes of 4 and 10 were joined

significantly more often than were single eggs (four: $F_{1, 71}$ = 13.0, p = 5.85 x 10^{-4}; ten: $F_{1, 78}$ =

8.76, p = 0.00407), though this trend was marginally non-significant for a cluster size of 7 ($F_{1,}$

$_{75}$ = 3.82, p = 0.0542).

Figure 3. **Females lay more eggs within existing clusters and lay eggs closer to existing egg clusters** (a) A higher proportion of females clustered at least one of their eggs with existing eggs, when those existing eggs were in clusters compared to singly laid eggs. Blue = singly laid eggs; pink = clustered eggs. (b). Females laid a higher proportion of their eggs in clusters with existing eggs, when those existing eggs were already in a cluster (of any size) compared to a singly laid egg. Means and standard errors are shown for the four treatment egg cluster sizes. Significant differences between treatment egg cluster sizes are represented by an overarching bar; * indicates a significant difference between paired treatments (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). (c). Mean distance of laid eggs from existing treatment eggs was greater when the number of existing eggs was one, compared to egg clusters of any size.

In addition to scoring whether a laid egg was physically in contact with an existing egg (i.e. clustered), we also measured the distance between each laid egg and treatment eggs already present on the substrate. This is because there could be benefits of laying eggs near to other eggs without the need to be in direct contact, and so we wanted to ensure we did not exclude nearby eggs by only scoring physically touching eggs. Females laid their eggs closer to existing clusters (H6; $F_{3, 145} = 9.86$, $p = 5.86 \times 10^{-6}$; Fig. 3c) with all three cluster sizes having shorter inter-egg distances than eggs from females exposed to single eggs (four: $F_{1, 67} = 14.8$, $p = 2.68 \times 10^{-4}$; seven: $F_{1, 70} = 11.4$, $p = 0.00122$; ten: $F_{1, 70} = 23.8$, $p = 6.57 \times 10^{-6}$). Hence, laid eggs that did not strictly join a cluster were still laid closer to existing clusters than to single eggs. The results are consistent with the finding that females respond plastically to the egg clustering patterns of eggs already present in an environment and lay their eggs preferentially within existing egg clusters (H7).

Discussion

The main findings were that females responded plastically to their social environment and to the presence of existing eggs within their oviposition environment. We found that egg laying patterns were significantly non-random (H1), except when females were held in social isolation. As the social group size of females increased, females laid eggs more quickly (H2), laid more eggs (H3), and laid more of them in clusters (H4). However, there was no evidence of a fitness benefit to clustered eggs in increased egg-adult viability (H5). Clustered eggs were usually of mixed maternity (H6) and though females did not prefer to lay their eggs within larger existing clusters in the environment, they did prefer egg laying in clusters overall (H7). Therefore, the majority of the hypotheses tested were supported and show that females show striking plasticity in their egg placement decisions, with a strong bias towards laying eggs in mixed maternity clusters. However, the fitness benefits of such behaviour remain elusive. These results are explored in more detail, below.

Egg-clustering decisions are plastic and non-random (H1)

When they are held in groups, we found that females were significantly more likely to cluster eggs than was predicted by the null model, demonstrating that they clustered their eggs more than expected by chance (H1). In contrast, solitary females laid eggs in patterns not significantly different from random. This provides evidence that females make active clustering decisions, and that egg-clustering responses are likely to be plastic and change in response to varying social environments. Increased clustering has the potential to increase local population density, especially at larval stages due to the limited mobility of this life stage in comparison to that of adults. This has the potential to have cumulative effects, as even relatively small-scale variation in density leading to increased competition may have consequences for fitness (58).

Presence of adult conspecifics increased the rate of oviposition and total number of eggs laid (H2, H3)

In larger groups, females laid more eggs and laid them more quickly, which is consistent with existing research (32) (H2-3). Following raffle theory, producing more eggs when competition is high ensures that mothers have a greater chance of at least some of their offspring surviving to adulthood (46). With a greater number of conspecifics present in the environment, there

should also be a higher chance of copying behaviours arising due to an increased number of

potential 'initiators' (37, 38, 40). Increased copying could increase laying rates, benefitting

mothers as it may reduce sampling times needed to locate optimal oviposition sites and thus

any associated costs. As predicted, in addition to the overall egg laying rate being faster, the

latency to lay the first egg (a stage that is required to initiate any such copying behaviours)

was also shorter at higher adult densities. However, in situations in which larval density is

high and resources are limiting, it is likely that later-laid eggs will have reduced survival due

to resource degradation and cannibalisation (13, 32, 59). Thus, in high density environments,

mothers should ensure they begin laying quickly to avoid increased offspring lethality.

Previous research (32) showed that light can inhibit egg laying in *Drosophila*, an effect

that can be over-ridden by being in a group (five females). Intriguingly, our results show

remarkably similar laying rates for isolated and paired females (whereas groups of four or

eight females matched laying times reported in (32)), suggesting that exposure to only one

other gravid female does not counteract the light-induced inhibition of egg production. This

is surprising given that in male *Drosophila*, exposure to one other rival is enough to facilitate

reproductive behavioural responses (60-62). This suggests that there are sex differences in

responses to local population density.

Females cluster more of their eggs in the presence of adult conspecifics (H4), and more commonly

form mixed-maternity clusters (H6)

As social density increased, proportion of eggs laid in clusters increased. Although there was

a higher chance of clustering when there are more eggs present in the environment, this

increase in clustering with increasing social density did not depend upon egg number (in

agreement with the null model comparison). It was not possible to determine the maternity

of the clusters in grouped treatments in our initial experiment, but our later experiment

(testing H6) showed that the clusters are often of mixed maternity. Clusters containing eggs

from multiple mothers could contain more diverse nutrients and microbiomes (63, 64) and

have a reduced risk of consisting of only cheaters under a potential public goods system (29,

30). These benefits would not be available to isolated females laying their eggs in single

maternity clusters, and they would only incur increased risks of sibling competition if food is

limited. Females may be opting for a bet-hedging strategy of laying in multiple sites thus

minimising predation/parasitism risk both by reducing transmission and egg visibility (65),

whilst adding to existing clusters to reap the benefits of any diffusible compounds present –

e.g., anticannibalism pheromones (28). It is not yet known what other compounds *Drosophila*

eggs may contain, but it is possible that they are also coated with compounds that minimise

transfer of pathogens such as antimicrobials (as seen in medfly (26)). If true, this could be a

further benefit of clustering decisions.

Females preferred to lay their eggs in existing egg clusters (H7)

To understand if females had a preference for existing cluster size, we manipulated exposure

to egg clusters via manual transfer of eggs. Our results were not consistent with the

hypothesis that females would show a preference for adding their eggs to larger clusters

compared to smaller clusters. Instead, our results suggest that females showed a strong

preference for laying with clusters of any size compared to with single eggs, by choosing to

cluster their eggs with existing clusters more frequently and by laying their eggs closer to

those clusters. The absence of a preference for adding eggs to larger clusters could be because

(i) there is no benefit to increasing cluster size (i.e., benefits to joining clusters are not

additive), or (ii) there is some trade-off to the benefits of clusters of these sizes. For example,

with larger clusters of eggs, there could be a higher chance that at least some eggs are older,

and given that eggs laid later are at a higher risk of cannibalisation (28, 59) the risk/cost of

this increases as cluster size increases.

We have shown that clusters are very often of mixed-maternity (at least 79%)

suggesting that there are greater benefits to females if eggs in an existing cluster have been

laid by more than one other female. It is possible that mothers could identify maternity of egg

clusters due to pheromones used to identify conspecific presence (41, 66, 67), and thus do

not rely on cluster size for information regarding maternity. Even if this were not true,

increased cluster size would not be a mixed maternity benefit in our experiment as in our

cluster size manipulations we ensured that all clusters were of mixed-maternity. Another

reason that we did not detect additive effects, could be that the cluster sizes chosen were not

sufficiently large enough to elicit benefits of larger cluster sizes. More work needs to be

completed to investigate whether females have a preference for egg clusters of any particular

size or if their choices are based upon other characteristics.

The strong aversion to isolated eggs that we observed raises the question of how

clusters form if females are only willing to lay eggs with existing clusters. It is possible that

when copying laying behaviours, females are searching for evidence of a repeated decision –

i.e., at least two eggs at the same site. Otherwise, a solitary egg could be present only because

the mother could no longer retain it, despite not finding an optimal site. It is not known if

females can detect whether an egg is infertile (although if laid by a virgin female pheromonal

cues differ (41)). However, if females cannot detect fertility, an isolated egg could be infertile

and thus could provide fewer clustering benefits. This result might suggest that in order for

clusters to form, the first two eggs need to be laid by the same mother before other gravid

females are induced to add to a cluster. However, the fact that clusters of two eggs laid by

different mothers were observed in our mixed-maternity investigation suggests further

studies on clustering initiation are required.

Contrary to our hypothesis of fitness benefits (H7), egg-adult viability did not increase

with increasing cluster proportions. It is not clear why we did not observe fitness effects, given

costs of maintaining plasticity (68, 69). It is possible that as mothers were able to choose a

clustering strategy, all individuals chose the optimal strategy and thus there were no negative

consequences associated with the differing clustering proportions. To test this, future work

could force mothers into making an incorrect choice via a mismatch of environments. It is also

possible that the *ad libitum* food provided obscured any costs of choosing the 'wrong' egg

laying strategy. It was necessary to remove females from the egg laying environments before

they had oviposited all fertile eggs, as eggs needed to be counted prior to larvae hatching

which takes place after 22 – 24 hours at 25°C (70, 71). Thus, it is also possible that differences

in egg-adult viability could have been detected if females were able to continue laying if the

benefit only occurs in clusters of mixed-age eggs.

Conclusion

Overall, we demonstrated that *D. melanogaster* mothers express non-random plasticity in

egg-clustering decisions and respond to fluctuations in social density from multiple life stages.

Exposure to increased adult social density resulted in females laying more eggs, at a faster

rate, and laying a higher proportion of those eggs in clusters. Exposure to existing egg clusters

also led to a higher frequency of clustering decisions, regardless of existing cluster size.

However, the potential costs and benefits of this plasticity and its overall ecological

importance, remain to be discovered. These results demonstrate a clear importance of egg

location decisions as well as the overall quantity of eggs produced. This could have strong

relevance in nature given that spatial distribution has significant ecological importance in

wild-living populations.

References

1.   West-Eberhard MJ. Phenotypic plasticity and the origins of diversity. Annual Review of Ecology and Systematics. 1989;20(1):249-78.

2.   Komers PE. Behavioural plasticity in variable environments. Canadian Journal of Zoology. 1997;75(2):161-9.

3.   Birkhead T, Møller A. Female control of paternity. Trends in Ecology & Evolution. 1993;8(3):100-4.

4.   Rodriguez-Enriquez CL, Tadeo E, Rull J. Elucidating the function of ejaculate expulsion and consumption after copulation by female *Euxesta bilimeki*. Behavioral Ecology and Sociobiology. 2013;67(6):937-46.

5.   Ward PI. Females influence sperm storage and use in the yellow dung fly *Scathophaga stercoraria*. Behavioral Ecology and Sociobiology. 1993;32(5):313-9.

6.   Elgar MA, Schneider JM, Herberstein ME. Female control of paternity in the sexually cannibalistic spider *Argiope keyserlingi*. Proceedings of the Royal Society of London Series B: Biological Sciences. 2000;267(1460):2439-43.

7.   Passera L, Aron S, Vargo EL, Keller L. Queen control of sex ratio in fire ants. Science. 2001;293(5533):1308.

8.   Sato A, Karino K. Female control of offspring sex ratios based on male attractiveness in the guppy. Ethology. 2010;116(6):524-34.

9.   Churchill ER, Dytham C, Bridle JR, Thom MDF. Social and physical environment independently affect oviposition decisions in *Drosophila*. Behavioral Ecology. 2021.

10.   Fowler EK, Leigh S, Rostant WG, Thomas A, Bretman A, Chapman T. Memory of social experience affects female fecundity via perception of fly deposits. BMC Biology. 2022;20(1):244.

11.   Takasuka A, Yoneda M, Oozeki Y. Disentangling density-dependent effects on egg production and survival from egg to recruitment in fish. Fish and Fisheries. 2019;20(5):870-87.

12.   Matsushima N, Kawata M. The choice of oviposition site and the effects of density and oviposition timing on survivorship in *Rana japonica*. Ecological Research. 2005;20(1):81-6.

13.   Khodaei L, Long TAF. Kin recognition and egg cannibalism by *Drosophila melanogaster* larvae. Journal of Insect Behavior. 2020;33(1):20-9.

14.   Perry JC, Roitberg BD. Games among cannibals: Competition to cannibalize and parent-offspring conflict lead to increased sibling cannibalism. Journal of Evolutionary Biology. 2005;18(6):1523-33.

15.   Kestemont P, Jourdan S, Houbart M, Mélard C, Paspatis M, Fontaine P, et al. Size heterogeneity, cannibalism and competition in cultured predatory fish larvae: Biotic and abiotic influences. Aquaculture. 2003;227(1):333-56.

16.   Novčić I, Aleksić D, Vidović Z. Copying of diving behavior in the Eurasian Coot *Fulica atra*. Journal of Ornithology. 2022;163(2):523-9.

17. Cocroft RB. Vibrational communication facilitates cooperative foraging in a phloem-
feeding insect. Proceedings of the Royal Society B: Biological Sciences. 2005;272(1567):1023-
9.

18. Fitzgerald TD, Peterson SCCFpdJ. Cooperative foraging and communication in
caterpillars. BioScience. 1988;38(1):20-5.

19. Smith CC, Sargent RC. Female fitness declines with increasing female density but not
male harassment in the western mosquitofish, *Gambusia affinis*. Animal Behaviour.
2006;71(2):401-7.

20. Zhao L, Wang X, Zhan J. Effects of female and male density on their mating
performance and female post-mating reproductive fitness in *Plagiodera versicolora*
(Laicharting) (Coleoptera: Chrysomelidae). The Coleopterists Bulletin. 2022;76(1):73-81.

21. Welsh DP, Fuller RC. Where to place your eggs: The effects of conspecific eggs and
water depth on oviposition decisions in bluefin killifish. Journal of Zoology. 2011;284(3):192-
7.

22. Riehl C. Evolutionary routes to non-kin cooperative breeding in birds. Proceedings of
the Royal Society B: Biological Sciences. 2013;280(1772):20132245.

23. Doody JS, Freedberg S, Keogh JS. Communal egg-laying in reptiles and amphibians:
Evolutionary patterns and hypotheses. The Quarterly Review of Biology. 2009;84(3):229-52.

24. Courtney SP. The evolution of egg clustering by butterflies and other insects. The
American Naturalist. 1984;123(2):276-81.

25. Ioannou C. Grouping and predation. In: Shackelford TK, Weekes-Shackelford VA,
editors. Encyclopedia of Evolutionary Psychological Science. Cham: Springer International
Publishing; 2017. p. 1-6.

26. Marchini D, Marri L, Rosetto M, Manetti AG, Dallai R. Presence of antibacterial
peptides on the laid egg chorion of the medfly *Ceratitis capitata*. Biochemical and biophysical
research communications. 1997;240(3):657-63.

27. Eisner T, Eisner M, Rossini C, Iyengar VK, Roach BL, Benedikt E, et al. Chemical defense
against predation in an insect egg. Proceedings of the National Academy of Sciences.
2000;97(4):1634-9.

28. Narasimha S, Nagornov KO, Menin L, Mucciolo A, Rohwedder A, Humbel BM, et al.
Drosophila melanogaster cloak their eggs with pheromones, which prevents cannibalism.
PLOS Biology. 2019;17(1):e2006012.

29. Olson M. The theory of collective action: Public goods and the theory of groups.
Harvard UniversityPress, Cambridge. 1965.

30. Archetti M. Cooperation as a volunteer's dilemma and the strategy of conflict in public
goods games. Journal of Evolutionary Biology. 2009;22(11):2192-200.

31. Archetti M, Scheuring I, Yu D. The non-tragedy of the non-linear commons. Preprints:
Preprints; 2020.

32. Bailly TPM, Kohlmeier P, Etienne RS, Wertheim B, Billeter J-C. Social modulation of
oogenesis and egg-laying in *Drosophila melanogaster*. Current Biology. 2023;33(14):2865-
77.e4.

33. Soto-Yéber L, Soto-Ortiz J, Godoy P, Godoy-Herrera R. The behavior of adult
Drosophila in the wild. PLOS ONE. 2018;13(12):e0209917.

34. Schwartz NU, Zhong L, Bellemer A, Tracey WD. Egg laying decisions in *Drosophila* are
consistent with foraging costs of larval progeny. PLOS ONE. 2012;7(5):e37910.

35. Lihoreau M, Clarke IM, Buhl J, Sumpter DJT, Simpson SJ. Collective selection of food
patches in *Drosophila*. The Journal of Experimental Biology. 2016;219(5):668.

36. Zhang L, Sun H, Grosse-Wilde E, Zhang L, Hansson BS, Dweck HKM. Cross-generation
pheromonal communication drives *Drosophila* oviposition site choice. Current Biology.
2023;33(10):2095-103.e3.

37. Golden S, Dukas R. The value of patch-choice copying in fruit flies. PLOS ONE.
2014;9(11):e112381.

38. Malek HL, Long TAF. On the use of private versus social information in oviposition site
choice decisions by *Drosophila melanogaster* females. Behavioral Ecology. 2020.

39. Battesti M, Moreno C, Joly D, Mery F. Spread of social information and dynamics of
social transmission within *Drosophila* groups. Current Biology. 2012;22(4):309-13.

40. Sarin S, Dukas R. Social learning about egg-laying substrates in fruitflies. Proceedings
of the Royal Society B: Biological Sciences. 2009;276(1677):4323-8.

41. Duménil C, Woud D, Pinto F, Alkema JT, Jansen I, Van Der Geest AM, et al. Pheromonal
cues deposited by mated females convey social information about egg-laying sites in
Drosophila melanogaster. Journal of Chemical Ecology. 2016;42(3):259-69.

42. Dombrovski M, Poussard L, Moalem K, Kmecova L, Hogan N, Schott E, et al.
Cooperative behavior emerges among *Drosophila* larvae. Current Biology. 2017;27(18):2821-
6.e2.

43. Fowler EK, Leigh S, Bretman A, Chapman T. Plastic responses of males and females
interact to determine mating behavior. Evolution. 2022;76(9):2116-29.

44. Yang C-h, Belawat P, Hafen E, Jan LY, Jan Y-N. *Drosophila*; Egg-laying site selection as
a system to study simple decision-making processes. Science. 2008;319(5870):1679.

45. Vijayan V, Wang Z, Chandra V, Chakravorty A, Li R, Sarbanes SL, et al. An internal
expectation guides *Drosophila* egg-laying decisions. Science Advances.8(43):eabn3852.

46. Parker GA. Sperm competition games: Raffles and roles. Proceedings of the Royal
Society B: Biological Sciences. 1990;242(1304):120-6.

47. Chess KF, Ringo JM. Oviposition site selection by *Drosophila melanogaster* and
Drosophila simulans. Evolution. 1985;39(4):869-77.

48. Moore JA. Competition between *Drosophila melanogaster* and *Drosophila simulans*. I.
Population cage experiments. Evolution. 1952;6(4):407-20.

49. Wolfram Research Inc. Mathematica. Champaign, Illinois: Wolfram Research, Inc.;
2023.

50. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nature methods. 2012;9(7):671-5.

51. R Core Team. R: A language and environment for statistical computing. 2019.

52.   Wickham H. ggplot2: Elegant graphics for data analysis: Springer-Verlag New York;
2016.

53.   Wilke CO. cowplot: Streamlined plot theme and plot annotations for 'ggplot2': R
package version 1.1.1; 2020.

54.   Hope RM. Rmisc: Ryan miscellaneous: R package version 1.5.1; 2022.

55.   Kleiber C, Zeileis A. Applied econometrics with R: Springer-Verlag New York; 2008.

56.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using
lme4. Journal of Statistical Software; Vol 1, Issue 1 (2015). 2015.

57.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed
Effects Models. Journal of Statistical Software. 2017;1(13).

58.   Churchill ER, Bridle JR, Thom MDF. Spatially clustered resources increase male
aggregation and mating duration in *Drosophila melanogaster*. Animal Behaviour.
2020;169:45-50.

59.   Vijendravarma RK, Narasimha S, Kawecki TJ. Predatory cannibalism in *Drosophila*
melanogaster larvae. Nature Communications. 2013;4(1):1789.

60.   Bretman A, Fricke C, Chapman T. Plastic responses of male *Drosophila melanogaster*
to the level of sperm competition increase male reproductive fitness. Proceedings of the
Royal Society B: Biological Sciences. 2009;276(1662):1705-11.

61.   Moatt JP, Dytham C, Thom MDCP. Exposure to sperm competition risk improves
survival of virgin males. Biol Lett. 2013;9(2):20121188.

62.   Bretman A, Westmancoat JD, Chapman T. Male control of mating duration following
exposure to rivals in fruitflies. Journal of Insect Physiology. 2013;59(8):824-7.

63.   Guilhot R, Xuéreb A, Lagmairi A, Olazcuaga L, Fellous S. Microbiota acquisition and
transmission in *Drosophila* flies. iScience. 2023;26(9):107656.

64.   Bakula M. The persistence of a microbial flora during postembryogenesis of
Drosophila melanogaster. Journal of Invertebrate Pathology. 1969;14(3):365-74.

65.   Turner GF, Pitcher TJ. Attack abatement: A model for group protection by combined
avoidance and dilution. The American Naturalist. 1986;128(2):228-40.

66.   Bartelt RJ, Schaner AM, Jackson LL. cis-Vaccenyl acetate as an aggregation pheromone
in *Drosophila melanogaster*. Journal of Chemical Ecology. 1985;11(12):1747-56.

67.   Verschut TA, Ng R, Doubovetzky NP, Le Calvez G, Sneep JL, Minnaard AJ, et al.
Aggregation pheromones have a non-linear effect on oviposition behavior in *Drosophila*
melanogaster. Nature Communications. 2023;14(1):1544.

68.   Auld JR, Agrawal AA, Relyea RA. Re-evaluating the costs and limits of adaptive
phenotypic plasticity. Proceedings of the Royal Society B: Biological Sciences.
2009;277(1681):503-11.

69.   DeWitt TJ. Costs and limits of phenotypic plasticity: Tests with predator-induced
morphology and life history in a freshwater snail. Journal of Evolutionary Biology.
1998;11(4):465-80.

70.   Markow TA, Beall S, Matzkin LM. Egg size, embryonic development time and
ovoviviparity in *Drosophila* species. Journal of Evolutionary Biology. 2009;22(2):430-4.

71.   Fernández-Moreno MA, Farr CL, Kaguni LS, Garesse R. *Drosophila melanogaster* as a
model system to study mitochondrial biology. Methods in molecular biology (Clifton, NJ).
2007;372:33-49.

Funding

This study was funded by NERC (NE/T007133/1).

Author Contributions

Conceptualisation: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, Marco Archetti, Douglas
W. Yu, Andrew F.G. Bourke, Tracey Chapman, and Amanda Bretman. *Data curation*: Emily R.
Churchill, Emily K. Fowler, and Lucy A. Friend. *Funding acquisition*: Emily K. Fowler, Marco
Archetti, Douglas W. Yu, Andrew F.G. Bourke, Tracey Chapman, and Amanda Bretman.
Investigation: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, and Marco Archetti.
Methodology: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, and Marco Archetti. *Project
administration*: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, Tracey Chapman, and
Amanda Bretman. *Resources*: Tracey Chapman and Amanda Bretman. *Software*: Emily R.
Churchill and Marco Archetti. *Visualisation*: Emily R. Churchill and Marco Archetti. *Writing –
original draft*: Emily R. Churchill. *Writing – review and editing*: Emily R. Churchill, Emily K.
Fowler, Lucy A. Friend, Marco Archetti, Douglas W. Yu, Andrew F.G. Bourke, Tracey Chapman,
and Amanda Bretman. All authors gave final approval for publication.

 Supplementary Information

 Methods

 Effects of adult social density on egg-clustering decisions (H1-5)

Figure S1. The experimental set up for testing the effects of social densities on egg laying. Day 0 – 4: males and

females housed in same-sex groups of 10. Day 5 – 6: females transferred to one of four social treatments (whilst

males remained in groups of 10). Day 6: males were added to female treatment vials in a ratio of 3:2 (2:1 for

isolated females) and given 2h to mate. Day 6 – 7: number of eggs, number of clusters and size of clusters were

counted at the six listed timepoints. Shown is the strategy for defining egg clusters (contact between the main

body of eggs).

Effects of existing egg location on egg-clustering decisions

(i) Set up of egg locations and adult density treatments

The rationale was to test isolated versus grouped females laying eggs in 3 types of 'egg

treatment' vials: no egg present (control), and 1 egg either placed in the centre or at the edge

of the vial. The non-focal eggs used to create the egg treatment vials were laid by groups of

10 gravid females allowed to lay overnight on Petri dishes filled with 40ml of SYA medium.

Only solitary eggs were chosen and transferred, using a mounted needle, to make the

experimental egg treatment vials (to reduce variation in uncontrolled factors). To enable

transfer of eggs without damage to the SYA medium, standard plastic vials were cut in half,

and the top half reattached after egg transfer using Sellotape. Test flies were collected from

standard density vials (as described above) and females were housed in solitude, and males

in groups of four. After six days, one male was transferred into the vial with each female, and

pairs were watched to ensure matings occurred. We then transferred females either alone or

in groups of 4 into vials containing the egg treatments (no egg, 1 edge egg, or 1 central egg)

described above (N = 15 - 42 vials per treatment).

(ii) Laying latency and egg location decisions

We then checked vials every 30 minutes until 21:30 (one count after lights off in the constant

environment room) and removed the female(s) if they had laid eggs. If they did not lay, they

were kept overnight, and egg counts resumed at 09:30 for three further hours, until 24 hours

after females were introduced to the vials. We calculated egg laying latencies and the number

of eggs laid by the focal females. We also noted whether eggs were laid in a cluster (defined

in Fig. S1) alongside the existing egg.

Results

Egg laying patterns of females were significantly non-random (H1)

Table S1. Comparison of the clustering proportion model outputs and empirical egg laying patterns observed in

D. melanogaster. Shown are Kolmogorov-Smirnov P value outputs for the comparison of empirical data to

expected clustering preference values (К = 0 to К = 1, in 0.1 increments) rounded to two decimal places. Values

that show the most likely clustering preference are highlighted in bold.

Social treatment	K values										
	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
Solitary N = 20 vials	0.09	0.34	0.44	**0.48**	0.46	0.29	0.17	0.06	0.02	0.03	0.01
Paired N = 25 vials	0.01	0.20	0.34	0.43	**0.43**	0.38	0.22	0.15	0.07	0.04	0.01
Groups of four N = 29 vials	0.00	0.08	0.16	0.25	**0.29**	0.29	0.23	0.13	0.05	0.02	0.00
Groups of eight N = 30 vials	0.00	0.00	0.07	0.15	0.20	**0.21**	0.15	0.08	0.03	0.01	0.00

803 *Females in larger social groups laid significantly more of their eggs in clusters (H4)*

805 Figure S2. Maximum cluster size increased with increasing female social density. Total number of vials with each
806 cluster size observed is displayed for each of the four social treatments.

808 Figure S3. A greater number of clusters were observed with increasing female social density. Total number of
809 clusters per vial is displayed for each of the four social treatments.

 Fitness effects of social environment (H5)

Figure S4. There was no effect of social group size on egg-adult viability. Means (large dot) and standard errors
are shown in black.

Edge effect egg laying decisions were not overridden by adult social group size or existing egg
locations

There was a strong preference to lay at the edge of the vial, and there was no effect of existing

egg location (X^2 = 1.32, d.f. = 113, p = 0.517) or group size (X^2 = 3.80, d.f. = 112, p = 0.0513) on

where in the environment females laid. There was also no effect of either egg location or

group size on whether females chose to cluster their eggs as no females clustered their eggs

(Table S2).

Table S2. Effect of social density and previous egg laying patterns on female fecundity and egg clustering. The

proportion of vials where eggs laid were observed to be clustered with the existing egg (main bodies in direct

contact) and laid at the edge of the vial (within 0.5cm from the vial circumference).

		Proportion of eggs laid at the edge (%)	Proportion of eggs clustered (%)
Social treatment	Egg location		
Solitary females	None (N = 33 vials)	100	-
	Edge (N = 22 vials)	100	0
	Centre (N = 15 vials)	100	0
Groups of four	None (N = 17 vials)	94.1	-
	Edge (N = 17 vials)	93.8	0
	Centre (N = 13 vials)	100	0

As before, we found that females housed in solitude took longer to initiate laying than those

in groups of four (X^2 = 4.38, d.f. = 126, p = 0.001; Fig. S5). There was no effect of existing egg

location on laying latency (X^2 = 1.30, d.f. = 125, p = 0.139; Fig. S5).

Figure S5. Females housed in solitude were slower to start laying than those housed in groups of 4, but there

was no effect of existing egg location on laying latency. Means and standard errors are shown for the six

treatments. Grey box indicates period of dark (21:00 – 09:00 GMT).