Female fruit flies use social cues to make egg clustering decisions

Emily R. Churchill[1], Emily K. Fowler[2], Lucy A. Friend[2], Marco Archetti[3], Douglas W. Yu[2], Andrew

F. G. Bourke[2], Tracey Chapman[2*] and Amanda Bretman[1*]

* Corresponding authors

1. School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, UK

2. School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

NR4 7TJ, UK

3. Department of Biology, Pennsylvania State University, University Park, PA 16802, USA

ORCID information:

Emily R. Churchill – 0000-0003-0910-8600

Emily K. Fowler – 0000-0001-5290-7630

Lucy A. Friend – 0000-0001-5728-9384

Marco Archetti – 0000-0002-8303-1632

Douglas W. Yu – 0000-0001-8551-5609

Andrew F. G. Bourke – 0000-0001-5891-8816

Tracey Chapman – 0000-0002-2401-8120

Amanda Bretman – 0000-0002-4421-3337

Keywords

Cooperative oviposition; Laying rate; Public goods; Density; Fitness; *Drosophila melanogaster*

Abstract

Background: The ability to respond plastically to environmental variation is a key determinant of fitness. Females may use cues to strategically place their eggs, for example adjusting the number or location of eggs according to whether other females are present, driving the dynamics of local competition or cooperation. The expression of plasticity in egg laying patterns within individual patches, i.e., in contact clusters or not, represents an additional, under-researched and potentially important opportunity for fitness gains. Clustered eggs might benefit from increased protection or defence, and clustering could facilitate cooperative feeding. However, increased clustering is also expected to increase the risk of over-exploitation through direct competition. These potential benefits and costs likely covary with the number of individuals present, hence egg clustering behaviour within resource patches should be socially responsive. We investigate this new topic using the fruit fly *Drosophila melanogaster*.

Results: Our mathematical model, parameterised by data, verified that females cluster their eggs non-randomly, and increase clustering as group size increases. We also showed that, as the density of adult females increased, females laid more eggs, laid them faster, and laid more eggs in clusters. Females also preferred to place eggs within existing clusters. Most egg clusters were of mixed maternity.

Conclusions: Collectively, the results reveal that females actively express plasticity in egg clustering according to social environment cues and prefer to lay in clusters of mixed

maternity, despite the potential for increased competition. These findings are consistent with

egg clustering plasticity being selected due to public goods-related benefits.

Background

In natural contexts, social environments and availability of resources are highly variable,

exposing populations to fluctuating competition. In response, phenotypic plasticity can play

a key role in ensuring individuals maximise their lifetime reproductive success (1, 2). Because

optimal strategies may vary with fluctuating levels of competition, individuals that respond

to the prevailing environment should show increased fitness. Plasticity in investment

decisions should be particularly important for females, who can generally exert greater

control over reproductive investment in progeny. For example, females can sometimes

control paternity (3-5), the number and sex ratio of offspring via ejaculate manipulation (6-

8), and the choice of nesting/oviposition site locations. For females that deposit eggs upon a

substrate, egg site location can be a vital determinant of hatching success. Consistent with

this, female aggregation and oviposition behaviours of many species appear to be plastic in

response to environmental variation (9-12).

Plasticity in aggregation and egg laying - competition, cooperation, and cheating among

females

At increasing population densities, there is expected to be a greater risk of direct competition

but also increased potential for cooperation, and the relative importance of the two is

expected to differ as individuals progress through life stages. For example, when food is

limiting and competition high, eggs and larvae are typically more vulnerable to cannibalism

than are adults, due to their more restricted mobility (13-15). On the other hand, in some

species, there may be benefits of joining with conspecifics when foraging or hunting. For

example, Eurasian Coots (*Fulica atra*) copy each other's behaviours to reduce foraging costs

(16), and larvae in metamorphosing species can benefit from increased efficiencies resulting

from communal feeding (17, 18). Females can use the presence of conspecifics as a cue to

modify reproductive investment in response to changing social conditions. For example,

increasing adult female density reduces offspring production and body size in Western

Mosquitofish (*Gambusia affinis*) (19). In contrast, in the Willow Leaf beetle (*Plagiodera*

versicolora) fitness effects of increasing adult female density can vary with development,

resulting in a decrease in egg hatching, but an increase in egg clutch number and duration of

laying period (20).

Among females of species that deposit their eggs upon substrates, it is also often

found that eggs are laid in conspecific clusters, e.g. in fish (21), birds (22), reptiles, amphibians

(23), and invertebrates (24). Egg clustering within and among kin may increase the local

population density of offspring and thus the potential not only for competition but also

cooperation at early life-history stages. Benefits from cooperation could include reduced

energy expenditure for ovipositing females by minimising search times, reduced egg

predation risk (the dilution effect' hypothesis (25)), increased protection from abiotic

challenges, or increased larval communal feeding.

However, cooperative interactions among non-kin can be exploited by cheater

individuals. Potential opportunities for cheating can occur in egg-laying species in which

mothers provide their eggs with diffusible defensive compounds. One example of defensive

compounds is found in the medfly (*Ceratitis capitata*) (26); females provision the surface of

their eggs with antimicrobial peptides, which spread into the surrounding environment,

creating a zone of microbial growth inhibition. Other examples include in the Green lacewing

moth (*Ceraeochrysa cubana*), in which eggs are endowed with an anti-predator alkaloid (27),

and in fruit flies (*Drosophila melanogaster*), in which eggs are coated with anticannibalism

compounds (28). In each case, these defensive substances represent potential public goods

(29, 30) as all nearby eggs could benefit, including those not provisioned with defensive

compounds. This raises the intriguing possibility that in egg clusters of mixed maternity, there

could be cooperator eggs coated with defensive compounds, and non-defended cheater eggs

that benefit from their placement within the defensive diffusion radius. In this scenario,

cheaters might benefit by avoiding the energetic expense of provisioning eggs with defensive

compounds. Egg clustering might also provide benefits as well as cheating opportunities

through communal feeding. For example, if eggs are laid on hard substrates there could be

benefits for all via the collective processing of food, while simultaneously allowing the

possibility for cheaters to benefit by utilising resources liberated by the processing of food by

their cluster mates. Such co-operators and cheats could stably coexist within a population if

the benefits of public goods have a non-linear relationship with fitness (30, 31).

The potential fitness benefits of joining an existing egg cluster within a food patch

could also depend on key factors such as number, quality, fertility, and age of eggs already

present. Based on existing evidence that females can vary the number of eggs they lay

according to social density (9, 10, 32) we predict, and test here in the *D. melanogaster* fruit

fly model system, the idea that females should use number of eggs and/or number of adult

conspecifics as cues to direct their egg-clustering patterns.

Drosophila melanogaster as a model for understanding plasticity in oviposition

In wild populations, *D. melanogaster* larvae and adults periodically occur at high densities

around fallen, fermenting fruit. For females, fruit is both a source of nutrition and potential

oviposition sites. In *Drosophila* and other species in which fruit resources are patchy,

ephemeral, and vary in nutrient availability (density, ripeness, and state of decomposition)

access to oviposition sites is likely to be shaped by the immediate social environment (number

of conspecific or heterospecific males and females utilising the same resource) (33). To

oviposit in a way that maximises fitness, females should assess and respond appropriately to

the nutritional quality, degree of interspecific competition, risk of pathogens and parasites

present at each oviposition site, as well as the ongoing search costs, and potential benefits

and costs for developing offspring (34).

Consistent with this reasoning, previous research shows that female *Drosophila* can

show attraction to conspecifics that are utilising specific oviposition resources (35, 36), which

they can detect directly or via pheromones, markings, or the presence of eggs or larvae (37-

41). Several studies show that gravid females can aggregate to lay their eggs on substrate

patches used by others according to a variety of physical and social environmental conditions

(9, 42-44). This attraction could arise because it allows females to copy the site-selection

choices of others (37, 38, 40) to enable cooperative feeding among larvae (45), or to gain

other potential public goods benefits. Existing evidence also suggests that female *Drosophila*

can adjust the number of eggs they lay according to their previous or current social

environments, laying fewer eggs after exposure to conspecifics before mating (9, 46) but

laying eggs more quickly if they are in larger social groups after mating (32).

Drosophila eggs can often be found in extremely close proximity, in large clusters (Fig.

S1). Hence, after deciding to lay on substrates with existing conspecific eggs, females must

also decide whether to lay eggs that join existing clusters. Females generally lay one egg at a

time (47), explore oviposition sites between laying, and can retain eggs until an optimal laying

site is found (48). Therefore, clustering of eggs suggests that females make repeated egg-

laying decisions and that females lay their eggs in non-random distributions. Whether the

temporal or spatial distribution of egg laying is significantly non-random, how laying location

choices vary *within* a patch, and whether the factors that drive female egg laying patterns and

choices vary *within* patches are not known. The pattern of egg laying within patches can be

examined by assessing egg clustering patterns, which we define as two or more eggs in direct

contact (Fig. 1). Egg clustering is expected to be important, as it offers females the opportunity

to optimise fitness by placing their eggs in manner that maximises benefits of cooperation

and the potential for public goods benefits (e.g., protection, defence, or cooperative

tunnelling) against potential costs of competition (e.g., over-exploitation of food). Clustering

by our definition, also potentially enables further benefits only transferred via eggs being in

direct contact.

144

Figure 1. **Egg clustering is defined as eggs in direct contact.** Eggs where the main bodies were in direct contact

(left) were classified as clustered, and those not in contact (or where respiratory appendages only overlap) were

classified as singly laid (right).

We tested these ideas with three hypotheses:

(1) that female *D. melanogaster* lay their eggs non-randomly within a food patch

(2) that changes to adult social density, and the presence of existing egg clusters, alter egg-

clustering patterns

(3) that females would lay eggs in clusters of mixed maternity at high frequency

Though our initial observations suggests that egg laying patterns *within* patches are

non-random, this has not yet been formally confirmed. Therefore, we first constructed a

mathematical model, parameterised by experimental data, to test the hypothesis that

females exhibit non-random egg placement (H1). We then conducted additional experiments

to determine key drivers of egg laying patterns observed. We examined whether egg-

clustering patterns responded plastically to adult female group size, or to the density of eggs

already present in the local environment (H2). Potential advantages of site copying and egg

clustering (9, 17, 18, 25-28, 32, 38, 45, 49, 50) predicted that females should increase egg

clustering in response to higher densities of adult females and already-laid eggs in the

environment. Females could potentially maximise public goods-related benefits under

increased clustering by laying eggs alongside those of others. To establish whether conditions

for this scenario exist, we examined whether females so indeed cluster eggs within patches

with those of conspecifics by using dyed eggs to distinguish eggs laid by different females

(H3).

Results

We first constructed a model, parameterised with data, to test the hypothesis that females

exhibit non-random egg placement (H1). Having confirmed that egg laying patterns were

significantly non-random, we then conducted experiments to test the effect of social group

size, and presence of existing egg clusters, on the rate and likelihood of egg laying, egg

clustering patterns and fitness (egg-adult viability as a function of egg clustering (H2)). The

final experiment was then conducted using females that produced dyed eggs, to measure the

frequency of egg clusters with mixed maternity (H3).

1. Egg laying patterns were significantly non-random, and clustering increased with increasing

group size (H1, H2)

Across the four social treatments, clustering preferences (values of Қ that, based on the

Kolmogorov-Smirnov test, yielded no significant difference between the simulated and

observed distributions) were between 0.3 and 0.5 (Table S2).

For all social treatments with two or more females, patterns of egg clustering in the

empirical data were significantly more clustered than those predicted by the null model (H1;

Wilcoxon Signed Rank test comparisons with simulations using Қ = 0; paired: $z_{(6)}$ = -2.29,

p = 0.0220, r = 0.866, N = 7 vials; groups of four: $z_{(25)}$ = -4.47, p = 7.95 x 10^{-6}, r = 0.876, N = 26

vials; groups of eight: $z_{(27)}$ = 4.63, p = 3.63 x 10^{-6}, r = 0.875, N = 28 vials). Therefore,

egg-clustering patterns of all grouped females followed a non-random distribution (Table S2,

Fig. 2) but those of solitary females did not ($z_{(2)}$ = -1.36, p = 0.174, r = 0.786, N = 3 vials),

suggesting that their laying patterns were random (though we note that parameters were

more difficult to estimate for the solitary social treatment as clustering was observed only

infrequently under those conditions). Clustering preferences increased with increasing

female group size: solitary and paired females had a significantly lower clustering preference

in comparison to females housed in groups of 4 or 8 (Table S2, Fig. 2). The findings support

H1, that eggs were laid in non-random patterns that became more non-random as social

density increased. Therefore, egg clustering patterns were plastic and responsive to the

female's social environment.

Figure 2. **Egg clustering is non-random and increases with increasing adult social density.** (a) Comparison of the clustering proportion model outputs and empirical egg

laying patterns observed in *D. melanogaster,* for four social group sizes (1, 2, 4, and 8 adult females per group). The average Kolmogorov-Smirnov P values at К (clustering

preference) values ranging from 0 to 1 in 0.1 increments – where the null distribution is К = 0 (no eggs clustered), and К = 1 (all eggs laid in one large cluster). The best-fit

values of К are those that produced the least significant (highest P value) difference between simulated and observed data. (b-e) The Kolmogorov-Smirnov P values for

individual replicates in the four social group sizes (1, 2, 4, and 8 adult females), with the values for each vial shown in the pale, thin lines and the average value indicated with

a bold line. The vertical grey dotted lines show clustering preference values.

2. Egg-clustering patterns responded both to the density of adult females and presence of

already laid eggs in the environment (H2)

(i) Females in larger social groups clustered their eggs more and, as expected, laid more eggs

and laid eggs at a faster rate

Using the empirical data shown in Fig. 2, we found that females in larger social groups were

significantly more likely to lay eggs (X^2 = 20.2, d.f. = 116, p = 1.51 x 10^{-4}). Eggs were present in

100% of vials housing eight females, a significantly higher proportion than for vials containing

a solitary female (66.7%, X^2 = 15.9, d.f. = 58, p = 6.76 x 10^{-5}) or those with two females (83.3%,

X^2 = 7.39, d.f. = 58, p = 0.00657). There were also significantly more vials containing eggs from

the four female treatment (96.7%) in comparison to the solitary female treatment (X^2 = 10.2,

d.f. = 58, p = 0.00140).

Females in larger social groups also laid more eggs per female than did the females

from the other treatments (solitary females: 7 ± 10 eggs per female (N = 30 vials); pairs: 9 ± 9

eggs per female (N = 30 vials); groups of four: 18 ± 8 eggs per female (N = 30 vials); groups of

eight: 19 ± 5 eggs per female (N = 30 vials); $F_{3, 116}$ = 265, p = 2.93 x 10^{-8}; Fig. 3a). Consistent

with this, post-hoc tests showed that females in solitary and paired social densities laid fewer

eggs than females in groups of four and eight (all p << 0.001). To control for the increased

sampling effort in treatments with larger groups of females, the analysis was repeated with a

randomised subset of data to make number of flies (rather than the number of vials) in each

treatment similar. The results of these analyses were comparable with the main analysis

(larger social groups showed significantly higher likelihood of egg laying: X^2 = 50.0, d.f. = 54,

p = 0.0271 and significantly higher numbers of eggs laid: $F_{3, 54}$ = 137, p = 0.00427).

An analysis of the data at the final timepoint (24h post-mating) supported the

clustering analysis described in the previous section. Females housed in larger groups laid a

higher proportion of their eggs in clusters (solitary females: 22.6% ± 26.9% (N = 20 vials);

pairs: 38.1% ± 28.9% (N = 25 vials); groups of four: 62.7% ± 19.2% (N = 29 vials); groups of

eight: 73.7% ± 13.3% (N = 30 vials); total number of eggs laid included in model: $F_{3, 99} = 28.4$,

$p = 2.53 \times 10^{-13}$; all pairwise comparisons $p < 0.05$; Fig. 3b). We included the total number of

eggs laid in the model as we found that there was a positive relationship between the number

of eggs and the proportion of eggs clustered ($F_{1, 99} = 6.66$, $p = 0.0113$), although this was not

the key driver of social density effects on clustering proportions. In all grouped female

treatments, the proportion of clustering also increased over time (individual vial included in

model as a random effect: pairs: $F_{1, 58} = 19.1$, $p = 5.26 \times 10^{-5}$; groups of four: $F_{1, 95} = 75.8$, $p =$

9.99×10^{-14}; groups of eight: $F_{1, 111} = 74.7$, $p = 4.45 \times 10^{-14}$). There was no significant effect of

time on the proportion of clusters in eggs laid by solitary females ($F_{1, 40} = 1.15$, $p = 0.289$).

Consistent with this, the number of clusters and size of the largest cluster increased

with both increasing social density (number: $F_{3, 376} = 463$, $p = 2.20 \times 10^{16}$; size: $F_{3, 299} = 58.3$,

$p = 2.20 \times 10^{-16}$; Figs. S3-4), and time (number: $F_{1, 376} = 631$, $p = 2.20 \times 10^{-16}$; size: $F_{1, 299} = 67.1$,

$p = 7.62 \times 10^{-15}$). These results were robust to differences in sampling effort as shown by an

additional analysis using the same number of individuals (rather than vials) at the final

timepoint. The effects remained: females in larger groups laid a higher proportion of their

eggs in clusters ($F_{3, 41} = 6.40$, $p = 0.00117$) and the number and size of clusters still increased

with increasing social density (number: $F_{3, 134} = 199$, $p = 2.20 \times 10^{16}$; size: $F_{3, 102} = 25.1$, $p = 2.96$

$\times 10^{-12}$).

Interestingly, there was no evidence for a fitness benefit for clustered eggs in terms

of increased egg-adult survival (Fig 2c). There was no effect of social density (1, 2, 4, or 8

females) on the proportion of eggs that survived to adulthood ($F_{3, 99}$ = 0.292, p = 0.831; Fig.

S5) and there was no correlation between the proportion of eggs laid in clusters and egg-adult

offspring viability (t = -0.485, d.f. = 101, p = 0.629; Fig. 3c).

Figure 3. **Females in larger groups laid more eggs, more quickly, with a higher proportion of clusters**. (a) Females housed in groups of four and eight were quicker to lay,

and laid more eggs per female, than those kept in solitude or in pairs (all N = 30 vials). Eggs laid per individual were calculated by dividing total number of observed eggs by

the number of females in the vial. Social treatment means and standard errors are shown for the four housing densities at each observation time point. The grey box indicates

period of dark (21:00 – 09:00 GMT). (b) Females housed in groups of four or eight laid a higher proportion of their eggs in clusters compared to those kept in solitude or in

pairs (all N = 30 vials). Proportion of eggs clustered was calculated by summing all cluster sizes and dividing by the total number of eggs counted. (c) There was no effect of

the proportion of egg clustering on egg-adult viability (linear regressions with 95% confidence intervals; solitary: N = 20 vials; paired: N = 24 vials; groups of 4: N = 29 vials;

groups of 8: N = 30 vials).

Overall, the results supported the hypothesis that the egg-clustering behaviour of

females within a food patch responds plastically to the social environment, with females

laying eggs more quickly, laying more eggs and laying more eggs in clusters, as the social group

size increases. However, there was no indication that eggs laid in clusters had higher egg-adult

viability.

(ii) Females preferred to lay eggs within existing clusters rather than with singly laid eggs

Female egg-clustering behaviour also showed plasticity in response to eggs already present

in the environment. Females were more likely to add at least one of their eggs to existing

clusters of 4, 7, or 10 eggs, than they were to lay with a singly placed existing egg ($X^2 = 12.5$,

d.f. = 156, p = 4.18 x 10^{-4}; Fig. 4a). The total proportion of eggs laid that joined existing eggs

was also higher when existing eggs were clustered ($F_{3, 154} = 4.75$, p = 0.00340; Fig. 4b).

Proportions were calculated from the number of eggs females were able to lay within the 30

min observation window (between 1 – 19 eggs per vial). Post-hoc tests showed that egg

cluster sizes of 4 and 10 were joined significantly more often than were single eggs (four: $F_{1, 71} = 13.0$, p = 5.85 x 10^{-4}; ten: $F_{1, 78} = 8.76$, p = 0.00407), though this trend was marginally non-

significant for a cluster size of 7 ($F_{1, 75} = 3.82$, p = 0.0542).

Figure 4. **Females lay more eggs within existing clusters and lay eggs closer to existing egg clusters** (a) A higher proportion of females clustered at least one of their eggs

with existing eggs, when those existing eggs were in clusters compared to singly laid eggs (singly laid eggs: N = 36; clustered eggs: N = 122. (b). Females laid a higher proportion

of their eggs in clusters with existing eggs, when those existing eggs were already in a cluster (of any size) compared to a singly laid egg (1 egg: N = 36; 4 eggs: N = 37; 7 eggs:

N = 41; 10 eggs: N = 44). Means and standard errors are shown for the four treatment egg cluster sizes. Significant differences between treatment egg cluster sizes are

represented by an overarching bar; * indicates a significant difference between paired treatments (* P < 0.05, ** P < 0.01, *** P < 0.001). (c). Mean distance of laid eggs from

existing treatment eggs was greater when the number of existing eggs was one, compared to egg clusters of any size (1 egg: N = 32; 4 eggs: N = 37; 7 eggs: N = 40; 10 eggs:

N = 40).

In addition to scoring whether a laid egg was physically in contact with an existing egg

(i.e. clustered), we also measured the distance between each newly-laid egg and existing

treatment eggs already present on the substrate. We did this to test for potential benefits of

close-proximity in addition to direct contact. Females laid their eggs closer to existing clusters

(H3; $F_{3, 145}$ = 9.86, p = 5.86 x 10^{-6}; Fig. 4c) with all three cluster sizes having shorter inter-egg

distances than eggs from females exposed to single eggs (four: $F_{1, 67}$ = 14.8, p = 2.68 x 10^{-4};

seven: $F_{1, 70}$ = 11.4, p = 0.00122; ten: $F_{1, 70}$ = 23.8, p = 6.57 x 10^{-6}). Hence, laid eggs that did not

strictly join a cluster were still laid closer to existing clusters than to single eggs. The results

show that females also respond plastically to the clustering patterns of eggs already present

in the environment and, consistent with the results above, lay their eggs preferentially within

existing egg clusters (H3).

3. Egg clusters were typically of mixed-maternity (H3)

In the final experiment we used a lipophilic dye to enable us to distinguish between eggs laid

by different females (51). We set up groups of 4 females comprising 1 standard-fed focal wild

type and 3 Sudan Black B dye-fed non-focals. We found that of the focal eggs that had been

laid in clusters, a mean of 79.5% of them were in mixed-maternity clusters (Fig. 5). This

showed that females do not preferentially choose to isolate their eggs by laying them solely

in single-maternity clusters, but instead lay eggs in clusters of mixed-maternity at high

frequency.

Figure 5. **High frequency of mixed maternity egg clusters.** Females showed no preference for laying their eggs

in single-maternity clusters, as 79.5% of egg clusters observed were of mixed maternity (N = 28 vials). Shown is

a frequency distribution of the proportion of egg clusters counted that had eggs laid by two or more gravid

females. The inset picture shows a cluster of four eggs: two undyed and two dyed with Sudan Black B dye.

Discussion

The main new findings of this study were that female egg-clustering behaviour within patches

responded plastically to the social environment and to the presence of existing eggs within

the oviposition environment. All three hypotheses were supported. 1) Egg-clustering patterns

were significantly non-random, for females in groups, but not for eggs laid by socially isolated

females. 2) As the social group size of females increased, females laid eggs more quickly, laid

more eggs, and laid more of them in clusters. However, there was no evidence of a fitness

benefit to clustered eggs in increased egg-adult viability. 3) Clusters of eggs were usually of

mixed maternity and females preferred to add their eggs to existing clusters. Overall, females

showed striking plasticity in their egg-clustering decisions, with a strong bias towards laying

eggs in mixed-maternity clusters. However, the fitness benefits of such behaviour remain

elusive. These results are explored in more detail, below.

Egg-clustering decisions are plastic and eggs are laid in non-random patterns

For all females except those held in social isolation egg clustering occurred more than the

chance outcomes predicted by the model. This provides evidence that females make active

egg location decisions, and that these responses are plastic and change in response to varying

social environments. Egg clustering shows that females make repeated egg-laying decisions,

as they generally explore potential laying sites between laying each individual egg, and can

retain their eggs until they find an optimal site (47, 48). Increased clustering has the potential

to increase local population density within patches, which has potential cumulative effects,

as even relatively small-scale variation in density leading to increased competition can have

consequences for fitness (52).

Females cluster more of their eggs in larger groups, and more commonly form mixed-maternity

clusters

The proportion of eggs that were laid in clusters increased as social density increased.

Although there was a higher chance of clustering when there are more eggs present in the

environment, this increase in clustering with increasing social density did not depend upon

egg number (in agreement with the null model comparison). It was not possible to determine

the maternity of the clusters in grouped treatments in the initial experiment, but our later

experiment showed that the clusters are usually of mixed maternity. Females exposed to

other (non-kin) females preferred to add their eggs to mixed-maternity clusters, and overall

socially isolated females cluster their eggs much less, which suggests that there are potential

benefits of laying in mixed-maternity clusters over and above laying in clusters *per se*. The

drivers of this are not yet known, however females could be balancing a bet-hedging strategy

of laying in multiple sites to reduce risks of sibling competition (if food is limited) or to

minimise predation/parasitism risk by reducing transmission and egg visibility (53), whilst

adding to existing clusters to reap potential public goods benefits of acquiring more diverse

microbiomes (54, 55) or diffusible defensive compounds such as anticannibalism pheromones

(28) or antimicrobials (as seen in medfly (26)), but see (51). Mixed-maternity egg clustering

could select for cheating among females under a public goods system (29, 30), a possibility

that would be interesting to test further.

Females preferred to lay their eggs in existing egg clusters

Females preferred to add their eggs, or lay them close to, existing clusters of any size.

Interestingly, the strength of this preference did not increase with increasing cluster size. This

could be because (i) there is no benefit to increasing cluster size (i.e., benefits to joining

clusters are not additive), (ii) the clusters here did not vary sufficiently in size, or (iii) there is

some trade-off to the benefits of clusters of these sizes. For example, eggs laid later may be

at a higher risk of cannibalisation (28, 56) and the risks and costs might increase with cluster

size. Costs of laying in mixed-maternity clusters are also possible if being surrounded by kin

provides benefits via communal feeding of larvae (13). Exactly how individuals might identify

maternity of egg clusters within patches, whether by detecting conspecific pheromones (41,

57, 58)) or absolute cluster size remain to be investigated. The strong aversion to laying with

isolated eggs raises the question of how clusters are initiated in the first place. Female choice

to copy and lay with other eggs could require a threshold of a repeated decision – i.e., at least

two eggs at the same site, because this is a more robust indicator of site quality than only

one. Females may avoid laying with existing single eggs if those have a higher probability of

infertility and thus offer potentially fewer benefits. However, we found no relationship

between cluster size and egg-to-adult viability, which argues against this idea. In addition,

whether females can detect infertile eggs is not known, although those laid by virgins have a

distinct pheromonal profile (41)). One scenario is that clusters form if the first two eggs are

laid by the same mother. Arguing against this idea is that clusters of two eggs laid by different

mothers were observed in our mixed-maternity investigation. This suggests that further

studies on clustering initiation are required.

Potential fitness benefits of egg clustering remain elusive

Interestingly, we detected no increase in egg-adult viability with increasing egg clustering. It

is not clear why we did not observe fitness benefits, given potential costs of both maintaining

plasticity (59, 60) and of increasing local competition (13, 28, 34, 52-56). It is possible that as

mothers were able to choose a clustering strategy, all individuals chose the optimal strategy.

To test this, future work could force mothers into making an incorrect choice via a mismatch

of environments. It is also possible that the *ad libitum* food provided obscured any costs of

choosing the 'wrong' egg laying strategy. It was necessary to remove females from the egg

laying environments before they had oviposited all fertile eggs, as eggs needed to be counted

prior to larvae hatching which takes place after 22 – 24 hours at 25°C (61, 62). Thus, it is also

possible that differences in egg-adult viability could have been detected if females were able

to continue laying if the benefit only occurs in clusters of mixed-age eggs.

Egg number and laying rate also increased as adult female density increased

As expected, based on existing research (32), females in larger social groups laid more eggs

and laid them more quickly. This is consistent with raffle theory, in which the production of

more eggs when competition is high ensures that mothers have a greater chance of at least

some of their offspring surviving to adulthood (63). With a greater number of conspecifics

present in the environment, there should also be a higher chance of copying behaviours

arising, because a greater number of potential 'initiators' are present (37, 38, 40). Increased

copying could increase laying rates, benefitting mothers as it may reduce sampling times

needed to locate optimal oviposition sites. Consistent with this, we observed that latency to

lay the first egg was also shorter at higher adult densities. In situations in which larval density

is high and resources are limiting, later-laid eggs could have reduced survival due to resource

degradation and cannibalisation (13, 32, 56). Thus, in high density environments, mothers

should ensure they begin laying quickly to avoid increased offspring lethality.

Previous research (32) showed that the daylight period can inhibit egg laying in

Drosophila, an effect that can be over-ridden by being in a group (five females). Intriguingly,

our results showed remarkably similar laying rates for isolated and paired females (whereas

groups of four or eight females matched laying times reported in (32)), suggesting that

exposure to only one other gravid female does not counteract the previously described light-

induced inhibition of egg production. This is surprising given that in male *Drosophila*, exposure

to one other rival is enough to facilitate reproductive behavioural responses (64-66). This may

suggest that there are sex differences in responses to local population density.

Conclusions

Overall, we demonstrated that *D. melanogaster* mothers express plasticity in egg-clustering

decisions, laying eggs in a significantly non-random manner and responding to differences in

social density from multiple life stages. Exposure to increased adult social density resulted in

females laying more eggs, at a faster rate, and, a new result from this study, laying a higher

proportion of those eggs in clusters. Exposure to existing egg clusters also led to a higher

frequency of clustering decisions, regardless of existing cluster size. We hypothesise that eggs

may be clustered to gain public goods benefits. However, the overall ecological importance,

and the costs and benefits of this plasticity remain unknown. These results demonstrate that

egg laying and clustering decisions are highly sophisticated and add to the growing evidence

that even non-social organisms such as *D. melanogaster* have unexpectedly rich social lives

(67, 68).

Methods

Fly husbandry and rearing of experimental flies

Fly rearing and all experiments were conducted at 25°C on a 12h light:dark cycle (09:00-21:00

GMT). Wild type flies originating from a large Dahomey laboratory population maintained in

cages with overlapping generations were used throughout. We collected eggs from cages

using Petri dishes (*Sarstedt #82.1473.001*) filled with grape juice agar-based medium (50g

agar (*Fisher Scientific #10048991*), 600ml red grape juice (*Young's Brew red wine enhancer*),

42ml Nipagin solution (methylparaben, 10% w/v solution, dissolved in 95% Ethanol) per 1.1l

H_2O). Once eggs had hatched, we transferred first instar larvae to a 40ml plastic vial (*Sarstedt*

#58.490) containing 7ml of a standard sugar-yeast-agar (SYA) medium (100g brewer's yeast

(*Buy Wholefoods Online*), 50g sugar (*Tate & Lyle*), 15g agar, 30ml Nipagin solution, and 3ml

propionic acid (*Fisher Scientific #10193190*) per litre of medium), at a standard density of 100

larvae per vial (i.e., 'standard density vials'). Experimental flies were then collected under light

ice anaesthesia, within six hours of eclosion to ensure virginity.

1. Development of mathematical model of egg-clustering to test for non-random egg

deposition (H1)

To ascertain whether the proportion of observed egg clustering differed from a null

expectation, we produced a model to calculate random distributions of eggs in a standard

vial. We defined a 'cluster' as any group of two or more eggs in which the main bodies of the

eggs were in direct contact (Fig. 1). In the model, given that the observed egg to dish area is

1/5770, we assume that: (i) there are 5770 positions available for egg laying, and (ii) there is

a total number (parameter $\mathcal{E}$) of eggs per vial. Assuming that all eggs are laid sequentially and

at random, we calculated the expected size of clusters (defined as the number of eggs per

cluster) for any number of total eggs laid ($\mathcal{E}$).

The preference for clustering eggs next to an existing egg (hereafter 'clustering

preference') is defined by the probability of a clustering decision. К is the probability that a

fly will lay an egg next to an already existing egg. After the first egg is placed at random, where

$\mathcal{E}$ > 1, the next egg is either laid in a random position (probability of probability of 1 - К) or

next to an existing egg (probability of К); in the latter case, the position of the egg is chosen

at random amongst only positions with existing eggs.

The expected number of eggs per cluster can be predicted based on К and $\mathcal{E}$. When

К = 0 (no preference for egg clustering), very few clusters with 2 eggs occur and virtually no

clusters with more than 2 eggs (2.5% with $\mathcal{E}$ = 233 (the maximum number of eggs observed in

our experiment); 1% with Ɛ = 100; no clusters for Ɛ below 70). Therefore, if Ɛ << 5770 (as is

the case in our experiment), randomly laid eggs form almost exclusively singly laid eggs. When

К = 1 (absolute preference for egg clustering), all eggs are clustered in one large cluster in a

single position.

At intermediate values of К, some variation in cluster size emerges. We simulated the

expected number and size of egg clusters for different values of К from 0 to 1 in intervals of

0.1 (using the average of 10 different simulations per К value). We then compared these

expected distributions to the empirically observed distributions of eggs laid by females from

the differing social density treatments (groups of 1, 2, 4, or 8) described below, to estimate

the most likely preference К leading to that distribution. For each vial, we performed a

Kolmogorov-Smirnov test (implemented in Mathematica 13. 1 (69)) between the expected

and observed distribution, for all К values. We considered the К value yielding the highest

(least significant) P values to be the most likely К value for that vial.

2. Testing for plasticity in egg-clustering behaviour in response to varying egg and adult density

(H2)

(i) Effects of adult social density on egg-clustering decisions and fitness

We investigated how variation in adult social group size affected egg laying and egg-clustering

decisions: specifically, the speed of egg laying, the number of eggs laid, egg clustering

patterns, and fitness (egg-adult viability of clustered versus non-clustered eggs). Some data

from this experiment were also used to parameterise the model above. Experimental flies

were collected and housed in same-sex groups of 10 in standard vials. At four days post-

eclosion, females were randomly assigned to one of the following social group treatments: 1,

2, 4, or 8 females and kept in these conditions for 48h (N = 30 vials each treatment; Fig. S1).

Males were held for six days until they were used for matings.

At 6 days post-eclosion, males were introduced to all the female social group

treatment vials for 2h at a density of 3:2 males:females (or 2:1 for isolated females) to enable

choice (Fig. S1). To ensure that this was a set up that would result in all females successfully

mating within the 2h period allowed, we separately recorded female mating frequency in an

identical experimental set up. In this we observed that 97.3% of females mated, with no

significant difference in mating frequency between treatments (X^2 = 1.79, d.f. = 3, p = 0.617).

Thus, we can assume that there is only a very low probability of females remaining unmated

by using this procedure.

After the 2h mating period in the main experiment, we transferred all mated females

to fresh standard vials to observe egg laying decisions. Every 2 – 3h, we counted the number

of eggs laid, number of egg clusters, and size of egg clusters. Observations were made at

14:00, 16:00, 19:00, and 22:00 on day 1 of laying, and 10:00 and 12:00 on day 2 (i.e., 2, 4, 7,

10, 22, and 24h after the end of the mating period, respectively). Differences in counts

(number of eggs, number of clusters, and size of clusters) between timepoints were calculated

to give overall egg laying latencies, numbers and clustering rates. We calculated the

proportion of eggs in clusters at each timepoint by summing the total number of eggs in each

cluster and dividing by the overall total number of eggs laid in the vial. All females were

removed after 12:00 on day 2 and vials were then kept for 14 days, to count the total number

of eclosed adults from each of the social exposure treatments.

(ii) Effects of existing egg clusters on female egg-clustering decisions

To assess whether the size of egg clusters already present in an environment alters

subsequent female egg-clustering decisions, we presented focal females with a laying

environment that already contained varying numbers of eggs and egg clusters. To create the

egg clusters, singly laid eggs were collected within 2 hours of laying from SYA medium-filled

Petri dishes and transferred into vials to create four different egg cluster sizes: 1, 4, 7, and 10

eggs per cluster. Females have a strong tendency to lay eggs at vial edges (Supplementary

Information Text; Table S1; (9, 49, 50)), hence we simulated this preference by placing all egg

clusters at the edge of the egg treatment vials.

To obtain the Dahomey females and males for the experiment, we collected virgin flies

from standard density vials. Prior to the tests, females were maintained alone, males in

groups of four per vial. At seven days post-eclosion, males and females were paired and

observed to ensure they had all successfully mated. After mating, groups of 4 females were

then transferred to each of the 4 types of egg cluster treatment vials (1 egg: N = 20 vials; 4

eggs: N = 17 vials; 7 eggs: N = 20 vials; 10 eggs: N = 21 vials). After the introduction of the focal

females, vials were checked at 30 min intervals from 13:00 until 21:30, at which point if

females had not laid eggs they were removed. If eggs were laid by the introduced focal

females, we counted the number of eggs laid and categorised them in relation to the existing

egg cluster present (clustered or not clustered). All vials were then frozen at -20°C for imaging.

Vials were imaged to record inter egg distances, using a video camera (Sony Handycam

HDR-CX405). Using ImageJ's multiple point selector tool (70) we captured x-y coordinates of

each egg and converted these into Euclidean pairwise distances. Selected coordinates were

taken from the edge points of eggs that were in closest proximity, resulting in clustered eggs

having a distance value of zero.

3. Testing for mixed maternity of egg clusters (H3)

In the final experiment, we determined the extent of egg clusters of single or mixed maternity.

In each treatment group there was one wild type (Dahomey) focal and 3 non-focal females

from the *scarlet* strain (recessive *scarlet* eye colour mutation backcrossed into the Dahomey

wild type > 4 times). These non-focal *scarlet* females were fed Sudan Black B dye that stained

laid eggs, allowing the eggs of focal versus non-focal females to be identified. First instar

larvae of Dahomey wild type and *scarlet* strains were raised in cultures of 100 larvae per vial

as described above. Focal Dahomey females were collected from cultures reared and

maintained on standard medium, whereas *scarlet* females were derived from cultures

containing Sudan Black B dye (100g brewer's yeast, 50g sugar, 15g agar, 30ml Nipagin

solution, 3ml propionic acid, and 1.4g Sudan Black B powder (*Sigma-Aldrich,* cat#199664)

dissolved in 14ml corn oil (*Mazola* 100% pure corn oil) per litre of medium). Focal Dahomey

virgin females were collected from the standard medium vials, and non-focal *scarlet* virgin

females were collected from Sudan Black B vials and maintained on that same medium until

use. Upon collection, all females were housed in groups of 10 and then Dahomey focal

females were housed in groups of 4 from 5 days post-eclosion. At 4 days post-eclosion, *scarlet*

females were wing clipped for identification, and also housed 4 per vial. Dahomey and *scarlet*

males for matings were collected from standard medium vials and housed in groups of 10

until use.

Seven days post-eclosion, males were transferred into the female vials for mating: 8

Dahomey males were added to each group of 4 Dahomey females, and 6 *scarlet* males for

each group of 4 *scarlet* females. The flies were given 4 hours to mate. One female from each

of the Dahomey vials was then transferred to fresh standard media alongside three *scarlet*

dye-fed females at 12:00 GMT (N = 29). We then counted the number of eggs laid at 10.5h

post-mating. Eggs were categorised as clustered or not clustered (Fig. 1) and according to

whether they were laid by a focal or non-focal female. This allowed us to calculate the

proportion of eggs that were clustered, and whether the cluster contained eggs from both

focal and non-focal females.

Statistical analysis

All statistical analyses were conducted using R v 4. 2. 2 (71), and mean values and standard

deviations are reported within the text. Graphs were produced using packages 'ggplot2',

'cowplot' and 'grid' (72, 73), with plotted standard errors calculated using 'Rmisc' (74).

We analysed the effect of egg and adult social density on the likelihood that females

laid eggs and produced adult offspring, by using generalised linear models (GLMs) with

binomial errors. We used the package 'AER' (75) to test for overdispersion, and where that

was found, we used GLMs with a quasi-Poisson distribution to account for this over-dispersion

and analyse the effect of adult density on the number of offspring produced. Linear models

were used to investigate the effect of adult density on egg-clustering patterns (egg laying and

clustering rates), and total eggs laid was included as a fixed effect to account for increased

likelihood of clustering when more eggs are present in the environment.

When analysing variables throughout multiple timepoints, we used linear mixed

effects models; the timepoint was included as a fixed effect in the model to account for

differences in rates and the test vial was included as a random effect – using functions in

packages 'lme4' (76) and 'lmerTest' (77). We tested for Pearson's product-moment

correlations between the proportion of eggs laid in clusters and subsequent egg-adult

viability. In all cases, post-hoc tests (models including only pairwise comparisons) were used

to identify significant differences in the six pairwise comparisons available.

In the first experiment investigating the effects of social density, there was variation

in sampling effort between treatments, due to the differences in female group sizes (and lack

of identification of focal females). To account for this, all significant results from the initial

analyses were reanalysed with a randomised subset of data so that the number of individual

flies (rather than number of vials) per treatment were similar. In this reduced subset, all vials

from solitary females were included (N = 30 females), half of those in pairs (N = 30 females),

eight of those from groups of 4 (N = 32 females), and four of those from groups of 8 (N = 32

females). These analyses were congruent with those of the initial analyses, suggesting the

initial analyses were robust.

References

1.   West-Eberhard MJ. Phenotypic plasticity and the origins of diversity. Annual Review
of Ecology and Systematics. 1989;20(1):249-78.

2.   Komers PE. Behavioural plasticity in variable environments. Canadian Journal of
Zoology. 1997;75(2):161-9.

3.   Birkhead T, Møller A. Female control of paternity. Trends in Ecology & Evolution.
1993;8(3):100-4.

4.   Rodriguez-Enriquez CL, Tadeo E, Rull J. Elucidating the function of ejaculate expulsion
and consumption after copulation by female *Euxesta bilimeki*. Behavioral Ecology and
Sociobiology. 2013;67(6):937-46.

5.   Ward PI. Females influence sperm storage and use in the yellow dung fly *Scathophaga
stercoraria*. Behavioral Ecology and Sociobiology. 1993;32(5):313-9.

6.   Elgar MA, Schneider JM, Herberstein ME. Female control of paternity in the sexually
cannibalistic spider *Argiope keyserlingi*. Proceedings of the Royal Society of London Series B:
Biological Sciences. 2000;267(1460):2439-43.

7.   Passera L, Aron S, Vargo EL, Keller L. Queen control of sex ratio in fire ants. Science.
2001;293(5533):1308.

8.   Sato A, Karino K. Female control of offspring sex ratios based on male attractiveness
in the guppy. Ethology. 2010;116(6):524-34.

9.   Churchill ER, Dytham C, Bridle JR, Thom MDF. Social and physical environment
independently affect oviposition decisions in *Drosophila*. Behavioral Ecology. 2021.

10.   Fowler EK, Leigh S, Rostant WG, Thomas A, Bretman A, Chapman T. Memory of social
experience affects female fecundity via perception of fly deposits. BMC Biology.
2022;20(1):244.

11.   Takasuka A, Yoneda M, Oozeki Y. Disentangling density-dependent effects on egg
production and survival from egg to recruitment in fish. Fish and Fisheries. 2019;20(5):870-
87.

12.   Matsushima N, Kawata M. The choice of oviposition site and the effects of density and
oviposition timing on survivorship in *Rana japonica*. Ecological Research. 2005;20(1):81-6.

13.   Khodaei L, Long TAF. Kin recognition and egg cannibalism by *Drosophila melanogaster*
larvae. Journal of Insect Behavior. 2020;33(1):20-9.

14.   Perry JC, Roitberg BD. Games among cannibals: Competition to cannibalize and
parent-offspring conflict lead to increased sibling cannibalism. Journal of Evolutionary
Biology. 2005;18(6):1523-33.

15.   Kestemont P, Jourdan S, Houbart M, Mélard C, Paspatis M, Fontaine P, et al. Size
heterogeneity, cannibalism and competition in cultured predatory fish larvae: Biotic and
abiotic influences. Aquaculture. 2003;227(1):333-56.

16.   Novčić I, Aleksić D, Vidović Z. Copying of diving behavior in the Eurasian Coot *Fulica
atra*. Journal of Ornithology. 2022;163(2):523-9.

17.   Cocroft RB. Vibrational communication facilitates cooperative foraging in a phloem-
feeding insect. Proceedings of the Royal Society B: Biological Sciences. 2005;272(1567):1023-
9.

18.   Fitzgerald TD, Peterson SCCFpdJ. Cooperative foraging and communication in
caterpillars. BioScience. 1988;38(1):20-5.

19.   Smith CC, Sargent RC. Female fitness declines with increasing female density but not
male harassment in the western mosquitofish, *Gambusia affinis*. Animal Behaviour.
2006;71(2):401-7.

20.   Zhao L, Wang X, Zhan J. Effects of female and male density on their mating
performance and female post-mating reproductive fitness in *Plagiodera versicolora*
(Laicharting) (Coleoptera: Chrysomelidae). The Coleopterists Bulletin. 2022;76(1):73-81.

21.   Welsh DP, Fuller RC. Where to place your eggs: The effects of conspecific eggs and
water depth on oviposition decisions in bluefin killifish. Journal of Zoology. 2011;284(3):192-
7.

22.   Riehl C. Evolutionary routes to non-kin cooperative breeding in birds. Proceedings of
the Royal Society B: Biological Sciences. 2013;280(1772):20132245.

23. Doody JS, Freedberg S, Keogh JS. Communal egg-laying in reptiles and amphibians:
Evolutionary patterns and hypotheses. The Quarterly Review of Biology. 2009;84(3):229-52.

24. Courtney SP. The evolution of egg clustering by butterflies and other insects. The
American Naturalist. 1984;123(2):276-81.

25. Ioannou C. Grouping and predation. In: Shackelford TK, Weekes-Shackelford VA,
editors. Encyclopedia of Evolutionary Psychological Science. Cham: Springer International
Publishing; 2017. p. 1-6.

26. Marchini D, Marri L, Rosetto M, Manetti AG, Dallai R. Presence of antibacterial
peptides on the laid egg chorion of the medfly *Ceratitis capitata*. Biochemical and biophysical
research communications. 1997;240(3):657-63.

27. Eisner T, Eisner M, Rossini C, Iyengar VK, Roach BL, Benedikt E, Meinwald J. Chemical
defense against predation in an insect egg. Proceedings of the National Academy of Sciences.
2000;97(4):1634-9.

28. Narasimha S, Nagornov KO, Menin L, Mucciolo A, Rohwedder A, Humbel BM, et al.
Drosophila melanogaster cloak their eggs with pheromones, which prevents cannibalism.
PLOS Biology. 2019;17(1):e2006012.

29. Olson M. The theory of collective action: Public goods and the theory of groups.
Harvard UniversityPress, Cambridge. 1965.

30. Archetti M. Cooperation as a volunteer's dilemma and the strategy of conflict in public
goods games. Journal of Evolutionary Biology. 2009;22(11):2192-200.

31. Archetti M, Scheuring I, Yu D. The non-tragedy of the non-linear commons. Preprints:
Preprints; 2020.

32. Bailly TPM, Kohlmeier P, Etienne RS, Wertheim B, Billeter J-C. Social modulation of
oogenesis and egg-laying in *Drosophila melanogaster*. Current Biology. 2023;33(14):2865-
77.e4.

33. Soto-Yéber L, Soto-Ortiz J, Godoy P, Godoy-Herrera R. The behavior of adult
Drosophila in the wild. PLOS ONE. 2018;13(12):e0209917.

34. Schwartz NU, Zhong L, Bellemer A, Tracey WD. Egg laying decisions in *Drosophila* are
consistent with foraging costs of larval progeny. PLOS ONE. 2012;7(5):e37910.

35. Lihoreau M, Clarke IM, Buhl J, Sumpter DJT, Simpson SJ. Collective selection of food
patches in *Drosophila*. The Journal of Experimental Biology. 2016;219(5):668.

36. Zhang L, Sun H, Grosse-Wilde E, Zhang L, Hansson BS, Dweck HKM. Cross-generation
pheromonal communication drives *Drosophila* oviposition site choice. Current Biology.
2023;33(10):2095-103.e3.

37. Golden S, Dukas R. The value of patch-choice copying in fruit flies. PLOS ONE.
2014;9(11):e112381.

38. Malek HL, Long TAF. On the use of private versus social information in oviposition site
choice decisions by *Drosophila melanogaster* females. Behavioral Ecology. 2020.

39. Battesti M, Moreno C, Joly D, Mery F. Spread of social information and dynamics of
social transmission within *Drosophila* groups. Current Biology. 2012;22(4):309-13.

40.   Sarin S, Dukas R. Social learning about egg-laying substrates in fruitflies. Proceedings
of the Royal Society B: Biological Sciences. 2009;276(1677):4323-8.

41.   Duménil C, Woud D, Pinto F, Alkema JT, Jansen I, Van Der Geest AM, et al. Pheromonal
cues deposited by mated females convey social information about egg-laying sites in
Drosophila melanogaster. Journal of Chemical Ecology. 2016;42(3):259-69.

42.   del Solar E. Selection for and against gregariousness in the choice of oviposition sites
by *Drosophila pseudoobscura*. Genetics. 1968;58(2):275-82.

43.   del Solar E, Palomino H. Choice of oviposition in *Drosophila melanogaster*. The
American Naturalist. 1966;100(911):127-33.

44.   del Solar E, Ruiz G. Behavioral analysis of the choice of oviposition site by single
females of *Drosophila melanogaster* (Diptera: Drosophilidae). Journal of Insect Behavior.
1992;5(5):571-81.

45.   Dombrovski M, Poussard L, Moalem K, Kmecova L, Hogan N, Schott E, et al.
Cooperative behavior emerges among *Drosophila* larvae. Current Biology. 2017;27(18):2821-
6.e2.

46.   Fowler EK, Leigh S, Bretman A, Chapman T. Plastic responses of males and females
interact to determine mating behavior. Evolution. 2022;76(9):2116-29.

47.   Yang C-h, Belawat P, Hafen E, Jan LY, Jan Y-N. *Drosophila*; Egg-laying site selection as
a system to study simple decision-making processes. Science. 2008;319(5870):1679.

48.   Vijayan V, Wang Z, Chandra V, Chakravorty A, Li R, Sarbanes SL, et al. An internal
expectation guides *Drosophila* egg-laying decisions. Science Advances.8(43):eabn3852.

49.   Chess KF, Ringo JM. Oviposition site selection by *Drosophila melanogaster* and
Drosophila simulans. Evolution. 1985;39(4):869-77.

50.   Moore JA. Competition between *Drosophila melanogaster* and *Drosophila simulans.* I.
Population cage experiments. Evolution. 1952;6(4):407-20.

51.   Fowler EK, Friend LA, Churchill ER, Yu DW, Archetti M, Bourke AFG, et al. Female
oviposition decisions are influenced by the microbial environment. bioRxiv.
2024:2024.07.03.601843.

52.   Churchill ER, Bridle JR, Thom MDF. Spatially clustered resources increase male
aggregation and mating duration in Drosophila melanogaster. Animal Behaviour.
2020;169:45-50.

53.   Turner GF, Pitcher TJ. Attack abatement: A model for group protection by combined
avoidance and dilution. The American Naturalist. 1986;128(2):228-40.

54.   Guilhot R, Xuéreb A, Lagmairi A, Olazcuaga L, Fellous S. Microbiota acquisition and
transmission in *Drosophila* flies. iScience. 2023;26(9):107656.

55.   Bakula M. The persistence of a microbial flora during postembryogenesis of
Drosophila melanogaster. Journal of Invertebrate Pathology. 1969;14(3):365-74.

56.   Vijendravarma RK, Narasimha S, Kawecki TJ. Predatory cannibalism in *Drosophila
melanogaster* larvae. Nature Communications. 2013;4(1):1789.

57. Bartelt RJ, Schaner AM, Jackson LL. cis-Vaccenyl acetate as an aggregation pheromone
in *Drosophila melanogaster*. Journal of Chemical Ecology. 1985;11(12):1747-56.

58. Verschut TA, Ng R, Doubovetzky NP, Le Calvez G, Sneep JL, Minnaard AJ, et al.
Aggregation pheromones have a non-linear effect on oviposition behavior in *Drosophila*
melanogaster. Nature Communications. 2023;14(1):1544.

59. Auld JR, Agrawal AA, Relyea RA. Re-evaluating the costs and limits of adaptive
phenotypic plasticity. Proceedings of the Royal Society B: Biological Sciences.
2009;277(1681):503-11.

60. DeWitt TJ. Costs and limits of phenotypic plasticity: Tests with predator-induced
morphology and life history in a freshwater snail. Journal of Evolutionary Biology.
1998;11(4):465-80.

61. Markow TA, Beall S, Matzkin LM. Egg size, embryonic development time and
ovoviviparity in *Drosophila* species. Journal of Evolutionary Biology. 2009;22(2):430-4.

62. Fernández-Moreno MA, Farr CL, Kaguni LS, Garesse R. *Drosophila melanogaster* as a
model system to study mitochondrial biology. Methods in molecular biology (Clifton, NJ).
2007;372:33-49.

63. Parker GA. Sperm competition games: Raffles and roles. Proceedings of the Royal
Society B: Biological Sciences. 1990;242(1304):120-6.

64. Bretman A, Fricke C, Chapman T. Plastic responses of male *Drosophila melanogaster*
to the level of sperm competition increase male reproductive fitness. Proceedings of the
Royal Society B: Biological Sciences. 2009;276(1662):1705-11.

65. Moatt JP, Dytham C, Thom MDCP. Exposure to sperm competition risk improves
survival of virgin males. Biol Lett. 2013;9(2):20121188.

66. Bretman A, Westmancoat JD, Chapman T. Male control of mating duration following
exposure to rivals in fruitflies. Journal of Insect Physiology. 2013;59(8):824-7.

67. Harrison LM, Churchill ER, Fairweather M, Smithson CH, Chapman T, Bretman A.
Ageing effects of social environments in 'non-social' insects. Philosophical Transactions of the
Royal Society B: Biological Sciences. 2024;379(1916):20220463.

68. Billeter JC, Bailly TPM, Kohlmeier P. The social life of *Drosophila melanogaster*.
Insectes Sociaux. 2024.

69. Wolfram Research Inc. Mathematica. Champaign, Illinois: Wolfram Research, Inc.;
2023.

70. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nature methods. 2012;9(7):671-5.

71. R Core Team. R: A language and environment for statistical computing. 2019.

72. Wickham H. ggplot2: Elegant graphics for data analysis: Springer-Verlag New York;
2016.

73. Wilke CO. cowplot: Streamlined plot theme and plot annotations for 'ggplot2': R
package version 1.1.1; 2020.

74. Hope RM. Rmisc: Ryan miscellaneous: R package version 1.5.1; 2022.

75. Kleiber C, Zeileis A. Applied econometrics with R: Springer-Verlag New York; 2008.

76. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using
lme4. Journal of Statistical Software; Vol 1, Issue 1 (2015). 2015.

77. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed
Effects Models. Journal of Statistical Software. 2017;1(13).

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and/or analysed during the current study are attached for review as

supplementary material and will be openly available from the Leeds Research Data Repository

on publication.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by NERC (NE/T007133/1).

Author's contributions

Conceptualisation: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, Marco Archetti, Douglas

W. Yu, Andrew F.G. Bourke, Tracey Chapman, and Amanda Bretman. *Data curation*: Emily R.

Churchill, Emily K. Fowler, and Lucy A. Friend. *Funding acquisition*: Emily K. Fowler, Marco

Archetti, Douglas W. Yu, Andrew F.G. Bourke, Tracey Chapman, and Amanda Bretman.

Investigation: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, and Marco Archetti.

Methodology: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, and Marco Archetti. *Project*

administration: Emily R. Churchill, Emily K. Fowler, Lucy A. Friend, Tracey Chapman, and

Amanda Bretman. *Resources*: Tracey Chapman and Amanda Bretman. *Software*: Emily R.

Churchill and Marco Archetti. *Visualisation*: Emily R. Churchill and Marco Archetti. *Writing –*

original draft: Emily R. Churchill. *Writing – review and editing*: Emily R. Churchill, Emily K.

Fowler, Lucy A. Friend, Marco Archetti, Douglas W. Yu, Andrew F.G. Bourke, Tracey Chapman,

and Amanda Bretman. All authors gave final approval for publication.

Acknowledgements

We would like to thank Paul Candon and Kerri Armstrong for technical assistance, and all

anonymous reviewers of this manuscript for their thoughtful and constructive feedback,

which great enhanced the quality of this manuscript.

Supplementary Information

Supplementary text 1. Determination of effect of natural egg placement and egg 'edge effects'

on subsequent egg laying behaviour

In this study, we tested the responses of females to the presence of existing eggs of various

cluster sizes in the environment. This required us to manually place eggs in vials to create the

required variation in egg clustering. Previous research shows that females naturally prefer to

lay eggs at the edges of substrates (9, 47, 48). Such edge biases could themselves influence

the decisions of subsequent females about whether to create egg clusters. To test for this,

and thus understand how best to place eggs in the main experiment, we conducted an initial

preliminary experiment to test whether proximity of existing eggs to the edge off the vial

affected the subsequent latency of females to lay eggs near them.

We tested isolated versus grouped females laying eggs in 3 types of 'egg treatment' vials: in

which there was (i) no egg present (control), and (ii) 1 egg either placed in the centre or (iii)

at the edge of the vial. The non-focal eggs used to create the egg treatment vials were laid by

groups of 10 gravid females allowed to lay overnight on Petri dishes filled with 40ml of SYA

medium. To standardise eggs, we selected only eggs laid naturally in isolation from these

dishes for subsequent transfer to create the egg treatment vials. To enable transfer of eggs

without damage to the SYA medium, elected eggs were transferred using a mounted needle,

into standard plastic vials that had been cut in half. Then the top half was reattached after

egg transfer using Sellotape. Test flies were collected from standard density vials (as in the

main experiments) and unmated females were housed in solitude, and males in groups of

four. After six days, one male was transferred into the vial with each female, and pairs were

watched to ensure matings occurred. We then transferred females in isolation or in groups of

4 into vials containing the 3 egg treatments (no egg, 1 edge egg, or 1 central egg; N = 15 - 42

vials per treatment).

We then checked vials every 30 minutes until 21:30 (one count after lights off in the constant

environment room) and removed the female(s) if they had laid eggs. If they did not lay, they

were kept overnight, and egg counts resumed at 09:30 for three further hours, until 24 hours

after females had initially been introduced into the vials. We calculated egg laying latencies

and the number of eggs laid by the focal females. We also noted whether eggs were laid in a

cluster (defined in Fig. 1) alongside the existing egg.

The results revealed a strong preference to lay at the edge of the vial, confirming previous

reports. There was no effect of existing egg location (X^2 = 1.32, d.f. = 113, p = 0.517) or group

size (X^2 = 3.80, d.f. = 112, p = 0.0513) on where in the environment females subsequently laid

their eggs. There was also no effect of either egg location or group size on whether females

chose to cluster their eggs as no females clustered their eggs (Table S1). Females housed in

solitude took longer to initiate laying than did those in groups of four (X^2 = 4.38, d.f. = 126, p

= 0.001; Fig. S2). There was no effect of existing egg location on laying latency (X^2 = 1.30,

d.f. = 125, p = 0.139; Fig. S2). Based on these results, we created egg clusters in the main

experiments by following the natural patterns by placing eggs at the vial edges.

     Figure S1. Experimental set up for testing the effects of social densities on egg laying

Figure S1. Summary of the experimental set up for testing the effects of social densities on egg laying. Day 0 –

4: males and females housed in same-sex groups of 10. Day 5 – 6: females transferred to one of four social

treatments (whilst males remained in groups of 10). Day 6: males were added to female treatment vials in a

ratio of 3:2 (2:1 for isolated females) and given 2h to mate. Day 6 – 7: number of eggs, number of clusters and

size of clusters were counted at the six listed timepoints. Shown is the strategy for defining egg clusters (contact

between the main body of eggs).

Figure S2. Existing egg location did not impact latency to lay

Figure S2. Females housed in solitude were slower to start laying than those housed in groups of 4, but there

was no effect of existing egg location on laying latency. Means and standard errors are shown for the six

treatments. Grey box indicates period of dark (21:00 – 09:00 GMT). N = 30 vials for all four social treatments.

Figure S3. Females in larger social groups laid significantly larger clusters

Figure S3. Maximum cluster size increased with increasing female social density. Shown is a frequency

distribution of the largest egg cluster for mated females placed in social groups of 1, 2, 4, and 8 for 24 hours.

Clusters were defined as in Fig. 1. Data are shown as the total number of vials with each cluster size observed

for each of the four treatments (solitary females: N = 11; pairs: N = 18; groups of 4: N = 29; groups of 8: N = 30

vials).

Figure S4. Females in larger social groups laid significantly more clusters

Figure S4. A greater number of clusters were observed with increasing female social density. Shown is a

frequency distribution of the egg clustering sizes for mated females placed in social groups of 1, 2, 4, and 8 for

24 hours. Clusters were defined as in Fig. 1. Data are shown as the total number of vials with each cluster size

observed for each of the four treatments (solitary females: N = 11; pairs: N = 18; groups of 4: N = 29; groups of

8: N = 30 vials).

Figure S5. No evidence of fitness effects of social environment

Figure S5. There was no effect of social group size on egg-adult viability (solitary: N = 20 vials; paired: N = 24

vials; groups of 4: N = 29 vials; groups of 8: N = 30 vials). Means (large dot) and standard errors are shown in

black.

Table S1. Edge effect egg laying location decisions were not overridden by adult social group

size or existing egg locations

Table S1. Effect of social density and previous egg laying patterns on female fecundity and egg clustering. The proportion of vials where eggs laid were observed to be clustered with the existing egg (main bodies in direct contact) and laid at the edge of the vial (within 0.5cm from the vial circumference).

		Proportion of eggs laid at the edge (%)	Proportion of eggs clustered (%)
Social treatment	Egg location		
Solitary females	None (N = 33 vials)	100	-
	Edge (N = 22 vials)	100	0
	Centre (N = 15 vials)	100	0
Groups of four	None (N = 17 vials)	94.1	-
	Edge (N = 17 vials)	93.8	0
	Centre (N = 13 vials)	100	0

869 *Table S2. Egg laying patterns of females were significantly non-random*

870 Table S2. Comparison of the clustering proportion model outputs and empirical egg laying patterns observed in

871 *D. melanogaster*. Shown are Kolmogorov-Smirnov P value outputs for the comparison of empirical data to

872 expected clustering preference values (К = 0 to К = 1, in 0.1 increments) rounded to two decimal places. Values

873 that show the most likely clustering preference are highlighted in bold.

Social treatment	K values										
	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
Solitary N = 20 vials	0.09	0.34	0.44	**0.48**	0.46	0.29	0.17	0.06	0.02	0.03	0.01
Paired N = 25 vials	0.01	0.20	0.34	0.43	**0.43**	0.38	0.22	0.15	0.07	0.04	0.01
Groups of four N = 29 vials	0.00	0.08	0.16	0.25	**0.29**	0.29	0.23	0.13	0.05	0.02	0.00
Groups of eight N = 30 vials	0.00	0.00	0.07	0.15	0.20	**0.21**	0.15	0.08	0.03	0.01	0.00