1   **Taxonomy, nomenclature, and identification of the giant hummingbirds (*Patagona spp.*)**

2   **(Aves: Trochilidae)**

Jessie L. Williamson[1–3*], Chauncey R. Gadek[3–5], Bryce W. Robinson[1,2], Emil Bautista[6],

Selina M. Bauernfeind[3], Matthew J. Baumann[3], Ethan F. Gyllenhaal[3,4], Peter P. Marra[7],

Natalia Ricote[8], Nadia D. Singh[9], Thomas Valqui[6,10], and Christopher C. Witt[3,4]

[1]Cornell Lab of Ornithology, Cornell University; Ithaca, 14850, USA.

[2]Department of Ecology and Evolutionary Biology, Cornell University; Ithaca, 14850, USA.

[3]Museum of Southwestern Biology, University of New Mexico; Albuquerque, 87131, USA.

[4]Department of Biology, University of New Mexico; Albuquerque, 87131, USA.

[5]Environmental Stewardship, Los Alamos National Laboratory; Los Alamos, 87545, USA.

[6]Centro de Ornitología y Biodiversidad (CORBIDI); Lima, 15064, Peru.

[7]The Earth Commons Institute; Department of Biology; McCourt School of Public Policy,

Georgetown University; Washington, D.C., 20057, USA.

[8]Facultad de Artes Liberales, Departamento de Ciencias, Universidad Adolfo Ibáñez;

Santiago, 7941169, Chile.

[9]Department of Biology, Institute of Ecology and Evolution, University of Oregon, Eugene,

97403, USA.

[10]Facultad de Ciencias Forestales, Universidad Nacional Agraria La Molina; Lima, 15024,

Perú.

Corresponding author: Jessie Williamson, 159 Sapsucker Woods Road, Ithaca, NY 14850;

707-287-8486; jlw432@cornell.edu

Abstract

Giant hummingbirds (*Patagona* spp.) are extraordinarily large hummingbirds whose

taxonomy has been muddled for two centuries. *Patagona* systematics were recently redefined

in a study of migration, physiology, and genomics, revealing two species: the Southern Giant

Hummingbird and Northern Giant Hummingbird. Here, we re-evaluate taxonomy and

nomenclature of the genus in light of its newly-clarified biology and species limits, analyzing

data from 608 specimens and wild-caught individuals spanning 1864–2023. The forms *gigas*

and *peruviana* were both described based on multiple syntypes. No adequate syntypes for *P.*

gigas are extant, so we designate a neotype for this taxon. We then critically consider the

identity and usage of *gigas* and *peruviana*, respectively, and examine identification

challenges that have fostered taxonomic uncertainty. We endorse the names *Patagona gigas*

for the Southern Giant Hummingbird and *P. peruviana* for the Northern Giant Hummingbird.

The genetic identity of the *peruviana* lectotype remains untested, but its plumage appears to

match the northern species. We found that ~33% of *Patagona gigas* specimens in major

museum collections are misidentified as *peruviana*; we provide this list to correct the

historical record. To facilitate identification and future study of these two cryptic species, we

provide comprehensive information on plumage, measurements, and seasonal ranges.

Key words: Andes, bird, biodiversity, cryptic species, genus revision, identification key,

intraspecific variation, nomenclature, morphological comparison, taxonomic revision.

 INTRODUCTION

Giant hummingbirds (*Patagona spp.*) comprise the most distinctive clade in the family

Trochilidae because of their extraordinary size. Within the genus *Patagona*, however,

morphological and plumage variation are subtle and it is difficult to disentangle geographic

variation from age, wear, molt, sex and/or individual variation. Seasonal overlap among

cryptic but distinct geographic breeding populations has exacerbated the challenge of

characterizing the differences among genetic lineages and appropriately defining taxonomy.

A recent study made it possible to accurately adjudicate taxonomy within *Patagona*

for the first time by presenting new information about genetics, migration, and phenotypes

(Williamson et al. 2024; Fig. 1). We now know that *Patagona* is a clade with two biological

species that diverged ~2–3 Mya and that differ in breeding latitude and migratory behavior.

The two lineages are largely allopatric during the breeding season. During the austral winter,

the southern lineage, which breeds from central Chile through northwestern Argentina and

likely throughout Bolivia, migrates to tropical latitudes of the central Peruvian Andes where

it overlaps extensively with the northern, high-elevation resident lineage. The difficulty of

distinguishing the two cryptic species based on plumage and/or measurements has, until now,

made it difficult to characterize their seasonal range overlap. However, the two lineages are

reproductively isolated and experience no gene flow, making their status as biological species

clear (Williamson et al. 2024). Each of the two species, the Northern Giant Hummingbird and

Southern Giant Hummingbird, contains shallow geographic population structure. The high-

elevation resident Northern Giant Hummingbird averages larger in bill length, wing chord,

tail length, and tarsus length, although there is substantial overlap in morphological

characteristics with the migratory Southern Giant Hummingbird—only ~65–85% of

individuals can be identified by measurements when comparing sexes separately. The two

species have near-identical body mass distributions. Similarly, subtle plumage differences

between the northern and southern species are useful for identification but not universally

applicable across the range of individual variation (Fig. 1).

With unambiguous species status and newly defined taxon limits, Williamson et al.

(2024) proposed a new species name for the Northern Giant Hummingbird, picaflor gigante

del norte in Spanish: *chaski* (*Patagona chaski* sp. nov.) and designated a holotype. The name

peruviana had been introduced by Boucard (1893) in a brief description, and has since been

applied at the subspecies level to all giant hummingbirds found from northwestern Argentina

and Bolivia northward. Williamson et al. (2024) argued that the name *peruviana* should not

continue to be used for two main reasons: (1) Historic and continued misalignment of the

name *peruviana* with the northern genetic lineage; and (2) The type series (syntypes)

appeared to have been comprised of a mix of the two giant hummingbird species.

However, the taxonomic description by Williamson et al. (2024), overlooked key

information regarding the syntypes of *peruviana* that came to the attention of the authors only

after publication; an omission that we rectify here. Namely, Williamson et al. (2024) did not

know that the likely identities of syntypes used by Boucard (1893) to designate *peruviana*

were known. Although Boucard did not provide identifiers for the specimens he referenced in

his description, he stated that they were "…collected by Whitely in Peru" (Boucard 1893). At

least 16 specimens were collected by Henry Whitely, Jr. in southern Peru from 1867–1873 or

later (Supplement; Table S1). Williamson et al. (2024) were not aware of two specimens still

at the Muséum National d'Histoire Naturelle in Paris (MNHN-ZO-MO-1989-340 and

MNHN-ZO-MO-1989-341) which are considered to be Boucard's (1893) *peruviana* syntypes

and are labeled accordingly (e.g., Zimmer 1930, Hellmayr 1932, Jouanin 1950; Fig. S1a-b;

Table S1). These syntypes appear to correspond to the Northern Giant Hummingbird based

on plumage features, although the female specimen is somewhat atypical. The identifications

of both should be considered tentative until confirmed with DNA sequencing. Shortly after

Williamson et al. (2024) was published, Areta et al. (2024) designated MNHN-ZO-MO-1989-

340 as a lectotype for *peruviana*. This quick taxonomic action by Areta et al. (2024) sought to

affirm the link between the Northern Giant Hummingbird and the name *peruviana*.

A guiding purpose of the International Code of Zoological Nomenclature (the Code)

is "to promote stability and universality in the scientific names of animals and to ensure that

the name of each taxon is unique and distinct", thereby ensuring precision and consistency in

the use of terms (ICZN 1999). Williamson et al. (2024) stated that they proposed the species

name *chaski* because the name *peruviana* had been misapplied to a mixture of two species,

both the Northern and Southern Giant Hummingbird, and that the species identities of the

specimens listed in the description were dubious. Because the newly-designated *peruviana*

lectotype appears to correspond to the Northern Giant Hummingbird, the names *chaski* and

peruviana should be considered synonyms, with *peruviana* taking precedent because it is

senior. Art. 23.1 of the Code, the Principle of Priority, states the general principle that "the

valid name of a taxon is the oldest name applied to it…" (ICZN 1999).

Here, we present the taxonomic history and nomenclature of *Patagona* from 1824 to

present, elaborating on Williamson et al. (2024) and Areta et al. (2024) and correcting errors

in both papers. We review new information about species limits, genetics, migration, and

identification that clarifies seasonal ranges and systematics of this genus. We additionally

evaluate the first species description by Vieillot (1824) and that of the genus *Patagona* Gray

1840, and examine those of subsequently introduced taxon names, *peruviana* and *boliviana*

by Boucard (1893), as well as the specimens that were key to historical treatments and

nomenclature. Finally, we designate a neotype to fix the identity of the Southern Giant

Hummingbird.

120   **MATERIALS AND METHODS**

     Abbreviations

AMNH, American Museum of Natural History, New York, USA; ANSP, Academy of Natural

Sciences of Drexel University, Philadelphia, USA; CM, Carnegie Museum of Natural

History, Pittsburgh, USA; CORBIDI, Centro de Ornitología y Biodiversidad, Lima, Peru;

CUMV, Cornell University Museum of Vertebrates, Ithaca, USA; FMNH, Field Museum of

Natural History, Chicago, USA; LSUMZ, Louisiana State University of Museum of Natural

Science, Baton Rouge, USA; Museum of Comparative Zoology at Harvard University,

Cambridge, USA; MNHN, Muséum National d'Histoire Naturelle, Paris, France; MSB,

Museum of Southwestern Biology, Albuquerque, USA; NHMUK, Natural History Museum,

Tring, UK; USNM, Smithsonian National Museum of Natural History, Washington, D.C.,

USA.

Taxonomic history of *Patagona gigas*

The species *Trochilus gigas* was described by Vieillot (1824) based on two syntypes from

"Brésil" (Brazil) (Vieillot 1824). Vieillot wrote that one specimen "…appears to us to be only

a young male; the old one is unknown to us; but we believe we recognize the female in an

individual in the possession of Mr. Becœur, naturalist…" (Vieillot 1824). Lesson (1829)

renamed the species *Ornismya tristis*, noting that its "homeland" was Chile (Lesson 1829). In

1840, Gray named the genus *Patagona*, listing *P. gigas* (ascribed to Vieillot, 1824) as the sole

species (Gray 1840). In spite of other proposed names (e.g., *Hylocharis gigas*, *Ornismya*

gigantea, *Trochilus tristis*, *Cynanthus tristis*, and *Hypermetra gigas*; see Elliot 1879, Porter

1897, Hartert 1900), from this point forward, *Patagona gigas* entered into widespread use

among ornithologists.

Vieillot's putative "young male" syntype, illustrated by Paul Louis Oudart in Plate

180 of *La Galerie des Oiseaux* and attributed to M. (Mr.; for "monsieur" in French) Portier of

the Ministry of the Navy (Vieillot 1824; Fig. S2) has not been located and its current

whereabouts are unknown to us. Vieillot's putative female syntype is believed to be

NHMUK-1933.11.14.36, a taxidermied mount at NHMUK (Warren 1966; Fig. S3). This

specimen was obtained by George Loddiges from Jean-Baptiste Bécouer, a pharmacist and

collector known for his arsenical soap method of specimen preparation, who died in 1777

(Rookmaaker et al. 2006). It is not known whether NHMUK-1933.11.14.36 was determined

to be a female during dissection or whether sex was presumed based on plumage (the latter

would be unreliable). This putative female syntype bears two sets of confusing labels that

make its identity as a syntype tentative; neither label is original, and both were written at

different times and in different hands (date and author unknown): One is a green tag that

appears to state "Trochilus gigas Vielliot female / gal. ois. 1.1e 29 p. 296 pl 180 / Loddiges

Coll / May be type of female" (note misspelling of "Vieillot"; Fig. S3). The other tag, a white

British Museum label, reads on the front, "Trochilus gigas female / Loddiges Colln.

(Purchased)" and on the back states, "1933-11-14-36. / ~~Not accepted as Viellot's type of~~

~~female but~~ may be. See Type catalogue" (again note misspelling of "Viellot"). The line

through the text "Not accepted as Viellot's type of female but" was done in blue ink at what

appears to be a later date (Mark Adams, *pers. comm.*; Fig. S3). Because specimens from the

Loddiges collection, purchased by NHMUK in 1933, bore no original labels, it is not possible

to verify that they specimens are, indeed, the type specimens of the species they represent;

thus, Loddiges specimens are listed by NHMUK as probable holotypes or syntypes (Warren

1966).

Despite the unknown collection dates and provenance (including both specific locality

and higher geography) of Vieillot's two putative syntypes, much later, in 1932, Austrian

ornithologist and neotropical bird expert Carl Hellmayr suggested that the (syn)type locality

of Brazil described by Vieillot (1824) was an "errore", stating "…we suggest Valparaíso,

Chile" (Hellmayr 1932). On pg. 231 of the *Birds of Chile*, Hellmayr justified his type locality

restriction, writing, "Oudart's plate of *T. gigas*, based on an evidently immature bird from

"Brésil" in the collection of 'M. Portier, attaché au ministère de la marine,' while none too

good, corresponds fairly well to certain bright-colored Chilean specimens, such as No.

61,676, Caldera, and accordingly I propose to restrict Vieillot's term to the small southern

form suggesting Valparaiso [, Chile] as type locality" (Hellmayr 1932). Hellmayr stated that

Oudart's plate illustrated an "immature" bird, but it actually depicts a juvenile (immature

birds are subadults which differ in plumage from juveniles and resemble adults in

measurements and plumage). The specimen "No. 61,676" referenced by Hellmayr is FMNH-

61676, an immature female from Caldera, Atacama Region, Chile, corresponding in plumage

to the Southern Giant Hummingbird (we confirmed this by examining the specimen at

FMNH). Thus, Hellmayr's decision to restrict the specific (syn)type locality for *Patagona*

gigas to Valparaíso, Chile, was based on his comparison of an immature specimen (FMNH-

61676) from Atacama Region, Chile (unconnected to the syntypes), with the Oudart's

illustration of a juvenile putative male from an unknown locality. This restriction of the type

locality based on visual evidence was unjustified by any known geographic variable trait.

Further confusing the identity of putative (syn)types, in 1893, Adolph Boucard, a

French ornithologist and specimen trader, wrote, "I have in my collection what I consider as

the type of Vieillot 'Ex Coll Riocour'" (Boucard 1893). Boucard references obtaining

Vieillot's type specimen from the Comte de Riocour collection, which he procured most of in

1889 (Renshaw 1905). The specimen he mentions is MNHN-ZO-2023-396 (sex and locality

unknown; collection date prior to 1842) at the MNHN, which bears multiple tags, including

Comte de Riocour tags with characteristic red outlines and Boucard Collection tags with

characteristic purple outlines (Fig. S4). The Comte de Riocour tag reads, "Ex. Coll Riocour /

Oiseau mouche geánt / *Trochilus giganteus* / Vieill". The front side of the Boucard Collection

tag reads, "Patagona gigas / Ex. Col. Riocour Type Vieillot ?" Boucard's tag includes a

question mark ("?") after the words "type Vieillot" (Patrick Bousses, *pers. comm.*), indicating

doubt about whether the specimen was, indeed, a Vieillot (syn)type; Boucard seems to have

additionally indicated this doubt in his text by writing "…what I consider the type…"

(Boucard 1893). However, we could find no evidence that MNHN-ZO-2023-396, which

Boucard believed was "Vieillot's type", is *the* type or even a syntype of *P. gigas*, nor could

we find any indication that this specimen was ever examined by Vieillot. Vieillot described *P.*

gigas based on two syntypes: the "young male" (whereabouts unknown) from M. (Mr.)

Portier of the Navy and a putative adult female (likely NHMUK-1933.11.14.36), obtained by

Jean-Baptiste Bécouer and then by George Loddiges before being purchased by NHMUK

(Warren 1966). The specimen MNHN-ZO-2023-396 was obtained by Boucard from the

Comte de Riocour collection and resides at the MNHN. There is no evidence that Boucard (or

Comte de Riocour before him) ever had access to George Loddiges' collection, which

remained within the Loddiges family after George Loddiges' death in 1846 (Mark Adams,

pers. comm.). Consistent with this history, the specimen NHMUK-1933.11.14.36 does not

bear original Boucard Collection tags, which would be expected had this specimen been "in

my collection", as Boucard described.

Thus, the entity formerly known as the Giant Hummingbird (*Patagona gigas*) and

currently known as the Southern Giant Hummingbird (*Patagona gigas, sensu* Williamson et

al. 2024) was described based on a putative type series consisting of two to three presently

unconfirmed syntypes with unknown collection dates, localities, and provenance, and whose

history is uncertain.

219   **Taxonomic history of *peruviana* and *boliviana***

In 1893, Boucard proposed the names *Patagona peruviana* and *Patagona boliviana* for

specimens in his collection, "…if the two should prove distinct species" (Boucard 1893).

Boucard wrote, "…I have also three specimens collected by Whitely in Peru, and in Bolivia

by Buckley" (Boucard 1893; Supplement; Fig. S1). In total, Henry Whitely collected at least

16 giant hummingbird specimens (not the 13 mentioned in Areta et al. 2024; Table S1).

Boucard's original text does not list specific identifiers, collection dates, collection localities,

or sexes for the specimens in his collection from Chile, Peru, or Bolivia. However, two

Whitely specimens at the MNHN (MNHN-ZO-MO-1989-340, a male from Tinta, Peru

collected 15 June 1868 and MNHN-ZO-MO-1989-34, a female from Tinta, Peru with no

date), were assigned equal status as *peruviana* syntypes, while MNHN-ZO-MO-1989-342

(undetermined sex from Bolivia with no date), collected by Clarence Buckley, was

designated as the holotype for *boliviana* (Jouanin 1950; Fig. S1; Table S1). The names

peruviana and *boliviana*, in addition to the nominate *gigas*, were henceforth recognized and

entered into use (e.g., Hartert 1900).

In 1921, Boucard's (1893) original description was contested by Eugène Simon, a

French naturalist and systematist from the MNHN. Simon wrote that *Patagona gigas* is a bird

of the high mountains "…without presenting sufficient local variation to be considered as

subspecies" (Simon 1921). Simon stated that Boucard (1893) had "…conditionally proposed

the name *Patagona peruviana* for the specimens from Peru from Whitely's trip and *Patagona*

boliviana for those brought back from Bolivia by Buckley". He then added, "…I have

ascertained, by examining the types, that the coloration characters [Boucard] gives (in Gen.

Humm. Birds. p. 61) are age-related or even completely illusory" (Simon 1921). In one

footnote, Simon noted the white borders on the wing feathers and wing coverts and "the

absence of a rufous part in front of the white uropygial band" on Boucard's types, which are

characteristics of *Patagona* juveniles and immatures (see Fig. 1). In a second footnote, Simon

clarified his description of the phrase "completely illusory", noting that he was discussing,

"…for example, the partly black throat of Peruvian birds" (Simon 1921). Simon did not

provide specific identifiers of the supposed types, but it is presumed that he examined the

same specimens referenced by Boucard (1893).

In 1930, American ornithologist John Zimmer provided a brief account of *Patagona*

gigas peruviana Boucard and *Patagona gigas boliviana* Boucard in *Birds of the Marshall*

field Peruvian expedition 1922-1923 (Zimmer 1930), recognizing Boucard's names as

subspecific identifiers. Zimmer described examining four recently-collected *Patagona*

specimens from Peru, compared against 23 previously-collected Peru specimens and 13 *gigas*

specimens from central and northern Chile and northwest Argentina. Zimmer wrote, "The

Peruvian birds, with one or two exceptions, are consistently larger than the Chilean skins,

with longer wings and longer, heavier bills. They also, with one or two exceptions, are more

rufescent on the under parts, sometimes markedly so. Since Dr. Hellmayr plans to discuss the

characters of these two races in a forthcoming paper on the birds of Chile, I will leave further

details to his able pen" (Zimmer 1930).

Two years later, Hellmayr published *The Birds of Chile* (Hellmayr 1932). In his

account for the giant hummingbird, he wrote, "Although the late Eugène Simon (Hist. Nat.

Troch., p. 157) questioned the possibility of discriminating any geographic races of the Giant

Humming bird, the study of between fifty and sixty properly labeled specimens from the

whole range clearly indicates the existence of two forms" (Hellmayr 1932). Hellmayr said

Simon's assessment that *P. peruviana* and *P. boliviana* were *nomina nuda* was a mistake, but

Hellmayr agreed with Simon when he noted "…several of the characters claimed by Boucard

prove to be individual" (Hellmayr 1932). Hellmayr went on to describe plumage differences

among populations from Chile compared to those from Ecuador, Peru, and Bolivia, noting

that the extent and intensity of rufous color to the underparts seemed to be "purely

individual" and unrelated to sex or age (Hellmayr 1932). Hellmayr concluded, "While the

palest examples of the northern form can be closely matched by one or two unusually rufous-

bellied birds from central Chile, the general run of the two series is easily told apart"

(Hellmayr 1932). His firm conclusion seemed to leave no doubt that there were two forms

that were easily distinguishable.

Based on his assessment, Hellmayr offered nomenclatural suggestions: "The larger

northern race is entitled to the name *P. gigas peruviana*, tentatively proposed by Boucard for

a specimen from Peru in his collection. The type, a male obtained by H. Whitely on June 15,

1868, at Tinta, Dept. Cuzco, agrees with the average of our Peruvian series, while *P.*

boliviana, was based on an individual variant with wholly cinnamon-rufous under parts…"

(Hellmayr 1932). Hellmayr thus cast doubt on *boliviana*, while buttressing *peruviana*.

Hellmayr noted that "…An adult male from Tilcara, Jujuy (Nov. 24, 1905; L. Dinelli), and a

couple from Lara, Tucuman (Feb. 12, 1903; G. A. Baer), all in the Tring Museum, are in

every particular typical of the large northern form" (Hellmayr 1932). Years later, Hellmayr

added that specimen no. 868/48 collected by L. Dinelli on August 10, 1916, from Colalao del

Valle, Tucumán, Argentina was "typical" of *peruviana* based on plumage and morphology

(Revilliod et al. 1942).

Hellmayr's 1932 account is largely regarded as the definitive taxonomic treatment of

Patagona in the literature. Based on his plumage and morphological analysis and

nomenclatural suggestions, subsequent authorities recognized *peruviana* as a valid subspecies

and ceased recognizing *boliviana*, folding *boliviana* into subsequent treatments of *P. g.*

peruviana (Peters 1945, Zimmer 1952)—despite the uncertainty surrounding the distribution

of forms.

In 1952, John Zimmer revisited *Patagona* taxonomy in *Studies of Peruvian Birds*

(Zimmer 1952). In his account of *P. g. peruviana*, he stated, "This northern form of the

species is fairly readily determinable in distinction from the southern *gigas gigas* when fully

adult birds are compared, but young examples can be confusing" (Zimmer 1952). Zimmer

went on to describe the difficulty of distinguishing juvenile birds by plumage. He noted, "All

the adult examples from [the Urubamba Valley of Peru] are clearly *peruviana*. I have seen no

Peruvian specimens that indicate the occurrence of *gigas gigas* in Perú, although there is

evidence that the subspecies has some migratory tendencies, as noted by Hellmayr…"

(Zimmer 1952). Zimmer presented a list of 83 specimens that he examined from numerous

localities and of both sexes. He wrote, "I would emend a statement made by Hellmayr (*loc.*

cit.) with reference to certain Argentina specimens that he found to be 'in every particular

typical of the large northern form.' Three females from Tilcara are fairly typical *peruviana*

but the others are identifiable only by their measurements, since they are immature and in

coloration resemble other young of both forms" (Zimmer 1952). Later, in 1974, Ecuadorian

biologist and naturalist Fernando Ortiz-Crespo supported previous scholars, noting "[The

race *peruviana*]…differs from the nominate race, occurring in central Chile, adjacent

Argentina, and the southern Chilean provinces, in having a larger body size, a stouter, longer

bill, and rufescent underparts" (Ortiz-Crespo 1974).

Thus, since ca. 1930, the giant hummingbird has been considered to contain two

forms: The nominate *P. g. gigas*, a suspected migrant and supposed smaller-bodied form with

a shorter bill and shorter wings, stated to range from approximately central Chile to central

Argentina; and *P. g. peruviana*, a supposed larger-bodied form with a longer bill and longer

wings, thought to range (and breed) from northwestern Argentina north through Bolivia,

northern Chile, Peru and Ecuador. Individuals of *peruviana* were believed to have more

rufescent underparts (although previous workers debated about whether the rufous coloration

might be related to age, sex, and/or individual variation). The two forms were described as

being "easily distinguishable" based on plumage and morphology. It is notable that numerous

historical treatments have parallels with the results of modern genomic and tracking studies,

despite those historical scholars having no access to DNA sequences. However, the early

treatments also introduced errors and greatly oversimplified the phenotypic distinction

between the two species, resulting in misidentification, errant inference of ranges, and failure

to recognize seasonal range overlap (see below).

Historical characterization of migration

The migration of Chilean giant hummingbirds has been suspected and discussed since at least

1834, but every aspect of the migratory journey had remained a mystery ever since.

Throughout history, naturalists puzzled over where Chilean populations "disappeared" to

after breeding (Goodall et al. 1946). Even recently, the *Atlas of the Breeding Birds of Chile*

2011-2016 called the winter whereabouts of Chilean-breeding giant hummingbirds an

"unsolved mystery" (Medrano et al. 2018). This paucity of migration data has contributed to

longstanding confusion about the ranges, taxonomy, and identification of the two giant

hummingbird forms.

Darwin was among the earliest to note migration of the Southern Giant

Hummingbird: During his voyage on *The Beagle*, he remarked, "At Valparaíso, in the year

1834, I saw several of these birds in the middle of August, and I was informed they had only

lately arrived from the parched deserts of the north" (Darwin 1839). German-Chilean botanist

Federico Johow wrote that giant hummingbirds are "rather tropical" and were observed in

central Chile only during the Chilean summer; he noted that their presence along the coast of

Chile coincided with the flowering of *Lobelia* and columnar cacti, proposing that they

wintered in the northern provinces of "…Peru, Ecuador, etc." (Johow 1910, 1921). Hellmayr

(1932) stated that *Patagona* is absent from central Chile during June–August, writing, "The

birds probably migrate northwards, but a few, at least, cross the Andes on their migration, as

is shown by specimens… which clearly pertain to the Chilean race" (Hellmayr 1932).

Concurring with this hypothesis, *Birds of the High Andes* posited that Chilean breeders

migrate across the Andes to adjacent western Argentina (Fjeldså and Krabbe 1990), while

others have suggested that they migrate to northwestern Argentina (Sánchez Osés 2003).

More recently, the guide "*Biodiversidad de Chile, patrimonio y desafíos*" asserted without

evidence that Chilean giant hummingbirds migrate to Ecuador for the austral fall (Ambiente

2008).

In 1952, Chilean naturalist Rafael Barros offered what is perhaps the most detailed

early treatment of *Patagona* migration: Barros described Chilean-breeding giant

hummingbirds as "…fleeing from the rainy season and the cold" (Barros 1952). He wrote, "It

is not well elucidated if, when it leaves its places of residence…it actually moves to the

provinces located to the north of Aconcagua (where my observations begin), or it crosses the

high mountain range of the Andes and passes to the Republic of Argentina, or if it goes to

winter in some regions of Bolivia or Peru. I know of no observations in this regard, and this

is a problem to be solved, of great scientific interest" (Barros 1952). Barros posited the

question: "Does it go up…to the heights of the Puna to cross the Andes?", later responding:

"Based on what I have been able to observe for many years… I do not believe that it crosses

the mountain range in the central part of the country, during its pilgrimages, either on its

outward journey or on its return, because in the mountainous region it first reaches the lower

parts, the pre-cordillera, the foothills of the Cordillera and fields beneath them, gaining

altitude afterwards" (Barros 1952). This same year, Zimmer cast doubt on the extent of

northward travel of southern temperate breeders, stating that he had seen no Peruvian

specimens suggestive of *P. gigas gigas* (Zimmer 1952).

In 1974, Fernando Ortiz-Crespo published a detailed treatment of the ecology of giant

hummingbirds in Ecuador. He noted that the giant hummingbird population in central Chile

"…builds up during the spring and summer", attributing their movement to northward

migration (Ortiz-Crespo 1974). Ortiz-Crespo credited Johow (1910) and Darwin (1839) as

his sources for migration information, both of whom had previously characterized Southern

Giant Hummingbird migration as unresolved. In this same study, Ortiz-Crespo analyzed

morphology between sexes and subspecies (which he assigned based on collection country),

stating that his results "...do not seem to confirm the possibility that Ecuador is regularly

visited on migration by birds which nest in central Chile, and whose winter quarters thus

remain to be discovered" (Ortiz-Crespo 1974). Twelve years later, in a different manuscript,

Ortiz-Crespo expressed continued uncertainty about *Patagona* migration, stating "…it is not

known for sure where they winter" (Ortiz-Crespo 1986). He described examining 180

specimens, including 156 specimens from outside of Chile, only four of which (ANSP-

145417 and FMNH-179394 from Bolivia, ANSP-167558 from Argentina, and FMNH-67518

from Peru) he said had measurements and coloration of the nominate form *gigas* (we

examined these same specimens at ANSP and FMNH, and have confirmed that they are

Southern Giant Hummingbirds). Ortiz-Crespo concluded, "This meager evidence suggests

that southern populations disperse northward after nesting" (Ortiz-Crespo 1986), providing

no additional specifics. He further suggested that examining specimens collected during the

austral fall and winter from northwest Argentina and Bolivia would be important for

understanding distribution and movement (Ortiz-Crespo 1986). Thus, although *Patagona*

migration had been discussed in the literature since ~1834, every aspect remained unknown

before Williamson et al. (2024).

Re-evaluating historical identification criteria for *peruviana*

Hellmayr's (1932) text is regarded as definitive for *Patagona* taxonomy, particularly with

regard to distinguishing *gigas* from *peruviana*. In it, Hellmayr examined morphological

characteristics (bill, wing, and tail lengths) from "…between fifty and sixty properly labeled

specimens from the whole range…". Hellmayr concluded that there are "clearly" two forms

and that "…the general run of the two series is easily told apart" (Hellmayr 1932). In his text,

he presented a table of measurements from 45 specimens from Ecuador (no localities), Peru,

Bolivia (one specimen; no locality), Argentina, and Chile. He suggested that *P. g. gigas* was

markedly smaller and did not overlap morphologically with *P. g. peruviana*, supporting his

conclusion that "the general run of the two series is easily told apart" (Hellmayr 1932). This

claim was reinforced by Areta et al. (2024), who stated that Hellmayr "provided a diagnosis

of *peruviana* on plumage and morphometry" that was "clearly mirrored" by the data provided

in Williamson et al. (2024).

However, Hellmayr's conclusions about the ease with which *gigas* and *peruviana*

could be distinguished contradict results of analyses by Ortiz-Crespo (1974) and Williamson

et al. (2024). Among 361 *Patagona* specimens from Ecuador, Peru, Bolivia, Argentina, and

Chile, Williamson et al. (2024) found substantial inter-species overlap in the same three

morphological characteristics studied by Hellmayr (Williamson et al. 2024). Northern and

Southern Giant Hummingbirds differed significantly, on average, in bill length, wing chord,

and tail length, respectively; however, average trait differences were extremely subtle, on the

scale of millimeters (e.g., ~1–5 mm)—size differences between the two species were not

typically distinguishable by eye, and the ranges of trait values for both species overlapped

substantially (Williamson et al. 2024). Similarly, within each giant hummingbird species,

sexes differed significantly—but not substantially—in nearly all measurements, despite lack

of sexually dimorphic plumage. When analyzing morphological data separately for each sex,

linear discriminant analyses using data from three traits (bill, wing, and tail) combined could

only distinguish adult female Northern from Southern Giant Hummingbird in 65.4–82.8% of

cases and adult males in 79.5–85.7% of cases (Williamson et al. 2024). The mass ranges of

the two species are near-identical (Williamson et al. 2024), indicating general body size

equivalence.

To understand Hellmayr's (1932) conclusion that the two forms are easily

distinguished, we compared Hellmayr's original data (*n*=45 adults) with published data from

Williamson et al. (2024), measurements from wild-caught Northern and Southern Giant

Hummingbirds captured in Peru during July–August 2023, and data from specimens

measured at ANSP, CM, FMNH, and USNM during June and August 2024 (*n*=608

individuals total). All individuals were measured by [ANONYMIZED AUTHOR INITIALS].

After removing juveniles and considering only individuals of known sex, locality, and age,

our dataset included *n*=448 adults.

We predicted that if Hellmayr's analysis of *peruviana* and *gigas* corresponded to the

northern and southern species, respectively, the properties of his data should be consistent

with trait distributions from our larger dataset of measurements from these two species. To

test this, we randomly subsampled our data without replacement to match Hellmayr's sample

sizes (i.e., we subsampled to *n*=8 for Southern males and *n*=4 for Southern females to match

Hellmayr's *gigas* sampling, and to *n*=25 for Northern males and *n*=8 for Northern females to

match Hellmayr's *peruviana* sampling), and generated 10,000 random datasets. For each of

the 10,000 resampled datasets, we calculated trait means and coefficients of variation (CVs),

generating expected distributions. We did this separately for each sex and species. We further

calculated the magnitudes of trait difference between the two giant hummingbird species,

generating expected distributions for differences in trait means. We then calculated the

probability that Hellmayr's data were compatible with each expected distribution, as would

be expected if specimens had been selected without bias from individuals of the two species.

444   **RESULTS**

A neotype for *Patagona gigas* (now the Southern Giant Hummingbird)

To fix the identity of the unique species now known as the Southern Giant Hummingbird

Patagona gigas (*sensu* Williamson et al. 2024), we designate as its neotype

MSB:Bird:56142; CHILE: Valparaíso Region, locality Algarrobo, 13°20.916' S, 71°38.294'

W, elevation 10 meters, collected 9 December 2018. Sex: female. Parts saved: partial

skeleton, lung, heart, muscle, liver, eye, brain, and blood. Prepared by Selina M. Bauernfeind;

personal catalog number SMB 516. Geolocator ID: BJ104, corresponding to annual migration

tracks (Williamson et al. 2024). Specimen record:

https://arctos.database.museum/guid/MSB:Bird:56142 (Fig. 2).

Why designate a neotype for *Patagona gigas*, rather than a lectotype from the

putative syntype series? Art. 72.2.1 states that the type series consists of "…all the specimens

on which the author established a nominal species-group taxon…in the absence of holotype

designation, or the designation of syntypes, or the subsequent designation of a lectotype, all

are syntypes and collectively they constitute the name-bearing type" (ICZN 1999). For *P.*

gigas, the putative syntypes that could be considered candidates for lectotype designation

may include: (1) Oudart's plate of a "young male" described by Vieillot (1824), the specimen

for which could not be located (Fig. S2); (2) the putative female voucher NHMUK-

1933.11.14.36 (catalogued by NHMUK as possibly being Vieillot's "adult female"; Fig. S3);

and (3) the voucher MNHN-ZO-2023-396 (which Boucard believed might be "Vieillot's

type"; Fig. S4).

There are several reasons to doubt that either of the vouchers NHMUK-1933.11.14.36

and MNHN-ZO-2023-396 is a syntype of *P. gigas*. First, neither specimen possesses original

tags—it is therefore impossible to confirm that they were used for Vieillot's (1824) original

description. It remains unknown whether NHMUK-1933.11.14.36 was the specimen

originally acquired by Bécouer and subsequently examined by Vieillot in 1824; and there is

no evidence that Vieillot ever examined MNHN-ZO-2023-396 (the latter specimen's history

and data are misaligned with Vieillot's original description: for example, MNHN-ZO-2023-

396 is sex unknown, yet Vieillot described an adult female; additionally, the MNHN does not

recognize this specimen as a syntype). Second, neither voucher has a date of collection.

NHMUK-1933.11.14.36 was obtained in 1933 after the purchase of George Loddiges'

collection; Loddiges, in turn, acquired the specimen from Jean-Baptiste Bécouer, who died in

1777, giving NHMUK-1933.11.14.36 an estimated but unconfirmed collection date prior to

1777. The MNHN-ZO-2023-396 specimen tags list no collection date for the skin, but the

specimen is catalogued with a date of "avant 1842" ("before 1842"); however, Vieillot

described *Patagona gigas* 18 years earlier (in 1824), and it is not possible to confirm that

MNHN-ZO-2023-396 was collected prior to this date. Third, both vouchers have completely

unknown provenance. There is no evidence that Bécouer ever traveled to Chile (or South

America) to acquire NHMUK-1933.11.14.36, nor any specimen in his collection

(Rookmaaker et al. 2006). The locality of MNHN-ZO-2023-396 is given only as "South

America". Hellmayr's (1932) restriction of the type locality of *P. gigas* to "Valparaíso, Chile"

was erroneous; it was based on the resemblance that he recognized between Oudart's plate of

a juvenile putative male and the immature female specimen FMNH-61676, collected

September 23, 1924 from Caldera, Atacama Region, Chile, which is ~830 km north of

Valparaíso, Chile. Neither Oudart's plate nor FMNH-61676 have any known link to

NHMUK-1933.11.14.36 or MNHN-ZO-2023-396 (and there is no evidence that Vieillot's

"young male" and "adult female" syntypes were collected at the same time or from the same

place). It is therefore not possible to confirm the identities of the putative syntype vouchers,

and thus, it is not possible to confirm that voucher syntypes are extant.

Oudart's plate of the "young male" is clearly indicated as part of Vieillot's syntype

series (Fig. S2). (Despite searching, we were unable to locate the voucher specimen of the

individual depicted in the plate). Art. 74.4 of the Code states, "designation of an illustration

or description of a syntype as a lectotype is to be treated as designation of the specimen

illustrated or described; the fact that the specimen no longer exists or cannot be traced does

not of itself invalidate the designation" (ICZN 1999). Oudart's plate of the "young male"

could therefore be considered an extant syntype and therefore a valid choice for a lectotype of

P. gigas. However, Oudart's plate depicts a juvenile and it is established that the Southern and

Northern Giant Hummingbird juveniles cannot be reliably distinguished by plumage

(Williamson et al. 2024). Furthermore, the plate, which Hellmayr (1932) noted was "…none

too good…", does not illustrate any diagnostic phenotypic characteristics of the Southern

Giant Hummingbird, the species which it is meant to denote. The locality, provenance, and

date of collection of the specimen voucher illustrated in the plate are additionally unknown,

so it cannot be verified whether it truly corresponds to the Southern Giant Hummingbird. The

collector of the specimen is also unknown. Thus, Oudart's "young male" (sex unconfirmed)

plate violates Recommendations 74C (and by extension, 73C) and 74E and is therefore

unsuitable as a lectotype for *P. gigas.* Art. 72.2 states, "…If no name-bearing type is believed

to be extant a neotype may be fixed…" (ICZN 1999, 2003).

In accordance with Art. 75.3 of the Code, there is an exceptional need to designate a

neotype for *P. gigas*. The designation of neotype specimen MSB:Bird:56142 satisfies the full

list of qualifying conditions of the Code: This designation is made with the express purpose

of clarifying both the taxonomic status *and* the type locality of the nominal taxon *P. gigas*

(Art. 75.3.1); the neotype specimen MSB:Bird:56142 is unequivocally identified as a

Southern Giant Hummingbird based on definitive nuclear and mitochondrial genomic,

physiological, morphological, and migration tracking data; these features clearly distinguish

MSB:Bird:56142 from individuals of the Northern Giant Hummingbird (Williamson et al.

2024; Art. 75.3.2); the archived specimen data and above neotype description are sufficient to

ensure recognition of the neotype, MSB:Bird:56142 (see Fig. 2; Art. 75.3.3); no name-

bearing types are believed to be extant for the aforementioned reasons (Art. 75.3.4); the

neotype MSB:Bird:56142 has phenotypic characteristics consistent with what is known of the

name-bearing syntypes originally described by Vieillot (1824) and subsequently described by

other authorities; the neotype MSB:Bird:56142 is additionally a female, matching Vieillot's

original description of one of the syntypes (Art. 75.3.5); the neotype MSB:Bird:56142 was

collected in Valparaíso Region, Chile, the type locality that Hellmayr (1932) intended to

designate, but which he had restricted in error (Art. 75.3.6); and, the neotype

MSB:Bird:56142 is housed at the Museum of Southwestern Biology at the University of New

Mexico, a recognized scientific and educational institution that maintains a research

collection and which possesses proper facilities for preserving name-bearing types that are

accessible for study (Art. 75.3.7). This neotype designation is in accordance with

Recommendation 75A. [The authors have sought feedback from the Working Group on Avian

Nomenclature to satisfy Recommendation 75B and will include the appropriate ZooBank

LSID after Working Group input] (ICZN 1999, 2003).

As stated in Art. 76A.2, "A statement of a type locality that is found to be erroneous

should be corrected". Art. 76.3 states, "The place of origin of the neotype becomes the type

locality of the nominal species-group taxon, despite any previously published statement of

the type locality". Thus, the type locality for the Southern Giant Hummingbird (*Patagona*

gigas) should henceforth be considered Algarrobo, Valparaíso Region, Chile, the collection

locality of the designated neotype, which corrects the previous erroneous type locality

restriction. The designation of MSB:Bird:56142 as the neotype fixes the identity of *Patagona*

gigas, ensuring nomenclatural stability (ICZN 1999, Jiménez-Mejías et al. 2024).

How an unknown migratory route led to a muddied taxonomy

Numerous published accounts, both historical and modern, have mentioned giant

hummingbird migration or presence/absence of *Patagona* at certain times of the year in Chile

with only sparse details (e.g., Gould 1861, Barros 1952, Aguirre 2001, González-Gómez and

Valdivia 2005, Estades et al. 2008, Speziale and Lambertucci 2009, Velásquez-Noriega et al.

2021, Medel et al. 2022, Rueda-Uribe et al. 2023). One publication stated that *Patagona*

migrates 2,000 km (Rodríguez-Flores et al. 2019), but no information about the migratory

route or sources for distance traveled were provided. Two previous studies used coarse-scale

citizen science data to estimate the magnitude of elevational shift experienced seasonally by

migratory populations (Williamson and Witt 2021a, Rueda-Uribe et al. 2023); both generated

what we now know are underestimates.

Lack of information about the migration of southern populations has led to persistent

uncertainty about the ranges of each giant hummingbird form. Throughout history, the giant

hummingbird range has generally been described as encompassing the Andes, but texts are

not consistent and have often omitted mention of Argentina (e.g., Gould 1861, Boucard

1893). In particular, the range limits of *peruviana* had never been described correctly.

Hellmayr (1932) wrote that "*P. g. peruviana* … replaces the typical form in the Andes of

Ecuador, Peru, Bolivia, and extreme northern Chile. It is probably also this race that nests in

northwestern Argentina, although I have not been able to examine undoubted breeding

specimens" (Hellmayr 1932). In 1945, Peters wrote, "The breeding bird of northwestern

Argentina (Jujuy, Catamarca, and Tucumán) is probably referable to [*P. g. peruviana*]"

(Peters 1945). In 1952, Zimmer identified several Southern Giant hummingbirds from

Argentina and Bolivia as *peruviana* (Zimmer 1952). The belief that the race *peruviana* occurs

in northwest Argentina has been supported by many key historical accounts (Barros 1952,

Zimmer 1952, Ruschí 1961, Koepcke 1983, Fjeldså and Krabbe 1990) and has continued to

persist into recent years (e.g., Sánchez Osés 2003, Pearman and Areta 2020, Velásquez-

Noriega et al. 2021). More recently, in *Birds of Chile: A Photo Guide*, Howell and Schmitt

(2018) stated that there is "no range overlap" between *peruviana* and *gigas* (Howell and

Schmitt 2018). However, we now know that northern Argentina contains only the Southern

Giant Hummingbird, and that the two species overlap substantially in range during the non-

breeding season.

Uncertainty about migration and confusion about the numbers of possible

subspecies/forms, and their respective distributions (e.g., Fjeldsa and Barbosa 1983), has

additionally resulted in misleading generalizations about elevations of residence. Woods

(1998) suggested that the two forms, *peruviana* and *gigas*, might be elevational replacements

and incompletely characterized the Southern Giant Hummingbird range, writing, "The

northern subspecies of Giant Hummingbird *P. g. peruviana* regularly reaches paramo and

puna habitats in Ecuador and Peru at 3,500-4,000 m, while the southern *P. g. gigas* occurs

seasonally to below 2,000 m in central Chile and adjacent Argentina" (Woods et al. 1998).

Ruschi (1961) stated that *P. g. peruviana* is found in the Andes, varying in altitude from

2,500 to 5,000 meters (Ruschí 1961) but he did not mention *P. g. gigas* or its range. In many

other texts, it is not possible to determine which form of giant hummingbird is being

discussed.

We now know that the giant hummingbirds are comprised of two phenotypically

cryptic species that overlap at the same sites and elevations during the southern hemisphere

winter, approximately May–September (Williamson et al. 2024). The Southern Giant

Hummingbird is a long-distance latitudinal migrant, traveling up to 8,300 km roundtrip in a

circular loop across the Andes (Williamson et al. 2024). At least some (i.e., Chilean) Southern

Giant Hummingbird populations shift >4,100 meters in elevation twice annually, making this

species one of ~100 taxa that undertake elevational niche-shift migration (ENSM), or

physiologically-taxing 'extreme' shifts in elevation between the breeding and non-breeding

seasons (Williamson and Witt 2021a, b; Williamson et al. 2024, Ivy and Williamson 2024).

Although Southern Giant Hummingbird populations from Argentina also spend the non-

breeding season at high elevations in central Peru, their migratory routes and degrees of

elevational shift have not yet been studied. Resolution of longstanding mysteries about giant

hummingbird migration was made possible with the use of miniaturized tracking devices and

genomic sequencing—technologies that were not available to Darwin, Johow, or the other

workers cited above—that has finally permitted clarification of taxonomy within this clade.

603   **The giant hummingbirds are not "easily" distinguished**

Contrary to Hellmayr's (1932) findings, our measurements of >600 specimens revealed

substantial overlap between species for bill, wing, and tail lengths within each sex (Fig. 3).

Hellmayr's *peruviana* measurements were significantly larger than our Northern Giant

Hummingbird measurements in sex-specific comparisons of wing and tail, respectively, and

female bill length (Fig. 3b–f). For each sex, Hellmayr's data also showed significant

differences between *gigas* and *peruviana* tail measurements, whereas we found none (Fig.

3c,f). When we conducted a Principal Components Analysis (PCA) using bill, wing, and tail

measurements separately for each sex, Hellmayr's data clustered tightly and were shifted to

the outer edges of the expected distributions (Fig. 4). In this same analysis, the MNHN male

peruviana lectotype (MNHN-ZO-MO-1989-340), an apparent Northern Giant Hummingbird,

and the *boliviana* holotype (MNHN-ZO-MO-1989-342), a Southern Giant Hummingbird of

unknown sex, both fell within Hellmayr's range for *peruviana*, and within a morphological

zone of overlap between male Northern and Southern Giant Hummingbirds; when analyzed

with female data, MNHN-ZO-MO-1989-342 was an outlier for both species (Fig. 4). The

MNHN *peruviana* female paralectotype (MNHN-ZO-MO-1989-341) fell more closely in line

with Southern Giant Hummingbird measurements, but its identity was not clearly identifiable

based on measurements (Fig. 4).

621    Hellmayr's differences in trait means between forms were noticeably larger, and CVs

in his dataset noticeably smaller, than in the expected distributions generated from 10,000

resampled datasets (Figs. S5–S6, Tables S2–S4). For all sex-trait combinations, his CVs and

differences in trait means were statistically improbable and unrealistically small (Figs. S5–

S6; Supplement). For example, the difference between Hellmayr's *peruviana* and *gigas* mean

wing measurements was ~2x greater for males and ~4x greater for females than the

respective differences we found between Northern and Southern Giant Hummingbirds (Fig.

3).

630      **Misidentification of specimens in historical texts and museum collections**

Many historical texts do not provide specimen identifiers but, in several instances, it is

possible to diagnose specimen identities using descriptive information provided. Hellmayr

(1932) writes of three specimens that "…are in every particular typical of the large northern

form…an adult male from Tilcara, Jujuy (Nov. 24, 1905; L. Dinelli), and a couple from Lara,

Tucumán (Feb. 12, 1903; G. A. Baer), all in the Tring Museum" (Hellmayr 1932). These

specimens were formerly at NHMUK but were moved to the AMNH in 1932 after Lord

Rothschild sold his collection (Lanyon 1995, Roselaar 2003; Mark Adams *pers. comm.*; Tom

Trombone, *pers. comm.*) The adult male from Tilcara described by Hellmayr is AMNH-

479595, definitively identified as a Southern Giant Hummingbird based on targeted sequence

capture of ultra-conserved elements (UCEs), known breeding season collection locality, and

diagnostic plumage characteristics (Williamson et al. 2024). The other two Lara, Tucumán

individuals mentioned are AMNH-479593 and AMNH-479594; both are juvenile Southern

Giant Hummingbirds, for which Williamson et al. (2024) also provide UCE data. The fact

that Hellmayr believed these individuals from Argentina were "in every particular typical" of

peruviana, when in fact, they are Southern Giant Hummingbirds ranging in age from

juveniles to adults highlights the difficulty of distinguishing the two forms by plumage.

Due to these challenges, many historic and modern Southern Giant Hummingbirds

(*gigas*) were misidentified as Northern Giant Hummingbirds (*peruviana*). The reverse does

not seem to be true: We have found no Northern Giant Hummingbirds misidentified as

Southern Giant Hummingbirds. At least eleven major museum collections whose specimens

we have examined possess giant hummingbird specimens from each Peru, Bolivia, Argentina,

and Chile that are catalogued or labeled as *P. g. peruviana*, and which contain a mix of both

the Northern and Southern Giant Hummingbird. Of the 603 total adult and juvenile

specimens with locality data whose catalog records and tags we analyzed, only 269

individuals (44.6%) are catalogued with subspecific identifiers. A total of 74 individuals

(32.74% of specimens) labeled "*peruviana*" are actually Southern Giant Hummingbirds (e.g.,

AMNH-139179, ANSP-120834, CM-52921, CUMV-52570, FMNH-67518, LSUMZ-37404,

MCZ-262196, USNM-529148; Fig. 5; Table S5). Two individuals (ANSP-145417 and CM-

94400) from Bolivia were labeled with both "*peruviana*" and "*gigas*" (on the tag of ANSP-

145417, pencil marks are visible where "*peruviana*" was erased and "*gigas*" was written; this

individual still bears "*peruviana*" on the back of its tag).

Typically, any individual collected in Ecuador, Peru, Bolivia, and northwestern

Argentina that appeared large and had reddish underparts was assigned to *peruviana* without

substantial scrutiny; by this measure, many rufous-toned *gigas* individuals, especially those

that seemed larger, would appear to correspond to *peruviana* (e.g., Fig. 1). The name

peruviana has thus been applied to specimens on the basis of oversimplified morphology,

plumage, and assumptions about range. However, the extent to which *gigas* specimens have

been misidentified (and thus, the name *peruviana* misapplied) in global collections is difficult

to ascertain because the vast majority of catalogued specimens (55.4%) do not have

subspecific identifiers.

Nomenclatural recommendations

The neotype designated here (MSB:Bird:56142) corresponds unambiguously to the Southern

Giant Hummingbird, and we thus recommend continued use of *Patagona gigas* for this

newly-recognized species. The name *peruviana* has a contentious history: Although it has

always intended to apply to the 'larger, northern form' of the giant hummingbird, there has

been longstanding uncertainty about how the 'larger, northern form' is defined, and about

how the name was applied to individuals of certain characteristics. Consequently, the name

peruviana has been applied, variably, to the two giant hummingbird species based on a

combination of erroneous or incomplete information about plumage, morphology, range, and

seasonal occurrence. The *peruviana* lectotype (MNHN-ZO-MO-1989-340) appears to

correspond in plumage to the Northern Giant Hummingbird. Contrary to Areta et al. (2024)'s

statement that MNHN-ZO-MO-1989-340 is "unambiguously" identifiable as the northern

taxon, there is reasonable room to doubt this plumage-only-based assessment and the identity

of the lectotype remains unconfirmed without genetic sequencing. However, following rules

set forth by the Code, the name *peruviana*, the current senior synonym, should be adopted for

the Northern Giant Hummingbird.

DISCUSSION

Historical oversimplification of complexity within and between *Patagona* species

Generations of ornithologists have pondered whether the giant hummingbirds comprise one

or more forms, and have attempted to characterize their distinguishing criteria. In his 1932

text, Hellmayr concluded that there were strong morphological differences "clearly"

indicating the existence of two forms. However, the analysis by Williamson et al. (2024), as

well as our present comparison, which included data from an additional 248 adults from

across the range, revealed substantial overlap in sex-specific mensural traits between

Northern and Southern Giant hummingbirds. Hellmayr's data did not show this overlap

between forms, even when controlling for sex (Fig. 3). Areta et al. (2024)'s assertion that

Hellmayr's results were "clearly mirrored" by Williamson et al. (2024) is an overstatement.

The *peruviana* measurements in Hellmayr's dataset were significantly greater than those in

our dataset for all comparisons except male bill length (Fig. 3). When compared with 10,000

resampled trait distributions, Hellmayr's results were highly improbable, suggesting that the

birds were identified, at least in part, based on their measurements and incorrect assumptions

about inter-taxon phenotypic variation. The differences between taxa in Hellmayr's data are

too large and the variability within taxa too small to have been produced by an unbiased

sampling of the two species. In this way, Hellmayr's measurements do not comprise valid

taxonomic evidence.

Why did Hellmayr's (1932) data appear so decisive in distinguishing two *Patagona*

taxa? One explanation is that Hellmayr included misidentified specimens in his dataset; we

know this occurred because at least several *P. gigas* specimens were included in his

"*peruviana*" series (e.g., AMNH-479593, AMNH-479594, and AMNH-479595). Including

misidentified individuals would diminish between-taxon differences and inflate variability

within taxa—the opposite of the pattern observed in Hellmayr's data. Alternatively, Hellmayr

may have used a combination of plumage assessment, measurements, and locality to assign

each specimen to a taxonomic category; thus, he may have started with the *a priori*

assumption that the two taxa were non-overlapping in size traits and range, and may then

have included or omitted data that fit with this hunch. Considering the magnitude of the

challenge inherent in identifying giant hummingbirds based on plumage, it may have been

Hellmayr's best option to make these educated guesses, seeking consilience of evidence as he

tried to fit a taxonomy to this difficult group. While this approach may have been reasonable

given the tools of the era, the resulting conclusions were therefore based on overly-generous

distinctions that consequently led to further misidentification. Hellmayr is not the only

worker to have studied the giant hummingbirds; however, his 1932 text is widely regarded as

definitive for giant hummingbird identification and it is perhaps the only known text to

provide raw data, permitting present-day re-analysis.

726   Robust analysis of morphology using linear discriminant analysis, principal

component analysis, and comparisons of means on our dataset spanning the entire *Patagona*

distribution has reinforced that individuals of each the Northern and Southern species may be

difficult to distinguish even with a suite of measurements used in concert (Figs. 3–4). It is

therefore unwise to characterize the two giant hummingbird species on the basis of any single

morphological trait, and it would be imprudent to use any single measurement to attempt to

identify individuals to species, as was done recently by Areta et al. (2024).

733   One underappreciated component of giant hummingbird identification is the extent of

individual plumage variation that exists *within* each species (Fig. 1). The authors' study

of >600 wild-caught individuals and museum specimens suggests that the extent of cinnamon

coloration appears to be related to age; even so, each species possesses a high degree of

plumage variation that appears unrelated to sex. Hellmayr (1932) and Zimmer (1952) each

noted this variation, suggesting that it might be random; Simon (1921) noted age-associations

with plumage color corroborated by our study. Even among adults, the range of variation in

the throat color and pattern of Southern Giant Hummingbirds overlaps that of Northern Giant

Hummingbirds. Although it is generally possible to distinguish the two species using

plumage characteristics, because within-species variation in throat color and pattern may

overlap, many individuals are best left unidentified when observed during locations and

seasons of co-occurrence. A clearer understanding of the seasonal elevational and latitudinal

ranges of both species will be essential to field identification and further understanding

migration, seasonal zones of overlap, and regions where the two species hybridize.

748   **Distribution and identification of the giant hummingbirds**

749 Here, we provide details about Northern and Southern Giant Hummingbird identification,

750 seasonal distributions, elevational ranges relevant to ornithologists and birders.

752   *Northern Giant Hummingbird (Patagona peruviana)*

Plumage: Adult Northern Giant Hummingbirds typically have whitish or buffy-colored

throats with dark blackish streaks (but note that throat color is variable, from very high to

more subtle contrast), a subtle cinnamon patch at the base of the throat (giving the throat base

a more rufous color), and a whitish eye-ring with post-ocular spot (Fig. 1). Wings and bill

average longer, but identification by measurements alone is often not possible.

Distribution and range: The Northern Giant Hummingbird is found year-round in extreme

southern Colombia, Ecuador, Peru and northern Chile (Arica and Parinacota Region); it is not

yet confirmed to occur in Bolivia, and there are no records (photographs, recordings, or

specimens) of this species in Argentina. It spans an elevational range of ~1,800-3,600 m in

Ecuador, ~1,900-4,300 in Peru, and ~1,900-3,800 m in Chile. In Peru, observers should

carefully scrutinize giant hummingbirds during periods of seasonal overlap (~May–Sept), as

the two *Patagona* species co-occur in many of the same sites and elevations. Individuals

observed throughout Peru from November–April are most likely Northern Giant

Hummingbirds; individuals observed in central to southern Peru from May–September could

plausibly be Northern or Southern Giant Hummingbirds. The Northern Giant Hummingbird

may appear very similar in plumage to the Southern Giant Hummingbird (Fig. 1), particularly

in poor lighting conditions or when seen in flight. This species is found in arid and semi-arid

habitats on the west slope of the Andes and in intermontane valleys, including montane scrub,

hedgerows, agricultural areas, open woods (including *Polylepis*), and gardens, often with

Agave, *Puya*, and Cactaceae spp. Across its range, it tends to associate with fast-flowing

streams.

Southern Giant Hummingbird (Patagona gigas)

Plumage: The Southern Giant Hummingbird tends to have a warm brown to cinnamon throat

with subtly contrasting grayish-green streaks (but note that throat contrast is variable and may

approach Northern in coloration), occasionally with a subtle cinnamon patch at the base;

some individuals may have extensive rufous throughout the throat and breast. The eye-ring

and post-ocular spot are subtle (buffy-colored, if present) to absent, giving Southern

individuals a 'blank-faced' appearance (Fig. 1). Wings and bill average shorter.

Distribution and range: The Southern Giant Hummingbird is found in central Chile, central

through northwest Argentina, Bolivia, and Peru (to central Ancash Department). This species

breeds from sea level to ~2,000 m in Chile, above ~2,500 m in Argentina, and ~2,100–4,150

m in Bolivia. After breeding, Southern Giant Hummingbirds in central Chile migrate

(beginning ~mid-February to March) to Peru, passing through the breeding grounds of

populations from northwest Argentina. At least some populations that breed in northwest

Argentina also migrate to central Peru, but details about migration are as yet unknown. In

Peru, the Southern Giant Hummingbird is found on the nonbreeding grounds, up to 4,300 m,

at the same sites and elevations as the Northern Giant Hummingbird. Southern Giant

hummingbirds are estimated to comprise ~20–40% of the nonbreeding population in Peru

during May–September (Williamson et al. 2024); however, further work is needed to refine

our understanding of this species migratory stopover sites and nonbreeding range. Individuals

observed in Bolivia should be carefully evaluated, particularly in the north, to rule out

presence of the Northern Giant Hummingbird. Individuals in Chile and Argentina may

overwinter. Observers should carefully scrutinize individuals in northern Chile (Arica and

Parinacota Region), particularly during ~August–October when Southern Giant

Hummingbirds migrate through the region (and may pass through the range of Northern

Giant Hummingbirds).

801   *Juvenile giant hummingbirds*

The plumage of juvenile giant hummingbirds (called juvenal plumage) of both species is

nearly identical (Fig. 1); previous workers have noted the difficulty of distinguishing

juveniles (e.g., Zimmer 1952, Ortiz-Crespo 1974). The juveniles of both species have deep

cinnamon throats, breasts, and bellies that fade to duller brown as birds molt into immature

plumage; emerald-toned heads, backs, and wings with buffy-tips to contour feathers; a

pronounced white rump patch that may extent mid-way up the back; and fresh white tips to

primaries and rectrices (Mulsant and Verreaux 1876; Fig. 1). Dark black streaking seems to

appear sooner on the throats of Northern Giant Hummingbird juveniles; presence of this

character may prove useful for identification. Season and locality should be carefully

evaluated when observing juveniles.

813   **Conclusions**

The migration-tracking, genomic, physiological, and morphological study conducted by

Williamson et al. (2024) demonstrated that elevational migrant and high-elevation resident

populations of the giant hummingbird comprise distinct species, now known as the Southern

and Northern Giant Hummingbird, respectively (Williamson et al. 2024). Before the

introduction of these novel data on migration and genetics, the taxonomy of giant

hummingbirds was impossible to disentangle, in part because of erroneous assumptions about

ranges and oversimplified identification criteria. The first described species in the genus,

Patagona gigas, and the taxon *peruviana,* have had confusing histories. The putative voucher

syntypes for *P. gigas* have unconfirmed identities, unknown collection dates, and dubious

provenance; lacking critical information, they cannot be confirmed as extant. The final

syntype, a plate depicting a juvenile putative male, does not possess any identifying features

of the Southern Giant Hummingbird and is thus unsuitable as a lectotype. To clarify the status

of the Southern Giant Hummingbird (*Patagona gigas*; *sensu* Williamson et al. 2024), we

designate a neotype for *P. gigas* by selecting a modern specimen of known provenance linked

to definitive nuclear and mitochondrial DNA sequence data, and for which annual migration

data exist (MSB:Bird:56142); this designation additionally corrects the previous erroneous

type locality restriction. The newly-designated *peruviana* lectotype appears to correspond to

the Northern Giant Hummingbird in plumage, and the name *peruviana* therefore has

nomenclatural precedence as a senior synonym. Thus, despite introduction of the name *chaski*

concomitantly with definitive delimitation of species limits, phenotypic characterizations,

physiological and behavioral differences, and year-round distributions for the two *Patagona*

species (Williamson et al., 2024), *chaski* is a junior synonym of *peruviana*. We recommend

DNA sequencing to confirm the identity of the *peruviana* lectotype.

The two cryptic giant hummingbird species are highly challenge to distinguish by

plumage and morphology alone, as confirmed by our analysis of 608 wild-caught individuals

and museum-vouchered natural history specimens spanning 1864–2023, including both

species across the Andean range. These findings contrast the longstanding idea that the two

forms were "easily" distinguished, additionally highlighting how inaccurate historical

diagnoses led to misidentification in the field and in collections. We clarify the historical

record by providing a list of specimens misidentified in museums. Finally, we offer detailed

species accounts with notes on plumage diagnosis, and seasonal latitudinal and elevational

ranges, to aid ornithologists and birders in further field study.

As ornithologist Ted Parker once noted, "A taxonomic puzzle illuminates everything

else about a bird. The thing to keep in mind—because some people here and elsewhere think

finding a new species is the ultimate thing—is how much we have learned about all of these

birds" (Stap 1991, pg. 160).

Acknowledgements

We are grateful to Frank Rheindt, Thomas Pape, Pamela Rasmussen, Oscar Johnson, Alex Figueroa, Glaucia Del-Rio, Shawn Billerman, Terry Chesser, and J. V. Remsen, Jr. for helpful discussion on taxonomy and nomenclature. Field assistance was provided by Kyana Montoya and Marlon Chagua, French language translation by Peter Truog, and distribution information by Leonardo Campagna. We are grateful to the museum curators, collections managers, staff, and students who facilitated this work: Mark Adams, Mike Andersen, Clare Atkinson, John Bates, Bentley Bird, Sharon Birks, Patrick Boussès, Serina Brady, Mariel Campbell, Eamon Corbett, Joel Cracraft, Charles Dardia, Carla Dove, Scott Edwards, Rob Faucett, Mary Margaret Ferraro, Jérôme Fuchs, John Gerwin, Gary Graves, Emily Griffith, Shannon Hackett, Andy Johnson, John Klicka, Andy Kratter, Terry Lott, Irby Lovette, Ben Marks, Nick Mason, Jon Merwin, Chris Milensky, Rob Moyle, Jose Antonio Otero, Nathan Rice, Mark Robbins, Sahid Robles-Bello, Vanya Rohwer, Sam Rutledge, Dave Steadman, Paul Sweet, Jeremiah Trimble, Tom Trombone, Hein van Grouw, Jason Weckstein, and J. V. Remsen, Jr. Collections access, data, and historical information were provided by the American Museum of Natural History, Academy of Natural Sciences of Drexel University, Carnegie Museum of Natural History, Centro de Ornitología y Biodiversidad, Cornell University Museum of Vertebrates, Field Museum of Natural History, Louisiana State University of Museum of Natural Science, Museum of Comparative Zoology at Harvard University, Muséum National d'Histoire Naturelle, Museum of Southwestern Biology, Natural History Museum at Tring, and the Smithsonian National Museum of Natural History. This manuscript is under review by the Working Group on Avian Nomenclature. Data from wild birds measured in Peru were collected under Servicio Nacional Forestal y de Fauna Silvestre RD-000056-2023-DGGSPFFS-DGSPFS. IACUC approval was granted under Cornell University protocol 2022-0127.

Author contributions

JLW and CCW conceived the idea. All authors collected the data. JLW curated the data and

obtained funding. JLW and CRG analyzed the data. JLW, CRG, and BWR made figures and

tables. JLW, TV, and CCW contributed critical resources. JLW wrote the initial draft. JLW

and CCW substantially revised the manuscript with input from all authors.

Conflict of interest

The authors declare no conflicts of interest.

Funding

This work was supported by an NSF Postdoctoral Research Fellowship in Biology (DBI-

2208924) and a Cornell Lab of Ornithology Edward W. Rose Postdoctoral Fellowship to

JLW. Collections visits were supported by an American Philosophical Society Franklin

Research Grant.

Data availability statement

Data from Williamson et al. (2024) analyzed here are available on Dryad:

https://datadryad.org/stash/dataset/doi:10.5061/dryad.44j0zpcnp. Unpublished data will be

uploaded to a repository upon manuscript acceptance.

References

References

Aguirre, J. (2001). Nidificacion patagonas gigas gigas en una quebrada costera de Chile central.
Boletín Chileno de Ornitología 8:10–12.

Ambiente, C. N. del M. (2008). Biodiversidad de Chile, patrimonio y desafios. Ocho Libros
Editores, Santiago, Chile.

Areta, J. I., M. R. Halley, G. M. Kirwan, H. V. Norambuena, N. K. Krabbe, and V. Q. Piacentini
(2024). The world's largest hummingbird was described 131 years ago. *Bulletin of the*
British Ornithologists' Club 144.

Barros, R. (1952). Apuntes sobre la historial natural de la Pinguera, Patagona gigas. *Anales de la*
Academia Chilena de Ciencias Naturales 17:135–144.

Boucard, A. (1893). Genera of Humming Birds. Pardy & Co. Printers, London, UK.

Elliot, D. G. (1879). A classification and synopsis of the Trochilidae. Smithsonian Institution,
Washington, D.C.

Estades, C. F., M. A. Vukasovic, J. A. Tomasevic, and N. Desert (2008). Giant Hummingbirds
(Patagona gigas) Ingest Calcium-rich Minerals. *The Wilson Journal of Ornithology*
120:651–653.

Fjeldsa, J., and C. E. Barbosa (1983). A Giant Hummingbird from Paramo de Chingaza,
Colombia. *The Wilson Bulletin* 95:661–662.

Fjeldså, J., and N. Krabbe (1990). Birds of the High Andes. Zoological Museum, University of
Copenhagen, Copenhagen, Denmark.

González-Gómez, P. L., and C. E. Valdivia (2005). Direct and Indirect Effects of Nectar Robbing
on the Pollinating Behavior of Patagona gigas (Trochilidae). *Biotropica* 37:693–696.

Goodall, J. D., A. W. Johnson, and R. A. P. Bañados (1946). Las Aves de Chile, su conocimiento
y sus custumbres.

Gould, J. (1861). An introduction to the trochilidae, or family of humming-birds. London.

Gray (1840). A List of the Genera of Birds, with an indicator of the Typical Species of each
Genus. London.

Hartert, E. (1900). Trochilidae. Berlin, R. Friedländer und sohn, 1900.

Hellmayr, C. H. (1932). The Birds of Chile. Field Museum of Natural History, Chicago, IL,
USA.

Howell, N. G., and F. Schmitt (2018). Birds of Chile: A Photo Guide. Princeton University Press,
Princeton, NJ, USA.

ICZN (1999). International Code of Zoological Nomenclature. International Commission on
Zoological Nomenclature, London, UK. 306 pp.

ICZN (2003). Declaration 44. Amendment of Article 74.7.3. *Bulletin of Zoological*
Nomenclature 60:263.

Ivy, C. M., and J. L. Williamson (2024). On the Physiology of High-altitude Flight and
Altitudinal Migration in Birds. *Integrative And Comparative Biology*:icae062.

Jiménez-Mejías, P., S. Manzano, V. Gowda, F.-T. Krell, M.-Y. Lin, S. Martín-Bravo, L. Martín-
Torrijos, G. Nieto Feliner, S. L. Mosyakin, R. F. C. Naczi, C. Acedo, et al. (2024).
Protecting stable biological nomenclatural systems enables universal communication: A
collective international appeal. *BioScience*:biae043.

Johow, F. (1910). Estudios de biología vejetal, I. Sobre algunos casos de ornitolofilía en la flora
Chilena. *Anales de la Universidad de Chile* 126:27–50.

Johow, F. (1921). Las cactacea de los alrededores de Zapallar. *Revista Chilena de Historia*
Natural 15:152–166.

Jouanin, C. (1950). Catalogue Systématique des types de trocitilidés du Muséum National
D'Histoire Naturelle de Paris. *Bulletin Du Muséum National D'Histoire Naturelle*.
Muséum National D'Histoire Naturelle, Paris, France.

Koepcke, M. (1983). The birds of the Department of Lima, Peru. Harrowood Books, Newtown
Square, Pennsylvania, USA.

Lanyon, W. E. (1995). Ornithology at the American Museum of Natural History. In
Contributions to the History of North American Ornithology (Jr. and J. A. J. William E.
Davis, Editor). Nuttall Ornithological Club, Cambridge, MA, pp. 113–144.

Lesson, R. P. (1829). Histoire naturelle des oiseaux-mouches. Arthus Bertrand, Paris, France.

Medel, R., M. López-Aliste, and F. E. Fontúrbel (2022). Hummingbird-plant interactions in
Chile: An ecological review of the available evidence. *Avian Research* 13.

Medrano, F., R. Barros, H. V. Norambuena, M. Ricardo, and S. Fabrice (2018). Atlas de las Aves
Nidificantes de Chile 2011 – 2016.

Mulsant, E., and E. Verreaux (1876). Histoire naturelle des oiseaux-mouches. Paris.

Ortiz-Crespo, F. I. (1974). The Giant Hummingbird Patagona gigas in Ecuador. *Ibis* 116:347–
359.

Ortiz-Crespo, F. I. (1986). Consideraciones sobre las migraciones de dos picaflores
neotropicales. *El Hornero* 12:298–300.

Pearman, M., and J. I. Areta (2020). Birds of Argentina. Princeton University Press, Princeton,
960   New Jersey, USA.

961   Peters, J. L. (1945). Check-list of Birds of the World, Volume V. Oxford University Press,
962   London.

963   Porter, C. E. (1897). Revista Chilena de Historia Natural.

964   Renshaw, G. (1905). The Reunion Starling. *The Zoologist* 4:418–422.

965   Revilliod, P., J. Carl, and R. de Lessert (1942). Revue Suisse de Zoologie.

966   Rodríguez-Flores, C. I., J. F. Ornelas, S. Wethington, and M. del C. Arizmendi (2019). Are
967   hummingbirds generalists or specialists: Using network analysis to explore the
968   mechanisms influencing their interaction with nectar resources. *PLoS ONE*:1–32.

969   Rookmaaker, L. C., P. A. Morris, I. E. Glenn, and P. J. Mundy (2006). The ornithological cabinet
970   of Jean-Baptiste Bécoeur and the secret of the arsenical soap. *Archives of Natural History*
971   33:146–158.

972   Roselaar, C. S. (2003). An inventory of major European bird collections. *Bulletin of the British
973   Ornithological Club*.

974   Rueda-Uribe, C., L. Herrera-Alsina, L. T. Lancaster, I. Capellini, K. K. S. Layton, and J. M. J.
975   Travis (2023). Citizen science data reveal altitudinal movement and seasonal ecosystem
976   use by hummingbirds in the Andes Mountains. *Ecography*:e06735.

977   Ruschí, A. (1961). Algumas observacoes sobre Patagona gigas peruviana Boucard. *Boletim Do
978   Museu De Biologia* 23.

979   Sánchez Osés, C. (2003). Taxonomy, phylogeny, and biogeography of the Andean hummingbird
980   genera Coeligena LESSON, 1832; Pterophanes GOULD, 1849; Ensifera LESSON 1843;
981   and Patagona GRAY, 1840 (Aves: Trochiliformes).

982   Simon, E. (1921). Histoire naturelle des Trochilidae (Synopsis et Catalogue). *Encyclopédie-
983   Roret*:i–vi, 1–416.

984   Speziale, K. L., and S. Lambertucci (2009). El picafor gigante (Patagona gigas) en la Provincia
985   de Río Negro. *Nuestras Aves* 54:39–40.

986   Stap, D. (1991). A Parrot Without a Name. First University of Texas Press, Austin, TX.

987   Velásquez-Noriega, P., V. Serrudo-Gonzáles, A. C. Paca-Condori, and D. M. Larrea-Alcazar
988   (2021). El picaflor gigante de los Andes. *Bolivia Ecológica* 97.

989   Vieillot, L. P. (1824). La Galerie des Oiseaux. Constant-Chantpie, Paris, France.

990   Warren, R. L. M. (1966). Type-specimens of birds in the British Museum (Natural History). Eyre
991   and Spottiswoode Limited, Portsmouth.

Williamson, J., E. F. Gyllenhaal, S. M. Bauernfeind, E. Bautista, M. J. Baumann, C. R. Gadek, P.
P. Marra, N. Ricote, T. Valqui, F. Bozinovic, N. D. Singh, and C. C. Witt (2024). Extreme
elevational migration spurred cryptic speciation in giant hummingbirds. *Proceedings of*
the National Academy of Sciences 121:1–11.

Williamson, J. L., and C. C. Witt (2021a). Elevational niche-shift migration: Why the degree of
elevational change matters for the ecology, evolution, and physiology of migratory birds.
Ornithology 138:1–26.

Williamson, J. L., and C. C. Witt (2021b). A lightweight backpack harness for tracking
hummingbirds. *Journal of Avian Biology*:1–9.

Woods, S., F. Ortiz-Crespo, and P. M. Ramsay (1998). Presence of giant hummingbird Patagona
gigas and Ecuadorian hillstar Oreotrochilus chimborazo jamesoni at the Ecuador -
Colombia border. *Cotinga* 10:37–40.

Zimmer, J. T. (1930). Birds of the Marshall field Peruvian expedition 1922-1923. Field Museum
of Natural History, Chigago, USA.

Zimmer, J. T. (1952). Studies of Peruvian Birds. No. 62 The hummingbird genera Patagona,
Sappho, Polyonymus, Ramphomicron, Metallura, Chalcostigma, Taphrolesbia, and
Aglaiocercus. *American Museum of Novitates* 1595:1–30.

1009

Northern

ADULT

throat variable, from very high to more subtle contrast

bill averages longer

noticeable eye-ring and post-ocular spot

can have darker rufous background color

throat contrasting with dark **blackish** streaks, most often set against whitish to light buffy background

wings average longer

JUVENILE

juvenal plumage similar, possibly indistinguishable between species

Southern

ADULT

throat variable, can be high contrast approaching Northern

bill averages shorter

subtle to absent eye-ring and post-ocular spot

can have extensive rufous throughout throat and breast

throat less contrasting with subtle dark **grayish-green** streaks, most often set against buffy background

wings average shorter

Within-species variation in traits important for identification, such as throat color and the boldness and extent of the eye-ring and post-ocular spot, may overlap. Although it is possible to distinguish between the two species using plumage characteristics, many individuals are best left unidentified when observed in locations and seasons where both co-occur.

1010

1011   **Figure 1.** Key for distinguishing Northern and Southern Giant Hummingbirds, highlighting

1012   important plumage characteristics described in Williamson et al. (2024), including throat color

1013   and pattern and extent of eye-ring and post-ocular spot, as well as morphological characteristics,

1014   such as wing length and bill length. Juvenile is represented for comparison (the term "juvenal" is

1015   a specific term corresponding to juvenile plumage). Illustrations: Bryce W. Robinson.

Figure 2. The *Patagona gigas* neotype, MSB:Bird:56142, housed at the Museum of

Southwestern Biology at the University of New Mexico. The specimen, an adult female, was

collected in Algarrobo, Valparaíso Región, Chile (13°20.916' S, 71°38.294' W) on 9 December

2018. Female. This specimen was unequivocally identified as a Southern Giant Hummingbird

based on definitive nuclear and mitochondrial genomic, physiological, morphological, and

migration tracking data (Williamson et al. 2024). **A** Ventral view. **B** Dorsal view. Specimen

record and details are available from https://arctos.database.museum/guid/MSB:Bird:56142.

Photos: [AUTHOR].

Figure 3. Comparison of morphological data (bill length, wing chord, and tail length; all in mm) between Hellmayr (1932) and Williamson et al. (present study). **A–C** Males. **D–F** Females. Hellmayr compared adults of the two suspected forms, *gigas* and *peruviana* (*n*=45 specimens total). Williamson et al. compared adults of the two species, the Southern and Northern Giant

1087   Hummingbird (*n*=448 individuals total). Sample sizes for each comparison are reported on the x-

1088   axis and significant *p*-values are indicated above each comparison (from t-tests or non-

1089   parametric Wilcoxon Rank Sum tests). Box plot horizontal lines represent median values. Each

1090   point is a value from a single individual.

1091

Figure 4. Principal Component Analysis (PCA) of bill, wing, and tail measurements for *gigas*

and *peruviana* individuals from Hellmayr (1932) and Southern and Northern Giant

Hummingbirds from Williamson et al. (present study) with MNHN *peruviana* specimens and the

boliviana holotype. Hellmayr's data were entered numerically, rather than by individual; we have

therefore included only records with single individuals per site. **A** Males (*n*=172 individuals); **B**

Females (*n*=174 individuals). The male *peruviana* lectotype (MNHN-ZO-MO-1989-340), female

peruviana paralectotype (MNHN-ZO-MO-1989-341), and *boliviana* holotype (MNHN-ZO-MO-

1989-342) are plotted for reference; the *boliviana* holotype is sex unknown and is thus included

in both PCAs. The female *peruviana* specimen is plotted despite the tip of its maxilla being

broken (Patrick Bousses, *pers. comm.*); however, *Patagona* maxilla and mandible lengths are

typically equal. For reference, bill measurements: *peruviana* male: 35.7 mm; *peruviana* female

35.8 mm (mandible only); *boliviana*: 35.8 mm.

1104

Figure 5. Visual representation of the *Patagona* specimen record as it pertains to previous versus current identifications (see Table S5 for full list). Of the 603 total adult and juvenile specimens with locality data whose catalog records and tags we analyzed from eleven major museum collections, a total of 269 individuals (~44%) are catalogued with subspecific identifiers. Of these, 29 specimens are labeled as *P. gigas* and pertain to the Southern Giant Hummingbird (*Patagona gigas*, sensu Williamson et al. 2024). A total of 226 specimens with known lineage are labeled as *P. g. peruviana*; of these, 152 specimens are correctly identified as Northern Giant Hummingbirds (*P. peruviana*) and 74 specimens are incorrectly identified as *P. g. peruviana* and are Southern Giant Hummingbirds (*P. gigas*).

1114 **SUPPLEMENT**

1115 **Taxonomy, nomenclature, and identification of the giant hummingbirds (*Patagona* spp.)**

1116 **(Aves: Trochilidae)**

1117

The Whitely series of *Patagona* specimens

Henry H. Whitely, Jr., a British naturalist and explorer particularly interested in ornithology

(Sclater 1893, Gladstone 1927), was a key figure in the history of *Patagona* taxonomy. For

many years, Whitely collected specimens in southern Peru, with numerous trips throughout

Arequipa and Cusco between the years of 1867 and 1873 (Sclater and Salvin 1867, 1869;

Sclater 1893, Chapman 1921, Stephens and Traylor 1983). During this time, Whitely was

credited with a number of new species discoveries, including the Bearded Mountaineer

(*Oreonympha nobilis*; Sclater 1893). Many of Whitely's specimens were dispersed through

the specimen trading business of his father, Henry Whitely, Sr. (Sharpe 1906, Rasmussen and

Prys-Jones 2003). Although Whitely Sr., and to a lesser extent, Whitely Jr.'s, reputations have

been questioned (Whitely, Sr. has been called 'reliably unreliable'; Rasmussen and Prys-

Jones 2003), none of the specific problems that have been found appear to concern the

localities of Whitely's *Patagona* specimens.

Whitely collected at least 16 *Patagona* specimens in Peru from 1867 to 1873 (Table

S1). In one letter about his 1867 Arequipa trip, Whitely noted, "… now commenced the worst

part of the journey…the way, however, was enlivened by the sight of numerous birds, and

especially, for some 2,000 feet, by the movements of the Giant Humming-bird (*Patagona*

gigas)" (Sclater and Salvin 1867). Whitely's *Patagona* specimens, particularly those from

Tinta, Cusco, Peru, have played a crucial role in history because they are thought to have

formed the basis for a subsequent type description by Boucard (Boucard 1893). Whitely

collected in Tinta on several trips, including May–June 1868 and January 1869 (Sclater and

Salvin 1869). The identities of at least three *Patagona* specimens from his 1868 trip are

known (MNHN-ZO-MO-1989-340, FMNH-45797, and FMNH-45798), and three others are

suspected (AMNH-37500, AMNH-37501, and MNHN-ZO-MO-1989-34; Table S1). One

specimen (AMNH-37501), housed at the AMNH was designated as the "Type of Fig. 33, p.

67" in Giraud Elliot's 1879 synthesis, "*A classification and synopsis of the Trochilidae*"

(Elliot 1879). This specimen, a male from Tinta, was noted in original ledgers as "figured in

synopsis", as was AMNH-37502, a female collected from Cachupata, Cusco, Peru on August

21, 1873; both specimens were denoted with equivalent status (Table 1). These specimens,

along with two other Whitely specimens, one each from Cachupata and Tinta (AMNH-37499

and AMNH-37500), were originally part of the Tring Collection of birds but were obtained

by the AMNH in 1932 when Lord Rothschild sold his collection (Lanyon 1995; Table S1).

Based on available data from gazetteers and related publications, AMNH-37500 and AMNH-

37501 are suspected to have been collected during Whitely's June–August 1868 trip to Tinta;

a point of clarification from Williamson et al. (2024). There are at least three other Whitely

specimens (ANSP-50447, NHMUK 1887.3.22.700, and NHMUK 1913.3.20.291) collected

during January 1869, and one additional specimen (ANSP-50448) is suspected to have been

collected in either May–June 1868 or January 1869 (Table S1). At present, Whitely's

Patagona specimens are known to reside at the AMNH, MNHN, ANSP, FMNH, and

NHMUK; it is possible that there are additional Whitely *Patagona* specimens in other

collections, as well.

A MNHN-ZO-MO-1989-340

B MNHN-ZO-MO-1989-341

C MNHN-ZO-MO-1989-342

Figure S1. A Lectotype for the taxon *peruviana*, MNHN-ZO-MO-1989-340, a male, collected June 15, 1868. **B** Paralectotype for the taxon *peruviana*, MNHN-ZO-MO-1989-341, a female, no date. Both specimens are from Boucard's (1893) collection and were collected by Henry H. Whitely from Tinta, Peru. **C** Holotype for the taxon *boliviana* from Boucard's (1893) collection, MNHN-ZO-MO-1989-342, undetermined sex, collected in 1875 by Clarence Buckley from Bolivia. Photos are unedited from originals. Photos: Patrick Bousses, Muséum National d'Histoire Naturelle (MNHN), Paris.

L'Oiseau-mouche géant Trochilus gigas.

P. Oudart d'après le dessin de M. Prête.   Litho de C. Motte.

1196   **Figure S2.** Vieillot's putative "young male" syntype, illustrated by Paul Louis Oudart in Plate

1197   180 of Vieillot's *La Galerie des Oiseaux* (Vieillot 1824). The individual depicted was

1198   attributed to M. (Mr.; for "monsieur" in French) Portier, "attaché" of the Ministry of the

1199   Navy. The specimen has not been located and its current whereabouts are unknown to us.

1200

Figure S3. The putative "adult female" syntype of Vieillot (1824) for the species *Patagona*

gigas, formerly known as the Giant Hummingbird and currently known as the Southern Giant

Hummingbird (*sensu* Williamson et al. 2024). The specimen shown is NHMUK-

1933.11.14.36, a female with no collection date and no specific locality (provenance

unknown), first acquired by Jean-Baptiste Bécouer before his death in 1777. This specimen, a

taxidermied mount, is thought to be one of two syntypes used to describe *Patagona gigas* by

Vieillot in 1824; but see specimen tag notes. **A** Ventral view showing the bird's throat, which

is a strong match for the Southern Giant Hummingbird; **B–C** Non-original specimen labels

(the specimen does not possess original labels) indicating uncertainty about whether

NHMUK-1933.11.14.36 was one of Vieillot's (1824) syntypes; **D** Dorsal view of the

specimen. Photos: Mark Adams, NHMUK.

Figure S4. Specimen MNHN-ZO-2023-396 (sex and locality unknown; collection date

unknown but prior to 1842) from the MNHN in Paris, believed by Boucard (1893) to be

"Vieillot's type". Boucard wrote, "I have in my collection what I consider as the type of

Vieillot 'Ex Coll Riocour'" (Boucard 1893). MNHN-ZO-2023-396 is not listed as a syntype

by MNHN. **A** Ventral view of specimen. **B** Side view of specimen; photos are unedited from

originals. **C** Front side of the Boucard Collection tag showing the question mark ("?") after

the words "type Vieillot" (Patrick Bousses, *pers. comm.*). **D** Back of Boucard Collection tag.

E Front of Comte de Riocour Collection tag. Photos: MNHN.

Figure S5. Expected distributions of trait mean differences between Northern and Southern

Giant Hummingbirds, based on 10,000 resampled datasets generated from Williamson et al.

(present study). **A–C** Expected male trait differences for each bill, wing, and tail; **D–F** Expected

female trait differences for each bill, wing, and tail. Red dashed vertical lines indicate the

differences between *gigas* and *peruviana* trait means from Hellmayr (1932) data, with the

estimated probabilities of obtaining Hellmayr's values (red text). Estimated probabilities were

calculated from 10,000 simulated datasets.

Figure S5. Expected distributions of coefficients of variation (CVs) for each Northern and

Southern Giant Hummingbirds, based on 10,000 resampled datasets generated from Williamson

et al. (present study). CVs were estimated separately for each species (Northern and Southern

Giant Hummingbird), each sex (males and females), and each trait (bill, wing, and tail). **A–C**

Northern males; **D–F** Northern females; **G–I** Southern males; **J–L** Southern females. Red dashed

vertical lines indicate the CVs from Hellmayr's (1932) original data, with the estimated

probabilities of obtaining Hellmayr's values (red text). Estimated probabilities were calculated

from 10,000 simulated datasets.

Specimen identifier	Museum	Country	Department	Specific Locality	Collection Date	Putative Identification	Evidence
AMNH-37500	AMNH	Peru	Cusco	Tinta	no date	Northern	Genomic data; plumage (suggestive)
AMNH-37501*	AMNH	Peru	Cusco	Tinta	no date	Northern	Genomic data; plumage (suggestive)
AMNH-37499	AMNH	Peru	Cusco	Cachupata	21 Aug 1873	Northern	Genomic data; plumage (suggestive)
AMNH-37502**	AMNH	Peru	Cusco	Cachupata	21 Aug 1873	Undetermined	Genomic data (Southern); plumage (Northern)
MNHN-ZO-MO-1989-340‡	MNHN	Peru	Cusco	Tinta	15 June 1868	Northern	Plumage (diagnostic)
MNHN-ZO-MO-1989-341‡^	MNHN	Peru	Cusco	Tinta	no date	Northern	Plumage (suggestive)
FMNH-45797	FMNH	Peru	Cusco	Tinta	25 May 1868	Northern	Plumage (diagnostic)
FMNH-45798	FMNH	Peru	Cusco	Tinta	20 June 1868	Northern	Plumage (diagnostic)

ANSP-50447	ANSP	Peru	Cusco	Tinta	4 Jan 1869	Northern	Plumage (diagnostic)
NHMUK 1887.3.22.700	NHMUK	Peru	Cusco	Tinta	4 Jan 1869	Northern	Plumage (diagnostic)
NHMUK 1913.3.20.291	NHMUK	Peru	Cusco	Tinta	4 Jan 1869	Northern	Plumage (diagnostic)
NHMUK 1913.3.20.293	NHMUK	Peru	unknown	W. Peru	no date	Northern	Plumage (diagnostic)
NHMUK 1887.3.22.701	NHMUK	Peru	Arequipa	Chiguata	10 June 1867	Unknown (juvenile)	–
NHMUK unregistered	NHMUK	Peru	Arequipa	Chiguata	10 June 1867	Unknown (juvenile)	–
NHMUK 1913.3.20.294	NHMUK	Peru	unknown	W. Peru	no date	Unknown (juvenile)	–

1249

1250 **Table S2.** Morphological summary statistics for *gigas* (Hellmayr 1932) and Southern Giant Hummingbird (Williamson et al., present

1251 study) males and females. For each measured parameter, we provide sample size (*n*), mean, standard deviation, range (minimum and

1252 maximum values), and coefficient of variation.

Dataset	Sex	Parameter (units)	*n*	Mean	Standard Deviation	Minimum	Maximum	Coefficient of Variation
Hellmayr 1932		Bill	8	34	0.76	33	35	0.02
		Wing	8	125.5	3.34	120	129	0.03
	Male	Tail	8	80.5	1.41	79	83	0.02
Williamson et al. (present study)		Bill	72	36.45	2.33	32.41	42.25	0.06
		Wing	69	127.81	6.71	111	141	0.05
		Tail	70	82.16	4.83	71	92	0.06
Hellmayr 1932		Bill	4	34.25	0.5	34	35	0.01
		Wing	4	118.75	1.5	118	121	0.01
	Female	Tail	4	75	3.37	70	77	0.04
Williamson et al. (present study)		Bill	72	36.56	2.48	30.91	41.74	0.07
		Wing	70	122.66	6.83	112	139	0.06
		Tail	71	79.06	4.11	71	90	0.05

1253

1254 **Table S3.** Morphological summary statistics for *peruviana* (Hellmayr 1932) and Northern Giant Hummingbird (Williamson et al.,

1255 present study) males and females. For each measured parameter, we provide sample size (*n*), mean, standard deviation, range

1256 (minimum and maximum values), and coefficient of variation.

Dataset	Sex	Parameter (units)	*n*	Mean	Standard Deviation	Minimum	Maximum	Coefficient of Variation
Hellmayr 1932		Bill	25	38.46	1.49	36	42	0.04
		Wing	25	136.36	2.02	133	140	0.01
	Male	Tail	25	85.88	2.33	83	90	0.03
Williamson et al. (present study)		Bill	112	38.53	1.71	34.76	42.03	0.04
		Wing	98	132.7	4.68	121	147	0.04
		Tail	109	81.31	4.23	66	91	0.05
Hellmayr 1932		Bill	8	39.31	1.03	38	41	0.03
		Wing	8	130.88	3.31	126	137	0.03
	Female	Tail	8	84	3.25	78	87	0.04
Williamson et al. (present study)		Bill	108	38.26	1.89	31.93	42.76	0.05
		Wing	102	125.75	5.32	109	141	0.04
		Tail	108	79.57	3.98	69	91	0.05

1257

1258   **Table S4.** Comparison of differences in taxon-mean trait values between Hellmayr (1932) and

1259   Williamson et al. (present study). Hellmayr's data compare differences between the taxa

1260   *peruviana* and *gigas*. Williamson et al. data compare differences between the species Northern

1261   Giant Hummingbird and Southern Giant Hummingbird.

		Difference between taxon-mean trait values (mm)		
Sex	**Source**	**Bill length**	**Wing chord**	**Tail length**
Male	Hellmayr 1932	4.46	10.86	5.38
	Williamson et al. (present study)	2.08	4.89	-0.85
Female	Hellmayr 1932	5.06	12.12	9.0
	Williamson et al. (present study)	1.70	3.09	0.51

1262

Table S5. List of museum specimens from Peru, Bolivia, Argentina, and Chile that are labeled *P.*

g. peruviana (or "*peruviana*" on tags) but that are actually Southern Giant Hummingbirds

(*Patagona gigas*). We analyzed the catalog records, tags, and skins of 603 adult and juvenile

specimens with locality data from eleven major museum collections. A total of 269 individuals

(~44%) are catalogued with subspecific identifiers; of these, 74 specimens (32.7%) are

incorrectly identified as *peruviana* (see Fig. 5). This list is intended to facilitate taxonomic

revision within the genus *Patagona* and aid museums in correcting the historical record.

Specimen identifier	Museum	Country	Department	Specific Locality	Collection Date	Sex
AMNH-139179	AMNH	Bolivia	Sucre	Incambo	20 Nov 1915	male
AMNH-139180	AMNH	Bolivia	Cochabamba	Totora	2 Oct 1915	male
AMNH-140718	AMNH	Argentina	Jujuy	Tilcara	10 Feb 1916	female
AMNH-140719	AMNH	Argentina	Jujuy	Tilcara	9 Feb 1916	female
AMNH-140720	AMNH	Argentina	Jujuy	Tilcara	10 Feb 1916	female
AMNH-145020	AMNH	Peru	Cusco	Huaracondo Canon	23 July 1916	female
AMNH-148279	AMNH	Bolivia	Cochabamba	Arque	5 Feb 1915	female
AMNH-37502	AMNH	Peru	Cusco	Cachupata	21 Aug 1873	female
AMNH-479593	AMNH	Argentina	Tucumán	Lara	12 Feb 1903	male
AMNH-479594	AMNH	Argentina	Tucumán	Lara	12 Feb 1903	female
AMNH-479595	AMNH	Argentina	Jujuy	Tilcara	24 Nov 1905	male
ANSP-120834	ANSP	Bolivia	La Paz	Calacoto	25 Jan 1935	male
ANSP-120835	ANSP	Bolivia	La Paz	Calacoto	24 Jan 1935	female
ANSP-120836	ANSP	Bolivia	La Paz	Calacoto	26 Jan 1935	male
ANSP-145414	ANSP	Bolivia	Cochabamba	Tiraque	27 Sept 1937	male
ANSP-145415	ANSP	Bolivia	Potosi	Oploca	27 Feb 1938	female
ANSP-145416	ANSP	Bolivia	Potosi	Oploca	28 Feb 1938	female
ANSP-145418	ANSP	Bolivia	Potosi	Oploca	28 Feb 1938	female

ANSP-145419	ANSP	Bolivia	Potosi	Oploca	28 Feb 1938	female
ANSP-145420	ANSP	Bolivia	Potosi	Oploca	2 Mar 1938	male
ANSP-145421	ANSP	Bolivia	Potosi	Oploca	2 Mar 1938	male
ANSP-145422	ANSP	Bolivia	Potosi	Oploca	28 Feb 1938	male
ANSP-145423	ANSP	Bolivia	Cochabamba	Tutimayu	4 Mar 1937	female
ANSP-145424	ANSP	Bolivia	Cochabamba	Tutimayu	3 Mar 1937	female
ANSP-145425	ANSP	Bolivia	La Paz	Viloco	1 Apr 1938	female
ANSP-145426	ANSP	Bolivia	La Paz	Viloco	31 Mar 1938	female
ANSP-167558	ANSP	Argentina	Catamarca	Andaluzala	20 Sept 1917	male
CM-120263	Carnegie	Bolivia	Cochabamba	Arani	12 Feb 1927	female
CM-120284	Carnegie	Bolivia	Cochabamba	Tiraque	26 Mar 1927	male
CM-52921	Carnegie	Argentina	Jujuy	Maimara	17 Nov 1911	male
CM-53156	Carnegie	Argentina	Jujuy	Maimara	16 Jan 1911	male
CM-81114	Carnegie	Bolivia	Cochabamba	Arani	31 Mar 1920	male
CM-81120	Carnegie	Bolivia	Cochabamba	Vacas	2 Apr 1920	male
CM-81121	Carnegie	Bolivia	Cochabamba	Vacas	2 Apr 1920	male
CM-81180	Carnegie	Bolivia	Cochabamba	Cochabamba	14 Apr 1920	female
CM-81182	Carnegie	Bolivia	Cochabamba	Cochabamba	14 Apr 1920	female
CM-81219	Carnegie	Bolivia	Cochabamba	Mollemolle	19 Apr 1920	male
CM-81271	Carnegie	Bolivia	Cochabamba	Mollemolle	23 Apr 1920	female
CM-81273	Carnegie	Bolivia	Cochabamba	Mollemolle	23 Apr 1920	female
CM-81275	Carnegie	Bolivia	Cochabamba	Mollemolle	23 Apr 1920	male
CM-81581	Carnegie	Bolivia	Cochabamba	Cochabamba	2 Oct 1920	male
CM-81585	Carnegie	Bolivia	Cochabamba	Cochabamba	3 Oct 1920	female
CM-94400	Carnegie	Bolivia	La Paz	Guaqui	26 Jan 1922	male
CUMV-52570	CUMV	Argentina	Jujuy	Volcan	5 Nov 2005	female
FMNH-179394	FMNH	Bolivia	Cochabamba	Vacas	15 Mar 1939	male
FMNH-179395	FMNH	Bolivia	Cochabamba	Vacas	16 Feb 1939	male

FMNH-179396	FMNH	Bolivia	Cochabamba	Vacas	16 Feb 1939	male
FMNH-179397	FMNH	Bolivia	Cochabamba	Arani	15 Dec 1940	male
FMNH-179398	FMNH	Bolivia	Cochabamba	Arani	15 Dec 1940	male
FMNH-179399	FMNH	Bolivia	Cochabamba	Arani	6 Dec 1940	female
FMNH-179400	FMNH	Bolivia	Cochabamba	Arani	6 Dec 1940	male
FMNH-179401	FMNH	Bolivia	Cochabamba	Arani	6 Dec 1940	male
FMNH-179402	FMNH	Bolivia	Cochabamba	Colomi	28 Jan 1938	female
FMNH-179403	FMNH	Bolivia	Cochabamba	Colomi	28 Jan 1938	female
FMNH-179404	FMNH	Bolivia	Cochabamba	Colomi	28 Jan 1938	male
FMNH-179405	FMNH	Bolivia	Cochabamba	Colomi	28 Jan 1938	female
FMNH-179406	FMNH	Bolivia	Cochabamba	Cochabamba	15 July 1939	female
FMNH-179407	FMNH	Bolivia	Cochabamba	Cochabamba	15 July 1939	male
FMNH-179408	FMNH	Bolivia	Cochabamba	Tiraque	5 Aug 1928	male
FMNH-293680	FMNH	Bolivia	Chuquisaca	Camargo	4 Feb 1973	female
FMNH-293681	FMNH	Bolivia	Chuquisaca	Camargo	5 Feb 1973	female
FMNH-293683	FMNH	Bolivia	Tarija	Tojo	25 Jan 1973	female
FMNH-57059	FMNH	Argentina	Tucumán	Colalao del Valle	14 Aug 1916	female
FMNH-61681	FMNH	Chile	Tarapacá	Putre	3 July 1924	male
FMNH-67517	FMNH	Peru	Lima	Matucana	3 May 1922	male
FMNH-67518	FMNH	Peru	Lima	Matucana	3 May 1922	female
LSUMZ-35863	LSUMZ	Bolivia	Cochabamba	Orcoma	21 July 1960	female
LSUMZ-37404	LSUMZ	Bolivia	Cochabamba	Cochabamba	10 Apr 1958	female
MCZ-262196	MCZ	Argentina	NA	Cumbres de Luracatao	NA Jan 1962	female
MCZ-287952	MCZ	Chile	NA	Chusmisa	NA Nov 1943	female
MCZ-95880	MCZ	Bolivia	La Paz	La Paz	NA Sept 1895	female
USNM-172374	USNM	Bolivia	La Paz	La Paz	10 Feb 1893	male
USNM-529148	USNM	Peru	Arequipa	Volcan Misti	8 July 1938	male
USNM-529149	USNM	Peru	Arequipa	Volcan Misti	8 July 1938	male

1270

References

1271

Boucard, A. (1893). Genera of Humming Birds. Pardy & Co. Printers, London, UK.

Chapman, F. M. (1921). The distribution of the bird life in the Urubamba Valley of Peru.
Washington Government Printing Office, Washington, D.C.

Elliot, D. G. (1879). A classification and synopsis of the Trochilidae. Smithsonian Institution,
Washington, D.C.

Gladstone, H. S. (1927). Henry Whitely, 1817-1898. British Birds XXII:290–293.

Gould, J. (1861). An introduction to the trochilidae, or family of humming-birds. London.

Hartert, E. (1900). Trochilidae. Berlin, R. Friedländer und sohn, 1900.

Lanyon, W. E. (1995). Ornithology at the American Museum of Natural History. In
Contributions to the History of North American Ornithology (Jr. and J. A. J. William E.
Davis, Editor). Nuttall Ornithological Club, Cambridge, MA, pp. 113–144.

Rasmussen, P., and R. Prys-Jones (2003). History vs. mystery: the reliability of museum
specimen data. History vs. mystery: the reliability of museum specimen data. Bull. Br.
Ornithol. 123A:66–94.

Rookmaaker, L. C., P. A. Morris, I. E. Glenn, and P. J. Mundy (2006). The ornithological
cabinet of Jean-Baptiste Bécoeur and the secret of the arsenical soap. Archives of Natural
History 33:146–158.

Sclater, P. L. (1893). The Ibis, A Quarterly Journal of Ornithology.

Sclater, P. L., and O. Salvin (1869). On Peruvian Birds collected by Mr. Whitley (1869).

Sclater and Salvin (1867). On Whitely's Arequipa Expeditions. Proceedings of the Zoological
Society of London:21.

Sharpe, R. B. (1906). The history of the collections contained in the Natural History Departments
of the British Museum, Birds.

Stephens, L., and M. A. Traylor (1983). Ornithological Gazetteer of Peru. President and Fellows
of Harvard College, Cambridge, MA.

Vieillot, L. P. (1824). La Galerie des Oiseaux. Constant-Chantpie, Paris, France.

1298