1 **Taxonomy and nomenclature of the giant hummingbirds (*Patagona spp*.) (Aves:**

2 **Trochilidae)**

4 Jessie L. Williamson[1–3*], Chauncey R. Gadek[3–5], Bryce W. Robinson[1,2], Emil Bautista[6], Selina M.
5 Bauernfeind[3], Matthew J. Baumann[3], Ethan F. Gyllenhaal[3,4], Peter P. Marra[7], Natalia Ricote[8],
6 Nadia D. Singh[9], Thomas Valqui[6,10], and Christopher C. Witt[3,4]
7
8
[1]Cornell Lab of Ornithology, Cornell University; Ithaca, 14850, USA.
[2]Department of Ecology and Evolutionary Biology, Cornell University; Ithaca, 14850, USA.
[3]Museum of Southwestern Biology, University of New Mexico; Albuquerque, 87131, USA.
[4]Department of Biology, University of New Mexico; Albuquerque, 87131, USA.
[5]Environmental Stewardship, Los Alamos National Laboratory; Los Alamos, 87545, USA.
[6]Centro de Ornitología y Biodiversidad (CORBIDI); Lima, 15064, Peru.
[7]The Earth Commons Institute; Department of Biology; McCourt School of Public Policy;
Georgetown University; Washington, D.C., 20057, USA.
[8]Facultad de Artes Liberales, Departamento de Ciencias, Universidad Adolfo Ibáñez; Santiago,
7941169, Chile.
[9]Department of Biology, Institute of Ecology and Evolution, University of Oregon, Eugene,
97403, USA.
[10]Facultad de Ciencias Forestales, Universidad Nacional Agraria La Molina; Lima, 15024, Perú.
22

23 Corresponding email: jlw432@cornell.edu

**Abstract**

Giant hummingbirds (*Patagona* spp.) are extraordinarily large but nondescript hummingbirds whose taxonomy was muddled for two centuries. *Patagona* systematics were redefined in a study of migration and genetics by Williamson et al. (2024), revealing two species with divergent genomes and extensive seasonal range overlap. The latter study introduced a new name, *chaski*, for the Northern Giant Hummingbird; however, its brief nomenclatural justification omitted a key historical detail. Here, we revisit that justification by reviewing the taxonomy and nomenclature of the genus *Patagona* in light of its newly-clarified biology and species limits. First, we provide comprehensive information on identification of these cryptic species and report the extent of misidentification present in major collections. Second, we examine the validity of the name *peruviana* (Boucard 1893), long considered to be a poorly defined subspecies, and its potential application to the Northern Giant Hummingbird. Boucard's two putative syntypes are in fact known, and they appear to correspond to the northern genetic lineage; therefore, we designate a lectotype to clarify the entity originally associated with *peruviana*. We show that the entire subsequent history of *peruviana* has involved misapplication to entities that comprise variable mixtures of the two giant hummingbird species. Redefining *peruviana* to match the newly recognized Northern Giant Hummingbird would shift the limits of the entity to which this name refers, sowing confusion due to its non-universal meaning. By contrast, the name *chaski* was introduced concomitantly with the definitive discovery of two *Patagona* species and their taxonomic limits, morphology, physiology, migration, genetics, and year-round distributions. The Principle of Priority (Art. 23) of the International Code of Zoological Nomenclature provides a mechanism to suppress problematic senior synonyms. We invoke Art. 23.9.3 and urge

46   suppression of *peruviana* and adoption of *chaski*, the only name that applies specifically and

47   uniquely to the Northern Giant Hummingbird.

49   **Key words:** Andes, *chaski*, elevation, giant hummingbirds, *Patagona*, taxonomy, migration,

50   Trochilidae

**Introduction**

Giant hummingbirds (*Patagona spp.*) comprise the most distinctive clade in the family

Trochilidae because of their extraordinary size. Within the genus *Patagona*, however,

morphological and plumage variation are subtle and it is difficult to disentangle geographic

variation from age, wear, molt, sex and/or individual variation. Seasonal overlap among cryptic

but distinct geographic breeding populations has exacerbated the challenge of characterizing the

differences among genetic lineages and appropriately defining taxonomy.

A recent study made it possible to accurately adjudicate taxonomy and classification

within *Patagona* for the first time by presenting new information about genetics, migration, and

phenotypes (Williamson *et al.* 2024; Fig. 1). We now know that *Patagona* is a clade with two

biological species that diverged ~2–3 Mya and that differ in breeding latitude and migratory

behavior. The two lineages are largely allopatric during the breeding season. During the austral

winter, the southern lineage, which breeds from central Chile through northwestern Argentina

and likely throughout Bolivia, migrates to tropical latitudes of the central Peruvian Andes where

it overlaps extensively with the northern, high-elevation resident lineage. The challenge of

distinguishing the two cryptic species based on plumage and/or measurements has, until now,

made it difficult to characterize their seasonal range overlap. However, the two lineages are

reproductively isolated and experience no gene flow, making their status as biological species

clear (Williamson *et al.* 2024). Each of the two species, the Northern Giant Hummingbird and

Southern Giant Hummingbird, contains shallow geographic population structure. Additionally,

the two species are vocally distinct (Williamson et al., *In. Prep*). The Northern Giant

Hummingbird, which is a high-elevation resident, averages larger in bill length, wing chord, tail

length, and tarsus length, although there is substantial overlap in morphological characteristics

with the Southern Giant Hummingbird and only ~65–85% of individuals can be identified by

measurements alone when comparing sexes separately. Similarly, subtle plumage differences

between northern and southern forms, which have been noted by previous scholars (e.g.,

Hellmayr 1932; Zimmer 1952) and clarified by Williamson et al. (2024), are useful for

identification but not universally applicable across the range of individual variation (Fig. 1).

With unambiguous species status and newly defined taxon limits for the northern and

southern forms, Williamson et al. (2024) evaluated what name should apply to the northern

species. The name *peruviana* was introduced by Boucard (1893) in a brief description, and has

since been applied, variably, as a subspecies comprising populations from northwestern

Argentina and Bolivia northward. Williamson et al. (2024) argued that the name *peruviana*

should not continue to be used for two main reasons: (1) Historic and persistent misalignment of

the name *peruviana* with the northern genetic lineage; and (2) The type series (syntypes)

appeared to have been comprised of a mix of the two giant hummingbird species. Williamson et

al. (2024) therefore proposed the new species name, *chaski* (*Patagona chaski* sp. nov., Northern

Giant Hummingbird, picaflor gigante del norte in Spanish) and designated a holotype.

The taxonomic description contained in Williamson, et al. (2024), omitted key

information regarding the syntypes of *peruviana* that came to the attention of the authors after

publication. We rectify this omission here. Williamson et al. (2024) did not know that the likely

identities of the type series (syntypes) used by Boucard (1893) to designate *peruviana* were, in

fact, known. Although Boucard did not provide identifiers for the specimens he referenced in his

description, he stated that they were "…collected by Whitely in Peru" (Boucard 1893). At least

16 specimens were collected by Henry Whitely, Jr. in southern Peru from 1867–1873 or later

(Table 1). At least one of these (AMNH-37502), which was assigned to the "northern" form,

corresponds to the Southern Giant Hummingbird based on genetic evidence; a finding that is

expected because we now know that the southern lineage occurs in Peru during the austral

winter. However, Williamson et al. (2024) were not aware of two specimens at the Muséum

National d'Histoire Naturelle in Paris (MNHN; MNHN-ZO-MO-1989-340 and MNHN-ZO-MO-

1989-341) that are considered to be Boucard's (1893) *peruviana* syntypes and are labeled

accordingly (e.g., Jouanin 1950; Fig. 2a,b; Table 1). These syntypes appear to correspond to the

northern lineage based on plumage features, although the female specimen is somewhat atypical.

Their identifications should be considered tentative until confirmed using DNA sequencing.

A guiding purpose of the International Code of Zoological Nomenclature (the Code) is

"to promote stability and universality in the scientific names of animals and to ensure that the

name of each taxon is unique and distinct", thereby ensuring precision and consistency in the use

of terms (ICZN 1999). Williamson et al. (2024) proposed the new species name, *chaski*,

specifically because the name *peruviana* has been persistently misapplied, in variable ways, to an

entity that is a mixture of both the northern and southern lineages of giant hummingbird. Based

on new recognition that the likely syntypes of *peruviana* correspond to the northern lineage, the

names *chaski* and *peruviana* should be considered synonyms, with *peruviana* senior and *chaski*

junior. Art. 23.1 of the Code, the Principle of Priority, states the general principle that "the valid

name of a taxon is the oldest name applied to it…" (ICZN 1999). However, the code allows for

exceptions. For example, Art. 23.9, the Reversal of Precedence, provides a mechanism to favor a

clear junior synonym over a confusingly applied older alternative. Art. 23.9.3 states, "If… an

author considers that the use of the older synonym… would threaten stability or universality or

cause confusion, and so wishes to maintain use of the younger synonym or homonym, he or she

must refer the matter to the Commission for a ruling under the plenary power [Art. 81]. While

the case is under consideration use of the junior name is to be maintained [Art. 82]" (ICZN

1999).

Here, we present the taxonomic history and nomenclature of *Patagona* from 1840 to

present, in light of the newly defined species and the new information about genetics and

migration that clarifies seasonal range limits and methods for identification. We review the

description of the genus *Patagona* Gray 1840, the original taxon (Vieillot 1824), and examine

those of subsequently introduced names, *peruviana* and *boliviana* (Boucard 1893), within the

genus, as well as the specimens key to historical identifications and naming. We further examine

available information about the giant hummingbirds' distributions and migration that is relevant

to our understanding of systematics. Next, we evaluate which name should be applied to the

northern genetic lineage, the species-level entity now known under Northern Giant

Hummingbird (*sensu* Williamson et al. 2024), with the aim of providing guidance to the

International Commission on Zoological Nomenclature (ICZN). To this end, we ask whether

instability, non-universality, or confusion would be caused by elevating the name *peruviana*, as is

indicated by its persistent and continuous misapplication to a mixture of species.

**Taxonomy of the genus *Patagona***

The type specimen of the genus *Patagona* was described by Vieillot (1824) as *Trochilus gigas*,

and was assigned to a type locality of "Brésil" (Brazil; Vieillot 1824). The type locality was later

corrected to Chile by Lesson, who renamed the bird *Ornismiya gigas* (Lesson 1832). In 1840,

Gray named the genus *Patagona*, listing *P. gigas* (ascribed to Vieillot, 1824) as the type species

(Gray 1840). In spite of other proposed names (e.g., *Trochilus tristis*, *Cynanthus tristis*,

*Ornismya gigantea*, *Hypermetra gigas*; see Elliot 1879; Hartert 1900), from this point forward,

*Patagona gigas* entered into widespread use among ornithologists and researchers.

**The Whitely series of *Patagona* specimens**

Henry H. Whitely, Jr., a British naturalist and explorer particularly interested in ornithology

(Gladstone 1927; Sclater 1893), was a key figure in the history of *Patagona* taxonomy. For many

years, Whitely collected specimens in southern Peru, with numerous trips throughout Arequipa

and Cusco between the years of 1867 and 1873 (Chapman 1921; Sclater 1893; Sclater & Salvin

1867, 1869; Stephens & Traylor 1983). During this time, Whitely was credited with a number of

new species discoveries, including the Bearded Mountaineer (*Oreonympha nobilis*; Sclater

1893). Many of Whitely's specimens were dispersed through the specimen trading business of

his father, Henry Whitely, Sr. (Rasmussen & Prys-Jones 2003; Sharpe 1906). Although Whitely

Sr., and to a lesser extent, Whitely Jr.'s, reputations have been questioned (Whitely, Sr. has been

called 'reliably unreliable'; Rasmussen & Prys-Jones 2003), none of the specific problems that

have been found have concerned the localities of Whitely's *Patagona* specimens.

Whitely collected at least 16 *Patagona* specimens in Peru from 1867 to 1873 (Table 1). In

one letter about his 1867 Arequipa trip, Whitely noted, "… now commenced the worst part of the

journey…the way, however, was enlivened by the sight of numerous birds, and especially, for

some 2,000 feet, by the movements of the Giant Humming-bird (*Patagona gigas*)" (Sclater &

Salvin 1867). Whitely's *Patagona* specimens, particularly those from Tinta, Cusco, Peru, have

played a crucial role in history because they are thought to have formed the basis for a

subsequent type description. Whitely collected in Tinta on several trips, including May–June

1868 and January 1869 (Sclater & Salvin 1869). The identities of at least three *Patagona*

specimens from his 1868 trip are known (MNHN-ZO-MO-1989-340, FMNH-45797, and

FMNH-45798), and three others are suspected (AMNH-37500, AMNH-37501, and MNHN-ZO-

MO-1989-34; Table 1). One specimen (AMNH-37501), housed at the American Museum of

Natural History (AMNH) was designated as the "Type of Fig. 33, p. 67" in Giraud Elliot's 1879

synthesis, "*A classification and synopsis of the Trochilidae*" (Elliot 1879). This specimen, a male

from Tinta, was noted in original ledgers as "figured in synopsis", as was AMNH-37502, a

female collected from Cachupata, Cusco, Peru on August 21, 1873 (Table 1). These specimens,

along with two other Whitely specimens, one each from Cachupata and Tinta (AMNH-37499

and AMNH-37500), were obtained by the AMNH in 1932 as part of the collection acquired from

Walter Rothschild (Lanyon 1995;Table 1). Based on available data from gazetteers and related

publications, AMNH-37500 and AMNH-37501 are suspected to have been collected during

Whitely's June–August 1868 trip to Tinta. There are at least three other Whitely specimens

(ANSP-50447, NHMUK 1887.3.22.700, and NHMUK 1913.3.20.291) collected during January

1869, and one additional specimen (ANSP-50448) is suspected to have been collected in either

May–June 1868 or January 1869 (Table 1). At present, Whitely's *Patagona* specimens are known

to reside at the AMNH, MNHN, Academy of Natural Sciences of Drexel University (ANSP),

Field Museum of Natural History (FMNH) and Natural History Museum at Tring (NHMUK),

and it is possible that there are additional Whitely *Patagona* specimens in other collections.

**Taxonomic history of the names *peruviana* and *boliviana***

In 1893, Adolph Boucard, a French ornithologist and specimen trader, proposed the names

*Patagona peruviana* and *Patagona boliviana* for specimens in his collection, "…if the two

should prove distinct species" (Boucard 1893). Boucard wrote, "I have in my collection what I

consider as the type of Vieillot 'Ex Coll Riocour.' My other specimens were collected in Chili by

Reed. I have also three specimens collected by Whitely in Peru, and in Bolivia by Buckley"

(Boucard 1893) (Fig. 2). Boucard appears to reference obtaining Vieillot's type specimen from

the Comte de Riocour collection, of which he obtained most in 1889 (Renshaw 1905). Boucard's

original text does not provide specific identifiers, collection dates, collecting localities, or sexes

for the specimens in his collection from Chile, Peru, or Bolivia. However, two Whitely

specimens at the Muséum National d'Histoire Naturelle in Paris (MNHN), MNHN-ZO-MO-

1989-340 (a male from Tinta, collected 15 June 1868) and MNHN-ZO-MO-1989-341 (a female

from Tinta, no date), were assigned equal status as syntypes for *peruviana*, while the specimen

MNHN-ZO-MO-1989-342 (undetermined sex, no date; from Bolivia), collected by Clarence

Buckley, was designated as the holotype *boliviana* (Jouanin 1950; Fig. 2; Table 1). The names

*peruviana* and *boliviana*, in addition to the nominate *gigas*, were henceforth recognized and

entered into use (e.g., Hartert 1900).

In 1921, Boucard's (1893) original description was contested by Eugène Simon, a

naturalist and systematist from the MNHN. Simon wrote that *Patagona gigas* is a bird of the

high mountains "…without presenting sufficient local variation to be considered as subspecies"

(Simon 1921). Simon stated that Boucard (1893) had "…conditionally proposed the name

*Patagona peruviana* for the specimens from Peru from Whitely's trip and *Patagona boliviana*

for those brought back from Bolivia by Buckley" (Simon 1921). Then, Simon added, "…I have

ascertained, by examining the types, that the coloration characters [Boucard] gives (in Gen.

Humm. Birds. p. 61) are age-related or even completely illusory" (Simon 1921). In one footnote,

Simon noted the white borders on the wing feathers and wing coverts and "the absence of a

rufous part in front of the white uropygial band" on Boucard's types, which are characteristics of

*Patagona* juveniles and immatures (see Fig. 1). In a second footnote, Simon clarified his

description of the phrase "completely illusory", noting that he was discussing, "…for example,

the partly black throat of Peruvian birds" (Simon 1921). Simon did not provide specific

identifiers of the supposed types, but it is presumed from his text that he examined the same

specimens referenced in Boucard's (1893) original description.

In 1930, American ornithologist John Zimmer provided a brief account of *Patagona*

*gigas peruviana* Boucard and *Patagona gigas boliviana* Boucard in *Birds of the Marshall field*

*Peruvian expedition 1922-1923* (Zimmer 1930), recognizing Boucard's names as subspecific

identifiers. Zimmer described examining four recently-collected *Patagona* specimens from Peru,

compared with 23 previously-collected Peru specimens and 13 *gigas* specimens from central

Chile and northwestern Argentina. Zimmer wrote, "The Peruvian birds, with one or two

exceptions, are consistently larger than the Chilean skins, with longer wings and longer, heavier

bills. They also, with one or two exceptions, are more rufescent on the under parts, sometimes

markedly so. Since Dr. Hellmayr plans to discuss the characters of these two races in a

forthcoming paper on the birds of Chile, I will leave further details to his able pen" (Zimmer

1930).

Two years later, Austrian ornithologist Carl Hellmayr published *The Birds of Chile*

(Hellmayr 1932). In his account for the giant hummingbird, Hellmayr wrote, "Although the late

Eugène Simon (Hist. Nat. Troch., p. 157) questioned the possibility of discriminating any

geographic races of the Giant Humming bird, the study of between fifty and sixty properly

labeled specimens from the whole range clearly indicates the existence of two forms" (Hellmayr

1932). Hellmayr said Simon's assessment that *P. peruviana* and *P. boliviana* were *nomina nuda*

was a mistake, yet he agreed with Simon when he noted that "…several of the characters claimed

by Boucard prove to be individual" (Hellmayr 1932). Hellmayr went on to describe plumage

differences of populations from Chile compared to those from Ecuador, Peru, and Bolivia, noting

that the extent and intensity of rufous color to the underparts seemed to be "purely individual"

and unrelated to sex or age (Hellmayr 1932). Hellmayr again agreed with Simon with respect to

certain plumage features. Hellmayr concluded, "While the palest examples of the northern form

can be closely matched by one or two unusually rufous-bellied birds from central Chile, the

general run of the two series is easily told apart" (Hellmayr 1932). Hellmayr's resolute

conclusion seemed to leave no doubt that there were two forms that were easily distinguished;

this would later prove false (see below).

Based on his analyses of plumage, Hellmayr offered nomenclatural suggestions: "The

larger northern race is entitled to the name *P. gigas peruviana*, tentatively proposed by Boucard

for a specimen from Peru in his collection. The type, a male obtained by H. Whitely on June 15,

1868, at Tinta, Dept. Cuzco, agrees with the average of our Peruvian series, while *P. boliviana*,

was based on an individual variant with wholly cinnamon-rufous under parts…" (Hellmayr

1932). Hellmayr thus cast doubt on the name *boliviana*, while bolstering the name *peruviana*.

Hellmayr noted that "…An adult male from Tilcara, Jujuy (Nov. 24, 1905; L. Dinelli), and a

couple from Lara, Tucuman (Feb. 12, 1903; G. A. Baer), all in the Tring Museum, are in every

particular typical of the large northern form" (Hellmayr 1932). Years later, Hellmayr would add

that specimen no. 868/48 collected by L. Dinelli on August 10, 1916, from Colalao del Valle,

Tucumán, Argentina was "typical" of *peruviana* based on plumage and morphology (Revilliod *et*

*al.* 1942).

Based on Hellmayr's account, largely regarded as definitive in the literature, subsequent

authorities recognized *peruviana* as a valid subspecies and ceased recognizing *boliviana*, folding

*boliviana* into descriptions for *P. g. peruviana* (Peters 1945; Zimmer 1952). In *Check-list of the*

*Birds of the World*, Peters wrote that "The breeding bird of northwestern Argentina (Jujuy,

Catamarca, and Tucumán) is probably referable to [*P. g. peruviana*]" (Peters 1945).

In 1952, John Zimmer revisited *Patagona* taxonomy in *Studies of Peruvian Birds*

(Zimmer 1952). Under his entry for *P. g. peruviana*, he stated, "This northern form of the species

is fairly readily determinable in distinction from the southern *gigas gigas* when fully adult birds

are compared, but young examples can be confusing" (Zimmer 1952). Zimmer went on to

describe the difficulty of distinguishing juvenile birds by plumage. He noted, "All the adult

examples from [the Urubamba Valley of Peru] are clearly *peruviana*. I have seen no Peruvian

specimens that indicate the occurrence of *gigas gigas* in Perú, although there is evidence that the

subspecies has some migratory tendencies, as noted by Hellmayr…" (Zimmer 1952). Zimmer

presented a list of 83 specimens, from numerous localities and of both sexes, that he examined as

part of his treatment. He wrote, "I would emend a statement made by Hellmayr (*loc. cit.*) with

reference to certain Argentina specimens that he found to be 'in every particular typical of the

large northern form.' Three females from Tilcara are fairly typical *peruviana* but the others are

identifiable only by their measurements, since they are immature and in coloration resemble

other young of both forms" (Zimmer 1952).

Historical descriptions of plumage and range by Hellmayr, Zimmer, and others, have

since been regarded as definitive, without detailed examination of variation between the

suspected races of *Patagona*. For example, in 1974, Fernando Ortiz-Crespo supported previous

scholars, noting "[The race *peruviana*]…differs from the nominate race, occurring in central

Chile, adjacent Argentina, and the southern Chilean provinces, in having a larger body size, a

stouter, longer bill, and rufescent underparts" (Ortiz-Crespo 1974).

Thus, since ca. 1930, the giant hummingbird has been considered to contain two forms:

The nominate *P. g. gigas*, a suspected migrant and apparently smaller-bodied bird with a shorter

bill and shorter wings, stated to range from approximately central Chile to central Argentina; and

*P. g. peruviana*, an apparently larger-bodied form with a longer bill and longer wings, thought to

range (and breed) continuously from northern Chile through Peru and Ecuador, and through

Bolivia and northwestern Argentina. Individuals of *peruviana* were believed to have more

rufescent underparts and the two forms were described as being "easily distinguishable" based

on plumage and morphology; we now know this is not the case (see below).

**Uncertainty about migration added to taxonomic confusion**

Inadequate information about migration contributed to longstanding uncertainty about *Patagona*

taxonomy. Migration of the southern temperate populations was long suspected, and naturalists

throughout history puzzled over where Chilean populations 'disappeared' to after breeding

(Goodall *et al.* 1946). Even recently, the *Atlas of the Breeding Birds of Chile 2011-2016* called

the winter whereabouts of Chilean-breeding giant hummingbirds an "unsolved mystery"

(Medrano *et al.* 2018).

Darwin was among the earliest to note migration of southern giants: During his voyage

on *The Beagle*, he remarked, "At Valparaíso, in the year 1834, I saw several of these birds in the

middle of August, and I was informed they had only lately arrived from the parched deserts of

the north" (Darwin 1839). German-Chilean botanist Federico Johow wrote that giant

hummingbirds are "rather tropical" and were observed in central Chile only during the Chilean

summer; he noted that their presence along the coast of Chile coincided with the flowering of

*Lobelia* and columnar cacti, proposing that they wintered in the northern provinces of "…Peru,

Ecuador, etc." (Johow 1910, 1921). Hellmayr (1932) stated that *Patagona* is absent from central

Chile during June–August, writing, "The birds probably migrate northwards, but a few, at least,

cross the Andes on their migration, as is shown by specimens… which clearly pertain to the

Chilean race" (Hellmayr 1932). Supporting this idea, *Birds of the High Andes* posited that

Chilean breeders migrate across the Andes to adjacent western Argentina (Fjeldså & Krabbe

1990), while others have suggested that they migrate to northwestern Argentina (Sánchez Osés

2003). More recently, and without citing evidence, the guide "*Biodiversidad de Chile,*

*patrimonio y desafíos*" stated that Chilean giant hummingbirds migrate to Ecuador for the austral

fall (Ambiente 2008).

Chilean naturalist Rafael Barros offered what is perhaps the most detailed early treatment

of *Patagona* migration. In 1952, Barros described Chilean-breeding giant hummingbirds as

"…fleeing from the rainy season and the cold" (Barros 1952). Barros wrote, "It is not well

elucidated if, when it leaves its places of residence…it actually moves to the provinces located to

the north of Aconcagua (where my observations begin), or it crosses the high mountain range of

the Andes and passes to the Republic of Argentina, or if it goes to winter in some regions of

Bolivia or Peru. I know of no observations in this regard, and this is a problem to be solved, of

great scientific interest" (Barros 1952). Barros posited the question: "Does it go up…to the

heights of the Puna to cross the Andes?", later responding: "Based on what I have been able to

observe for many years… I do not believe that it crosses the mountain range in the central part of

the country, during its pilgrimages, either on its outward journey or on its return, because in the

mountainous region it first reaches the lower parts, the pre-cordillera, the foothills of the

Cordillera and fields beneath them, gaining altitude afterwards" (Barros 1952). In this same year,

Zimmer cast doubt on the extent of northward travel of migratory temperate breeders, stating that

he had seen no Peruvian specimens indicating the occurrence of *P. gigas gigas* in Peru (Zimmer

1952).

In 1974, Ecuadorian ornithologist Fernando Ortiz-Crespo published a detailed treatment

of the ecology of giant hummingbirds in Ecuador (Ortiz-Crespo 1974). Ortiz-Crespo briefly

noted that the giant hummingbird population in central Chile "…builds up during the spring and

summer", attributing their movement to northward migration (Ortiz-Crespo 1974). Ortiz-Crespo

credited Johow (1910) and Darwin (1839) as his sources for migration information. In this same

study, Ortiz-Crespo analyzed morphology between sexes and *Patagona* subspecies (which he

assigned based on collection country), stating that his results "...do not seem to confirm the

possibility that Ecuador is regularly visited on migration by birds which nest in central Chile, and

whose winter quarters thus remain to be discovered" (Ortiz-Crespo 1974). Twelve years later, in

a different manuscript about austral hummingbird migration, Ortiz-Crespo expressed similar

uncertainty about *Patagona* migration, stating "…it is not known for sure where they winter"

(Ortiz-Crespo 1986). He described examining 180 specimens, including 156 specimens from

outside of Chile, only four of which (ANSP-145417 and FMNH-179394 from Bolivia, ANSP-

167558 from Argentina, and FMNH-67518 from Peru) he said had measurements and coloration

of the nominate form *gigas* (we examined these same specimens at ANSP and FMNH, and have

confirmed that they are Southern Giant Hummingbirds). Ortiz-Crespo concluded, "This meager

evidence suggests that southern populations disperse northward after nesting" (Ortiz-Crespo

1986), providing no additional specifics. He further suggested that examining specimens

collected during the austral fall and winter from northwest Argentina and Bolivia would be

important for understanding distribution and movement (Ortiz-Crespo 1986).

Since Darwin's mention of migration in 1839, many published accounts have mentioned

*Patagona* migration or presence/absence of *Patagona* at certain times of the year in Chile with

few additional details (e.g., (Aguirre 2001; Barros 1952; Estades *et al.* 2008; González-Gómez &

Valdivia 2005; Gould 1861; Medel *et al.* 2022; Rueda-Uribe *et al.* 2023; Speziale & Lambertucci

2009; Velásquez-Noriega *et al.* 2021)). One publication stated that *Patagona* migrates 2,000 km

(Rodríguez-Flores *et al.* 2019), but no information about the migratory route or sources for

distance traveled were provided. The many, pre-2024 sources cited above corroborated the

longstanding idea that Chilean *Patagona* populations migrate, but none provided detailed or

definitive information about where these populations travel or how they get there.

Largely owing to a paucity of information about the migration of southern *Patagona*

populations, the ranges and distributions of the putative giant hummingbird subspecies remained

a source of confusion for their entire history. Gould (1861), though he did not mention forms,

confirmed *Patagona* presence in Ecuador, Peru, Bolivia, and Chile, with no mention of

Argentina (Gould 1861). Later, in his 1893 description, Boucard mentioned that the three forms

(*gigas, peruviana, and boliviana*) were found in Ecuador, Peru, Bolivia, and Chile (Boucard

1893), providing no mention of Argentina. Hartert (1900), using Boucard's proposed names,

noted that *P. peruviana* and *P. boliviana* were found in the "Andes from southern Ecuador to

Chili and Argentina" (Hartert 1900). Hellmayr (1932) wrote that "*P. g. peruviana* … replaces the

typical form in the Andes of Ecuador, Peru, Bolivia, and extreme northern Chile. It is probably

also this race that nests in northwestern Argentina, although I have not been able to examine

undoubted breeding specimens" (Hellmayr 1932). The idea that the race *peruviana* occurs in

northwest Argentina was subsequently corroborated by many key historical accounts (Barros

1952; Fjeldså & Krabbe 1990; Koepcke 1983; Ruschí 1961; Zimmer 1952). In one, several

Southern Giant hummingbirds from Argentina and Bolivia were identified as *peruviana* (Zimmer

1952)*.* This information continues to persist into recent years (e.g., Pearman & Areta 2020;

Sánchez Osés 2003; Velásquez-Noriega *et al.* 2021). In *Birds of Chile: A Photo Guide*, Howell

and Schmitt (2018) state that there is "no range overlap" between *peruviana* and *gigas* (Howell

& Schmitt 2018). We now know northern Argentina contains only Southern Giant

Hummingbirds, and that there is substantial range overlap of the two giant hummingbird species

during the non-breeding season.

Uncertainty about migration, and confusion about the numbers of possible

subspecies/forms and their respective distributions (e.g., Fjeldsa & Barbosa 1983), has

additionally resulted in misleading generalizations about their elevations of residence. Woods

(1998) suggested that the two forms, *peruviana* and *gigas*, might be elevational replacements and

incompletely characterized the Southern Giant Hummingbird range, writing, "The northern

subspecies of Giant Hummingbird *P. g. peruviana* regularly reaches paramo and puna habitats in

Ecuador and Peru at 3,500-4,000 m, while the southern *P. g. gigas* occurs seasonally to below

2,000 m in central Chile and adjacent Argentina" (Woods *et al.* 1998). Woods does not mention

latitudinal migration in southern temperate breeders. In a text about *P. g. peruviana*, Ruschi

(1961) states that it is found in the Andes, varying in altitude from 2,500 to 5,000 meters (Ruschí

1961), but he does not mention *P. g. gigas* or its range. In many other texts, it is not possible to

determine which form of giant hummingbird is being discussed.

The migration of Chilean giant hummingbirds has been suspected and discussed since at

least 1834, but every aspect of the migratory journey had remained a mystery ever since, and no

published works described population genetic connectivity until 2024. Two previous studies used

open citizen science data to coarsely estimate the magnitude of elevational shift experienced

seasonally by Chilean-breeding populations (Rueda-Uribe *et al.* 2023; Williamson & Witt

2021b), and both studies underestimated the magnitude of elevational shift. Based on findings

from Williamson et al. (2024), we now know that the conspicuous giant hummingbirds are

comprised of two phenotypically cryptic species that overlap at the same sites and elevations

during the southern hemisphere winter, approximately May–September (Williamson *et al.* 2024).

The Southern Giant Hummingbird is a long-distance latitudinal migrant, traveling up to 8,300

km roundtrip in a circular loop across the Andes (Williamson *et al.* 2024). These data are even

more striking when put into a global context: The Southern Giant Hummingbird has one of the

two longest hummingbird migrations in the world, as its journey is similar to the typical distance

traveled by the Rufous Hummingbird (*Selasphorus rufus*), which flies up to ~8,200 km roundtrip

annually (Supp *et al.* 2015). At least some Southern Giant Hummingbird populations

shift >4,100 meters in elevation twice annually, making this species one of ~100 taxa that

undertake Elevational Niche-Shift Migration, or 'extreme' shifts in elevation between the

breeding and non-breeding seasons (Williamson *et al.* 2024; Williamson & Witt 2021b; a). These

discoveries, made possible with the use of miniaturized tracking devices and genomic

sequencing—technologies that were not available to Darwin, Johow, or any of the workers cited

above—have finally permitted the resolution of longstanding mysteries about migration and

distribution, clarifying taxonomy within this clade.

**Was a lectotype ever designated for *peruviana*?**

In their recent paper, Williamson et al. (2024) designated a holotype for the unique species

*Patagona chaski* (CORBIDI AV-012403, MSB:Bird:33080; PERU: dept. Cusco, Prov.

Urubamba, district Ollantaytambo, loc. Choquechaca, 13°11.620' S, 72°13.972' W, elev. 4,030

meters, 8 March 2009), based on diagnostic whole nuclear genomic and mitochondrial genomic

DNA-sequence data, confirmed locality, plumage, measurements, and physiological

characteristics. The name *chaski* is available and was registered on ZooBank (LSID:

urn:lsid:zoobank.org:pub:D2D498B7-1427-4A56-81F9-BFEA8C9CB6F5). Williamson et al.

(2024) introduced the name *chaski* on the grounds that the name *peruviana* was *nomen dubium*

and had never been applied consistently with the northern genetic lineage. An alternative path

would have been to designate a lectotype for *peruviana*. This raises the question of whether a

lectotype for *peruviana* has already been appropriately designated under the Code. Boucard's

(1893) mention of multiple specimens in the original description causes those specimens to

comprise syntypes, making it necessary to designate a single type, or lectotype. If Boucard had

instead designated a holotype for *peruviana*, the lectotype concept would have been unnecessary.

Confusion about the existence of a *peruviana* lectotype stems from pg. 231 in Hellmayr's

(1932) *Birds of Chile*, and possible ambiguity in the relevant section of the Code. In Boucard's

(1893) original description for *peruviana* and *boliviana*, he mentioned the existence of multiple

types ("…the specimens from Peru…") (Boucard 1893). However, Hellmayr noted that the name

*P. g. peruviana* was tentatively proposed by Boucard for "…a specimen from Peru in his

collection". Hellmayr called this individual, "the type", which he described as "…a male

obtained by H. Whitely on June 15, 1868, at Tinta, Dept. Cuzco…" (Hellmayr 1932). We cannot

be certain whether Hellmayr himself examined Boucard's putative syntypes, and no ledgers or

records exist at the MNHN to confirm that Hellmayr visited the collection (MNHN, *pers.*

*comm.*). Hellmayr wrote only "the type examined in the Paris Museum…" (Hellmayr 1932).

Under one interpretation of the Code, Hellmayr's assertion of the identity of "the type"

might constitute a lectotype designation. Art. 74.5 states that "…either the term "lectotype", or

an exact translation or equivalent expression (e.g. "the type"), must have been used or the author

must have unambiguously selected a particular syntype to act as the unique name-bearing type of

the taxon" (ICZN 1999). Furthermore, Art. 74.1.3 "…permanently deprives all other specimens

that were formerly syntypes of that nominal taxon of the status of syntype". Under this

interpretation, MNHN-ZO-MO-1989-340 was designated as a lectotype by Hellmayr in 1932 and

the female from Boucard's syntype series (MNHN-ZO-MO-1989-341) would therefore be a

paralectotype with no name-bearing function.

However, under an alternative interpretation of the Code, Hellmayr may not have

designated a lectotype. This is because Hellmayr's assertion that there was a single "type"

contradicted Boucard's original description of multiple "types" (Boucard 1893); if this was a

selection by Hellmayr, rather than an error, it was not made 'unambiguously'. Art 74.6 of the

Code allows authors to restrict a type series to one holotype under certain conditions ("When it

has been accepted that a nominal species-group taxon was based on a single specimen and the

original description neither implies nor requires that there were syntypes, and if it is considered

subsequently that the original description was based on more than one specimen, the first author

to have published before 2000 the assumption that the species-group taxon was based upon a

single type specimen is deemed to have designated that specimen as the lectotype.") Article 74.6

would therefore allow Hellmayr's (1932) publication to constitute a lectotype designation if the

original description of the name *peruviana* (Boucard 1893) had never referred to a type series.

However, given that Boucard's original description referenced "…the specimens from Peru…"

in the plural, he made it unequivocally clear that there was not only a single type specimen, a fact

confirmed by the MNHN (Jouanin 1950). Additionally, Art. 72.4.1 states that all specimens

mentioned in the original description are type specimens, unless explicitly excluded ("72.4.1.

The type series of a nominal species-group taxon consists of all the specimens included by the

author in the new nominal taxon (whether directly or by bibliographic reference), except any that

the author expressly excludes from the type series [Art. 72.4.6], or refers to as distinct variants

(e.g. by name, letter or number), or doubtfully attributes to the taxon."). Thus, under this

interpretation, Hellmayr's (1932) statements do not constitute designation of a lectotype, and the

taxon *peruviana*, as validly described by Boucard (1893), continues to be based on a type series

consisting of multiple syntypes.

**A lectotype for the taxon *peruviana***

In order to fix the identity on the taxonomic concept associated with the name *peruviana*, we re-

designate or designate—depending on the interpretation of the Code—as its lectotype the male

specimen (MNHN-ZO-MO-1989-340; Fig. 2a) collected in Tinta, Cusco, Peru, on June 15, 1868,

by Henry Whitely, Jr. This is the same specimen that was subsequently singled out by Hellmayr

(1932). The specimen is currently housed at the Muséum National d'Histoire Naturelle in Paris

(MNHN). This designation satisfies Arts. 74.7.1, 74.7.2 and 74.7.3 (both original and amended

versions; (ICZN 1999, 2003)), and is in accordance with Recommendations 74A and 74C.

Importantly, this designation fixes the identity of *peruviana* to the individual described in

Hellmayr (1932), thereby fulfilling a primary objective of the ICZN to promote stability of

scientific names (ICZN 1999). Selection of MNHN-ZO-MO-1989-340 as the lectotype results in

MNHN-ZO-MO-1989-341 (female, no date; Fig. 2b) becoming a paralectotype of *peruviana*.

Selecting the female specimen as the lectotype would have needlessly contradicted Hellmayr's

(1932) text, which has been subsequently referenced by other authorities (e.g., (Peters 1945;

Zimmer 1952).

By selecting the specimen MNHN-ZO-MO-1989-340, which appears to correspond to

the northern giant hummingbird lineage, as the *peruviana* lectotype, it could be argued that

*peruviana* has priority over *chaski* as the scientific name of the northern genetic lineage.

However, we argue that *peruviana* has been applied with substantial taxonomic confusion and, as

a result, it should be suppressed. To retain *peruviana* despite its tumultuous history and the

inconsistency of its previous applications to mixtures of two distinct biological species would be

promoting the continuation of instability and confusion rather than embracing the scientific

advances that have finally clarified the taxonomy of giant hummingbirds. The single specimen

that now underpins the name *peruviana* matches the northern genetic lineage due, in no small

part, to chance. It was collected in an area and season where both lineages occur, and

ornithological experts throughout history have never proven able to distinguish the two lineages

consistently and without error. Because the name was misapplied for over a century, an

adjustment in its meaning would comprise taxonomic instability and would cause the name to

have multiple definitions throughout the literature (i.e., to be non-universal). All of this would

lead to additional confusion. The Principle of Priority (Art. 23), a pillar of the Code, provides a

remedy for this situation by allowing suppression of senior synonyms whose meaning has been

similarly muddled.

**Unreliable historical identification of "*peruviana*" by morphology**

Hellmayr's (1932) account is often regarded as the definitive descriptive text for *Patagona*

taxonomy and has been subsequently cited by numerous authorities, particularly in reference to

distinguishing the nominate *gigas* from *peruviana*. Hellmayr examined "…between fifty and

sixty properly labeled specimens from the whole range…", writing that there are "clearly" two

forms and that "…the general run of the two series is easily told apart" (Hellmayr 1932). In his

text, Hellmayr presented a table of measurements from 45 specimens, from Ecuador (no

localities given), Peru, Bolivia (one specimen; no locality), Argentina, and Chile. Hellmayr's

data suggested that *P. g. gigas* was markedly smaller and did not overlap morphologically with *P.*

*g. peruviana*, supporting his conclusion that there were two forms that were "clearly" and

"easily" distinguished by morphology and plumage.

However, Hellmayr's conclusions about the ease with which *gigas* and *peruviana* could

be distinguished are problematic when examined in detail. They directly contradict analyses

conducted by Ortiz-Crespo (1974) and Williamson et al. (2024). Among 361 *Patagona*

specimens from Ecuador, Peru, Bolivia, Argentina, and Chile, Williamson et al. (2024) found

substantial overlap in morphological characteristics—including the same three measured traits

studied by Hellmayr—between the two giant hummingbird species (Williamson *et al.* 2024).

Williamson et al. (2024) found that Northern and Southern Giant Hummingbirds differed

significantly, on average, in bill length, wing chord, and tail length, respectively; however,

average trait differences were extremely subtle, on the scale of millimeters (e.g., ~1-5 mm)—

size differences between the two species are not typically distinguishable by eye, and the ranges

of trait values are similar for both species (Williamson *et al.* 2024). Similarly, within each giant

hummingbird species, sexes differed significantly but not substantially in nearly all size

measurements, although they are not sexually dimorphic in plumage. When analyzing

morphological data separately for each sex, a Linear Discriminant Analyses using data from

three traits (bill, wing, and tail) combined could distinguish Northern from Southern Giant

Hummingbird adult females in only 65.4–82.8% of cases and adult males in 79.5–85.7% of

cases, depending on the model set (Williamson *et al.* 2024). Additionally, Williamson et al.

(2024) found that the mass ranges of the two species are nearly identical. Thus, it would be

unwise to characterize the two giant hummingbird species on the basis of any single

morphological trait, and it would be imprudent to use any single measurement to attempt to

identify individuals to species. In light of this and the substantial degree of individual variation in

plumage (e.g., Fig. 1), the two giant hummingbird species are challenging to distinguish.

To understand how Hellmayr (1932) reached the conclusion that the two forms are easily

distinguished, we compared Hellmayr's original data (*n*=45 adults) with our own data, which

included published data from Williamson et al. (2024), measurements from wild-caught Northern

and Southern Giant Hummingbirds captured in Peru from July–August 2023, and data from

specimens at ANSP and FMNH (*n*=526 individuals total). After removing juveniles and filtering

only for individuals of known sex, locality, and age, our dataset included *n*=393 adults.

Hellmayr's data showed stark differences in bill, wing, and tail measurements for each

sex and species, with almost no overlap in trait distributions; these results sharply contrasted our

own, which showed substantial overlap in distributions for bill, wing, and tail measurements of

both sexes (Fig. 3). Hellmayr found significant differences between *gigas* and *peruviana* tail

measurements, for each males and females, where we found none between Southern and

Northern Giant Hummingbirds (Fig. 3c,f). Additionally, Hellmayr's *peruviana* measurements

were significantly greater than our Northern Giant Hummingbird measurements in comparisons

of male and female wing and tail, respectively (Fig. 3b–f). When we conducted a Principal

Components Analysis (PCA) using bill, wing, and tail measurements separately for each sex, we

found that Hellmayr's data clustered tightly and were shifted to the outer edges of the

distributions expected based on our much larger data set (Fig. 4). In this same analysis, the

MNHN male *peruviana* lectotype (MNHN-ZO-MO-1989-340), an apparent Northern Giant

Hummingbird, and the MNHN *boliviana* holotype (MNHN-ZO-MO-1989-342), a Southern

Giant Hummingbird, both fell within Hellmayr's range for *peruviana*, and within a

morphological zone of overlap between male Northern and Southern Giant Hummingbirds; when

analyzed with female data, MNHN-ZO-MO-1989-342 was an outlier for both species (Fig. 4).

The MNHN *peruviana* female paralectotype (MNHN-ZO-MO-1989-341), fell more closely in

line with measurements for Southern Giant Hummingbirds (Fig. 4). Taken together, these data

reinforce that individuals of each the Northern and Southern species may be difficult to

distinguish even with a suite of measurements used in concert.

The differences in trait means between forms in Hellmayr's dataset were noticeably

larger, and Hellmayr's CVs noticeably smaller, than those observed between species in our

dataset (Tables 2–4). For example, the difference between Hellmayr's *peruviana* and *gigas* mean

wing measurements was 2.64x greater for males and 4.21x greater for females than the

respective differences between Northern and Southern Giant Hummingbirds in our data (Fig. 3).

Hellmayr's tail measurements were particularly anomalous: Hellmayr's difference in tail lengths

between female *peruviana* and *gigas* was 19.57x greater than difference in tail lengths between

female Northern and Southern Giant Hummingbirds in our dataset (Fig. 3).

If Hellmayr's two groups, *peruviana* and *gigas*, corresponded to the northern and

southern species, respectively, the properties of his data should be consistent with trait

distributions from our much larger dataset of measurements from these two species. To test this,

we randomly subsampled our data, without replacement, to match Hellmayr's sample sizes, and

generated 10,000 random datasets. We did this separately for each sex and form (i.e., we

subsampled to $n=8$ for Southern males and $n=4$ for Southern females to match Hellmayr's *gigas*

sampling, and to $n=25$ for Northern males and $n=8$ for Northern females to match Hellmayr's

*peruviana* sampling). For each trait and sex, and for each of the 10,000 datasets generated, we

calculated trait means and CVs, resulting in expected distributions against which we could

compare Hellmayr's data. Using these 10,000 resampled data sets, we further calculated the

magnitudes of trait difference between the two giant hummingbird species, separately for each

sex and trait, yielding expected distributions for differences in trait means.

The CVs were smaller and magnitudes of trait difference were larger in Hellmayr's

(1932) data than in the expected distributions generated from our resampled datasets (Figs. 5–6,

Tables 2–5). For all sex-trait combinations, Hellmayr's data were statistically improbable (Figs.

5–6). For all traits and in both sexes, Hellmayr's differences in trait means fell on the outer right

tail of the expected distribution, based on 10,000 simulated datasets (Fig. 5). Across six trait-sex

combinations, the estimated probabilities of obtaining a difference of trait means as large or

larger than Hellmayr's ranged from 0.0001–0.014 probability (Fig. 5). Hellmayr's CVs were

realistic for some sex-trait combinations and unrealistically small for others, as they generally

fell on the outer left tails of the expected trait distributions (Fig. 6). The probabilities of obtaining

CVs as extreme or more extreme than Hellmayr's ranged from 0.001–0.301 for bill length,

0.000–0.178 for wing chord, and 0.0005–0.361 for tail length (Fig. 6).

While Hellmayr's (1932) original data suggested strong morphological differences that

"clearly" indicated the existence of two forms, these data contradict robust comparisons by

Williamson et al. (2024), and our present comparison, which included data from an additional

165 individuals from across the *Patagona* range. Williamson et al. (2024) showed—and we

again demonstrate here—that there is substantial overlap in mensural traits between Northern

and Southern Giant hummingbirds, and between males and females of each species. Hellmayr's

data do not show the expected pattern of trait distribution overlap between giant hummingbird

forms, even when controlling for sex (Fig. 3). Hellmayr's *peruviana* measurements were

consistently greater than those in our larger dataset, sometimes significantly so, for all

comparisons except male bill length (Fig. 3). Importantly, Hellmayr's data are statistically

improbable: For example, based on 10,000 resampled datasets, there was only a 0.004

probability of finding as large of a taxon difference in male bill length as Hellmayr recovered

(Fig. 5). Similarly, there were only 0.002 and 0.007 probabilities of finding Hellmayr's

magnitudes of difference in male and female wing lengths, respectively, between his *gigas* and

*peruviana* categories. In sum, throughout Hellmayr's data, the differences between taxa are too

large and the variability within taxa is too low to have been produced by a random sampling of

the two genetic lineages (species) that we now know to exist.

Why were Hellmayr's (1932) data improbably decisive in distinguishing two *Patagona*

taxa in the paper in which he sought to resurrect the name *peruviana*? First, we must consider the

possibility that Hellmayr had included misidentified specimens in his dataset; we know this to be

true because at least several *P. gigas* were included in his "*peruviana*" series (see below).

Inclusion of misidentified individuals would have had the effect of diminishing between-taxon

differences and inflating the variability within taxa—the opposite of the pattern that we observed

in Hellmayr's data. The only remaining explanation is that Hellmayr did not report complete and

unaltered data. Rather than this being a case of scientific fraud, we speculate that Hellmayr may

have used a combination of plumage assessment, measurements, and locality to assign each

specimen to a taxonomic category; in other words, he may have started with the *a priori*

assumption that the two taxa were non-overlapping in size traits and range, and may then have

selectively included or omitted data that fit with his preconceived assumptions. Considering the

magnitude of the challenge inherent in identifying giant hummingbirds based on plumage, it may

have been Hellmayr's only option to make these educated guesses, seeking consilience of

evidence as he tried to fit a taxonomy to this difficult group. However, his data cannot be

considered valid scientific evidence because of the flawed selection process by which they

appear to have been derived. Hellmayr's taxonomic conclusion that there were "clear"

morphological differences between giant hummingbird forms held sway for nearly a century, but

the distinction that he made was both scientifically invalid and taxonomically false.

**The subspecies name *peruviana* has been misapplied for 130 years**

Could the taxon name *peruviana* apply to the Northern Giant Hummingbird, a distinct and

unique genetic lineage identified by Williamson et al. (2024), and to which its lectotype

corresponds? Since Boucard's (1893) original description, there has been persistent uncertainty

surrounding the diagnostic criteria and subspecies limits for *P. g. peruviana*, as well as the

nominate *P. g. gigas*. While the work of scholars like Hellmayr and Zimmer was overall

admirable and befitting of an era when neither genomic tools or miniature migration tracking

devices existed, the characteristics presumed to be useful in diagnosing the two giant

hummingbird forms have been found to be flawed (in the case of morphology); "illusory", as

noted in 1921 by Simon (in the case of plumage), or misinformed (in the case of using

misleading ideas about morphology and plumage in concert to assume latitudinal and elevational

range limits). The name *peruviana* has been asserted to apply to "the large northern race" of

giant hummingbird; yet, it has not been applied consistently to the taxon to which it is meant to

denote. Instead, *peruviana* has been continuously and variably applied to a mixture of two

distinct species on the basis of erroneous diagnoses of plumage characteristics, morphological

measurements, and ranges. Widespread misclassification of the two *Patagona* taxa as subspecies

and misapplication of the name *peruviana* to both taxa has marked nearly a century of flawed

taxonomy in this group. To continue using *peruviana* by altering its definition, concomitantly

with its rank, would be confusing and would cause the name to have a non-universal definition

pervading the literature, contrary to the spirit of the Principle of Priority (Art. 23).

Hellmayr's text is widely regarded as the definitive treatment for the entity *peruviana*.

Although Hellmayr did not provide specimen identifiers, in a few cases he provides sufficient

information to diagnose specimen identities. He describes three specimens that "…are in every

particular typical of the large northern form…an adult male from Tilcara, Jujuy (Nov. 24, 1905;

L. Dinelli), and a couple from Lara, Tucumán (Feb. 12, 1903; G. A. Baer), all in the Tring

Museum" (Hellmayr 1932). These three specimens, formerly at Tring, were moved to the AMNH

when Walter Rothschild sold his collection in 1932 (NHMUK, *pers. comm.*; AMHH, *pers.*

*comm.*) The adult male from Tilcara described by Hellmayr is AMNH-479595, definitively

identified as a Southern Giant Hummingbird, based on targeted sequence capture of ultra-

conserved elements (UCEs), known breeding season collection locality, and definitive plumage

characteristics (Williamson *et al.* 2024). The other two Lara, Tucumán individuals mentioned by

Hellmayr, are AMNH-479593 and AMNH-479594—these two individuals are both juvenile

Southern Giant Hummingbirds, for which Williamson et al. (2024) also provide UCE data. The

fact that Hellmayr stated that these individuals were "in every particular typical" of the form

*peruviana*, when in fact, they are Southern Giant Hummingbirds ranging in age from juveniles to

adults, highlights that—despite claiming the two forms were "easily told apart"—he could not

actually distinguish them consistently.

Importantly, the *Patagona* series collected by Henry Whitely from Tinta, Peru—of

historical importance for its significance in the description of types—is a mixed series. The

Whitely specimens held at AMNH contain three Northern Giant Hummingbirds (AMNH-37400,

AMNH-37500, AMNH-37501) and one Southern Giant Hummingbird (AMNH-37502), despite

all being labeled "*peruviana*". That Boucard's apparent Whitely syntypes at the MNHN both

appear to be Northern Giant Hummingbirds, based on plumage features (genetic identification is

as yet unknown), has little bearing on whether the name *peruviana* has been correctly applied in

the past, as the identities of these specimens are purely by chance: There is no evidence that any

ornithologist or taxonomist of the 19th or 20th centuries was able to distinguish these two species

or had any idea that they occurred together seasonally.

Misapplication of the name *peruviana* has persisted into the present. At least thirteen

major museum collections whose specimens we have examined possess giant hummingbird

specimens from each Peru, Bolivia, and Argentina that are labeled as *P. g. peruviana*, and which

contain a mix of both the Northern and Southern Giant Hummingbird. Of the 557 specimens

with locality data that we have examined, a total of 222 individuals (40%) are catalogued with

subspecific identifiers. A total of 54 individuals (30.34% of specimens) that are labeled

"*peruviana*" are actually Southern Giant Hummingbirds (e.g., AMNH-139179, LSUMNS-37404,

CUMV-52570, MCZ-262196, FMNH-67518; Fig. 7). One individual (ANSP-145417) from

Bolivia was labeled with both "*peruviana*" and "*gigas*" (and pencil marks are visible from where

"*peruviana*" was erased and where "*gigas*" was instead written; this individual still bears

"*peruviana*" on the back of its tag). Typically, individual *Patagona* collected in Ecuador, Peru,

Bolivia, and northwestern Argentina that appeared large and had reddish underparts were

assigned the name *peruviana* without greater scrutiny; by this measure, many reddish-toned

*gigas* individuals appear to correspond to *peruviana* (e.g., Fig. 1). Thus, the name *peruviana* has

been applied to specimens on the basis of erroneous information about morphology, plumage,

and assumptions about range. Perhaps as a consequence of this confusion, the extent to which

*peruviana* has been applied (or misapplied) in global collections appears to be limited, because

the vast majority of catalogued specimens (~60%) were not assigned subspecific identifiers.

**The world's largest hummingbird needs a robust name**

The migration-tracking, genomic, physiological, and morphological study conducted by

Williamson et al. (2024) demonstrated that high-elevation resident populations of the giant

hummingbird comprise a distinct species, a northern genetic lineage to which they assigned the

name *Patagona chaski* (Williamson *et al.* 2024). This was a novel taxonomic distinction and the

first time that analytical evidence had been presented supporting more than one species of giant

hummingbird. Although the authors were not aware at the time of evidence pertaining to the

identities of specific individual specimens that comprised the syntypes of *peruviana*, the fact that

those syntypes appear to correspond to the northern genetic lineage implies that *chaski* could

now be considered a junior synonym. However, based on the evidence presented above and Art.

23 of the Code, we conclude that use of *peruviana* should be suppressed because its application

has never aligned with the northern genetic lineage; as a result, its continued use would

perpetuate taxonomic and nomenclatural confusion and non-universality.

In accordance with the purpose of the Principle of Priority (Art. 23.2), "prevailing usage

must be maintained when the following conditions are both met (Art. 23.9.1): 1) the senior

synonym or homonym has not been used as a valid name after 1899 (Art. 23.9.1.1); and 2) the

junior synonym or homonym has been used for a particular taxon, as its presumed valid name, in

at least 25 works, published by at least 10 authors in the immediately preceding 50 years and

encompassing a span of not less than 10 years" (Art. 23.9.1.2) (ICZN 1999). However, the Code

provides a mechanism to address situations of undue confusion and instability, such as this one:

"If the conditions of 23.9.1 are not met but nevertheless an author considers that the use of the

older synonym or homonym would threaten stability or universality or cause confusion, and so

wishes to maintain use of the younger synonym or homonym, he or she must refer the matter to

the Commission for a ruling under the plenary power [Art. 81]. While the case is under

consideration use of the junior name is to be maintained [Art. 82]" (Art. 23.9.3).

In light of the evidence presented here, we recommend a reversal of precedence under

Art. 23.9.3. We urge that the name *chaski* Williamson et al. 2024 take precedence and that the

name *peruviana* Boucard 1893 be suppressed. In accordance with guidelines set forth by the

Code (Art. 23.9.3), while under consideration of the plenary power of the Commission, the

junior synonym, *chaski* in the present case, shall be maintained.

**Conclusions**

Before the introduction of data on migration and genetics in 2024, the taxonomy of giant

hummingbirds was impossible to disentangle, in part because of faulty assumptions about ranges

and identification criteria. The taxon name *peruviana* has had a confusing history. To clarify its

status, we (re)designated a lectotype for this taxon by selecting the same individual referenced by

Hellmayr (1932) as "the type". By analyzing the descriptions and data presented by Hellmayr

(1932), we uncovered that the morphological and plumage features thought to be key for

distinguishing *peruviana* from the nominate *gigas* were erroneous; Hellmayr's data showing that

the two giant hummingbird forms were "easily" discriminable were possibly derived from

selective omission of data that did not fit with the overly optimistic assumption that the two taxa

did not overlap in size. As a result, Hellmayr himself misidentified *gigas* specimens as

"*peruviana*", and in his attempt to resurrect the latter taxon, he erred in both its range and

phenotypic diagnosis. The name *peruviana* has been misapplied throughout history, in variable,

non-universal ways, to a mixture of two different giant hummingbird species. Applications of the

name *peruviana* have always been misaligned with the unique species, the Northern Giant

Hummingbird, that was identified and delimited by Williamson et al. (2024). There has been

widespread and ongoing confusion as a result. Continued use of the name *peruviana* would

foster a continuation of taxonomic confusion. The junior synonym, *chaski*, was introduced

concomitantly with definitive discoveries of species limits, phenotypic characterizations,

physiological and behavioral differences, and year-round distributions for the two *Patagona*

species. It is the only name that has been applied specifically and uniquely to the genetically

distinct entity that we now know as the Northern Giant Hummingbird. We therefore urge the

continued use of *Patagona chaski* for the Northern Giant Hummingbird.

**Acknowledgements**

We thank Frank Rheindt, Thomas Pape, Pamela Rasmussen, Alex Figueroa, Oscar Johnson,

Glaucia Del-Rio, Shawn Billerman, Matthew Halley, Vítor Piacentini, and J. V. Remsen, Jr. for

helpful discussion on taxonomy and nomenclature. Assistance in the field was provided by

Kyana Montoya and Marlon Chagua. We are grateful to the museum curators, collections

managers, staff, and students who facilitated this work: Mark Adams, Clare Atkinson, John

Bates, Bentley Bird, Sharon Birks, Patrick Boussès, Mariel Campbell, Eamon Corbett, Joel

Cracraft, Charles Dardia, Scott Edwards, Rob Faucett, Mary Margaret Ferraro, Jérôme Fuchs,

John Gerwin, Gary Graves, Emily Griffith, Andy Johnson, John Klicka, Andy Kratter, Terry Lott,

Irby Lovette, Ben Marks, Nick Mason, Jon Merwin, Chris Milensky, Rob Moyle, Jose Antonio

Otero, Nathan Rice, Mark Robbins, Vanya Rohwer, Sam Rutledge, Dave Steadman, Paul Sweet,

Jeremiah Trimble, Tom Trombone, Hein van Grouw, Jason Weckstein, and J. V. Remsen, Jr.

Collections access, data, and historical information were provided by the Museum of

Southwestern Biology, Muséum National d'Histoire Naturelle in Paris, Field Museum of Natural

History, Academy of Natural Sciences of Drexel University, American Museum of Natural

History, Museum of Comparative Zoology at Harvard University, Louisiana State University

Museum of Natural Science, Smithsonian National Museum of Natural History, Cornell

University Museum of Vertebrates, The Burke Museum at the University of Washington, Kansas

Biodiversity Institute, Florida Museum of Natural History, and Centro de Ornitología y

Biodiversidad (CORBIDI). This work was supported by an NSF Postdoctoral Research

Fellowship in Biology (DBI-2208924) and a Cornell Lab of Ornithology Edward W. Rose

Postdoctoral Fellowship to JLW. Collections visits were supported by an American Philosophical

Society Franklin Research Grant. Data from wild birds measured in Peru were collected under

Servicio Nacional Forestal y de Fauna Silvestre RD-000056-2023-DGGSPFFS-DGSPFS.

IACUC approval was granted under Cornell University protocol 2022-0127.

**References**

**References**

Aguirre, J. (2001) Nidificacion patagonas gigas gigas en una quebrada costera de Chile central.

*Boletín Chileno de Ornitología* 8, 10–12.

Ambiente, C.N. del M. (2008) *Biodiversidad de Chile, patrimonio y desafios*. 2nd ed. M. S.

Jaime Rovira, Jaime Ugalde (Ed). Ocho Libros Editores, Santiago, Chile, 640 pp.

Barros, R. (1952) Apuntes sobre la historial natural de la Pinguera, Patagona gigas. *Anales de la*

*Academia Chilena de Ciencias Naturales* 17, 135–144.

Boucard, A. (1893) *Genera of Humming Birds*. Pardy & Co. Printers, London, UK, 288 pp.

Chapman, F.M. (1921) *The distribution of the bird life in the Urubamba Valley of Peru*.

Washington Government Printing Office, Washington, D.C., 1–168 pp.

Elliot, D.G. (1879) *A classification and synopsis of the Trochilidae*. Smithsonian Institution,

Washington, D.C., 1–300 pp. Available from:

https://archive.org/details/cu31924000050777/page/n61/mode/2up (April 13, 2022)

Estades, C.F., Vukasovic, M.A., Tomasevic, J.A. & Desert, N. (2008) Giant Hummingbirds

(Patagona gigas) Ingest Calcium-rich Minerals. *The Wilson Journal of Ornithology* 120,

651–653. https://doi.org/10.1676/07-054.1

Fjeldsa, J. & Barbosa, C.E. (1983) A Giant Hummingbird from Paramo de Chingaza, Colombia.

*The Wilson Bulletin* 95, 661–662.

Fjeldså, J. & Krabbe, N. (1990) *Birds of the High Andes*. Zoological Museum, University of

Copenhagen, Copenhagen, Denmark, 1–876 pp.

Gladstone, H.S. (1927) Henry Whitely, 1817-1898. *British Birds* XXII, 290–293.

González-Gómez, P.L. & Valdivia, C.E. (2005) Direct and Indirect Effects of Nectar Robbing on

the Pollinating Behavior of Patagona gigas (Trochilidae). *Biotropica* 37, 693–696.

Goodall, J.D., Johnson, A.W. & Bañados, R.A.P. (1946) *Las Aves de Chile, su conocimiento y*
*sus custumbres*.

Gould, J. (1861) *An introduction to the trochilidae, or family of humming-birds.* London, 1–226
pp.

Gray (1840) *A List of the Genera of Birds, with an indicator of the Typical Species of each*
*Genus*. London.

Hartert, E. (1900) *Trochilidae*. Berlin, R. Friedländer und sohn, 1900, 1–275 pp.

Hellmayr, C.H. (1932) *The Birds of Chile*. Field Museum of Natural History, Chicago, IL, USA,
1–488 pp.

Howell, N.G. & Schmitt, F. (2018) *Birds of Chile: A Photo Guide*. Princeton University Press,
Princeton, NJ, USA, 1–240 pp.

ICZN (1999) *International Code of Zoological Nomenclature*. 4th ed. International Trust for
Zoological Nomenclature, London, UK. 306 pp. Available from: https://www.iczn.org/the-
code/the-code-online/ (April 13, 2022)

ICZN (2003) Declaration 44. Amendment of Article 74.7.3. *Bulletin of Zoological Nomenclature*
60, 263.

Johow, F. (1910) Estudios de biología vejetal, I. Sobre algunos casos de ornitolofilía en la flora
Chilena. *Anales de la Universidad de Chile* 126, 27–50.

Johow, F. (1921) Las cactacea de los alrededores de Zapallar. *Revista Chilena de Historia*
*Natural* 15, 152–166.

Jouanin, C. (1950) XXII Bulletin Du Muséum National D'Histoire Naturelle *Catalogue*
*Systématique des types de trocitilidés du Muséum National D'Histoire Naturelle de Paris*.
2nd ed. Muséum National D'Histoire Naturelle, Paris, France, 1–32 pp.

Koepcke, M. (1983) *The birds of the Department of Lima, Peru*. Third. Harrowood Books,

Newtown Square, Pennsylvania, USA, 1–144 pp.

Lanyon, W.E. (1995) Ornithology at the American Museum of Natural History. In: J. and J. A. J.

William E. Davis (Ed), *Contributions to the History of North American Ornithology*. Nuttall

Ornithological Club, Cambridge, MA, pp. 113–144.

Lesson, R.P. (1832) Index géneral et synoptique des oiseaux du genre Trochilus. In: *Index*

*géneral et synoptique des oiseaux du genre Trochilus*. Paris.

Medel, R., López-Aliste, M. & Fontúrbel, F.E. (2022) Hummingbird-plant interactions in Chile:

An ecological review of the available evidence. *Avian Research* 13.

https://doi.org/10.1016/j.avrs.2022.100051

Medrano, F., Barros, R., Norambuena, H. V, Ricardo, M. & Fabrice, S. (2018) *Atlas de las Aves*

*Nidificantes de Chile 2011 – 2016*. 1–672 pp.

Ortiz-Crespo, F.I. (1974) The Giant Hummingbird Patagona gigas in Ecuador. *Ibis* 116, 347–

359.

Ortiz-Crespo, F.I. (1986) Consideraciones sobre las migraciones de dos picaflores neotropicales.

*El Hornero* 12, 298–300.

Pearman, M. & Areta, J.I. (2020) *Birds of Argentina*. Princeton University Press, Princeton, New

Jersey, USA, 1–480 pp.

Peters, J.L. (1945) *Check-list of Birds of the World, Volume V*. Oxford University Press, London,

1–324 pp.

Rasmussen, P. & Prys-Jones, R. (2003) History vs. mystery: the reliability of museum specimen

data. *History vs. mystery: the reliability of museum specimen data. Bull. Br. Ornithol.* 123A,

66–94.

Renshaw, G. (1905) The Reunion Starling. *The Zoologist* 4, 418–422.

Revilliod, P., Carl, J. & Lessert, R. de (1942) *Revue Suisse de Zoologie*. 1–646 pp.

Rodríguez-Flores, C.I., Ornelas, J.F., Wethington, S. & Arizmendi, M. del C. (2019) Are

hummingbirds generalists or specialists: Using network analysis to explore the mechanisms

influencing their interaction with nectar resources. *PLoS ONE*, 1–32.

Rueda-Uribe, C., Herrera-Alsina, L., Lancaster, L.T., Capellini, I., Layton, K.K.S. & Travis,

J.M.J. (2023) Citizen science data reveal altitudinal movement and seasonal ecosystem use

by hummingbirds in the Andes Mountains. *Ecography*, e06735.

https://doi.org/10.1111/ecog.06735

Ruschí, A. (1961) Algumas observacoes sobre Patagona gigas peruviana Boucard. *Boletim Do*

*Museu De Biologia* 23.

Sánchez Osés, C. (2003) Taxonomy, phylogeny, and biogeography of the Andean hummingbird

genera Coeligena LESSON, 1832; Pterophanes GOULD, 1849; Ensifera LESSON 1843;

and Patagona GRAY, 1840 (Aves: Trochiliformes). University of Bonn.

Sclater, P.L. (1893) 5 *The Ibis, A Quarterly Journal of Ornithology*. 1–709 pp.

Sclater, P.L. & Salvin, O. (1869) On Peruvian Birds collected by Mr. Whitley (1869).

Sclater & Salvin (1867) On Whitely's Arequipa Expeditions. *Proceedings of the Zoological*

*Society of London*, 21.

Sharpe, R.B. (1906) *The history of the collections contained in the Natural History Departments*

*of the British Museum, Birds*. 1–451 pp.

Simon, E. (1921) Histoire naturelle des Trochilidae (Synopsis et Catalogue). *Encyclopédie-*

*Roret*, i–vi, 1–416.

Speziale, K.L. & Lambertucci, S. (2009) El picafor gigante (Patagona gigas) en la Provincia de

Río Negro. *Nuestras Aves* 54, 39–40.

Stephens, L. & Traylor, M.A. (1983) *Ornithological Gazetteer of Peru*. President and Fellows of

Harvard College, Cambridge, MA, 284 pp.

Supp, S.R., La Sorte, F.A., Cormier, T.A., Lim, M.C.W., Powers, D.R., Wethington, S.M.,

Goetz, S. & Graham, C.H. (2015) Citizen-science data provides new insight into annual and

seasonal variation in migration patterns. *Ecosphere* 6. https://doi.org/10.1890/ES14-00290.1

Velásquez-Noriega, P., Serrudo-Gonzáles, V., Paca-Condori, A.C. & Larrea-Alcazar, D.M.

(2021) El picaflor gigante de los Andes. *Bolivia Ecológica* 97.

Vieillot, L.P. (1824) *Colibris et Oiseaux-Mouches*. Chez-Desray, Paris, France.

Williamson, J., Gyllenhaal, E.F., Bauernfeind, S.M., Bautista, E., Baumann, M.J., Gadek, C.R.,

Marra, P.P., Ricote, N., Valqui, T., Bozinovic, F., Singh, N.D. & Witt, C.C. (2024) Extreme

elevational migration spurred cryptic speciation in giant hummingbirds. *Proceedings of the*

*National Academy of Sciences* 121, 1–11. https://doi.org/10.1073/pnas.2313599121

Williamson, J.L. & Witt, C.C. (2021a) A lightweight backpack harness for tracking

hummingbirds. *Journal of Avian Biology*, 1–9. https://doi.org/10.1111/jav.02802

Williamson, J.L. & Witt, C.C. (2021b) Elevational niche-shift migration: Why the degree of

elevational change matters for the ecology, evolution, and physiology of migratory birds.

*Ornithology* 138, 1–26. https://doi.org/10.1093/ornithology/ukaa087

Woods, S., Ortiz-Crespo, F. & Ramsay, P.M. (1998) Presence of giant hummingbird Patagona

gigas and Ecuadorian hillstar Oreotrochilus chimborazo jamesoni at the Ecuador -

Colombia border. *Cotinga* 10, 37–40.

Zimmer, J.T. (1930) *Birds of the Marshall field Peruvian expedition 1922-1923*. Field Museum

of Natural History, Chigago, USA.

Zimmer, J.T. (1952) Studies of Peruvian Birds. No. 62 The hummingbird genera Patagona,

Sappho, Polyonymus, Ramphomicron, Metallura, Chalcostigma, Taphrolesbia, and

Aglaiocercus. *American Museum of Novitates* 1595, 1–30.

https://doi.org/10.1029/eo070i046p01481-03

**Figure 1.** Key for distinguishing between Northern and Southern Giant Hummingbirds, highlighting important plumage characteristics described in (Williamson *et al.* 2024), including throat color and pattern and extent of eye-ring and post-ocular spot, as well as morphological characteristics, such as wing length and bill length. Juvenile is represented for comparison. Illustrations: Bryce W. Robinson.

**Figure 2. A–B,** Syntypes for the taxon *peruviana* from Boucard's (1893) collection described in Boucard (1893). **A** MNHN-ZO-MO-1989-340, a male, collected June 15, 1868. **B** MNHN-ZO-MO-1989-341, a female, no date. Both specimens were collected by Henry H. Whitely from Tinta, Peru. **C** Holotype for the taxon *boliviana* from Boucard's (1893) collection, MNHN-ZO-MO-1989-342, undetermined sex, collected in 1875 by Clarence Buckley from Bolivia. Photos: Muséum National d'Histoire Naturelle (MNHN), Paris.

**Figure 3.** Comparison of morphological data (bill length, wing chord, and tail length; all in mm) between Hellmayr (1932) and Williamson et al. (present manuscript). **A–C** Males. **D–F** Females. Hellmayr compared adults of the two suspected forms, *gigas* and *peruviana*. Williamson et al. compared adults of the two species, the Southern and Northern Giant Hummingbird. Sample sizes for each comparison are reported on the x-axis and significant p-values are indicated above each comparison (from t-tests or non-parametric Wilcoxon Rank Sum tests). Box plot horizontal lines represent median values. Each point is a value from a single individual.

**Figure 4.** Principal Component Analysis (PCA) of bill, wing, and tail measurements for *gigas* and *peruviana* individuals from Hellmayr (1932) and Southern and Northern Giant Hummingbirds from Williamson et al. (present study) with MNHN *peruviana* syntypes and *boliviana* holotype. Because Hellmayr's data were organized numerically, rather than by individual, we have included only his records of single individuals per site to ensure accuracy. **A** Males (*n*=165 individuals); **B** Females (*n*=162 individuals). The male *peruviana* lectotype (MNHN-ZO-MO-1989-340), female *peruviana* paralectotype (MNHN-ZO-MO-1989-341), and *boliviana* holotype (MNHN-ZO-MO-1989-342) are plotted for reference; the *boliviana* holotype is of unknown sex and is therefore included in both PCAs. Although the female *peruviana* specimen is plotted, it should be noted that its maxilla is broken (MNHN, *pers. comm.*); however, *Patagona* maxilla and mandible lengths are typically equal. For reference, bill measurements: *peruviana* male: 35.7 mm; *peruviana* female 35.8 mm (mandible only); *boliviana*: 35.8 mm.

**Figure 5.** Expected distributions of trait mean differences between Northern and Southern Giant Hummingbirds, based on 10,000 resampled datasets generated from Williamson et al. (present study). **A–C** Expected male trait differences for each bill, wing, and tail; **D–F** Expected female trait differences for each bill, wing, and tail. Red dashed vertical lines indicate the differences between *gigas* and *peruviana* trait means from Hellmayr (1932) data, with the estimated probabilities of obtaining Hellmayr's values (red text). Estimated probabilities were calculated from 10,000 simulated datasets.

**Figure 6.** Expected distributions of coefficients of variation (CVs) for each Northern and Southern Giant Hummingbirds, based on 10,000 resampled datasets generated from Williamson et al. (present study). CVs were estimated separately for each species (Northern and Southern Giant Hummingbird), each sex (males and females), and each trait (bill, wing, and tail). **A–C** Northern males; **D–F** Northern females; **G–I** Southern males; **J–L** Southern females. Red dashed vertical lines indicate the CVs from Hellmayr's (1932) original data, with the estimated probabilities of obtaining Hellmayr's values (red text). Estimated probabilities were calculated from 10,000 simulated datasets.

**Figure 7.** Visual representation of the *Patagona* specimen record as it pertains to previous versus current identifications. A total of 557 specimens were examined from 13 museum collections. Of these, 30 specimens are labeled as *P. g. gigas* and pertain to the Southern Giant Hummingbird (*Patagona gigas*). A total of 178 specimens are labeled as *P. g. peruviana*; of these, 124 specimens pertain to the Northern Giant Hummingbird (*Patagona chaski*) and 54 specimens pertain to the Southern Giant Hummingbird (*Patagona gigas*).

**Table 1.** Summary of *Patagona* specimens collected from southern Peru by Henry H. Whitely, Jr. One additional individual (collected ca. July 20, 1867 in Arequipa; known from Whitely's letter to Philip Lutley Sclater and Osbert Salvin, 1867; (Sclater & Salvin 1867)) is suspected but remains unidentified. Additionally, ANSP-50448 appears to be a Whitely specimen, but its origins and provenance are unknown (ANSP, *pers. comm.*) Museums: AMNH = American Museum of Natural History; ANSP = Academy of Natural Sciences of Drexel University; FMNH = Field Museum of Natural History; MNHN = Muséum National d'Histoire Naturelle; NHMUK = The Natural History Museum at Tring. [*]Indicates status as "type of fig" in Elliot 1879. [**]Equal status in AMNH ledgers with AMNH-37501. [‡]Indicates status as a *peruviana* syntype (Jouanin 1950). [^]MNHN-ZO-MO-1989-341 is listed in the MNHN database as having a collection date of "ca. 1868", which was estimated based on Whitely's Peru collection dates. However, the specimen has no date information, but was likely collected during 1868–1878 (MNHN, *pers. comm.*), and definitively before Whitely's death in 1892.

| Specimen identifier | Museum | Country | Department | Specific Locality | Collection Date | Putative Identification | Evidence |
|---|---|---|---|---|---|---|---|
| AMNH-37500 | AMNH | Peru | Cusco | Tinta | no date | Northern | Genomic data; plumage not diagnostic |
| AMNH-37501[*] | AMNH | Peru | Cusco | Tinta | no date | Northern | Genomic data; plumage not diagnostic |
| AMNH-37499 | AMNH | Peru | Cusco | Cachupata | 21 Aug 1873 | Northern | Genomic data; plumage not diagnostic |
| AMNH-37502[**] | AMNH | Peru | Cusco | Cachupata | 21 Aug 1873 | Southern | Genomic data; plumage not diagnostic |
| MNHN-ZO-MO-1989-340[‡] | MNHN | Peru | Cusco | Tinta | 15 June 1868 | Northern | Plumage (diagnostic) |
| MNHN-ZO-MO-1989-341[‡,^] | MNHN | Peru | Cusco | Tinta | no date | Northern | Plumage (suggestive) |
| FMNH-45797 | FMNH | Peru | Cusco | Tinta | 25 May 1868 | Northern | Plumage (diagnostic) |
| FMNH-45798 | FMNH | Peru | Cusco | Tinta | 20 June 1868 | Northern | Plumage (diagnostic) |
| ANSP-50447 | ANSP | Peru | Cusco | Tinta | 4 Jan 1869 | Northern | Plumage (diagnostic) |
| NHMUK 1887.3.22.700 | NHMUK | Peru | Cusco | Tinta | 4 Jan 1869 | Northern | Plumage (diagnostic) |
| NHMUK 1913.3.20.291 | NHMUK | Peru | Cusco | Tinta | 4 Jan 1869 | Northern | Plumage (diagnostic) |
| NHMUK 1913.3.20.293 | NHMUK | Peru | unknown | W. Peru | no date | Northern | Plumage (diagnostic) |
| NHMUK 1887.3.22.701 | NHMUK | Peru | Arequipa | Chiguata | 10 June 1867 | Unknown (juvenile) | – |
| NHMUK unregistered | NHMUK | Peru | Arequipa | Chiguata | 10 June 1867 | Unknown (juvenile) | – |
| NHMUK 1913.3.20.294 | NHMUK | Peru | unknown | W. Peru | no date | Unknown (juvenile) | – |

**Table 2.** Morphological summary statistics for *gigas* (Hellmayr 1932) and Southern Giant Hummingbird (Williamson et al., present study) males and females. For each measured parameter, we provide sample size (*n*), mean, standard deviation, range (minimum and maximum values), and coefficient of variation.

| Dataset | Sex | Parameter (units) | *n* | Mean | Standard Deviation | Minimum | Maximum | Coefficient of Variation |
|---|---|---|---|---|---|---|---|---|
| Hellmayr 1932 | | Bill | 8 | 34 | 0.76 | 33 | 35 | 0.02 |
| | | Wing | 8 | 125.5 | 3.34 | 120 | 129 | 0.03 |
| | Male | Tail | 8 | 80.5 | 1.41 | 79 | 83 | 0.02 |
| Williamson et al. (present study) | | Bill | 59 | 36.27 | 2.4 | 32.41 | 42.25 | 0.07 |
| | | Wing | 59 | 127.64 | 6.74 | 111 | 141 | 0.05 |
| | | Tail | 57 | 81.88 | 4.86 | 71 | 92 | 0.06 |
| Hellmayr 1932 | | Bill | 4 | 34.25 | 0.5 | 34 | 35 | 0.01 |
| | | Wing | 4 | 118.75 | 1.5 | 118 | 121 | 0.01 |
| | Female | Tail | 4 | 75 | 3.37 | 70 | 77 | 0.04 |
| Williamson et al. (present study) | | Bill | 64 | 36.5 | 2.51 | 30.91 | 41.74 | 0.07 |
| | | Wing | 65 | 122.77 | 7.01 | 112 | 139 | 0.06 |
| | | Tail | 63 | 79.17 | 4.19 | 71 | 90 | 0.05 |

**Table 3.** Morphological summary statistics for *peruviana* (Hellmayr 1932) and Northern Giant Hummingbird (Williamson et al., present study) males and females For each measured parameter, we provide sample size (*n*), mean, standard deviation, range (minimum and maximum values), and coefficient of variation.

| Dataset | Sex | Parameter (units) | *n* | Mean | Standard Deviation | Minimum | Maximum | Coefficient of Variation |
|---|---|---|---|---|---|---|---|---|
| Hellmayr 1932 | | Bill | 25 | 38.46 | 1.49 | 36 | 42 | 0.04 |
| | | Wing | 25 | 136.36 | 2.02 | 133 | 140 | 0.01 |
| | Male | Tail | 25 | 85.88 | 2.33 | 83 | 90 | 0.03 |
| Williamson et al. (present study) | | Bill | 102 | 38.71 | 1.65 | 34.76 | 42.03 | 0.04 |
| | | Wing | 102 | 131.76 | 6 | 111 | 147 | 0.05 |
| | | Tail | 99 | 81.42 | 4.38 | 66 | 91 | 0.05 |
| Hellmayr 1932 | | Bill | 8 | 39.31 | 1.03 | 38 | 41 | 0.03 |
| | | Wing | 8 | 130.88 | 3.31 | 126 | 137 | 0.03 |
| | Female | Tail | 8 | 84 | 3.25 | 78 | 87 | 0.04 |
| Williamson et al. (present study) | | Bill | 94 | 38.41 | 1.85 | 31.93 | 42.76 | 0.05 |
| | | Wing | 94 | 125.65 | 5.74 | 109 | 141 | 0.05 |
| | | Tail | 95 | 79.64 | 4.04 | 69 | 91 | 0.05 |

1 **Table 4.** Difference between taxon-mean trait values from Hellmayr (1932) (i.e., between
2 '*peruviana*' and *gigas*) and Williamson et al. (i.e., between the Northern Giant Hummingbird and
3 Southern Giant Hummingbird).

| Sex | Source | Difference between species means (mm) | | |
|---|---|---|---|---|
| | | **Bill length** | **Wing chord** | **Tail length** |
| Male | Hellmayr 1932 | 4.46 | 10.86 | 5.38 |
| | Williamson et al. (present study) | 2.44 | 4.12 | -0.45 |
| Female | Hellmayr 1932 | 5.06 | 12.12 | 9.0 |
| | Williamson et al. (present study) | 1.91 | 2.88 | 0.46 |