Life Cycle and Morphogenetic Differentiation in Heteromorphic Cell Types of a Cosmopolitan Marine Microalga

Laurie Bousquet[1,2], Shai Fainsod[1,2,°],Johan Decelle[3], Omer Murik[4], Fabien Chevalier[3] , Benoit Gallet[3], Rachel Templin[3], Yannick Schwab[3], Yoav Avrahami[1,2], Gil Koplovitz[1,2],Chuan Ku[5], Miguel J. Frada[1,2]

[1]Department of Ecology, Evolution and Behavior, Silberman Institute of Life Sciences, The Hebrew University of Jerusalem, Jerusalem, Israel; [2]The Interuniversity Institute for Marine Sciences in Eilat, Eilat, Israel; [3]Laboratoire Physiologie Cellulaire et Végétale, Univ. Grenoble Alpes, CNRS, CEA, INRAE, IRIG-DBSCI-LPCV, 38000 Grenoble, France; [4]Translation Genomics Lab and Medical Genetics Institute, Shaare Zedek Medical Center, 93722, Jerusalem, Israel; [5]Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan.

°Current address: Faculty of Biology, Technion – Israel Institute of Technology, Haifa 3200003, Israel.

Author for correspondence

Miguel J. Frada

Email: miguel.frada@mail.huji.ac.il

Summary

- *Gephyrocapsa huxleyi* is a prevalent, bloom-forming phytoplankton species in the oceans. It exhibits a complex haplo-diplontic life cycle, featuring a diploid-calcified phase, a haploid phase, and a third 'decoupled' phase produced during viral infection. Decoupled cells display a haploid-like phenotype, but are diploid.

- Here, we investigated the fate of decoupled cells during culture observations and we compared the transcriptome profiling and the cellular architecture in three dimensions of the three cell types.

- We found that decoupled cells can revert to the calcified form in the absence of viral pressure, revealing the transient nature of this cell type. Ultrastructural analyses showed distinct nuclear organisation with variations in chromatin volume. Transcriptomic analyses revealed gene expression patterns specific to each life phase. These included multiple regulatory functions in chromatin remodelling, broader epigenetic mechanisms and life cycling, which likely contributed to cell differentiation. Finally, the exploration of available host-virus transcriptomes supports life cycle transition during viral infection.

- This study provides cellular and molecular foundations for nuclear remodelling and cell differentiation in coccolithophores and the identification of gene markers for studying coccolithophore life cycles in natural populations.

Key words: Cell differentiation, Coccolithophore, Chromatin organization, Epigenetic regulation, Life cycle, Transcriptome, Viral infection.

Introduction

Eukaryotes exhibit complex life cycles comprising multiple phases interlinked by sexual mechanisms (syngamy and meiosis). They involve gene recombination, variations in ploidy levels (haploid,1n and diploid, 2n), as well as asexual mechanisms of cell differentiation. These transitions are crucial for the developmental and reproductive cycles of organisms, as well as for the adaptation to environmental changes through the differentiation of physiologically fit or resistant phenotypes (Mable & Otto, 1998; Von Dassow & Montresor, 2011).

Coccolithophores are a prevalent group of marine unicellular algae that produce distinct $CaCO_3$ plates, coccoliths, coating the cell surface (Monteiro *et al.*, 2016). Due to photosynthesis and calcification, these microorganisms play a pivotal role in the global carbon cycle (Balch, 2018). Coccolithophores have complex sexual haplodiplontic life cycles, composed of 1n- and calcified 2n-cell phases, each capable to independent cell division and expressing dissimilar morphology (heteromorphic). Such life cycles are proposed to widen niche-space and facilitate the adaptation to environmental variability (Houdan *et al.*, 2004; Frada *et al.*, 2019; de Vries *et al.*, 2021).

Gephyrocapsa huxleyi (formerly *Emiliania huxleyi*, (Bendif *et al.*, 2023)), is the most prevalent coccolithophore in the ocean and forms extensive seasonal blooms in high latitude regions, impacting marine ecology and biogeochemistry (Balch, 2018).The haplodiplontic, heteromorphic life cycle of *G. huxleyi* features diploid calcified cells (2n-calcified) that are non-motile and haploid non-calcified, biflagellated cells, coated by simpler organic scales (1n-flagellated) (Klaveness, 1972; Green *et al.*, 1996). Although 1n cells have been detected in 2n cultures, controlled manipulation of sexual transitions (syngamy or meiosis) has not yet been achieved. Transcriptome analyses have identified vast differences between diploid and haploid cells, particularly in transcripts linked to calcification, membrane trafficking and motility genes expressed in 2n-calcified cells and 1n-flagellated cells, respectively (von Dassow *et al.*, 2009; Rokitta *et al.*, 2011; Rokitta & Rost, 2012). However, despite extensive research on the ecological and biogeochemical roles of the 2n-calcified cells, the ecology of 1n-flagellated cells remains poorly understood (Frada *et al.*, 2012).

A major influence on *G. huxleyi* blooms are specific lytic viruses (*Emiliania huxleyi-*

specific viruses, EhV) that infect diploid cells and drive bloom decay (e.g. Bratbak *et*

al., 1993; Vardi *et al.*, 2012). Strikingly, culture experiments have demonstrated that

1n cells are resistant to EhV (Frada *et al.*, 2008). In addition, during infection of 2n-

calcified cells, cells displaying a haploid-like phenotype (biflagellated, organic scale-

bearing) and resistant to EhV are produced during viral infections (Frada et al. 2017).

These cells are however diploid, thus referred to as the 'decoupled ' cell type (2n-

flagellated) (Frada *et al.*, 2017), whose production resembles apospory reported in

plants and algae, where diploid gametophytes (typically haploid) develop without

meiosis (Coelho *et al.*, 2007). Such differentiation of 2n-flagellated cells may confer

an escape strategy from EhV, potentially setting the stage for post-bloom recovery

(Frada *et al.*, 2017).

Life cycle transitions typically involve layers of mechanisms, notably epigenetic

processes. These include chemical modifications of chromatin via the addition or

removal of chemical moieties on histone tails (histone post-translational

modifications, hPTMs) driving chromatin ultrastructural remodelling and gene

expression regulation (Lawrence *et al.*, 2016). Some hPTMs and associated proteins

are implicated in life cycle regulation. Notably, the Polycomb Repressive Complex 2

(PRC2) mediates the silencing of key three-amino acids loop extension (TALE)

homeodomain transcription factors (KNOX- and BELL-like) that in turn, regulate

haploid-to-diploid transitions in land plants (Sakakibara *et al.*, 2013; Horst *et al.*,

2016; Dierschke *et al.*, 2021) and in multiple algae (e.g., Lee *et al.*, 2008; Thangavel

& Nayar, 2018; Arun *et al.*, 2019; Mikami *et al.*, 2019; Hirooka *et al.*, 2022).

Specificities in these mechanisms are associated to different species. However,

alterations of these life cycle regulators can result in the decoupling of ploidy and life

phase generation (Schmidt, 2020), underlying either apospory (defined above) or

apogamy where instead haploid sporophytes (typically diploid) develop without

gamete fusion (e.g., Zhao *et al.*, 2001; Sakakibara *et al.*, 2013; Coelho *et al.*)

Here, we surveyed the fate of 2n-flagellated (decoupled) cells in culture of the

cosmopolitan microalga *G. huxleyi*. Moreover, we examined their relation to 2n-

calcified and 1n-flagellated forms, from morphological and gene expression

perspectives using 3D-microscopy (FIB-SEM, Focused ion beam scanning electron

microscopy) and transcriptome analyses. Finally, we also probed published *G.*

huxleyi-EhV transcriptomes to examine 2n-flagellated cell differentiation during viral

infection. Our results provide novel insights on the regulatory mechanisms driving life

cycle differentiation and response to viruses in *G. huxleyi*.

Material and Methods

Algal strains, maintenance and viral infections:

We used *Gephyrocapsa huxleyi* strains RCC1216 (2n-calcified, Houdan *et al.*, 2005),

LC5-11A (1n-flagellated), and LC5-12A (2n-flagellated) in this study. LC5-11A was

isolated from RCC1216 via single-cell pipetting following a spontaneous life cycle

change in culture, 6 months prior the study. LC5-12A and other 2n-flagellated strains

were isolated from RCC1216 during an EhV 201 infection (Schroeder *et al.*, 2002) 4

to 6 months prior the study, as described in Frada *et al.*, 2017. Cultures, maintained

in K/2 (-Tris, -Si) medium (Keller *et al.*, 1987), were not axenic. Genome sizes were

validated by flow cytometry of isolated nuclei (von Dassow *et al.*, 2009). Ten

additional decoupled strains were isolated from 3 independent RCC1216 infection

experiments with EhV 201. Over 2 years, routine inspection of decoupled cultures

was performed using optical microscopy. Sensitivity to EhV201 infection was

assessed during the exponential growth phase with a multiplicity of infection of 0.2.

Cell and viral concentrations were determined by flow cytometry (Attune NxT Life

Technology).

Algal growth for transcriptome analyses:

The 3 strains were grown in triplicate in 4L borosilicate vessels, under gentle air-

bubbling at 18°C, illuminated with 80 µmol photons $m^{-2}s^{-1}$ from warm-white LED

panels on a 14:10h light/dark cycle. Cultures were initiated at 5-8 x 10^4 cells mL^{-1}

and monitored for 48h by flow cytometry. The maximum photochemical quantum

yields of PSII (Fv/Fm), were measured by fluorescence induction and relaxation

system (Satlantic Inc., Halifax, NS, Canada; Gorbunov and Falkowski, 2005) (Table

S1). Four hours after dawn, 200 mL of each culture was harvested and pelleted by

centrifugation (8000 g, 10 min, 4°C). Cell pellets were flash-frozen in liquid N_2 and

stored at -80°C until RNA extraction.

RNA isolation, libraries preparation and sequencing:

RNA was extracted from cell pellets using the RNeasy Plant Kit (Qiagen) and treated

with DNase (Turbo DNA-free kit, Ambion). RNA concentration and integrity were

measured by Nanodrop (Thermo Fisher Scientific) and Tapestation (Agilent

Genomics). cDNA libraries were prepared according to the Illumina RNA sample

preparation kit. mRNA was isolated using poly(T) oligo-attached magnetic beads,

fragmented into 200-500 bp pieces and RNA fragments were reverse transcribed

with SuperScript II reverse transcriptase (Life Technology). Adapters ligation,

amplification and purifications preceded the sequencing of the cDNA libraries on an

Illumina NextSeq platform into 75-bp single-end reads.

Transcriptome curation, assembly and annotation:

FastQC (www.bioinformatics.babraham.ac.uk) and Trimmomatic (Bolger *et al.*, 2014)

were used for quality assessment and adapter trimming. Reads from the 9 *G. huxleyi*

libraries (triplicates for RCC1216, LC5-10A, LC5-12A) were aligned to the *G. huxleyi*

RCC1217 genome (Kao *et al.*, 2024), a haploid strain isolated from RCC1216.

Alignment was performed with STAR v2.7.10a (Dobin *et al.*, 2013). Transcripts were

reconstructed and merged into a synthetic life phase transcriptome using StringTie

v2.2.1 (Pertea *et al.*, 2015), Cuffmerge v2.2.1 (Trapnell *et al.*, 2012), and gffread

v0.12.7 (Pertea & Pertea, 2020), yielding 60930 transcripts. Transcriptome

completeness was evaluated with BUSCO v5.2.2 (Manni *et al.*, 2021) against the

eukaryota_odb10 database. Redundancy was reduced with CD-HIT-EST v4.8.1 (Fu

et al., 2012), resulting in 56092 non-redundant transcripts. Coding sequences were

identified with TransDecoder v5.5.0 . Functional annotation combined InterProScan

v5.52-86.0, Diamond blastP v2.0.14.152 against the nr database and EggNOG

mapper (Huerta-Cepas *et al.*, 2019; Blum *et al.*, 2021; Buchfink *et al.*, 2021). Results

were consolidated in OmicsBox, and Gene Ontology annotations were retrieved

using Blast2GO (Götz *et al.*, 2008). Parameters for each software are detailed in

Methods S1. Life phase transcriptome information is provided in Methods S2.

Histones and chromatin associated transcripts were examined using two

complementary strategies: (1) identification of close homologs through Blast

similarity searches against local databases, and (2) detection of conserved protein

domains with HMMER v3.3.2. Only shared hits were retained for subsequent

analyses. Databases were assembled from Thiriet-Rupert *et al.*, 2016; Grau-Bové *et*

al., 2022 and https://planttfdb.gao-lab.org. Details are provided in Methods S3.

Transcriptome Quantification, Differential Expression and Clustering:

RSEM v1.3.3 (Li & Dewey, 2011) with the parameter --bowtie2 was used for gene-

level abundance estimation. Features with fewer than 3 raw counts in at least 3

samples were filtered out. Transcripts per Million (TPM) normalization was applied,

and features with a mean expression > 0.01 TPM were considered expressed.

Differentially expressed transcripts were identified using DESeq2 (Love *et al.*, 2014),

with those showing a log2fold change > |2| and FDR < 0.05 clustered using K-

means. The optimal number of clusters was determined through complementary

statistical methods and visual inspection for homogeneity and separation.

Enrichment analysis was performed in R (v4.2.2; R Core Team 2021) using the

hypergeometric (HG) test. Null p-values were adjusted by adding an epsilon value

(smallest non-zero p-value divided by 1000) and applying the Benjamini-Hochberg

(BH) correction (Benjamini & Hochberg, 1995). Data analysis and visualization were

conducted in R, Unix, Python. Software parameter settings are detailed in Methods

S1. Validation of differential expression between cell types was performed by qRT-

PCR using various life phase-specific genes (Methods S4, Table S2).

Analyses of available *G. huxleyi* transcriptomes during viral infection:

Data from studies on the transcriptomic dynamics of *G. huxleyi* during viral infection

(Feldmesser *et al.*, 2014; Rosenwasser *et al.*, 2014) were used to examine the

expression patterns of specific life phase transcripts. *G. huxleyi* CCMP2090 was

infected with lytic viral strain EhV 201 at a multiplicity of infection of ~1:1. Samples

were collected at 1h and 24h post-infection, with non-infected cultures as controls,

as indicated in the original studies. In the present study, life phase transcripts were

blasted against CCMP2090 transcripts to identify homologs, retaining those with

100% identities over 80% alignment length. We focused on "flagellated-specific"

transcripts (expressed exclusively in flagellated forms - haploid and decoupled cells).

Abundance estimations from CCMP2090 were used to track the expression of these

transcripts over the infection timeline (see Table S3 for functional annotation and

clustering information of CCMP2090 homologs to flagellated transcripts).

Focused Ion Beam-Scanning Electron Microscopy (FIB-SEM): sample preparation and analysis:

G. huxleyi life stages were maintained under similar conditions as described above and harvested during exponential growth. Cells were centrifuged at 3000 g for 2 min, cryo-fixed using high-pressure freezing (HPM100, Leica), and freeze-substituted (EM ASF2, Leica) as in Decelle *et al.*, 2019; Uwizeye *et al.*, 2021 and Gallet *et al.*, 2024. Samples were embedded in 100% resin, mounted on SEM stubs, gold sputter-coated (Quorum Q150RS; 180 s at 30 mA), and imaged with a FIB-SEM (Crossbeam 540, Carl Zeiss). Atlas3D software (Fibics Inc. and Carl Zeiss) was used for sample preparation and 3D acquisitions, with SEM images recorded at an 8 nm pixel size. Segmentation of organelles (plastids, mitochondria, nucleus) and cellular compartments was performed using 3D Slicer software (Oyama *et al.*, 2014) using a manually-curated, semi-automatic pixel clustering mode (Decelle *et al.*, 2021; Uwizeye *et al.*, 2021). Colors were assigned to segmented regions, and image intensity thresholds were adjusted. Morphometric analyses were conducted with the "segmentStatistics" module in 3D Slicer, converting values to μm^3 or μm^2 based on an 8 nm voxel size. Five diploid-calcified cells, five haploid cells, six 2n-flagellated cells (non-calcified), and four reverted calcifying cells were analyzed. Further methodological details are provided in Methods S5.

Results

The life cycle of *G. huxleyi* includes three cell types representing distinct phases: a 2n-calcified (RCC1216), 1n-flagellated (LC5-10A) and 2n-flagellated (LC4-12A) cell type (Fig. 1a). Nuclear DNA content analyses by flow cytometry confirmed the ploidy of the three cell-types (Fig. S1a).The fate of decoupled cells and the morphogenetic differentiation between the three life cycle cell types was examined in our study.

Life cycle reversal: detection of calcified cells in 2n-flagellated cultures:

A total of eleven 2n-flagellated (decoupled) strains (LC4-12A-G and LC5-12A-D) were isolated during EhV -infections of RCC1216. Between 9 to 12 months of isolation, calcified cells were detected by microscopy in all 2n-flagellated cultures. As example, in LC4-12A, the fraction of calcified cells accounted for 21% of the total

cells as determined by flow cytometry (Fig. 1b, c). The new calcified cells are

referred to as reverted cells. Subsequent addition of the virus EhV 201 to all 2n-

flagellated cultures lead to the selective removal of reverted cells by lysis, but not 2n-

flagellated cells, and resulted in viral production (Fig. 1b, c and d). The decoupled

strain LC4-12A was randomly used in ensuing detailed 3D-microscopy and

transcriptome analyses to further examine the characteristics of the 2n-flagellated life

phase.

Figure 1 *Gephyrocapsa huxleyi* life cycle, reversion and response to viral infection. (a)

Schematic illustration of the life cycle of *G. huxleyi* and role of virus on life cycle transitions.

Sexual reproduction (meiosis and syngamy) drives the alternation between haploid,

uncalcified, biflagellated cells (1n-flagellated, grey) and diploid calcified cells (2n-calcified,

orange). During infection by specific *Emiliania huxleyi* viruses (EhV), 2n-calcified cells can

produce 2n-flagellated cells (red) that are phenotypically similar to 1n-flagellated cells. This

phenotype is reversible in absence of viral pressure (see results). (b) Microscopy and flow

cytometry depicting the emergence of calcified cells in 2n-decoupled cultures in the absence

of EhV. Left panel: light microscopy images of a 2n-flagellated (light grey arrow) and

reverted (dark grey arrow) cells, co-isolated in a clonal culture of 2n-flagellated cells in

absence of viral pressure. Central and right panels: Gating of uncalcified, bi-flagellated (light

grey ellipses) and calcified populations (reverted cells, dark grey ellipses) (chlorophyll

autofluorescence/side scatter) in a clonal culture of 2n-flagellated cells. (c) Removal of

reverted cells in 2n-flagellated cultures following EhV infections (120 h post-infection). The

fraction (%) of 2n-flagellated and reverted cells is indicated (n = 3). (d) EhV production in 2n-

flagellated cultures with reverted cells (n = 3). In c and d, non-infected controls were used for
comparison.

Cell and nuclear ultrastructural differences between cell types:

The cellular ultrastructure of 2n-calcified, 1n-flagellated, 2n-flagellated and reverted
calcifying cells were analysed by FIB-SEM. Cells from each life phase were
segmented and volumetrics of organelles (chloroplast, mitochondria and nuclear
compartments, such as electron-dense chromatin and nucleolus) were measured.
(Fig. 2a).

FIB-SEM results showed that cell volume was higher in 1n-flagellated and 2n-
flagellated (data not shown). Strikingly, reverted calcifying cells were 3-fold more
voluminous compared to 2n-calcified cells and coccolith formation was observed in
the vacuole in some cells (fig.2a). In 2n-calcified and 1n-flagellated cells,
chloroplasts occupied a similar cellular volume (35.8% ± 1.7% and 35.5% ± 3.4% of
the cell, respectively). Despite their higher cellular volume, chloroplasts in reverted
cells accounted for similar proportions (34.0% ± 2.9%). By contrast, in 2n-flagellated
cells chloroplast only represented 25.6% ± 4.1% of the cellular volume. Mitochondria
were largest in 1n-flagellated cells, occupying 7.5% ± 0.8% of the cell volume,
compared to about 5 % in other cell types (2n-flagellated, reverted and 2n-clacified
cells).

The nucleus was proportionally larger in 2n-calcified cells (14.6% ± 1.5% of cell
volume) compared to 11.9% ± 2.7% in 2n-flagellated cells, 10.3% ± 1.8% in reverted
cells, and 9.2% ± 1.4% in 1n cells (fig.2b). Within the nucleus, variations in electron
density patterns were detected, namely high and low density DNA (chromatin) and
nucleolus (Fig. 2 a,c). We therefore assessed the volume occupancy of these
nuclear compartments (% of the nuclear volume) (fig. 2c). The 2n-calcified and
reverted cells exhibited similar density patterns, with an average occupancy of ~49%
for high-density chromatin, ~47% of low-density chromatin and between 2.4% to 3%
for the nucleolus. In contrast, haploid cells (1n-flagellated) displayed an average
occupancy of 27% for high-density chromatin (two-fold less than 2n calcified cells),
nearly 68% for low-density chromatin and about 5% for nucleolus. Finally, 2n-
flagellated cells displayed an intermediate profile, with average occupancies of

nearly 40% for high-density chromatin, ~57% for low-density chromatin and 3% for the nucleolus.

Figure 2 Ultrastructural differentiation of the different life stages of *Gephyrocapsa huxleyi.* (a) 3D reconstitution with FIB-SEM (focused ion beam scanning electron microscopy) of 2n-calcified, 1n-flagellated, 2n-flagellated (decoupled) and reverted (calcifying) cells. Chloroplasts, mitochondria, nuclear areas and coccoliths are painted in different colors. The scale bar represents 2µm. 1n and 2n: haploid and diploid respectively (b) Relative proportions (% of cell volume) of cytoplasm, chloroplast, mitochondria and nucleus in the four cell types, were calculated from the 3D reconstructions. Significant differences ($p < 0.05$) identified by Tukey HSD tests are indicated with different letters above the bars (a,b, ab).(c) Relative proportions (% of nuclear occupancy) of high-density and low-density areas of the nucleus and the nucleolus in the four cell types. Significant differences

(p < 0.05) identified by Tukey HSD tests are indicated with different letters above the bars

(a,b, ab).Statistical analysis was performed using ANOVA type II and Tukey HSD tests to

identify significant differences across conditions for each organelle type. Measurements

were based on five 2n-calcified cells, five 1n-flagellated cells, six 2n-flagellated cells, and

four reverted cells.

Cell growth parameters and global transcriptome features:

Gene diversity and expression differences between life phases were examined by

transcriptome analyses. We note that transcriptome analyses were undertaken prior

to the production of reverted cells in LC4-12A, thus solely 2n-calcified, 1n-flagellated

and 2n-flagellated cells are included. Inspection by light microscopy validated that

each culture contained only one cell type. Cells were grown in replete medium and

harvested during exponential growth (Fig. S1b and Table S1). Variations in growth

rates were detected one day prior to sampling for the transcriptomes. The 2n-

calcified cells displayed higher growth rates (1.33 ± 0.04 day^{-1}), followed by 1n-

flagellated cells (0.72 ± 0.02 day^{-1}) and then 2n-flagellated cell with lower growth rate

dynamics (0.47 ± 0.04 day^{-1}). All cultures maintained high photosynthetic quantum

yield of photosystem II (Fv/Fm) throughout cell growth. At the time of biomass

harvesting Fv/Fm equalled ~0.80 in all strains (Fig. S1c).

To evaluate transcriptional profiles, we defined as transcriptome richness the ratio of

expressed transcripts to the total number of transcripts within this 'synthetic life

phase transcriptome'. It is important to note that we set a conservative expression

threshold (> 0.01 TPM) to define a transcript as expressed or not. Therefore, we do

not rule out the presence, in this core of transcripts, of sequences presenting only

residual expression levels. Transcripts with less than 3 raw counts across the 9

libraries were filtered-out, yielding 27350 expressed transcripts. Among these, an

average of 23307 (85.2%) were expressed in the 2n-calcified cells, 25263 (92.4%) in

the 1n-flagellated cells and 25543 (93.4%) in the 2n-flagellated ones (Fig. 3a),

aligning with the observed nuclear patterns. Irrespective of expression magnitude or

category-specific differential expression, >78% (21434) of gene-transcripts were

common to all three life phases (Fig. 3b). A larger overlap was detected between

flagellated cell types (1n-flagellated and 2n-flagellated,11.6%, 3176) than between

either of the diploid (2n-calcified and 2n-flagellated ;2.7%, 750) or the non-decoupled

stages (1n-flagellated and 2n-calcified, 1.3%, 344). Only a small subset of transcripts was unique to each phase: 1.6% (431) in 1n-flagellated, 3.3% (910) in 2n-calcified cells, and only 1.2% (316) in the 2n-flagellated cells. Independent analyses by RT-qPCR on a set of 19 gene-transcripts validated the specify of expression of the targeted transcripts (Table S2).

Principal Component Analysis (PCA) delineated the variance between the three life phases (Fig. 3c), with PC1 accounting for 85% of the variation, separating flagellated (haploid and decoupled) from 2n-calcified cells. PC2 encompassed 11.7% of the variation, with 1n-flagellated and 2n-flagellated cells well separated. The top 50 transcripts predominantly influencing PC1 were mostly associated to a morphotype: 34 unique to flagellated forms, two to calcified, nine to diploid, and two commonly detected across all conditions, albeit with expression differing by more than tenfold between flagellated and calcified states (see Table S4). Among these, 34 transcripts annotated with KEGG, KOG or GO terms, were linked to energy metabolism (including nitrogen), amino acid and inorganic metabolism, as well as genetic information processing (transcription machinery, chaperone and folding catalysts, chromosomal proteins, and the ubiquitin system). In contrast, the transcripts central to PC2 demonstrated different specificities: two associated with flagellated morphotypes, 19 with diploid-specific expressions, 13 with decoupled cells, and ten as core transcripts with significant differential gene expression (DGE) across conditions. Additionally, two transcripts were exclusive to the haploid phase, and four were associated with non-decoupled phases (Table S4). These transcripts were related to genetic information processing, including transcription, RNA processing and modification, replication, recombination, and post-transcriptional modifications, as well as lipid biosynthesis and photosynthesis.

Figure 3 Global properties and differentiation of the transcriptomes of life cycle

phases of *Gephyrocapsa huxleyi*. (a) Transcriptome richness of the 2n-calcified, 2n-

flagellated and 1n-flagellated. Richness was defined as the ratio of expressed transcripts to

the total number of transcripts. A transcript was considered expressed if > 0.01 TPM. (b)

Venn diagram of transcript specificity across life phases, in the 1n-flagellated, 2n-calcified

and 2n-flagellated. Number of transcripts in each category, and corresponding relative

proportions (%) are provided. (c) Principal Component Analysis (PCA) of transcriptional

expression profiles and relative separation between different life phases along first 2

principal components (PCs). Each point represents a sample, colored according to its life

phase. Arrows represent the contribution of individual transcripts to the PCA, with the color

gradient indicating the magnitude of their contribution (blue: low contribution, purple: high

contribution). x and y axes: standardized values of the PCs (arbitrary units).

Transcriptional differentiation and functional enrichments:

We identified transcriptional differences between 2n-calcified, 1n-flagellated and 2n-
flagellated cells. Abundance estimations were performed at the gene level and
23048 transcripts (84.3%) were statistically differentially expressed (False Discovery
Rate (FDR) < 0.05). Of these, 9114 transcripts exhibited at least a fourfold change in
expression in one pairwise comparison (log2 fold-change) (Table S5). Cluster
analysis with k-means resulted in 10 clusters with unique patterns of expression (Fig.
4, Table S6) examined for functional enrichments (GO, KEGG and COG). In order to
reduce terms redundancy of the presented Biological Processes (GO- BP, Fig. 4),
significantly enriched terms were grouped under their direct GO ancestor term. Refer
Table S7 for complete results concerning GO-molecular functions, KEGG and COG.
Below are described the enrichments identified in each cluster across all annotation
systems.

Clusters 1 and 5, accounted for over half (52%) of the differentially expressed genes
and contained the most influential transcripts in the PCA displayed in Fig. 3c (and
Supporting Information Tables S4, 6). These clusters, exhibited expression patterns
specific to morphotypes (calcified vs non-calcified, flagellated cell types). Cluster 1
(3097 transcripts) was overexpressed in flagellated forms (1n and 2n), featuring
enrichments in cytoskeleton and flagellar processes (cilium or flagellum dependent
cell motility, microtubule and cilium formation, microtubule-based transport, dynein
and catenin binding), as well as metabolism and transport pathways such as
carbohydrates (glycosaminoglycan, fucose), amino-acids and nucleotides and DNA
mediated transposition and DMSP-dethiomethylase activity (Table S7). By contrast,
cluster 5 (1644 transcripts) with transcripts overexpressed in calcified cells, featured
enrichments for transport and metabolism of inorganic ions (NO_3^-, NO_2^-), as well as
RNA transcription, processing and modification and in cell cycle control and
chromosome partitioning (Table S7). Necrotic cell death, regulation of reactive
oxygen species as well as sialic acid transferase were also associated to cluster 5.

Clusters 4 (900 transcripts) and 10 (823 transcripts) represented transcripts
overexpressed in flagellated cells (1n and 2n). Cluster 4 was associated with
coenzyme and nucleotide transport and metabolism (Supporting Information Table
8). Cluster 10 (with higher expression in 2n-flagellated than in 1n-flagellated cells)

was enriched in cytoskeleton and cellular architecture organization (actin filament, calmodulin binding), in lipid and secondary metabolism associated to energy production and conversion (glucose metabolism), calcium transmembrane transport as well as RNA processing, modifications and chromatin structure and dynamic (Table S7).

Clusters 2 (569 transcripts) and 8 (414 transcripts) comprised transcripts overexpressed in 2n-calcified and 2n-flagellated. Cluster 2, was associated with eukaryotic clusters of orthologous proteins (COGs) related to transcription, processing and modification of RNA and signal transduction mechanisms (Table S7). Cluster 8 (414 transcripts), with more elevated expression in the 2n-calcified phase, was enriched in the DNA-related functions replication, recombination and repair (DNA replication, post replication repair), as well as control of the cell cycle, cell division and chromosome partitioning (including control of mitotic cell cycle, nuclear chromosome segregation and sister chromatin cohesion). Processes related to the structure of the chromatin and its dynamic were also enriched in cluster 8 (chromatin binding, DNA conformation change, chromosome condensation) as well as mating type determination, switching and reproductive process in single-cell organism) and response to various stimuli (UV, carbohydrates).

Clusters 6 (337 transcripts) and 9 (481 transcripts), represented non-decoupled cell transcripts, overexpressed in both 1n-flagellated and 2n-calcified cells. Cluster 6 was associated with carbohydrate biosynthesis and organic cyclic compound binding. It was additionally associated to intracellular trafficking and DNA replication, recombination and repair and post-translational modifications (Table S7). Cluster 9 (more prevalent in 2n-calcified cells), displayed enrichments in carbohydrate metabolism, polar nucleus fusion and cinnamyl and coniferyl-alcohol dehydrogenases activity (Table S7).

Cluster 3 (353 transcripts) was the only group specific to 2n-flagellated (decoupled) cells. It was associated to transcription, replication, recombination, repair and chromatin structure (Table S7), as well as phosphatidyl inositol binding, response and regulation to calcium ion and aluminum.

Finally, cluster 7 (496 transcripts) was haploid specific and included enrichments in photosynthesis-related functions (chlorophyll and pigment binding) and processes,

as well as signal transduction, energy production and conversion, carbohydrate and

lipid metabolism and intracellular trafficking and vesicular transport.

Figure 4 Expression profiles of transcripts in the life phases of *Gephyrocapsa huxleyi*.
(a) K-mean (10) clustering of transcripts with significant variations of expression (log2 fold change > 2) between life phases. Expression is standardized by transcript (z-scale). Dark green, high expression levels; brown, low expression levels. b. Most significantly enriched GO terms (Biological Processes), (hypergeometric test, adjusted p value < 0.05) in the clusters presented in a. Terms were grouped under parental term to reduce redundancy and facilitate visualization. For a complete list of enriched GO terms in each cluster, see Supporting information table S8.

Expression of histone, chromatin associated transcripts and transcription factors:

FIB-SEM observations, highlighting marked differences in nuclear ultrastructure

between life cycle phases, stimulated the bioinformatic examination of genetic

functions potentially involved in chromatin remodeling and likely implicated in life

cycling. Alongside, several functional enrichments, particularly those related to

general DNA replication and repair, chromosome conformation and regulation of

gene expression, underscored the potential role of epigenetic regulation in

differentiating life phases. Thus, we zoomed on the transcripts associated with the

general chromatin machinery: histones, readers, writers, remodelers, chaperones, as

well as transcription factors (TFs) (Fig. 5).

Ten clusters were defined, each with highly heterogeneous gene family composition.

Clusters 1 and 2 contained more than a third of the transcripts, segregating cells by

their morphotypes. Cluster 1 (146 transcripts), primarily expressed in flagellated

cells, was enriched in TFs and protein domains commonly associated with the

chromatin machinery ('common'), though no specific family was identified. Cluster 2

(111 transcripts), highly expressed in calcified cells, was enriched in chromatin

specificmethyltransferases, TFs, and chromatin readers.

The other clusters were considerably smaller and associated to specific life phases.

Cluster 9 (69 transcripts), expressed in both flagellated conditions, was more

abundant in the 2n-decoupled cells compared to the 1n-calcified cells. This cluster

was associated with TFs, proteins involved in deubiquitination processes, and

'common' domains. Clusters 4, 5, and 10 (54, 43, and 47 transcripts, respectively)

were expressed in diploid cells. Cluster 4, characterized by clear heterogeneity

among its replicates and higher expression in diploids, was enriched in readers,

chaperones, and remodelers. Clusters 5 and 10, more abundant in the 2n-calcified

and 2n-flagellated cells respectively, had no significant enrichments. Cluster 3 (53

transcripts) was the only group characterized by an overexpression exclusive of the

1n-flagellated cells and was associated with TFs and deubiquitination proteins.

Cluster 8 (47 transcripts), the only group overexpressed in the decoupled cells, was

not associated with any significant enrichments. Finally, clusters 6 and 7 (35 and 53

transcripts, respectively) were more expressed in non-decoupled cells. Cluster 6,

with higher expression in 1n-flagellated cells, was not associated with significant

enrichments. Cluster 7, with transcripts more expressed in the 2n-calcified than in

the 1n-flagellated cells, was associated with the Polycomb protein complex.

Twenty-two transcripts were associated to canonical histones or their variants in our

dataset: 10 histones H2A, 2 H2B, 11 histones H3, 1 H4, 1 transcript had homologies

with the linker histone H1 and 3 transcripts were associated to the histone family

without a clear identification (Table S5). 5 histones H2A were highly expressed in all

cells and 2 of them were strongly depleted in the 2n-flagellated and 1n-flagellated

respectively. One H2B was over 50% more expressed in 2n-calcified cells. Four

histones H3 were abundant and strongly overexpressed in one life phase. One was

expressed only in the haploid cells and 1 was expressed in the diploid phases and

more abundant in the 2n-flagellated. Two other histones H3 were overexpressed in

the 2n-calicified cells compared to the 2 other conditions. (Table S5). Finally, histone

H1 was also more detected in 2n cells compared to the1n cells (Table S5).

One TALE-HD TF was identified and expressed only in the haploid cells This

specificity was validated by RT-qPCR (Table S2).

Figure 5 Expression profiles of chromatin associated transcripts related to epigenetic

functions in the life phases of *Gephyrocapsa huxleyi*. (a) K-mean (10) clustering of

transcripts associated to statistically significant variations of expression (FDR < 0.05,

log2fold change >|0.5|) between life phases. Expression is standardized by transcript (z-

scale). Dark green, high expression levels; brown, low expression levels. (b) Significantly

enriched epigenetic families, (hypergeometric test, adjusted p value < 0.05) in clusters

presented in (a). "common": transcripts for which only one protein domain, commonly found

associated to chromatin-related proteins and processes, could be identified. In the lack of

other identifications, transcripts family could not be determined (conserved domains: Dpy-30,

Kelch, WD-40, zf-C2H2 and zf-CXXC).

Flagellated cell-specific expression during viral infection:

Earlier work showed that decoupled cells are produced during Ehv infection (Frada

et al. 2017). To test life cycle transition during viral infection and gain insight on

mechanisms of 2n-flagellated production, we re-examined published transcriptomes

of the diploid *G. huxleyi* strain CCMP2090 interacting with EhV201 (1 and 24 hr post

infection) (Rosenwasser *et al.*, 2014). We targeted transcripts that demonstrated a

high homology level (see methods) with flagellated-specific transcripts identified in

our work.

We identified 176 gene homologs in this dataset. Through k-means clustering, four

co-expressed groups were identified and functional enrichments were examined

(Fig. 6a; see Table S3 for complete list of clustered transcripts, associated

annotations and Table S7 for enrichments). Clusters 1 and 2 grouped transcripts

with higher expression levels in control (non-infected) conditions. Cluster 1 (13

transcripts) was found to be enriched in organelle organisation process and to

general metabolism annotations, as well as with protein families involved in genetic

information processing (mRNA biogenesis, chaperones and folding catalysts).

Cluster 2 (22 transcripts) was enriched in general metabolism pathways

(carbohydrate, sugar, lipid). Its transcripts were also related to proteins involved in

genetic information processing (transcription and translation machinery, chromatin

structure). Cluster 3 (22 transcripts) comprised transcripts that were upregulated at

1h post-infection but downregulated by 24h. This cluster was enriched in protein

modification process, and its transcripts associated to the metabolism of amino

acids, the ubiquitin system and cilium and its associated proteins. Cluster 4,

containing 68% (119 transcripts) of the total clustered transcripts was markedly

upregulated 24h post-infection. It was enriched in orthologous groups and functions

associated to the cell cycle control and division and partitioning of chromosomes, to

signal transduction and cytoskeleton mechanisms (tubulin binding, microtubule

cytoskeleton organisation), as well as post-translational modifications, chaperones

and protein turnover (histone methyltransferase activity, nucleotide/nucleoside

biosynthetic processes and binding) and intracellular trafficking and carbohydrate

transport and metabolism.

Figure 6 Expression profiles of "flagellated-specific" genes in the *Gephyrocapsa*

***huxleyi* strain CCMP2090 during viral infection by EhV201 and non-infected control.**

(a). K-means clustering of homologs in CCMP2090 to flagellated transcripts in absence

(control, ctl) or presence (+EhV) of viral infection, 1h and 24h post infection. Expression is

standardized by transcript (z-scale). Dark pink; high expression level, Blue, low expression

level. (b) Significantly enriched GO terms (1left panel -molecular functions, right panel-

biological processes), (hypergeometric test, adjusted p value < 0.05) in cluster presented in

(a). abbreviations: microtub.: microtubule, dvt.: development, sign.: signal, bioS.:

biosynthetic, proc.: processes, loc.: localization, mbne.: membrane, proj.: projection, ass.:

assembly, transp.: transport, cytosk.: cytoskeleton, org.: organization, intracell.: intracellular,

modif.: modification

Discussion

The life cycle, encompassing all cellular stages, is a fundamental attribute of

eukaryotes (Mable & Otto, 1998), underpinning their range of biological and

ecological functions, and their capacity to adapt to environmental changes through

cell differentiation. In marine phytoplankton, the understanding of life cycle versatility

and regulation remains limited. Here, we examined the life cycle of the cosmopolitan,

bloom-forming coccolithophore *G. huxleyi*, which includes 2n-calcified cells, 1n-

flagellated cells and 2n-flagellated cells produced during viral infections (Frada *et al.*,

2017). We specifically investigated the fate of the virus-resistant 2n-flagellated cells

during culture observations and the morphogenetic differentiation between 2n-

calcified, 1n-flagellated and 2n-flagellated cell types through ultrastructural and

transcriptome analyses.

A key finding of this study is the ability of 2n-flagellated cells to revert back to a

calcifying morphotype in the absence of viral pressure. Given the low proportion of

newly calcifying cells (reverted) within the 2n-flagellated cultures (20%), it is

reasonable to assume that only a small subset of cells undergoes flagellate-to-

calcified switch. Reverted cells subsequently proliferate mitotically, alongside

flagellated cells, becoming detectable by microscopy perhaps only weeks post-

transition.

Morphological transformations into a resistant phenotype in response to viral

infections, followed by a reversal to the original cell phenotype, have been observed

in other protists. For example, certain diatom and raphidophyte species can produce

viral-resistant cysts during infections, which germinate back to the vegetative cell-

type in the absence of viruses (Dingman, Jennifer, 2015; Pelusi *et al.*, 2021).

However, the role of viral infection in triggering life cycle transitions and the

subsequent ability to reverse this process, has been to our knowledge uniquely

described here, in *G. huxleyi*. The production of decoupled cells resembling

apospory observed in algae and plants (e.g.,(Cock *et al.*, 2014) is thought to

represent a natural escape strategy from EhV infection (Frada *et al.*, 2008, 2017).

However, the reversal to a calcifying state suggests a potential cost for diploid cells

in maintaining a haploid-like phenotype. The cost of viral resistance is a common

phenomenon in bacteria and protists (e.g, Lenski & Levin, 1985; Tomaru *et al.*, 2009;

Thomas *et al.*, 2011; Dingman, Jennifer, 2015). In *G. huxleyi*, as corroborated by a

previous study (Frada *et al.*, 2017), 2n-flagellated (decoupled) cells display reduced

growth rate and carrying capacity compared to both 2n-calcified and 1n-flagellated

(haploid) cells. This may be explained by their inherently larger cell volume, revealed

here with electron microscopy, and ensuing higher nutrient requirements for cell

division. In addition, FIB-SEM measurements indicate a relative proportion of

chloroplasts ~25% lower in the 2n-flagellated cells compared to other cell types,

suggesting a reduced capacity to generate energy, while the mitochondrial volume in

these cells, does not change. Thus, the phenotype reversion (2n-flagellated to 2n-

calcified) likely restores the metabolic and physiological advantages.

We also observed that the reversion of 2n-flagellated cell in *G. huxleyi*, is associated

with a loss of the viral resistance. Calcified reverted cells, once re-exposed to EhV,

were eliminated from the cultures. Mechanisms underlying *G. huxleyi*'s sensitivity

and resistance to EhV may involve a variety of metabolic processes, reflecting the

complex host-virus 'arms-race' dynamics (Bidle & Vardi, 2011; Bidle, 2015). Previous

work linked specific sphingolipid molecules (sialic acid sphingolipids, sGSL) to EhV

sensitivity in diploid cells (Fulton *et al.*, 2014). sGSLs were proposed to play an

intrinsic role in coccolith biosynthesis and serve as ligands for viral attachment and

entry. These sGSLs are not biochemically detected in haploid cells, which are

insensitive to EhV (Hunter *et al.*, 2015). Enrichment for sialylation transcripts (cluster

5, Fig. 4) was observed only in diploid-calcified cells and not in flagellated cell types

(Table S7). This suggests that sGSLs are likely absent in 2n-flagellated cells but may

reappear in reverted cells, potentially restoring viral sensitivity.

Ultrastructural analyses enabled the quantitative assessment of significant

differences in the nuclear architecture between cell-types. Notably, nuclei of 2n-

calcified and reverted cells displayed nearly equal parts of low- and high-density

chromatin, accounting for 47-49% of the nuclear volume, respectively. In contrast,

nuclei of 1n-flagellated were predominantly of low-density chromatin (> 67%), while

the nuclei of 2n-flagellated cells displayed an intermediate phenotype (Fig. 2c). Specific approaches are needed to confirm the nature of high- and low-density areas ('Chromatin accessibility profiling methods', 2021). However, it is reasonable to assume that they correspond to heterochromatin (densely packed and usually transcriptionally inactive), and euchromatin (relaxed and generally associated with a more active transcriptional state of the DNA), respectively (Huisinga *et al.*, 2006). In eukaryotes, chromatin packaging is highly dynamic and central for spatiotemporal regulation of transcription in many cellular processes, including cell differentiation, development and life cycle in animals, plants and algae (Zheng & Xie, 2019; Vigneau & Borg, 2021). Rearrangements in chromatin ultrastructure is often mediated by extensive epigenetic reprogramming, involving hPTMs and the activity of TFs that modulate transcription landscapes (Morgan *et al.*, 2005; Thiriet-Rupert *et al.*, 2016; Vigneau & Borg, 2021). Thus, the observed nuclear ultrastructural patterns across the different phases of the life cycle in *G. huxleyi* through unprecedented volume quantification, suggest vast divergence of gene expression between cell types, with 2n-flagellated cells maybe displaying an intermediary transcriptional profile.

Complementary comparative transcriptome analyses enabled to test this hypothesis, while gaining insights on wider functional attributes of 2n-calcified, 1n-flagellated and 2n-flagellated cells during exponential growth. Conservative identification of common and phase specific transcripts highlighted that the strong majority of transcripts were shared between the three life phases (>78%), regardless of expression levels (Fig. 3b). A more important overlap was observed between flagellated morphotypes (11.6%) compared to the diploid cells (2.7%) (Fig. 3b). Moreover, as revealed by the PCA (Fig.3c), the main factor of separation between samples (>85% of the variance) was attributed to the cell morphology (flagellated vs calcified), while only 11.7% of the variance equally separated 1n and 2n-flagellated from 2n-calcified(Fig. 3c). This separation might lay in the mechanisms of production of these cells, haploid cells being produced by meiosis while decoupled cells would originate from apospory like mechanisms. Thus, from a transcriptomic perspective, 2n-flagellate cells do not represent an intermediary level of expression but are in fact much closer to the haploid profile. Regulation of cell morphology, rather than the ploidy level itself, is the main factor of differentiation between cell types in *G. huxleyi,* as similarly detected in brown algae, displaying a haplodiplontic life cycle and apomictic variants (Arun *et al.*,

2019). Examination of DGE and enrichment profiles further highlighted the contrasts

between flagellated and 2n-calcified cells. Namely, enrichments in functions related

to cell motility were identified in flagellated cell types. By contrast, enrichments in

functions related to the transport and metabolism of inorganic ions were associated

in 2n-calcified cells. This can be associated with flagella biosynthesis, signalling, and

with calcification, respectively, in broad agreement with previous transcriptome

comparisons of 1n-flagellated and 2n-calcified cells (von Dassow *et al.*, 2009; Rokitta

et al., 2011).

Significant differences between cell types were also detected in functions associated

to chromatin remodelling and transcriptional regulation (including transcription

factors, TFs). At this level, morphotype is still the main differentiating factor of

transcript expression (Fig.5). A majority of transcripts were either over-expressed in

the calcified cells or in the flagellated cells, further emphasizing phenotype as the

main manifestation of gene regulation in *G. huxleyi* life phases, independently of

ploidy levels. We detected for example an over expression in the 2n-calcified cells

associated to histones methyltransferases (Fig. 5a, cluster 2), which can align with

the high chromatin compaction (Chen & Dent, 2014) detected in these cells (Fig. 2b).

Additionally, ubiquitin removal (deubiquitination) enrichments were associated to

transcripts with clear DE between cell types, a function associated with chromatin

relaxation and increased transcriptional activity (Mikulski *et al.*, 2017). It fits overall

patterns of chromatin, less dense in flagellated cells compared to 2n-calcified cells

(Fig. 5a, clusters 3 and 9). Enrichments observed in cluster 4 (Fig. 5a) were also

relevant. We observed an heterogeneous gradient of expression from 2n-calcified to

2n-flagellated cells, in transcripts associated to chromatin "architects" (remodelers,

readers, and chaperones) (Clapier & Cairns, 2009). suggesting intermediate level of

expression and recruitment. We think it might further be associated to a transitional

level of chromatin organization between the two types of diploid cells. We also

identified enrichments for the Polycomb-Complex, associated with transcripts over-

expressed in 2n-calcified cells compared to 1n-flagellated cells, a TALE-

homeodomain TF (KNOX) specific to 1n cells and histone H3 variants with marked

DGE between life phases. All these gene-functions are conserved across eukaryotes

and play central roles in the regulation of life cycle transitions (Horst *et al.*, 2016;

Mikami *et al.*, 2019; Dierschke *et al.*, 2021; Lee *et al.*). Polycomb associated

transcripts in our dataset, likely correspond to the Polycomb Repressive Complex 2

(PRC2,(Grau-Bové *et al.*, 2022)). PRC2 mediates the trimethylation of histones 3

(H3) on lysine 27 (H3K27me3), associated to the silencing of genes encoding TALE-

HD TF (KNOX and BELL). These transcription factors are crucial for haploid-to-

diploid transitions in plants and algae (Yelagandula *et al.*, 2014; Wollmann *et al.*,

2017; Thangavel & Nayar, 2018; Loppin & Berger, 2020; Vigneau & Borg, 2021;

Hirooka *et al.*, 2022) (Loppin & Berger 2020, Wollman et al. 2017, Yelagandula et al.

2014, Vigneau and Borg, 2021). Overall, these results suggest the involvement of

conserved key mechanisms in the life cycle differentiation in *G. huxleyi*.

Finally, we screened for homologs of flagellated-specifc transcripts (expressed only

in the 1n- and 2n-flagelled cells) in the *G. huxleyi* strain CCMP2090 during EhV

infection (Rosenwasser *et al.*, 2014). A variety of transcripts mainly involved in

metabolic functions were detected in control (non-infected) conditions, indicating that

life cycle specificity of various functional genes can change between *G. huxleyi*

strains, and likely as well between growth conditions and external environmental

stimuli. However, most of the flagellated-cell homologs identified were over-

expressed 24h post-infection, strongly supporting life cycle transition during EhV

infections (Frada *et al.*, 2017). Among flagellated-cell homologs, we detected a wide

set of motility-related genes and cell architecture, as well as signaling transduction

mechanisms. We note that CCMP2090 does not form flagella, albeit a strong up-

regulation of some flagellar functions upon infection (Frada *et al.*, 2017) . This

morphological impairment likely results from the genome erosion of this strain and

subsequent loss of functional flagella biosynthesis genes(von Dassow *et al.*, 2015).

In any case, CCMP2090 can form viral resistant cells that display organic-body

scales, only featured in flagellated cells (Frada *et al.*, 2017). We further detected the

association of these flagellated specific transcripts with functions involved in gene

expression, including a strong enrichment in H3K79 specific methyltransferase that

regulates diverse cellular processes, such as development, reprogramming,

differentiation, and proliferation (Feng *et al.*, 2010; Takahashi *et al.*, 2011; Tatum &

Li, 2011). It is generally associated to active transcription and maintains chromatin

accessibility for histone acetylation and transcription factor binding (Godfrey *et al.*,

2019). Although limited, these lines of evidence further recall the involvement of the

upstream epigenetic regulation during the differentiation of 2n-flagellated cells in *G. huxleyi* during viral infection.

In conclusion, our study provides detailed comparative assessment of three life-cycle cell types in *G. huxleyi*, evidence for life cycle versatility during viral interactions and novel insight on chromatin remodeling and transcription regulators, which likely play a central role in mediating life cycle transitions. These results should provide the basis for further development of future studies on role epigenetic control during cell differentiation and life cycle in coccolithophores. Finally, one of the greatest gaps limiting comprehensive understanding about *G. huxleyi* is the lack of knowledge of the occurrence, distribution, and role of flagellated cells in natural populations and during blooms where viral infections are recurrent. The analyses presented in this study enable to refine useful gene markers for the three life phases of *G. huxleyi* that can be used to interrogate coccolithophore assemblages in the oceans.

Acknowledgements

This study was supported by the Israel Science Foundation (2921/20) attributed to MJF. JD was supported by CNRS and ATIP-Avenir program funding. We thank Prof. Aharon Kaplan for hosting SF and Prof. Assaf Vardi to enable access to his lab for molecular biology work. We thank The Life Sciences Core Facilities from The Weizmann Institute of Science for the preparation and transcriptome sequencing as well as Esther Feldmesser for support and feedback on bioinformatic analyses. We thank Guy Schoehn and Christine Moriscot, and the electron microscope facility at IBS, which is supported by the Rhône-Alpes Region, the Fondation Recherche Medicale (FRM), the fonds FEDER, the Center National de la Recherche Scientifique (CNRS), the CEA, the University of Grenoble, EMBL, and the GIS Infrastructures en Biologie Sante et Agronomie (IBISA). CK was supported by Academia Sinica (AS-CDA-110-L01) and NSTC, Taiwan (111-2611-M001-008-MY3).

Author contributions

Conceptualization: MJF, LB, SF; Methodology: MJF, JD, LB, SF, OM; Investigation: LB, SF, YA,GK, FC, BG, RT, YS, MJF; Formal Analysis: LB, YD, SF; Resources:

CK, FC, MJF; Data curation: LB, JD, BG, RT, YS; Visualization: LB, JD, RT, YS;

Writing (original draft): LB, MJF; Writing (review & editing): LB, MJF, SF, JD and CK;

Supervision: MJF, OM; Funding acquisition: MJF, CK.

Competing interests:

All authors declare they have no competing interests.

References

Arun A, Coelho SM, Peters AF, Bourdareau S, Pérès L, Scornet D, Strittmatter M, Lipinska AP, Yao H, Godfroy O, *et al.* 2019. Convergent recruitment of TALE homeodomain life cycle regulators to direct sporophyte development in land plants and brown algae (S McCormick and CS Hardtke, Eds.). *eLife* **8**: e43101.

Balch WM. 2018. The Ecology, Biogeochemistry, and Optical Properties of Coccolithophores. *Annual Review of Marine Science* **10**: 71–98.

Bendif EM, Probert I, Archontikis OA, Young JR, Beaufort L, Rickaby RE, Filatov D. 2023. Rapid diversification underlying the global dominance of a cosmopolitan phytoplankton. *The ISME Journal* **17**: 630–640.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**: 289–300.

Bidle KD. 2015. The Molecular Ecophysiology of Programmed Cell Death in Marine Phytoplankton. *Annual Review of Marine Science* **7**: 341–375.

Bidle KD, Vardi A. 2011. A chemical arms race at sea mediates algal host–virus interactions. *Current Opinion in Microbiology* **14**: 449–457.

Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, Nuka G, Paysan-Lafosse T, Qureshi M, Raj S, *et al.* 2021. The InterPro protein families and domains database: 20 years on. *Nucleic Acids Research* **49**: D344–D354.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Bratbak G, Egge JK, Heldal M. 1993. Viral mortality of the marine alga *Emiliania huxleyi* (Haptophyceae) and termination of algal blooms. *Marine Ecology Progress Series* **93**: 39–48.

Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nature Methods* **18**: 366–368.

Chen T, Dent SYR. 2014. Chromatin modifiers and remodellers: regulators of cellular differentiation. *Nature Reviews Genetics* **15**: 93–106.

Chromatin accessibility profiling methods. 2021. *Nature Reviews Methods Primers* **1**: 11.

Clapier CR, Cairns BR. 2009. The Biology of Chromatin Remodeling Complexes. *Annual Review of Biochemistry* **78**: 273–304.

Cock JM, Godfroy O, Macaisne N, Peters AF, Coelho SM. 2014. Evolution and regulation of complex life cycles: a brown algal perspective. *Current Opinion in Plant Biology* **17**: 1–6.

Coelho SM, Godfroy O, Arun A, Corguillé GL, Peters AF, Cock JM. OUROBOROS is a
master regulator of the gametophyte to sporophyte life cycle transition in the brown alga
Ectocarpus. : 6.

Coelho SM, Peters AF, Charrier B, Roze D, Destombe C, Valero M, Cock JM. 2007.
Complex life cycles of multicellular eukaryotes: New approaches based on the use of model
organisms. *Gene* **406**: 152–170.

von Dassow P, John U, Ogata H, Probert I, Bendif EM, Kegel JU, Audic S, Wincker P,
Da Silva C, Claverie J-M, *et al.* 2015. Life-cycle modification in open oceans accounts for
genome variability in a cosmopolitan phytoplankton. *The ISME Journal* **9**: 1365–1377.

von Dassow P, Ogata H, Probert I, Wincker P, Da Silva C, Audic S, Claverie J-M, de
Vargas C. 2009. Transcriptome analysis of functional differentiation between haploid and
diploid cells of *Emiliania huxleyi*, a globally significant photosynthetic calcifying cell. *Genome*
Biology **10**: R114.

Decelle J, Stryhanyuk H, Gallet B, Veronesi G, Schmidt M, Balzano S, Marro S,
Uwizeye C, Jouneau P-H, Lupette J, *et al.* 2019. Algal Remodeling in a Ubiquitous
Planktonic Photosymbiosis. *Current Biology* **29**: 968-978.e4.

Decelle J, Veronesi G, LeKieffre C, Gallet B, Chevalier F, Stryhanyuk H, Marro S,
Ravanel S, Tucoulou R, Schieber N, *et al.* 2021. Subcellular architecture and metabolic
connection in the planktonic photosymbiosis between *Collodaria* (radiolarians) and their
microalgae. *Environmental Microbiology* **23**: 6569–6586.

Dierschke T, Flores-Sandoval E, Rast-Somssich MI, Althoff F, Zachgo S, Bowman JL.
2021. Gamete expression of TALE class HD genes activates the diploid sporophyte program
in *Marchantia polymorpha*. *eLife* **10**: e57088.

Dingman, Jennifer. **2015**. Virus induced cell death, evasion and resistance in the harmful
bloom-forming alga, *Heterosigma akashiwo*.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. **2013**. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Feldmesser E, Rosenwasser S, Vardi A, Ben-Dor S. **2014**. Improving transcriptome
construction in non-model organisms: integrating manual and automated gene definition in
Emiliania huxleyi. *BMC Genomics* **15**: 148.

Feng S, Cokus SJ, Zhang X, Chen P-Y, Bostick M, Goll MG, Hetzel J, Jain J, Strauss
SH, Halpern ME, *et al.* 2010. Conservation and divergence of methylation patterning in
plants and animals. *Proceedings of the National Academy of Sciences* **107**: 8689–8694.

Frada MJ, Bendif EM, Keuter S, Probert I. **2019**. The private life of coccolithophores.
Perspectives in Phycology **6**: 11–30.

Frada MJ, Bidle KD, Probert I, de Vargas C. **2012**. In situ survey of life cycle phases of the
coccolithophore *Emiliania huxleyi* (Haptophyta): *Emiliania huxleyi* life cycling. *Environmental*
Microbiology **14**: 1558–1569.

Frada M, Probert I, Allen MJ, Wilson WH, de Vargas C. **2008**. The "Cheshire Cat" escape
strategy of the coccolithophore *Emiliania huxleyi* in response to viral infection. *Proceedings*
of the National Academy of Sciences **105**: 15944–15949.

Frada MJ, Rosenwasser S, Ben-Dor S, Shemi A, Sabanay H, Vardi A. **2017**.

Morphological switch to a resistant subpopulation in response to viral infection in the bloom-forming coccolithophore *Emiliania huxleyi* (JS Weitz, Ed.). *PLOS Pathogens* **13**: e1006775.

Fu L, Niu B, Zhu Z, Wu S, Li W. **2012**. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**: 3150–3152.

Fulton JM, Fredricks HF, Bidle KD, Vardi A, Kendrick BJ, DiTullio GR, Van Mooy BAS. **2014**. Novel molecular determinants of viral susceptibility and resistance in the lipidome of *Emiliania huxleyi*. *Environmental Microbiology* **16**: 1137–1149.

Gallet B, Moriscot C, Schoehn G, Decelle J. **2024**. Cryo-fixation and resin embedding of biological samples for electron microscopy and chemical imaging v1.

Godfrey L, Crump NT, Thorne R, Lau I-J, Repapi E, Dimou D, Smith AL, Harman JR, Telenius JM, Oudelaar AM, *et al.* 2019. DOT1L inhibition reveals a distinct subset of enhancers dependent on H3K79 methylation. *Nature Communications* **10**: 2803.

Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, Robles M, Talón M, Dopazo J, Conesa A. **2008**. High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Research* **36**: 3420–3435.

Grau-Bové X, Navarrete C, Chiva C, Pribasnig T, Antó M, Torruella G, Galindo LJ, Lang BF, Moreira D, López-Garcia P, *et al.* 2022. A phylogenetic and proteomic reconstruction of eukaryotic chromatin evolution. *Nature Ecology & Evolution* **6**: 1007–1023.

Green JC, Course PA, Tarran GA. **1996**. The life-cycle of *Emiliania huxleyi*: A brief review and a study of relative ploidy levels analysed by flow cytometry. *Journal of Marine Systems* **9**: 33–44.

Hirooka S, Itabashi T, Ichinose TM, Onuma R, Fujiwara T, Yamashita S, Jong LW, Tomita R, Iwane AH, Miyagishima S. **2022**. Life cycle and functional genomics of the unicellular red alga *Galdieria* for elucidating algal and plant evolution and industrial use. *Proceedings of the National Academy of Sciences* **119**: e2210665119.

Horst NA, Katz A, Pereman I, Decker EL, Ohad N, Reski R. **2016**. A single homeobox gene triggers phase transition, embryogenesis and asexual reproduction. *Nature Plants* **2**.

Houdan A, Billard C, Marie D, Not F, Sáez AG, Young JR, Probert I. **2004**. Holococcolithophore-heterococcolithophore (Haptophyta) life cycles: Flow cytometric analysis of relative ploidy levels. *Systematics and Biodiversity* **1**: 453–465.

Houdan A, Probert I, Van Lenning K, Lefebvre S. **2005**. Comparison of photosynthetic responses in diploid and haploid life-cycle phases of *Emiliania huxleyi* (Prymnesiophyceae). *Marine Ecology Progress Series* **292**: 139–146.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR, Letunic I, Rattei T, Jensen LJ, *et al.* 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research* **47**: D309–D314.

Huisinga KL, Brower-Toland B, Elgin SCR. **2006**. The contradictory definitions of heterochromatin: transcription and silencing. *Chromosoma* **115**: 110–122.

Hunter JE, Frada MJ, Fredricks HF, Vardi A, Van Mooy BAS. **2015**. Targeted and untargeted lipidomics of *Emiliania huxleyi* viral infection and life cycle phases highlights molecular biomarkers of infection, susceptibility, and ploidy. *Frontiers in Marine Science* **2**.

Kao T-T, Lai M-W, Wang T-H, Yang C-L, Frada MJ, Ku C. **2024**. Haplotype-aware
multiomics unveils the regulatory basis of haplodiplontic life-cycle differentiation in a
cosmopolitan marine alga. : 2024.05.26.595999.

Keller MD, Selvin RC, Claus W, Guillard RRL. **1987**. Media for the Culture of Oceanic
Ultraphytoplankton1,2. *Journal of Phycology* **23**: 633–638.

Klaveness D. **1972**. *Coccolithus huxleyi* (Lohm.) Kamptn II. The flagellate cell, aberrant cell
types, vegetative propagation and life cycles. *British Phycological Journal* **7**: 309–318.

Lawrence M, Daujat S, Schneider R. **2016**. Lateral Thinking: How Histone Modifications
Regulate Gene Expression. *Trends in Genetics* **32**: 42–56.

Lee J-H, Lin H, Joo S, Goodenough U. **2008**. Early Sexual Origins of Homeoprotein
Heterodimerization and Evolution of the Plant KNOX/BELL Family. *Cell* **133**: 829–840.

Lenski RE, Levin BR. **1985**. Constraints on the Coevolution of Bacteria and Virulent Phage:
A Model, Some Experiments, and Predictions for Natural Communities. *The American*
Naturalist **125**: 585–602.

Li B, Dewey CN. **2011**. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. : 16.

Loppin B, Berger F. **2020**. Histone Variants: The Nexus of Developmental Decisions and
Epigenetic Memory. *Annual Review of Genetics* **54**: 121–149.

Love MI, Huber W, Anders S. **2014**. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology* **15**.

Mable BK, Otto SP. **1998**. The evolution of life cycles with haploid and diploid phases.
BioEssays **20**: 453–462.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. **2021**. BUSCO Update: Novel
and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for
Scoring of Eukaryotic, Prokaryotic, and Viral Genomes (J Kelley, Ed.). *Molecular Biology and*
Evolution **38**: 4647–4654.

Mikami K, Li C, Irie R, Hama Y. **2019**. A unique life cycle transition in the red seaweed
Pyropia yezoensis depends on apospory. *Communications Biology* **2**: 299.

Mikulski P, Komarynets O, Fachinelli F, Weber APM, Schubert D. **2017**. Characterization
of the Polycomb-Group Mark H3K27me3 in Unicellular Algae. *Frontiers in Plant Science* **8**:
607.

Monteiro FM, Bach LT, Brownlee C, Bown P, Rickaby REM, Poulton AJ, Tyrrell T,
Beaufort L, Dutkiewicz S, Gibbs S, *et al.* 2016. Why marine phytoplankton calcify. *Science*
Advances **2**: e1501822.

Morgan HD, Santos F, Green K, Dean W, Reik W. **2005**. Epigenetic reprogramming in
mammals. *Human Molecular Genetics* **14**: R47–R58.

Oyama R, Haba G, Kaido Y, Kanasugi T, Isurugi C, Kikuchi A, Sugiyama T, Jakab M,
Pujol S, Kikinis R. **2014**. P03.15: Towards improved ultrasound based analysis and 3D
visualisation of the fetal brain using 3DSlicer. *Ultrasound in Obstetrics & Gynecology* **44**:
200–200.

Pelusi A, De Luca P, Manfellotto F, Thamatrakoln K, Bidle KD, Montresor M. **2021**.

Virus-induced spore formation as a defense mechanism in marine diatoms. *New Phytologist* **229**: 2251–2259.

Pertea G, Pertea M. **2020**. GFF Utilities: GffRead and GffCompare. *F1000Research* **9**: 304.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. **2015**. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology* **33**: 290–295.

Rokitta SD, de Nooijer LJ, Trimborn S, de Vargas C, Rost B, John U. **2011**. Transcriptome Analyses Reveal Differential Gene Expression Patterns Between the Life-Cycle Stages of *Emiliania Huxleyi* (haptophyta) and Reflect Specialization to Different Ecological Niches1. *Journal of Phycology* **47**: 829–838.

Rokitta SD, Rost B. **2012**. Effects of CO_2 and their modulation by light in the life-cycle stages of the coccolithophore *Emiliania huxleyi*. *Limnology and Oceanography* **57**: 607–618.

Rosenwasser S, Mausz MA, Schatz D, Sheyn U, Malitsky S, Aharoni A, Weinstock E, Tzfadia O, Ben-Dor S, Feldmesser E, *et al.* 2014. Rewiring Host Lipid Metabolism by Large Viruses Determines the Fate of *Emiliania huxleyi* , a Bloom-Forming Alga in the Ocean. *The Plant Cell* **26**: 2689–2707.

Sakakibara K, Ando S, Yip HK, Tamada Y, Hiwatashi Y, Murata T, Deguchi H, Hasebe M, Bowman JL. **2013**. KNOX2 Genes Regulate the Haploid-to-Diploid Morphological Transition in Land Plants. *Science* **339**: 1067–1070.

Schmidt A. **2020**. Controlling Apomixis: Shared Features and Distinct Characteristics of Gene Regulation. *Genes* **11**: 329.

Schroeder DC, Oke J, Malin G, Wilson WH. **2002**. Coccolithovirus (Phycodnaviridae): Characterisation of a new large dsDNA algal virus that infects *Emiliana huxleyi*. *Archives of Virology* **147**: 1685–1698.

Takahashi Y-H, Schulze JM, Jackson J, Hentrich T, Seidel C, Jaspersen SL, Kobor MS, Shilatifard A. **2011**. Dot1 and Histone H3K79 Methylation in Natural Telomeric and HM Silencing. *Molecular Cell* **42**: 118–126.

Tatum D, Li S. **2011**. Evidence That the Histone Methyltransferase Dot1 Mediates Global Genomic Repair by Methylating Histone H3 on Lysine 79. *Journal of Biological Chemistry* **286**: 17530–17535.

Thangavel G, Nayar S. **2018**. A Survey of MIKC Type MADS-Box Genes in Non-seed Plants: Algae, Bryophytes, Lycophytes and Ferns. *Frontiers in Plant Science* **9**: 510.

Thiriet-Rupert S, Carrier G, Chénais B, Trottier C, Bougaran G, Cadoret J-P, Schoefs B, Saint-Jean B. **2016**. Transcription factors in microalgae: genome-wide prediction and comparative analysis. *BMC Genomics* **17**: 282.

Thomas R, Grimsley N, Escande M, Subirana L, Derelle E, Moreau H. **2011**. Acquisition and maintenance of resistance to viruses in eukaryotic phytoplankton populations: Viral resistance in *Mamiellales*. *Environmental Microbiology* **13**: 1412–1420.

Tomaru Y, Mizumoto H, Nagasaki K. **2009**. Virus resistance in the toxic bloom-forming dinoflagellate *Heterocapsa circularisquama* to single-stranded RNA virus infection. *Environmental Microbiology* **11**: 2915–2923.

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL,

Rinn JL, Pachter L. **2012**. Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. *Nature Protocols* **7**: 562–578.

Uwizeye C, Decelle J, Jouneau P-H, Flori S, Gallet B, Keck J-B, Bo DD, Moriscot C,
Seydoux C, Chevalier F, *et al.* **2021**. Morphological bases of phytoplankton energy
management and physiological responses unveiled by 3D subcellular imaging. *Nature*
Communications **12**: 1049.

Vardi A, Haramaty L, Van Mooy BAS, Fredricks HF, Kimmance SA, Larsen A, Bidle KD.
2012. Host-virus dynamics and subcellular controls of cell fate in a natural coccolithophore
population. *Proceedings of the National Academy of Sciences* **109**: 19327–19332.

Vigneau J, Borg M. **2021**. The epigenetic origin of life history transitions in plants and algae.
Plant Reproduction.

Von Dassow P, Montresor M. **2011**. Unveiling the mysteries of phytoplankton life cycles:
patterns and opportunities behind complexity. *Journal of Plankton Research* **33**: 3–12.

de Vries J, Monteiro F, Wheeler G, Poulton A, Godrijan J, Cerino F, Malinverno E,
Langer G, Brownlee C. **2021**. Haplo-diplontic life cycle expands coccolithophore niche.
Biogeosciences **18**: 1161–1184.

Wollmann H, Stroud H, Yelagandula R, Tarutani Y, Jiang D, Jing L, Jamge B, Takeuchi
H, Holec S, Nie X, *et al.* **2017**. The histone H3 variant H3.3 regulates gene body DNA
methylation in Arabidopsis thaliana. *Genome Biology* **18**: 94.

Yelagandula R, Stroud H, Holec S, Zhou K, Feng S, Zhong X, Muthurajan UM, Nie X,
Kawashima T, Groth M, *et al.* **2014**. The Histone Variant H2A.W Defines Heterochromatin
and Promotes Chromatin Condensation in Arabidopsis. *Cell* **158**: 98–109.

Zhao H, Lu M, Singh R, Snell WJ. **2001**. Ectopic expression of a *Chlamydomonas* mt+-
specific homeodomain protein in mt− gametes initiates zygote development without gamete
fusion. *Genes & Development* **15**: 2767–2777.

Zheng H, Xie W. **2019**. The role of 3D genome organization in development and cell
differentiation. *Nature Reviews Molecular Cell Biology* **20**: 535–550.