Sulfated mannan helps diatoms domesticate their microbiome

J. Krull[1,2,6], C.J. Crawford[4], C. Sidhu[3], V. Solanki[3], M. Bligh[3], L. Rößler[2,3], R.K. Singh[4],

G. Huang[2,3], C.S. Robb[2,3], H. Teeling[3], P.H. Seeberger[4], T. Schweder[5,6], J-H. Hehemann[1,2,3]*

[1] BIOM, Faculty of Chemistry & Biology, University of Bremen, Germany

[2] MARUM, University of Bremen

[3] Max Planck Institute for Marine Microbiology, Bremen

[4] Max Planck Institute of Colloids and Interfaces, Potsdam, Germany

[5] Institute of Pharmacy, University of Greifswald, Germany

[6] Institute of Marine Biotechnology, Greifswald

*jhhehemann@marum.de

Significance statement:

Eukaryotes host microbial partners that shape their health, yet how they selectively assemble

beneficial microbes remains unclear. Using diatom microalgae as a model, we show they

exude a sulfated mannan that nourishes highly adapted bacteria tracking them across the

global ocean. Our findings suggest that single-celled eukaryotes can "domesticate"

prokaryotes—analogous to how humans have domesticated animals—albeit on a microscopic

scale. Dominating much of Earth's aquatic surface, diatoms drive ~20% of global

photosynthesis. We propose that sulfated mannan contributes to this success by helping

diatoms shape microbial partnerships that underpin planetary energy balance and atmospheric

chemistry.

Abstract

Diatoms are a keystone phylum in Earth's ecosystems, specializing in oxygen production and carbohydrate fixation that fuels global food webs. Diatoms host a microbiome, but how they preferentially collect bacteria with complementary traits remains unknown. Here we show that diatoms exude a C6-sulfated α-1,3-mannan that serves as a selective carbon source for adapted bacteria. Its structure was resolved by NMR spectroscopy, chromatography, chemical synthesis, and enzymatic dissection. Biochemical, physiological, and structural analyses revealed that specialized *Bacteroidota* employ a four-enzyme pathway to metabolize this mannan. Metagenomic and transcriptomic data indicate that the mannan globally selects for bacteria carrying these enzymes and associated traits. Because the mannan provides only carbon, oxygen, sulfur, and hydrogen, bacteria must obtain other essential elements from alternative sources, reinforcing metabolic interdependence. We propose that diatoms use sulfated mannans to attract beneficial partners and exclude competitors, thereby engineering a microbiome that enhances their productivity and underpins carbon cycling.

Main

Domestication (lat. *domesticus,* belonging to the house) is a process where an organism with sufficient access to energy can maintain an environment that hosts other species with complementary traits (1). The process involves reciprocal exchange of molecules and activities enabled through cognate genetic adaptations in the interacting species. Over time, sustained environmental control can lead to the emergence of symbiosis (2) In multicellular eukaryotes, such as animals and plants, slow reproduction rates delay the detection of genetic changes in offspring, making it challenging to detect molecular signatures of domestication (3). As a result, the underlying process of domestication remains largely unknown, especially at the molecular level.

To investigate this process, we considered single celled eukaryotes, diatoms and their associated prokaryotes. In the sunlight fueled surface ocean, diatoms can produce several hundred thousand cells within days in just a milliliter of seawater (1, 4). They photosynthesize organic molecules that sustain the carbon metabolism of associated prokaryotes (4, 5). Importantly, both partners can be cultivated, which facilitates identifying coevolving genes and the molecules that mediate interactions. Some functions resemble feed, fertilizers, herbicides and fences. For example, diatoms release carbohydrates used by bacteria as a carbon source (6, 7). They also release antibiotics (5, 8, 9) that suppress bacteria with incompatible traits, analogous to fences deterring predators or herbicides suppressing weeds. Fed, protected, and relieved of competition, the retained bacteria can concentrate nutrients (10), and return them, along with hormones and vitamins (8, 10, 11), to the auxotrophic algal host. Hence, diatoms and bacteria form a model system to study the process of domestication.

Carbohydrates, such as the abundant carbon energy molecule glucose (100 Gt C annual synthesis rate (18), can mediate both positive and negative interactions. In closed ring form glucose provides energy, but in the open ring form, with an exposed aldehyde group, glucose is toxic. This aldehyde group is one reason why insulin must activate muscle cells to remove excess glucose, as overexposure will cause host cells to die (19). Similar dual effects of

glucose on microbes have been described (20)(21). Thus, the same molecule can serve as

both nutrient and toxin, offering hosts a mechanism to reward some partners while excluding

others. Such "functional flips" provide a molecular basis for domestication, challenging our

positively biased view (22) of how complex communities assemble. Extracting this ancient

insight from science, fiction (23), and philosophy (24), and applying it to organic molecules

thought to benefit microbes, may help to uncover unknown rules that regulate biomes,

including those central to the carbon cycle.

Diatoms exude substantial quantities of organic carbon in the form of complex carbohydrates

(glycans) (25-27). These glycans, alongside many other molecules, likely shape the

reproducible communities of bacteria attracted to or repulsed by diatoms. These include

Alphaproteobacteria (e.g., *Candidatus* Pelagibacter ubique; SAR11), *Roseobacteraceae,*

Gammaproteobacteria (e.g. Vibrionaceae), Bacteroidota (e.g. Polaribacter spp., *Formosa*

spp.), *Cyanobacteria, Verrucomicrobiota,* and many other taxa (28-30). The chemical

structures and mechanisms that govern glycan effects on bacteria remain difficult to resolve.

A glycan chain with over 1000 monomers, about the length of a bacterial cell, can be highly

anionic. Sulfate and carboxyl groups render these glycans extremely surface-active, promoting

coagulation with cations, minerals, bacterial cells, proteins, and other molecules (27, 31, 32).

The resulting gels impede separation and purification, hindering structural determination.

Without defined structures, the ecological effects of glycans on bacteria, with or without the

enzymes to metabolize, evade, or extract nutrients, remain obscure.

Here we resolved the structure of a sulfated mannan secreted by diatoms collectively

responsible for 20% of global primary production (18). This figure rises to 70% in coastal

upwelling zones and ~50% in temperate subpolar regions during spring and summer (33). T

The mannan selectively recruits bacteria that harbor a genetic island for mannan utilization,

consistently associated with diatoms across the global ocean microbiome. The mannan may

mediate association and disassociation between specific bacterial taxa, traits, functions. From

a eukaryotic perspective, prokaryotes are rapidly growing, abundant, mobile and accessible

sources of essential nutrients. Here we deduce how diatoms may "domesticate" and farm

prokaryotes.

Results

Diatoms may cultivate adapted bacterial taxa by exudation of sulfated mannan

During the analysis of diatom bloom-associated metagenomes and bacterioplankton genomes,

we noted the recurrence of bacterial taxa with a putative polysaccharide utilization locus (PUL)

for mannan that consistently co-occurred with diatoms (**Fig. 1**). The PUL is found in marine

Bacteroidota including *Formosa* spp., *Ochrovirga* spp., and *Polaribacter* spp., which are which

are globally distributed and often highly abundant, like the diatoms they frequently associate

with. Some of these species can account for ~10% of the bacterioplankton cells surrounding

diatoms (34-36). We reasoned that, among many organic molecules, a mannan may contribute

to a stable microbiome (37-40). To uncover this hypothetical mannan, we worked together with

the centric diatom *Conticribra weissflogii* (formerly *Thalassiosira weissflogii*), a diatom model

species (41).

Figure 1: *Bacteroidota* with putative mannan utilization enzymes follow diatoms during algal blooms. Shown

are data from six years from the long-term ecological research station (LTER) on Helgoland. The diatoms cells are

followed by certain *Bacteroidota* taxa that have a polysaccharide utilization locus (PUL, **Fig. S2**) with putative

enzyme and related genes that indicate utilization of a hypothetical, sulfated mannan secreted by the diatoms. The

DNA sequence of the mannan PUL of *Polaribacter* sp. Hel1_33_49 was mapped against metagenomes collected

previously from the LTER, Helgoland Roads. Grey shows diatom cell counts obtained by microscopy. Green shows

the PUL abundance over years calculated in terms of Reads Per Kilobase per Million mapped reads (RPKM). For

more information, see Materials and Methods.

A C6-sulfated α-1,3-mannan secreted by a diatom

We previously used anionic exchange chromatography (AEX) to show that several diatom

species, including *C. weissflogii*, secrete a fucose-containing sulfated polysaccharide (FCSP)

of unknown structure (27). By refining AEX chromatography to achieve higher molecular

resolution, we found that the material secreted by *C. weissflogii* comprises at least three

anionic glycans: the FCSP, a glucuronomannan (42, 43) and a previously unknown

homomannan. The FCSP and the glucuronomannan eluted at lower salt concentrations (0.5-

1.2 M NaCl) and contained diverse monosaccharides, whereas, the homomannan eluted only

at 2 M NaCl **(Fig. S1 A-C)**. Acid hydrolysis coupled with HPAEC-PAD (high-performance anion

exchange chromatography with pulsed amperometric detection) revealed a composition of

~95 mol % mannose. Additional ion chromatography revealed that the mannan contains an

almost equimolar ratio of sulfate to mannose **(Fig. S1 D)**.

During chromatography, concentration, and dialysis, the mannan remained soluble in both

seawater and deionized water. It reached solubility up to 20 mg mL^{-1}, the highest tested,

whereas other the algal polysaccharides (agar, κ-carrageenan, alginate and pectate) formed

gels under seawater concentrations of calcium and magnesium ions (44). These insoluble

glycans may target, or be targeted by, distinct bacterial populations (39), in contrast to the

soluble mannan. From 160 L of diatom culture, we obtained between 15-20 mg of the mannan

(~100 µg L^{-1}).

For configuration and connectivity, we used one- and two-dimensional NMR **(Fig. 2 A, B)**.

Details are provided in the supplementary information. Briefly, the major component was α-

1,3-D-mannopyranose with the C6-carbon de-shielded at 67.5 ppm, consistent with 6-*O*-

sulfation and confirming the equimolar mannose-to-sulfate ratio **(Fig. 2B, Supplementary**

Text). For further verification, we synthesized a mannan oligosaccharide and used it as a

reference for comparative NMR.

Figure 2: Diatoms synthesize a homomannan made of α-1,3-mannose-6-sulfate. In addition to

chromatographic structure elucidation **(Fig. S1 B, D)** we used automated glycan assembly (AGA) to synthesize a

mannan oligosaccharide of the NMR-hypothesized structure of the diatom mannan and compared the two spectra.

(**A**) [1]H-NMR overlay of an automated glycan assembly derived, synthetic α-1,3-mannose-6-sulfate

hexaoligosaccharide (green) and the mannan purified from *Conticribra weissflogii* culture supernatant (gray). (**B**)

Shown is an overlay of [1]H-[13]C HSQC spectra from the synthetic mannan hexamer (green) and the purified, diatom

mannan (gray). NMR spectra at 700 MHz were collected at 20 °C. HSQC-TOCSY and [1]H-[13]C HMBC can be found

in supplementary data. (**C**) Scheme of chemical synthesis of mannan oligosaccharides. L1: free reducing end solid-

phase linker (45). 1: Dibutyl 2-*O*-benzoyl-4-*O*-benzyl-6-*O*-(9-fluorenylmethoxycarbonyl)-3-*O*-levulinoyl-1-

phosphate-α-D-mannopyranoside.

We used automated glycan assembly to synthesize a reference mannan oligosaccharide. As

a building block we prepared ethyl-2-*O*-benzoyl-4-*O*-benzyl-6-*O*-(9-

fluorenylmethoxycarbonyl)-3-*O*-levulinoyl-1-thio-α-D-mannopyranoside, which was converted

into a dibutyl phosphate donor **(1) (Fig. 2C, Supplementary Text).** This enabled the

automated assembly of a hexa-α-1,3-mannose bearing one sulfate on each C6. Comparison

of NMR spectra from natural and synthetic mannan showed deviations of <0.03 ppm in the

proton and <0.1 ppm in the carbon dimension **(Fig. 2 A, B, Table S1)**, confirming the structure

as a C6-sulfated α-1,3-mannan. **(Fig. S2)**.

Polaribacter **spp. can metabolize the mannan**

We tested with *Polaribacter* strains - Hel1_33_49, _78 and _96 whether the PUL (46) enables

mannan metabolism. These strains have 100% identical 16S RNA genes and 98.5% genome-

wide average nucleotide identity (ANI) (47), providing three biological replicates. They have

the PUL, indicating a similar ability to metabolize the mannan. Accordingly, when the mannan

was provided as a limited carbon source, the three strains grew better **(Fig. S3-6, Table S2)**.

As a negative control, we used *Polaribacter* sp. Hel1_88 (36, 46), which lacks the mannan-

PUL (37). The Hel1_88 strain grew less on mannan in comparison to Hel1_33_49, _78 and

_96. The mannan remained unconsumed in the culture of *Polaribacter* sp. Hel1_88 **(Fig. S4)**.

In 2020 we showed *Polaribacter* transcribe this PUL in situ (38), the North Sea. To confirm

transcription and translation of enzymes capable of depolymerizing the mannan, we used cell

lysates of *Polaribacter* sp. Hel1_33_78 for activity tests. Enzymes were extracted from

bacterial cultures grown with or without the mannan **(Fig. S5,6)**. The enzymes in the lysates

from cells grown with the mannan digested the mannan. These expression and activity assays

suggest these enzymes of adapted *Polaribacter* spp. are for the metabolism of the mannan.

The in situ enzyme expression indicates bacteria respond to and interact with the mannan

released by diatoms into their environment.

The bacterial enzymes can hydrolyze the mannan

To verify enzyme activity and specificity, we used recombinant gene expression, enzymology

and structural biology **(Fig. 3)**. The PUL comprises 23 genes **(Fig. S2),** 21 of which encode

putative enzymes. We cloned these genes from *Polaribacter* sp. Hel1_33_78 into vectors,

transformed and expressed them in *Escherichia coli,* and conducted activity tests using purified

mannan, synthesized mannan oligosaccharides, and para-nitrophenyl-labeled substrates

(PNP-substrates). The products of these substrates digested with the enzymes were analyzed

by fluorophore-assisted carbohydrate gel electrophoresis (FACE) and reducing sugar assays

(Table S3). To further substantiate the findings, we obtained homologous enzymes from

another marine *Bacteroidota*, *Ochrovirga pacifica* S85, which was isolated from the Pacific

Ocean (48) with 69.8% ANI with the *Polaribacter* sp. Hel1_33_78 genome.

O. pacifica has a distantly related version of the PUL (49) with 68.8% ANI to the mannan PUL

of *Polaribacter* sp. Hel1_33_78. We found that the four enzymes are present in both PULs

(*Polaribacter*: PbSulf1-GHx, PbGH92_1, PbSulf2; *O. pacifica*: OpSulf1, OpGHx, OpGH92,

OpSulf2). These enzymes, comprising one unknown family GHx, one GH92 and two

sulfatases, hydrolyzed the mannan into mannose and sulfate **(Fig. 3A)**. The remaining 18

putative enzymes were cloned and tested but appeared inactive on the mannan **(Table S3)**.

For one sulfatase (OpSulf1) the validity of biochemical experiments could be supported by X-

ray crystallography analysis. For the other enzymes, which did not crystallize, we used *in silico*

generated AlphaFold2 (50) structure models to falsify the biochemical data.

199
200
Figure 3: Four bacterial enzymes in three genes hydrolyze the sulfated α-mannan from diatoms. (**A**).

Suggested degradative pathway for sulfated α-mannan. (**B**) Enzymatic digestions of the mannan via sequentially

added PbSulf1-GHx, PbGH92_1 and PbSulf2. Top: Product separation by fluorophore-assisted carbohydrate

electrophoresis using 2-aminoacridone as fluorophore. Middle: Free sulfate quantification. Bottom: Reducing end-

assay. Bars represent mean of n=3 and circles display individual data points. n.e. = no enzyme. (**C**) Digestion of

oligosaccharides produced by PbSulf1-GHx through PbGH92_1 were analyzed using LC-MS. Extracted ion

chromatograms at 259.013 *m/z* of the digestion compared with synthesized or commercially purchased mannose-

sulfate standards. Man-2S: Mannose-2-Sulfate; Man-3S: Mannose-3-Sulfate; Man-4S: Mannose-4-Sulfate; Man-

6S: Mannose-6-Sulfate. (**D**) Mannose release after a sequential digestion of the mannan by PbSulf1-GHx,

PbGH92_1 and PbSulf2 detected using HPAEC-PAD. (**E**-**F**) Structural model of enzymes involved in the

degradative cascade were generated using AlphaFold2. Active site regions (green color) were identified using

structural superimpositions with previously characterized enzymes. A simplified version of the degradative cascade

is depicted according to the symbol nomenclature for glycans (51). (**E**) The PbSulf1-GHx tandem enzyme with the

rotated active site region of the Sulfatase- and GHx-domain shown in top left and right, respectively. Grey arrows

indicate the rotation axis of each domain towards the full-length protein (below). (**F**) Exo-acting PbGH92_1 and

PbSulf2 have a pocket shape active site.

Mannan degradation is accomplished in four enzymatic steps. The first two steps are

completed by the multimodular enzyme PbSulf1-GHx. The enzyme contains a T9SS domain

for extracellular secretion. The sulfatase and GHx domain in PbSulf1-GHx work sequentially

to catalyze steps one and two of the cascade. The sulfatase domain belongs to the S1 family,

subfamily 51 (52). It is the first described activity of this subfamily and the first known sulfatase

with this specificity. The GHx domain is distantly related to endo α-1,2 mannanases of the

GH99 family and may therefore classify as a new family (53). Separating and visualizing

fluorophore-labeled reaction products on FACE gels showed hydrolysis of the mannan by

PbSulf1-GHx, producing a ladder-like pattern of oligosaccharides typical for endo-acting

enzymes (54) **(Fig. 3B).** The tandem enzyme released sulfate and oligosaccharides as shown

by ion chromatography and reducing sugar assays **(Fig. 3B)**. Liquid chromatography mass

spectrometry (LC-MS) showed that the released oligosaccharides constituted sulfated hexose

oligomers with degrees of polymerization between three and seven **(Fig. S7)**. The

oligosaccharides were consistently missing one sulfate group, e.g. a mannose tetrasaccharide

had only three sulfate groups, further indicating that both catalytic domains were active on the

mannan.

We reproduced the results with enzymes from *O. pacifica,* which encodes the two catalytic

domains as two separate genes. The enzymes OpSulf1 and OpGHx exhibit 53% and 36%

amino acid identity to the respective PbSulf1-GHx domains. The separated enzymes showed

that prior sulfatase activity is indeed necessary for GHx activity, suggesting that this enzyme

requires a desulfated mannose for productive binding **(Fig. S8)**. The AlphaFold2 model of

PbSulf1-GHx showed two appended catalytic domains with open, extended clefts **(Fig. 3E)**

typical for endo-acting glycoside hydrolases (54) and sulfatases (55), corroborating the

biochemical experiments. We obtained diffraction data from a crystal structure for the OpSulf1

at 1.49 Å resolution (**Fig. S9, Table S4**). The crystal structure confirmed an extended substrate

binding cleft compatible with the elongated mannan chain. The conserved catalytic residues

present in the center may remove one sulfate from the center of the bound mannan chain,

which then becomes accessible to the GHx **(Fig. S9B, C)**. In conclusion, we identified and

characterized two new enzyme activities, one endo C6-sulfate α-1,3-mannanase forming a

new CAZy family, and an endo C6-sulfate α-1,3-mannan sulfatase. These predicted outer

membrane enzymes may produce oligosaccharides of suitable size and shape for import into

the periplasm.

The third step in the mannan degradation cascade was catalyzed by the periplasmic

PbGH92_1, a mannosidase that removed C6-sulfated mannose from the oligosaccharides.

Periplasmic localization of PbGH92_1, which belongs to the GH92 family (56), was predicted

by the Sec/SPI signal peptide (57). The GH92 family includes Ca^{2+}-dependent exo-acting α-

1,2/3/4/6 mannosidases, but an enzyme acting on, specific for sulfated mannan remains

unknown. Hydrolysis of oligosaccharides into mannose-6-sulfate and a minor amount of

mannose was shown by FACE, reducing sugar assays and HPAEC-PAD **(Fig. 3B-D)**. The

mannose-6-sulfate product was verified using LC-MS **(Fig. 3C)**. For this analysis, we

synthesized mannose-2/3/4-sulfate isomers **(Supplementary Text, Data S1, Fig. S10)**. The

retention times revealed that only the mannose-6-sulfate aligned with the enzyme product peak

and therefore confirmed this isomer as the product of the PbGH92_1 **(Fig. S11)**. The

AlphaFold2 model showed a pocket topology of size and shape typical for an exo-acting GH

(Fig. 3F) binding one monomer in the -1 subsite (54). Conclusively, PbGH92_1 is a 6-sulfate

α-1,3-mannosidase, representing a new specificity within the GH92 family.

The fourth step was catalyzed by PbSulf2 that desulfates mannose-6-sulfate to yield mannose.

A periplasmatic location was predicted by a Sec/SPII signal peptide. PbSulf2 is classified into

sulfatase family S1, subfamily 11. Enzymes in this subfamily have been described to act on

glucosamine-6-sulfate (58) and 6-sulfated porphyran oligosaccharides (59), but not on

mannose-6-sulfate. PbSulf2 removes a sulfate group from mannose-6-sulfate as shown by

appearance of the final product mannose in HPAEC-PAD measurements **(Fig. 3D)** and the

disappearance of sulfated mannose in a complementary FACE experiment **(Fig. 3B).** The

AlphaFold2 model aligned with the two previously described subfamily members (PDB ID

Codes 5G2V / 7LJ2, rmsd: 0.613 Å / 0.845 Å) and showed a pocket topology typical for exo-

acting sulfatases (55). PbSulf2 is a mannose-6-sulfate sulfatase, representing a new

specificity within this enzyme family.

In summary, we elucidated a minimal enzymatic cascade consisting of four enzymes that

achieve complete hydrolysis of the mannan into mannose and sulfate. The oligo- and

monosaccharide structures produced by the enzymes confirmed the mannan structure solved

with NMR.

Biocatalytic quantification of mannan synthesis and degradation

To verify and quantify the synthesis of the mannan by diatom cells and its consumption by

adapted *Polaribacter* spp. we used the here characterized enzymes to develop a quantitative

assay. The assay involved a combination of chromatography and enzyme digest steps for

structure-specific quantification. The enzymes specifically produce mannose that provides a

much stronger signal than the undigested polysaccharide with different quantitative

approaches, e.g., reducing sugar assay or HPAEC-PAD **(Fig. S12A)**. Chromatographic

separation and enrichment of the mannan were crucial for enzyme digestion. Without this

purification step, we found the enzymes remained inactive, potentially due to sulfate groups

from FCSP or other sulfated glycans binding to and interfering with the enzymes in the mixture

(60). After chromatographic isolation of the mannan, the biocatalytic assay detected up to

842 ± 62 µg L^{-1} in the diatom culture supernatant **(Fig. 4A).** The concentration was positively

correlated with the number of diatom cells ($y = 3.97 \times 10^{-3} x - 2.94 \times 10^{-2}$, $P = 1 \times 10^{-23}$, $R^2 =$

0.95). Each individual *C. weissflogii* cell produced on average a net amount of 3.9 fmol carbon

in form of mannan **(Fig. 4B).** Substantially less mannan was present in diatom cells or particles

(Fig. S12B,C). Notably, this new biocatalytic assay detects almost ten-fold higher

concentrations (842 ± 62 µg L^{-1}) compared to our initial assessment (~100 µg L^{-1}) based on

the preparative mannan purification yield, highlighting the analytic value of enzyme-based

assays.

Next, we conducted growth experiments with *Polaribacter* sp. Hel1_33_49 and applied the

biocatalytic assay to quantify mannan consumption **(Fig. 4C)**. The bacteria grew to an OD$_{600}$

of 0.2 when providing 0.1% (w/v) of the mannan as the limiting carbon source. Mannan

concentration and bacterial abundance were negatively correlated ($y = -5.62 x + 1.09$, $P = 1 \times$

10^{-18}, $R^2 = 0.96$) **(Fig. 4D)**. Conclusively, under nutrient-replete conditions, the bacteria can

remove the mannan within a few days.

To test whether other diatoms also release this type of mannan, we adapted the biocatalytic

assay into a qualitative version that rapidly identifies the structure in different species. We

tested *Thalassiosira pseudonana*, *Thalassiosira rotula* and *Chaetoceros affinis,* genera of

global ecological and biogeochemical relevance (34). We used AEX to enrich the mannan from

each culture supernatant, applied the enzymes and analyzed the products using FACE **(Fig.**

4E, F). For each diatom the first two enzymes (PbSulf1-GHx) produced the same set of

oligosaccharides that we identified above as oligomers of C6-sulfated α-1,3-mannose missing

one sulfate group. PbGH92_1 released mannose-6-sulfate, showing that the same mannan

structure is produced by different diatoms common around the globe.

Figure 4: Synthesis of mannan by diatoms and consumption by bacteria quantified with the mannan degrading enzymes. (**A**). Growth analysis of *Conticribra weissflogii* cultures using cell counts (black) and enzymatic quantification of sulfated α-mannan (green) in the anionic fraction of the culture supernatant. Data points were modeled using a logistic growth function employing the non-linear least squares method for fitting. n=4. (**B**) Linear regression analysis of diatom cell counts vs. mannan C production. The net production of mannan per cell was 3.9 fmol C cell^{-1} with an R²=0.95. (**C**) *Polaribacter* sp. Hel1_33_49 grown with purified mannan for one week at 18 °C. The black line represents optical density measured at 600 nm. The green line shows the mannan consumption as detected via enzymatic quantification. Data points were modeled using a logistic growth function employing the non-linear least squares method for fitting. n=3. (**D**) Linear regression analysis of mannan concentration vs. optical density of the growth curve. (**E**-**F**) Mannan identification in anionic exudates purified from 0.5-5 M NaCl from AEX from *Conticribra weissflogii*, *Thalassiosira pseudonana*, *Thalassiosira rotula* and *Chaetoceros affinis*. Products were digested with PbSulf1-GHx (**E**) and PbGH92_1 (**F**) and analyzed via fluorophore-assisted carbohydrate electrophoresis using 8-aminonaphthalene-1,3,6-trisulfonic acid (**E**) or 2-aminoacridone (**F**) as fluorophore. Negative controls (-) contained heat inactivated enzyme.

Mannan draws adapted bacteria towards diatoms around the globe

To test the hypothesis that the mannan selects for bacteria with these enzymes we quantified the mannan PUL within the *Tara* Oceans metaG dataset (61). As negative and positive controls for glycan structure-based selection of adapted microbiome members, we used bacterial PULs specific for known glycans from red, green and brown algal phyla. Preconditions for the PULs used as controls were that their target glycans are soluble and therefore accessible to

planktonic bacteria, that the glycan structure has been verified by NMR and/or other methods,

and that the structure and function of the PUL or of a closely related variant has been

empirically verified with biochemical experiments. As negative control we used porphyran, a

mixed linked α-, β-galactan that is also sulfated on C6, and the corresponding PUL from the

flavobacterium *Zobellia galactanivorans* Dsij[T] (62, 63). Compared to diatoms, porphyran-

producing red macroalgae *Porphyra* spp., are sessile, require rocks or other hard substrates,

grow along coasts. This range restriction makes the corresponding porphyran-PUL a suitable

negative control.

As a positive control, we used laminarin (chrysolaminarin), a β-glucan that is also produced by

diatoms, and the corresponding PUL from a metagenome-assembled genome (MAG

C_MB344, accession: GCA_943788335) of a flavobacterial *Aurantivirga* species (38, 64, 65).

Finally, we used ulvan, a sulfated rhamnogalacturonan produced by green macroalgal species

including *Ulva* spp. *Ulva* spp. grows attached and planktonic, moves via currents between

coastal and offshore regions (66), hence whether they, corresponding PUL can serve as

negative or positive control was unclear. The corresponding PUL (67) is from the

flavobacterium *Formosa agariphila* KMM 3901[T] (68, 69).

Mannan PUL abundances ranged from 0.0025 to 0.41 RPKM in the 45 sampled sites including

open and coastal ocean areas of the Atlantic, Pacific and Indian Ocean (**Fig. S13A**). Higher

mannan PUL abundances were detected at upwelling sites (Benguela: 0.41 RPKM, Chile: 0.06

RPKM, Peru: 0.03 RPKM, California: 0.04 RPKM) where higher nutrient concentrations

enhance diatom-dominated algal blooms (70, 71). The positive control, laminarin PUL showed

comparable abundances between 0.006 and 2.27 RPKM **(Fig. S13C)**. The porphyran PUL

showed the lowest RPKM values of the tested glycans and was absent in the open ocean **(Fig.**

S13E), as expected for this negative control. The ulvan PUL showed abundances between

0.002 and 0.08 RPKM but at different locations compared to the mannan and laminarin PULs

(Fig. S13D). The observed differences between the compared groups were significant across

the dataset (Friedmann-Test: $P = 6 \times 10^{-21}$).

Metagenomics of bacterioplankton and eukaryotic 18S rRNA-based taxonomy revealed that

the mannan PUL is associated with diatoms. To verify that diatoms were the main

phytoplankton associated with the large RPKM values of PULs at the time of sampling for the

Tara Ocean metaG dataset, we used previously reported total V9 read numbers as a proxy for

diatoms and correlated those with PUL reads (71). This approach was previously verified by

showing that V9 reads correlate with diatom algal cell counts (72). At the sites where

microalgae were measured, diatoms were a major phytoplankton group (71). Linear regression

analyses showed that the abundance of the mannan PUL correlated with diatom read numbers

($y = 7.7 \times 10^{-5}\ x + 6.2 \times 10^{-3}$, $P = 1.5 \times 10^{-3}$, $R^2 = 0.44$) **(Fig. 5A)**. The laminarin PUL also

significantly correlated with diatom abundance ($y = 7.3 \times 10^{-4}\ x + 1.7 \times 10^{-3}$, $P = 6.0 \times 10^{-4}$, R^2

$= 0.49$) **(Fig. 5B)**. In contrast, we did not observe correlation between the ulvan PUL and

diatom cell abundances ($P = 0.35$; $R^2 = 0.05$), which is consistent with the synthesis of this

glycan by green algae instead of diatoms **(Fig. C)**.

Diatoms are the only known microalgae that synthesize both, the mannan and the laminarin.

Hence, we asked whether the PULs for those two glycans co-occurred, which would further

support that diatoms are the source of these carbon-rich structures. We found a strong linear

relationship between mannan and laminarin PULs ($y = 0.158\ x + 5.35 \times 10^{-3}$, $P = 4 \times 10^{-24}$, $R^2 =$

0.91, **Fig. 5F**). In contrast, the PULs for mannan and ulvan did not correlate ($P = 0.414$; $R^2 =$

0.02). Finally, we queried for gene expression of the Polaribacter GH92 and found transcripts

in the *Tara* Ocean metaT dataset (61). The presence of transcripts suggests associated

bacteria degrade the mannan released by diatoms globally (**Fig. 5E)**. Spatially, temporally and

phylogenetically separated micro- and macroalgae produce distinguished glycans.

Conclusively, these diatom glycans, which select for specifically adapted bacteria, may

contribute to the formation of a functional and predictable mosaic of host specific microbiomes

in the sunlit ocean.

Figure 5: Bacteria with mannan degrading enzymes follow diatoms around the globe. Global abundance of the mannan utilization locus from *Polaribacter* sp. Hel1_33_49 from *Tara* Oceans database. The DNA sequence of the Mannan PUL was mapped against the raw metagenome reads of *Tara* Oceans surface water samples filtered at sizes ranging 0.2-3 µm across 45 different stations (61). Read abundances were calculated in terms of RPKM and normalized by length of the PUL and displayed as filled circles in SI. Corresponding plots for the laminarin PUL from *Aurantivirga* MAG C_MB344 (38), the ulvan PUL from *Formosa agariphila* KMM 3901[T] (67) and the porphyran PUL from *Zobellia galactanivorans* Dsij[T] (62) can be found in SI. (**A**) Raw read abundance of the clade of Bacillariophyta within the V9 rDNA metabarcoding dataset from *Tara* Oceans surface water samples filtered at sizes ranging 5-20 µm across 35 stations. Please note: No diatom data were available for the remaining stations. (**B-D**) Linear regression analyses of raw read abundance of Bacillariophyta in V9 rDNA metabarcoding dataset from *Tara* Oceans versus RPKM of different PULs across the same stations. Outliers where identified individually for each plot using the studentized residuals method with a threshold of three and excluded from the linear regression. Plots display the correlation analysis of Bacillariophyta vs. (**A**) mannan PUL, (**B**) laminarin PUL and (**C**) ulvan PUL. (**D**) Linear regression analyses of mannan PUL and Laminarin PUL RPKM within the *TARA* Oceans metagenomes. **(E)**

400   The GH92 sequence from the *Polaribacter* cascade was searched against the Tara Oceans OM-RGCv2 database,

401   which includes meta-transcriptomic (metaT) data, using BLASTp with an E-value cutoff of ≤ 1e-20 through the

402   Ocean Gene Atlas v2 web platform. The resulting hits were used to calculate relative abundance, expressed as the

403   percentage of mapped reads.

Discussion

Here we show that diatoms secrete a system of anionic glycans that shape microbiomes in multiple ways. The structurally complex FCSP may protect against enzymatic degradation and invasion (73). The carboxyl groups of glucuronomannan may chelate rare trace metals (74). In contrast, the structurally simple mannan provides an easily accessible carbon source for adapted bacteria such as *Polaribacter* spp. which in return supply diatoms with missing nutrients, metabolites and vitamins (75). The characterized enzyme cascade can now be applied to quantify this mannan and assess its role in the carbon cycle (76). The global abundance of mannan degrading enzymes **(Fig. 5)** underscores its ecological importance.

The mannan can influence microbes both positively and negatively because sulfation renders glycans protein surface-active. Negatively charged sulfate groups bind to positively charged amino groups of proteins, and potentially interfere with their function (60). C6-sulfation extends the sulfate group farther from the sugar backbone, allowing it to reach positively charged residues even in protein pockets (62). Indeed, cation exchange resins designed to bind diverse protein surfaces with high capacity often use sulfated polysaccharides (77). C6-sulfation may be one of the reasons explaining why, of the three anionic diatom glycans, the mannan was the strongest binder of the surface-exposed amines **(Fig. S1 A-C).**

Prokaryotes depend on proteins to degrade, sense, bind, infect, import, move, replicate, transfer, and communicate (78). Many protein structures contain surface-exposed arginine, lysine or histidine residues that can be targeted by sulfate groups (77). This hypothesis aligns with Wheeler and colleagues who uncovered that mucilage glycans exuded by animal epithelia "tame" bacteria by binding to virulence proteins (79). By analogy, diatom-derived mannans may bind proteins across bacteria, eukaryotes, or even viruses, reducing their invasive potential.

During algal blooms, sulfated glycans secreted by diatoms, including the mannan **(Fig. S1 A-C)**, coagulate into aggregates that entrap bacterial cells, proteins, minerals and other molecules (27, 80-82). Bacteria can bind those glycans, or be bound by them. This interaction

may especially capture cells lacking evasive mechanisms, promoting their incorporation into

sinking particles. The outcome is a relative increase of some taxa and a decline of other in the

sunlit ocean. As dominant algal taxa shift, their exuded sulfated glycans also change, selecting

for new bacterial traits **(Fig. 5)** and driving host-specific microbiomes (83, 84).

Trade-offs may shape adaptive strategies used by bacteria to evade surface-active glycans.

By digesting the mannan the *Polaribacter* spp. "domesticated" by diatoms obtain both carbon

and energy, while also removing the interference caused by sulfation. In contrast, bacteria,

lacking these enzymes minimize interaction as an alternative defense strategy. For instance

SAR11 *Alphaproteobacteria*, among the most successful groups of bacterioplankters around

diatoms (85), have a "non-stick" cell surface that bind fewer surface-active glycans (86). This

reduced binding is thought to underlie their greater resistance to grazing (86), but it comes at

the cost of being unable to access glycan-derived carbon. Similar trade-offs are seen in

Roseobacteraceae (75), which focus on smaller organic molecules as carbon sources (87). By

avoiding competition for glycans, they maintain a free-living, planktonic lifestyle that

nonetheless supports a "domesticated" association with diatoms.

This "competition-dispersal" tradeoff has been described in *Vibrio cyclitrophicus,* where closely

related strains show signs of recent sympatric speciation into two groups: strong vs. weak

binders (88). Strains with surface pili, including the mannose sensitive hemagglutinin (MSH)

type involved in host infection (89), bind and exploit surface glycans more effectively than the

strains lacking MSH genes. However, non-piliated strains, though less efficient at glycan

binding and exploitation, can avoid being irreversibly trapped by glycans (90). Surface-active,

adhesive glycans (27) may therefore disproportionately bind bacteria that rely on MSH pili or

other attachment proteins, constraining those most poised to intimately interact with eukaryotic

hosts. Similar mechanisms operate in animals: epithelial cells secrete the heavily glycosylated

immunoglobulin A (IgA) to constrain pathogens. IgA enchains and immobilizes bacteria in

aggregates, facilitating their clearance from the intestine (91). Cells trapped by glycans or

glycoproteins can be expelled or separated from microbiomes by multiple processes, including

bowel movements (91), sinking (83), filtration (27) and consumption (86).

In summary, glycans can simultaneously support (92, 93) and suppress (79, 94) bacterial traits.

Highly defensive types such as SAR11, with cell surfaces that minimize interaction, or

Polaribacter spp., equipped with enzymes to deconstruct glycans, can avoid being trapped. In

contrast, offensive bacteria such as Vibrio, rich in surface exposed proteins for attachment,

degradation, or infection, are more easily trapped and may therefore remain rare (95, 96).

Alternatively, they may be "tamed" by glycans, downregulating interactive surface proteins,

while upregulating dispersal proteins such as flagella (79). This short-term adaptation, imposed

by surface-active glycans, enhances dispersal, reduces competition (90), and protects the host

with its domesticated partners. While we highlighted only a few cases, such interactions likely

affect every cell to some extent. Through this dual capacity to nourish and constrain, glycans

contribute to the emergence of beneficial microbiomes and productive hosts.

To conclude, with sufficient light, CO_2, and nutrients, bacteria cannot escape the abundance

of photosynthesis-derived surface-active glycans. Around diatoms, these glycans enforce a

"domesticated" state through long-term genetic adaptation, or a non-virulent, tamed phase

through short-term physiological shifts. Improved molecular understanding will help resolve

uncertainties in the large-scale application of such glycans. Future work should focus on the

mechanisms by which surface-active glycans protect the collective of domesticated

prokaryotes and their eukaryotic hosts against competition, predation, and infection.

This process of "domestication" is important in today's context. By fixing CO_2 into simple

glycans, diatoms sustain rapidly growing bacteria that in turn boost diatom productivity. In

response, diatoms invest in more complex glycans that resist degradation, locking away

carbon. Through these reciprocal interactions, mutualistic and antagonistic alike, CO_2 is fixed

into glycans in ways that reinforce climate stability. Thus, by secreting sulfated glycans that

both nourish and constrain, diatoms domesticate their microbiomes, a microscopic strategy

with planetary consequences.

Materials and Methods

Mannan PUL abundance in *TARA* Oceans dataset

Metagenome data of bacterioplankton during spring phytoplankton blooms was generated in previous studies (29, 38-40, 97). For the global analysis, we used metagenome data from the *TARA* Oceans Expedition (61). The surface water metagenomes (0.2 to 3 µm) from 45 different stations were included in the analysis. Mapping of PULs was done with Bowtie2 (98) in the SqueezeMeta v1.6.2 pipeline. PUL abundances as reads per kilobase per million mapped reads (RPKM) were calculated using the formula: (metagenome reads mapped to the PUL*10^6)/(length of PUL in kbp × total reads in a metagenome). For correlation of PULs to diatom abundance, we extracted the read numbers from 25 surface water samples (5-20 µm) from the eukaryotic V9 metabarcoding data (71).

Mannan purification

C. weissflogii was cultured non-axenic in NEPCC medium (99). Diatom growth was monitored by cell counting using a Neubauer counting chamber (Marienfeld, Königshofen, Germany). Cells were separated from culture the supernatant using a continuous flow centrifuge (Thermo Fisher Scientific, Walthan, MA, USA) set at 150 mL min^{-1} and 6800 x g. The culture supernatant was filtered using a peristaltic pump through Whatman® glass microfiber filters (Grade GF/F, 0.70 µm). 160 L of *C. weissflogii*, cell and debris free, culture supernatant were loaded onto a preparative scale, self-assembled AEX column. Elution occurred with appropriate NaCl concentrations. Samples were dialyzed against MilliQ water (Merck) and freeze-dried prior to monosaccharide composition and sulfate analysis.

Monosaccharide composition and sulfate content analyses

Purified mannan was dissolved in MilliQ water at 1 mg mL^{-1} and hydrolyzed by 1 M HCl (100 °C, 24 h) in in pre-combusted glass vials. HCl was removed by vacuum or diluted 1:1000 in MilliQ water (Merck) prior to further analyses. Monosaccharide measurements occurred as described previously using a Dionex ICS-5000+ system (Thermo Fisher Scientific) (100) and

512   for sulfate measurements we used a Metrohm 761 compact ion chromatograph (Metrohm,

513   Herisau, Switzerland). All samples for monosaccharide and sulfate composition were prepared

514   in triplicates.

Cloning of enzyme genes into plasmids, recombinant expression and purification

Genes were cloned using the USER cloning or Gibson assembly methodologies (New England

Biolabs, Ipswich, NE, USA) with genomic DNA from *Polaribacter* sp. Hel1_33_78. Primer pairs

are provided in supplementary table S2. USER cloning constructs were cloned into the pet28A

vector (101) without signal peptides (102). Clones were validated by DNA sequencing.

Plasmids transformed into *E. coli* BL21 (DE3) were induced in 1 L lysogeny broth (LB) medium

containing kanamycin and isopropyl-β-D-1-thiogalactopyranoside (IPTG). For sulfatase

activation, the formylglycine generating enzyme plasmid (Addgene plasmid repository,

Watertown, MA, USA) (103), which was co-transformed, was induced with L-arabinose and

ampicillin added right before cooling to 12 °C. Protein purification was conducted using

standard procedures (104).

Enzyme activity and specificity assessment

Enzyme reactions were carried out in 10 mM MOPS pH 7.5 and 500 mM NaCl with 0.1% (w/v)

of mannan and enzymes were added in excess (0.01-0.1 mg mL^{-1}). 3.5% Sea Salts (Sigma

Aldrich, Burlington, MA, USA) were also added. Enzyme reactions were monitored using

fluorophore-assisted carbohydrate electrophoresis (FACE) with carbohydrates labeled with 2-

aminoacridone as described in (105). Reducing-end measurements were carried out using

PAHBAH as detection reagent as described previously (7). The release of sulfate and

mannose from enzymatic reactions were measured using ion chromatography and HPAEC-

PAD. Enzymatic assays were carried out as technical replicates in triplicates unless otherwise

stated.

Crystallization, data collection, structure solution, refinement and AlphaFold2

Crystallization was performed in two drop 96-well crystallization plates in sitting drop format

using commercial crystallization screens. Diffraction data was collected on DESY beamline

P11. Measured reflections intensities were indexed, integrated and scaled using the autoproc

pipeline available at the beamline facility. OpSulf1 structure was solved using molecular

replacement with the AlphaFold2 predicted model as reference in PHASER (106). For

automatic model building we used BUCCANEER (107). Refinement was done in REFMAC5

(108) and COOT (109). Model and structure factors for OpSulf_1 were deposited in the Protein

Data Bank (PDB) with accession 9FVT. Corresponding data-processing and refinement

statistics are summarized in Table S4. The structural comparison of OpSulf with homologs

were performed using the PyMOL v.2.3.2 (Schrödinger, New York, NY, USA). For enzymes

that did not crystallize we used models generated with AlphaFold2 for structural analyzes (50).

Liquid chromatography-mass spectrometry of mannose-sulfate and mannan digests

Standards and digests were diluted in 90% UHPLC-grade and 10% (v/v) MilliQ water (Merck)

for LC-MS measurements. 5 µL were injected to an Accucore-150-amide HILIC column

(Thermo Fisher Scientific) held at 60 °C on a Thermo Fisher Vanquish Horizon UHPLC

coupled to a Thermo Fisher Q-Exactive Plus MS. Mobile phases A and B consisted of 10 mM

ammonium formate at pH 5 and UHPLC-grade acetonitrile, respectively. A gradient started

after 1 min at 90% B and decreased from 90% to 40% B over 40 min at a flow rate of 0.4 mL

min^{-1}. The column was equilibrated at 90% for 16 min at the end of each run. Mono- and

oligosaccharides was detected in negative mode. The detailed MS conditions are provided in

the SI material.

Bacteria cultivation

Growth experiments with *Polaribacter* spp. Hel1_33_49/78/96 and Hel1_88 were carried out

in HaHa-medium supplemented with 0.1 mg mL^{-1} yeast extract and 2 mg mL^{-1} mannan from

either *C. weissflogii* or *C. affinis* (110). We used *Polaribacter* sp. Hel1_33_78 for in depth

physiological experiments. Cells were harvested by centrifugation, the pellets chemically lysed.

Lysed cells, enzymes were activity tested on *C. weissflogii* and *C. affinis* mannan. Digests

occurred overnight using 3.5% Sea Salts (Sigma), 10 mM MOPS pH 7.5 and 0.5 mg mL^{-1}

mannan and were analyzed using carbohydrate-polyacrylamide gel electrophoresis (C-

PAGE). C-PAGE analysis was carried out as described in (111) with modifications described

in the SI. *Polaribacter* sp. Hel1_33_49 was grown in 3.5% Sea Salts (Sigma) with supplements

(110). The mannan was used as limiting carbon source at a concentration of 1 mg mL^{-1}. Growth

curves (absorbance 600 nm) were plotted with a logistic growth function using the non-linear

least squares method. Substrate consumption was verified qualitatively using C-PAGE.

Enzyme assisted quantification of the mannan

Cultures of *Conticribra weissflogii* were grown in NEPCC medium as described above.

Subsamples of 200 mL of the culture were filtered to remove cells and debris. The mannan in

the filtrate was purified using AEX. After desalting the mannan was digested PbSulf1-GHx and

PbGH92_1. Undigested sample served as negative control. Detection occurred with the

PAHBAH assay. Results were compared with a standard curve containing defined amounts of

mannan. The recovery was determined 80 ± 5% (n=3) by comparison to untreated mannan

(fig S7C).

Extraction of mannan from cellular, diatom biomass

After diatom cultivation the cell pellet was freeze-dried and sequentially extracted using MilliQ

water (Merck), 0.4 M EDTA (pH 8) and 4 M NaOH with 0.1% NaBH4 (27). For 1 g of biomass

20 ml of sample were used. In each extraction step the sample was sonicated and centrifuged

at 14.000 x g for 10 minutes. NaOH extracts were neutralized to pH 7.5 using HCl and diluted

in MilliQ to < 500 mM NaCl. The mannan content in all extracts was determined as described

above.

Biocatalytic detection of mannan in multiple diatom species

The diatom strains *Thalassiosira pseudonana*, *Chaetoceros affinis* and *Thalassiosira rotula*

were cultivated as described above. 20 L of culture supernatant were concentrated after

filtration using GF/F. The filtrate was concentrated by ultrafiltration and then dialyzed to remove

salts. The dialyzed concentrate was purified by anionic exchange chromatography. The

purified mannan was again dialyzed and then lyophilized. The dialyzed and freeze-dried

samples were used as substrates for enzymatic assays using PbSulf1-GHx_1 and PbGH92_1.

Digested samples were analyzed with FACE (116).

Nuclear magnetic resonance and chemical synthesis of oligosaccharides

All chemicals used were reagent grade and used as supplied unless otherwise noted. A

detailed description of the structural elucidation and glycan synthesis processes is provided in

the supplementary information. The automated syntheses were performed on a custom-built

synthesizer (Potsdam-Golm, Germany)(112-115). More synthesis and analysis details are

provided in the SI.

Acknowledgements:

We thank Manual Liebeke for access to LC-MS devices and advise. We also thank C. Eschen, P. Subramanya, P. Schoppmeier for large-scale cultivation of *Conticribra weissflogii* and K. Imhoff for advice and support during sulfate quantification and T. Trautmann, A. Bolte and K. Föll for technical support. We thank Hagen Buck-Wiese for help with the LC-MS analysis, S. Niggemeier for assistance in cloning and expression and A. Sichert for help with genome assembly. We thank André Scheffel for providing the *C. weissflogii* culture. We further thank M. Schultz-Johansen for valuable discussions and his substantial help with the sampling campaigns.

Funding: J-H.H. was funded by the German Research Foundation through the Heisenberg program Grant (HE 7217/5-1). The work was supported by the European Research Council, ERC grant, C-Quest, (Grant 101044738) to J-H.H. The work was supported by the PriME collaboration of the Simons Foundation, Grant 970824 to J-H.H. This study was supported by the German Federal Ministry of Education and Research (BMBF) in the frame of the BaMS-BALI project (031B0915D1/2) and the German Research Foundation (DFG) for the Research Unit FOR 2406 "Proteogenomics of Marine Polysaccharide Utilization" (POMPU) by Grants to J-H.H. (HE 7217/2-3), T.S. (SCHW 595/10-3) and H.T (TE 813/2-3). The work was supported by the Cluster of Excellence initiative (EXC-2077–390741603). C.J.C. was funded by MSCA grant: MARINEGLYCAN (101029842). The Max-Planck Society supported P.H.S., J-H.H, C.J.C. and R.K.S.

Author contributions: J.K. and J-H.H. planned and analyzed the experiments with support by T.S.. J.K. and J-H.H. designed the artwork with help of M.B.. J.K. cloned and expressed the genes, conducted all activity assays, purified the glycans and cultured marine diatoms. P.H.S., C.J.C. and R.K.S. synthesized the oligosaccharide and mannose-sulfate standards and conducted all NMR experiments. V.S. provided structural models and refined the crystal structure. H.T., C.S., J.K. and J-H.H. analyzed the metagenomes. L.R. purified the proteins, set up crystallization experiments and edited the manuscript. M.B. conducted LC-MS

experiments. J-H.H. and J.K. wrote the manuscript with input from all authors. **Competing**

interests: All authors declare that they have no competing interests. **Data and materials**

availability: All data are available in the manuscript or the supplementary materials.

List of Supplementary Materials:

Supplementary Text

Figs. S1 to S31

Scheme S1

Table S1 to S6

Data S1 Chemical Synthesis

References and Notes

1. M. D. Purugganan, What is domestication? *Trends in Ecology & Evolution* **37**, 663-671 (2022).
2. M. A. Zeder, Core questions in domestication research. *Proceedings of the National Academy of Sciences* **112**, 3191-3198 (2015).
3. K. M. Olsen, J. F. Wendel, A Bountiful Harvest: Genomic Insights into Crop Domestication Phenotypes. *Annual Review of Plant Biology* **64**, 47-70 (2013).
4. E. V. Armbrust, The life of diatoms in the world's oceans. *Nature* **459**, 185-192 (2009).
5. J. R. Seymour, S. A. Amin, J.-B. Raina, R. Stocker, Zooming in on the phycosphere: the ecological interface for phytoplankton–bacteria relationships. *Nature Microbiology* **2**, 17065 (2017).
6. F. Unfried *et al.*, Adaptive mechanisms that provide competitive advantages to marine bacteroidetes during microalgal blooms. *The ISME Journal* **12**, 2894-2906 (2018).
7. S. Becker, J.-H. Hehemann, Laminarin Quantification in Microalgae with Enzymes from Marine Microbes. *Bio-protocol* **8**, e2666 (2018).
8. N. Barak-Gavish *et al.*, Bacterial lifestyle switch in response to algal metabolites. *Elife* **12** (2023).
9. M. R. Seyedsayamdost, R. J. Case, R. Kolter, J. Clardy, The Jekyll-and-Hyde chemistry of Phaeobacter gallaeciensis. *Nature Chemistry* **3**, 331-335 (2011).
10. E. E. Clerc *et al.*, Strong chemotaxis by marine bacteria towards polysaccharides is enhanced by the abundant organosulfur compound DMSP. *Nature Communications* **14**, 8080 (2023).
11. M. T. Croft, A. D. Lawrence, E. Raux-Deery, M. J. Warren, A. G. Smith, Algae acquire vitamin B12 through a symbiotic relationship with bacteria. *Nature* **438**, 90-93 (2005).
12. M. Uchimiya, W. Schroer, M. Olofsson, A. S. Edison, M. A. Moran, Diel investments in metabolite production and consumption in a model microbial system. *The ISME Journal* **16**, 1306-1317 (2022).
13. D. Muratore *et al.*, Complex marine microbial communities partition metabolism of scarce resources over the diel cycle. *Nature Ecology & Evolution* **6**, 218-229 (2022).
14. W. Petroski, D. M. Minich, Is There Such a Thing as "Anti-Nutrients"? A Narrative Review of Perceived Problematic Plant Compounds. *Nutrients* **12** (2020).
15. Y. Idota *et al.*, Relationship between Physical Parameters of Various Metal Ions and Binding Affinity for Alginate. *Biological and Pharmaceutical Bulletin* **39**, 1893-1896 (2016).
16. J. Li *et al.*, Microbial iron limitation in the ocean's twilight zone. *Nature* **633**, 823-827 (2024).
17. J.-H. Hehemann *et al.*, Adaptive radiation by waves of gene transfer leads to fine-scale resource partitioning in marine microbes. *Nature Communications* **7**, 12860 (2016).
18. C. B. Field, M. J. Behrenfeld, J. T. Randerson, P. Falkowski, Primary Production of the Biosphere: Integrating Terrestrial and Oceanic Components. *Science* **281**, 237 (1998).
19. M. F. White, C. R. Kahn, Insulin action at a molecular level – 100 years of progress. *Molecular Metabolism* **52**, 101304 (2021).
20. J. Monod, From Enzymatic Adaptation to Allosteric Transitions. *Science* **154**, 475-483 (1966).
21. A. Y. Valiakhmetov, A. V. Kuchin, N. E. Suzina, A. N. Zvonarev, A. O. Shepelyakovskaya, Glucose causes primary necrosis in exponentially grown yeast Saccharomyces cerevisiae. *FEMS Yeast Research* **19**, foz019 (2019).
22. G. S. Adams, B. A. Converse, A. H. Hales, L. E. Klotz, People systematically overlook subtractive changes. *Nature* **592**, 258-261 (2021).
23. A. Reichstein, Batman — An American Mr. Hyde? *Amerikastudien / American Studies* **43**, 329-350 (1998).
24. X. Jiang, Chinese Dialectical Thinking—the Yin Yang Model. *Philosophy Compass* **8**, 438-446 (2013).

693   25.   S. Myklestad, O. Holm-Hansen, K. M. Vårum, B. E. Volcani, Rate of release of
694   extracellular amino acids and carbohydrates from the marine diatom Chaetoceros
695   affinis. *Journal of Plankton Research* **11**, 763-773 (1989).

696   26.   L. I. Aluwihare, D. J. Repeta, A comparison of the chemical characteristics of oceanic
697   DOM and extracellular DOM produced by marine algae. *Marine Ecology Progress*
698   *Series* **186**, 105-117 (1999).

699   27.   S. Vidal-Melgosa *et al.*, Diatom fucan polysaccharide precipitates carbon during algal
700   blooms. *Nat Commun* **12**, 1150 (2021).

701   28.   F. Vincent, C. Bowler, Diatoms Are Selective Segregators in Global Ocean Planktonic
702   Communities. *mSystems* **5**, 10.1128/msystems.00444-00419 (2020).

703   29.   H. Teeling *et al.*, Recurring patterns in bacterioplankton dynamics during coastal spring
704   algae blooms. *Elife* **5**, e11888 (2016).

705   30.   D. M. Needham, J. A. Fuhrman, Pronounced daily succession of phytoplankton,
706   archaea and bacteria following a spring bloom. *Nature Microbiology* **1**, 16005 (2016).

707   31.   W.-C. Chin, M. V. Orellana, P. Verdugo, Spontaneous assembly of marine dissolved
708   organic matter into polymer gels. *Nature* **391**, 568-572 (1998).

709   32.   A. Engel, S. Thoms, U. Riebesell, E. Rochelle-Newall, I. Zondervan, Polysaccharide
710   aggregation as a potential sink of marine dissolved organic carbon. *Nature* **428**, 929-
711   932 (2004).

712   33.   J. Uitz, H. Claustre, B. Gentili, D. Stramski, Phytoplankton class-specific primary
713   production in the world's oceans: Seasonal and interannual variability from satellite
714   observations. *Global Biogeochemical Cycles* **24** (2010).

715   34.   S. Malviya *et al.*, Insights into global diatom distribution and diversity in the world's
716   ocean. *Proceedings of the National Academy of Sciences* **113**, E1516-E1525 (2016).

717   35.   J. T. Staley, J. J. Gosink, Poles apart: biodiversity and biogeography of sea ice bacteria.
718   *Annu Rev Microbiol* **53**, 189-215 (1999).

719   36.   B. Avcı, K. Krüger, B. M. Fuchs, H. Teeling, R. I. Amann, Polysaccharide niche
720   partitioning of distinct Polaribacter clades during North Sea spring algal blooms. *The*
721   *ISME Journal* **14**, 1369-1383 (2020).

722   37.   L. Kappelmann *et al.*, Polysaccharide utilization loci of North Sea Flavobacteriia as
723   basis for using SusC/D-protein expression for predicting major phytoplankton glycans.
724   *ISME J* **13**, 76-91 (2019).

725   38.   C. Sidhu *et al.*, Dissolved storage glycans shaped the community composition of
726   abundant bacterioplankton clades during a North Sea spring phytoplankton bloom.
727   *Microbiome* **11**, 77 (2023).

728   39.   F.-Q. Wang *et al.*, Particle-attached bacteria act as gatekeepers in the decomposition
729   of complex phytoplankton polysaccharides. *Microbiome* **12** (2024).

730   40.   T. B. Francis *et al.*, Changing expression patterns of TonB-dependent transporters
731   suggest shifts in polysaccharide consumption over the course of a spring phytoplankton
732   bloom. *The ISME Journal* **15**, 2336-2350 (2021).

733   41.   J. Hevia-Orube *et al.*, Molecular and morphological analyses of solitary forms of
734   brackish Thalassiosiroid diatoms (Coscinodiscophyceae), with emphasis on their
735   phenotypic plasticity. *European Journal of Phycology* **51**, 11-30 (2016).

736   42.   C. Ford, E. Percival, 1299. Carbohydrates of Phaeodactylum tricornutum. Part II. A
737   sulphated glucuronomannan. *Journal of the Chemical Society (Resumed)*, 7042-7046
738   (1965).

739   43.   T. Le Costaouëc, C. Unamunzaga, L. Mantecon, W. Helbert, New structural insights
740   into the cell-wall polysaccharide of the diatom Phaeodactylum tricornutum. *Algal*
741   *Research* **26**, 172-179 (2017).

742   44.   B. Kloareg, Y. Badis, J. M. Cock, G. Michel, Role and Evolution of the Extracellular
743   Matrix in the Acquisition of Complex Multicellularity in Eukaryotes: A Macroalgal
744   Perspective. http://dx.doi.org/10.3390/genes12071059.

745   45.   K. Le Mai Hoang *et al.*, Traceless Photolabile Linker Expedites the Chemical Synthesis
746   of Complex Oligosaccharides by Automated Glycan Assembly. *Journal of the American*
747   *Chemical Society* **141**, 9079-9086 (2019).

46. R. L. Hahnke *et al.*, Dilution cultivation of marine heterotrophic bacteria abundant after
a spring phytoplankton bloom in the North Sea. *Environmental Microbiology* **17**, 3515-
3526 (2015).
47. M. Richter, R. Rosselló-Móra, F. Oliver Glöckner, J. Peplies, JSpeciesWS: a web
server for prokaryotic species circumscription based on pairwise genome comparison.
Bioinformatics **32**, 929-931 (2016).
48. Y. K. Kwon *et al.*, Ochrovirga pacifica gen. nov., sp. nov., a novel agar-lytic marine
bacterium of the family Flavobacteriaceae isolated from a seaweed. *Curr Microbiol* **69**,
445-450 (2014).
49. C. S. Robb, A. A. Mystkowska, J. H. Hehemann, Crystal structure of a marine glycoside
hydrolase family 99-related protein lacking catalytic machinery. *Protein Sci* **26**, 2445-
2450 (2017).
50. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021).
51. A. Varki *et al.*, Symbol Nomenclature for Graphical Representations of Glycans.
Glycobiology **25**, 1323-1324 (2015).
52. T. Barbeyron *et al.*, Matching the Diversity of Sulfated Biomolecules: Creation of a
Classification Database for Sulfatases Reflecting Their Substrate Specificity. *PLOS*
ONE **11**, e0164846 (2016).
53. E. Drula *et al.*, The carbohydrate-active enzyme database: functions and literature.
Nucleic Acids Res **50**, D571-D577 (2022).
54. G. Davies, B. Henrissat, Structures and mechanisms of glycosyl hydrolases. *Structure*
3, 853-859 (1995).
55. A. G. Hettle *et al.*, The Molecular Basis of Polysaccharide Sulfatase Activity and a
Nomenclature for Catalytic Subsites in this Class of Enzyme. *Structure* **26**, 747-
758.e744 (2018).
56. Y. Zhu *et al.*, Mechanistic insights into a Ca2+-dependent family of alpha-
mannosidases in a human gut symbiont. *Nat Chem Biol* **6**, 125-132 (2010).
57. F. Teufel *et al.*, SignalP 6.0 predicts all five types of signal peptides using protein
language models. *Nature Biotechnology* **40**, 1023-1025 (2022).
58. A. Cartmell *et al.*, How members of the human gut microbiota overcome the sulfation
problem posed by glycosaminoglycans. *Proceedings of the National Academy of*
Sciences **114**, 7037-7042 (2017).
59. C. S. Robb *et al.*, Metabolism of a hybrid algal galactan by members of the human gut
microbiome. *Nature Chemical Biology* **18**, 501-510 (2022).
60. K. T. Kim, L. E. Rioux, S. L. Turgeon, Alpha-amylase and alpha-glucosidase inhibition
is differentially modulated by fucoidan obtained from Fucus vesiculosus and
Ascophyllum nodosum. *Phytochemistry* **98**, 27-33 (2014).
61. C. Tara Oceans Consortium, P. Tara Oceans Expedition (2017) Registry of all samples
from the Tara Oceans Expedition (2009-2013). (PANGAEA).
62. J.-H. Hehemann *et al.*, Biochemical and Structural Characterization of the Complex
Agarolytic Enzyme System from the Marine Bacterium Zobellia galactanivorans*.
Journal of Biological Chemistry **287**, 30571-30584 (2012).
63. G. Correc, J. H. Hehemann, M. Czjzek, W. Helbert, Structural analysis of the
degradation products of porphyran digested by Zobellia galactanivorans beta-
porphyranase A. *Carbohyd Polym* **83**, 277-283 (2011).
64. S. Becker, A. Scheffel, M. F. Polz, J. H. Hehemann, Accurate quantification of laminarin
in marine organic matter with enzymes from marine microbes. *Appl Environ Microb* **83**,
e03389 (2017).
65. F. Unfried *et al.*, Adaptive mechanisms that provide competitive advantages to marine
bacteroidetes during microalgal blooms. *The ISME Journal* **12** (2018).
66. V. Smetacek, A. Zingone, Green and golden seaweed tides on the rise. *Nature* **504**,
84-88 (2013).
67. L. Reisky *et al.*, A marine bacterial enzymatic cascade degrades the algal
polysaccharide ulvan. *Nat Chem Biol* **15**, 803-812 (2019).

68. O. I. Nedashkovskaya *et al.*, Formosa agariphila sp. nov., a budding bacterium of the
family Flavobacteriaceae isolated from marine environments, and emended description
of the genus Formosa. *Int J Syst Evol Microbiol* **56**, 161-167 (2006).
69. A. J. Mann *et al.*, The genome of the alga-associated marine flavobacterium Formosa
agariphila KMM 3901T reveals a broad potential for degradation of algal
polysaccharides. *Appl Environ Microbiol* **79**, 6813-6822 (2013).
70. E. V. Armbrust, The life of diatoms in the world's oceans. *Nature* **459**, 185-192 (2009).
71. C. de Vargas *et al.*, Eukaryotic plankton diversity in the sunlit ocean. *Science* **348**,
1261605 (2015).
72. C. d. Vargas *et al.* (2015) First Tara Oceans V9 rDNA metabarcoding dataset.
73. M. Bligh, N. Nguyen, H. Buck-Wiese, S. Vidal-Melgosa, J. H. Hehemann, Structures
and functions of algal glycans shape their capacity to sequester carbon in the ocean.
Curr Opin Chem Biol **71**, 102204 (2022).
74. C. S. Hassler, V. Schoemann, C. M. Nichols, E. C. V. Butler, P. W. Boyd, Saccharides
enhance iron bioavailability to Southern Ocean phytoplankton. *Proceedings of the*
National Academy of Sciences **108**, 1076-1081 (2011).
75. F. X. Ferrer-González *et al.*, Resource partitioning of phytoplankton metabolites that
support bacterial heterotrophy. *The ISME Journal* **15**, 762-773 (2021).
76. S. Becker *et al.*, Laminarin is a major molecule in the marine carbon cycle. *Proc Natl*
Acad Sci U S A **117**, 6599-6607 (2020).
77. J. Kurreck, J. W. Engels, F. Lottspeich, *Bioanalytik* (Springer Spektrum Berlin,
Heidelberg, 2021), vol. 4.
78. H. M. Berman *et al.*, The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242 (2000).
79. K. M. Wheeler *et al.*, Mucin glycans attenuate the virulence of Pseudomonas
aeruginosa in infection. *Nature Microbiology* **4**, 2146-2154 (2019).
80. C. Cisternas-Novoa, C. Lee, A. Engel, Transparent exopolymer particles (TEP) and
Coomassie stainable particles (CSP): Differences between their origin and vertical
distributions in the ocean. *Marine Chemistry* **175**, 56-71 (2015).
81. E. A. Pelve, K. M. Fontanez, E. F. DeLong, Bacterial Succession on Sinking Particles
in the Ocean's Interior. *Frontiers in Microbiology* **8** (2017).
82. S. Thiele, M. Fuchs Bernhard, R. Amann, H. Iversen Morten, Colonization in the Photic
Zone and Subsequent Changes during Sinking Determine Bacterial Community
Composition in Marine Snow. *Appl Environ Microb* **81**, 1463-1471 (2015).
83. H. Teeling *et al.*, Substrate-Controlled Succession of Marine Bacterioplankton
Populations Induced by a Phytoplankton Bloom. *Science* **336**, 608-611 (2012).
84. D. M. Needham *et al.*, Dynamics and interactions of highly resolved marine plankton
via automated high-frequency sampling. *The ISME Journal* **12**, 2417-2432 (2018).
85. S. J. Giovannoni, SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Annual*
Review of Marine Science **9**, 231-255 (2017).
86. A. Dadon-Pilosof *et al.*, Surface properties of SAR11 bacteria facilitate grazing
avoidance. *Nature Microbiology* **2**, 1608-1615 (2017).
87. J. Sun *et al.*, One carbon metabolism in SAR11 pelagic marine bacteria. *PLoS One* **6**,
e23973 (2011).
88. B. J. Shapiro *et al.*, Population genomics of early events in the ecological differentiation
of bacteria. *Science* **336**, 48-51 (2012).
89. A. S. Utada *et al.*, Vibrio cholerae use pili and flagella synergistically to effect motility
switching and conditional surface attachment. *Nature Communications* **5**, 4913 (2014).
90. Y. Yawata *et al.*, Competition-dispersal tradeoff ecologically differentiates recently
speciated marine bacterioplankton populations. *Proc Natl Acad Sci U S A* **111**, 5622-
5627 (2014).
91. K. Moor *et al.*, High-avidity IgA protects the intestine by enchaining growing bacteria.
Nature **544**, 498-502 (2017).
92. J. M. Grondin, K. Tamura, G. Dejean, D. W. Abbott, H. Brumer, Polysaccharide
utilization loci: fueling microbial communities. *J Bacteriol* **199**, e00860-00816 (2017).
93. N. M. Koropatkin, E. A. Cameron, E. C. Martens, How glycan metabolism shapes the
human gut microbiota. *Nature Reviews Microbiology* **10**, 323-335 (2012).

94. R. B. Cassie *et al.,* Carbohydrate-active enzymes from Akkermansia
muciniphila breakdown mucin O-glycans to completion. *bioRxiv*
10.1101/2024.03.27.586211, 2024.2003.2027.586211 (2024).

95. O. Gainza, J. Romero, Effect of mannan oligosaccharides on the microbiota and
productivity parameters of Litopenaeus vannamei shrimp under intensive cultivation in
Ecuador. *Scientific Reports* **10**, 2719 (2020).

96. A. M. Martin-Platero *et al.*, High resolution time series reveals cohesive but short-lived
communities in coastal plankton. *Nature Communications* **9**, 266 (2018).

97. G. Reintjes *et al.*, Short-term changes in polysaccharide utilization mechanisms of
marine bacterioplankton during a spring phytoplankton bloom. *Environmental*
Microbiology **22**, 1884-1900 (2020).

98. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nature*
Methods **9**, 357-359 (2012).

99. P. J. Harrison, R. E. Waters, F. J. R. Taylor, A BROAD SPECTRUM ARTIFICIAL SEA
WATER MEDIUM FOR COASTAL AND OPEN OCEAN PHYTOPLANKTON1. *Journal*
of Phycology **16**, 28-35 (1980).

100. A. Engel, N. Händel, A novel protocol for determining the concentration and
composition of sugars in particulate and in high molecular weight dissolved organic
matter (HMW-DOM) in seawater. *Marine Chemistry* **127**, 180-191 (2011).

101. P. J. Shilling *et al.*, Improved designs for pET expression plasmids increase protein
production yield in Escherichia coli. *Communications Biology* **3**, 214 (2020).

102. J. J. Almagro Armenteros *et al.*, SignalP 5.0 improves signal peptide predictions using
deep neural networks. *Nature Biotechnology* **37**, 420-423 (2019).

103. I. S. Carrico, B. L. Carlson, C. R. Bertozzi, Introducing genetically encoded aldehydes
into proteins. *Nat Chem Biol* **3**, 321-322 (2007).

104. V. Solanki *et al.*, Glycoside hydrolase from the GH76 family indicates that marine
Salegentibacter sp. Hel_I_6 consumes alpha-mannan from fungi. *ISME J* **16**, 1818-
1830 (2022).

105. J. H. Hehemann *et al.*, Transfer of carbohydrate-active enzymes from marine bacteria
to Japanese gut microbiota. *Nature* **464**, 908-912 (2010).

106. A. J. McCoy *et al.*, Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674
(2007).

107. K. Cowtan, The Buccaneer software for automated model building. 1. Tracing protein
chains. *Acta Crystallogr D Biol Crystallogr* **62**, 1002-1011 (2006).

108. G. N. Murshudov *et al.*, REFMAC5 for the refinement of macromolecular crystal
structures. *Acta Crystallogr D Biol Crystallogr* **67**, 355-367 (2011).

109. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501 (2010).

110. R. L. Hahnke, J. Harder, Phylogenetic diversity of Flavobacteria isolated from the North
Sea on solid media. *Syst Appl Microbiol* **36**, 497-504 (2013).

111. M. Robb, J. K. Hobbs, A. B. Boraston, Separation and Visualization of Glycans by
Fluorophore-Assisted Carbohydrate Electrophoresis. *Methods Mol Biol* **1588**, 215-221
(2017).

112. C. J. Crawford *et al.*, Automated Synthesis of Algal Fucoidan Oligosaccharides. *Journal*
of the American Chemical Society **146**, 18320-18330 (2024).

113. C. Crawford, S. Oscarson, Optimized Conditions for the Palladium-Catalyzed
Hydrogenolysis of Benzyl and Naphthylmethyl Ethers: Preventing Saturation of
Aromatic Protecting Groups. *European Journal of Organic Chemistry* **2020**, 3332-3337
(2020).

114. O. J. Plante, E. R. Palmacci, P. H. Seeberger, Automated Solid-Phase Synthesis of
Oligosaccharides. *Science* **291**, 1523-1527 (2001).

115. O. J. Plante, R. B. Andrade, P. H. Seeberger, Synthesis and Use of Glycosyl
Phosphates as Glycosyl Donors. *Organic Letters* **1**, 211-214 (1999).