Modelling determinants of region-specific dopamine dynamics in the striatum

Aske L. Ejdrup[1, †], Jakob K. Dreyer[2], Matthew D. Lycas[1], Søren H. Jørgensen[1], Trevor W. Robbins[3,4], Jeffrey W. Dalley[3,4,5], Freja Herborg[1, †, *], Ulrik Gether[1, †, *]

[1]Department of Neuroscience, Faculty of Health and Medical Sciences, Maersk Tower 7.5, University of Copenhagen, DK-2200 Copenhagen, Denmark

[2]Department of Bioinformatics, H Lundbeck A/S, Valby, Denmark

[3]Behavioural and Clinical Neuroscience Institute, University of Cambridge, Cambridge CB2 3EB, UK

[4]Department of Psychology, University of Cambridge, Cambridge CB2 3EB

[5]Department of Psychiatry, University of Cambridge, Cambridge CB2 0SZ, UK

* These authors contributed equally to the work

† To whom correspondence should be addressed: ejdrup@sund.ku.dk, frejahh@sund.ku.dk or gether@sund.ku.dk

Abstract

Striatal dopamine (DA) release regulates reward-related learning, motivation, and behavioural activation, and is believed to consist of a short-lived *phasic* and continuous *tonic* component. Here, we build a large-scale three-dimensional model of extracellular DA dynamics in the dorsal (DS) and ventral striatum (VS) based on experimentally determined biological parameters. The model predicts rapid dynamics in DS with little-to-no basal DA, whereas lower uptake capacity in VS slows signalling and enables build-up of a *tonic* DA level. Receptor binding simulations reveal that DA D1 receptor occupancy follows extracellular DA concentration with millisecond delay, while DA D2 receptors integrate DA signal over seconds. The modelling further predicts that VS extracellular levels are highly sensitive to DA transporter (DAT) uptake capacity and posit prevalent nanoclustering of DAT in VS as a regulator of DA uptake. Our simulations substantiate regional differences in extracellular DA dynamics and challenge prevailing paradigms of striatal DA signalling.

Introduction

Striatal dopamine (DA) release is essential for regulating reward-related learning, incentive motivation, and motor function[1,2]. DA exerts these roles over a broad range of time scales, yet DA primarily operates as a volume transmitter that targets metabotropic receptors located within a micrometre range from the sites of release[3–7]. The temporal and spatial dynamics of DA release in striatum, however, remain a highly contested topic. Classically, DA release has been divided into *tonic* release, driven by pacemaker-like spontaneous firing, and *phasic* release from coordinated bursts of firing across neurons[5,8,9]. However, this sharp distinction in release modes, as well as the existence of a basal DA level has recently been challenged[1,10–13].

The picture is further complicated by major regional differences across striatal sub domains. These include differences in Ca^{2+}-channel and nicotinic acetylcholine receptor (nAChR) expression profiles on DA terminals, as well differential regulation and expression of the DA transporter (DAT)[14–18]. In addition, we and others have found remarkable differences in extracellular DA release dynamics between the dorsal (DS) and ventral striatum (VS)[12,19,20]. Fibre photometry recordings in the DS in mice using the DA sensor dLight1.3b during self-paced exploratory activity showed a rapidly fluctuating signal, whereas we observed up to minutes-long DA dynamics in VS that correlated with behavioural output[12]. Concurrent measurements of extracellular DA by microdialysis and fibre photometry have furthermore corroborated the lack of tonic levels of DA in DS while supported its presence in VS[12 13]. Despite these reported differences in striatal DA dynamics, electrophysiological recordings suggest that DA neurons from the primary innervators of DS, substantia nigra par compacta (SNc) and VS, ventral tegmental area (VTA), have remarkably similar firing patterns at rest[21]. We therefore set out to better understand the fundamental

principles governing extracellular DA dynamics by constructing a new computational model

of the striatal DA system.

Extracellular DA dynamics have been modelled before; either one-dimensionally or

with a primary focus on single release events or post-synaptic receptor binding[22–27]. Here, we

present a three-dimensional model of tens of thousands of release sites, focused on larger-

scale signalling and based on experimentally observed biological parameters. The model

faithfully replicates experimentally observed results as well as the difference in DA dynamics

between DS and VS. Importantly, it offers compelling evidence that these differences do not

primarily reflect different release phenomena but rather arise from differential expression and

possibly nanoscale localization of the DA transporter (DAT).

Results

Construction of a model of DA dynamics in the striatum

We constructed a novel model of DA release using experimentally determined parameters from DS, including release, uptake, and cytoarchitecture[5,10,22,23,28,29]. DA release sites on axons projecting from the midbrain were randomly simulated as uniformly distributed discrete points in a three-dimensional space (Fig. 1a). The release events themselves were simulated as point source events[4] driven by action potentials (AP). We then modelled DA release for each voxel in the simulation containing a release site as function of three key parameters; firing rate, release probability and quantal size:

(1)

$$\text{release}_{n,t} = \text{Poisson}(f_{rate}\, dt)_n\, P(R_\%)_t\, Q\, dt$$

where $\text{Poisson}(f_{rate}\, dt)_n$ is a Poisson distribution of action potentials (AP) for the given neuron (n) with the firing rate f_{rate}[23], $P(R_\%)_t$ is the probability of release at the individual terminal (t) for each AP, while Q is the number of DA molecules released per event (dopaminergic quantal size) and dt the time step. Changing f_{rate} can be used to model both pacemaker firing, typically reported at 2-10 Hz, and burst firing, which can exceed 20 Hz[5].

DA reuptake in the striatum is almost exclusively mediated by the DA transporter (DAT)[30], which is widely distributed along DA axons and varicosities[31–33]. As reuptake follows concentration-dependent Michaelis-Menten kinetics[34], we simulated uptake as follows:

93   (2)

94
$$\text{uptake} = \frac{V_{max}\ [DA]}{K_m\ +\ [DA]}\ dt$$

where [DA] is the concentration of DA for each voxel in the model, V_{max} is the maximal

uptake capacity in the region and K_m is the concentration of DA at which half of V_{max} is

reached.

The spatial distribution of released DA is a complex interplay between release, uptake

and diffusion. Diffusion in an open 3D space can be simulated for each voxel with a

Laplacian operator:

(3)

102
$$\text{diffusion} = \frac{\partial DA_{x,y,z,t}}{\partial t} = D_a\ dt \left(\frac{\partial^2 DA_{x,y,z,t}}{\partial x^2} + \frac{\partial^2 DA_{x,y,z,t}}{\partial y^2} + \frac{\partial^2 DA_{x,y,z,t}}{\partial z^2} \right)$$

103   where D_a is a corrected diffusion coefficient and *dt* is the timestep. As the extracellular space

104   of the striatum is tortuous, we modified the conventional diffusion coefficient *D* to an

105   apparent diffusion coefficient (D_a) to correct for the tortuosity (λ) of the striatum[4,35]:

106   (4)

107
$$D_a = \frac{D}{\lambda^2}$$

108   Combining equation 1 through 4, we model DA changes in each voxel with a single

109   conceptual equation:

110   (5)

111
$$\frac{dDA}{dt} = \text{release} - \text{uptake} + \text{diffusion}$$

112   We first compared our 3D model of DA dynamics to the analytical solution of a single

113   release event[3,4]. To do this, we simulated a quantal event of 3000 DA molecules and

114   calculated DA concentrations across space at three separate time points (Fig. 1b, c). The

115   analytic solution and our model predicted almost identical results. Slight differences were

116 introduced as the analytical solution assumes linear uptake from DAT while our model

117 incorporate non-linear Michaelis-Menten kinetics. These differences, however, were almost

118 negligible. The main difference between the two models lies in scalability across both space

119 and time. Summarized, the model enables a dynamic incorporation of the surrounding DA

120 concentration, release events and uptake, and can be scaled to cover DA dynamics of a large

121 3D space, whose size and granularity is only limited by computing power (see Code

122 Availability Section for the Python code with numerical implementations of the equations

123 listed and scripts to run the simulations and plot the main figures).

124   We also tested the validity of our model, by examining the response to electrical

125 stimulation. Importantly, our model faithfully mirrored DA release seen with fast-scan cyclic

126 voltammetry (FSCV) recordings upon direct stimulation of striatal slices when we corrected

127 for kinetics of the typical FSCV recording setup (Extended Data Fig. 1b, c and discussion in

128 Supplementary Information)[36–39].

130 Simulating large scale DA dynamics of the dorsal striatum

131 To better understand the extracellular DA dynamics that arise from the balance between

132 dopaminergic pacemaker activity and uptake, we simulated DA dynamics in the DS

133 generated by pacemaker activity (4 Hz) of 150 neurons in the midbrain in a 100x100x100 μm

134 space (Fig. 1d, see parameters in Table 1). Our simulations yielded a pattern of partially

135 segregated DA hot spots with large fractions of the simulated space devoid of DA, suggesting

136 that release events in DS only elevate DA in the immediate surroundings with DAT-

137 dependent clearance preventing a larger spread in space (Fig. 1d). This was also illustrated by

138 a cross-section in time (Fig. 1e). In line with our recent in vivo microdialysis experiments, the

139 average DA concentration in the simulations during pacemaker activity was approximately

140 10 nM[40]. Further, when we calculated the average concentration of a larger area across time,

which fibre photometry conceivably does, the results resembled a tonic DA concentration

(Extended Data Fig. 1a). However, our model predicted a spatial distribution that is highly

heterogenous and devoid of pervasive resting or tonic DA levels (Fig. 1d-f).

To ensure our simulations were performed within a sufficiently large space to yield

consistent results, we tested different sizes of the simulated area and found a diameter of 50

µm to faithfully mimic the results of larger simulations (Extended Data Fig. 2a, b).

Additionally, we tested our simulations at different granularity (0.1, 0.5, 1 and 2 µm). The

finer the spatial grain, the higher detail close to a release event; however, at a spatial

granularity of 1 µm, [DA] deviated by <2% across most percentiles and only by >1 nM above

the 99.5th percentile (Extended Data Fig. 2c-f), leading us to use this voxel size for our

simulations.

Burst firing and receptor occupancy

DA cell bodies are known to fire short bursts of APs, which is a phenomenon strongly linked

to reward-prediction error and learning[8]. To gain insights into accompanying extracellular

DA dynamics, we simulated three different firing scenarios for a group of terminals within a

10x10x10 µm field encompassing roughly 40 release sites from the randomly simulated 150

neurons: 3 pulses at 10 Hz, 6 pulses at 20 Hz and 12 pulses at 40 Hz (Fig. 1g – burst

properties matched to be the same duration). Unsurprisingly, peak DA concentration was

reached at the end of the bursts (Fig. 1g). The 3 APs/10 Hz bursting scenario generated no

significant spill-over of DA outside the region of activity, whereas the 6 APs/20 Hz and 12

APs/40 Hz bursting scenarios markedly overwhelmed uptake (Fig. 1g). The relationship

between firing rate and the sphere of influence by DA became further evident when plotting

maximal concentration of the surrounding space (Fig. 1h) and the volume of space with a DA

concentration above 100 nM (Extended Data Fig. 1d). We found that the 3 APs/10 Hz

stimulation produced DA responses that largely resembled that of a single pulse. In both

cases, DA was mostly cleared after 100 ms and the volume exposed to greater than 100 nM

was similar (Fig. 1g and Extended Data Fig. 1d). In contrast, the high bursting activities

caused a frequency-dependent spill-over, where the areas exposed to a DA concentration

above 100 nM were 10 and 30 times larger than the terminal origin for 6 APs/20 Hz and 12

APs/40 Hz, respectively. Even after 100 ms, a considerable amount of DA remained in the 12

APs/40 Hz scenario (Fig. 1g).

To understand how these DA dynamics could affect the postsynaptic response, we

modelled receptor binding. D1 receptors (-Rs) were assumed to have a half maximal effective

concentration (EC_{50}) of 1000 nM and we extrapolated the reverse rate constant (K_{off}) to 19.5

s^{-1} based on a linear fit of the recently characterized DA receptor-based sensors (Extended

Data Fig. 1e) [41]. We set the EC_{50} of D2Rs to 7 nM and K_{off} to 0.2 s^{-1} based on the DA sensor

kinetic fit (Extended Data Fig. 1e), which matches a recent binding study[42]. To determine

how these receptors would respond to our predicted DA dynamics, we simulated pacemaker

activity at 4 Hz with an added burst of 6 APs/20 Hz. Fig. 1i shows a representative trace of

DA concentration and occupancy of the D1R and D2R within 1 μm of a release site. During

pacemaker activity, D1R showed an occupancy close to 0, whereas D2R occupancy was

approximately 0.65 (Fig. 1i). Both D1Rs and D2Rs occupancies were largely invariant to

individual release events caused by pacemaker activity. However, upon coordinated burst

firing, the occupancy rapidly increased (Fig. 1i). D1R receptor occupancy closely tracked

extracellular DA with a delay of only ~50 ms (Extended Data Fig. 1f). In contrast, it took at

least 5 seconds before the burst-induced increase in D2R occupancy had declined to baseline

levels (Fig. 1i), and, perhaps more surprisingly, the D2R occupancy only fell from 0.55 to

0.45 when simulating a full second pause in firing due to the slow off kinetics (Fig. 1j).

These simulations suggest that the dopaminergic architecture of the DS limits DA

overflow during physiologically relevant bursting activity. Further, DA receptors had a

temporally mostly uniform response to DA release caused by pacemaker activity with D1R

occupancy responding rapidly to both onset and offset extracellular DA concentrations

following bursts, while D2R showed seconds-long delays in offset.

Ventral striatum maintains pervasive DA tone

Mounting evidence points to considerable differences in DA dynamics across striatal

subregions[12,43], which might reflect differences in the cytoarchitectural and/or molecular

dopaminergic make up. In line with this, most studies report lower dopaminergic density in

the VS than in DS regardless of methodological modality with a median value of ~90% in VS

relative to DS (Extended Data Table 1). Further, DAT-mediated uptake capacity is reported

to be lower in VS with a median capacity at ~30% of DS (Extended Data Table 2).

Consistently, we observed a clear dorsoventral gradient for DAT expression when analysing

immunostainings in striatal mouse brain slices from a previous publication (Sørensen et al.,

2021[44]) (Extended Data Fig. 3a, b). By contrast, the VMAT2 staining only decreased slightly

from VS to DS (Extended Data Fig. 3a-c).

We simulated DA release during pacemaker activity in both DS and VS. DS values

were set as previously described (25 μm^3 per terminal, uptake capacity of 6.0 $\mu M\ s^{-1}$), but for

VS we reduced the terminal density to 90% (27.8 μm^3 per terminal) and DAT uptake

capacity to 33% (2.0 $\mu M\ s^{-1}$) (Extended Tables 1 and 2). The remaining parameters were kept

identical. With these two differences, our simulations revealed markedly different

spatiotemporal DA distributions during pacemaker activity. While DS formed segregated

domains with low DA concentrations in the inter-domain space (Fig. 2a), DA diffused further

throughout the simulated space in VS, before being cleared by DAT. This gave rise to what

may be considered a tonic DA level with hotspots of higher DA concentrations (Fig. 2a, b).

We compared our model of the two regions with existing experimental data. In an

earlier study by May and Wightman, 120 stimulus pulses were delivered in the medial

forebrain bundle (MFB) at either 10, 30 or 60 Hz and DA responses were recorded by FSCV

in both caudate-putamen (CPu) and nucleus accumbens (NAc)[45]. To mirror this, we simulated

120 action potentials at similar frequencies (10, 30, and 60 Hz) at 6% release probability and

ran the result through convolution, as in Extended Data Fig. 1b and c, to generate an FSCV

read-out (Fig. 2e). Since May and Wightman reported no significant difference in DA

released per electrically delivered pulse ($[DA]_p$) between VS and DS, we applied equal

quantal size and $R_\%$ for DS and VS in our simulations, while uptake capacity in VS was kept

to a third of DS and terminal density was set to 90% as specified above. Importantly, our

simulated FSCV data closely resembled the earlier findings, with VS reaching considerably

higher DA levels for all three stimulation frequencies (Fig. 2e). This regional difference

presumably arises from differences in DAT capacity between DS and VS, as the lower

terminal density in VS would have the opposite effect (see below) and the remaining

parameters were held identical.

To compare with our results for DS, we tested how VS responded during simulated

burst activity. Using firing patterns identical to the DS simulations (Fig. 1g), we found a

larger spill-over of DA into the surrounding areas in VS (Fig. 2f and Extended data Fig. 3d,

e). Significant amounts of extracellular DA also remained 100 ms after the physiologically

relevant 6APs/20Hz firing stimulus. At the receptor level, D1R occupancy in VS showed a

similar response to that in DS during the burst (Fig. 1i and 2g). By contrast, D2R occupancy

during pacemaker activity was higher in VS than DS (~0.8 versus ~0.65 in DS), in

accordance with the higher prevailing basal DA concentration. Additionally, the larger DA

overflow in VS after a burst caused a higher relative increase in receptor occupancy further

away from the area actively bursting than compared to DS (Fig. 2h). A pause in firing had the

same effect on D2R as in DS (Extended Data Fig. 3f).

Testing sensitivity of model parameters

To test the robustness of the results, we performed simulations across wide ranges of the

variable key parameters on which the model is based. First, we varied the number of

varicosities actively releasing DA by setting the varicosity density to one site per 9 µm^3 [28]

and simulating 4 Hz pacemaker activity with the release fraction ranging from 5% to 100%

(reported values range from 20% to virtually all)[10,46–48] (Fig. 3a). As the fraction of active

sites increased, DA concentrations increased at both the median level (50th percentile), which

we consider a measure of tonic or baseline DA levels, and at peak levels (99.5th percentile) in

both DS and VS (Fig. 3b, see Extended Data Fig. 4a for schematic of tonic and peak DA).

We then used the 99.5th/50th percentile ratio as a measure of the focality of the DA

distribution (i.e. hotspot DA relative to baseline DA). Quantifying this across the percentage

of active terminals showed that the focality of the DA distribution dropped as the active

fraction increased in both regions (Fig. 3c). However, the percentage of active sites in VS

needed to drop to 5% to reach a relative distribution resembling the DS at a full 100% active

sites, underscoring a marked difference in the spatial confinement of DA signals in VS and

DS.

The predicted total DA content of a vesicle and the fraction of content released per

fusion event is reported to range from 1,000 to 30,000 molecules[49–52] (Fig. 3d). As expected,

tonic and peak concentrations increased in both DS and VS as quantal size was increased

(Fig. 3e). Also as expected, the focality of the DA distributions dropped for both regions as

quantal size increased (Fig. 3f). The relative difference in tonic DA, however, remained

persistently higher in VS and even increased as quantal size increased, indicating a tendency

for VS to maintain basal levels of DA regardless of release content (Fig. 3g). The higher end

of quantal sizes, however, resulted in median concentrations far beyond what is typically

reported (Fig. 3e)[5]. We observed largely similar pattern when changing either release

probability or firing rate (Extended Data Fig. 4b-g).

DAT activity is governed by two parameters: K_m and V_{max}[53]. To mimic competitive

inhibition of DAT by e.g., cocaine, we ran a simulation across various K_m values (Fig. 3h)

showing that increasing K_m caused a linear increase in DA levels, consistent with DAT

uptake rate responding almost linearly to increases in [DA] below K_m (Fig. 3i). Of note, most

microdialysis studies have reported that cocaine increases [DA] to the same degree in both

DS and VS; however, these quantifications are usually derived as a ratio of the absolute

baseline level[54,55]. If we divide our simulations of increasing K_m with the basal levels

estimated in Fig. 3a-c a similar response for DS and VS is found (Extended Data Fig. 5a).

Further, we observe a convergence on a two-fold difference in the absolute values at both

tonic and peak levels, which matches reports from earlier FSCV studies (Extended Data Fig.

5b)[56]. This regionally differential response to cocaine matches our observations in a previous

biosensor-based study[12].

Finally, we changed V_{max} by $\pm$ 50% in both regions and observed a smaller change in

for tonic level in DS (11 nM) than in VS (38 nM) (Fig. 3k). This suggests modulation of V_{max}

has higher impact in VS than DS. Further, the cusp-like behaviour of V_{max} in VS was

independent of both Q and $R_{\%}$ within values typically reported in the literature (Extended

Data Fig. 5c, d). In contrast to the changes in tonic levels, the relative effect V_{max} had on peak

levels was much more modest (Fig. 3l).

Changes to uptake rate may be mediated by DAT internalization pathways but to our

knowledge there is no in vivo evidence of differential release-uptake balances between

animals that could lead to varying tonic DA levels across animals. We therefore reanalysed

data from our previously published comparison of fibre photometry and microdialysis and

found evidence of natural variations in the release-uptake balance of the mice (Extended Data

Fig. 5f, g), which may underlie different tonic levels of DA in the striatum between animals.

Simulations of DAT nanoclustering

Our simulations highlight DAT V_{max} as an effective regulator of extracellular DA levels in

VS (Fig. 3k). Internalization of DAT can serve as a mechanism for this control but is a

relatively slow process operating on the order of minutes[53]. Interestingly, our recent studies

have provided evidence that DAT move laterally in the plasma membrane and transition from

a clustered to an unclustered nanoscale distribution in response to excitatory drive and other

inputs[57,58]. This led us to hypothesize that DAT nanoclustering serve as a mechanism for

regulating DAT activity on a faster time scale. We speculated that dense nanoclusters of DAT

would produce domains of low [DA] due to uptake overpowering diffusion (Fig. 4a). As

uptake rate is concentration dependent, this would reduce uptake efficiency (Fig. 4b). To

address this hypothesis, we simulated a single ellipsoid varicosity of 1.5 μm in length and

800 nm thick with surrounding extracellular space[48]. The surface was unfolded to a square of

equal area (Fig. 4c), and as 9-16.4% of terminals in the striatum are estimated to be

dopaminergic (Hökfelt, 1968; Tennyson et al., 1974), we set the volume of the surrounding

space to seven times the varicosity volume. On the surface of the varicosity, we randomly

distributed eight DAT nanoclusters (Fig. 4c) and ran simulations of how DAT clustering

density influenced the DA clearance from the surrounding space. The observed DA

concentration in the space surrounding the varicosity shown in Fig. 4c is illustrated by the

cross-section shown in Fig. 4d. Mean DA uptake capacity of the entire space was kept

constant at 4 μM s^{-1} (between the values observed for DS and VS) throughout the

simulations, representing a constant amount of DAT molecules on the surface of the

varicosity. We only changed the fraction of the surface of the varicosity that was uptake

competent by altering the cluster size from small clusters of high density to large clusters of

lower density. We ran simulations of eight clusters at 20, 40, 80 or 160 nm in diameter to

mirror experimentally observed cluster sizes on DA varicosities, as well as a scenario with

DAT fully dispersed (Fig. 4e)[57,58]. We then set the extracellular space to a [DA] of 100 nM

and simulated uptake for the four cluster sizes and an unclustered distribution of DAT. As

shown in Fig. 4f, the distribution of DAT into clusters of increasing density resulted in

progressively slower DA clearance. Thus, the unclustered scenario reduced [DA] to 10 nM in

just above 200 ms, whereas the 20 nm clusters took ~400 ms. As cluster size increased and

DAT density decreased, clearance time approached that of the fully unclustered scenario (Fig.

4g). The resulting hyper-local low-[DA] environment that came from dense DAT clusters

was apparent when we plotted the [DA] at the centre of a cluster compared to the mean [DA]

of the extracellular space (Fig. 4h). For the 20 nm cluster-scenario, the clearance was almost

entirely limited by how quickly DA diffused to the nanocluster, as the local [DA] dropped to

near zero. We originally hypothesized the effect would mainly be due to a depression in [DA]

at the cluster centre, but a concentration profile of the 80 nm cluster-scenario revealed the

entirety of the cluster was enveloped in a low-[DA] environment (Fig. 4i).

Summarized, the data supports that dense DAT nanoclusters produce domains of low

[DA] as uptake outcompetes diffusion. With transporter uptake being concentration

dependent, uptake efficiency is reduced. The simulations therefore posit DAT nanoclustering

as an efficient way to regulate the extracellular levels of tonic DA, particularly in VS.

DAT clusters more in the ventral striatum

As our simulations suggest that nanodomain clustering is a way of regulating uptake, and this may be particularly efficient at controlling extracellular DA in VS, we hypothesized nanoscale clustering would be more prominent in VS. To investigate this, we reanalysed data from a previous publication[44], where we acquired super-resolution images of coronally sliced striatal sections from mice stained for DAT. For the present study, images were split in DS and VS based on where in the slices they were taken (Fig. 4j). DA terminals were identified by vesicular monoamine transporter 2 (VMAT2) expression, and qualitatively DAT appeared more clustered in VS than DS (Fig. 4k). To quantify this, we applied the clustering algorithm Density-Based Spatial Clustering of Applications with Noise (DBSCAN) (Fig. 4l). This confirmed a regional difference with more DAT localized to mean cluster size of 80 nm in VS compared to DS (Fig. 4m), and we observed a similar regional difference across the range of sizes typically reported (Fig. 4n)[57,58], supporting the conclusion that DAT nanoclustering is more prevalent in VS than DS.

Discussion

We developed a three-dimensional, finite-difference computational model to investigate

spatiotemporal DA dynamics of striatal subregions in detail. Leveraging prior experimental

information on regional differences in dopaminergic innervation density and DAT uptake

capacity, our model predicts important differences in dopaminergic dynamics between DS

and VS. Strikingly, our simulations suggest that large areas of the DS are effectively devoid

of a basal level of DA at pacemaker activity, whereas VS maintain a more homogenous tonic-

like basal DA concentration with only small changes in uptake activity powerfully regulating

the extracellular DA tone. Furthermore, we modelled receptor binding kinetics and found that

D1R binding faithfully followed recently described rapid DA dynamics of the striatum[12,13,61],

while D2R, with an off-rate of ~5 seconds, appeared better suited for detecting background

tone, ramps and integrating prolonged activity[12,62]. Collectively, these observations have

important implications for our understanding of striatal function in behaviour, including

decoding of inputs from the prefrontal cortex (PFC) and limbic system as well as the

influential phasic-tonic model of dopaminergic signalling[8,9,63].

It has been assumed for long that there are tonic levels of DA in the striatum[5,8,9],

although the phenomenon has no clear definition[1]. Our simulation of DA dynamics in DS

during pacemaker-activity showed no evidence for a homogenous extracellular distribution.

Rather, elevated [DA] was transiently present around release sites during pacemaker activity

with the remaining space mostly depleted of DA. The absence of a general tonic DA level in

DS predicted by our model directly supports the notion that DA release sites in DS establish

distinct and only partially overlapping DA domains rather than diffuse, tonic DA levels (see

Liu et al., 2021[10]). This conclusion aligns with recent data where we found that DA

concentrations measured by microdialysis correlates with the average of rapid activity

recorded with fibre photometry rather than a baseline DA tone[13].

In contrast to the DS, our model predicted VS to hold a considerable basal level of DA even in spaces without an immediately adjacent release site. This is conceivably what most refer to as tonic DA. In this study, we quantified tonic DA as the median concentration of the entire space (50th percentile), which appeared significantly higher in VS than DS because of the lower VS uptake capacity. Importantly, this matches the results of our direct in vivo comparison of DS and VS in freely moving mice[12]. To challenge our model, we performed our simulations across a wide range of parameters. Only changes to V_{max} for uptake generated differential responses in the two regions. With release and uptake parameters at values from the literature, VS was at a critical point where minor changes to uptake significantly impacted the tonic levels without any major effect on peak concentrations.

Contrary to our observations, some previous microdialysis experiments have suggested higher basal DA levels in the DS compared to VS (Kuczenski & Segal, 1992; Shen et al., 2004 have reported 2-4 times higher [DA] in DS compared to VS[64,65]). However, there are disparate observations in the literature (e.g. Carboni et al., 2001 reported 20% higher [DA] in VS[54]). Moreover, it is important to note that regional comparisons in microdialysis might be confounded by the considerably higher uptake rate in DS. This will increase the extraction fraction and possibly lead to a significant overestimation of the extracellular concentration as compared to a region with lower uptake rate, such as the VS[66].

Our three-dimensional simulations highlight DAT-mediated reuptake as a key mechanism governing striatal DA dynamics and as a key mediator of regional-specific DA dynamics. A physiologically relevant way to regulate uptake capacity is moving DAT to-and-from the plasma membrane. Indeed, DAT is subject to such regulation and some of these mechanisms may even be exclusive to the ventral region, including protein kinase C-induced DAT internalization and Vav2 regulation of DAT surface expression[67,68]. Chemogenetic Gq-

coupled DREADD activation of DA neurons also result in differential DAT trafficking in the

two regions[69]. The findings position DAT regulation as an excellent candidate for changing

tonic DA levels in VS, which has been proposed to selectively attenuate afferent drive from

the PFC through D2R activation[63]. Recent studies of D2R expressing spiny projection

neurons (SPNs) in the VS also suggest that the receptor is not fully saturated under basal

firing[70], matching both our simulations of receptor binding and the notion that tonic DA can

be manipulated to alter D2R activation.

If changing uptake capacity is to have a behavioural relevance on a fast timescale, a

mechanism to regulate DAT function faster than internalization must exist. Importantly, the

transporter does not only move to-and-from the surface, but also laterally in the plasma

membrane. We have reported that DAT forms nanoclusters in the plasma membrane that

dynamically reshape based on excitatory and inhibitory input[57,58]. Moreover, we have

previously shown that cocaine, which both competitively inhibits DAT and reorganizes the

transporter nanodomains[58], changes the DA signal of the DS to dynamics akin to the VS[12].

Importantly, our simulations showed that nanoclustering may be an effective way to

sequester DAT in a dense domain where uptake overpowers diffusion and as a result brings

down effective uptake speed through microdomain DA depletion. We also found that the

nanoclustering phenomenon was considerably more prevalent in VS than in DS. Taken

together, these data point to DAT nanoclustering as a way to shape tonic levels of DA in the

striatum – particularly in the VS.

Our incorporation of receptor binding was inspired by previous data[22,23,71]. Dreyer &

colleagues assumed instantaneous equilibrium between extracellular DA and receptor

occupation while Hunger & colleagues based their kinetics on early pharmacological

studies[72–75]. However, we based our receptor kinetics on newer pharmacological experiments

in live cells[42] and properties of the recently developed DA receptor-based biosensors[41].

Indeed, these sensors are mutated receptors but only on the intracellular domains with no

changes of the binding site [41]. Overall, they exhibit kinetics that are well aligned with both

modelled and experimentally reported extracellular DA dynamics using non-biosensor-based

methods[3,37,76].

The presented simulations suggested that receptor binding was largely invariant to

single release events during pacemaker activity, while bursts of activity rapidly changed

occupancy. Both D1R and D2R immediately responded to burst onset; however, while D1R

occupancy rapidly declined to zero within approximately 50 ms, the slow D2R kinetics

resulted in an occupancy decline over ~5 seconds, returning to the baseline maintained by

tonic firing. This means that D1R is better suited to discriminate inputs in rapid succession

and allow for postsynaptic decoding of the fast-paced in vivo dynamics described in

particularly for DS[12,13,43]. By contrast, our analysis shows that D2Rs integrate DA signals

over several seconds. As D1R occupancy is negligible during pacemaker activity and D2R

kinetics are too slow to pick up rapid changes in DA concentration, our simulations moreover

suggest pauses in firing of less than 1 second is not an effective way of signalling for the

striatal dopaminergic system.

In conclusion, we have developed a three-dimensional model for DA release

dynamics and receptor binding that integrates a wealth of experimentally determined

parameters and generates responses to electrical and pharmacological input that fits robustly

with literature observations. The model offers an important theoretical framework and a

predictive tool that can serve as the basis for future experimental endeavours and help guide

the interpretation of new as well as older empirical findings on DA signalling dynamics under

both physiological conditions and in disease.

Methods

Three-dimensional finite difference model

DA release sites in the striatum were simulated at a density of one site per 25 μm^3 in DS and

one site per 27.8 μm^3 in VS (except for Fig. 4b and c). As a single medium spiny neuron in

the striatum is estimated to be influenced by axons from 95 to 194 dopaminergic neurons[77],

we simulated DA release sites as originating from 150 separate cell bodies. These were set to

fire independently from each other at 4 Hz generated by a Poisson distribution. We set the

release probability in response to an action potential at the individual terminal levels to 6%[23],

and if a release occurred the DA concentration of the voxel was elevated by the equivalent of

3000 DA molecules as per Equation 1. The volume of each voxel was corrected for an

extracellular volume fraction of 0.21[78].

We did not incorporate paired-pulse depression in our model as most in vivo FSCV

studies have not reported this phenomenon[45,79–81]. In addition, we could explain the gradual

blunt of the response during prolonged stimulation (Fig. 2e) as the result of an equilibrium

between release and uptake rather than a depression in release. Newer in vivo biosensor-

based studies observe the same "blunting" phenomenon during stimulation trains[82]. A low

initial release probability that scales well with frequency has also been reported for VMAT2-

positive release sites in dopaminergic cultures where the axons are not severed from their cell

bodies as in acute striatal slices[83]. Note that we did not explicitly model autoreceptor

inhibition at pacemaker activity. However, this value should be implicitly reflected in the in

vivo estimates we base our model on.

When simulating bursts or trains of stimulations, action potentials were added as

described for each simulation, but release probability was kept the same unless otherwise

noted. Uptake was evenly distributed throughout the simulation space with a V_{max} at 6 $\mu m\ s^{-1}$

for DS and 2 $\mu m\ s^{-1}$ for VS (see Extended Data Table S2) and a K_m at 210 nM[84] and

calculated for each voxel as per Equation 2. For each time step, *dt*, diffusion calculated with a

finite difference implementation of the Laplacian operator described in Equation 3 using a

diffusion coefficient corrected for tortuosity at 321.7 μm^2 s^{-1}. Periodic boundaries were used

to avoid edge issues with a simulation size of 50x50x50 μm unless otherwise noted. The

Python-based implementation of the model can be found as described in the Code and Data

Availability section.

Immunohistochemistry analysis

Analysed widefield images were obtained from (Sørensen et al., 2021[44]). The mouse striatum

slices were stained for either DAT or VMAT2 (see Sørensen et al., 2021 for slicing, fixation,

staining and imaging protocol[44]). Fluorescence intensities were extracted in a single

dorsoventral line parallel to the midline through the horizontally centre of the anterior

commissure using ImageJ v1.53q. for each slice. Data was vertically centred around the

anterior commissure. No background subtraction was performed.

FSCV modelling

FSCV recordings were modelled by convoluting a simulated DA response to a pseudo FSCV

trace. The impulse function used was as described in Venton et al. 2002[76]. In brief, the

impulse function was defined as $IF_{FSCV} = e^{-(t+1)(k_{-1}*t_s + k_{-2}*t_o)}$, where t is time, t_s is scan

time, t_o is oxidization time, and k_{-1} and k_{-2} desorption constants. t_s was set to 100 ms as

reported in May & Wightman, 1989[45] and t_o to 4 ms, which is half of the typically reported

symmetrical waveform length. The desorption constants (k_{-1} = 1.2 s^{-1} and k_{-2} = 12 s^{-1}) were

taken directly from Venton et al., 2002[76].

Nanocluster uptake modelling

Nanoclustering of DAT was simulated in Python. An ellipsoid-shaped varicosity of 1.5 μm in

length and 800 nm in width was modelled as a flat, square surface with a width of 1.8 μm to

match the total surface area[57,58]. As 9-16.4% of terminals in the striatum are estimated to be

dopaminergic[59,60], we set the height of the simulation space to 3.52 μm to generate an

extracellular volume seven times the size of the varicosity. Axons and cell bodies are likely a

non-negligible part of the striatal volume but omitted in this analysis due to a lack of

information in the literature (however, decreasing the share of striatal space taken up by DA

varicosities works in the direction of our hypothesis). Spatial granularity was set to 20 nm,

and we randomly placed eight non-overlapping locations for clusters on the simulated

varicosity. Uptake capacity per voxel for each clustering scenario was set so the mean uptake

of the entire space was 4 μm s^{-1}, which resulted in clusters with increased DAT density with

decreasing cluster size. We set the initial DA concentration of the entire space to 100 nM and

ran the simulation with a time step of 1.875e^{-7} s to avoid numerical instability. Diffusion

corrected for tortuosity was set to 321.7 μm^2 s^{-1} and calculated with a Laplacian operation as

described in Equation 3. Periodic boundaries were used to account for the varicosity being

circular and avoid issues at the edge of the simulation space. The full implementation in

Python can be found as described in the Code and Data Availability section.

Super-resolution clustering analysis

Super-resolved fluorescence microscopy dual color images from Sørensen et al., 2021[44] of

striatal brain slices from mice were analyzed based on their position in the dorsal or ventral

regions (Fig. 4j). Localizations were fitted using DAOSTORM[85] on the originally acquired

direct stochastic optical reconstruction microscopy (dSTORM) videos with the following

settings: background sigma of 8, maximum likelihood estimation as the fitting error model,

20 peak identification iterations, an initial sigma estimate of 1.5 and a threshold adjusted to

each imaging session. Each image was analysed for percentage of localizations within a

cluster using density-based spatial clustering of applications with noise (DBSCAN)[86] using

the Python package Scikit-learn 0.22[87]. To uncover broader trends, we scanned across a range

of parameters: radius was run from 10 nm to 200 nm and number of localizations from 10 to

200. For Fig. 4m a diameter of 80 nm was chosen to match the size reported in Rahbek-

Clemmensen et al. 2017[57].

Statistical Analysis

The statistical analyses performed can be found in the legends of each figure. Statistical

analyses were performed using open-source python packages SciPy v1.5.2 and NumPy

v1.18.1. Boxplots show the 25th and 75th percentile range and whiskers indicate up to 1.5

times the interquartile range. The remaining data points are plotted as individual outliers.

Code and data availability

Numerical implementation and code to run the simulations are available at:

https://github.com/GetherLab/striatal-dopamine-modelling. All experimental data are from

Ejdrup & Wellbourne-Wood et al., 2023[13] or Sørensen et al., 2021[44]. Corresponding authors

on these papers should be consulted for data availability.

Acknowledgements

We thank Dr. Kenneth L. Madsen, University of Copenhagen for general input on our modelling and Dr. Nicolas Tritsch, New York University Grossman School of Medicine, for a fruitful discussion on the postsynaptic receptor properties and occupation. The work was supported by the Lundbeck Foundation grants R266-2017-4331 (UG), R276-2018-792 (UG), R359-2020-2301 (UG), R181-2014-3090 (FH) and R303-2018-3540 (FH).

Author contributions

ALE, FH and UG conceptualized the study. ALE built the model with help from JKD. ALE ran the simulations. MDL and SHJ provided technical input on experimental data. ALE drafted the manuscript. ALE, FH and UG wrote the final version of the manuscript with comments and suggestions from JKD, TWR and JWD. All authors critically evaluated the manuscript. FH and UG supervised the project.

Competing Interests

JKD is employed at H. Lundbeck A/S

References

1. Berke, J. D. What does dopamine mean? *Nature Neuroscience 2018 21:6* **21**, 787–793 (2018).

2. Klaus, A., Alves Da Silva, J. & Costa, R. M. What, If, and When to Move: Basal Ganglia Circuits and Self-Paced Action Initiation. *Annu Rev Neurosci* **42**, 459–483 (2019).

3. Gonon, F. *et al.* Geometry and kinetics of dopaminergic transmission in the rat striatum and in mice lacking the dopamine transporter. *Prog Brain Res* **125**, 291–302 (2000).

4. Cragg, S. J. & Rice, M. E. DAncing past the DAT at a DA synapse. *Trends Neurosci* **27**, 270–277 (2004).

5. Sulzer, D., Cragg, S. J. & Rice, M. E. Striatal dopamine neurotransmission: regulation of release and uptake. *Basal Ganglia* **6**, (2016).

6. Borroto-Escuela, D. O. *et al.* Brain Dopamine Transmission in Health and Parkinson's Disease: Modulation of Synaptic Transmission and Plasticity Through Volume Transmission and Dopamine Heteroreceptors. *Front Synaptic Neurosci* **10**, (2018).

7. Agnati, L. F., Bjelke, B. & Fuxe, K. Volume versus wiring transmission in the brain: a new theoretical frame for neuropsychopharmacology. *Med Res Rev* **15**, (1995).

8. Schultz, W. Multiple dopamine functions at different time courses. *Annu Rev Neurosci* **30**, (2007).

9. Niv, Y., Daw, N. D., Joel, D. & Dayan, P. Tonic dopamine: opportunity costs and the control of response vigor. *Psychopharmacology (Berl)* **191**, 507–520 (2007).

10. Liu, C., Goel, P. & Kaeser, P. S. Spatial and temporal scales of dopamine transmission. *Nat Rev Neurosci* (2021) doi:10.1038/s41583-021-00455-7.

11. Sippy, T. & Tritsch, N. X. Unraveling the dynamics of dopamine release and its actions on target cells. *Trends Neurosci* (2023) doi:10.1016/J.TINS.2022.12.005.

12. Jørgensen, S. H. *et al.* Behavioral encoding across timescales by region-specific dopamine dynamics. *Proceedings of the National Academy of Sciences* **120**, e2215230120 (2023).

13. Ejdrup, A. L. *et al.* Within-Mice Comparison of Microdialysis and Fiber Photometry-Recorded Dopamine Biosensor during Amphetamine Response. *ACS Chem Neurosci* **14**, (2023).

14. Cardozo, D. L. & Bean, B. P. Voltage-dependent calcium channels in rat midbrain dopamine neurons: modulation by dopamine and GABAB receptors. *J Neurophysiol* **74**, 1137–1148 (1995).

15. Threlfell, S. *et al.* Striatal muscarinic receptors promote activity dependence of dopamine transmission via distinct receptor subtypes on cholinergic interneurons in ventral versus dorsal striatum. *J Neurosci* **30**, 3398–3408 (2010).

16. Richards, T. L. & Zahniser, N. R. Rapid substrate-induced down-regulation in function and surface localization of dopamine transporters: rat dorsal striatum versus nucleus accumbens. *J Neurochem* **108**, 1575–1584 (2009).

17. Brown, H. D., McCutcheon, J. E., Cone, J. J., Ragozzino, M. E. & Roitman, M. F. Primary food reward and reward-predictive stimuli evoke different patterns of phasic dopamine signaling throughout the striatum. *Eur J Neurosci* **34**, (2011).

18. Kearney, P. J. *et al.* Presynaptic Gq-coupled receptors drive biphasic dopamine transporter trafficking that modulates dopamine clearance and motor function. *J Biol Chem* **299**, (2023).

19. Salinas, A. G. *et al.* Distinct sub-second dopamine signaling in dorsolateral striatum
measured by a genetically-encoded fluorescent sensor. *Nature Communications 2023*
14:1 **14**, 1–16 (2023).

20. Mohebi, A., Wei, W., Pelattini, L., Kim, K. & Berke, J. D. Dopamine transients follow
a striatal gradient of reward time horizons. *Nature Neuroscience 2024* 1–10 (2024)
doi:10.1038/s41593-023-01566-3.

21. Dodson, P. D. *et al.* Representation of spontaneous movement by dopaminergic
neurons is cell-type selective and disrupted in parkinsonism. *Proc Natl Acad Sci U S A*
113, E2180-8 (2016).

22. Dreyer, J. K. & Hounsgaard, J. Mathematical model of dopamine autoreceptors and
uptake inhibitors and their influence on tonic and phasic dopamine signaling.
https://doi.org/10.1152/jn.00502.2012 (2013) doi:10.1152/jn.00502.2012.

23. Dreyer, J. K., Herrik, K. F., Berg, R. W. & Hounsgaard, J. D. Influence of phasic and
tonic dopamine release on receptor activation. *Journal of Neuroscience* **30**, 14273–
14283 (2010).

24. Beyene, A. G., McFarlane, I. R., Pinals, R. L. & Landry, M. P. Stochastic Simulation
of Dopamine Neuromodulation for Implementation of Fluorescent Neurochemical
Probes in the Striatal Extracellular Space. *ACS Chem Neurosci* **8**, 2275–2289 (2017).

25. Wiencke, K., Horstmann, A., Mathar, D., Villringer, A. & Neumann, J. Dopamine
release, diffusion and uptake: A computational model for synaptic and volume
transmission. *PLoS Comput Biol* **16**, e1008410 (2020).

26. Venton, B. J. *et al.* Real-time decoding of dopamine concentration changes in the
caudate-putamen during tonic and phasic firing. *J Neurochem* **87**, 1284–1295 (2003).

27. Dreyer, J. K., M, V. W. C., Lovic, V. & Aragona, B. J. Functionally Distinct
Dopamine Signals in Nucleus Accumbens Core and Shell in the Freely Moving Rat. *J*
Neurosci **36**, (2016).

28. Doucet, G., Descarries, L. & Garcia, S. Quantification of the dopamine innervation in
adult rat neostriatum. *Neuroscience* **19**, (1986).

29. Olson, L., Seiger, Å. & Fuxe, K. Heterogeneity of striatal and limbic dopamine
innervation: Highly fluorescent islands in developing and adult rats. *Brain Res* **44**,
283–288 (1972).

30. Jones, S. R. *et al.* Profound neuronal plasticity in response to inactivation of the
dopamine transporter. *Proc Natl Acad Sci U S A* **95**, 4029–4034 (1998).

31. Eriksen, J., Jørgensen, T. N. & Gether, U. Regulation of dopamine transporter function
by protein-protein interactions: new discoveries and methodological challenges. *J*
Neurochem **113**, 27–41 (2010).

32. Eriksen, J. *et al.* Visualization of dopamine transporter trafficking in live neurons by
use of fluorescent cocaine analogs. *J Neurosci* **29**, 6794–6808 (2009).

33. Block, E. R. *et al.* Brain Region-Specific Trafficking of the Dopamine Transporter. *J*
Neurosci **35**, 12845–12858 (2015).

34. Nicholson, C. Interaction between diffusion and Michaelis-Menten uptake of
dopamine after iontophoresis in striatum. *Biophys J* **68**, 1699–1715 (1995).

35. Nicholson, C. Diffusion from an injected volume of a substance in brain tissue with
arbitrary volume fraction and tortuosity. *Brain Res* **333**, 325–329 (1985).

36. Xie, K. *et al.* Organic electrochemical transistor arrays for real-time mapping of
evoked neurotransmitter release in vivo. *Elife* **9**, (2020).

37. Atcherley, C. W. *et al.* Improved Calibration of Voltammetric Sensors for Studying
Pharmacological Effects on Dopamine Transporter Kinetics in Vivo. *ACS Chem*
Neurosci **6**, 1509–1516 (2014).

38. Brimblecombe, K. R. *et al.* Calbindin-D28K Limits Dopamine Release in Ventral but
Not Dorsal Striatum by Regulating Ca2+ Availability and Dopamine Transporter
Function. *ACS Chem Neurosci* **10**, 3419–3426 (2019).
39. Stuber, G. D., Hnasko, T. S., Britt, J. P., Edwards, R. H. & Bonci, A. Dopaminergic
Terminals in the Nucleus Accumbens But Not the Dorsal Striatum Corelease
Glutamate. *Journal of Neuroscience* **30**, 8229–8233 (2010).
40. Ejdrup, A. L. *et al.* Within-animal comparison of microdialysis and fiber photometry
in amphetamine-exposed mice. *bioRxiv* 2022.10.27.513984 (2022)
doi:10.1101/2022.10.27.513984.
41. Labouesse, M. A. & Patriarchi, T. A versatile GPCR toolkit to track in vivo
neuromodulation: not a one-size-fits-all sensor. *Neuropsychopharmacology 2021*
46:12 **46**, 2043–2047 (2021).
42. Ågren, R., Stepniewski, T. M., Zeberg, H., Selent, J. & Sahlholm, K. Dopamine d2
receptor agonist binding kinetics—role of a conserved serine residue. *Int J Mol Sci* **22**,
(2021).
43. Wei, W., Mohebi, A. & Berke, J. D. A Spectrum of Time Horizons for Dopamine
Signals. *bioRxiv* 2021.10.31.466705 (2022) doi:10.1101/2021.10.31.466705.
44. Sørensen, G. *et al.* Disruption of the PDZ domain-binding motif of the dopamine
transporter uniquely alters nanoscale distribution, dopamine homeostasis, and reward
motivation. *J Biol Chem* **297**, (2021).
45. May, L. J. & Wightman, R. M. Effects of D-2 antagonists on frequency-dependent
stimulated dopamine overflow in nucleus accumbens and caudate-putamen. *J*
Neurochem **53**, 898–906 (1989).
46. Pereira, D. B. *et al.* Fluorescent false neurotransmitter reveals functionally silent
dopamine vesicle clusters in the striatum. *Nature Neuroscience 2016 19:4* **19**, 578–586
(2016).
47. Liu, C., Kershberg, L., Wang, J., Schneeberger, S. & Kaeser, P. S. Dopamine
Secretion Is Mediated by Sparse Active Zone-like Release Sites. *Cell* **172**, 706-
718.e15 (2018).
48. Ducrot, C. *et al.* Dopaminergic neurons establish a distinctive axonal arbor with a
majority of non-synaptic terminals. *The FASEB Journal* **35**, e21791 (2021).
49. Garris, P. A., Ciolkowski, E. L., Pastore, P. & Wightman, F. M. Efflux of Dopamine
from the Synaptic Cleft in the Nucleus Accumbens of the Rat Brain. (1994).
50. Staal, R. G. W., Mosharov, E. V. & Sulzer, D. Dopamine neurons release transmitter
via a flickering fusion pore. *Nat Neurosci* **7**, 341–346 (2004).
51. Sulzer, D. & Pothos, E. N. Regulation of quantal size by presynaptic mechanisms. *Rev*
Neurosci **11**, 159–212 (2000).
52. Pothos, E. N., Davila, V. & Sulzer, D. Presynaptic recording of quanta from midbrain
dopamine neurons and modulation of the quantal size. *J Neurosci* **18**, 4106–4118
(1998).
53. Kristensen, A. S. *et al.* SLC6 neurotransmitter transporters: structure, function, and
regulation. *Pharmacol Rev* **63**, 585–640 (2011).
54. Carboni, E. *et al.* Cocaine and Amphetamine Increase Extracellular Dopamine in the
Nucleus Accumbens of Mice Lacking the Dopamine Transporter Gene. *Journal of*
Neuroscience **21**, RC141–RC141 (2001).
55. Maisonneuve, I. M. & Glick, S. D. Interactions between ibogaine and cocaine in rats:
in vivo microdialysis and motor behavior. *Eur J Pharmacol* **212**, 263–266 (1992).
56. Wu, Q., Reith, M. E. A., Kuhar, M. J., Carroll, F. I. & Garris, P. A. Preferential
increases in nucleus accumbens dopamine after systemic cocaine administration are

704		caused by unique characteristics of dopamine neurotransmission. *J Neurosci* **21**, 6338–
705		6347 (2001).
706	57.	Rahbek-Clemmensen, T. *et al.* Super-resolution microscopy reveals functional
707		organization of dopamine transporters into cholesterol and neuronal activity-dependent
708		nanodomains. *Nat Commun* **8**, 1–14 (2017).
709	58.	Lycas, M. D. *et al.* Nanoscopic dopamine transporter distribution and conformation are
710		inversely regulated by excitatory drive and D2 autoreceptor activity. *Cell Rep* **40**,
711		111431 (2022).
712	59.	Tennyson, V. M., Heikkila, R., Mytilineou, C., Côté, L. & Cohen, G. 5-
713		Hydroxydopamine 'tagged' neuronal boutons in rabbit neostriatum: interrelationship
714		between vesicles and axonal membrane. *Brain Res* **82**, 341–348 (1974).
715	60.	Hökfelt, T. In vitro studies on central and peripheral monoamine neurons at the
716		ultrastructural level. *Z Zellforsch Mikrosk Anat* **91**, 1–74 (1968).
717	61.	Markowitz, J. E. *et al.* Spontaneous behaviour is structured by reinforcement without
718		explicit reward. *Nature 2023 614:7946* **614**, 108–117 (2023).
719	62.	Howe, M. W., Tierney, P. L., Sandberg, S. G., Phillips, P. E. M. & Graybiel, A. M.
720		Prolonged dopamine signalling in striatum signals proximity and value of distant
721		rewards. *Nature* **500**, 575–579 (2013).
722	63.	Grace, A. A., Floresco, S. B., Goto, Y. & Lodge, D. J. Regulation of firing of
723		dopaminergic neurons and control of goal-directed behaviors. *Trends Neurosci* **30**,
724		220–227 (2007).
725	64.	Kuczenski, R. & Segal, D. S. Differential effects of amphetamine and dopamine
726		uptake blockers (cocaine, nomifensine) on caudate and accumbens dialysate dopamine
727		and 3-methoxytyramine. *Journal of Pharmacology and Experimental Therapeutics*
728		**262**, (1992).
729	65.	Shen, H. W. *et al.* Regional Differences in Extracellular Dopamine and Serotonin
730		Assessed by In Vivo Microdialysis in Mice Lacking Dopamine and/or Serotonin
731		Transporters. *Neuropsychopharmacology 2004 29:10* **29**, 1790–1799 (2004).
732	66.	Chefer, V. I., Thompson, A. C., Zapata, A. & Shippenberg, T. S. Overview of Brain
733		Microdialysis. *Curr Protoc Neurosci* **47**, 7.1.1-7.1.28 (2009).
734	67.	Fagan, R. R. *et al.* Dopamine transporter trafficking and Rit2 GTPase: Mechanism of
735		action and in vivo impact. *J Biol Chem* **295**, 5229–5244 (2020).
736	68.	Zhu, S. *et al.* Identification of a Vav2-dependent mechanism for GDNF/Ret control of
737		mesolimbic DAT trafficking. *Nat Neurosci* **18**, 1084–1093 (2015).
738	69.	Kearney, P. J. *et al.* Presynaptic Gq-coupled receptors drive biphasic dopamine
739		transporter trafficking that modulates dopamine clearance and motor function. *J Biol*
740		*Chem* **299**, 102900 (2023).
741	70.	Lee, S. J. *et al.* Cell-type-specific asynchronous modulation of PKA by dopamine in
742		learning. *Nature 2020 590:7846* **590**, 451–456 (2020).
743	71.	Hunger, L., Kumar, A. & Schmidt, R. Abundance Compensates Kinetics: Similar
744		Effect of Dopamine Signals on D1 and D2 Receptor Populations. *J Neurosci* **40**, 2868–
745		2881 (2020).
746	72.	Nishikori, K., Noshiro, O., Sano, K. & Maeno, H. Characterization, Solubilization, and
747		Separation of T w o Distinct Dopamine Receptors in Canine Caudate Nucleus*.
748		*Journal of Biological Chemistry* **255**, 10909–10915 (1980).
749	73.	Maeno, H. Dopamine receptors in canine caudate nucleus. *Mol Cell Biochem* **43**, 65–
750		80 (1982).
751	74.	Sano, K., Noshiro, O., Katsuda, K., Nishikori, K. & Maeno, H. Dopamine receptors
752		and dopamine-sensitive adenylate cyclase in canine caudate nucleus. Characterization
753		and solubilization. *Biochem Pharmacol* **28**, 3617–3627 (1979).

75. Burt, D. R., Creese, I. & Snyder, S. H. Properties of [3H]Haloperidol and
[3H]Dopamine Binding Associated with Dopamine Receptors in Calf Brain
Membranes. *Mol Pharmacol* **12**, (1976).
76. Venton, B. J., Troyer, K. P. & Wightman, R. M. Response times of carbon fiber
microelectrodes to dynamic changes in catecholamine concentration. *Anal Chem* **74**,
539–546 (2002).
77. Matsuda, W. *et al.* Single Nigrostriatal Dopaminergic Neurons Form Widely Spread
and Highly Dense Axonal Arborizations in the Neostriatum. *Journal of Neuroscience*
(2009) doi:10.1523/JNEUROSCI.4029-08.2009.
78. Rice, M. E. & Nicholson, C. Diffusion characteristics and extracellular volume
fraction during normoxia and hypoxia in slices of rat neostriatum.
https://doi.org/10.1152/jn.1991.65.2.264 **65**, 264–272 (1991).
79. Garris, P. A. & Wightman, R. M. Different Kinetics Govern Dopaminergic
Transmission in the Amygdala, Prefrontal Cortex, and Striatum: An in vivo
Voltammetric Study. *The Journal of Neuroscience* **14**, 442–450 (1994).
80. Hamid, A. A. *et al.* Mesolimbic dopamine signals the value of work. *Nat Neurosci* **19**,
117–126 (2015).
81. Chergui, K., Suaud-Chagny, M. F. & Gonon, F. Nonlinear relationship between
impulse flow, dopamine release and dopamine elimination in the rat brainin vivo.
Neuroscience **62**, 641–645 (1994).
82. Mohebi, A., Collins, V. L. & Berke, J. D. Cholinergic Interneurons Drive Motivation
by Promoting Dopamine Release in the Nucleus Accumbens. *bioRxiv*
2022.11.06.515335 (2022) doi:10.1101/2022.11.06.515335.
83. Silm, K. *et al.* Synaptic Vesicle Recycling Pathway Determines Neurotransmitter
Content and Release Properties. *Neuron* **102**, 786-800.e5 (2019).
84. Hovde, M. J., Larson, G. H., Vaughan, R. A. & Foster, J. D. Model Systems for
Analysis of Dopamine Transporter Function and Regulation. *Neurochem Int* **123**, 13
(2019).
85. Babcock, H., Sigal, Y. M. & Zhuang, X. A high-density 3D localization algorithm for
stochastic optical reconstruction microscopy. *Opt Nanoscopy* **1**, 1–10 (2012).
86. Ester, M., Kriegel, H.-P., Sander, J. & Xu, X. A Density-Based Algorithm for
Discovering Clusters in Large Spatial Databases with Noise. in *Proceedings of the 2nd
International Conference on Knowledge Discovery and Data Mining* (1996).
87. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine
Learning Research* **12**, 2825–2830 (2021).
88. Paladini, C. A., Robinson, S., Morikawa, H., Williams, J. T. & Palmiter, R. D.
Dopamine controls the firing pattern of dopamine neurons via a network feedback
mechanism. *Proc Natl Acad Sci U S A* **100**, 2866–2871 (2003).
89. Zhou, F. M., Liang, Y. & Dani, J. A. Endogenous nicotinic cholinergic activity
regulates dopamine release in the striatum. *Nat Neurosci* **4**, 1224–1229 (2001).
794
795

Table 1 – List of variables used in simulation of the dorsal striatum.

Variable	Abbreviation	Value	Reference
Firing rate		4 Hz	[88]
Release probability		6%	[23]
DA molecules per vesicle		3000	[4]
Diffusion coefficient		763 μm^2 s^{-1}	[34]
Tortuosity		1.54	[78]
Vmax		6.0 μm s^{-1}	See Table S2
Km		210 nM	[84]
Active terminal density	-	0.04 μm^{-3}	[10]
Extracellular volume fraction		0.21	[78]
Number of neurons in simulation space	-	150	[77]

Fig. 1 – Large-scale 3D model of the dorsal striatum

a, Self-enveloped simulation space of 100^3 μm^3 with approximately 40.000 release sites from 150 neurons. Colours of individual release sites are not matched to neurons. **b,** Simulation of a single release event after 5 ms and 10 ms. Colour-coded by DA concentration. **c,** Comparison of analytical solution and simulation of diffusion after a single release event at three different time points. **d,** Representative snapshot of steady state DA dynamics at 4 Hz tonic firing with parameters mirroring the dorsal striatum. **e,** Cross-section of temporal dynamics for data shown in (d). **f,** Histogram of DA concentrations ([DA]) across the entire space in (d). **g,** DA release during three burst activity scenarios for all release sites in a 10x10x10 μm cube (black boxes) and spill-over into the surrounding space. Burst simulated as increase in firing rate on top of continued tonic firing of the surrounding space. Traces on top are average DA concentrations for the marked cubes, with bursts schematized by coloured lines below. First image row is at the end of the burst, second row another 100 ms after. Scale bars for traces are 200 ms and 500 nM. Scale bar for the images is 20 μm. **h,** Peak DA concentration reached at different distances from area with phasic activity for the three firing scenarios. **i,** Top: representative [DA] trace 1 μm away from a release site during pacemaker and burst activity. Bottom: Occupancy of D1Rs and D2Rs for same site. **j,** Effect of complete pause in firing for 1 second on both average [DA] and D1R and D2R occupation.

Fig. 2 – Regional difference in uptake greatly impact DA dynamics

a. Representative snapshots of steady state dynamics at 4 Hz tonic firing with parameters mirroring the dorsal (left) and ventral striatum (right). **b,** Cross-section of temporal dynamics for data shown in a. Bottom row show concentrations of the dashed lines in the top panels **c,** Normalized density of DA concentration of simulations in (a). Thick lines are for the entire space, thin lines are across time for five randomly sampled locations. Dashed red line is for simulation of the ventral striatum with lowest reported innervation density in the literature. **d,** Same data as in (c), but for concentration percentiles. Note that even the lowest percentiles of VS were above 10 nM in [DA]. **e,** Convoluted model response (extended data Fig. 1c) to mimic FSCV measurements mirroring the experimentally tested stimulation paradigm in [45] for the dorsal (left) and ventral striatum (right) **f,** DA release during three burst activity scenarios for all release sites in a 10x10x10 µm cube (black boxes) and spill-over into the surrounding space. Burst simulated as increase in firing rate on top of continued tonic firing of the surrounding space. Traces on top are average DA concentrations for the marked cubes, with bursts schematized by coloured lines below. First image row is at the end of the burst, second row another 100 ms after. Scale bars for traces are 200 ms and 500 nM. Scale bar for the images is 20 µm. **g,** Top: representative [DA] trace 1 µm away from a release site during pacemaker and burst activity. Bottom: Occupancy of D1Rs and D2Rs for same site. Occupancy data from corresponding DS simulation on Fig. 1k shown in as dotted line. **h,** Peak occupancy at different distances from the area bursting normalized to maximal and minimum occupancy.

Fig. 3 – Sensitivity of the model to parameter changes

a, Schematic of the fraction of active release sites. Black dots are inactive sites and green dots
indicate actively releasing sites. **b,** Effect of changing fraction of active release sites on DA
concentrations. Blue line, DS peak DA concentration (99.5th percentile); Red line, VS peak DA
concentration (99.5th percentile); Dotted blue line, DS tonic DA concentration (50th percentile);
Dotted red line, VS tonic DA concentration (50th percentile).**c,** Ratio between peak (99.5th
percentile) and tonic (50th percentile) concentrations across fractions of active release sites in
the DS (blue line) and VS (red line) as a measure of DA signal focality. **d,** Schematic of
changing quantal size (Q). **e,** Effect of changing quantal size on tonic and peak DA
concentrations in DS (blue lines) and VS (red lines). **f,** Ratio between peak and tonic
concentrations across various quantal sizes in in DS (blue line) and VS (red line). **g,** Relative
difference between the DS and VS for peak (black line) and tonic DA (dotted line) at different
quantal sizes. **h,** Schematic of changing DAT K_m. **i,** Effect of changing DAT K_m on DA
concentrations in DS (blue lines) and VS (red lines). **j,** Schematic of changing DAT V_{max}. **k,**
Effect of changing DAT V_{max} on DA concentrations. Shaded areas are median V_{max} of the two
regions (DS and VS) as found in the literature shown in Table S2 $\pm$ 50%. **l,** Effect of changing
DAT V_{max}, with tonic (50th percentile) and peak (99.5th percentile) DA concentrations
normalized to their value at 2 µm s^{-1} (median value for VS). Shaded area indicates median V_{max}
for VS found in the literature shown in Table S2 $\pm$ 50%.

Fig. 4 – DAT nanoclustering reduces uptake and shows regional variation
a. Schematic of dense DA cluster. White dots represent individual DAT molecules, and colour gradient the surrounding DA concentration. **b,** Effective transport rate dependent on local concentration. **c,** Top view of unfolded DA varicosity. Black shapes denote clusters of DAT. Dashed white line indicates placement of cross-section shown in (d). **d,** Cross section showing DA concentration in space surrounding varicosity unfolded in c. Grey line at the bottom is surface of the varicosity. Colour-coded for DA concentration. **e,** Top view from (c), but colour coded for DA concentration immediately above membrane surface at different DAT cluster sizes. **f,** Mean [DA] over time for different DAT cluster sizes and unclustered (Un.). **g,** Time to reach a mean DA concentration of 10 nM for different DAT cluster sizes and unclustered (Un.). **h,** Difference between DA concentration at the centre of clusters and mean concentration throughout the simulation for different cluster sizes and unclustered (Un.). **i,** Concentrations across a cross section of a surface with 80 nm diameter clusters. Shaded areas highlight cluster locations. **j,** Location of images of the dorsal (DS) and ventral striatum (VS) in striatal slices from mice as imaged in (Sørensen et al., 2021[44]) with direct stochastic optical reconstruction microscopy (dSTORM). **k,** Two representative DA varicosities from DS and VS with VMAT2 in white and DAT in magenta. Images are 1.5x2 µm. **l,** Individual DAT localizations from images in (k) coloured by clustering. Black indicates localization identified as clustered based on DBSCAN with parameters 80 nm diameter and 40 localizations. Grey indicates unclustered localizations. **m,** Quantification of clustering across all images in (j) with parameters in (l). Welch's two-sample t-test, P = 0.012(*), n = 12 (DS) and 13 (VS). **n,** Absolute difference in percentage of clustering as assessed with DBSCAN across a range of parameters. VS has a higher propensity to cluster across cluster sizes typically reported for DAT clusters.

Extended Data Fig. 1 – Average concentration and receptor kinetics
a. Average DA concentration across a 100x100x100 μm volume of simulated DS at
pacemaker activity. **b,** Mean concentration change in response to single event with 60%
release probability (see supplementary notes on electrical stimulation) in all neurons at time
zero. Blue trace is output directly from simulation, grey trace is the predicted FSCV
measurement. **c,** Modelling of predicted FSCV measurement and representative snapshot of
simulation space just after a release event. **d,** Volume of space exposed to greater than 100
nM DA after firing relative to volume of space where terminals actively burst. **e,** Least-square
fit linear regression of the dLight and GRAB$_{DA}$ sensors based on reported kinetics[41] and
newest experimental characterization of D2R[42]. Shaded are indicates 95% C.I. **f,** Zoom on a
DA burst as on Fig. 1i, with [DA] in blue and D1R occupancy in teal. Shaded area indicates
period of bursting with 6 APs at 20 Hz.

Extended Data Fig. 2 – Simulation size and granularity
a. Schematic of different simulation sizes. **b,** Concentration percentiles at different simulation
diameters. Results are not robust until a diameter close to 50 µm is reached (line runs behind
100 µm line). **c,** Schematic of simulation granularity. **d,** Effect of simulation voxel diameter
on concentration percentiles. Virtually no difference in concentration profiles below the
99.9[th] percentile of [DA], with the 1.0 µm voxel size still following 0.1 and 0.5 µm well
above that level. Inset shows 99.75[th] to 100[th] percentile with y-axis matching main y-axis. **e,**
Absolute difference in [DA] between simulations at 0.1 µm voxel diameter and 0.5, 1.0 and
2.0 µm. Inset shows 99[th] to 100[th] percentile with y-axis matching main y-axis. **f,** Percentage
difference in [DA] between simulations at 0.1 µm voxel diameter and 0.5, 1.0 and 2.0 µm.

Extended Data Fig. 3 – Histochemical gradient of DAT and VMAT2 fluorescence

a. Representative image of the mouse striatal slices analysed in b. Dashed white line indicates the quantified dorsoventral gradient. **b,** Relative intensity of the DAT and VMAT2 immunosignal in the dorsoventral axis of striatal mouse brain slices from [44]. All slices show a drop at the anterior commissure (AC). Shaded areas around lines denote S.E.M.. **c,** Mean relative intensity of the DAT and VMAT2 signal before and after AC. Two-sided t-test, VS-DAT:VMAT2, P = 0.012(*), n = 4 mice; one-sided t-tests, DAT-DS:VS, P = 0.0021(**), VMAT2-DS:VS, P = 0.0086(**), n = 4 mice. **d,** Peak DA concentration reached at different distances from area with phasic activity for the three firing scenarios in VS. **e,** Volume of space in VS exposed to greater than 100 nM after firing relative to volume of space where terminals actively burst. **f,** Effect of complete pause in firing in VS for 1 second on both average [DA] and D1R and D2R occupation.

Extended Data Fig. 4 – Model parameter testing

a. Schematic of our definitions of tonic (50^{th} percentile/median, dashed lines) and peak (99.5^{th} percentile, solid line) DA for both the dorsal and ventral striatum. **b,** Effect of changing release probability ($R_\%$) on DA concentrations. **c,** Relative difference between the ventral and dorsal striatum at different percentiles for different release probabilities. **d,** Ratio between 99.5^{th} and 50^{th} percentile as a measure of focality for both regions. As $R_\%$ increases the concentrations becomes more homogeneous. **e,** Effect of changing firing rate on DA concentrations. **f,** Relative difference between the ventral and dorsal striatum at different percentiles for different firing rates. **g,** Ratio between 99.5^{th} and 50^{th} percentile for both regions. As firing rate increases the concentrations becomes more homogeneous.

Extended Data Fig. 5 – Fold change during inhibition, V$_{max}$-sensitivity at different release parameters and release-uptake balance.
a. Fold change over baseline (K$_m$ of 210 nM) for mean DA concentration in the dorsal (DS) and ventral striatum (VS) with changing DAT K$_m$. **b,** Relative difference between the dorsal and ventral striatum for both phasic and tonic DA at different K$_m$ values. **c,** Effect of changing DAT V$_{max}$ on DA concentrations for three different quantal sizes (Q). [DA] normalized to highest values within each Q. Shaded area indicates median V$_{max}$ for DS and VS as found in the literature shown in Extended Data Table S2 with ± 50%. **d,** Effect of changing DAT V$_{max}$ on DA concentrations for three different release probabilities (R$_\%$). [DA] normalized to highest values within each R$_\%$. Shaded area indicates median V$_{max}$ for DS and VS as found in Table S2 with ± 50%. **e,** Difference between VS and DS for both mean and median concentrations across different Qs for DS (VS held at 3,000). Shaded area indicates range of Q where median is higher in VS but mean is higher in DS. **f,** Least-square fit linear regression between release rate and autocorrelation decay rate (τ) from (Ejdrup & Wellbourne-Wood et al., 2023[13]). Shaded area highlights 95% C.I. **g,** Partial regression plot (error of the regression in (f) and error between DA response to amphetamine as measured by microdialysis and the release rate from (Ejdrup & Wellbourne-Wood et al., 2023[13]) to show that the less release and uptake correlate, the less release rate can explain the microdialysis response, suggesting release and uptake are partially independent of each other. Shaded area highlights 95% C.I.

Measure	Ratio	DS	VS	Units	Species	Source
TH expression density	100 %	~90	~90	A.U.	Mouse	Alberquilla et al., 2020
TH immunoreactivity	95 %	~68	~64	A.U.	Mouse	Kuroda et al., 2009
TH protein content	150 %	0.07	0.11	ng TH/µg prot.	Mouse	Salvatore et al., 2016
DA content	90 %	~155	~140	ng DA/mg prot.	Mouse	Salvatore et al., 2016
TH immunoreactivity	75 %	2.8	2.1	A.U.	Rat	Huang et al., 2019
TH protein content	66 %	0.36	0.24	ng TH/µg prot.	Mouse	Salvatore et al., 2005
FSCV - [DA]$_p$	91 %	57	52	nM	Rat	May & Wightman, 1989
FSCV - [DA]$_p$	76 %	89.3	67.5	nM	Rat	Garris & Wightman, 1994
Median	90 %	-	-	-	-	-

967 **Extended Data Table 1 – Overview of reports on dopaminergic density and release in
968 the striatum.** Only studies that assessed both regions in rodents are included. Studies that
969 stimulate directly in the striatum are omitted due to the large activation of nicotinic receptors
970 on DA terminals [47,89]. A.U. = arbitrary units, DS = dorsal striatum, VS = ventral striatum, TH
971 = tyrosine hydroxylase.

Method	VS/DS Ratio	DS (uM/s)	VS (uM/s)	Species	Source
FSCV	29 %	7.0	2.0	Mouse	Calipari et al, 2012
FSCV	28 %	6.0	1.7	Rat	Calipari et al, 2012
FSCV	47 %	3.0	1.4	Rat	May & Wightman, 1989
FSCV	31 %	6.5	2.0	Mouse	Siciliano et al., 2014
FSCV	44 %	6.0	2.2	Rat	Ferris et al., 2014
Median	31 %	6.0	2.0	-	-

975 **Extended Data Table 2 – Overview of reported V$_{max}$ values for DA uptake in the
976 striatum.** Only studies that assessed both regions in rodents are included. DS = dorsal
977 striatum, VS = ventral striatum, FSCV = fast-scan cyclic voltammetry.