Fat body-derived cytokine Upd2 controls disciplined migration of tracheal stem cells in *Drosophila*

Pengzhen Dong[1,2], Yue Li[1,2,3], Yuying Wang[1,2], Qiang Zhao[1,2], Tianfeng Lu[6], Jian Chen[1,2], Tianyu Guo[1,2], Jun Ma[7,8], Bing Yang[5,*], Honggang Wu[3,*], Hai Huang[1,2,4,*]

Second Affiliated Hospital, and Department of Cell Biology, Zhejiang University School of Medicine. Hangzhou, Zhejiang Province 310058.

State Key Laboratory of Transvascular Implantation Devices, Hangzhou 310009, China.

Zhejiang Key Laboratory of Precision Diagnosis and Therapy for Major Gynecological Diseases, Women's Hospital, Zhejiang University School of Medicine. Hangzhou, Zhejiang Province 310058.

Zhejiang Provincial Key Laboratory of Genetic & Developmental Disorders, Zhejiang University School of Medicine. Hangzhou 311121.

MOE Laboratory of Biosystem Homeostasis and Protection and Life Sciences Institute, Zhejiang University, Hangzhou, 310058, China.

Department of Developmental Biology & Neuroscience, Washington University in St. Louis, Missouri, United States, 63108.

Center for Genetic Medicine, the Fourth Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang 310058, China.

Institute of Genetics, Zhejiang University International School of Medicine, Hangzhou, Zhejiang 310058, China.

*Correspondence: bingyang@zju.edu.cn; honggangwu@zju.edu.cn; haihuang@zju.edu.cn

Abstract

Coordinated activation and directional migration of adult stem cells are essential for

maintaining tissue homeostasis. *Drosophila* tracheal progenitors are adult stem cells

that migrate posteriorly along the dorsal trunk to replenish degenerating branches that

disperse the FGF mitogen. However, it is currently unknown how the overall

anterior-to-posterior directionality of such migration is controlled. Here we show that

individual progenitor cells migrate together in a concerted, disciplined manner, a

behavior that is dependent on the neighboring fat body. We identify the fat body-

derived cytokine, Upd2, in targeting and inducing JAK/STAT signaling in tracheal

progenitors to maintain their directional migration. Perturbation of either Upd2

production in fat body or JAK/STAT signaling in trachea causes aberrant bidirectional

migration of tracheal progenitors. We show that JAK/STAT signaling promotes the

expression of genes involved in planar cell polarity leading to asymmetric localization

of Fat in progenitor cells. We provide evidence that Upd2 transport requires Rab5-

and Rab7-mediated endocytic sorting and Lbm-dependent vesicle trafficking. Our

study thus uncovers an inter-organ communication in the control of disciplined

migration of tracheal progenitor cells, a process that requires vesicular trafficking of

fat body-derived cytokine Upd2 and JAK/STAT signaling-mediated activation of PCP

genes.

Introduction

Adult stem cells are multipotent cell populations which inhabit their niche but mobilize to initiate tissue reconstruction during organismal growth and regeneration. An intriguing feature of stem cells is their capability of migrating in a disciplined directionality towards locations undergoing reconstruction (Li and Clevers, 2010). Such a highly disciplined movement is critical for maintaining tissue homeostasis and is influenced by various niche-intrinsic signals and external stimuli, and its aberrancy causes diseases such as hypertrophy (Zhou et al., 2024). The damaged tissue or distant organs that elicit systemic signals promote the migration of adult stem cells (Jones and Wagers, 2008). In addition, interactions with other cell types, soluble factors (e.g. cytokines, growth factors, and hormones) and tissue stiffness collectively bolster the mobilization of stem cells (Fuchs and Blau, 2020). Despite growing appreciation of adult stem cells as a primary source for tissue regeneration, the mechanism governing directional stem cell migration remains yet to be elucidated.

Drosophila tracheal progenitors are a population of adult stem cells that rebuild the degenerating trachea during metamorphosis. The progenitor cells reside in Tr4 and Tr5 metameres and start to move along the tracheal branch towards sites of regeneration (Chen and Krasnow, 2014; Pitsouli and Perrimon, 2010). Movement of these progenitor cells follows a stereotypical anterior-to-posterior axis (Figure 1A), thus representing a suitable system to investigate mechanisms controlling the directionality of stem cell migration. The activation of tracheal progenitors is stimulated by the morphogen Branchless (Bnl), fly homolog of fibroblast growth factor (FGF) (Chen and Krasnow, 2014), and the insulin hormone (Li et al., 2022). Intercellular communication and synergy between organs also contribute to the branching morphogenesis (Perochon et al., 2021; Schottenfeld et al., 2010; Tamamouna et al., 2021). The functional role of the interactions between trachea and other organs in modulating tracheal progenitor behavior has been largely unknown.

Drosophila fat body is the functional analog of mammalian adipose tissue and the major organ sensing various hormonal and nutritional signals to orchestrate systemic growth, metabolism and stem cell maintenance (Sriskanthadevan-Pirahas et al., 2022).

Fat body produces regulatory molecules known as fat body signals (FBSs), which

remotely affect the activity of other organs (Ingaramo et al., 2020; Zheng et al., 2016).

For instance, the fat body-to-brain signals modulate insulin-like peptides production

(Rajan and Perrimon, 2012), visual attention and sleep behavior (Ertekin et al., 2020).

The *Drosophila* family of interleukin-6 (IL-6)-like cytokines consist of Unpaired

(Upd, also called Outstretched), Upd2 and Upd3, and serve as mediators of systemic

signaling. Whereas Upd1 and Upd3 derive from fly brain and plasmatocytes (Beshel

et al., 2017; Woodcock et al., 2015), Upd2 is primarily produced by the fat body

(Rajan et al., 2017), although muscle-derived Upd2 is also reported (Zhao and

Karpac, 2017). The Upd proteins act as ligands which bind to a common GP130-like

receptor, Domeless (Dome) on target cells (Agaisse et al., 2003; Chen et al., 2002).

Upon association of ligands, the Dome receptors dimerize and recruit the non-receptor

tyrosine kinase JAKs leading to their subsequent transactivation via phosphorylation.

The transactivated JAKs then phosphorylate the tyrosine residues of their substrates,

including the bound receptors and cytosolic STATs. The phosphorylation of STATs

promotes their dimerization and nuclear translocation to activate transcriptional

program (Darnell, 1997). JAK/STAT signaling requires the IL-6 cytokines (Heinrich

et al., 2003), and is implicated in numerous cellular events including cell proliferation,

differentiation, migration and apoptosis (O'Shea et al., 2002).

Here, we investigate molecular basis underlying directional stem cell migration using

the *Drosophila* tracheal progenitors as a model. Our results identify a cytokine-

mediated inter-organ communication between fat body and the progenitor cells that is

necessary for their disciplined, directional migration. The directional migration of the

progenitors relies on JAK/STAT signaling and its downstream targets of planar cell

polarity (PCP) components. Importantly, the Upd2 cytokines derived from fat body

are transported through vesicular trafficking to induce JAK/STAT signaling in tracheal

progenitors. Our study reveals that tracheal progenitors establish migratory

directionality as they exit their niches and that the disciplined migration of the

progenitors depends on an inter-organ signaling originating from the fat body.

Results

Dependence of tracheal progenitors on the fat body

The fly tracheal progenitors are activated and move posteriorly along the dorsal trunk
(DT) at the onset of pupariation (Figure 1A). We set out to delve into the underlying
mechanisms of directional progenitor cell movement and tentatively surveyed organs
that may coordinate this process. In *Drosophila*, the fat body resides anatomically in
proximity with trachea (Figure 1B, C; Video 1) and is the principal reservoir for
energy consumption. To determine whether the integrity of fat body is required for
tracheal progenitors, we perturbed larval or pupal fat body by expressing pro-
apoptotic cell death genes, *hid* and *reaper* (*rpr*), under the control of a fat body-
specific driver, *lsp2*-Gal4 (Cherbas et al., 2003). Expression of *hid* and *rpr* in L3 stage
impaired fat body integrity and adipocyte abundance, and generated slender larvae
and pupae (Figure 1-figure supplement 1A-D). In these animals, the tracheal
progenitors exhibited a sign of undisciplined migration and tended to move
bidirectionally (Figure 1D-F), although their migration rate, cell number and
proliferation remained unchanged (Figure 1-figure supplement 1E-H and Figure 3-
figure supplement 1). The undisciplined bidirectional migration behavior of tracheal
progenitors in fat body-defective animals is in stark contrast to control animals where
the progenitors migrated unambiguously towards posterior (Figure 1D-D'' and Video
2). To gain a quantitative view of progenitor cell migration, we traced the movement
of individual cells by time-lapse confocal imaging. At 2hr APF, tracheal progenitors
from fat body deficit animals displayed a symmetrical distribution relative to the
junction between dorsal trunk (DT) and transverse connective (TC), compared with an
L-shape localization of niche-associated and migratory progenitors established by a
posterior movement in control (Figure 1G-H'). Gauging the vector denoting the
movement of each progenitor (Figure 1I-I') revealed that the traces of progenitor
groups in fat body-depleted animals exhibited a fan-shaped pattern (Figure 1J, J').
Owing to this undisciplined movement, entropy of the system was notably elevated
upon increased inconsistency of migration vectors (Figure 1K). The bidirectional
migratory progenitors displayed longer territory as assessed by binarized direction

(Figure 1L). Collectively, these results suggest that fat body has an integral role in

maintaining the discipline of tracheal progenitor movement.

<u>Upd2-JAK/STAT signaling between fat body and trachea</u>

Since fat body impacts the behavior of tracheal progenitors, we next attempted to

investigate the signal between these two interdependent organs. For this purpose, we

first performed RNA sequencing (RNA-seq) analysis of tracheal progenitors from

aforementioned fat body defective flies. The results revealed a dramatical alteration of

transcriptional program in tracheal progenitors upon the perturbation of fat body

(Figure 2-figure supplement 1A, B). Interestingly, the functional cluster of 'cytokine

activity' showed prominent enrichment in the differentially expressed genes (DEGs)

in progenitors from *lsp2>rpr.hid* pupae (Figure 2A). This raised the possibility that

certain cytokine-responsive signaling was induced in tracheal progenitors and the

signaling was compromised by impairment of fat body. Therefore, we proceeded to

analyze the expression of genes responsive to cytokine signaling. Analyzing the RNA-

seq data revealed that the cytokine-dependent JAK/STAT and Dpp signaling were

notably up-regulated upon the activation of progenitors (Figure 2B). Importantly, fat

body depletion led to suppression of target genes of JAK/STAT, PI3K and Dpp

signaling in tracheal progenitors, suggesting their dependence on the function of fat

body (Figure 2C).

To evaluate the roles of these signaling proteins, we perturbed their expression in fat

body by the expression of RNAi constructs. Knockdown of candidates including some

cytokines specifically in fat body did not affect the direction of tracheal progenitor

migration (Figure 2-figure supplement 1C-J), except for *upd2*, whose depletion

phenocopied fat body ablation-induced bidirectional movement of tracheal

progenitors (Figure 2D-F and Video 3). These results suggest a role of fat body-

produced Upd2 in remotely regulating the tracheal progenitors.

Then, we performed surface proteome *in vivo* (Li et al., 2020) to investigate the

spectrum of molecules received by trachea (Figure 2-figure supplement 2). The

trachea-associated proteins were biotinylated through a reaction mediated by a

membrane-tethered horse radish peroxidase (HRP-CD2) (Figure 2-figure supplement

2A-D). Of the 1684 streptavidin-precipitated proteins captured by mass spectrometry

(Figure 2G), a functional cluster enriched for receptor signaling via JAK/STAT was

identified (Figure 2H). The JAK/STAT pathway is one of the principal cellular

signaling that responds to Upd2 ligand (Hombria et al., 2005). *Drosophila* JAK/STAT

signaling is well-conserved (Arbouzova and Zeidler, 2006; Zeidler et al., 2000) and

comprises a single JAK (Hopscotch, Hop) and one STAT (Stat92E), in contrast to a

handful of homologues (four JAK and seven STAT genes) found in mammals.

Domeless (Dome), the receptor for JAK/STAT pathway, exhibited pronounced

expression in the tracheal progenitors (Figure 2-figure supplement 2E). To test if these

Dome receptors actively interact with their ligands, we adapted a technique to monitor

ligand-receptor interaction *in vivo* (Michel et al., 2011) and constructed a Dome

variant (DIPF) which only fluoresces in the ligand-binding and phosphorylated state

(Figure 2I). The signal of this DIPF reporter was detected in both larval fat body and

salivary gland (Figure 2-figure supplement 2F,H), which is consistent with active JAK

signaling implicated in the development of the tissues (Chakrabarti et al., 2016;

Krautz et al., 2020). When expressed in the tracheal system, DIPF displayed robust

fluorescent signal in the tracheal progenitors (Figure 2J). These data suggest that

receptor signaling of JAK/STAT is active in the tracheal progenitors.

To analyze the functional importance of JAK/STAT signaling in tracheal progenitors,

we perturbed the principal components of this signaling, namely the receptor Dome,

signal transducer Hop or the downstream transcription factor Stat92E, by *btl*-Gal4-

driven expression of RNAi constructs. Under these conditions in which JAK/STAT

pathway is compromised, the tracheal progenitors aberrantly migrated anteriorly,

which is reminiscent of *upd2* loss-of-function in the fat body (Figure 3A-E, Video 4

and Figure 3-figure supplement1A-D). The aberrant anterior migration of tracheal

progenitors upon perturbation of JAK/STAT components led to incomplete

regeneration of airway and impairment of tracheal integrity and caused melanization

in the trachea (Figure 3-figure supplement 1E-I). In agreement with genetic

perturbation of JAK/STAT signaling, pharmacological inhibition of JAK by a small-

molecule inhibitor, Tofacitinib (Palmroth et al., 2021), also triggered bidirectional

movement of tracheal progenitors (Figure 3F-H). The bidirectional movement was not

due to excessive progenitors or crowding (Figure 3-figure supplement 2A).

Concurrently, the activity of JAK/STAT pathway, as assessed by the Stat92E-GFP

reporter (Bach et al., 2007), was substantially impaired when components of the

pathway were depleted (Figure 3-figure supplement 2B-E, J). To determine whether

the tracheal JAK/STAT signaling depends on fat body-derived Upd2, we depleted

Upd2 in fat body and observed that Stat92E-GFP signal in tracheal progenitors was

severely decreased, suggesting that JAK/STAT signaling in the trachea requires fat

body-produced Upd2 (Figure 3-figure supplement 2F, G, J). Consistently, inhibition of

JAK/STAT signaling using Tofacitinib reduced the expression of Stat92E-GFP (Figure

3-figure supplement 2H-J). Taken together, these observations suggest that Upd2-

responsive JAK/STAT signaling in the trachea is essential for the disciplined

migration of progenitors.

Genes regulated by JAK/STAT signaling in the trachea

To gain a comprehensive understanding of the molecular details underlying the

discipline of tracheal progenitor migration, we conducted genomic chromatin

immunoprecipitation (ChIP-seq) to identify loci bound by Stat92E which functions as

the transcription factor of JAK/STAT pathway. This revealed a total of 21312 Stat92E

binding peaks, ~95.7% of which located within 2 kb of transcription start sites (TSSs)

of annotated genes (Figure 4-figure supplement 1A). In particular, 86% of the peaks

(18328 peaks) were enriched either in promoter regions or within gene bodies, and

66.1% of the peaks (13490 peaks) resided near the 5′ ends of annotated genes, namely

in the promoter regions, first exons and first introns (Figure 4-figure supplement 1B,

C). GO analysis of putative target genes of Stat92E identified one cluster associated

with establishment of planar polarity (Figure 4A). In line with this, the functional

class associated with establishment of planar polarity was also abundantly represented

among the differentially expressed genes (DEGs) upon the activation of tracheal

progenitors in larval-pupal transition (Figure 4B). Notably, Stat92E binding was

detected in the promoters and intronic regions of genes functioning in distal-to-

proximal signaling (Cho and Irvine, 2004), such as *dachsous* (*ds*), *four-jointed* (*fj*), *fz*,

stan, *Vang* and *fat2* (Figure 4C). Additionally, Stat92E occupied in the promoter

regions of *crb* and *yurt,* two genes involved in apical-basal polarity and tracheal tube

growth (Laprise et al., 2006; Schottenfeld-Roames and Ghabrial, 2012; Schottenfeld-

Roames et al., 2014) (Figure 4-figure supplement 1D). The enrichment of Stat92E in

the promoters and/or regulatory regions of these putative targets was confirmed by

ChIP-qPCR (Figure 4-figure supplement 1E).

To further validate these putative Stat92E targets and investigate their dependence on

JAK/STAT signaling, we analyzed their expression from several fosmid transgenes

which have a GFP tag fused to *ds*, *fj* or *ft* and express at endogenous levels. Ds and Fj

were abundant in the progenitor cells, but were vastly reduced upon depletion of

dome, *hop* or *stat92E*, suggesting that they are regulated by JAK/STAT pathway

(Figure 4D-M). Furthermore, it is reported that the function of Ft is influenced by

cell-autonomous increase of Ds level and its protein level is enhanced by Ds reduction

(Ambegaonkar et al., 2012; Matakatsu and Blair, 2004), which is also evidenced by

our analysis using *dsRNAi* and *UAS-ds* in the tracheal progenitors (Figure 4-figure

supplement 2A-D). In accordance with this notion, the level of Ft, as assayed by the

Ft-GFP reporter, was elevated by the reduction of JAK/STAT signaling (Figure 4N-

R). We also analyzed GFP-tagged fosmid transgenes of *fat2, crb* and *yurt* and found

that they were discernably reduced upon impairment of JAK/STAT signaling,

suggesting that they are also regulated by JAK/STAT pathway (Figure 4-figure

supplement 1F-T). Additionally, the transcription of *ds*, *fj*, *ft*, *fat2*, *crb*, and *yurt* was

compromised by expression of *stat92ERNAi* (Figure 4-figure supplement 1U). In sum,

these results suggest that JAK/STAT promotes components involved in the

establishment of polarity in tracheal cells.

The roles of JAK/STAT targets in the disciplined migration

To evaluate the functional roles of the polarity proteins in tracheal progenitor

migration, we perturbed their expression in the tracheal progenitors by expressing

RNAi against *ds*, *ft* or *fj*, which were identified by ChIP-seq as the targets of

JAK/STAT. In these flies, tracheal progenitors exhibited bidirectional movement,

which is reminiscent of the impairment of JAK/STAT signaling (Figure 5A-E and

Video 5). Similar observations were obtained by over-expression of *ft* or *ds* in the

trachea (Figure 5-figure supplement 1A-D), consistent with previous reports that both

loss- and gain-of-function of PCP components disrupt the planar cell polarity (Adler

et al., 2000; Tree et al., 2002; Vinson and Adler, 1987). The disciplined migration of

tracheal progenitors was also impaired by the expression of *fat2RNAi*, *crbRNAi*,

yurtRNAi or *scbRNAi* (Figure 5-figure supplement 1E-I), but was not affected by

perturbation of molecules involved in cell adhesion such as Enabled (Ena), Fak, E-

cadherin and Robo2 (Figure 5-figure supplement 1J).

Migratory cells generate protrusions at the leading edge to initiate movement (Cetera

et al., 2014). The normal posteriorly migrating tracheal progenitors extend protrusions

towards the migratory directions (Figure 5-figure supplement 2A), but in the

bidirectionally moving progenitors in which *upd2* in fat body was perturbed,

extensive filopodia were projected from both the anterior and posterior fronts (Figure

5-figure supplement 2B, C), indicating that the aberrantly anteriorly moving

progenitors may adopt the identity as those moving posteriorly. Bidirectionally

migrating progenitors induced by perturbation of JAK/STAT signaling did not alter

the expression of the tracheal inducer, *branchless* (*bnl*) (Figure 5-figure supplement

2D, E). Further analysis revealed that the progenitors exhibited elevated levels of Ft at

the leading edge where they attached to dorsal trunk (DT) (Figure 5F-G).

Accordingly, progenitors that underwent bidirectional movement exhibited

pronounced abundance of Ft at both the anterior and posterior frontal edges (Figure

5H-J). To further evaluate the functional roles of Ft-Ds-Fj module in disciplined

migration, we utilized the high-mobility carcinoma cells, SKOV-3, and found that

perturbation of Fj that phosphorylates the extracellular cadherin domains of both Ft

and Ds and modifies their heterophilic binding (Thomas and Strutt, 2012), Ft or Ds

concurrently displayed compromised directionality and reduced consistency of

movement in a two-dimensional culture (Figure 5K-M and Figure 5-figure

supplement 3). Together with the results in previous sections, these observations

suggest that the activated tracheal progenitors establish a disciplined migration

through the asymmetrical distribution of polarity proteins which is directed by an

Upd2-JAK/STAT signaling stemming from the remote organ of fat body.

Upd2 in the fat body-produced vesicle

Besides the JAK/STAT signaling, another functional class enriched for vesicle-

mediated transport was prominent from our surface proteome analysis of the trachea

(Figure 2H). A series of components that function in vesicle trafficking were

identified. It has been reported that IL-6 cytokines tend to be encapsulated in

secretory vesicles (Kandere-Grzybowska et al., 2003; Verboogen et al., 2018). To

visualize Upd2 production and investigate its transportation kinetics, an upd2-

mCherry transgene was developed and expressed under the control of *lsp2*-Gal4,

which enabled tracking the dynamics of Upd2 in fat body (Figure 6A). In agreement

with Upd2 being transported through vesicles, administration of L3 larvae with

Brefeldin A (BFA), which pharmacologically inhibits vesicle formation and transport,

sequestered Upd2 proteins in fat body (Figure 6B, C). To track the destination of the

Upd2-containing vesicles, we examined mCherry signals in adjacent tissues and

detected considerable amount of Upd2 puncta in the tracheal progenitors (Figure 6D).

BFA treatment reduced Upd2-mCherry puncta in the tracheal progenitors, suggesting

that tracheal progenitors receive vesicular Upd2 from the fat body (Figure 6E, F).

Perturbation of Grasp65, a Golgi reassembly stacking protein previously implicated in

Upd2 secretion (Rajan et al., 2017), also led to sequestration of Upd2-containing

vesicles in fat body (Figure 6G, H, J). The vesicle formation, function and

extracellular movement are dependent on the tetraspanin superfamily proteins

(Andreu and Yanez-Mo, 2014). We surveyed all the tetraspanin orthologs in fly for

potential roles in Upd2 vesicle formation and transport. When expressing *lbmRNAi* in

fat body, Upd2-containing vesicles were vastly increased (Figure 6I, J). Meanwhile,

perturbation of vesicle secretion or transport by expressing *grasp65RNAi* or *lbmRNAi*

in fat body eliminated the presence of fat body-origin Upd2 in the trachea, suggesting

that fat body-produced Upd2-containing vesicles function cell non-autonomously and

contribute to other tissues/organs (Figure 6K-N). It should be noted that knockdown

of *upd2* in the trachea did not alter the discipline of tracheal progenitor migration

(Figure 2-figure supplement 1K-M). Collectively, these results suggest that fat body-

produced Upd2 undergoes vesicle-mediated trafficking.

<u>The vesicular transport in JAK/STAT signaling</u>

The results in previous section suggest that the ligand of JAK/STAT signaling is

transported in a manner that depends on vesicle trafficking. To validate the role of fat

body-produced vesicles in inter-organ signaling, we used genetic and pharmacological

tools to perturb different processes of vesicle trafficking in fat body and monitored

JAK/STAT signaling in the tracheal progenitors. Expression of *grasp65RNAi* in fat

body reduced the activity of JAK/STAT signaling in the trachea, as assessed by the

Stat92E-GFP reporter (Figure 7A, B). Similarly, RNAi targeting expression of *lbm* in

fat body vanished JAK/STAT signal transduction in the trachea (Figure 7C). Rab

GTPases coordinate vesicle trafficking and production (Stenmark, 2009) and have

been shown to play pivotal roles in the regulation of intracellular trafficking of FGFR

and EGFR (Letizia et al., 2023; Olivares-Castineira and Llimargas, 2017), and were

identified in the surface proteome analysis. Consistently, knockdown of *rab5* or *rab7*

in fat body reduced the activity of JAK/STAT signaling in the progenitor cells (Figure

7D-F). Corroborating the genetic manipulations, BFA treatment that impeded

vesicular transport also resulted in impairment of JAK/STAT signaling in trachea

(Figure 7G-I). Taking advantage of the aforementioned DIPF reporter to assess the

response of receiving cells to ligands, we found that the fluorescent signal of DIPF

was compromised upon the presence of BFA, but was unaffected by inhibitors that

target the downstream JAK protein (Figure 7J-M), suggesting that signaling ligands

are less abundant in the recipient progenitor cells and that the vesicle-mediated

transport of ligands is essential for JAK/STAT signaling.

Phenotypically, the tracheal progenitors exhibited bidirectional migration in BFA-

treated flies, which phenocopies JAK/STAT loss-of-function (Figure 7N-P). In

concord with this observation, depletion of *grasp65* or *lbm* also led to bidirectional

movement (Figure 7Q-S'', and Video 6). Similar observations were made in tracheal

progenitors with either *rab5* or *rab7* knockdown (Figure 7T-V), whereas perturbation

of neither *rab2* nor *rab3* affected the disciplined progenitor migration (Figure 7-figure

supplement 1A-C). Taken together, these results suggest that JAK/STAT signaling in

the trachea is dependent on the vesicle-mediated transport of its ligands from fat body.

<u>The interaction between Upd2 and endocytic machinery</u>

Our results described thus far suggest that Upd2 emanating from fat body signals to

JAK/STAT signaling in the trachea. To further explore the molecular basis underlying

the vesicular transport of Upd2, we monitored Rab5-GFP and Rab7-GFP in fat body,

which mark early and late endosomes, respectively (Vonderheit and Helenius, 2005).

The fat body-produced Upd2 appeared vesicular (Figure 8A) and both Rab5 and Rab7

were found adjacent to the Upd2-harboring vesicles, suggesting that both Rab

GTPases function in the transport of Upd2 (Figure 8A'-B''). In contrast, Rab3

exhibited non-overlapping distribution with Upd2 (Figure 7-figure supplement 1E-

E''). Furthermore, we observed that Grasp65 was in close proximity to Upd2-

containing vesicles, indicating its integral roles in these vesicles (Figure 8C-C''). In

addition, Upd2 was observed to colocalize with the tetraspanin, Lbm (Figure 8D-D'').

At higher resolution, Upd2 and Lbm showed close association in a supramolecular

configuration (Figure 8D'''and 8D''''), corroborating its role in the transport of Upd2.

To determine if Upd2 interacts with the coordinators of vesicle trafficking, we

employed the Duolink in situ proximity ligation assay (PLA) which revealed strong

interactions between Upd2 and Rabs, such as Rab5 and Rab7 (Figure 8E-H), as well

as Lbm (Fig 8I, J). The interaction was further validated by the revelation that Upd2

co-immunoprecipitated with Rab5 and Rab7 (Figure 8K, L). The presence of Upd2 in

Lbm-containing vesicles was also evidenced in S2 cells (Figure 8-figure supplement

1A-C'') and co-IP experiment showed that Lbm physically associated with Upd2 in

both fat body and S2 cells (Figure 8M and Figure 8-figure supplement 1F). To further

understand the biogenesis of Lbm-containing vesicles that transport Upd2, we

conducted electron microscopic analysis of the Lbm-containing vesicles through the

expression of an HRP-fused Lbm in the fat body (Figure 8-figure supplement 1D).

The interaction between Lbm and Upd2 as simulated by Alphafold2 supported their

direct association (Figure 8-figure supplement 1E). Then, we generated an Lbm

chimera tagged with a pH-sensitive GFP variant, pHluorin (Yoshihara et al., 2005).

PHluorin fluorescence is squelched at the low pH domain such as in intravesicular

compartments, but becomes detectable when exposed to the extracellular

environment, thus enabling detection of exocytosis and endocytosis. Fat body

expressing Lbm-pHluorin produced GFP puncta at the plasma membrane (Figure 8N),

and the GFP signal was also detected in the trachea, suggesting the reception and

internalization of Lbm-containing vesicles by tracheal cells (Figure 8O). However, the

GFP fluorescence in both fat body and trachea was dramatically decreased by BFA

treatment, suggesting that Lbm-containing vesicles are diminished (Figure 8-figure

supplement 1G-J). Accordingly, the signals of Lbm-pHluorin in both fat body and

responding tracheoblasts were apparently compromised when Rab5 or Rab7 was

perturbed (Figure 8P-S), suggesting that the biogenesis and production of Lbm-

containing vesicles depend on Rab-mediated vesicle trafficking. Taken together, these

results suggest that fat body-derived Upd2 interacts with Rab-mediated endocytic

trafficking system to control the disciplined movement of tracheal progenitors.

Discussion

Resident stem cells and progenitors are mobilized to regenerate damaged or degenerated tissue. Despite the large distance between the niche where stem cells interact with their microenvironments and the destination for reconstruction, their commitment to a stereotyped track implicates sophisticated mechanism that controls disciplined migration as stem cells are activated and move out of the niche. While primary inducers expressed by damaged tissues are in play to coordinate the newly generated architecture with the degenerated counterpart (Chen and Krasnow, 2014), the present study elucidates an integral role of Upd2-JAK-STAT pathway in regulating the expression of polarity-related genes and maintaining the disciplined migration of tracheal progenitors (Figure 8T). The transport of Upd2 from fat body to trachea suggests intensive inter-organ communication during the migration of tracheal progenitors.

Several possibilities could account for the JAK/STAT-dependent polarity. The signaling components of JAK/STAT pathway could exhibit polarized localization (Sotillos et al., 2008). Alternatively, this signaling may activate genes controlling cell polarity and adhesion (Mallart et al., 2024; Tsurumi et al., 2011). Functional interplay between JAK/STAT signaling and cell polarity has been observed in various contexts (Chatterjee et al., 2023; Zeidler et al., 1999). Our results support a role of JAK/STAT signaling in promoting expression of genes with established roles in planar polarity, which may hallmark the route for the migration of the tracheal progenitor cells. Epithelial cells exhibit two aspects of polarity: apical-basal polarity and planar cell polarity (PCP). The latter refers to the collective alignment of cell polarity within the plane of an epithelial sheet (Zallen, 2007). Molecularly, PCP is generated by the asymmetry of a group of proteins (PCP proteins) that mediate communication between neighboring cells (Barlan et al., 2017a; Matis and Axelrod, 2013; Williams et al., 2022). In *Drosophila*, the components of PCP are considered to group functionally into two core modules. The seven-pass transmembrane protein Frizzled (Fz), the cytosolic proteins Dishevelled (Dsh), Diego (Dgo), the four-pass transmembrane protein Strabismus (Stbm, also known as Van Gogh (Vang)) and the cytosolic protein

Prickle (Pk) belong to the first module. The second module consists of Fat (Ft; also

known as cadherin-related tumor suppressor), Dachsous (Ds), Four-jointed (Fj) and

Atro (a transcription repressor) (Peng and Axelrod, 2012). Intensive functional

interplay occurs between these two modules (Ayukawa et al., 2014). Aberrant activity

of the core PCP proteins leads to misoriented hairs and complex swirling patterns (Ma

et al., 2003; Wong and Adler, 1993). In addition to arrangement of epithelial

appendages, PCP pathway is also required for collective and directed cell movements

(Munoz-Soriano et al., 2012). The migratory cell cohort is polarized into 'pioneer'

cells that lead the trailing followers (Vitorino and Meyer, 2008).

Our data indicate that expression of Ds, Fj, Fz, Stan and Fat2, core components or

regulators of PCP, depends on JAK/STAT pathway. The interaction between atypical

cadherin Fat (Ft) and its ligand, Dachsous (Ds) directs core protein asymmetry (Strutt

and Strutt, 2021; Yang et al., 2002). Phenotypically, aberrancy of PCP protein

abundance, either excessive core protein or deficit in expression gradients, gives rise

to similar morphological abnormality (Adler et al., 2000; Casal et al., 2002; Taylor et

al., 1998; Tree et al., 2002). Consistent with this notion, gain-of-function of Fj or Ds

phenocopies that of perturbation of PCP proteins. JAK/STAT signaling promotes the

expression of Ds, but reduces Ft expression (Figure 4). Thereby, perturbation of

JAK/STAT signaling disrupts the Ds-Ft system.

A precedent for JAK/STAT signaling in directional cell movement is border cell

migration from anterior to posterior compartment during *Drosophila* oogenesis.

Migration of the border cells is guided by a gradient of PDGF and VEGF chemokines

(Duchek et al., 2001). Loss of either *hop* (encoding JAK) or *stat* in the border cells

impinges their recruitment into the cluster and subsequent migration (Silver et al.,

2005). Our results suggest that JAK/STAT signaling does not serve as a guidance cue

for tracheal progenitors, but rather directs the directionality of cell movement. The

downstream PCP components may contribute to either polarity of progenitors or cell-

cell interactions between the progenitors and tracheal cells that they track along. It

remains unknown how individual progenitor cells perceive directional information

and convert it into group choreography.

We identified the fly fat body as the major source for the JAK/STAT signaling ligand,

Upd2 production. Fat body is functionally equivalent to the mammalian liver which

stores proteins, lipids and sugars and functions as an energy reservoir (Li et al., 2019).

It supplies proteins and/or hormones that are utilized by other organs, and thereby

serves as an interchange center to disperse systemic hormonal and nutritional signals.

For instance, it generates collagen IV to decorate imaginal discs and produces

xanthine dehydrogenase for eye pigmentation (Pastor-Pareja and Xu, 2011; Reaume et

al., 1989). The transport between fat body and trachea has been reported on a secreted

chitin deacetylase, Serpentine (Serp), which is expressed by fat body and contributes

to tracheal morphogenesis (Dong et al., 2014). Our results reveal that fat body also

signals to regulate the disciplined migration of tracheal progenitors through the

dispersion of Upd2 cytokines. These studies collectively suggest that fat body

orchestrates systemic tissue growth and patterning and that metabolic regulation is

critical for adult stem cells.

Proteins that are locally produced can execute systemic function in distant organs. A

possible route of transport is through the hemolymph or bloodstream and taken up by

the target tissues. The signaling proteins such as cytokines can be packaged in

extracellular vesicles with various dimensions (Buzas, 2023; Javeed et al., 2021). A

precedent of vesicular transport of signaling molecules is reported in migrasomes

whose diameter exceeds 500 nm (Jiang et al., 2019).

These extracellular vesicles mediate cell-to-cell communication (Colombo et al.,

2014), perhaps at a distance (Hood et al., 2011) and even traverse between organs

(Corrigan et al., 2014). The vesicular Upd2 is able to signal at recipient cells,

suggesting that the activity of Upd2 is preserved in the vesicle, and it is released upon

vesicle fusion. Compared with conventional extracellular vesicles such as exosomes,

the Upd2-containing vesicles possess larger dimension. Its production and trafficking

depend on GRASP-mediated unconventional secretion and interaction with Lbm.

Lbm belongs to the tetraspanin protein family that contains four transmembrane

domains. The mammalian homologs of tetraspanins, CD9, CD63, CD81 or CD37 are

principal constituents of extracellular vesicles. The Lbm-containing vesicles are

regulated by GRASP-mediated secretion and are sensitive to pharmacological

inhibition of EV transport.

It has been proposed that tetraspanins facilitate regeneration and wound healing in

cultural cells and single-cell plasma membrane. Tetraspanin-enriched macrodomains

are assembled into a ring-like structure (Huang et al., 2022), which is recruited to

large membrane wounds and promotes membrane repair (Wang et al., 2022). The

present study adds another dimension to the roles of tetraspanin proteins in tissue

regeneration which can be ascribed to transport of signaling proteins and modulation

of stem cells.

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-GFP (1:400)	Invitrogen	#A11122
Mouse anti-GFP (1:1000)	Abclonal	#AE012
Mouse anti-mCherry (1:1000)	Abclonal	#AE002
Rabbit anti-tubulin (1:5000)	Baoke	#BK7010
Phalloidin Alexa Fluor 640 (1:50)	Biotum	#00050
HRP-conjugated Streptavidin (1:5000)	Proteintech	#SA00001-0
Alexa Fluor 488 (1:200)	Abclonal	#AS053
Alexa Fluor 555 (1:200)	Abclonal	#AS007
DAPI	VECTASHIELD	#H1200
Reagents		
Halocarbon oil 700	Sigma	#H8898
RNeasy Micro Kit	QIAGEN	#74004
SMART-Seq v4 Ultra low input RNA Kit	Takara	#634889
AMPure XP	Beckman Coulter	#A63882
TruePrep DNA Library Prep Kit V2	Vazyme	#TD501
Anti-FLAG M2 Magnetic Beads	Millipore	#8823
BXXP	APEXBIO	#A8012
Experimental models: Organisms/strains		
D. melanogaster:UAS-upd2RNAi	Tsinghua Stock Center	THU1331, THU1288
D. melanogaster:UAS-domeRNAi	Tsinghua Stock Center	THU0574, THU5825
D. melanogaster:UAS-hopRNAi	Tsinghua Stock Center	THU5762, TH201501042.S
D. melanogaster:UAS-stat92ERNAi	Tsinghua Stock Center	THU0573, THU1915
D. melanogaster:UAS-lbmRNAi	Tsinghua Stock Center	THU2602, BDSC27278
D. melanogaster:UAS-grasp65RNAi	Tsinghua Stock Center	TH04282.N, THU1429
D. melanogaster:UAS-rab5RNAi	Tsinghua Stock Center	TH02192.N, THU0671
D. melanogaster:UAS-rab7RNAi	Tsinghua Stock Center	TH02539.N, THU2437
D. melanogaster:UAS-fjRNAi	Tsinghua Stock Center	THU201500988.S, THU1538
D. melanogaster:UAS-fat2RNAi	Tsinghua Stock Center	THU4120, VDRC27113
D. melanogaster:UAS-yurtRNAi	Tsinghua Stock Center	THU1740, VDRC28674
D. melanogaster:UAS-crbRNAi	Tsinghua Stock Center	THU2783, THU5212
D. melanogaster:UAS-scbRNAi	Tsinghua Stock Center	THU3905, THU2707
D. melanogaster:UAS-dsRNAi	Tsinghua Stock Center	THU2846, VDRC36219
D. melanogaster:UAS-ftRNAi	Tsinghua Stock Center	TH201500989.S, VDRC9396
D. melanogaster:10XStat92E-GFP	Tsinghua Stock Center	THJ0273
D. melanogaster:Stat92E-GFP	BDSC	#38670
D. melanogaster:UAS-grasp65-GFP	BDSC	#8507
D. melanogaster:Ds::GFP	BDSC	#59425
D. melanogaster:Yurt::GFP	VDRC	#318067
D. melanogaster:Crb::GFP	VDRC	#318384
D. melanogaster:Ft::GFP	VDRC	#318477
D. melanogaster:Fj::GFP	VDRC	#318457
D. melanogaster:UAS-dsRNAi	VDRC	#36219
D. melanogaster:UAS-ftRNAi	VDRC	#9396
D. melanogaster:UAS-ft	(Brittle et al., 2010)	
D. melanogaster:UAS-ds	(Brittle et al., 2012)	
D. melanogaster: fat2-GFP	(Barlan et al., 2017b)	
D. melanogaster:UAS-GFP-Rab7	BDSC	#9779
D. melanogaster:UAS-GFP-Rab5	BDSC	#24616
D. melanogaster:hop^{Tum}/FM7C	BDSC	#8492
D. melanogaster:stat92E^{F}	BDSC	#24757
D. melanogaster:dome^{G0264}	Kyoto	#111866

500 **Materials and methods**

Primer α-tubulin84b Forward:CACACCACCCTGGAGCATTC	This paper	N/A
Primer α-tubulin84b Reverse:CCAATCAGACGGTTCAGGTTG	This paper	N/A
Primer upd2 Forward:TCAATCCGTATCGCGGTCTG	This paper	N/A
Primer upd2 Reverse:AGAAGAGTCGCAGGTTGTGG	This paper	N/A
Primer ds Forward:ACAACCGAACTCGAACCGAA	This paper	N/A
Primer ds Reverse:AGTAGCATCACACACAAGTGA	This paper	N/A
Primer ft Forward:CTGGATCGAGAGCAGCAGAG	This paper	N/A
Primer ft Reverse:GACGGTAAATTCTCGCGCAC	This paper	N/A
Primer fj Forward:ATTACTCAAGCGGTTGGGGG	This paper	N/A
Primer fj Reverse:CGGTTCCTGTTCCTGTCTCC	This paper	N/A
Primer fat2 Forward:TATCTGCGCCCATACGCATT	This paper	N/A
Primer fat2 Reverse:TCTCATCGGCCTTGCTTTGT	This paper	N/A
Primer yurt Forward:GGTCAGCTCAGGGTGACTATC	This paper	N/A
Primer yurt Reverse:ATTGGTAAGCTTGGCGTTGC	This paper	N/A
Primer crb Forward:CAGCAGTGTTTGAACGGTGG	This paper	N/A
Primer crb Reverse:AGGCAGTGACCAATGGGG	This paper	N/A
Software and algorithms		
Fiji / ImageJ	NIH	RRID:SCR_002285
GraphPad Prism 8.0	GraphPad Software	https://www.graphpad.com/scientific-software/prism/
Zen 3.1	Zeiss	https://www.zeiss.com.cn/corporate/home.html
PCA-flow	Bradski, G.[79]	https://www.drdobbs.com/open-source/the-opencv-library
Deposited data		
Raw and processing Data	This paper	GSE256177 https://www.iprox.cn/page/PSV023.html;?url=170832392127 7ZyfC qvupguyijvyndyf

502   **Key Resources Table**

Fly lines and husbandry

All flies were reared on normal cornmeal and agar medium at 25°C unless noted.
UAS-upd2RNAi (THU1331, THU1288), UAS-domeRNAi (THU0574), UAS-
hopRNAi (THU5762), UAS-stat92ERNAi (THU0573), UAS-lbmRNAi (THU2602),
UAS-grasp65RNAi (TH04282.N), UAS-rab5RNAi (TH02192.N), UAS-rab7RNAi
(TH02539.N), UAS-fjRNAi (THU201500988.S), UAS-fat2RNAi (THU4120), UAS-
yurtRNAi (THU1740), UAS-crbRNAi (THU2783) and UAS-scbRNAi (THU3905)
were ordered from Tsinghua Stock Center. Stat92E-GFP (BSDC#38670), UAS-
grasp65-GFP (BSDC#8507), Ds::GFP (BSDC#59425) were from Bloomington
Drosophila Stock Center. Yurt::GFP (v318067), Crb::GFP (v318384), Ft::GFP
(v318477), Fj::GFP (v318457), UAS-dsRNAi (v36219, THU2846), and UAS-ftRNAi
(v9396) were obtained from VDRC. UAS-ft was kindly provided by Dr. Xianjue Ma,
UAS-ds was kindly provided by Dr. Xing Wang and UAS-fat2-GFP was from Dr.
Shunfan Wu. UAS-GFP-rab5 and UAS-GFP-rab7 were kindly provided by Dr.
Xiaohang Yang.

Plasmid construction and transgenic flies

To generate UAS-upd2-mCherry and UAS-lbm-GFP, UAS-lbm-HRP transgenic flies,
the coding sequence of upd2 or lbm was PCR amplified from a fly cDNA library and
cloned into a pUAST vector with C-terminal mCherry, GFP or HRP.

The DIPF reporter was generated by first fusing cpYFP and *Drosophila* FKBP12 and
subsequently ligating to the dome cDNA via a GTG linker. The above product was
then cloned into a pUAST vector and verified by DNA sequencing, and injected into
y[1] M{vasint.Dm}ZH-2A w[*]; P{CaryP}attP2 recipient flies or *w1118* following
standard *Drosophila* transformation injection procedures (Core Facility of Drosophila
Resource and Technology, SIBCB, CAS).

Cell culture and transfection

S2 cells were grown in Schneider Drosophila Medium (Gibco, #21720024) supplemented with 10% (v/v) Fetal Bovine Serum (FBS, Gibco, #10099141C) and 1% (v/v) Penicillin Streptomycin (Pen/Strep, Life Technologies) at 28°C with 0.2% CO_2. Transfection was conducted with 5 μg plasmids (act-GAL4, UAS-lbm-GFP, UAS-upd2-mCherry) using Effectene Transfection Reagent (QIAGEN, #301425).

SKOV3 cells were cultured in DMEM medium (CR#12800) containing 10% FBS (Gibco, #10099141C) in an incubator with 5% CO2 at 37 °C. Cells with ~80% confluency were infected with lentivirus loaded with siRNAs for gene knockdown or gRNAs for knockout, and the medium was replaced 24 hours post-infection. 72 hours after lentivirus infection, bright-field images were taken every 5 minutes for 12 hours using a confocal microscope.

Quantitative reverse transcription PCR

Larval or pupal trachea were dissected in cold PBS, and then transferred to RNA extraction reagent (AG21101). Next, reverse transcription was performed using qPCR RT Mix with gDNA Remover reagent (AG11706). qPCR was performed using the Universal SYBR Select Master Mix (AG11701) with a Bio-Rad system. The foldchange of target gene expression was normalized to that of α-tubulin. The primers are listed in Key Resources Table.

Western Blotting and Co-immunoprecipitation

Total protein was extracted from cells or tissues by RIPA buffer supplemented with a protease inhibitor cocktail (Merck, #11836170001) and phenylmethanesulfonyl fluoride (Beyotime, #ST507), separated by 10% SDS-PAGE gels and transferred to PVDF membrane (Millipore, #IPVH00010). Blots were detected with an ECL Western Blotting detection system (Bio-Rad). For co-immunoprecipitation, lysates of larval fat body or transfected S2 cells were incubated overnight at 4°C with protein A Magnetic beads (Thermo Scientific, #2736141) pre-coated with GFP antibody (Invitrogen, #A11122). Immunoprecipitates were eluted in SDS-containing loading

559 buffer for subsequent immunoblotting analysis. Antibodies for immunoblotting

560 include: α-tubulin (Baoke, #BK7010), α-HRP-conjugated Streptavidin (Proteintech,

561 #SA00001-0), α-GFP (Abclonal, #AE012, 1:1000); α-mCherry (Abclonal, #AE002,

562 1:1000).

Proximity Ligation Assay

Proximity ligation assay (PLA) was carried out with Duolink® In Situ Detection

Reagents Far Red (Sigma Aldrich, #DUO92013) according to manufacturer's

instructions, using the probes anti-rabbit PLUS (Sigma Aldrich, #DUO92002) and

anti-mouse MINUS (Sigma Aldrich, #DUO92004). Briefly, larval fat body was

dissected and fixed in 4% formaldehyde. The animals not expressing upd2-mCherry

served as controls. After permeabilization, the samples were incubated with primary

antibodies overnight at 4°C. Then, the samples were washed with PLA buffer A,

hybridized with PLA probes, ligated, and amplified. Samples were washed twice with

PLA buffer B (Sigma Aldrich, #DUO82049) and fluorescence images were taken with

an LSM Zeiss 900 inverted confocal laser scanner microscope.

Immunofluorescence

Trachea from white pupae (0hr APF) were dissected in PBS and fixed with 4%

formaldehyde for 25 min at room temperature. After washes, trachea samples were

permeabilized with 1% TritonX-100 in PBS, and then blocked in 10% goat serum.

Incubation with primary antibody (GFP, 1:400; lacZ, 1:40) was performed at 4°C with

gentle rotation for overnight. Then, the samples were incubated with secondary

antibodies conjugated to Alexa Fluor 488 or 555 (1:200) and Phalloidin (1:50) for 2

hrs. After washing, samples were mounted in antifade mounting medium with DAPI

(VECTASHIELD) and imaged under an LSM Zeiss 900 inverted confocal laser

scanner microscope.

Live imaging of pupal trachea stem cells

White pupae (0hr APF) were briefly washed in double distilled water and mounted in

halocarbon oil 700 (Sigma, #H8898). The pupae were positioned with forceps to bring

a single dorsal trunk of the trachea up for optimal imaging of Tr4 and Tr5 metameres.

Then, pupae were immobilized by a 22x30 mm No.1.5 high precision coverslip

spaced by vacuum grease. Time-lapse images were captured by an LSM Zeiss 900

inverted confocal laser scanner microscope. For migration distance measurement, we

took sequential snapshots of the moving progenitors of pupae staged at 0hr, 1hr, 2hr

and 3hr APF. The migration distance was measured as the distance from the starting

position (the junction of TC and DT) to the leading edge of progenitor groups. The

migration velocity was calculated by v=d (micrometer)/t (min).

RNA sequencing of tracheal progenitors

Total RNA was isolated from the Tr4 and Tr5 metamere progenitors dissected from

1hr APF pupae using RNeasy Micro Kit (QIAGEN, #74004). SMART-Seq v4 Ultra

low input RNA Kit (Takara, #634889) was used for first-strand and second-strand

cDNA synthesis and double-stranded cDNA end repair. Double-stranded cDNAs were

cleaned using AMPure XP (Beckman Coulter, #A63882). Then cDNAs were

subjected to tagmentation and ligation to adaptors to generate the sequencing libraries

using TruePrep DNA Library Prep Kit V2 for Illumina kit (Vazyme, #TD501). The

quality and concentration of the libraries were assessed using the Agilent High

Sensitivity DNA Kit and Bioanalyzer 2100 (Agilent Technologies) and submitted to

150 bp paired-end high throughput sequencing using Hiseq4000 (Illumina).

Analysis of RNA-seq data was performed using a computer system equipped with

multiple processors. Clean reads were mapped to the *Drosophila* genome sequence

using Hisat2 with default parameters. Successfully mapped reads were counted using

FeatureCounts. Differential gene expression analysis was performed using the

DESeq2 package. Adjusted *p*-value < 0.05 was used as the threshold to identify the

differentially expressed genes. Gene ontology and KEGG pathway enrichment

analyses for the differentially expressed genes were conducted using the Database for

Annotation, Visualization and Integrated Discovery (DAVID).

Chromatin Immunoprecipitation

Third instar larval trachea from Stat92E-Flag (BDSC, #38670) were fixed in 1%

formaldehyde. The fixation reaction was terminated by adding glycine (125 mM).

Trachea were washed and resuspended in lysis buffer, and sonicated to generate 200-

600 bp DNA fragments. Procedures of immunoprecipitation and ChIP sequencing

library construction were as previously described (Li et al., 2022). Anti-FLAG M2

Magnetic Beads (Millipore, #8823) were used for enriched DNA binding to

transcription factor Stat92E.

Immunoprecipitated DNA was subjected to next-generation sequencing using the

Epicenter Nextera DNA Sample Preparation Kit or to real-time PCR. Library

construction was performed using the High Molecular Weight tagmentation buffer

(source), and tagmented DNA was linearly amplified by PCR. The libraries were then

sequenced on a Novaseq according to manufacturer's standard protocols. The

sequences were processed using *Fastqc* and low-quality bases and adaptor

contamination were trimmed by *cutadapt*. Filtered reads were mapped to *Drosophila*

genome using *BWA mem* algorithm. Peaks were called using macs2 *callpeak* (Zhao et

al., 2019) and plotted using pyGenomeTracks. GO analysis of biological processes

was conducted by DAVID.

Cell-surface proteomics of fly trachea

Trachea from white pupae were dissected in pre-cooled Schneider Medium (Gibco)

and collected in 1.5 mL low-binding tube (Axygen) containing 500 μL Schneider

Medium. The samples were washed with 500 μL fresh medium and incubated with

100 μM BXXP (APEXBIO, #A8012) for 1 hr on ice with occasional pipetting.

Labeling reaction was initiated by adding 1 mM (0.03%) H_2O_2 to the sample-

containing medium and proceeded for 7 minutes at room temperature. The reaction

was immediately quenched by five thorough washes with PBS containing 10 mM

sodium ascorbate (Aladdin, #S105024) and 5 mM Trolox (APEXBIO, #C3183). For

biochemical characterization or proteomic sample preparation, the quenching solution

was drained, and the trachea in minimal residual quenching solution were quickly

frozen in liquid nitrogen and stored at 80°C. LC-MS/MS analysis was performed

using a Q Exactive HF-X instrument (Thermo Fisher) coupled with Easy-nLC 1200

system. The acquired MS raw data were processed using MaxQuant version 2.0.1.0

(Max Planck Institute of Biochemistry, Germany). Label-free quantification (LFQ)

was set with a default parameter and iBAQ was selected.

Transmission electron microscopy

Fat body of third instar larvae were dissected and fixed in 0.12 M Na-cacodylate

buffer (pH 7.4) containing 2.5% glutaraldehyde for 1 hr on ice. Then the samples

were rinsed in 0.12 M Na-cacodylate buffer (6 × 5 min, on ice). The dissected fat

body were pre-incubated with DAB (10 mg/20 ml) in 0.12 M Na-cacodylate buffer

(containing 0.1% saponin) for 30 minutes with agitation in the dark. Then 30% H_2O_2

was quickly mixed in DAB solution to a 0.03% v/v concentration and reacted for 30

min at RT. The fat body were transferred to 0.12 M Na-cacodylate buffer (6 × 5 min,

RT). To increase the electron density of the HRP/DAB product, samples were

transferred into 0.01% OsO4 in 0.12 M Na-cacodylate buffer (pH 7.4) for 10 min at

RT, then rinsed in 0.12 M Na-cacodylate buffer (3 × 10 min, on ice). 0.1%

thiocarbohydrazide in 0.12 M Na-cacodylate buffer (pH 7.4) was used for 10 min at

RT, then rinsed in 0.12 M Na-cacodylate buffer (3 × 10 min, on ice). After post-

fixation for 1 hour at RT with 1% OsO4 in 0.12 M Na-cacodylate buffer (pH 7.4), the

samples were rinsed with MilliQ water (3 × 5 min, RT) and dehydrated in a series of

15 min with 10%, 30%, 50%, 70%, 90% and 100% (3×) ethanol. Infiltration was

conducted at RT with a mixture of acetone and resin 1:1 for 1.5 h, 1:2 for 3 h and 1:3

overnight. The tissues were then dissected from the carcasses and placed in block

molds filled with resin for hardening at 60°C during 48 h. 70 nm ultrathin section

from the hardened blocks were cut on a Leica EM UC7 ultramicrotome using an Ultra

45° diamond knife and imaged in a thermos scientific Talos L120C electron

microscope.

Numeration of tracheal progenitors

The pupae at indicated stages were dissected and then fixed with 4% PFA for 25

minutes. The nuclei were labeled by DAPI. The samples were imaged by an LSM

Zeiss 900 inverted confocal laser scanner microscope. The number of progenitor cells

was scored from image stacks.

Optic Flow Analysis

<u>Motion Correlation</u>

Time-lapse images of trachea progenitor cells were captured every 5 minutes over a

total duration of 2 hrs using an LSM Zeiss 900 microscope. The movies and images

were subjected to a three-step motion collection using ImageJ (Schneider et al., 2012)

as follows:

(1) Images were transformed to grey scale.

(2) Automated stabilization was performed on grey images by image stabilizer.

(3) The feature points were mapped and affine transformation was applied using big

warp (Bogovic et al., 2015).

<u>Optical flow</u>

Optical flow represents the pattern of motion of pixels in a sequence of images.

Between two consecutive frames $I(x, y, t)$ and $I(x, y, t + \Delta t)$, the optical flow vector

$\mathbf{v} = \left(v_x, v_y\right)$ represents the motion of pixels during this time. The optical flow

constraint equation is shown below:

$$I_x v_x + I_y v_y + I_t = 0$$

We adopted the PCA-flow algorithm (Bradski, 2000) in openCV library

(cv::optflow::OpticalFlowPCAFlow) in which the sparse optical flow vectors within

each small region of the image sequence are calculated before training optical flow

fields via principal component analysis (PCA) (Wulff and Black, 2015). The vectors

were assembled to generate a smooth vector field of optic flow using the learning

linear models of flow. The PCA-flow was validated for the efficiency and robustness.

<u>Variance in 1D axis</u>

We developed a robust estimator to evaluate the variance of optic flow projection

along the 1D migration axis between samples. The vector length is normalized by a

frame-specific normalization factor κ to fit with scale of optic flow computed among

different frames.

$$\kappa = Q_{0.9}\{||v||_2 \mid v \in vectorfield\}$$

where Q is the quantile function. The variance is computed as

$\text{Var}(\{||v||_2 > Q_{0.2} \mid v \in vectorfield\})$ and scaled to 0-1 by $\text{Var}(x)/\big(\text{Var}(x) +$

$\text{Var}(y)\big)$.

<u>Random Variable</u>

The direction of optic flow in each volume was assigned to 'left' or 'right'. Then the

distribution of the binarized directions is modeled as a Bernoulli random variable

X ~ Bernoulli(p) with PMF

$$P(X = x) = \begin{cases} p & \text{if } x = 1, \\ 1 - p & \text{if } x = 0. \end{cases}$$

708 <u>Binary Entropy</u>

709 The entropy for Bernoulli random variable X ~ Bernoulli(p) is defined as

$$H_{binary}(X) = -p\log(p) - (1 - p)\log(1 - p)$$

710   The entropy evaluates the information contained in the random variable (also called

711   uncertainty). In this case, when $p = 0.5$, it reaches the maxima 1; if the variable is

712   determinate (i.e. p=0 or 1), the entropy is zero. More directed cell migration leads to a

713   lower entropy in optic flow since the certainty is high for the migration direction. We

714   estimated p by computing ratio of left and right direction

$$p = \frac{\#left}{\#left + \#right}$$

Image and statistical analysis

Confocal z-stack images were analyzed to extract information of fluorescent intensity

of stat92E-GFP and DIPF, and the number of Upd2-mCherry-containing vesicles. z

slices of fluorescent intensity for Ds, Ft and Fj were measured. The number of

particles for DIPF and Upd2-mCherry-containing vesicles was analyzed using

ImageJ. All statistical analysis was conducted using GraphPad Prism 8.0. Mean and

standard error of the mean (SEM) were shown. Unpaired *t*-tests with Benjamin's

correction were used to evaluate statistical significance between groups.

Acknowledgements

We thank Drs Thomas Kornberg, David Strutt, Xiaohang Yang, Markus Affolter, Mark Krasnow,

Stefan Luschnig, Kai Yuan, Xing Wang, Zhouhua Li, Xianjue Ma, Sarah Bray, Yiming Zheng

and Shunfan Wu for generously providing reagents; Core Facility of Drosophila Resource and

Technology, SIBCB, CAS for injection service; Bloomington Drosophila Stock Center, Kyoto

Stock Center, Vienna stock center, Tsinghua Stock Center for fly stocks; Developmental Studies

Hybridoma Bank for antibodies; all members of Huang lab and Kornberg lab for discussions and

constructive suggestions. This work has been financially supported by NSFC92168101,

NSFC32070784 and Thousand Young Talent Program to H.H., NSFC32000574 to H.G.W.,

National Key R&D Program of China (2022YFF0608402) and Chinese National Natural Science

Funds (22374128) to B.Y. and NSFC32300699, LQ24C120001 and Postdoctoral Fellowship

Foundation 2023M733098 to Y.L..

Author contributions

Conceptualization: H.H., P.Z.D.; Methodology: B.Y., P.Z.D., T.F.L., T.Y.G;

Investigation: P.Z.D., Y.L., Y.Y.W., Q.Z., T.F.L., H.G.W; Supervision: J.M., H.H.,

H.G.W, B.Y.; Writing: J.M., H.H., P.Z.D, H.G.W.

Declaration of interests

The authors declare no competing interests.

References

Adler, P.N., Taylor, J., and Charlton, J. (2000). The domineering non-autonomy of frizzled and van Gogh clones in the Drosophila wing is a consequence of a disruption in local signaling. Mech Dev *96*, 197-207.

Agaisse, H., Petersen, U.M., Boutros, M., Mathey-Prevot, B., and Perrimon, N. (2003). Signaling role of hemocytes in Drosophila JAK/STAT-dependent response to septic injury. Dev Cell *5*, 441-450.

Ambegaonkar, A.A., Pan, G., Mani, M., Feng, Y., and Irvine, K.D. (2012). Propagation of Dachsous-Fat planar cell polarity. Curr Biol *22*, 1302-1308.

Andreu, Z., and Yanez-Mo, M. (2014). Tetraspanins in extracellular vesicle formation and function. Front Immunol *5*, 442.

Arbouzova, N.I., and Zeidler, M.P. (2006). JAK/STAT signalling in Drosophila: insights into conserved regulatory and cellular functions. Development *133*, 2605-2616.

Ayukawa, T., Akiyama, M., Mummery-Widmer, J.L., Stoeger, T., Sasaki, J., Knoblich, J.A., Senoo, H., Sasaki, T., and Yamazaki, M. (2014). Dachsous-dependent asymmetric localization of spiny-legs determines planar cell polarity orientation in Drosophila. Cell Rep *8*, 610-621.

Bach, E.A., Ekas, L.A., Ayala-Camargo, A., Flaherty, M.S., Lee, H., Perrimon, N., and Baeg, G.H. (2007). GFP reporters detect the activation of the Drosophila JAK/STAT pathway in vivo. Gene Expr Patterns *7*, 323-331.

Barlan, K., Cetera, M., and Horne-Badovinac, S. (2017a). Fat2 and Lar Define a Basally Localized Planar Signaling System Controlling Collective Cell Migration. Dev Cell *40*, 467-477 e465.

Barlan, K., Cetera, M., and Horne-Badovinac, S. (2017b). Fat2 and Lar Define a Basally Localized Planar Signaling System Controlling Collective Cell Migration. Dev Cell *40*, 467-477.e465.

Beshel, J., Dubnau, J., and Zhong, Y. (2017). A Leptin Analog Locally Produced in the Brain Acts via a Conserved Neural Circuit to Modulate Obesity-Linked Behaviors in Drosophila. Cell Metab *25*, 208-217.

Bogovic, J.A., Hanslovsky, P., Wong, A., and Saalfeld, S. (2015). Robust registration of calcium images by learned contrast synthesis. 2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI), 1123-1126.

Bradski, G. (2000). The OpenCV library. Dr Dobb's J Software Tools *25*, 120-125.

Brittle, A., Thomas, C., and Strutt, D. (2012). Planar Polarity Specification through Asymmetric Subcellular Localization of Fat and Dachsous. Current Biology *22*, 907-914.

Brittle, A.L., Repiso, A., Casal, J., Lawrence, P.A., and Strutt, D. (2010). Four-jointed modulates growth and planar polarity by reducing the affinity of dachsous for fat. Curr Biol *20*, 803-810.

Buzas, E.I. (2023). The roles of extracellular vesicles in the immune system. Nat Rev Immunol *23*, 236-250.

Casal, J., Struhl, G., and Lawrence, P.A. (2002). Developmental compartments and planar polarity in Drosophila. Curr Biol *12*, 1189-1198.

Cetera, M., Ramirez-San Juan, G.R., Oakes, P.W., Lewellyn, L., Fairchild, M.J., Tanentzapf, G., Gardel, M.L., and Horne-Badovinac, S. (2014). Epithelial rotation promotes the global alignment of contractile actin bundles during Drosophila egg chamber elongation. Nat Commun *5*, 5511.

Chakrabarti, S., Dudzic, J.P., Li, X., Collas, E.J., Boquete, J.P., and Lemaitre, B. (2016). Remote Control of Intestinal Stem Cell Activity by Haemocytes in Drosophila. PLoS Genet *12*, e1006089.

Chatterjee, D., Cong, F., Wang, X.F., Machado Costa, C.A., Huang, Y.C., and Deng, W.M. (2023). Cell

polarity opposes Jak/STAT-mediated Escargot activation that drives intratumor heterogeneity in a

Drosophila tumor model. Cell Rep *42*, 112061.

Chen, F., and Krasnow, M.A. (2014). Progenitor outgrowth from the niche in Drosophila trachea is

guided by FGF from decaying branches. Science *343*, 186-189.

Chen, H.W., Chen, X., Oh, S.W., Marinissen, M.J., Gutkind, J.S., and Hou, S.X. (2002). mom identifies

a receptor for the Drosophila JAK/STAT signal transduction pathway and encodes a protein distantly

related to the mammalian cytokine receptor family. Genes Dev *16*, 388-398.

Cherbas, L., Hu, X., Zhimulev, I., Belyaeva, E., and Cherbas, P. (2003). EcR isoforms in Drosophila:

testing tissue-specific requirements by targeted blockade and rescue. Development *130*, 271-284.

Cho, E., and Irvine, K.D. (2004). Action of fat, four-jointed, dachsous and dachs in distal-to-proximal

wing signaling. Development *131*, 4489-4500.

Colombo, M., Raposo, G., and Thery, C. (2014). Biogenesis, secretion, and intercellular interactions of

exosomes and other extracellular vesicles. Annu Rev Cell Dev Biol *30*, 255-289.

Corrigan, L., Redhai, S., Leiblich, A., Fan, S.J., Perera, S.M., Patel, R., Gandy, C., Wainwright, S.M.,

Morris, J.F., Hamdy, F.*, et al.* (2014). BMP-regulated exosomes from Drosophila male reproductive

glands reprogram female behavior. J Cell Biol *206*, 671-688.

Darnell, J.E., Jr. (1997). STATs and gene regulation. Science *277*, 1630-1635.

Dong, B., Miao, G., and Hayashi, S. (2014). A fat body-derived apical extracellular matrix enzyme is

transported to the tracheal lumen and is required for tube morphogenesis in Drosophila. Development

141, 4104-4109.

Duchek, P., Somogyi, K., Jekely, G., Beccari, S., and Rorth, P. (2001). Guidance of cell migration by

the Drosophila PDGF/VEGF receptor. Cell *107*, 17-26.

Ertekin, D., Kirszenblat, L., Faville, R., and van Swinderen, B. (2020). Down-regulation of a cytokine

secreted from peripheral fat bodies improves visual attention while reducing sleep in Drosophila. PLoS

Biol *18*, e3000548.

Fuchs, E., and Blau, H.M. (2020). Tissue Stem Cells: Architects of Their Niches. Cell Stem Cell *27*,

532-556.

Heinrich, P.C., Behrmann, I., Haan, S., Hermanns, H.M., Muller-Newen, G., and Schaper, F. (2003).

Principles of interleukin (IL)-6-type cytokine signalling and its regulation. Biochem J *374*, 1-20.

Hombria, J.C., Brown, S., Hader, S., and Zeidler, M.P. (2005). Characterisation of Upd2, a Drosophila

JAK/STAT pathway ligand. Dev Biol *288*, 420-433.

Hood, J.L., San, R.S., and Wickline, S.A. (2011). Exosomes released by melanoma cells prepare

sentinel lymph nodes for tumor metastasis. Cancer Res *71*, 3792-3801.

Huang, Y., Zhang, X., Wang, H.W., and Yu, L. (2022). Assembly of Tetraspanin-enriched

macrodomains contains membrane damage to facilitate repair. Nat Cell Biol *24*, 825-832.

Ingaramo, M.C., Sanchez, J.A., Perrimon, N., and Dekanty, A. (2020). Fat Body p53 Regulates

Systemic Insulin Signaling and Autophagy under Nutrient Stress via Drosophila Upd2 Repression. Cell

Rep *33*, 108321.

Javeed, N., Her, T.K., Brown, M.R., Vanderboom, P., Rakshit, K., Egan, A.M., Vella, A., Lanza, I., and

Matveyenko, A.V. (2021). Pro-inflammatory beta cell small extracellular vesicles induce beta cell

failure through activation of the CXCL10/CXCR3 axis in diabetes. Cell Rep *36*, 109613.

Jiang, D., Jiang, Z., Lu, D., Wang, X., Liang, H., Zhang, J., Meng, Y., Li, Y., Wu, D., Huang, Y.*, et al.*

(2019). Migrasomes provide regional cues for organ morphogenesis during zebrafish gastrulation. Nat

Cell Biol *21*, 966-977.

Jones, D.L., and Wagers, A.J. (2008). No place like home: anatomy and function of the stem cell niche.
Nat Rev Mol Cell Biol *9*, 11-21.

Kandere-Grzybowska, K., Letourneau, R., Kempuraj, D., Donelan, J., Poplawski, S., Boucher, W.,
Athanassiou, A., and Theoharides, T.C. (2003). IL-1 induces vesicular secretion of IL-6 without
degranulation from human mast cells. J Immunol *171*, 4830-4836.

Krautz, R., Khalili, D., and Theopold, U. (2020). Tissue-autonomous immune response regulates stress
signaling during hypertrophy. eLife *9*, e64919.

Laprise, P., Beronja, S., Silva-Gagliardi, N.F., Pellikka, M., Jensen, A.M., McGlade, C.J., and Tepass,
U. (2006). The FERM protein Yurt is a negative regulatory component of the Crumbs complex that
controls epithelial polarity and apical membrane size. Dev Cell *11*, 363-374.

Letizia, A., Espinas, M.L., Giannios, P., and Llimargas, M. (2023). The TNFR Wengen regulates the
FGF pathway by an unconventional mechanism. Nat Commun *14*, 5874.

Li, J., Han, S., Li, H., Udeshi, N.D., Svinkina, T., Mani, D.R., Xu, C., Guajardo, R., Xie, Q., Li, T.*, et*
al. (2020). Cell-Surface Proteomic Profiling in the Fly Brain Uncovers Wiring Regulators. Cell *180*,
373-386 e315.

Li, L., and Clevers, H. (2010). Coexistence of quiescent and active adult stem cells in mammals.
Science *327*, 542-545.

Li, S., Yu, X., and Feng, Q. (2019). Fat Body Biology in the Last Decade. Annu Rev Entomol *64*, 315-
333.

Li, Y., Dong, P., Yang, Y., Guo, T., Zhao, Q., Miao, D., Li, H., Lu, T., Xia, F., Lyu, J.*, et al.* (2022).
Metabolic control of progenitor cell propagation during Drosophila tracheal remodeling. Nat Commun
13, 2817.

Ma, D., Yang, C.H., McNeill, H., Simon, M.A., and Axelrod, J.D. (2003). Fidelity in planar cell
polarity signalling. Nature *421*, 543-547.

Ma, J., Chen, T., Wu, S., Yang, C., Bai, M., Shu, K., Li, K., Zhang, G., Jin, Z., He, F.*, et al.* (2019).
iProX: an integrated proteome resource. Nucleic Acids Res *47*, D1211-D1217.

Mallart, C., Netter, S., Chalvet, F., Claret, S., Guichet, A., Montagne, J., Pret, A.M., and Malartre, M.
(2024). JAK-STAT-dependent contact between follicle cells and the oocyte controls Drosophila
anterior-posterior polarity and germline development. Nat Commun *15*, 1627.

Matakatsu, H., and Blair, S.S. (2004). Interactions between Fat and Dachsous and the regulation of
planar cell polarity in the
wing. Development *131*, 3785-3794.

Matis, M., and Axelrod, J.D. (2013). Regulation of PCP by the Fat signaling pathway. Genes Dev *27*,
2207-2220.

Michel, M., Raabe, I., Kupinski, A.P., Perez-Palencia, R., and Bokel, C. (2011). Local BMP receptor
activation at adherens junctions in the Drosophila germline stem cell niche. Nat Commun *2*, 415.

Munoz-Soriano, V., Belacortu, Y., and Paricio, N. (2012). Planar cell polarity signaling in collective
cell movements during morphogenesis and disease. Curr Genomics *13*, 609-622.

O'Shea, J.J., Gadina, M., and Schreiber, R.D. (2002). Cytokine signaling in 2002: new surprises in the
Jak/Stat pathway. Cell *109 Suppl*, S121-131.

Olivares-Castineira, I., and Llimargas, M. (2017). EGFR controls Drosophila tracheal tube elongation
by intracellular trafficking regulation. PLoS Genet *13*, e1006882.

Palmroth, M., Kuuliala, K., Peltomaa, R., Virtanen, A., Kuuliala, A., Kurttila, A., Kinnunen, A.,
Leirisalo-Repo, M., Silvennoinen, O., and Isomaki, P. (2021). Tofacitinib Suppresses Several JAK-

STAT Pathways in Rheumatoid Arthritis In Vivo and Baseline Signaling Profile Associates With

Treatment Response. Front Immunol *12*, 738481.

Pastor-Pareja, J.C., and Xu, T. (2011). Shaping cells and organs in Drosophila by opposing roles of fat

body-secreted Collagen IV and perlecan. Dev Cell *21*, 245-256.

Peng, Y., and Axelrod, J.D. (2012). Asymmetric protein localization in planar cell polarity:

mechanisms, puzzles, and challenges. Curr Top Dev Biol *101*, 33-53.

Perochon, J., Yu, Y., Aughey, G.N., Medina, A.B., Southall, T.D., and Cordero, J.B. (2021). Dynamic

adult tracheal plasticity drives stem cell adaptation to changes in intestinal homeostasis in Drosophila.

Nat Cell Biol *23*, 485-496.

Pitsouli, C., and Perrimon, N. (2010). Embryonic multipotent progenitors remodel the Drosophila

airways during metamorphosis. Development *137*, 3615-3624.

Rajan, A., Housden, B.E., Wirtz-Peitz, F., Holderbaum, L., and Perrimon, N. (2017). A Mechanism

Coupling Systemic Energy Sensing to Adipokine Secretion. Dev Cell *43*, 83-98 e86.

Rajan, A., and Perrimon, N. (2012). Drosophila cytokine unpaired 2 regulates physiological

homeostasis by remotely controlling insulin secretion. Cell *151*, 123-137.

Reaume, A.G., Clark, S.H., and Chovnick, A. (1989). Xanthine dehydrogenase is transported to the

Drosophila eye. Genetics *123*, 503-509.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image

analysis. Nat Methods *9*, 671-675.

Schottenfeld-Roames, J., and Ghabrial, A.S. (2012). Whacked and Rab35 polarize dynein-motor-

complex-dependent seamless tube growth. Nat Cell Biol *14*, 386-393.

Schottenfeld-Roames, J., Rosa, J.B., and Ghabrial, A.S. (2014). Seamless tube shape is constrained by

endocytosis-dependent regulation of active Moesin. Curr Biol *24*, 1756-1764.

Schottenfeld, J., Song, Y., and Ghabrial, A.S. (2010). Tube continued: morphogenesis of the Drosophila

tracheal system. Curr Opin Cell Biol *22*, 633-639.

Silver, D.L., Geisbrecht, E.R., and Montell, D.J. (2005). Requirement for JAK/STAT signaling

throughout border cell migration in Drosophila. Development *132*, 3483-3492.

Sotillos, S., Diaz-Meco, M.T., Moscat, J., and Castelli-Gair Hombria, J. (2008). Polarized subcellular

localization of Jak/STAT components is required for efficient signaling. Curr Biol *18*, 624-629.

Sriskanthadevan-Pirahas, S., Turingan, M.J., Chahal, J.S., Thorson, E., Khan, S., Tinwala, A.Q., and

Grewal, S.S. (2022). Adipose mitochondrial metabolism controls body growth by modulating systemic

cytokine and insulin signaling. Cell Rep *39*, 110802.

Stenmark, H. (2009). Rab GTPases as coordinators of vesicle traffic. Nat Rev Mol Cell Biol *10*, 513-

525.

Strutt, H., and Strutt, D. (2021). How do the Fat-Dachsous and core planar polarity pathways act

together and independently to coordinate polarized cell behaviours? Open Biol *11*, 200356.

Tamamouna, V., Rahman, M.M., Petersson, M., Charalambous, I., Kux, K., Mainor, H., Bolender, V.,

Isbilir, B., Edgar, B.A., and Pitsouli, C. (2021). Remodelling of oxygen-transporting tracheoles drives

intestinal regeneration and tumorigenesis in Drosophila. Nat Cell Biol *23*, 497-510.

Taylor, J., Abramova, N., Charlton, J., and Adler, P.N. (1998). Van Gogh: a new Drosophila tissue

polarity gene. Genetics *150*, 199-210.

Thomas, C., and Strutt, D. (2012). The roles of the cadherins Fat and Dachsous in planar polarity

specification in Drosophila. Dev Dyn *241*, 27-39.

Tree, D.R., Shulman, J.M., Rousset, R., Scott, M.P., Gubb, D., and Axelrod, J.D. (2002). Prickle

mediates feedback amplification to generate asymmetric planar cell polarity signaling. Cell *109*, 371-381.

Tsurumi, A., Xia, F., Li, J., Larson, K., LaFrance, R., and Li, W.X. (2011). STAT is an essential activator of the zygotic genome in the early Drosophila embryo. PLoS Genet *7*, e1002086.

Verboogen, D.R.J., Ter Beest, M., Honigmann, A., and van den Bogaart, G. (2018). Secretory vesicles of immune cells contain only a limited number of interleukin 6 molecules. FEBS Lett *592*, 1535-1544.

Vinson, C.R., and Adler, P.N. (1987). Directional non-cell autonomy and the transmission of polarity information by the frizzled gene of Drosophila. Nature *329*, 549-551.

Vitorino, P., and Meyer, T. (2008). Modular control of endothelial sheet migration. Genes Dev *22*, 3268-3281.

Vonderheit, A., and Helenius, A. (2005). Rab7 associates with early endosomes to mediate sorting and transport of Semliki forest virus to late endosomes. PLoS Biol *3*, e233.

Wang, Y., Yang, Q., Meng, X., Wijaya, C.S., Ren, X., and Xu, S. (2022). Recruitment of tetraspanin TSP-15 to epidermal wounds promotes plasma membrane repair in C. elegans. Dev Cell *57*, 1630-1642 e1634.

Williams, A.M., Donoughe, S., Munro, E., and Horne-Badovinac, S. (2022). Fat2 polarizes the WAVE complex in trans to align cell protrusions for collective migration. Elife *11*.

Wong, L.L., and Adler, P.N. (1993). Tissue polarity genes of Drosophila regulate the subcellular location for prehair initiation in pupal wing cells. J Cell Biol *123*, 209-221.

Woodcock, K.J., Kierdorf, K., Pouchelon, C.A., Vivancos, V., Dionne, M.S., and Geissmann, F. (2015). Macrophage-derived upd3 cytokine causes impaired glucose homeostasis and reduced lifespan in Drosophila fed a lipid-rich diet. Immunity *42*, 133-144.

Wulff, J., and Black, M.J. (2015). Efficient sparse-to-dense optical flow estimation using a learned basis and layers. Paper presented at: 2015 IEEE Conference on Computer Vision and Pattern Recognition (CVPR).

Yang, C.H., Axelrod, J.D., and Simon, M.A. (2002). Regulation of Frizzled by fat-like cadherins during planar polarity signaling in the Drosophila compound eye. Cell *108*, 675-688.

Yoshihara, M., Adolfsen, B., Galle, K.T., and Littleton, J.T. (2005). Retrograde signaling by Syt 4 induces presynaptic release and synapse-specific growth. Science *310*, 858-863.

Zallen, J.A. (2007). Planar polarity and tissue morphogenesis. Cell *129*, 1051-1063.

Zeidler, M.P., Bach, E.A., and Perrimon, N. (2000). The roles of the Drosophila JAK/STAT pathway. Oncogene *19*, 2598-2606.

Zeidler, M.P., Perrimon, N., and Strutt, D.I. (1999). Polarity determination in the Drosophila eye: a novel role for unpaired and JAK/STAT signaling. Genes Dev *13*, 1342-1353.

Zhao, Q., Wirka, R., Nguyen, T., Nagao, M., Cheng, P., Miller, C.L., Kim, J.B., Pjanic, M., and Quertermous, T. (2019). TCF21 and AP-1 interact through epigenetic modifications to regulate coronary artery disease gene expression. Genome Med *11*, 23.

Zhao, X., and Karpac, J. (2017). Muscle Directs Diurnal Energy Homeostasis through a Myokine-Dependent Hormone Module in Drosophila. Curr Biol *27*, 1941-1955 e1946.

Zheng, H., Yang, X., and Xi, Y. (2016). Fat body remodeling and homeostasis control in Drosophila. Life Sci *167*, 22-31.

Zhou, B., Feng, C., Sun, S., Chen, X., Zhuansun, D., Wang, D., Yu, X., Meng, X., Xiao, J., Wu, L.*, et al.* (2024). Identification of signaling pathways that specify a subset of migrating enteric neural crest cells at the wavefront in mouse embryos. Dev Cell.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium (https://www.iprox.cn/page/PSV023.html;?url=1708323921277ZyfC) via the iProX partner repository (Ma et al., 2019) with accession number gfnR. The RNA-Seq data generated and analyzed in this study have been deposited in the NCBI database under accession number GSE256177 with token code: clyfasuctboflat. The SMART-Seq data (L3, 0hr APF and 2hr APF) analyzed in this study have been deposited in the NCBI database under accession number GSE184856 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE184856]. The ChIP-Seq data generated and analyzed in this study have been deposited in the NCBI database under accession code 256176 with token: qvupguyijvyndyf.

     Figure legends

     Figure 1 Fat body affects disciplined migration of tracheal progenitors. (A)

     Schematic cartoon showing the migration of tracheal progenitors (red) and

     degenerative tracheal branches (dashed grey lines) in pupae. Fat body is shown in

     beige. Arrows denote anterior-posterior (A-P) axis. (B, C) Frontal section (B) and

     sagittal view (C) showing the relative position of fat body and tracheal progenitors.

     (D-J') Migration of tracheal progenitors in control and fat body perturbation flies. (D-

     D") Migration of tracheal progenitors (red) upward from transverse connective (blue

     dashed lines) and along the dorsal trunk (white dotted lines) at 0hr APF (D), 1hr APF

     (D'), and 3hr APF (D"). (E-E") Bidirectional movement of tracheal progenitors in fat

     body-depleted (*lsp2>rpr.hid*) flies. 0hr APF (E), 1hr APF (E'), 3hr APF (E"). Arrows

     point to anterior movement of tracheal progenitors. (F) Bar graph showing the

     migration distance of tracheal progenitors. The top chart of column represents the

     migration distance of anterior-most stem cells, and the lower chart of column

     represents the migration distance of posterior-most stem cells. Error bars represent

     SEM, n=6. (G-G') The distribution of progenitors at 2hr APF. (H-H") The distribution

     of progenitors in fat body-depleted flies at 2hr APF. (I-J') Computer simulation

     depicting trajectories of progenitor migration. (I, J) Confocal images of tracheal

     progenitors. (I', J') Vectors of progenitor migration. (K) Bar graph plots the binary

     entropy that represents the disorderedness of migration direction of tracheal

     progenitors. Error bars represent SEM, n=10. (L) The Bernoulli random variable X

     showing optic flow distribution of the binarized directions in each group. Error bars

     represent SEM, n=10. N.S. indicates not significant. Scale bar: 100 μm (B, C, G), 200

     μm (D-E"). Genotypes: (B, C) *UAS-mCD8-GFP/+; lsp2-Gal4,P[B123]-RFP-moe/+*;

    (D-D", G, G') *Gal80^{ts}/+;lsp2-Gal4,P[B123]-RFP-moe/+* ; (E-E", H, H') *UAS-rpr-*

    hid/+;Gal80^{ts}/+;lsp2-Gal4,P[B123]-RFP-moe/+.

    Figure 2 Dependence of tracheal progenitors on cytokines from fat body. (A) Top

    functional clusters among the differentially expressed genes of progenitors between

control and fat body-depleted pupae. Gene ratio refers to the proportion of genes in a

dataset that are associated with a particular biological process, function, or pathway.

Count indicates the number of genes from an input gene list that are associated with a

specific GO term. (B) Heatmap depicting expression levels of principal target genes of

signaling pathways in L3 larvae, 0hr APF pupae and 2hr APF pupae. (C) Heatmap

showing the differential expression of target genes of signaling pathways between

control and fat body-depleted pupae. (D-D'') Migration of tracheal progenitors along

the dorsal trunk at 0hr APF (D), 1hr APF (D'), and 3hr APF (D''). The white dashed

line shows transverse connective. (E-E''). Migration of tracheal progenitors in

upd2RNAi flies. (F) Bar graph plots the migration distance of tracheal progenitors.

Error bars represent SEM, n=6. (G) Volcano plot showing surface proteomics of

tracheal epithelium (up-regulated genes with ten-fold or higher changes in red; down-

regulated genes with ten-fold or higher changes in blue). (H) Top functional classes

among the surface proteomics of trachea. (I) Schematic diagram depicting the

working principle of the DIPF reporter. (J) The signal of DIPF reporter in tracheal

progenitors. The progenitors are outlined by dashed lines. N.S. indicates not

significant. Scale bar: 200 μm (D-E''), 50 μm (J). Genotypes: (A, C) *lsp2-*

Gal4,P[B123]-RFP-moe/+ for control*, UAS-rpr-hid/+;Gal80^{ts}/+;lsp2-Gal4,P[B123]-*

RFP-moe/+; (B) *P[B123]-RFP-moe/+*. (D-D'') *lsp2-Gal4,P[B123]-RFP-moe/+*; (E-

E'') *lsp2-Gal4,P[B123]-RFP-moe/UAS-upd2RNAi*; (J) *btl-Gal4/UAS-DIPF*.

1025

Figure 3 JAK/STAT pathway is required for the discipline of tracheal progenitor

migration. (A-D'') Migration of tracheal progenitors along the dorsal trunk at 0hr

APF, 1hr APF and 3hr APF. The white dashed line shows transverse connective. The

progenitors of control (A-A''), *domeRNAi* (B-B''), *hopRNAi* (C-C''), and

stat92ERNAi (D-D'') flies. (E) Bar graph showing migration distance of progenitors.

Error bars represent SEM, n=6. (F-G'') JAK inhibition causes bidirectional movement

of progenitors. Migration of tracheal progenitors in the absence (DMSO-fed) (F-F'')

or in the presence of Tofacinib (JAK inhibitor) (G-G''). (H) Bar graph showing the

distance of anterior movement. Error bars represent SEM, n=6. Scale bar: 200 μm (A-

D", F-G"). Genotypes: (A-A", F-G") *btl-Gal4/+;P[B123]-RFP-moe/+*; (B-B") *btl-*

Gal4/+;P[B123]-RFP-moe/UAS-domeRNAi; (C-C") *btl-Gal4/+;P[B123]-RFP-*

moe/UAS-hopRNAi; (D-D") *btl-Gal4/+;P[B123]-RFP-moe/UAS-stat92ERNAi*.

1038

Figure 4 Identification of gene targets of Stat92E in *Drosophila* tracheal

progenitors. (A) Bubble plot represents the top functional clusters among gene

targets. The establishment of planar polarity denoted in red solid box is identified with

high enrichment score. (B) Top functional classes among the differentially expressed

genes in larval-pupal transition. (C) ChIP-seq peaks at loci regulated by Stat92E.

Scale bar: 20kb (*ds*, *fz*, *stan*), 5kb (*fat2*, *vg*), 1kb (*fj*). (D-H) Validation of gene targets

of Stat92E ChIP-seq. (D-G) The expression of Ds-GFP in the tracheal progenitors of

control (D), *domeRNAi* (E), *hopRNAi* (F) and *stat92ERNAi* (G). The progenitors are

outlined by dashed lines. (H) The bar graphs plot the relative level of Ds. Error bars

represent SEM, n=6. (I-M) The expression of Fj in tracheal progenitors. The

expression of Fj-GFP in the tracheal progenitors of control (I), *domeRNAi* (J),

hopRNAi (K) and *stat92ERNAi* (L). Dashed lines outline tracheal progenitors. (M)

The bar graphs plot the relative level of Fj. Error bars represent SEM, n=6. (N-R) The

level of Ft-GFP in the tracheal progenitors of control (N), *domeRNAi* (O), *hopRNAi*

(P) and *stat92ERNAi* (Q). (R) The bar graphs plot the relative level of Ft. Error bars

represent SEM, n=4. Scale bar: 50 μm (D-G, I-L, N-Q). Genotypes: (D) *btl-Gal4,Ds-*

GFP/+; (E) *btl-Gal4,Ds-GFP/+;UAS-domeRNAi/+*; (F) *btl-Gal4,Ds-GFP/+;UAS-*

hopRNAi/+; (G) *btl-Gal4,Ds-GFP/+;UAS-stat92ERNAi/+*; (I) *btl-Gal4,Fj-GFP/+*; (J)

btl-Gal4,Fj-GFP/+;UAS-domeRNAi/+; (K) *btl-Gal4,Fj-GFP/+;UAS-hopRNAi/+*; (L)

btl-Gal4,Fj-GFP/+;UAS-stat92ERNAi/+; (N) *btl-Gal4/+;Ft-GFP/+*; (O) *btl-*

Gal4/+;Ft-GFP/UAS-domeRNAi; (P) *btl-Gal4/+;Ft-GFP/UAS-hopRNAi*; (Q) *btl-*

Gal4/+;Ft-GFP/UAS-stat92ERNAi.

1061

1062   **Figure 5 Disciplined migration requires planar cell polarity system.** (A-E)

1063   Migration of tracheal progenitors. The migration of progenitors in control (A-A"),

1064   *dsRNAi* (B-B"), *ftRNAi* (C-C") and *fjRNAi* (D-D") flies. (E) Bar graph plots the

1065	migration distance of anterior movement. Error bars represent SEM, n=4. (F-I) Level
1066	of Ft in tracheal progenitors of control (F, G) and *stat92ERNAi* (H-J) flies. The
1067	images show progenitors at 1hr APF (H, H') and 2hr APF (I, I'). Ft-GFP (green) (F,
1068	H, I), phalloidin (magenta), Hoechst (blue), and merged images (F', H', I'). (G, J)
1069	Profile plots showing the level of Ft-GFP in control (G) and *stat92ERNAi* (J) flies,
1070	n=5. ANOVA test: $p < 0.0001$. The levels of Ft were measured along the dotted lines
1071	in F' or I'. Anterior (A) and posterior (P). (K) Representative traces plot the migration
1072	distance relative to the origin, n=12. The x-axis represents the number of captured
1073	images. Individual frame is captured every 5 minutes. (K') Rose plot depicting the
1074	direction of cell movement. (L) Representative traces showing the movement of
1075	individual *fj*-KO cells relative to their origin, n=11. The x-axis represents the number
1076	of captured images. Individual frame is captured every 5 minutes. (L') Rose plot
1077	depicting the movement direction of *fj*-KO cells. (M) Scatter plots represent the ratio
1078	(*d*/*D*) of straight-line length displacement (*d*) relative to the length of the migration
1079	track (*D*) of individual cell. Error bars represent SEM. N.S. indicates not significant.
1080	Scale bar: 200 μm (A-D''), 50 μm (F, F'), 100 μm (H-I'). Genotypes: (A-A'') *btl-*
1081	*Gal4/+;P[B123]-RFP-moe/+*; (B-B'') *btl-Gal4/UAS-dsRNAi;P[B123]-RFP-moe/+*;
1082	(C-C'') *btl-Gal4/+;P[B123]-RFP-moe/UAS-ftRNAi*; (D-D'') *btl-Gal4/UAS-*
1083	*fjRNAi;P[B123]-RFP-moe/+*; (F, F') *btl-Gal4/+;Ft-GFP/+*; (H, H', I, I') *btl-*
1084	*Gal4/+;Ft-GFP/UAS-stat92ERNAi.*
1085	
1086	**Figure 6 The production and transport of Upd2 from fat body.** (A, B) The
1087	number of Upd2-mCherry-containing vesicles in fat body of control (DMSO-fed, A)
1088	and BFA-treated L3 larvae (B). Larger view in lower magnification is provided in
1089	Figure 6-figure supplement 1. (C) Bar graph plots the number of Upd2-mCherry-
1090	containing vesicles in fat body. Error bars represent SEM, n=6. (D, E) The confocal
1091	image showing the number of Upd2-containing vesicles in progenitors of DMSO-fed
1092	control (D) and BFA-treated flies (E). (F) Bar graph plots the number of Upd2-
1093	mCherry-containing vesicles in progenitors. Error bars represent SEM, n=4. (G) The
1094	number of Upd2-mCherry containing vesicles (red) in fat body. (H, I) Upd2

accumulation in fat body was increased in the presence of *grasp65RNAi* (H) and

lbmRNAi (I). (J) Bar graph plots the number of Upd2-mCherry-containing vesicles.

Error bars represent SEM, n=4. (K-M) The Upd2 vesicles (red) in tracheal progenitors

(DAPI) in control (K), *grasp65RNAi* (L) and *lbmRNAi* (M) flies. Dashed lines outline

tracheal progenitors. (N) Bar graph plots the number of Upd2-mCherry-containing

vesicles in progenitors. Error bars represent SEM, n=4. Scale bar: 20 μm (A, B, G-I),

50 μm (D, E, K-M). Genotypes: (A-G, K) *UAS-upd2-mCherry/+;lsp2-Gal4/+*; (H, L)

UAS-upd2-mCherry/UAS-grasp65RNAi;lsp2-Gal4/+; (I, M) *UAS-upd2-*

mCherry/+;lsp2-Gal4/UAS-lbmRNAi.

Figure 7 The dependence of tracheal progenitors on vesicle trafficking. (A-E)

The expression of Stat92E-GFP in tracheal progenitors of control (A), *grasp65RNAi*

(B), *lbmRNAi* (C), *rab5RNAi* (D) and *rab7RNAi* (E) flies. The progenitors are outlined

by dashed lines. (F) Bar graph plots the relative expression of Stat92E-GFP. Error bars

represent SEM, n=7. (G, H) The expression of Stat92E-GFP in tracheal progenitors in

DMSO-fed (G) and BFA-treated (H) flies. (I) Bar graph plots the relative expression

of Stat92E-GFP. Error bars represent SEM, n=5. (J) The signal of DIPF reporter in

tracheal progenitors. (K) The effects of Tofacinib (JAK inhibitor) on DIPF reporter in

progenitors. (L) The effects of Brefeldin A on DIPF reporter in progenitors. Dashed

lines outline tracheal progenitors. (M) Bar graphs showing the signal of DIPF

reporter. Error bars represent SEM, n=4. (N-O'') Migration of tracheal progenitors in

DMSO-fed flies (N-N'') and BFA-treated flies (O-O''). (P) Bar graph plots migration

distance of anterior movement. Error bars represent SEM, n≥3. (Q-U'') Migration of

tracheal progenitors at 0hr APF (Q), 1hr APF (Q'), 3hr APF (Q''). The confocal

images showing the tracheal progenitors in control (Q-Q''), *grasp65RNAi* (R-R''),

lbmRNAi (S-S''), *rab5RNAi* (T-T'') and *rab7RNAi* (U-U'') flies. (V) Bar graph plots

the migration distance of anterior movement. Error bars represent SEM, n=4. Scale

bar: 50 μm (A-E, G, H, J-L), 200 μm (N-O'', Q-U''). Genotypes: (A, G, H) *lsp2-*

Gal4,Stat92E-GFP/+; (B) *UAS-grasp65RNAi/+;lsp2-Gal4,Stat92E-GFP/+*; (C) *lsp2-*

Gal4,Stat92E-GFP/UAS-lbmRNAi; (D) *UAS-rab5RNAi/+;lsp2-Gal4,Stat92E-GFP/+*;

1125   (E) *UAS-rab7RNAi/+;lsp2-Gal4,Stat92E-GFP/+*; (J-L) *btl-Gal4/UAS-DIPF*; (N-Q")

1126   *lsp2-Gal4,P[B123]-RFP-moe/+*; (R-R") *lsp2-Gal4,P[B123]-RFP-moe/UAS-*

1127   *grasp65RNAi*; (S-S") *lsp2-Gal4,P[B123]-RFP-moe/+;UAS-lbmRNAi/+*; (T-T")

1128   *UAS-rab5RNAi/+;lsp2-Gal4,P[B123]-RFP-moe/+*; (U-U") *UAS-rab7RNAi/+;lsp2-*

1129   *Gal4,P[B123]-RFP-moe/+*.

1130

Figure 8 The roles of endocytic trafficking system in the transport of Upd2. (A-

D''') The confocal images showing the colocalization between Upd2 (red) (A, B, C,

D) and Rab5 (GFP) (A'), Rab7 (GFP) (B'), Grasp65 (GFP) (C') or Lbm (GFP). (A",

B", C", D") Merged images. (D''') 3D high-magnification view of the boxed inset in

D". (D'''') The Pearson's correlation coefficient depicting colocalization between

Lbm and Upd2 in fat body cells. (E-J) The PLA (Proximity Ligation Assay) assay

showing the interaction between Upd2 and Rab5 (E, F), Rab7 (G, H) or Lbm (I, J).

(K-M) Co-immunoprecipitation assay showing physical interaction between Upd2

and Rab5 (K), Rab7 (L) or Lbm (M) in larval fat body. (N-T) The expression of Lbm-

pHluorin in larval fat body and progenitors of control (N, O), *rab5RNAi* (P, Q) and

rab7RNAi (R, S) flies. Dashed lines outline tracheal progenitors. Arrowheads point to

Lbm-pHluorin puncta. DAPI signal indicates nuclei. (T) Schematic diagram depicting

Upd2-operated disciplined migration of tracheal progenitors. Scale bar: 5 μm (A-D",

E-J), 10 μm (N, P, R), 20 μm (O, Q, S). Genotypes: (A-A", E, F) *UAS-upd2-*

mCherry/+;lsp2-Gal4/UAS-GFP-rab5; (B-B", G, H) *UAS-upd2-mCherry/+;lsp2-*

Gal4/UAS-GFP-rab7; (C-C") *UAS-upd2-mCherry/UAS-grasp65-GFP;lsp2-Gal4/+*;

(D-D'''', I-L) *UAS-upd2-mCherry/UAS-lbm-GFP;lsp2-Gal4/+*; (N, O) *UAS-lbm-*

pHluorin/+;lsp2-Gal4/+; (P, Q) *UAS-lbm-pHluorin/UAS-rab5RNAi;lsp2-Gal4/+*; (R,

S) *UAS-lbm-pHluorin/UAS-rab7RNAi;lsp2-Gal4/+*.

1150

1151   **Figure 1-figure supplement 1 Genetic perturbation of fat body by expression of**

1152   **pro-apoptotic genes, *rpr* and *hid*.** Confocal images of larval fat body in control (A)

1153   and *lsp2>rpr.hid* (B) animals. Microscopic pictures of control (C) and *lsp2>rpr.hid*

1154   larvae (D). (E) Scatter plot showing migration velocity of tracheal progenitors in

control and *lsp2>rpr.hid* animals. Error bars represent SEM, n=6. (F, G) The

incorporation of EdU in the tracheal progenitors of control (F) and *lsp2>rpr.hid*

animals (G). (H) Bar graph depicting the number of EdU incorporation. Error bars

represent SEM, n=11. Scale bar: 100 μm (A, B), 20 μm (C, D), 50 μm (F, G).

Genotypes: (A, C, F) *lsp2-Gal4,P[B123]-RFP-moe/+*; (B, D) *UAS-rpr.hid/+;lsp2-*

Gal4,P[B123]-RFP-moe/+; (G) *UAS-rpr.hid/+;Gal80^{ts}/+;lsp2-Gal4,P[B123]-RFP-*

moe/+.

Figure 2-figure supplement 1 Identification of genes that contribute to

progenitors migration. (A) Differentially expressed genes in tracheal progenitors of

lsp2>rpr.hid pupa compared with control. (B) Volcano plot of RNA-seq showing

differentially regulated genes with 2-fold or higher changes (up-regulated genes in

red; down-regulated genes in blue) in *lsp2>rpr.hid* compared with control. (C-I'') The

migration of progenitors at 0hr APF, 1hr APF, or 3hr APF in control (C-C'') and

upd1RNAi (D-D''), *upd3RNAi* (E-E''), *pvf1RNAi* (F-F''), *pvf2RNAi* (G-G''), *pvf3RNAi*

(H-H''), and *hhRNAi* (I-I''). (J) Bar graph represents the migration distance of anterior

movement. Error bars represent SEM, n=4. (K) The migration of progenitors in

control (K-K'') and *btl>upd2RNAi* (L-L''). (M) Bar graph plots the migration distance

of anterior movement. Error bars represent SEM, n=4. N.S. indicates not significant.

Scale bar: 200 μm (C-I'', K-L''). Genotypes: (C-C'') *lsp2-Gal4,P[B123]-RFP-moe/+*;

(D-D'') *lsp2-Gal4,P[B123]-RFP-moe/UAS-upd1RNAi*; (E-E'') *lsp2-Gal4,P[B123]-*

RFP-moe/UAS-upd3RNAi; (F-F'') *lsp2-Gal4,P[B123]-RFP-moe/UAS-pvf1RNAi*; (G-*

G'') lsp2-Gal4,P[B123]-RFP-moe/UAS-pvf2RNAi; (H-H'') *lsp2-Gal4,P[B123]-RFP-*

moe/UAS-pvf3RNAi; (I-I'') *lsp2-Gal4,P[B123]-RFP-moe/UAS-hhRNAi*; (K-K'') *btl-*

Gal4/+;P[B123]-RFP-moe/+; (L-L'') *btl-Gal4/+;P[B123]-RFP-moe/UAS-upd2RNAi*.

Figure 2-figure supplement 2 Surface proteomics of *Drosophila* trachea. (A-C)

Representative confocal images showing the trachea of *btl>CD2-HRP* flies after

oxidation reaction in the presence of both BXXP and H_2O_2 (A), BXXP only (B), or

H_2O_2 only (C). (D) The western blot showing the biotinylated proteins labeled by an

HRP-catalyzed reaction with both BXXP and H_2O_2. (E) The confocal image showing

the trachea of *dome>GFP* flies. The expression domain of *dome* is visualized by

dome-Gal4-controlled GFP expression. The progenitors labeled by a progenitor-

specific enhancer, P[B123]-RFP-moe are outlined by dashed lines. The overlapping

expression of GFP and P[B123]-RFP-moe. (F, G) The expression of DIPF in fat body

(F) and salivary gland (G). Scale bar: 50 μm (E-H). Genotypes: (A-C) *btl-Gal4/UAS-*

CD2-HRP; (E) *dome-Gal4/+;UAS-GFP/P[B123]-RFP-moe*; (F,G) *tub-Gal4/UAS-*

DIPF.

1193

Figure 3-figure supplement 1 The roles of JAK/STAT pathway in tracheal

development. (A-E) Migration of tracheal progenitors along the dorsal trunk at 0hr

APF, 1hr APF and 3hr APF. The white dashed line shows transverse connective. The

progenitors of control (A-A''), *dome* mutants (B-B''), *hop* mutants (C-C''). (D) Bar

graphs represent migration distance of anterior movement. Error bars represent SEM,

n≥3. N.S. indicates not significant. (E-I) The pupal trachea in control (E), *upd2 RNAi*

(F), *dome^{G0264}* (G), *hop^{tum}* (H), and *stat92E^F* (I). Scale bar: 200 μm (A-E), 20mm (K-

L''). Genotypes: (A, E) *lsp2-Gal4,P[B123]-RFP-moe/+*(control); (B) *dome^{G0264}; lsp2-*

Gal4,P[B123]-RFP-moe/+; (C) *hop^{tum}; lsp2-Gal4,P[B123]-RFP-moe/+*; (F) *lsp2-*

Gal4,P[B123]-RFP-moe/upd2RNAi. (G) *dome^{G0264}*. (H) *hop^{tum}*. (I) *stat92E^F*.

1204

Figure 3-figure supplement 2 The dependence of tracheal progenitors on

JAK/STAT pathway. (A) Statistics of the number of progenitor cells. Error bars

represent SEM, n≥9. N.S. indicates not significant. (B-E) The expression of Stat92E-

GFP in tracheal progenitors of control (B), *domeRNAi* (C), *hopRNAi* (D) and

stat92ERNAi (E) flies. The progenitors are outlined by dashed lines. (F, G) The

expression of Stat92E-GFP in tracheal progenitors of control (F) or upd2RNAi flies

(G). (H, I) The expression of Stat92E-GFP in tracheal progenitors of DMSO-fed

(control, H) or Tofacitinib-treated (I) flies. Dashed lines outline tracheal progenitors.

(J) Bar graph plots relative intensity of Stat92E-GFP reporter. Error bars represent

SEM, n=6. N.S. indicates not significant. Scale bar: 50 μm (B-I). Genotypes: (A) *btl-*

Gal4/+;P[B123]-RFP-moe/+(control); *btl-Gal4/+;P[B123]-RFP-moe/UAS-*

domeRNAi; *btl-Gal4/+;P[B123]-RFP-moe/UAS-hopRNAi*; *btl-Gal4/+;P[B123]-RFP-*

moe/UAS-stat92ERNAi; (B) *btl-Gal4,Stat92E-GFP/+*; (C) *btl-Gal4,Stat92E-*

GFP/+;UAS-domeRNAi/+; (D) *btl-Gal4,Stat92E-GFP/+;UAS-hopRNAi/+*; (E) *btl-*

Gal4,Stat92E-GFP/+;UAS-Stat92ERNAi/+; (F, H, I) *Stat92E-GFP/+; lsp2-Gal4/+*;

(G) *Stat92E-GFP/+;lsp2-Gal4/UAS-upd2RNAi*.

1221

Figure 4-figure supplement 1 Annotation and analysis of genomic occupancy of

Stat92E. (A-C) Pie chart (A, B) and histogram (C) depicting the distribution of ChIP-

seq peaks relative to the nearby annotated genes. (D) Peaks of Stat92E association in

the promotor regions of *yurt* and *crb*. Scale bar: 2kb for *yurt*, 5kb for *crb*. (E) Relative

enrichment of JAK/STAT target genes analyzed by ChIP-qPCR. (F-I) The expression

of Fat2-GFP in the tracheal progenitors of control (F), *domeRNAi* (G), *hopRNAi* (H)

and *stat92ERNAi* (I). The progenitors are outlined by dashed lines. (J) The bar graphs

plot the relative level of Fat2. Error bars represent SEM, n=6. (K-N) The expression

of Yurt in tracheal progenitors. The expression of Yurt-GFP in the tracheal progenitors

of control (K), *domeRNAi* (L), *hopRNAi* (M) and *stat92ERNAi* (N). The progenitors

are outlined by dashed lines. (O) The bar graphs plot the relative level of Yurt. Error

bars represent SEM, n=6. (P-S) The expression of Crb-GFP in the tracheal progenitors

of control (P), *domeRNAi* (Q), *hopRNAi* (R) and *stat92ERNAi* (S). The progenitors are

outlined by dashed lines. (T) The bar graphs plot the relative level of Crb. Error bars

represent SEM, n=6. (U) The expression of JAK/STAT targets in the trachea of

stat92ERNAi flies. Quantitative RT-PCR was performed in triplets to analyze the

mRNA level. The expression levels in *stat92ERNAi* larvae were presented relative to

Gal4 control. Scale bar: 50 μm (F-I, K-N, P-S). Genotypes: (F) *btl-Gal4,Fat2-GFP/+*;

(G) *btl-Gal4,Fat2-GFP/+;UAS-domeRNAi/+*; (H) *btl-Gal4,Fat2-GFP/+;UAS-*

hopRNAi/+; (I) *btl-Gal4,Fat2-GFP/+;UAS-stat92ERNAi/+*; (J) *btl-Gal4/+;Yurt-*

GFP/+; (K) *btl-Gal4/+;Yurt-GFP/UAS-domeRNAi*; (L) *btl-Gal4/+;Yurt-GFP/UAS-*

hopRNAi; (M) *btl-Gal4/+;Yurt-GFP/UAS-stat92ERNAi*; (P) *btl-Gal4/+;Crb-GFP/+*;

(Q) *btl-Gal4/+;Crb-GFP/UAS-domeRNAi*; (R) *btl-Gal4/+;Crb-GFP/UAS-hopRNAi*;

(S) *btl-Gal4/+;Crb-GFP/UAS-stat92ERNAi*.

1246

1247   **Figure 4-figure supplement 2 Ds alters the level of Ft.** (A-C) The level of Ft-GFP

1248   in control (A), UAS-ds (B), and dsRNAi (C) flies. (D) Bar graph represents the signal

1249   of Ft-GFP. Error bars represent SEM, n=4. Scale bar: 50 µm (A-C). Genotypes: (A)

1250   *btl-Gal4/+;Ft-GFP/+*; (B) *btl-Gal4/+;Ft-GFP/UAS-ds*; (C) *btl-Gal4/+;Ft-GFP/UAS-*

1251   *dsRNAi*.

1252

Figure 5-figure supplement 1   Apico-basal polarity proteins contribute to the

migration polarity of progenitors. (A-H'') The migration of progenitors at 0hr APF

(A), 1hr APF (A'), 3hr APF (A'') in control (A-A''), *UAS-ds* (B-B''), *UAS-ft* (C-C''),

fat2RNAi (E-E''), *crbRNAi* (F-F''), *yurtRNAi* (G-G'') and *scbRNAi* (H-H''). (D,I,J)

Bar graph plots the migration distance of anterior movement. Error bars represent

SEM, n≥3. N.S. indicates not significant. Scale bar: 200 µm (A-C'', E-H'').

Genotypes: (A-A'') *btl-Gal4/+;P[B123]-RFP-moe/+*; (B-B'') *btl-*

Gal4/+;P[B123]/UAS-ds; (C-C'') *btl-Gal4/UAS-ft;P[B123]/+*; (E-E'') *btl-*

Gal4/+;P[B123]-RFP-moe/UAS-fat2RNAi; (F-F'') *btl-Gal4/+;P[B123]-RFP-*

moe/UAS-crbRNAi; (G-G'') *btl-Gal4/+;P[B123]-RFP-moe/UAS-yurtRNAi*; (H-H'')

btl-Gal4/+;P[B123]-RFP-moe/UAS-scbRNAi. (J) *btl-Gal4/+;P[B123]-RFP-*

moe/+(control); *btl-Gal4/+;P[B123]-RFP-moe/UAS-enaRNAi*; *btl-Gal4/ UAS-*

fakRNAi;P[B123]-RFP-moe/+; *btl-Gal4/+;P[B123]-RFP-moe/UAS-E-cadRNAi*; *btl-*

Gal4/+;P[B123]-RFP-moe/UAS-robo2RNAi.

1267

Figure 5-figure supplement 2 Protrusions extending from the leading edge of

progenitors. (A, B) The confocal image showing the filopodia at the leading edge of

migrating tracheal progenitors in control (A) and *upd2RNAi* flies (B). Arrows denote

anterior-posterior (A-P) axis. Dashed lines indicate dorsal trunk. (C) Bar graph

depicting the number of filopodia extending from leading edge. Error bars represent

SEM, n=8. (D, E) Expression of *bnl*-lacZ in the trachea of control (D) and

stat92ERNAi (E) flies. Scale bar: 100 µm (A, B), 500 µm (D, E). Genotypes: (A) *lsp2-*

Gal4,P[B123]-RFP-moe/+; (B) *lsp2-Gal4,P[B123]-RFP-moe/UAS-upd2RNAi*.

1276

Figure 5-figure supplement 3 The dependence of migration directionality on

Ft/Ds system. (A) Representative traces plot the migration distance relative to the

origin, n≥10. (B) Rose plot depicting the direction of cell movement. (C)

Representative traces showing the movement of individual *dsRNAi* cells relative to

their origin. (D) Rose plot depicting the movement direction of *dsRNAi* cells. (E)

Representative traces showing the movement of individual *ftRNAi* cells relative to

their origin. (F) Rose plot depicting the movement direction of *ftRNAi* cells. (G)

Scatter plots represent the ratio (*d/D*) of straight-line length displacement (*d*) relative

to the length of the migration track (*D*) of individual cell.

1286

1287 **Figure 6-figure supplement 1 Microscopic image of a preparation of fly fat body.**

1288 Scale bar: 50 um.

1289

Figure 7-figure supplement 1 The roles of Rabs in transport of Upd2. (A-C'')

Migration of tracheal progenitors. The migration of progenitors at 0hr APF, 1hr APF,

3hr APF in control (A-A''), *rab2RNAi* (B-B''), *rab3RNAi* (C-C''). (D) Bar graph plots

the migration distance of anterior movement. Error bars represent SEM, n≥3. (E-E'')

The confocal images showing the colocalization between Upd2 (red) (E) and Rab3

(GFP) (E'). (E'') Merged images. Scale bar: 200 μm (A-C''), 5 μm (E-E'').

Genotypes: (A-A'') *lsp2-Gal4,P[B123]-RFP-moe/+*; (B-B'') *lsp2-Gal4,P[B123]-*

RFP-moe/UAS-rab2RNAi; (C-C'') *lsp2-Gal4,P[B123]-RFP-moe/UAS-rab3RNAi*; (E-

E'') *UAS-upd2-mCherry/UAS-rab3-YFP;lsp2-Gal4/+*.

1299

1300 **Figure 8-figure supplement 1 Vesicular transport of Upd2.** (A-C'') Co-

1301 localization and interaction between Lbm and Upd2. Co-transfection of act-Gal4,

1302 UAS-lbm-GFP (A) and UAS-upd2-mCherry (A') in S2 cells. (A'') Merged image. (B-

1303 C'') Confocal images showing that Lbm co-localizes with Upd2 in vesicles along the

1304 protrusions of S2 cell. (D) Electron microscopic image of Lbm-HRP vesicles (dark) in

fat body. (E) A predicted interaction between Lbm (blue) and Upd2 (green). (F) Co-immunoprecipitation assay showing the interaction between Lbm and Upd2 in S2 cells. (G, H) The expression of Lbm-pHluorin in control fat body (G) and BFA-treated flies (H). (I, J) The expression of Lbm-pHluorin in control tracheal progenitors (I) and BFA-treated flies (J). The tracheal progenitors were outlined by dashed lines. Scale bar: 10 μm (A-A''), 20 μm (G, H). Genotypes: (G-J) *UAS-lbm-pHluorin/+;lsp2-Gal4/+*.

Video 1 3D-view of confocal image stack of tracheal progenitors and fat body. Scale bar: 100 μm. Genotype: *UAS-mCD8-GFP/+;lsp2-Gal4,P[B123]-RFP-moe/+*.

Video 2 The movement of tracheal progenitors in control and *rpr.hid* flies. Scale bar: 100 μm. Genotypes: *Gal80^{ts}/+;lsp2-Gal4,P[B123]-RFP-moe/+* (control) and *UAS-rpr-hid/+;Gal80^{ts}/+;lsp2-Gal4,P[B123]-RFP-moe/+*.

Video 3 The movement of tracheal progenitors in control and *upd2RNAi* flies. Scale bar: 100 μm. Genotypes: *lsp2-Gal4,P[B123]-RFP-moe/+* (control) and *lsp2-Gal4,P[B123]-RFP-moe/UAS-upd2RNAi*.

Video 4 The movement of tracheal progenitors in control and JAK/STAT pathway-perturbed flies. Scale bar: 100 μm. Genotypes: *btl-Gal4/+;P[B123]-RFP-moe/+* (control), *btl-Gal4/+;P[B123]-RFP-moe/UAS-domeRNAi*, *btl-Gal4/+;P[B123]-RFP-moe/UAS-hopRNAi* and *btl-Gal4/+;P[B123]-RFP-moe/UAS-stat92ERNAi*.

Video 5 The movement of tracheal progenitors in control and PCP component aberrant flies. Scale bar: 100 μm. Genotypes: *btl-Gal4/+;P[B123]-RFP-moe/+*(control), *btl-Gal4/UAS-dsRNAi;P[B123]-RFP-moe/+* and *btl-Gal4/+;P[B123]-RFP-moe/UAS-ftRNAi*.

1335 **Video 6 The movement of tracheal progenitors in control and *lbmRNAi* flies.**

1336 Scale bar: 100 μm. Genotypes: *lsp2-Gal4,P[B123]-RFP-moe/+* (control) and *lsp2-*

1337 *Gal4,P[B123]-RFP-moe/+;UAS-lbmRNAi/+.*

1338

1339

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A Upd2-mCherry **A'** GFP-Rab5 **A''** merge **E** PLA(Rab5+Upd2) **F** control

B **B'** GFP-Rab7 **B''** **G** PLA(Rab7+Upd2) **H** control

C **C'** Grasp65-GFP **C''** **I** PLA(Lbm+Upd2) **J** control

D **D'** Lbm-GFP **D''** **D'''** **D''''** Pearson's R value = 0.8

Lbm-GFP   Upd2-mCherry

K

	IP:α-GFP		Input	
Rab5-GFP	+	–	+	–
Upd2-mCherry	+	+	+	+	kDa
IB:α-mCherry					–72
IB:α-GFP					–57

L

	IP:α-GFP		Input	
Rab7-GFP	+	–	+	–
Upd2-mCherry	+	+	+	+	kDa
IB:α-mCherry					–72
IB:α-GFP					–50

M

	IP:α-GFP		Input	
Lbm-GFP	+	–	+	–
Upd2-mCherry	+	+	+	+	kDa
IB:α-mCherry					–72
IB:α-GFP					–48

N control — fat body cell / lbm-pHluorin

O tracheal progenitor / DAPI

P *rab5RNAi*

Q

R *rab7RNAi*

S

T

fat body   Upd2   tracheal progenitors   disciplined   undisciplined