Title :

kcnb1 loss-of-function in zebrafish causes neurodevelopmental and epileptic disorders
associated with GABA dysregulation

Lauralee Robichon[1,3], Claire Bar[1,2,3], Anca Marian[1,3], Lisa Lehmann[1,3], Solène Renault[1,3],

Edor Kabashi[1,3,*], Sorana Ciura[1,3], Rima Nabbout[1,2,3]

Affiliations:
9
[1] Laboratory of Translational Research for Neurological Disorders, INSERM UMR 1163,

Institut des maladies génétiques, Imagine Institute, Université Paris Cité, Paris, France

[2] APHP, Reference center for rare epilepsies, Department of Pediatric Neurology, Hôpital

Necker-Enfants Malades, member of EPICARE, Paris, France

[3] Université Paris Cité – Sorbonne Université, Paris, France

Authors e-mail addresses:

lauralee.robichon@institutimagine.org; claire.bar@chu-bordeaux.fr;

anca.marian@institutimagine.org; lisa.lehmann@etu.unistra.fr;

solene.renault@institutimagine.org; edor.kabashi@institutimagine.org;

sorana.ciura@institutimagine.org; rima.nabbout@aphp.fr

20
Corresponding author *:

Edor KABASHI

Laboratory of Translational Research for Neurological Disorders

Imagine Institute

24 Boulevard du Montparnasse

75015 Paris

France

01 42 75 46 48

Email : edor.kabashi@institutimagine.org

Declaration of interest: None of the authors has any conflict of interest to disclose

KEY POINTS

- kcnb1 is expressed in distinct cell subtypes and various regions of the central nervous system in zebrafish

- Brain anatomy and neuronal circuits are not disrupted in the *kcnb1* loss-of-function zebrafish model

- Loss of *kcnb1* leads to altered behavior phenotype, light and sound-induced locomotor impairments

- *kcnb1* knock-out zebrafish exhibit increased locomotor sensitivity to PTZ and elevated expression of epileptogenesis-related genes

- *kcnb1*$^{-/-}$ larvae show spontaneous and provoked epileptiform-like electrographic activity associated with disrupted GABA regulation

ABSTRACT

Objective: *KCNB1* encodes an α-subunit of the delayed-rectifier voltage-dependent potassium channel K$_v$2.1. *De novo* pathogenic variants of *KCNB1* have been linked to developmental and epileptic encephalopathies (DEE), diagnosed in early childhood and sharing limited treatment options. Loss-of-function (LOF) of *KCNB1* with dominant negative effects has been proposed as the pathogenic mechanism in these disorders. Here, we aim to characterize a knock-out (KO) zebrafish line targeting *kcnb1 (kcnb1*$^{+/-}$ *and kcnb1*$^{-/-}$*)* for investigating DEEs.

Methods: This study presents the phenotypic analysis of a *kcnb1* knock-out zebrafish model, obtained by CRISPR/Cas9 mutagenesis. Through a combination of immunohistochemistry, behavioral assays, electrophysiological recordings, and neurotransmitter quantifications, we have characterized the expression, function, and impact of this *kcnb1* LOF model at early stages of development.

Results: In wild-type larval zebrafish, kcnb1 was found in various regions of the central nervous system and in diverse cell subtypes including neurons, oligodendrocytes and microglial cells. Both *kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$ zebrafish displayed impaired swimming behavior and "epilepsy-like" features that persisted through embryonic and larval development, with variable severity. When exposed to the chemoconvulsant pentylenetetrazol (PTZ), both mutant models showed elevated locomotor activity. In addition, PTZ-exposed *kcnb1*$^{-/-}$ larvae exhibited higher bdnf mRNA expression and activated c-Fos positive neurons in the telencephalon. This same model presents spontaneous and provoked epileptiform-like electrographic activity associated with disrupted GABA regulation. In this KO model, neuronal circuit organization remained unaffected.

Significance: We conclude that *kcnb1* knock-out in zebrafish leads to early-onset phenotypic features reminiscent of DEEs, affecting neuronal functions and primarily inhibitory pathways in developing embryonic and larval brains. This study highlights the relevance of this model for investigating developmental neuronal signaling pathways in *KCNB1*-related DEEs.

Keywords: neurodevelopment, epilepsy, KCNB1, zebrafish, loss-of-function

1. INTRODUCTION

KCNB1 encodes the pore-forming α1 subunit of the voltage□gated potassium channel subfamily 2 (K$_v$2.1). K$_v$2.1 is extensively expressed across the central nervous system (CNS), and predominantly localized in clusters on neuronal soma, proximal dendrites, and axonal initial segment of neurons (1–4), generating a delayed-rectifier outward potassium current (1,2,5,6). Moreover, the channel modulates neuronal excitability through activity-dependent regulation of K$_v$2.1 phosphorylation in various neuronal subtypes (1,2,5,6). Beyond its role in

ionic conduction, $K_v2.1$ channels participate in intracellular protein trafficking and calcium

signaling (5,7,8).

De novo pathogenic variants of *KCNB1* have been reported in patients with

developmental and epileptic encephalopathy (DEE), as well as developmental encephalopathy

(DE) without epilepsy or with late-onset, severe and pharmacoresponsive epilepsy (9–12). All

patients reported in the largest series exhibit poor long-term outcome associated with a wide

phenotypic spectrum including severe intellectual disability, attention disorders and autism

spectrum disorder (ASD) (9,12,13).

In vitro studies of *KCNB1* variants have revealed various degrees of loss-of-function

(LOF) exhibiting dominant negative effects such as reduced potassium conductance, altered

ion selectivity, and diminished $K_v2.1$ channel expression at the cell surface (14–17). These

findings are supported by rodent models, including *Kcnb1* knock-out (KO) and knock-in (KI)

mice, which exhibit altered behaviors such as locomotor hyperactivity, reduced anxiety-like

behavior, and lower seizure thresholds when exposed to chemoconvulsants (18–20).

Spontaneous seizures have been reported in homozygous KI $Kcnb1^{R/R}$ ($Kcnb1^{G379R/G379R}$) (18)

and the KI $Kcnb1^{R312H}$ mouse models (21).

The underlying pathogenic mechanisms of *KCNB1*-related DEEs remain largely

uncharacterized in rodent models, and face limitations, particularly regarding drug screening.

Zebrafish has gained prominence for studying human neurological disorders, including

epilepsy, due to its genetic manipulability, rapid development, and suitability for high-

throughput drug screening (22–25). Larval zebrafish, when exposed to the proconvulsant

pentylenetetrazol (PTZ), manifest behaviors akin to seizures, making them a valuable model

for epilepsy investigations (26) such as Dravet syndrome related to *SCN1A* (26–28) and

epilepsies associated with *DEPDC5* (29,30). In addition, drug screening in larval zebrafish

has been performed in both genetic and chemically induced epilepsy models, demonstrating

the translational potential of this model by identifying compounds capable of reducing the

severity of epileptic seizures (31–34).

The zebrafish orthologue of *KCNB1, kcnb1,* shares 67% sequence homology with the

human gene. The expression of *kcnb1* starts at 19 hours post-fertilization (hpf), increases

during zebrafish development, and is mainly localized in the brain of adult zebrafish with

tissue-specific expression patterns comparable to mammals (35). The first *kcnb1* KO

zebrafish model (*kcnb1^{-/-}*) was generated using CRISPR/Cas9 mutagenesis, leading to a

premature stop codon between the N-terminal cytoplasmic domain and the first

transmembrane domain of the protein (36). This *kcnb1^{-/-}* model revealed developmental

abnormalities such as a small proportion of gastrulation defect and reduced brain ventricles

(36) along with disruption of inner ear development (35), emphasizing *kcnb1*'s role in

developmental signaling pathways.

However, the effects of *kcnb1* loss-of-function on brain in development, in the context of

epilepsy, remain unexplored. In this study, we aim to characterize the behavioral,

electrophysiological and molecular consequences of this *kcnb1^{-/-}* zebrafish model at various

developmental stages. This analysis will shed light on the key features of *KCNB1*-related

DEEs, contributing to a better understanding of the pathophysiological mechanisms

underlying this disorder.

2.   MATERIALS AND METHODS

2.1. Housing conditions of zebrafish

All procedures were approved by the Institutional Ethics Committees at the Research

Centers of IMAGINE Institute (INSERM U1163, Paris, France) and were performed in

accordance with the European Union Directive (2010/63/EU). Adult zebrafish (Danio Rerio)

were housed in a conventional animal facility. Experiments were performed on wild-type AB

and TU strains (Tübingen, Germany) and *kcnb1* mutant zebrafish lines (*kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$)

staged from 0- to 6-days post-fertilization (dpf) (37). Fertilized eggs were collected by natural

spawning. Embryos and larvae were maintained at 28 ± 1 °C in a non-CO_2 incubator (VWR)

with a 14h/10h light/dark cycle in embryo medium (Instant Ocean).

2.2. kcnb1 mutant zebrafish lines

Heterozygous *kcnb1* mutant embryos (*kcnb1*$^{+/-}$) were provided by Dr. Vladimir Korzh

(Warsaw, Poland). The *kcnb1* mutant line was generated using CRISPR/Cas9 mutagenesis,

resulting in a premature stop codon (36). *kcnb1*$^{+/-}$ mutants were raised to adulthood and

crossed to produce *kcnb1*$^{-/-}$ zebrafish for experiments (*kcnb1*$^{sq301/sq301}$, ZDB-ALT-170417-2)

(36). Genomic DNA was extracted from adult zebrafish fins. Samples were amplified by PCR

(Bio-Rad) using DreamTaq Hot Start PCR Master Mix (Thermo Scientific) and primers

targeting *kcnb1* (10 mM): F_5'-TGTGACGACTACAACCTGGA-3' and R_5'-

CTCCTCGTTCATCTGCTCCT-3'. DNA samples were sent for sequencing to GATC

Biotech, Eurofins Genomics.

2.3. Survival assay and morphological analysis

Zebrafish embryos and larvae were maintained at 28 ± 1 °C non-CO_2 incubator (VWR),

and were fed daily from 0 to 15 dpf. After 6 dpf, larvae were transferred to tanks with dripped

water flux. Survival was assessed based on the number of living individuals relative to the

total population. Zebrafish were photographed using a stereomicroscope (SZX16, Olympus

Life Science). Measurements of the body length and the head surface were performed

manually with ImageJ software (National Institutes of Health, NIH).

2.4. Reverse transcription quantitative polymerase chain reaction (RT-qPCR)

cDNA from larval zebrafish at 6 dpf (30 per pool) was synthesized using 5X All-In-One

RT MasterMix (abm, Canada). qPCR on *kcnb1* and *bdnf* was performed with BlasTaq 2X

qPCR Master Mix (abm) on a BioRad CFX384 System (Bio-Rad). Relative gene expression

was determined by the $2^{-\Delta\Delta Ct}$ method, normalized to *β-actin* or *ef1α*, with *kcnb1*$^{+/+}$ serving as

the reference (relative fold change = 1). The following primers were used for

qPCR : *β-actin* (F_5'-CGAGCTGTCTTCCCATCCA-3', R_5'-

TCACCAACGTAGCTGTCTTTCTG-3'); *ef1α* (F_5'-CTGGAGGCCAGCTCAAACAT-3',

R_5'-ATCAAGAAGAGTAGTACCGCTAGCATTAC-3'); *kcnb1* (F_5'-

TGAAGTTCCGGGAGAGTGTT-3', R_5'-CAGGTTGGCGATGTCGTTCT-3'); *bdnf (F_5'-*

GACTCGAAGGACGTTGACCTGTA-3', R_5'-CGGCTCCAAAGGCACTTG-3').

2.5. Immunoprecipitation (IP) and western blot of kcnb1

Tissues from larval zebrafish at 6 dpf (30 per pool) were homogenized in lysis buffer

supplemented with protease inhibitors (Roche). Lysates were centrifugated at $12,000 \times g$ for

15 minutes at 4°C followed by a BCA Protein Assay Kit (Invitrogen). Immunoprecipitation

was performed to detect kcnb1 by using the Immunoprecipitation Kit dynabeads protein G

(Invitrogen), following the manufacturer's protocol. 200 µg of the supernatant with 4 µg of

anti-kcnb1 (ProteinTech, #19963-1-AP) or anti-Pan actin (Thermo Scientific, #ms-1295) were

used and IgG antibody served as a negative control. Membranes from western blotting were

detected using an ECL reagent (Cytiva) and the ChemiDoc™ Imaging Systems (Bio-Rad).

Densities of kcnb1 bands were normalized to Pan-actin expression using ImageJ (NIH).

2.6. Locomotion assessment

Premotor activity. Tail coiling activity in zebrafish embryos was recorded every hour

from 24 hpf to 36 hpf. Sixty-second movies were obtained under a stereomicroscope with

darkfield illumination (SZX16, Olympus Life Science), recorded at 30 frames per second

(fps). Premotor activity was counted manually using ImageJ software (NIH).

Touch-evoked escape response (TEER). The tail of 48 hpf-embryos was mechanically

stimulated, and their swimming trajectories were recorded at 30 fps using SpinView

V2.0.0.147 software (FLIR Systems lnc.). The trajectory was traced using the Manual

Tracking Plug-in in ImageJ software (NIH), and analyzed for distance, velocity, and time

spent in motion (see **Figure 3A**).

Spontaneous locomotion and pentylenetetrazol (PTZ)-induced seizures. Larvae were

acclimatized in a 48-well plate (Fisher Scientific) with lights off for 15 min into a Zebrabox

equipped with Zebralab 438 software (Viewpoint Life Sciences). Larvae were submitted to

two different protocols: 1) a light-induced protocol of 60 minutes recording split into 10 min

light/dark conditions repeated three times and 2) a sound-induced protocol of four repeated

audio stimuli (450 hertz, 80 decibels, 1 second/audio). The 5 seconds following audio stimuli

were analyzed. To evaluate the effect of PTZ on locomotor activity, larvae were first recorded

for 30 min to determine baseline activity levels. Fresh 5 mM PTZ (Sigma-Aldrich) was then

added and zebrafish were recorded for a further 30 min (see **Figure 4A**).

2.7. Electrophysiological analysis

Larval zebrafish at 5 and 6 dpf were embedded in 1% low-melting-point agarose (Sigma-

Aldrich) and covered with artificial cerebro□spinal fluid (ACSF, pH 7.8, osmolarity to 290-

295 mOsm/l). A microelectrode (2–7 MΩ) was filled with ACSF and implanted into the optic

tectum, using an Olympus microscope (Infinity 3S, 10X magnification). Local field potential

(LFP) recordings were obtained using a MultiClamp700B amplifier (Molecular devices)

coupled with an Axon Digidata 1550 (Molecular devices) and the Clampex V11.1 software

(Molecular devices). Baseline recordings of 30 min duration were assessed to larvae before

adding 40 mM PTZ (Sigma-Aldrich). After 5 min of treatment, the recording continued for 30

min. Negative spikes were automatically analyzed using Clampfit V11.2 software (Molecular

devices) during a 10 min interval, using a low-pass Gaussian filter of 560 Hz and a digital

reduction of 10 (see **Figure 5A**).

2.8. Immunohistochemistry on slices

Zebrafish aged from 48 hpf to 6 dpf were fixed as previously described (30). Brain slices of zebrafish (20 µm) were obtained using the cryostat CM3050S (Leica). After permeabilization for 30 minutes, the slices were incubated overnight at 4°C in blocking solution containing primary antibodies : Kcnb1 (1:100, Tebubio, #PAB7569), NeuN (1:100, Merck, #ABN90), Olig2 (1:100, DSHB, #PCRP-OLIG2-1E9-s), CX3CR1 (1:100, ProteinTech, #13885-1-AP), Ankyrin G (1:100, Proteintech, #27980-1-AP) and HuC (1:100, Tebubio, #FNab04072). Secondary antibodies (Alexa Fluor 488, Alexa Fluor 568 and Alexa Fluor 647, Thermo Scientific) were incubated in blocking solution for 2h at room temperature (RT). Slices were incubated in DAPI (Invitrogen, #D3571) followed by mounting on glass slide in Immu-Mount mounting medium (Epredia, #9990402). Images were captured using a Spinning disk Zeiss system (Carl Zeiss). Colocalization, represented in white, was determined using Z-stack projection on IMARIS V10.1.0 software (Oxford Instruments).

2.9. Whole-mount immunohistochemistry

Whole-mount immunohistochemistry on 48 hpf and 6 dpf zebrafish was performed according to a previously established protocol (38). Embryos were treated with 0,003% 1-Phenyl-2-thiourea (PTU) (Sigma-Aldrich) to prevent pigmentation of the skin. Briefly, zebrafish were fixed with 4% formaldehyde (Sigma-Aldrich) for 2h at RT. A dehydration with methanol (Sigma-Aldrich) was followed by a gradual rehydration. Zebrafish were permeabilized in fresh acetone (Sigma-Aldrich) for 20 min and blocked with 10% Bovine standard-albumin (BSA) (Eurobio Scientific) overnight at 4°C. They were incubated with the primary antibody (1% BSA / 0,1%PBST) at 4°C for two days. Primary antibodies used at 1:100 were: acetylated tubulin (Sigma-Aldrich, #T7451), 3A10 (DSHB, #3A10) and c-Fos (Santa Cruz, #sc-166940). Secondary antibodies were incubated in 1% BSA / 0,1% PBST overnight at 4°C using Alexa Fluor 488 and Alexa Fluor 647 (Thermo Scientific) at 1:250. After passing through increased percentages of glycerol (Sigma-Aldrich), image acquisition

was performed under a Spinning disk Zeiss system (Carl Zeiss). The same parameters were

applied to all images using ImageJ (NIH).

2.10. GABA and Glutamate ELISA assays

Based on the PTZ treatment protocol, the head of 6 dpf larvae (50 per pool) was collected

at the end of each 30 min-period recording of the basal activity and 5 mM PTZ treatment (see

Figure 5A). Same reagents were used as the IP protocol for tissue homogenizing. Enzyme-

linked immunosorbent assay (ELISA) was performed to quantify gamma-aminobutyric acid

(GABA, immusmol, #BA E-2500) and glutamate (immusmol, #BA E-2400) from 20 µg of

supernatant, following the manufacturer's protocol.

2.11. Statistical analysis

Statistical analysis was performed using GraphPad Prism V.8 software (GraphPad

Software). Statistical tests used are specified in the legend of each figure. Values represent the

mean ± standard error of the mean (SEM). Results were significant when the p-value was <

0.05. For all experiments, at least 3 single experiments were performed (N) with a certain

number of embryos or larvae per condition (n).

3.   RESULTS

3.1. kcnb1 is expressed in diverse cell-subtypes and regions of the CNS in Wild-type

larval zebrafish

Previous studies have shown significant expression of *kcnb1* in the eyes, ears, and central

nervous system (CNS) of wild-type (WT) larval zebrafish using whole-mount *in situ*

hybridization (36). To further investigate the distribution of the kcnb1 protein in the brain,

we performed immunohistochemistry on WT larvae at 6 dpf (**Figure 1A-1F, Supporting**

Information 1A-1B). Our findings revealed broad expression of kcnb1 throughout multiple

regions of the CNS, such as the diencephalon, midbrain, telencephalon and hindbrain

(Figure 1A-1C, Supporting Information 1A) including the spinal cord **(Figure 1C)**,

although the protein was difficult to detect in the eyes. Further analysis demonstrated that

kcnb1 is expressed in various CNS cell subtypes. Notably, kcnb1 is localized in neurons as

evidenced by colocalization with NeuN, a neuronal nuclear marker, and further confirmed

by Ankyrin G marker, expressed on the axonal initial segment of neurons **(Figure 1D,**

Supporting Information 1B). kcnb1 was also found in oligodendrocytes, marked by Olig2,

a transcription factor **(Figure 1E)**, and in microglial cells, identified by CX3CR1, a

fractalkine receptor marker **(Figure 1F)**. These results provide the first characterization of

kcnb1 expression in distinct CNS cell subtypes in zebrafish.

3.2. kcnb1 knock-out expression and morphological analyses

Using a *kcnb1* knock-out zebrafish model generated by Shen et collaborators (2016), we

investigated loss-of-function effects of *kcnb1* in the context of DEEs. We first confirmed the

genomic sequence of *kcnb1*$^{-/-}$ fish by Sanger sequencing, identifying a distinct 2-G bases

deletion along with a 14-bp insertion on chromatograms (Δ14bp) **(Figure 2A)**. Real time

qPCR analysis revealed a significant reduction in *kcnb1* transcript expression by 44% in

kcnb1$^{+/-}$ and by 56% in *kcnb1*$^{-/-}$ zebrafish larvae compared to WT at 6 dpf **(Figure 2B)**.

Despite this reduction in transcript levels, kcnb1 protein expression remained comparable

across all genotypes **(Figure 2C)**. The survival rates of both *kcnb1*$^{+/-}$ *and kcnb1*$^{-/-}$ lines from 0

to 15 dpf were similar to those of WT **(Figure 2D and Table 1)**. Measurements of body

length and head surface of *kcnb1* mutants (*kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$) showed no significant

morphological differences compared to WT at 48 hpf and 6 dpf **(Figure 2E-2G)**, corroborated

by using the pan-neuronal marker HuC **(Supporting information 1C)**. Visualization of

neuronal fibers with an acetylated tubulin marker, revealed no major neuronal loss or

organizational defects due to *kcnb1* loss-of-function during early development **(Figure 2H)**,

supported by similar Mauthner cell body length observed across conditions at 48 hpf (M-cell,

3A10 marker, **Supporting information 1D-1E**). These results indicate that *kcnb1* loss-of-

function does not impact the normal growth of fish during early development.

3.3. Loss of kcnb1 leads to altered behavior phenotype, light and sound-induced

locomotor impairments

KCNB1 mutations in patients are associated with a range of locomotor disabilities

including hyperactivity, myoclonia, ataxia and hypotonia (9). First evidence of motor

impairments was observed in both *kcnb1* mutants showing a drastic decreased of tail coiling

activity compared to WT zebrafish from 24 to 36 hpf **(Supporting information 2A-2B)**.

Using the touch-evoked escape responses (TEER) test at 48 hpf **(Figure 3A)**, *kcnb1$^{+/-}$* and

kcnb1$^{-/-}$ zebrafish exhibited rapid circular swimming, in contrast to the straight-line swimming

observed in WT condition **(Figure 3B)**, significantly increased in terms of total distance

swam, velocity and time spent in motion **(Figure 3C-3E)**. Notably, within the same condition,

we observed a variability in the swimming behavior of embryos classified into two groups:

those with a severe phenotype (completing a minimum of two swim circles) and those with a

mild phenotype (grouping other trajectories) **(Table 2)**. We found that 30% of *kcnb1$^{+/-}$*

embryos displayed a severe phenotype compared to 52% of *kcnb1$^{-/-}$* zebrafish, with an

increase of studied parameters **(Table 2, Supporting information 2C-2F)**. While all three

conditions showed low locomotor activity in response to light changes at 3 and 4 dpf, this

activity significantly increased in all conditions at later stages **(Figure 3H)**. However, starting

at 5 dpf, *kcnb1$^{-/-}$* larvae displayed pronounced locomotor hyperactivity compared to WT and

kcnb1$^{+/-}$ zebrafish **(Figure 3H)**. This finding is aligned with results from **Figure 3G and 3F**,

showing significant locomotor hyperactivity in response to light/dark transitions. In a second

protocol, both *kcnb1$^{+/-}$* and *kcnb1$^{-/-}$* mutants exhibited a significant decrease in locomotor

activity after each audio stimulus compared to the WT condition **(Figure 3I)**. These results

suggest that the locomotor phenotype may be due to dysregulation of electrical neuronal

activity affecting different sensorimotor pathways.

3.4. kcnb1$^{-/-}$ zebrafish exhibit increased sensitivity to PTZ-induced seizures and elevated

expression of epileptogenesis-related genes

Previous studies have shown that exposure of zebrafish larvae to pentylenetetrazol (PTZ)

increases seizure-like behavior in a concentration-dependent manner (26). We recorded the

baseline locomotor activity of 6 dpf larvae for 30 minutes, followed by a 30-minute exposure

to 5 mM PTZ **(Figure 4A)**. Low baseline locomotor activity of *kcnb1* mutants was similar to

that of WT larvae **(Figure 4B-4C)**. However, after PTZ exposure, *kcnb1* mutants (*kcnb1$^{+/-}$*

and *kcnb1$^{-/-}$*) exhibited significantly higher swimming activity, characterized by fast circles

(Figure 4B-4C). To explore potential molecular disruptions, we measured *bdnf* (Brain-

Derived Neurotrophic Factor) mRNA expression, a neurogenesis and epileptogenesis-related

gene, before and after PTZ treatment **(Figure 4A and 4D)**. *bdnf* expression was similar

between conditions during baseline activity, although significantly increased after PTZ

treatment in the *kcnb1$^{-/-}$* condition as compared to *kcnb1$^{+/-}$* and WT zebrafish **(Figure 4D)**. In

addition, we assessed c-Fos expression, an early gene marker of epileptic seizures, in the

telencephalon of 6 dpf larvae **(Figure 4A, 4E-4G, Supporting information 3)**. The number

of c-Fos-positive neurons was similar across all conditions, both before and after PTZ

treatment **(Figure 4F)**. We analyzed the distribution of activated neurons according to the

fluorescence intensity value of c-Fos, reflecting the level of neuronal activation divided in:

low (0-25%), moderately low (25-50%), moderately high (50-75%) and high (75-100%)

neuronal activation **(Figure 4G, Supporting information 3B)**. *kcnb1$^{-/-}$* fish showed a shift

towards higher levels of neuronal activation, with a significant increase in moderately low and

moderately high activation levels after PTZ treatment **(Figure 4G, Supporitng information**

3B). These results suggest that the *kcnb1* LOF model exhibits neurogenesis impairments in

the developing brain of zebrafish.

3.5. kcnb1 knock-out zebrafish model show spontaneous and provoked "epileptic"-like

seizures associated with disrupted GABA regulation

Abnormal electrographic activity has been observed in zebrafish models with chemically

provoked-epileptic seizures (26,31,34). We recorded local field potentials (LFPs) from the

optic tectum of 6 dpf larvae during a 30-minute "basal activity" phase, followed by a 40 mM

PTZ treatment, recorded for an additional 30 minutes **(Figure 5A-5B)**. Prior to PTZ

treatment, *kcnb1*$^{-/-}$ larvae showed significantly increased spontaneous neuronal activity,

reflected by the number of spikes, compared to WT **(Figure 5Ca-5D)**. In contrast, *kcnb1*$^{+/-}$

larvae did not exhibit spontaneous seizures, displaying a similar profile to untreated WT

zebrafish **(Figure 5Ca-5D)**. In response to 40 mM PTZ exposure, WT larvae exhibited a

significant increase in number of spikes compared to the non-treated condition **(Figure 5Ca-**

5D). The *kcnb1*$^{+/-}$ model showed a similar response profile, although the duration of seizure

events was significantly longer than in PTZ-treated WT larvae **(Figure 5Ca-5E)**.

Interestingly, *kcnb1*$^{-/-}$ larvae had a significantly higher number of PTZ-induced events,

although event duration was comparable to WT-treated zebrafish **(Figure 5Ca-5E)**. This

kcnb1$^{-/-}$ model also displayed various « epileptic »-like signals seen in *KCNB1*-related DEE

patients, including polyspike discharges **(Figure 5Cb)**, 'ictal'-like activity **(Figure 5Cc)**, and

large amplitude spikes **(Figure 5Cd)**. To further confirm dysregulated neuronal activity in the

kcnb1 LOF model, we measured gamma-aminobutyric acid (GABA) and glutamate

concentrations in the heads of 6 dpf zebrafish following the same protocol and exposed to 5

mM PTZ **(Figure 5A)**. During the pre-treatment period, GABA levels were similar in

kcnb1$^{+/-}$ and WT larvae, while *kcnb1*$^{-/-}$ larvae showed a significant increase **(Figure 5F)**.

Although PTZ treatment significantly increased GABA concentration in WT zebrafish, it did

not alter GABA levels in the *kcnb1$^{+/-}$* model, which remained low, or in the *kcnb1$^{-/-}$* model,

showing similar high concentration of GABA as WT-treated larvae **(Figure 5F)**. Glutamate

levels were similar between all genotypes before PTZ exposure and remained unchanged in

post-treatment **(Figure 5G)**. These results suggest that the *kcnb1$^{-/-}$* model exhibits

spontaneous ans chemically induced epileptiform-like electrographic activity, along with

disrupted GABA regulation.

4. DISCUSSION

In this study, *kcnb1* knock-out in zebrafish results in early-onset phenotypes mimicking

key features of *KCNB1*-related DEE. This LOF impacts neuronal functions in particular

inhibitory pathways in developemental brains.

4.1. kcnb1 expression in the CNS

The study of kcnb1 expression revealed the presence of the protein across various regions

of the zebrafish CNS, including the diencephalon, midbrain, telencephalon, and hindbrain,

consistent with previously reported *in situ* hybridization of *kcnb1* by Shen and collaborators

(2016). This broad expression supports previous findings that kcnb1 is crucial for maintaining

neuronal excitability, starting from 19 hpf (35). Notably, the presence of kcnb1 in multiple

cell subtypes suggests its diverse functional roles, ranging from neuronal signaling to

potential involvement in glial cell function and neuroinflammation in zebrafish.

4.2. kcnb1 knock-out and developmental consequences

Despite a significant reduction in *kcnb1* transcript levels in both heterozygous and

homozygous mutants, the protein expression was similar between mutant models and the WT

condition, suggesting a potential genetic compensation. We observed no major

morphological, brain anatomical abnormalities or impaired survival rates in *kcnb1$^{-/-}$* zebrafish

during the early development. This finding is consistent with clinical observations in *KCNB1*-

related DEE patients, where normal brain morphology is reported in most cases despite severe

neurological symptoms (9). This suggests that *kcnb1* LOF does not impact brain structural

development, although it may influence more neuronal function aspects.

4.3. Altered behavioral phenotypes and sensorimotor dysregulation

Behavioral assays revealed that $kcnb1^{-/-}$ zebrafish exhibit significant locomotor

impairments, including hyperactivity, altered swimming patterns, and exaggerated responses

to sensory stimuli such as light and sound. Indeed, we showed that $kcnb1^{+/-}$ and $kcnb1^{-/-}$

embryos had reduced tail coiling activity from 25 hpf to 31hpf. This early stereotyped

behavior is supported by synchronized spinal locomotor circuits (39). Previous reports on

models of epilepsy have revealed a corkscrew-like trajectory swimming characteristic of

epileptogenic-like activity (22). We found a similar circular pattern of swim trajectories in

kcnb1 mutant models at 48 hpf, but revealed a huge variability within the same genotype.

These results might reflect the variable spectrum of behavioral and cognitive impairments

observed in patients (12). Furthermore, $kcnb1^{-/-}$ mutants exhibited increased response of

locomotor activity to a light-dark stimulus as compared to $kcnb1^{+/-}$ and WT lines, but

presented a decreased locomotor activity after a succession of audio stimuli. This last result is

in correlation with the data obtained by Jedrychowska and collaborators (2021). These

phenotypes closely mimic the motor dysfunctions observed in patients with *KCNB1*

mutations, such as ataxia and hyperactivity and suggest a dysregulation of sensorimotor

pathways.

4.4. Seizure susceptibility and electrophysiological abnormalities

The seizure-like behavior in $kcnb1^{+/-}$ and $kcnb1^{-/-}$ models is supported by their increased

locomotor activity characterized by fast circles trajectories induced by 5 mM PTZ compared

to wild-type. These results are concomitant with the stage II of the "seizure-like behavior

score" described by Baraban and collaborators (2005). Our electrophysiological recordings

further demonstrated that *kcnb1*$^{-/-}$ zebrafish exhibit spontaneous and chemically induced

epileptiform seizure-like activity. The absence of spontaneous seizures in the *kcnb1*$^{+/-}$ line

could reflect the existence of functional compensatory pathways masking the phenotypic

features due to reduced levels of *kcnb1* mRNA but similar protein expression as WT larvae.

Epileptiform seizure-like activity was confirmed by c-Fos acute neuronal hyperactivation in

the telencephalon of *kcnb1*$^{-/-}$ zebrafish. Moreover, the quantification of brain-derived

neurotrophic factor *(bdnf)* mRNA in larvae, a neurotrophin crucial for brain development and

synaptic plasticity, revealed a significant upregulation of its expression level within the *kcnb1*$^{-}$

$^{/-}$ line. This finding indicates deficits in neurogenesis within the developing brain of *kcnb1*$^{-/-}$

zebrafish. Similar results were obtained by identifying *c-Fos* and *bdnf* as genes associated

with seizures in zebrafish, with increased expression levels observed after exposure to 20 mM

PTZ 20mM (40). These findings are particularly relevant to understanding the

pathophysiology of DEEs, where patients often present with refractory seizures and abnormal

electroencephalogram patterns.

4.5. Neurotransmitter dysregulation in kcnb1$^{-/-}$ *zebrafish*

Our study also highlights the dysregulation of GABA in *kcnb1*$^{-/-}$ zebrafish, with

significantly elevated GABA levels observed in both baseline and post-PTZ conditions. This

disrupted GABAergic signaling likely contributes to the observed seizure phenotypes, as

GABA is a key inhibitory neurotransmitter involved in maintaining the balance of excitatory

and inhibitory signals in the CNS. The lack of significant changes in glutamate levels suggests

that *kcnb1* LOF primarily affects inhibitory pathways, leading to an imbalance that favors

neuronal hyperexcitability.

5. CONCLUSION

Our investigation suggests that the *kcnb1* loss-of-function zebrafish model provides a

valuable model to reproduce key features of *KCNB1*-related DEE, including early behavioral

disturbances, increased-susceptibility to epileptic seizures and neurotransmitter dysregulation.

Notably, this model could be used for discovering of new therapeutic compounds that may

improve the long-term prognosis of individuals with *KCNB1*-related DEE.

433 AUTHOR CONTRIBUTIONS

Lauralee Robichon designed and conceptualized the study, performed the experiments,

analyzed and interpreted the data, and wrote the manuscript. Claire Bar designed some parts

of the study, performed some behavioral assays and wrote the manuscript. Anca Marian

performed some behavioral assays and was responsible for technical support. Lisa Lehmann

performed some locomotor activity experiments and analyzed some data. Solène Renault was

responsible for technical support. Edor Kabashi supervised the work, revised the paper and

approved the manuscript. Sorana Ciura supervised the work, revised the paper and approved

the manuscript. Rima Nabbout supervised the work, revised the paper and approved the

manuscript. All authors revised and approved the final version of the manuscript.

444 ACKNOWLEGMENTS

This work was supported by grants from the Agence Nationale de la Recherche under

"Investissements d'avenir" program (ANR☐10IAHU☐01), the Fondation Bettencourt

Schueller (Rima Nabbout and Claire Bar), the Ligue Française Contre l'Épilepsie (Claire Bar),

the ERC Consolidator Grant (Edor Kabashi). Rima Nabbout and Lauralee Robichon are

supported by the Chair Geen-DS funded by FAMA fund hosted by Swiss Philanthropy

Foundation. Lauralee Robichon is recipient of a grant from the Fondation pour la Recherche

Médicale (grant number: PLP202009012460, 2021). The work was supported by the

"Association KCNB1 France".

We are grateful of the team of Dr. Vladimir Korzh (International Institute of Molecular

and Cell Biology of Warsaw, Poland) for kindling provided us the *kcnb1* knock-out transgenic

zebrafish line. We appreciate the help of the LEAT Zebrafish facilities and Cell Imaging

Facility of Imagine Institute for respectively fish maintenance and expert technical help. The

help of Nicolas Goudin, responsible of the Image analysis center of SFR Necker (Paris,

France) was highly appreciated for immunofluorescence quantification. We thank DSHB for

the antibodies used for immunohistochemistry: 3A10 was deposited to the DSHB by Jessell,

T.M. / Dodd, J. / Brenner-Morton, S. (DSHB Hybridoma Product 3A10), PCRP-OLIG2-1E9

was deposited to the DSHB by Common Fund – Protein Capture Reagents Program (DSHB

Hybridoma Product PCRP-OLIG2-1E9).

DISCLOSURE

None of the authors has any conflict of interest to disclose.

We confirm that we have read the Journal's position on issues involved in ethical publication

and affirm that this report is consistent with those guidelines.

REFERENCES

1. Bishop HI, Guan D, Bocksteins E, Parajuli LK, Murray KD, Cobb MM, et al. Distinct

Cell- and Layer-Specific Expression Patterns and Independent Regulation of Kv2

Channel Subtypes in Cortical Pyramidal Neurons. J Neurosci. 2015 Nov 4;35(44):14922–

42.

2. Du J, Tao-Cheng JH, Zerfas P, McBain CJ. The K+ channel, Kv2.1, is apposed to

astrocytic processes and is associated with inhibitory postsynaptic membranes in

hippocampal and cortical principal neurons and inhibitory interneurons. Neuroscience. 1998 Feb;84(1):37–48.

3. Sarmiere PD, Weigle CM, Tamkun MM. The Kv2.1 K+ channel targets to the axon initial segment of hippocampal and cortical neurons in culture and in situ. BMC Neurosci. 2008 Dec;9(1):112.

4. Trimmer JS. Immunological identification and characterization of a delayed rectifier K+ channel polypeptide in rat brain. Proc Natl Acad Sci. 1991 Dec 1;88(23):10764–8.

5. Antonucci DE, Lim ST, Vassanelli S, Trimmer JS. Dynamic localization and clustering of dendritic Kv2.1 voltage-dependent potassium channels in developing hippocampal neurons. Neuroscience. 2001 Dec;108(1):69–81.

6. Mohapatra DP, Misonou H, Sheng-Jun P, Held JE, Surmeier DJ, Trimmer JS. Regulation of intrinsic excitability in hippocampal neurons by activity-dependent modulation of the K $_V$ 2.1 potassium channel. Channels. 2009 Jan;3(1):46–56.

7. Johnson B, Leek AN, Solé L, Maverick EE, Levine TP, Tamkun MM. Kv2 potassium channels form endoplasmic reticulum/plasma membrane junctions via interaction with VAPA and VAPB. Proc Natl Acad Sci. 2018 Jul 31;115(31). Available from: https://pnas.org/doi/full/10.1073/pnas.1805757115

8. Vierra NC, Kirmiz M, van der List D, Santana LF, Trimmer JS. Kv2.1 mediates spatial and functional coupling of L-type calcium channels and ryanodine receptors in mammalian neurons. eLife. 2019 Oct 30;8:e49953.

9. Bar C, Barcia G, Jennesson M, Le Guyader G, Schneider A, Mignot C, et al. Expanding the genetic and phenotypic relevance of KCNB1 variants in developmental and epileptic encephalopathies: 27 new patients and overview of the literature. Hum Mutat. 2020 Jan;41(1):69–80.

10. de Kovel CGF, Syrbe S, Brilstra EH, Verbeek N, Kerr B, Dubbs H, et al. Neurodevelopmental Disorders Caused by De Novo Variants in *KCNB1* Genotypes and Phenotypes. JAMA Neurol. 2017 Oct 1;74(10):1228.

11. Marini C, Romoli M, Parrini E, Costa C, Mei D, Mari F, et al. Clinical features and outcome of 6 new patients carrying de novo *KCNB1* gene mutations. Neurol Genet. 2017 Dec;3(6):e206.

12. Bar C, Kuchenbuch M, Barcia G, Schneider A, Jennesson M, Le Guyader G, et al. Developmental and epilepsy spectrum of KCNB1 encephalopathy with long-term outcome. Epilepsia. 2020 Nov;61(11):2461–73.

13. Bar C, Breuillard D, Kuchenbuch M, Jennesson M, Le Guyader G, Isnard H, et al. Adaptive behavior and psychiatric comorbidities in KCNB1 encephalopathy. Epilepsy Behav. 2022 Jan;126:108471.

14. Kang SK, Vanoye CG, Misra SN, Echevarria DM, Calhoun JD, O'Connor JB, et al. Spectrum of Kv2.1 Dysfunction in KCNB1 Associated Neurodevelopmental Disorders. Ann Neurol. 2019 Dec;86(6):899–912.

15. Saitsu H, Akita T, Tohyama J, Goldberg-Stern H, Kobayashi Y, Cohen R, et al. De novo KCNB1 mutations in infantile epilepsy inhibit repetitive neuronal firing. Sci Rep. 2015 Oct 19;5(1):15199.

16. Thiffault I, Speca DJ, Austin DC, Cobb MM, Eum KS, Safina NP, et al. A novel epileptic encephalopathy mutation in KCNB1 disrupts Kv2.1 ion selectivity, expression, and localization. J Gen Physiol. 2015 Nov 1;146(5):399–410.

17. Torkamani A, Bersell K, Jorge BS, Bjork RL, Friedman JR, Bloss CS, et al. De novo KCNB1 mutations in epileptic encephalopathy: KCNB1 Mutations. Ann Neurol. 2014 Oct;76(4):529–40.

18. Hawkins NA, Misra SN, Jurado M, Kang SK, Vierra NC, Nguyen K, et al. Epilepsy and

neurobehavioral abnormalities in mice with a dominant-negative KCNB1 pathogenic

variant. Neurobiol Dis. 2021 Jan;147:105141.

19. Kang SK, Hawkins NA, Thompson CH, Baker EM, Echevarria-Cooper DM, Barse L, et

al. Altered neurological and neurobehavioral phenotypes in a mouse model of the

recurrent KCNB1-p.R306C voltage-sensor variant. Neurobiol Dis. 2024

May;194:106470.

20. Speca DJ, Ogata G, Mandikian D, Bishop HI, Wiler SW, Eum K, et al. Deletion of the

Kv2.1 delayed rectifier potassium channel leads to neuronal and behavioral

hyperexcitability: Kv2.1 deletion and hyperexcitability. Genes Brain Behav. 2014

Apr;13(4):394–408.

21. Bortolami A, Yu W, Forzisi E, Ercan K, Kadakia R, Murugan M, et al. Integrin-KCNB1

potassium channel complexes regulate neocortical neuronal development and are

implicated in epilepsy. Cell Death Differ. 2022 Oct 7; Available from:

https://www.nature.com/articles/s41418-022-01072-2

22. Gawel K, Langlois M, Martins T, van der Ent W, Tiraboschi E, Jacmin M, et al. Seizing

the moment: Zebrafish epilepsy models. Neurosci Biobehav Rev. 2020 Sep;116:1–20.

23. Kolesnikova TO, Demin KA, Costa FV, Zabegalov KN, De Abreu MS, Gerasimova EV,

et al. Towards Zebrafish Models of CNS Channelopathies. Int J Mol Sci. 2022 Nov

12;23(22):13979.

24. Sakai C, Ijaz S, Hoffman EJ. Zebrafish Models of Neurodevelopmental Disorders: Past,

Present, and Future. Front Mol Neurosci. 2018 Aug 29;11:294.

25. Yaksi E, Jamali A, Diaz Verdugo C, Jurisch Yaksi N. Past, present and future of

zebrafish in epilepsy research. FEBS J. 2021 Dec;288(24):7243–55.

26. Baraban SC, Taylor MR, Castro PA, Baier H. Pentylenetetrazole induced changes in

zebrafish behavior, neural activity and c-fos expression. Neuroscience. 2005

Jan;131(3):759–68.

27. Tiraboschi E, Martina S, Ent W, Grzyb K, Gawel K, Cordero Maldonado ML, et al. New

insights into the early mechanisms of epileptogenesis in a zebrafish model of Dravet

syndrome. Epilepsia. 2020 Mar;61(3):549–60.

28. Zhang Y, Kecskés A, Copmans D, Langlois M, Crawford AD, Ceulemans B, et al.

Pharmacological Characterization of an Antisense Knockdown Zebrafish Model of Dravet

Syndrome: Inhibition of Epileptic Seizures by the Serotonin Agonist Fenfluramine.

Herault Y, editor. PLOS ONE. 2015 May 12;10(5):e0125898.

29. de Calbiac H, Dabacan A, Marsan E, Tostivint H, Devienne G, Ishida S, et al. Depdc5

knockdown causes mTOR-dependent motor hyperactivity in zebrafish. Ann Clin Transl

Neurol. 2018 May;5(5):510–23.

30. Swaminathan A, Hassan-Abdi R, Renault S, Siekierska A, Riché R, Liao M, et al. Non-

canonical mTOR-Independent Role of DEPDC5 in Regulating GABAergic Network

Development. Curr Biol. 2018 Jun;28(12):1924-1937.e5.

31. Afrikanova T, Serruys ASK, Buenafe OEM, Clinckers R, Smolders I, de Witte PAM, et

al. Validation of the Zebrafish Pentylenetetrazol Seizure Model: Locomotor versus

Electrographic Responses to Antiepileptic Drugs. Burgess HA, editor. PLoS ONE. 2013

Jan 14;8(1):e54166.

32. Baraban SC, Dinday MT, Hortopan GA. Drug screening in Scn1a zebrafish mutant

identifies clemizole as a potential Dravet syndrome treatment. Nat Commun. 2013 Sep

3;4(1):2410.

33. Li J, Nelis M, Sourbron J, Copmans D, Lagae L, Cabooter D, et al. Efficacy of

Fenfluramine and Norfenfluramine Enantiomers and Various Antiepileptic Drugs in a

Zebrafish Model of Dravet Syndrome. Neurochem Res. 2021 Sep;46(9):2249–61.

34. Shin JN, Lee KB, Butterworth W, Park SK, Kim JY, Kim S. Zebrafish EEG predicts the

efficacy of antiepileptic drugs. Front Pharmacol. 2022 Dec 8;13:1055424.

35. Jedrychowska J, Gasanov EV, Korzh V. Kcnb1 plays a role in development of the inner

ear. Dev Biol. 2021 Mar;471:65–75.

36. Shen H, Bocksteins E, Kondrychyn I, Snyders D, Korzh V. Functional antagonism of

alpha-subunits of Kv channel in developing brain ventricular system. Development. 2016

Jan 1;dev.140467.

37. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic

development of the zebrafish. Dev Dyn. 1995 Jul;203(3):253–310.

38. Santos D, Monteiro SM, Luzio A. General Whole-Mount Immunohistochemistry of

Zebrafish (Danio rerio) Embryos and Larvae Protocol. In: Félix L, editor. Teratogenicity

Testing. New York, NY: Springer New York; 2018. p. 365–71. (Methods in Molecular

Biology; vol. 1797). Available from: http://link.springer.com/10.1007/978-1-4939-7883-

0_19

39. Saint-Amant L, Drapeau P. Time course of the development of motor behaviors in the

zebrafish embryo. J Neurobiol. 1998 Dec;37(4):622–32.

40. Orellana-Paucar AM, Afrikanova T, Thomas J, Aibuldinov YK, Dehaen W, De Witte

PAM, et al. Insights from Zebrafish and Mouse Models on the Activity and Safety of Ar-

Turmerone as a Potential Drug Candidate for the Treatment of Epilepsy. Kalueff AV,

editor. PLoS ONE. 2013 Dec 13;8(12):e81634.

41. Hermann A, Sitdikova G, Weiger T. Oxidative Stress and Maxi Calcium-Activated

Potassium (BK) Channels. Biomolecules. 2015 Aug 17;5(3):1870–911.

606   **TABLES**

Conditions	Number of alive fish		% survival (0 vs 15 dpf)	Log-rank test at 15 dpf (mutant vs WT)
	0 dpf	15 dpf		
kcnb1$^{+/+}$	137	116	84,67	/
kcnb1$^{+/-}$	121	109	82,65	P = 0,7677
kcnb1$^{-/-}$	169	156	88,17	P = 0,3302

609   **Table 1: Survival table between 0 and 15 days post-fertilization (dpf).** Results show a

610   similar growth of *kcnb1* mutant fish (*kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$) as compared to control fish

611   (*kcnb1*$^{+/+}$) at early stages development (N = 3 repeats; n = 121 - 169 ZF/genotype, Log-rank

612   test, ns: non-significant).

	Total number of embryos	Mild phenotype (MP)	* Severe phenotype (SP)	% Severe phenotype
$kcnb1^{+/+}$	73	/	/	/
$kcnb1^{+/-}$	88	61	27	30,68%
$kcnb1^{-/-}$	50	24	26	52%

Table 2: Diverse phenotypes were observed during the Touch-Evoked Escape Response (TEER) of 48 hpf-mutant embryos. Within the same genotype, embryos were divided in severe phenotype (*: swimming trajectory with at least two swim circles; SP) and mild phenotype (other trajectories; MP) indicating a huge variability in swimming behavior after a TEER test of $kcnb1^{+/-}$ and $kcnb1^{-/-}$ fish (N = 3 repeats; n = 50-88 ZF/genotype).

Condition	*kcnb1*$^{+/+}$			*kcnb1*$^{+/-}$					*kcnb1*$^{-/-}$				
Animal	1	2	3	1	2	3	4	5	1	2	3	4	5
Number of spikes in basal activity per 10 min recording	0	5	4	12	6	17	2	0	68	20	21	66	26
Number of spikes after 40 mM of PTZ per 10 min recording	158	148	105	238	115	213	291	157	508	510	179	239	415

Table 3: Individual data of negative spikes number during electrophysiological

recordings into the optic tectum of zebrafish. Neuronal activity of fish was recorded during

30 minutes of basal activity followed by 30 minutes of 40 mM PTZ treatment. A period of 10

minutes in the middle of each period (basal activity and provoked seizures) was used to

analyze the number of negative spikes. The table shows a higher number of spikes for *kcnb1*$^{-/-}$

larvae in pre- and post-PTZ treatment as compared to WT fish. The following parameters

were applied for each recording: Resistance: 2-7 MΩ; Low-pass Gaussian filter: 560 Hz; Data

reduction: 10. n = 3-5 ZF/genotype.

FIGURE LEGENDS

Figure 1. kcnb1 is expressed in distinct cell subtypes and various regions of the central

nervous system in 6 dpf Wild-type zebrafish. A and B. Horizontal and transversal sections

of Wild-type (WT) zebrafish expressing cells (DAPI, blue) labelled with anti-kcnb1 (green),

showing a large expression of the protein in the central nervous system (CNS) at 6-day post-

fertilization (dpf). The protein is expressed in the telencephalon, diencephalon, midbrain

including the optic tectum, and the hindbrain comprising the cerebellum and the spinal cord

(**A:** scale bar 50 µm; magnification10x ; **B:** scale bar 30 µm; magnification 20x). **C.**

Horizontal section of a WT zebrafish at 6 dpf showing the presence of kcnb1 along the spinal

cord (scale bar 10 µm; magnification 63x). **D-F.** Horizontal sections of WT zebrafish

expressing cells (DAPI, blue), kcnb1 (green) and specific cell subtype markers, respectively

(D) a neuronal nuclear marker (NeuN, red, scale bar 30 µm; magnification 20x); **(E)** an

oligodendrocyte transcription factor 2 (Olig2, red, scale bar 10 µm; magnification 63x) and

(F) CX3C motif chemokine receptor 1 expressed in microglial cells (CX3CR1, red, scale bar

10 µm; magnification 63x). Images demonstrate the colocalization between kcnb1 and the 3

different cell subtypes markers, represented in white and marked by arrows. These results

indicate the presence of kcnb1 in neurons, oligodendrocytes and microglial cells.

Colocalization was determined using Z-stack projection on IMARIS V10.1.0 software

(Oxford Instruments). In figures, CNS regions are delimited by dotted lines. CX3CR1: CX3C

motif chemokine receptor 1; D: Diencephalon; DAPI: 4,6☐diamidino☐2☐phenylindole; E:

Eyes; H: Hindbrain; kcnb1: Potassium voltage-gated channel subfamily B member 1; M:

Midbrain; NeuN: Neuronal nuclear antigen; Olig2: Oligodendrocyte transcription factor; T:

Telencephalon. n = 3-4 fish/section.

Figure 2. Genotypic characterization of a *kcnb1* knock-out zebrafish model that is not

affected by brain anatomical defects. A. Schematic illustration of the generation of a *kcnb1*

knock-out zebrafish model obtained by Shen et al. (2016). The representation of the α-subunit

kcnb1 is adapted from a previous schematic picture (41). Insertion of a 14 base-pairs (bp)-

nucleotide sequence (identified in red) and of a 2 bp-nucleotide deleted (identified in black

bold type) into the first exon of the gene introducing a premature stop (Δ14bp) between the N-

terminal region and the first transmembrane domain of the protein, using the CRISPR-Cas9

system (targeted sequence: GGAGCTGGACTACTGGGGAG in *kcnb1* exon 1; ID zfin:

ZDB-ALT-170417-2; indel mutation; line: Kcnb1$^{sq301/sq301}$). **B.** RT-qPCR analysis of total

kcnb1 at 6 dpf demonstrating a significant decrease of *kcnb1* mRNA in *kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$

compared with *kcnb1*$^{+/+}$ fish (N = 4; n = 30/sample; One-Way ANOVA with Bonferroni post-

hoc test; **p<0,01). Data are normalized to actin mRNA expression and the condition

kcnb1$^{+/+}$ is considered as the reference value (relative fold change = 1). **C.**

Immunoprecipitation (IP) of kcnb1 in 6 dpf zebrafish demonstrating similar profile of protein

expression between the three genotypes (N = 3; n = 30/sample; One-Way ANOVA with

Bonferroni post-hoc test; ns: non signifcant). Input was used as a control of Western blot and

IgG antibody served as a negative control of IP. Data are normalized to Pan-actin protein

expression, with *kcnb1*[+/+] serving as the reference (relative fold change = 1). **D.** Kaplan-Meier

survival curve between 0 and 15 dpf showing that the partial or complete loss of *kcnb1* do not

impact the normal growth of fish at early stages of development (see **Table 1**; N = 3 repeats;

n = 121-169/genotype; Log-rank test; ns: non significant). **E.** Images showing that *kcnb1*[+/-]

and *kcnb1*[-/-] embryos and larvae do not show gross morphological changes at 48 hours post-

fertilization (hpf) and 6 dpf (scale bar: 300 µm). **F-G.** Quantification of major morphological

aspects at 48 hpf and 6 dpf including measurement of **(F)** body length and **(G)** head surface.

The *kcnb1* mutant zebrafish models (*kcnb1*[+/-] and *kcnb1*[-/-]) do not show any significant change

in each parameter at different developmental stages as compared to *kcnb1*[+/+] (N = 4 repeats; n

= 28-38 ZF/genotype; one-Way ANOVA with Bonferroni post-hoc test; ns: non-significant).

H. Whole-mount images of embryonic (48 hpf) and larvae (6 dpf) zebrafish immunostained

with anti-acetylated tubulin marker to identify global circuits of neuronal fibers (lateral and

dorsal view; 3D reconstruction; magnification 20x and scale bar at 100 µm; magnification 40x

and scale bar at 20 µm). The partial or complete loss of *kcnb1* does not affect the neuronal

brain density at different early stages of development (n = 5-7 ZF/genotype/developmental

stage).

Figure 3. Loss of *kcnb1* leads to altered behavior phenotype, light and sound-induced

locomotor impairments. A. Schematic representation of the touch evoked-escape response

(TEER) test. The tail of 48 hpf-embryos is mechanically stimulated, and the swimming

trajectory of zebrafish is recorded. **B.** Representative traces of individual swimming episodes

at 48 hpf showing the typical tortuous trajectory of *kcnb1* mutant models (*kcnb1*[+/-] and *kcnb1*[-

/-]) as compared to the straight-line trajectory of *kcnb1*[+/+] zebrafish (5 trajectories/genotype).

C-E. The quantification of the swimming trajectory tortuosity shows a significant increase of

different studied parameters in both *kcnb1* mutant conditions as compared to WT zebrafish, including an increase of **(C)** the distance swam, **(D)** the velocity and **(E)** the time spent in motion. Furthermore, within the same genotype, a huge variability in the swimming behavior has been observed and fish were divided into two distinct phenotypes: severe (at least two swim circles) and mild (other trajectories). The subdivision of both phenotypes is represented in **Table 2** and **Supporting information 2C-2F** (N = 3 repeats; n = 50-88 ZF/genotype; one-way ANOVA with Bonferroni post-hoc test; ns: non-significant; *p<0,05; **p<0,01; ***p<0,001; ****p<0,0001). **F.** Schematic representation of individual trajectory of 3 larvae zebrafish per genotype obtained with ViewPoint software (Zebrabox). Between 3 and 6 dpf, a protocol was applied to fish with 3 repetitions of 10 minutes in the light followed by 10 minutes in the dark (green lines: slow movements <8 mm/sec; red lines: fast movements >8 mm/sec). **G.** Average distance swam by larvae zebrafish at 6 dpf during the whole light-dark protocol described previously. Spontaneous locomotor hyperactivity was observed in *kcnb1*$^{-/-}$ larvae in the light and maintained significantly increased during the dark phase as compared to WT and *kcnb1*$^{+/-}$ larvae (N = 3 repeats; n = 35 ZF/genotype; one-way ANOVA with Bonferroni post-hoc test; *p<0,05; **p<0,01; ***p<0,001; ****p<0,0001). **H.** Representation of average distance travelled by larvae at different days of development (from 3 to 6 dpf) following the 10 min light-dark protocol descrived. At 3 and 4 dpf, controls and *kcnb1* mutant fish (*kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$) present low locomotor activity that was significantly increased from 5 to 6 dpf. A significant gap was observed starting from 5 dpf with a locomotor hyperactivity for *kcnb1*$^{-/-}$ larvae as compared to WT and *kcnb1*$^{+/-}$ fish, reflecting the result obtained in **Figure 3G** (N = 3 repeats; n = 35 ZF/genotype; one-way ANOVA with Bonferroni post-hoc test; the locomotor activity of zebrafish from the same genotype was compared from 4 to 6 dpf with a reference value at 3 dpf and the three genotypes were also compared between them; ns: non-significant; ***p<0,001; ****p<0,0001). **I.** Quantification

of the distance swam during the 5 seconds following each of the four audio stimuli applied

(450 Hertz; 80 decibel; 1 second) to larvae at 6 dpf. Both mutant conditions ($kcnb1^{+/-}$ and

$kcnb1^{-/-}$) present a significant decrease of the locomotor activity in response to audio stimuli

as compared to the WT condition (N = 3 repeats; n = 48-64 ZF/genotype; one-way ANOVA

with Bonferroni post-hoc test; *p<0,05).

Figure 4. *kcnb1* knock-out zebrafish exhibit increased locomotor sensitivity to PTZ and

elevated expression of epileptogenesis-related genes. A. Schematic representation of the

protocol followed to identify impact of PTZ on locomotion, bdnf and c-Fos expression. To

test seizure susceptibility in the *kcnb1* LOF model, we recorded the baseline locomotor

activity of 6 dpf larvae for 30 minutes, followed by a 30-minute exposure to 5 mM PTZ

(Zebrabox, ViewPoint). We then quantified two epileptogenesis-related genes: *bdnf* (Brain-

Derived Neurotrophic Factor) by qPCR and c-Fos by immunofluorescence. **B.** Schematic

representation of individual trajectory of 3 larvae zebrafish per genotype obtained with

ViewPoint software (Zebrabox). Fast swimming circles were observed for both mutant larvae

conditions ($kcnb1^{+/-}$ and $kcnb1^{-/-}$) after 30 minutes of 5 mM Pentylenetetrazol (PTZ)

treatment, a pro-convulsant. Green lines : slow movements (<8 mm/sec), red lines : fast

movements (>8 mm/sec). **C.** Global locomotor activity of 6 dpf-zebrafish into the dark was

recorded by applying a protocol of 30 minutes of basal activity followed by 30 minutes of

chemically induced-seizures using PTZ treatment at 5 mM. Both $kcnb1^{+/-}$ and $kcnb1^{-/-}$ larvae

show a significant increase in distance travelled during the 30 min-recording after the

chemical treatment as compared to WT zebrafish (N = 3 repeats; n = 27 ZF/genotype; one-

way ANOVA with Bonferroni post-hoc test; ns: non-significant; +/- and -/- vs +/+ after PTZ

treatment: **p<0,01; ****p<0,0001). **D.** RT-qPCR analysis of total *bdnf* at 6 dpf, a

neurogenesis and epileptogenesis-related gene, before and after 5 mM PTZ treatment (N = 3;

n = 30/sample; One-Way ANOVA with Bonferroni post-hoc test; ns: non significant;

****p<0,0001). Data are normalized to *ef1α* mRNA expression and *kcnb1$^{+/+}$* non-treated fish

are considered as the reference value (relative fold change = 1). The *kcnb1$^{-/-}$* presents a

tendency to an increased bdnf expression during the basal locomotor activity which is

confirmed by a significant increased expression after chemically induced-seizures as

compared to *kcnb1$^{+/+}$ and Kcnb1$^{+/-}$*. **E.** Whole-mount images of 6 dpf-zebrafish

immunostained with anti-c-Fos, an acute neuronal activation marker, obtained 30 minutes

after a basal activity period and a provoked-seizures period due to 5 mM PTZ treatment. The

telencephalon was the major region activated after the chemical treatment for each genotype

(n=8-11 ZF/condition, dorsal view, 3D reconstruction, scale bar: 50 µm, magnification 20x).

F. Quantification of the number of c-Fos positive neurons normalized to the volume (μm^3) of

the telencephalon of 6 dpf larvae-zebrafish during their basal activity or chemically treated

using the IMARIS V10.1.0 software (Oxford Instruments) (see **Supporting Information**

3A). The results did not show any difference in the number of activated neurons in both

mutant conditions *(kcnb1$^{+/-}$* and *kcnb1$^{-/-}$*) as compared to the WT condition (n = 8-11

fish/condition; Mann Whitney test; ns: non-significant). **G.** Distribution (in %) of activated

neurons according to the fluorescence intensity value of c-Fos in the telencephalon of fish at 6

dpf. Neuronal activation that was divided in four equal shares: low (0-25%), moderately low

(25-50%), moderately high (50-75%) and high (75-100%) neuronal activation. The three non-

treated conditions were presenting a similar distribution with a majority of low neuronal

activation of c-Fos positive neurons. However, chemically treated *kcnb1$^{-/-}$* zebrafish were

presenting a global higher activation with a significant shift to a moderately low and

moderately high neuronal activation (see **Supporting information 3B**). n = 8-11

fish/condition; Mann Whitney test; ns: non significant; *p<0,05; T: Telencephalon.

Figure 5. *kcnb1*[-/-] **larvae show spontaneous and provoked epileptiform-like electrographic activity associated with disrupted GABA regulation. A and B.** Schematic representation of the protocol applied for electroencephalographic recordings of 5 and 6 dpf zebrafish. Neuronal activity in the optic tectum of zebrafish was recorded by applying a protocol of 30 minutes of basal activity followed by 30 minutes of 40 mM PTZ treatment. 10 minutes in the middle of each period (basal activity and provoked seizures) was used to analyze the total number of negative spikes and duration of events for each genotype. Enzyme-linked immunosorbent assay (ELISA) was performed to quantify gamma-aminobutyric acid (GABA) and glutamate following a 30 minutes pre- and post-PTZ treatment at 5 mM. **C. (a)** Representative traces of electroencephalographic recordings in the optic tectum of WT and *kcnb1* knock-out larvae showing spontaneous and provoked epileptiform-like electrographic activity in the *kcnb1*[-/-] model characterized by **(b)** polyspike discharges, **(c)** 'ictal'-like activity and **(d)** large amplitude spikes. **D-E.** Quantification of electrophysiological recordings by analyzing **(D)** the total number of negative spikes and **(E)** the duration of events, over 10 minutes as described in **Figure 5A**. *kcnb1*[-/-] larvae showed significantly increased spontaneous and provoked neuronal activity, reflected by a significant elevation of the number of spikes, although the duration of events was similar to the WT condition in pre- and post-PTZ treatment. *kcnb*[+/-] larvae presented similar profile as the WT condition in term of number of spikes but showed a signifiant increase of event duration (see **Table 3**; n = 3-5 ZF/genotype; unpaired t-test; ns: non significant; **p<0,01; ***p<0,001; ****p<0,0001). **F-G.** Quantification of **(F)** GABA and **(G)** glutamate by ELISA assays in the head of 6 dpf larvae following the 30-minutes pre- and post-5 mM PTZ treatment. *kcnb1*[-/-] zebrafish present significantly elevated GABA levels observed in both baseline and post-PTZ conditions conversely to *kcnb1*[+/-] larvae, compared to the WT line. We observed a lack of significant changes in glutamate levels in both *kcnb1* mutant models in pre- and post-PTZ

conditions (N = 3-5 repeats; n = 50 heads/sample; unpaired t-test; ns: non significant;

*p<0,05; **p<0,01). For figures 5D to 5G, colored statistic indications correspond to mutant

conditions compared to the WT condition in pre- or post-PTZ treatment.

SUPPORTING INFORMATIONS

Supporting information 1. Brain anatomy and neuronal circuits are not disrupted in the

kcnb1 **loss-of-function zebrafish model. A.** Horizontal section of WT zebrafish expressing

cells (DAPI, blue) labelled with anti-kcnb1 (green), showing a large expression of the protein

in various regions of the central nervous system (CNS) at 6-day post-fertilization (dpf) (scale

bar 50 µm; magnification10x; n = 3-4 fish/section). **B.** Horizontal sections of WT zebrafish

expressing cells (DAPI, blue), kcnb1 (green) and Ankyrin G expressed at the axonal initial

segment of neurons (AnkG, purple, scale bar 30 µm; magnification 20x; n = 3-4 fish/section).

The image demonstrate the colocalization between kcnb1 and AnkG, represented in white and

marked by arrows, indicating the presence of kcnb1 in neurons. Colocalization was

determined using Z-stack projection on IMARIS V10.1.0 software (Oxford Instruments). In

figures, CNS regions are delimited by dotted lines. **C.** Transversal slices of 48 hpf and 4 dpf

larvae expressing cells (DAPI, blue) labelled with anti-HuC (red), a pan-neuronal marker,

showing no major difference of anatomical brain regions between mutants and wild-type fish

(scale bar: 50 µm; magnification 20x; n = 3-4 fish/section). **D.** Whole-mount images of

embryonic (48 hpf) and larvae (6 dpf) zebrafish immunostained with anti-3A10 marker to

identify Mauthner cells (dorsal view; 3D reconstruction; scale bar: 50 µm; magnification 20x;

n = 3-5 fish/section). **E.** Quantification of Mauthner cell body length (M-cell) of fish at 48 hpf

between the two extremities as represented with the arrow. The average value of the two M-

cell bodies of each fish was used for quantification. Both *kcnb1* knock-out zebrafish models

(*kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$) do not present difference in the length of M-cell body at 48 hpf as

compared to the WT condition (n = 3 - 5 ZF/genotype; one-Way ANOVA with Bonferroni

post-hoc   test;   ns: non-significant).   C: Cerebellum;   D: Diencephalon;   DAPI:

4,6□diamidino□2□phenylindole; E: Eyes; H: Hindbrain; kcnb1: Potassium voltage-gated

channel subfamily B member 1; M: Midbrain; OT: Optic tectum; SC: Spinal cord; T:

Telencephalon.

Supporting information 2: *kcnb1* mutant zebrafish models present a premotor

impairment and variability in the swimming behavior. A. Locomotor activity heatmap

reflecting the average number of movements per minute of embryos in their chorion between

24 hpf and 36 hpf. **B.** A significant decrease in general spontaneous movements (coiling and

twitching) of *kcnb1* mutant models (*kcnb1*$^{+/-}$ and *kcnb1*$^{-/-}$) was observed as compared to the

WT condition (*kcnb1*$^{+/+}$) during the whole first day of development (N = 3 repeats; n = 80-

104 ZF/genotype; one-way ANOVA with Bonferroni post-hoc test; ns: non-significant;

*p<0,05; **p<0,01; ****p<0,0001). **C.** Within the same genotype, the swimming behavior of

fish has been divided into two distinct phenotypes: severe (at least two swim circles, severe

phenotype, SP) and mild (other trajectories, mild phenotype, MP) during the 2nd dpf. The

subdivision of both swimming behavior (severe and mild phenotype) per genotype is

represented as traces of individual swimming episodes showing a huge variability of

trajectories (4 trajectories/phenotype). 30,68% of *Kcnb1*$^{+/-}$ fish and 52% of *kcnb1*$^{-/-}$ fish were

identified with a severe phenotype (see **Table 2**). **D-F.** The variability between both

swimming phenotypes was quantified following different parameters including **(D)** the

distance swam, **(E)** the velocity and **(F)** the time spent in motion. The results show similar

values between both mutant conditions presenting a mild phenotype as compared to the WT

condition. Nevertheless, fish with a severe phenotype present a significant increase of the

studied parameters for both genotypes (*kcnb1*[+/-] and *kcnb1*[-/-]). N = 3 repeats; n = 50-88

ZF/genotype; one-way ANOVA with Bonferroni post-hoc test; **p<0,01; ***p<0,001;

****p<0,0001.

Supporting information 3: Shift of c-Fos positive-neuron intensity distribution from a

global low neuronal activation to a higher neuronal activation in the telencephalon

region of *kcnb1*[-/-] fish at 6 dpf after chemically provoked-seizures. A. Whole larvae-fish at

6 dpf were fixed 30 minutes after a basal activity period and a provoked-seizures period due

to PTZ treatment at 5 mM followed by the labelling of c-Fos, an acute neuronal activation

marker. The number of c-Fos positive-neurons and the fluorescence intensity value of c-Fos

per neuron were quantified using the software IMARIS by analyzing the telencephalon, the

major region activated after the chemical treatment. The contour of the region of interest has

been designed manually to obtain the volume (μm^3) of the telencephalon, neurons were then

identified automatically within the region following identical parameters for spot detection

between conditions and visualized in 3 dimensions to avoid false negative spots. Values were

normalized according to the volume of the telencephalon (see **Figure 4E-4F**; n = 8-11

fish/condition). **B.** Distribution (in %) of c-Fos positive-neurons according to the fluorescence

intensity value of c-Fos in the telencephalon of fish at 6 dpf, reflecting the level of neuronal

activation that was divided in four equal shares (from 0 to 100%): low (0-25%), moderately

low (25-50%), moderately high (50-75%) and high (75-100%) neuronal activation. During the

basal activity, the distribution of activated neurons is similar between each conditions with a

majority of neurons presenting a low neuronal activation. However, a shift to a moderately

low and moderately high neuronal activation was observed for each condition after a PTZ

treatment, with a global higher activation in the *kcnb1*[-/-] condition (see **Figure 4G**). n = 8-11

fish/condition; T: Telencephalon.

FIGURE 1.

FIGURE 2.

FIGURE 3.

FIGURE 4.

FIGURE 5.