Copy number variation contributes to parallel local adaptation in an invasive plant

Jonathan Wilson[1], Vanessa C. Bieker[2], Lotte van Boheemen[1], Tim Connallon[1], Michael D. Martin[2], Paul Battlay[1, 3], Kathryn A. Hodgins[1, 3]

[1] School of Biological Sciences, Monash University, Melbourne, Victoria, Australia

[2] Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

[3] These authors contributed equally: Paul Battlay, Kathryn A. Hodgins

Authors declare no competing interest.

Abstract

Adaptation is a critical determinant of the diversification, persistence, and geographic range limits of species. Yet the genetic basis of adaptation is often unknown and potentially underpinned by a wide range of mutational types – from single nucleotide changes to large-scale alterations of chromosome structure. Copy number variation (CNV) is thought to be an important source of adaptive genetic variation, as indicated by decades of candidate gene studies that point to CNVs underlying rapid adaptation to strong selective pressures. Nevertheless, population-genomic studies of CNVs face unique logistical challenges not encountered by other forms of genetic variation. Consequently, few studies have systematically investigated the contributions of CNVs to adaptation at a genome-wide scale. We present a genome-wide analysis of CNV contributing to the adaptation of an invasive weed, *Ambrosia artemisiifolia* – the first such study in an invasive pest. CNVs show clear signatures of parallel local adaptation between North American (native) and European (invaded) ranges, implying widespread reuse of CNVs during adaptation to shared heterogeneous patterns of selection. We used a local principal component analysis to genotype CNV regions in whole-genome sequences of samples collected over the last two centuries. We identified 16 large CNV regions of up to 11.85 megabases in length, six of which show signals of rapid evolutionary change, with pronounced frequency shifts between historic and modern populations. Our results provide compelling genome-wide evidence that

29 copy number variation underlies rapid adaptation over contemporary timescales of natural

30 populations.

Significance Statement

Using a population-genomic approach, we identified copy number variants – stretches of DNA

that can be either present, absent, or in multiple copies – displaying parallel signatures of local

adaptation across the native and introduced ranges of the invasive weed *Ambrosia artemisiifolia*.

We further identified 16 large copy number variants, some associated with ecologically

important traits including sex allocation and height, that show strong signatures of selection over

space, along with dramatic temporal changes over the past several decades. These results

highlight the importance of an often-overlooked form of genomic variation in both local

adaptation and rapid adaptation of invasive species.

Introduction

Understanding how populations adapt and persist in the face of rapid environmental change is one of the most pressing challenges of our time. Fundamental to this goal is determining the genetic basis of adaptive evolution. But despite considerable empirical and theoretical work in this area, many questions remain unresolved. For example, does adaptation typically rely on new and beneficial mutations or on standing genetic variation? Does adaptation generally result in the removal or maintenance of genetic variation affecting fitness? Do mutations contributing to adaptation have uniformly small phenotypic effects, or are large-effect mutations important as well? Do populations exposed to similar environments evolve using the same or different genetic variants?

The evolution of quantitative traits was traditionally thought to almost exclusively depend on evolutionary changes at many polymorphic loci with individually small phenotypic effects (1, 2). However, comparatively recent theoretical and empirical studies demonstrate that large-effect variants can also play important roles in adaptation (3–5). Large-effect mutations are particularly likely to contribute to the initial stages of a population's evolutionary response to a sudden shift in the environment (6), and to facilitate stable adaptive genetic differentiation among populations connected by migration. Such large-effect variants promote local adaptation by resisting the swamping effects of gene flow (7), including cases in which the alleles carry substantial pleiotropic costs (8).

Genomic structural variants, which include inversions, translocations, duplications and deletions, are predicted to have both large phenotypic effects and strong potential to contribute to adaptation (9, 10). Chromosomal inversions have a long history of evolutionary study, initially facilitated by classical cytogenetics (e.g., polytene chromosome studies [(11, 12)]), and more recently through advances in genome sequencing and analysis, which have produced compelling new evidence that inversions often underpin adaptation to environmental change (13–15). These findings have reinvigorated interest in the role of inversions in adaptation, yet other types of structural variation have not garnered the same level of attention.

Several case studies show that copy number variants (CNVs) – structural changes that include
gene duplications, deletions and variation in transposable element abundance – have facilitated
adaptation in well-characterized systems such as *Drosophila melanogaster* (16), *Anopheles*
gambiae (17), and *Arabidopsis thaliana* (18). Prominent examples include the repeated
contribution of CNVs to pesticide resistance (19–22), evident in the parallel evolution of CNVs
in the agricultural weed *Amaranthus tuberculatus* as a response to glyphosate exposure (23), and
amplification of cytochrome P450 family genes in Antarctic killifish and fall armyworm
populations exposed to toxins (24, 25). These studies demonstrate the immense adaptive
potential of CNVs, yet most are candidate-driven analyses that cannot resolve the broader
contributions of CNVs to adaptation. Few studies have systematically characterized the genome-
wide contributions of CNVs to adaptive divergence across a range of environmental conditions
and stresses (13, 26, 27).
83
Invasive species have several unique features that make them powerful systems for studying
adaptation in nature and uncovering its genetic basis. First, because recently introduced
populations are likely to be initially maladapted to local conditions in the new range, there is
strong scope for rapid adaptive evolution that is observable within decades (28, 29). Second,
some plant invasions have extensive documentation in geo-referenced herbarium collections that
can be phenotyped and sequenced to identify and track evolutionary changes over time – an
approach rarely possible in natural populations (15, 30, 31). Third, invasive species typically
occupy climatically diverse native and invasive ranges, promoting adaptive evolution to local
environmental conditions (28, 29, 32). In particular, those with broad ranges further enable tests
of the predictability of evolution, since local adaptation across native and invasive ranges may
stem from parallel or unique genetic solutions to similar environmental challenges (15, 32, 33).
Moreover, they are well suited for evaluating contributions of CNVs, which have been predicted
to be important in invasive species adaptation (34, 35). Yet to our knowledge, this hypothesis has
never been tested at a genome-wide scale.
98
Over the last 200 years, the North American native plant *Ambrosia artemisiifolia* (common
ragweed) has become a widespread pest on all continents except Antarctica (36). This wind-
pollinated, outcrossing species produces highly allergenic pollen that accounts for ~50% of hay-

fever cases in Europe (37). It is also an agricultural pest (38), with glyphosate resistance – a

phenotype associated with CNVs in other species (19)– reported in some *A. artemisiifolia*

populations (39). Furthermore, climate change is predicted to exacerbate this weed's impacts,

with increased pollen production due to an elongated flowering season (40), as well as reducing

the future geographic overlap with key biocontrol agents (41). Previous studies show that *A.*

artemisiifolia has established strong signals of local adaptation to climate across its native range,

and in introduced ranges in Europe, Asia, and Australia, with rapid local adaptation following

each introduction (15, 33, 41–45). We have previously demonstrated a significant contribution of

both SNPs and large-effect structural variants – in the form of chromosomal inversions – to

climate adaptation in Europe and North America (15). This raises the question of whether other

types of genome structural variants (i.e., CNVs) have similarly contributed to adaptive

divergence following *A. artemisiifolia*'s range expansion.

In this study, we leveraged a large temporally and spatially resolved dataset to investigate the

genome-wide contributions of CNVs to local adaptation in *A. artemisiifolia*. Over 600 whole-

genome sequences from individuals collected across the native North American range and the

introduced European range, including herbarium samples dating back to 1830, provided a unique

opportunity to detect signals of adaptation over space and time. We first investigated whether

putative CNVs display signatures of divergent selection in both Europe and North America. By

comparing the selection signatures of putative CNVs in each range, we then assessed the degree

to which these shared variants evolve in parallel between them. Second, using a subset of 121

phenotyped individuals, we tested for associations between putative CNVs and ecologically

important traits, such as flowering time and size. The typically low-coverage and error-prone

nature of herbarium sequences renders many existing CNV detection methods unsuitable for

these data. As such, we developed a novel approach that combines read-depth and local principal

component analysis (PCA) methods to genotype large CNV regions in both modern and low-

coverage historic samples, enabling us to identify CNVs, estimate temporal changes in their

frequencies, and thus directly track CNV evolution in these populations.

Results

CNV identification

To identify CNVs, we calculated read-depth in non-overlapping 10 kbp windows (normalized by the average coverage of each individual sample) across 311 modern re-sequenced genomes spanning both the North American and European ranges of *A. artemisiifolia*. We defined a window as a CNV if at least 5% of samples had average standardized read-depths either greater, or less, than two standard deviations from the window mean across samples. Out of the total 105,175 genomic windows, this resulted in 17,855 candidate CNVs retained for subsequent analyses.

CNV selection analysis

Local adaptation to spatially heterogeneous environments is expected to leave a signal of extreme allele frequency differentiation among populations (46). As our measures of read-depth for identifying candidate CNV windows were continuous, we used a Q_{ST}-F_{ST} outlier test to identify windows with population differentiation in excess of neutral expectations. We first calculated an F_{ST} distribution of neutral SNPs (Figure 1B, D) using the method described by Weir & Cockerham (47). Under neutrality, F_{ST} and Q_{ST} values should have similar distributions, whereas an enrichment of Q_{ST} values within the upper tail of this null F_{ST} distribution provides evidence for local adaptation, with Q_{ST} outliers representing local adaptation candidates (48).

In North America, 1,382 CNV windows exhibited Q_{ST} values at or above the top 1% threshold of neutral F_{ST} values: a 7.7-fold enrichment relative to the neutral expectation that 1% of Q_{ST} windows will fall within this tail (p < 2.2e-16; binomial test; Figure 1A, B). In Europe, 339 CNV windows exhibited Q_{ST} values exceeding the top 1% of the F_{ST} distribution: a 1.9-fold enrichment (Figure 1C, D; p < 2.2e-16; binomial test). Using an equation adapted from the McDonald-Kreitman test (49), this excess of outliers is consistent with a true positive rate of 87% for the North American CNV candidate windows, and 47% true positives for European outlier windows (see methods). Of the CNV windows displaying differentiation, 111 were outliers in both ranges (32% of European outliers; p=1.17e-40, hypergeometric test; Figure 2A), a highly significant excess indicative of parallelism in the same CNVs subject to divergent selection in both ranges. In contrast, there was no overlap between the top 1% of neutral SNP

F_{ST} values for the two ranges, suggesting that neutral processes cannot explain the parallelism

observed in CNVs.

Variation in recombination rate across the genome may interfere with the identification of

signatures of selection (50). To account for potential effects of local recombination rate on

patterns of CNV differentiation, we separated Q_{ST} windows into three recombination rate bins

based on the genetic map described in Prapas et al. (51): low (<0.5 cM/Mbp), medium (0.5-2

cM/Mbp) and high (>2 cM/Mbp). When Q_{ST}-F_{ST} analyses were repeated within each

recombination rate bin separately, 98.9% and 98.3% of the original Q_{ST} outliers remained

significant in North America and Europe, respectively. This demonstrates the minimal impact of

recombination rate on the divergent patterns of read depth within CNV windows in this dataset

(Supplementary Figure 2). We also investigated the possibility that non-independence between

10 kbp windows drives the observed patterns of divergence and repeatability. To do so, we

merged windows with correlated variation in read depth ($R^2 > 0.6$) within 1 Mbp of one another.

After merging windows, the number of candidate CNVs was reduced from 17,855 to 11,877,

with the largest window measuring 11.85 Mbp on chromosome 4. We then repeated the Q_{ST}-F_{ST}

analysis on these merged windows. Q_{ST} values remained elevated relative to neutral F_{ST}

distributions in both North America and Europe (6-fold and 1.3-fold respectively), with 41

outlier windows shared between ranges: far more than expected by chance (hypergeometric test

p=2.736e-16).

To identify candidate CNVs associated with climate, we estimated correlations between each

candidate CNV window and the six bioclimatic variables (Supplementary Figure 5; 6) that were

selected after filtering highly correlated variables from the original 19 WorldClim variables (52).

Of the 1,382 significant Q_{ST} windows in North America, 315 (22.7%) were associated with at

least one of the six climate variables, whereas only 12 of 339 significant windows in Europe

(3%) correlated with climate variables (Supplementary Figures 5; 6), suggesting the primary

selective forces driving differentiation of CNVs in Europe are likely not climate-related.

To determine the putative biological functions of the adaptation candidates, we used gene

ontology (GO) enrichment analyses of annotated genes residing within the outlier Q_{ST} windows.

Candidate CNVs in North America are enriched for biotic and abiotic stress response genes, with

significant GO terms including "systemic acquired resistance," "response to oomycetes", and

"response to freezing" (Supplementary Figure 3A; Supplementary Table 1). In Europe,

significant GO enrichments include the hormonal stress response pathways "response to abscisic

acid" and "response to jasmonic acid" (Supplementary Figure 3B; Supplementary Table 2).

Overlapping Q_{ST} candidates between the ranges exhibited GO term-enrichment for the defense

response terms "defense response to virus" and "response to abscisic acid" (Figure 2B;

Supplementary Table 3).

CNV-trait associations

We tested for relationships between genome-wide CNVs and 29 ecologically important traits

phenotyped in 121 of our samples, each reared in a common garden experiment (phenotype data

were previously reported in van Boheemen, Atwater & Hodgins [(43)]). Eighteen traits

(Supplementary Table 5) were significantly associated (using a Bonferroni-corrected 0.05

threshold) with normalized read-depth in at least one of the 17,855 candidate CNV windows

(Supplementary Figure 4). Of these trait-associated windows, 17 and 4 overlapped with Q_{ST}

outlier windows in North America and Europe, respectively. With a more relaxed significance

threshold of FDR = 0.05 using the Benjamini-Hochberg method, these overlaps were increased

to 76 in North America and 10 in Europe. Of particular interest, two nearby windows on

chromosome 14 (h1s14:17180001-17190000, h1s14:18470001-18480000) were associated with

flowering onset, dichogamy, and sex allocation (defined as female reproductive biomass/male

reproductive biomass; Supplementary Figure 4). These traits display strong latitudinal clines in

A. artemisiifolia, with overall earlier flowering, much earlier male flowering compared to female

flowering, and female-biased sex allocation occurring at high absolute latitudes (43). These two

windows flank an annotated gene, AGL-104, that is linked to pollen production in Arabidopsis

(53). Moreover, one of these windows (h1s14:18470001-18480000) is a Q_{ST} outlier in North

America and Europe, suggestive of parallel divergent selection.

Large CNV region identification (CNVr)

The continuous measures of read-depth in 10 kbp windows, while reliable in modern samples,

were inaccurate when using low-coverage historic data. Since we had previously obtained

accurate measures of genotypes for large inversions in these historic samples (15), we

implemented a similar approach to identify large CNV regions (CNVr) in order to perform

temporal comparisons between the historic and contemporary samples. Furthermore, we would

expect many CNVs to be larger than 10 kbp. We therefore used the same linkage disequilibrium-

based approach as stated above to identify and merge adjacent windows which appeared to be

components of a larger CNV. To corroborate the presence of large segregating CNVs within our

dataset, we performed a local PCA of genotype likelihoods in 100 kbp windows across the

genome. We determined potential segregating structural variants as regions with at least three

adjacent windows that were outliers for distortions in local population structure relative to the

rest of that chromosome. This resulted in a minimal size cutoff of 300 kbp for CNVrs. As such,

we defined CNV regions as those in which merged read-depth windows (greater than 300 kbp)

overlapped with at least three adjacent windows exhibiting variation in local population

structure. This approach identified 52 candidate CNVrs.

We then genotyped individuals for CNVrs in our population-genomic data, including low-

coverage historic samples. To do this, we used a combination of normalized read depth across

the genomic location of each candidate CNVr alongside a PCA calculated within that same

region to cluster samples into genotypes differing in both read-depth and PC1 (Figure 3). With

this approach, we were able to identify distinct clusters corresponding to genotypes in both

modern and historic samples for 16 out of 52 CNVrs. These 16 CNVrs, which ranged in size

from 0.3 - 11.85 Mbp, accounted for 8.1% of the 17,885 10 kbp windows identified in the

contemporary samples, including 22.6% of outlier Q_{ST} windows in North America and 7.2% of

outlier Q_{ST} windows in Europe. Fifteen of these CNVrs exist as heterozygotes within the diploid

reference genome, with haplotype 1 containing presence variants, meaning they can be

corroborated with alignments between each haplotype of the diploid assembly (Supplementary

Figure 9). The high heterozygosity of CNVrs in the reference was likely due to our genotyping

method favoring loci with absence alleles that are common in our samples, yet the presence

alleles needed to be found in the reference haplotype in order for the CNV to be identified. This

is consistent with the low frequencies (mean = 0.196; range: 0.027 −0.479) of all CNVr presence

variants (Supplementary Table 6). GO analysis of annotated genes indicates that the 16 CNVrs

were enriched for several biological processes, including "methylglyoxal catabolism",

"peroxisome fission", "pollen tube adhesion", and "glyphosate metabolism" (Supplementary

Table 4).

The largest CNVr that we identified was cnv-chr4a, which we estimated to be 11.85 Mbp in

length. Closer inspection of this region within the diploid reference reveals that large regions on

haplotype 1 are absent on haplotype 2, but are interspersed by three smaller inversions (Figure 3;

Supplementary Figure 8; 9). Analysis of coverage depth across chromosome 4 of three closely

related Ambrosia species sequences mapped to the *A. artemisiifolia* reference revealed the

absence haplotype as the likely ancestral state (Supplementary Figure 10). The derived insertion

variant contains an excess of transposable elements (TEs) relative to the remainder of

chromosome 4 (87.21% versus 70.48%). TE family Ty1/Copia accounts for 32.45% of the TEs

within cnv-chr4a, compared to just 14.87% throughout the rest of this chromosome. Repetitive

elements display greater density towards the beginning of this region (Supplementary Figure 11),

where gene density is very low. The region toward the end of the cnv-chr4a, which exhibits

greater gene density, shows strong synteny with chromosome 2 and aligns with inversions

present in the reference alignment (Supplementary Figure 8; 11). Most of chromosome 4

displays synteny with chromosome 2, suggesting they are homoeologous chromosomes. The

largest gap in syntenic blocks corresponds to the gene-depleted and TY1/Copia-enriched region

of cnv-chr4a (Supplementary Figure 11). It is therefore likely that this complex structural variant

consists of a series of inversions which have subsequently been separated by a large TE

expansion. Recombination is likely strongly suppressed within this structural variant, as the

coverage windows exhibit strong linkage disequilibrium across the region. Candidate CNV

windows within cnv-chr4a are Q_{ST} outliers (Figure 1A; C) and associated with the bioclimatic

variable mean diurnal range (Supplementary Figure 5), consistent with the CNVr contributing to

local adaptation.

Another noteworthy CNVr, cnv-chr8a, contains an ortholog of the *Arabidopsis thaliana* EPSPS

locus. This CNVr lies within a large inversion, hb-chr8, previously described in Battlay et al.

(15). While the frequency of cnv-chr8a is not strongly correlated with the frequency of this

inversion (R^2=0.006), cnv-chr8a presence alleles occur exclusively on the common, and

presumably ancestral, orientation of the inverted region.

Spatio-temporal CNVr modeling

We used whole genome sequences derived from >284 historical herbarium samples (dating back to 1830) and generalized linear models (GLMs) to uncover how the 16 CNVr alleles may have changed in frequency over both space and time. These GLMs predicted genotype as a function of range (native or introduced), latitude and year of collection, and each model was reduced to remove any non-significant interactions between these variables. To correct for population structure, we also included the first principal component of genetic variation (calculated from 10,000 neutral SNPs) as a covariate in each model. Twelve of the 16 CNVrs displayed at least one significant predictor variable (time, range, latitude or interactions). Nine CNVrs exhibited significant associations with latitude (cnv-chr4a, cnv-chr4b, cnv-chr5a, cnv-chr9a, cnv-chr14a, cnv-chr17b, cnv-chr17c and cnv-chr18a) (Supplementary Table 7; Supplementary Figure 12). Models of three CNVrs (cnv-chr10a, cnv-chr14a and cnv-chr17a) contained significant three-way interactions (Figure 4; Supplementary Table 7; 8; Supplementary Figure 12). Of note, both cnv-chr14a and cnv-chr17a displayed clinal patterns in North America regardless of year, whereas this same latitudinal pattern was present only in modern European samples — strong evidence of clinal reformation following an initial period of maladaptation in the introduced range (Figure 4; Supplementary Table 7; 8). The large cnv-chr4a insertion displays latitudinal clines in modern populations across both ranges, with the insertion at higher frequencies at lower latitudes. In the native range, this appears to be driven by increasing frequencies over time in more central and southern populations of North America (Supplementary Table 8). Correspondingly, this CNV overlaps with a previous SNP-based result consistent with a selective sweep in the St. Louis population (15).

To further analyze the potential adaptive significance of these CNVrs, we assessed associations between each variant and the same 29 phenotypes analyzed above (43). The cnv-chr10a variant displayed Bonferroni-significant associations with total biomass, root weight and total number of male inflorescences, whilst cnv-chr14a was significantly associated with sex allocation (Supplementary Table 9).

Discussion

CNVs are increasingly recognized as important in local adaptation (13, 24, 26, 27, 54). However, previous empirical evidence is predominantly limited to examples of pesticide resistance (18, 25) and candidate gene studies (55, 56). Genome-wide analyses of CNVs at the population scale are rare beyond model organisms (54). While many methods exist for identifying CNVs from resequenced genomes (57, 58), these often rely on long reads, or short reads with higher coverage than available for many population genomic datasets, including our historic specimens. We therefore combined local read-depth, linkage disequilibrium, and deviations in population structure along the chromosome, to identify CNVs in modest-coverage samples collected over the past 190 years.

We implemented a genome-wide discovery approach to identify CNVs and examined their potential contributions to local adaptation across the native and an invaded range of *A. artemisiifolia*. CNV windows were enriched for signatures of local adaptation, which occurred disproportionately in parallel between native and invasive ranges. As this signal was not replicated in putatively neutral SNP loci, this implies CNV-driven local adaptation across Europe takes place via standing variation inherited from North America. Furthermore, large CNV regions identified in modern and historical samples, such as cnv-chr14a and cnv-chr17a (Figure 4), show evidence of rapid local adaptation across the short timescale (ca. 150 years) of *A. artemisiifolia*'s invasion in Europe.

Ambrosia artemisiifolia CNV windows exhibited extensive signals of local adaptation, including elevated geographic divergence in fold coverage, relative to neutral expectations, in both North America and Europe (Figure 1). These windows contain an over-representation of genes involved in abiotic and biotic stress responses in North America (Supplementary Figure 3A, Supplementary Table 4) and pathogen defense in Europe (Supplementary Figure 3B, Supplementary Table 3) – consistent with SNP-based F_{ST} outlier windows in Europe being enriched for defense related functions (30). The overall weaker patterns of CNV differentiation in Europe relative to North America are also consistent with previous SNP-based analyses showing fewer differentiated outlier SNP windows in Europe (15). Some CNVs were associated with traits important to local adaptation, including a CNV on chromosome 16 associated with

mature plant height (Supplementary Figure 4), which also overlaps an annotated NB-ARC

domain. NB-ARC domains, occurring in most plant resistance (R) genes, are involved in

nucleotide binding and recognition (59), and duplications of such genes underlie the evolution of

resistance to pathogens (60, 61). CNVs also appear to influence phenological traits. For example,

the CNV windows on chromosome 14 are associated with flowering time (Supplementary Figure

4). Overall, our data point to important roles of CNVs in the local adaptation of *A. artemisiifolia*,

which aligns with evidence from other species that structural variants widely contribute to

adaptation (62–64).

One third of adaptive CNV windows in Europe were also candidates for local adaptation across

North America. Previous work in *A. artemisiifolia* has revealed similar patterns of repeatability,

or parallelism, with respect to SNPs (15, 33), inversions (15), and genes affecting locally adapted

traits like flowering time and sex allocation (43). Invasive species are expected to evolve in

parallel when responding to analogous selection pressures, as observed in our system and in

others, such as *Drosophila suzukii* and European starling (65, 66). Such parallelism is promoted

when standing genetic variation involved in local adaptation in the native range is recruited as a

source of adaptive variation within the invasive range (67, 68). Multiple introductions from

several genetically diverse source populations from North America to Europe presumably

facilitated repeatability by ensuring that most of the important standing variants successfully

made the journey to the new range (30, 33). That all CNVrs were present in historical European

populations indicates that they were introduced into Europe during the early stages of invasion

(Supplementary Table 6).

The 16 large CNV regions that we detected using a combination of read-depth, linkage

disequilibrium and local population structure analyses (Figure 3A, Figure 3B) contained 22.6%

of Q_{ST} outlier windows from North America and 7.2% of the outlier windows from Europe. Yet

these CNVrs comprised only 0.23% of the genome, demonstrating their disproportionate

contribution to these signals of local adaptation. Remarkably, 15 of the CNVrs co-localize with

segregating presence/absence variants in the highly heterozygous diploid reference, supporting

our detection method. Closer analysis of the largest CNVr (cnv-chr4a) within the reference

reveals that our detection method may lack sensitivity in fully revealing structural complexity

within CNVrs (Figure 3). Multiple abutting inversions and CNVs within this region appear to

segregate together as a single, complex structural variant. Nevertheless, the regions we identified

demonstrate the general adaptive potential of structural variants, in which CNVs are predominant

features. Since our method of detection for CNVrs was biased towards identifying large CNVrs

with presence variants on haplotype 1 of the reference, and our genotyping method was biased

towards identifying loci with common absence alleles, our results represent a lower bound on the

prevalence and adaptive significance of CNVrs in *A. artemisiifolia*. Investigations using

pangenomics (69) to elucidate a more complete picture of the contribution of CNVs to adaptation

are therefore warranted.

Our study goes beyond the traditional population-genomic approach of detecting signals of local

adaptation using contemporary samples alone. Our use of historical sequence data also allowed

us to track temporal change in CNV frequencies across nearly two centuries – a period that spans

the establishment and spread of *A. artemisiifolia* within Europe (36, 70), and significant

environmental upheaval in both ranges, owing to industrialization, agriculture, and climate

change (71). CNVs are rarely genotyped in historic genomes because sequence quality is poor

(72). However, by focussing only on large CNVs identified using modern data we were able to

confidently assign genotypes in historic samples. This use of modern data to validate historic

sequence variant calls is common in temporal genomic studies (30, 73). Eight CNVrs display

clear frequency shifts over time, consistent with rapid adaptation over its recent evolutionary

history. For example, while cnv-chr14a and cnv-chr17a both exhibit a consistent latitudinal cline

in historical and modern North American populations, these clines are only evident in modern

European populations (Figure 4A; Supplementary Table 7; 8), which is consistent with the rapid

cline reformation in Europe following an initial period of post-introduction maladaptation. The

cnv-chr14a CNVr is associated with sex allocation (Supplementary Table 9), and candidate CNV

windows within the region exhibit associations with flowering time, sex allocation and

dichogamy (Supplementary Figure 4), traits which show parallel clines in Europe and North

America (43). Furthermore, cnv-chr14a lies within 20 kbp of the AGL-104 gene, which is

involved in pollen development in Arabidopsis (53). Flowering time is a complex trait that is

affected by diverse forms of genomic variation (74) – our results indicate that SNPs, inversions

(15), and CNVs each play important roles in the rapid adaptation of this important trait in *A.*
artemisiifolia populations.

Many well-characterized CNVs in other species are associated with the evolution of pesticide
resistance (19, 20, 25). We identified a CNVr (cnv-chr8a) potentially involved in herbicide
resistance. An ortholog of the *Arabidopsis thaliana* EPSPS locus, the molecular target of
glyphosate herbicides, lies within the cnv-chr8a region. CNVs confer resistance to glyphosate in
numerous other weed species, where the increased gene expression caused by EPSPS gene
amplification ameliorates the herbicide's toxic effects (19, 55). Glyphosate resistance has been
documented in some *A. artemisiifolia* populations (39), and while we do not know which
populations in our study might be glyphosate resistant, this CNVr is a strong candidate for future
study.

Previous population-genomic analyses of *A. artemisiifolia* provide strong evidence that SNPs
and putative chromosomal inversions contribute to local adaptation; here we provide evidence of
a similar role for CNVs. Although CNVs are known to have large effects on traits (17), we
cannot be sure that the variants we have identified are the direct targets of selection – they may
instead be in linkage disequilibrium with other variants that are the actual targets. Assessing
relationships between CNVs and nearby SNPs is fraught, because CNVs disrupt SNP calling
(75). Two CNVrs overlap inversions identified in Battlay et al. (15), but are not in strong LD
with the inversion genotypes (R^2 = 0.006-0.01). However, smaller CNVs may exhibit stronger
associations with other SVs as part of coadapted gene complexes (76) or neutral hitchhikers.
Large insertion-deletion variants result in may hemizygous regions of reduced recombination
which may collect and bind together multiple variants (77). Functional assays such as RNA-seq
experiments are required to understand the mechanistic effect of these CNVs on traits and fitness
(78). Further efforts to determine the functional effects of CNVs, alongside greater sample sizes,
would also help uncover the likely epistatic interactions between CNVs and other adaptive
variants. The potential existence and nature of these interactions are pertinent questions in
evolutionary biology lacking empirical investigations on genome-wide scales (79). Future work
should also consider the role of other forms of genomic variation, for example transposable
element abundance and genome size which have been linked with local adaptation and

aggressive range expansion (74), alongside investigating the roles of CNVs in biotic interactions,

such as pathogen response.

Our study highlights the importance of copy number variation in the evolution of a widely

distributed and rapidly adapting invasive weed. While CNVs have previously been implicated in

adaptation in response to specific selection pressures in other species (23, 24), our genome-wide

discovery approach was able to identify candidate genomic regions that are more broadly

representative of the contribution of CNVs to local adaptation. We have linked several of these

candidates with traits ranging from flowering time to pathogen resistance. Along with previous

studies showing that SNPs and chromosomal inversions underlie local adaptation during *A.*

artemissifolia's expansion across vast environmental gradients, these new findings make it clear

that CNVs account for a significant and previously unrecognized component of this plant's past

success and are consequential for its invasive capacity wherever it may be introduced in the

future.

Methods

Samples and alignments

Analyses were conducted on 613 whole-genome *Ambrosia artemisiifolia* sequences described in

Bieker et al. (30) and a chromosome-level, phased, diploid *Ambrosia artemisiifolia* genome

assembly (15). Alignments and SNP calls against the primary haplotype of the diploid reference

(haplotype 1) were generated by Battlay et al. (15), using the Paleomix v1.2.13.4 (80) pipeline

and GATK UnifiedGenotyper v3.7 (81). Modern and historic samples were sequenced from

across the species' native North American (155 modern and 92 historic samples) and introduced

European (156 modern and 192 historic samples) ranges. Modern samples were collected

between 2014 and 2018 and sequenced to a mean coverage of 2.9x. Historic samples were

sequenced from herbarium samples collected between 1830 and 1973 with a median collection

date of 1905 (Supplementary Figure 1), and sequenced to a mean coverage of 1.4x. Present-day

population samples, whose geographic coordinates were recorded during sampling, included

between $n = 1$ to $n = 10$ individuals, however populations where $n = 1$ were removed from

analyses requiring population level information. Historic individuals were grouped into

populations according to their proximity (15, 30). Cases where only one sample was obtained

from a geographic location were excluded from analyses in which population information was a

requirement. Additionally, we aligned sequences of three other Ambrosia species (30) to the

primary haplotype of the diploid reference using the Paleomix pipeline, as described in Battlay et

al. (15).

Depth of coverage analysis

In order to identify copy number variation within our resequenced common ragweed individuals,

we analyzed depth of coverage in non-overlapping 10 kbp windows using Samtools v1.9 depth

(82) on alignment bam files. In the initial analyses of modern samples we only used reads with

mapping quality > 30 (-Q 30). Subsequent read-depth analyses of historic samples used reads

with mapping quality > 5 (-Q 5) in order to accommodate their poorer mapping quality. Each

window was then normalized by dividing window depth by the genome-wide coverage for the

sample. To apply a population frequency-based filter to this dataset, we kept only windows

which had at least 5% of samples greater than or less than 2 standard deviations from the

population mean. This filtering procedure resulted in 17,855 of 105,175 windows (5.8%) being

classified as copy number variants (CNVs).

Q_{ST}-F_{ST} analysis

Genomic loci that have responded to spatially heterogeneous selection are expected to show

elevated differentiation among populations, relative to neutrally evolving loci. SNPs associated

with local adaptation can be detected as outliers of genome-wide scans of F_{ST} or similar statistics

(47, 83, 84). However, unlike SNP data, candidate CNVs have been identified by depth of

coverage represented on a continuous scale. As such, coverage can be viewed as a phenotypic

measurement that is analogous to measurements of continuous traits. Tests for trait responses to

divergent selection are often inferred using a Q_{ST}-F_{ST} analysis (85, 86). Theory predicts that Q_{ST}

values for neutrally evolving traits should follow the same distribution as F_{ST} for neutrally

evolving loci (48). We used coverage data to measure Q_{ST} values for each window (analyses

were conducted separately for the European and North American ranges), adapting the

relationship between Q_{ST} and population variation from Whitlock (48):

$$Q_{ST} = \frac{VA,among}{VA,among + 2VA,within} \tag{1}$$

where VA,among represents the among-population variation in coverage for a given window,

and VA,within represents within-population coverage variation.

We performed linear mixed models, using the lme4 package in R (87) on populations from each

range to extract variance components attributed to within- and among-population variation for

each coverage window. Unlike other analyses performed in this study, we used unnormalised

coverage for each window in the model, and accounted for variation in individual sample depth

by including individual median coverage as a covariate in the model, and population was

included as a random effect. The variance among populations (VA,among) was extracted as the

variance component attributed to the population, and within-population variance (VA,within)

corresponded to the model's residual variance component.

To identify Q_{ST} values with divergence in excess of neutral expectations, the distribution of Q_{ST}

values in each range was compared to the distribution of neutral F_{ST} values from the same

populations. F_{ST} distributions were calculated in VCFtools (88) using 10,000 putatively neutral

and independently segregating LD-pruned SNPs, randomly sampled from outside both genic

regions and known structural variants (15). Outliers were classified as windows with Q_{ST} values

greater than 99% of neutral F_{ST} values. Under neutrality, 1% of Q_{ST} values are expected to fall

above this 99% threshold, and we therefore tested whether there was a significant excess of

windows above this null expectation. To identify the rate of true positives, we used the following

calculation

$$\frac{m}{M} = 1 - \frac{n * 0.01}{M} \tag{2}$$

where n represents the total number of windows analysed, M represents the total number of

outliers, m represents the number of true positives, and therefore M - m represents the number of

false positives. Analyses were performed using modern data from North America and European

ranges separately. We used a hypergeometric test to identify significant overlaps in both Q_{ST} and

F_{ST} loci between both ranges.

Environmental-CNV Associations

Correlations between genetic variation and environment provide further evidence for local

adaptation on the genomic level. CNV-environment associations were identified by measuring

Spearman's correlation coefficients (Rho) between normalized coverage of each 10 kbp window

and climate variables. Climatic variables were extracted from WorldClim (52) for each of the

geographic coordinates of our sample of individuals. We excluded highly correlated variables

($R^2 > 0.7$) from the analysis, resulting in six variables: BIO1: annual mean temperature, BIO2:

mean diurnal range, BIO8: mean temperature of wettest quarter, BIO9: mean temperature of

driest quarter, BIO12: annual precipitation, and BIO15: precipitation seasonality. Outliers were

identified as the top 1% of the empirical p-value of the Rho distribution. Coverage windows that

were in the 1% tails of the CNV-environment distribution and outliers in our Q_{ST} analysis were

considered CNV-climate adaptation candidates.

Identifying large CNV regions

Since larger CNVs will affect coverage in multiple adjacent windows, we used a linkage

disequilibrium-based approach to merge nearby windows. We merged windows within 1 Mbp

(and all windows within this region) that had sample depths that were correlated at $R^2 > 0.6$. By

generating heatmaps of LD between CNV windows, we visualized the presence of correlated

read-depth, indicative of larger CNVs (Supplementary Figure 7). This resulted in 11,877

windows ranging in size from 10 kbp to 11.85 Mbp.

Local principal component analysis has previously been used to detect population-genomic

signatures of chromosomal inversions (15, 62, 89). Here we employed this method to identify

distortions in local relatedness due to CNVs using two modifications. Due to low-coverage and

bias in historic samples, we performed these steps initially on modern samples only, including

historic samples for genotyping later (see below). Firstly, we calculated local covariance

matrices for each window using ANGSD (90) and PCAngsd (91). Secondly, we did not include a

filter for missing data in our ANGSD command, which meant that missingness would cause

distortion in the local PCAs. The analysis was performed in non-overlapping windows of 100

kbp, and multidimensional scaling axes 1-4 (calculated from local PCAs across each

chromosome) were examined for blocks of outliers indicative of large structural variants. While

individual MDS outlier windows are not necessarily due to structural variations, SVs are the

most likely cause of signals that are consistent across adjacent windows. Therefore, we retained

candidates that included at least three adjacent windows that were outliers, so that the lower limit

of CNVr size was 300 kbp. As such, we identified candidate CNVr as those where MDS

candidates overlapped the merged read-depth-based CNV windows which were also greater than

300 kbp in length. These 52 candidate CNVrs corresponded to 68% of the merged CNV

windows greater than 300 kbp in length.

CNV region genotyping

We attempted to genotype modern and historical samples, separately, for each of the 52

candidate CNVr. To do so, we performed another local PCA across the length of each candidate

CNVr using PCAngsd (91). We then compared PC1 of the candidate CNVr against the average

normalized read-depth across that CNVr to identify whether individuals clustered by read-depth

and PC1. To genotype these samples according to these clusters, we used k-means clustering

from the ClusterR package in R (92), followed by manual annotation of genotypes. Samples

appeared to either cluster into groups of two or three genotypes. CNVr candidates displaying two

clear genotypes were likely to have one rare homozygote, or the heterozygote and one

homozygote class were indistinguishable. As such, we assigned k-values of either 2 or 3

depending on whether visual inspection indicated the presence of two or three segregating

genotypes. To test for association between CNVrs overlapping or neighboring chromosomal

inversions, we used the cor function in R (R team) to calculate correlations between CNVr and

the genotypes of overlapping inversions that were previously identified by Battlay et al. (15). To

visualize CNVrs that were heterozygous between haplotypes of the diploid reference genome,

we aligned both reference haplotypes using minimap2 v2.1.8 (-k19 -w19 -m200) (93) and

generated dotplots of the alignments. To call segregating SVs in the diploid reference, we

aligned both haplotypes using nucmer (-- maxmatch -c 100 -b 500 -l 50) within the mummer

v3.23 software package (94). We then used SyRI (95) to identify and plotsr (-s 300000) (96) to

visualize structural variants greater than 300 kbp in length. This method of identifying large

segregating CNVs is constrained in that it can only identify CNVrs that contain a presence

variant on haplotype 1 of the reference. Since haplotype 1 is larger than haplotype 2, it is more

likely to contain more presence variants. Secondly, to create a strong signature of divergence

within the local PCA, a large number of individuals must contain the absence variant, meaning

that presence variants will also be rare. Furthermore, closer inspection of segregating CNVrs

within the reference indicated that some CNVrs may consist of complex SVs, containing

inversions and translocations, which our identification method using WGS failed to identify.

Since cnv-chr4a was surprisingly large and segregating in our diploid reference, we conducted

further analyses to understand its genomic makeup. Firstly, we analyzed TE content within this

region. We identified TEs using EDTA (97) and used RepeatMasker v4.1.1 (98) to obtain a

summary of various TE families within this region relative to the rest of chromosome 4. We then

performed a synteny analysis to determine if the small number of genes in that region were

collinear with other genomic regions and were therefore an ancestral arrangement, or if they

were novel combinations of genes. We used McScanX v97e74f4 (99) to determine syntenic gene

groups resulting from a self-alignment of protein sequences on haplotype 1 using blastp (-evalue

1e-10) in BLAST v2.7.1 [(100)]). We calculated read-depth on chromosome 4 for three aligned

Ambrosia outgroup species (30). Calculating the mean read-depth exclusively within the cnv-

chr4a region allowed us to ascertain the likely ancestral state of this structural variant.

Temporal CNV changes

To analyze shifts in CNV frequency over both time and space, we used generalized linear models

with the glm function in R (101) to predict presence/absence counts in populations from range

(Europe or North America), sample year and sample latitude, including significant interactions

between the three variables. PC1, which was calculated from a covariance matrix of 10,000

SNPs randomly sampled from outside of genes and putative inversions identified by Battlay et

al. (15), was included in each model to control for population structure. Model significance was

tested with the anova function using a type-3 test (car v3.1-2 package [(102)]). The models were

reduced in a stepwise fashion, removing non-significant interactions until all remaining

interactions (if any) were significant ($p < 0.05$). The emtrends function (emmeans v1.10.2

package [(103)]) was used to test directionality and obtain confidence intervals within interacting

predictors.

CNV-trait associations

To investigate associations between copy number variants and traits, we measured associations

between 29 traits (in 121 samples for which trait data were available [(43)]) and the coverage

windows described above. For each window, we fit linear models, using lm in R (101), between

each sample's normalized window depth and each trait value. To account for population

structure, the first principal component of neutral covariance was added to the model. As

previously, this was obtained by calculating a covariance matrix on 10,000 putatively neutral

sites that were outside gene and inversion regions using PCangsd. A PCA was performed on this

covariance matrix using prcomp in R (101). We assessed significance using a Bonferroni-

corrected significance threshold: 0.05 divided by the number of windows tested (17,855). We

performed CNVr-trait associations using EMMAX (v.beta-7Mar2010 [(104)]) to identify

associations between the CNV regions, using a covariance matrix consisting of the same 10,000

putatively neutral SNPs in previous analyses in this study to correct for population structure.

Gene Ontology analyses

Gene ontology enrichment analyses were performed using the R package topGO (105), using GO

terms from A. thaliana TAIR 10 BLAST results. Fisher's exact test, using a significance

threshold of $p < 0.05$, was used to identify GO terms enriched within candidate gene lists relative

to Q_{ST} outliers, as well as genes within CNVr.

Acknowledgements

We thank Stephen Wright for his advice in the early stages of this project, and we thank Keyne

Monro for her advice and expertise in developing the Q_{ST}-F_{ST} analysis. We also thank Claire

Mérot, an anonymous reviewer, and the editor for their thoughtful comments which greatly

improved the manuscript.

Data Accessibility

Sequences used in reference genome assembly and annotation are available from NCBI under

BioProject ID PRJNA819156. The phased diploid genome assembly is available from NCBI

under BioProject IDs PRJNA929657 and PRJNA929658. The haplotype 1 gene annotation GFF

file is available from Figshare [https://doi.org/10.6084/m9.figshare. 19672710.v1]. Individual

sample resequencing data are available from ENA under BioProject IDs PRJEB48563,

PRJNA339123 and PRJEB34825.

Code availability

All code for analyses performed in this work is available from Github

[https://github.com/jonrobwil/ragweed_cnv]

References

1. R. A. Fisher, XV.—The Correlation between Relatives on the Supposition of Mendelian Inheritance. *Earth Environ. Sci. Trans. R. Soc. Edinb.* **52**, 399–433 (1919).
2. J. K. Pritchard, J. K. Pickrell, G. Coop, The Genetics of Human Adaptation: Hard Sweeps, Soft Sweeps, and Polygenic Adaptation. *Curr. Biol.* **20**, R208–R215 (2010).
3. A. E. Van't Hof, *et al.*, The industrial melanism mutation in British peppered moths is a transposable element. *Nature* **534**, 102–105 (2016).
4. A. J. Jensen, *et al.*, Large-effect loci mediate rapid adaptation of salmon body size after river regulation. *Proc. Natl. Acad. Sci.* **119**, e2207634119 (2022).
5. T. Connallon, K. A. Hodgins, Allen Orr and the genetics of adaptation. *Evolution* **75**, 2624–2640 (2021).
6. H. A. Orr, The Population Genetics of Adaptation: The Distribution of Factors Fixed During Adaptive Evolution. *Evolution* **52**, 935–949 (1998).
7. S. Yeaman, M. C. Whitlock, The genetic architecture of adaptation under migration-selection balance. *Evol. Int. J. Org. Evol.* **65**, 1897–1911 (2011).
8. P. Battlay, S. Yeaman, K. A. Hodgins, Impacts of pleiotropy and migration on repeated genetic adaptation. [Preprint] (2023). Available at: https://www.biorxiv.org/content/10.1101/2021.09.13.459985v5 [Accessed 2 June 2024].
9. M. Wellenreuther, L. Bernatchez, Eco-Evolutionary Genomics of Chromosomal Inversions. *Trends Ecol. Evol.* **33**, 427–440 (2018).
10. M. Chakraborty, J. J. Emerson, S. J. Macdonald, A. D. Long, Structural variants exhibit widespread allelic heterogeneity and shape variation in complex traits. *Nat. Commun.* **10**, 4872 (2019).
11. A. H. Sturtevant, A Case of Rearrangement of Genes in Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **7**, 235–237 (1921).
12. T. Dobzhansky, *Genetics of the Evolutionary Process* (Columbia University Press, 1970).
13. H. Cayuela, *et al.*, Thermal adaptation rather than demographic history drives genetic structure inferred by copy number variants in a marine fish. *Mol. Ecol.* **30**, 1624–1641 (2021).
14. C. K. Tepolt, E. D. Grosholz, C. E. de Rivera, G. M. Ruiz, Balanced polymorphism fuels rapid selection in an invasive crab despite high gene flow and low genetic diversity. *Mol. Ecol.* **31**, 55–69 (2022).

15. P. Battlay, *et al.*, Large haploblocks underlie rapid adaptation in the invasive weed
Ambrosia artemisiifolia. *Nat. Commun.* **14**, 1717 (2023).

16. M. M. Magwire, F. Bayer, C. L. Webster, C. Cao, F. M. Jiggins, Successive Increases in
the Resistance of Drosophila to Viral Infection through a Transposon Insertion Followed
by a Duplication. *PLOS Genet.* **7**, e1002337 (2011).

17. E. R. Lucas, *et al.*, Whole-genome sequencing reveals high complexity of copy number
variation at insecticide resistance loci in malaria mosquitoes. *Genome Res.* **29**, 1250–
1261 (2019).

18. S. DeBolt, Copy number variation shapes genome diversity in Arabidopsis over
immediate family generational scales. *Genome Biol. Evol.* **2**, 441–453 (2010).

19. T. A. Gaines, *et al.*, Gene amplification confers glyphosate resistance in Amaranthus
palmeri. *Proc. Natl. Acad. Sci.* **107**, 1029–1034 (2010).

20. P. J. Daborn, *et al.*, A single p450 allele associated with insecticide resistance in
Drosophila. *Science* **297**, 2253–2256 (2002).

21. R. H. Ffrench-Constant, The molecular genetics of insecticide resistance. *Genetics* **194**,
807–815 (2013).

22. C. Zhang, *et al.*, Investigating the mechanisms of glyphosate resistance in goosegrass
(*Eleusine indica*) population from South China. *J. Integr. Agric.* **14**, 909–918 (2015).

23. J. M. Kreiner, *et al.*, Multiple modes of convergent adaptation in the spread of glyphosate-
resistant Amaranthus tuberculatus. *Proc. Natl. Acad. Sci.* **116**, 21076–21084 (2019).

24. N. M. Reid, *et al.*, The genomic landscape of rapid repeated evolutionary adaptation to
toxic pollution in wild fish. *Science* **354**, 1305–1308 (2016).

25. S. Gimenez, *et al.*, Adaptation by copy number variation increases insecticide resistance
in the fall armyworm. *Commun. Biol.* **3**, 664 (2020).

26. J. Prunier, *et al.*, Gene copy number variations involved in balsam poplar (Populus
balsamifera L.) adaptive variations. *Mol. Ecol.* **28**, 1476–1490 (2019).

27. Y. Dorant, *et al.*, Copy number variants outperform SNPs to reveal genotype–temperature
association in a marine species. *Mol. Ecol.* **29**, 4765–4782 (2020).

28. K. A. Hodgins, D. G. Bock, L. H. Rieseberg, "Trait Evolution in Invasive Species" in
Annual Plant Reviews Online, (John Wiley & Sons, Ltd, 2018), pp. 459–496.

29. R. I. Colautti, S. C. H. Barrett, Rapid Adaptation to Climate Facilitates Range Expansion
of an Invasive Plant. *Science* **342**, 364–366 (2013).

30. V. C. Bieker, *et al.*, Uncovering the genomic basis of an extraordinary plant invasion. *Sci.*
Adv. **8**, eabo5115 (2022).

31. A. McGaughran, *et al.*, Genomic Tools in Biological Invasions: Current State and Future
Frontiers. *Genome Biol. Evol.* **16**, evad230 (2024).

32. D. L. Stern, The genetic causes of convergent evolution. *Nat. Rev. Genet.* **14**, 751–764
(2013).

33. L. A. van Boheemen, K. A. Hodgins, Rapid repeatable phenotypic and genomic
adaptation following multiple introductions. *Mol. Ecol.* **29**, 4102–4117 (2020).

34. K. M. Dlugosch, S. R. Anderson, J. Braasch, F. A. Cang, H. D. Gillette, The devil is in the
details: genetic variation in introduced populations and its contributions to invasion. *Mol.*
Ecol. **24**, 2095–2111 (2015).

35. A. Estoup, *et al.*, Is There a Genetic Paradox of Biological Invasion? *Annu. Rev. Ecol.*
Evol. Syst. **47**, 51–72 (2016).

36. F. Essl, *et al.*, Biological Flora of the British Isles: Ambrosia artemisiifolia. *J. Ecol.* **103**,
1069–1098 (2015).

37. U. Schaffner, *et al.*, Biological weed control to relieve millions from Ambrosia allergies in
Europe. *Nat. Commun.* **11**, 1745 (2020).

38. M. J. Cowbrough, R. B. Brown, F. J. Tardif, Impact of common ragweed (Ambrosia
artemisiifolia) aggregation on economic thresholds in soybean. *Weed Sci.* **51**, 947–954

(2003).
39. C. E. Brewer, L. R. Oliver, Confirmation and Resistance Mechanisms in Glyphosate-
Resistant Common Ragweed (Ambrosia artemisiifolia) in Arkansas. *Weed Sci.* **57**, 567–
573 (2009).
40. L. Hamaoui-Laguel, *et al.*, Effects of climate change and seed dispersal on airborne
ragweed pollen loads in Europe. *Nat. Clim. Change* **5**, 766–771 (2015).
41. Y. Sun, G. K. Roderick, Rapid evolution of invasive traits facilitates the invasion of
common ragweed, Ambrosia artemisiifolia. *J. Ecol.* **107**, 2673–2687 (2019).
42. L. A. van Boheemen, *et al.*, Multiple introductions, admixture and bridgehead invasion
characterize the introduction history of Ambrosia artemisiifolia in Europe and Australia.
Mol. Ecol. **26**, 5421–5434 (2017).
43. L. A. van Boheemen, D. Z. Atwater, K. A. Hodgins, Rapid and repeated local adaptation
to climate in an invasive plant. *New Phytol.* **222**, 614–627 (2019).
44. A. J. Gorton, D. A. Moeller, P. Tiffin, Little plant, big city: a test of adaptation to urban
environments in common ragweed (Ambrosia artemisiifolia). *Proc. R. Soc. B Biol. Sci.*
285, 20180968 (2018).
45. B. V. McGoey, J. R. Stinchcombe, Introduced populations of ragweed show as much
evolutionary potential as native populations. *Evol. Appl.* **14**, 1436–1449 (2021).
46. R. C. Lewontin, J. Krakauer, DISTRIBUTION OF GENE FREQUENCY AS A TEST OF
THE THEORY OF THE SELECTIVE NEUTRALITY OF POLYMORPHISMS. *Genetics* **74**,
175–195 (1973).
47. B. S. Weir, C. C. Cockerham, Estimating F-Statistics for the Analysis of Population
Structure. *Evolution* **38**, 1358–1370 (1984).
48. M. C. Whitlock, Evolutionary inference from QST. *Mol. Ecol.* **17**, 1885–1896 (2008).
49. J. H. McDonald, M. Kreitman, Adaptive protein evolution at the Adh locus in Drosophila.
Nature **351**, 652–654 (1991).
50. T. R. Booker, S. Yeaman, M. C. Whitlock, Variation in recombination rate affects
detection of outliers in genome scans under neutrality. *Mol. Ecol.* **29**, 4274–4279 (2020).
51. D. Prapas, *et al.*, Quantitative trait loci mapping reveals an oligogenic architecture of a
rapidly adapting trait during the European invasion of common ragweed. *Evol. Appl.* **15**,
1249–1263 (2022).
52. S. E. Fick, R. J. Hijmans, WorldClim 2: new 1-km spatial resolution climate surfaces for
global land areas. *Int. J. Climatol.* **37**, 4302–4315 (2017).
53. B. J. Adamczyk, D. E. Fernandez, MIKC* MADS Domain Heterodimers Are Required for
Pollen Maturation and Tube Growth in Arabidopsis. *Plant Physiol.* **149**, 1713–1723
(2009).
54. D. R. Schrider, M. W. Hahn, D. J. Begun, Parallel Evolution of Copy-Number Variation
across Continents in Drosophila melanogaster. *Mol. Biol. Evol.* **33**, 1308–1316 (2016).
55. E. L. Patterson, D. J. Pettinga, K. Ravet, P. Neve, T. A. Gaines, Glyphosate Resistance
and EPSPS Gene Duplication: Convergent Evolution in Multiple Plant Species. *J. Hered.*
109, 117–125 (2018).
56. R. T. Good, *et al.*, The Molecular Evolution of Cytochrome P450 Genes within and
between Drosophila Species. *Genome Biol. Evol.* **6**, 1118–1134 (2014).
57. A. Abyzov, A. E. Urban, M. Snyder, M. Gerstein, CNVnator: An approach to discover,
genotype, and characterize typical and atypical CNVs from family and population genome
sequencing. *Genome Res.* **21**, 974–984 (2011).
58. S. Zarate, *et al.*, Parliament2: Accurate structural variant calling at scale. *GigaScience* **9**,
giaa145 (2020).
59. G. van Ooijen, *et al.*, Structure-function analysis of the NB-ARC domain of plant disease
resistance proteins. *J. Exp. Bot.* **59**, 1383–1397 (2008).
60. A. Żmieńko, A. Samelak, P. Kozłowski, M. Figlerowicz, Copy number polymorphism in

plant genomes. *Theor. Appl. Genet.* **127**, 1–18 (2014).

61. A. Dolatabadian, D. A. Patel, D. Edwards, J. Batley, Copy number variation and disease
resistance in plants. *Theor. Appl. Genet.* **130**, 2479–2490 (2017).

62. M. Todesco, *et al.*, Massive haplotypes underlie ecotypic differentiation in sunflowers.
Nature **584**, 602–607 (2020).

63. C. Mérot, *et al.*, Genome assembly, structural variants, and genetic differentiation
between lake whitefish young species pairs (Coregonus sp.) with long and short reads.
Mol. Ecol. **32**, 1458–1477 (2023).

64. T. Hämälä, *et al.*, Genomic structural variants constrain and facilitate adaptation in natural
populations of Theobroma cacao , the chocolate tree. *Proc. Natl. Acad. Sci.* **118**,
e2102914118 (2021).

65. J. B. Koch, *et al.*, Population genomic and phenotype diversity of invasive Drosophila
suzukii in Hawai'i. *Biol. Invasions* **22**, 1753–1770 (2020).

66. N. R. Hofmeister, S. J. Werner, I. J. Lovette, Environmental correlates of genetic variation
in the invasive European starling in North America. *Mol. Ecol.* **30**, 1251–1263 (2021).

67. T. Lin, P. G. L. Klinkhamer, K. Vrieling, Parallel evolution in an invasive plant: effect of
herbivores on competitive ability and regrowth of Jacobaea vulgaris. *Ecol. Lett.* **18**, 668–
676 (2015).

68. K. C. Stuart, W. B. Sherwin, R. J. Edwards, L. A. Rollins, Evolutionary genomics: Insights
from the invasive European starlings. *Front. Genet.* **13** (2023).

69. J. Ebler, *et al.*, Pangenome-based genome inference allows efficient and accurate
genotyping across a wide spectrum of variant classes. *Nat. Genet.* **54**, 518–525 (2022).

70. B. Chauvel, F. Dessaint, C. Cardinal-Legrand, F. Bretagnolle, The historical spread of
Ambrosia artemisiifolia L. in France from herbarium records. *J. Biogeogr.* **33**, 665–673
(2006).

71. M. J. Case, K. A. Stinson, Climate change impacts on the distribution of the allergenic
plant, common ragweed (Ambrosia artemisiifolia) in the eastern United States. *PLOS*
ONE **13**, e0205677 (2018).

72. A. S. Kim, *et al.*, Temporal collections to study invasion biology. *Mol. Ecol.* **32**, 6729–6742
(2023).

73. J. M. Alves, *et al.*, Parallel adaptation of rabbit populations to myxoma virus. *Science* **363**,
1319–1326 (2019).

74. J. M. Kreiner, S. Hnatovska, J. R. Stinchcombe, S. I. Wright, Quantifying the role of
genome size and repeat content in adaptive variation and the architecture of flowering
time in Amaranthus tuberculatus. *PLOS Genet.* **19**, e1010865 (2023).

75. B. E. Stranger, *et al.*, Relative Impact of Nucleotide and Copy Number Variation on Gene
Expression Phenotypes. *Science* **315**, 848–853 (2007).

76. M. Kirkpatrick, B. Barrett, Chromosome inversions, adaptive cassettes and the evolution
of species' ranges. *Mol. Ecol.* **24**, 2046–2055 (2015).

77. G. Potente, *et al.*, Comparative Genomics Elucidates the Origin of a Supergene
Controlling Floral Heteromorphism. *Mol. Biol. Evol.* **39**, msac035 (2022).

78. P. Battlay, *et al.*, Structural variants underlie parallel adaptation following global invasion.
[Preprint] (2024). Available at:
https://www.biorxiv.org/content/10.1101/2024.07.09.602765v1 [Accessed 12 December
2024].

79. M. Wellenreuther, C. Mérot, E. Berdan, L. Bernatchez, Going beyond SNPs: The role of
structural genomic variants in adaptive evolution and species diversification. *Mol. Ecol.*
28, 1203–1209 (2019).

80. M. Schubert, *et al.*, Characterization of ancient and modern genomes by SNP detection
and phylogenomic and metagenomic analysis using PALEOMIX. *Nat. Protoc.* **9**, 1056–
1082 (2014).

81.   M. A. DePristo, *et al.*, A framework for variation discovery and genotyping using next-
846   generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011).

847   82.   P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008
848   (2021).

849   83.   G. Coop, D. Witonsky, A. Di Rienzo, J. K. Pritchard, Using Environmental Correlations to
850   Identify Loci Underlying Local Adaptation. *Genetics* **185**, 1411–1423 (2010).

851   84.   M. Gautier, Genome-Wide Scan for Adaptive Divergence and Association with
852   Population-Specific Covariates. *Genetics* **201**, 1555–1579 (2015).

853   85.   K. Spitze, Population structure in Daphnia obtusa: quantitative genetic and allozymic
854   variation. *Genetics* **135**, 367–374 (1993).

855   86.   T. Prout, J. S. Barker, F statistics in Drosophila buzzatii: selection, population size and
856   inbreeding. *Genetics* **134**, 369–375 (1993).

857   87.   D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using
858   lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

859   88.   P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
860   (2011).

861   89.   H. Li, P. Ralph, Local PCA Shows How the Effect of Population Structure Differs Along
862   the Genome. *Genetics* **211**, 289–304 (2019).

863   90.   T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation
864   Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).

865   91.   J. Meisner, A. Albrechtsen, Inferring Population Structure and Admixture Proportions in
866   Low-Depth NGS Data. *Genetics* **210**, 719–731 (2018).

867   92.   L. Mouselimis, *et al.*, ClusterR: Gaussian Mixture Models, K-Means, Mini-Batch-Kmeans,
868   K-Medoids and Affinity Propagation Clustering. (2023). Deposited 4 December 2023.

869   93.   H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
870   3100 (2018).

871   94.   S. Kurtz, *et al.*, Versatile and open software for comparing large genomes. *Genome Biol.*
872   **5**, R12 (2004).

873   95.   M. Goel, H. Sun, W.-B. Jiao, K. Schneeberger, SyRI: finding genomic rearrangements
874   and local sequence differences from whole-genome assemblies. *Genome Biol.* **20**, 277
875   (2019).

876   96.   M. Goel, K. Schneeberger, plotsr: visualizing structural similarities and rearrangements
877   between multiple genomes. *Bioinformatics* **38**, 2922–2926 (2022).

878   97.   S. Ou, *et al.*, Benchmarking transposable element annotation methods for creation of a
879   streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

880   98.   Smit, AFA, Hubley, R & Green, P., RepeatMasker Open-4.0. 2013-2015
881   <http://www.repeatmasker.org>.

882   99.   Y. Wang, *et al.*, MCScanX: a toolkit for detection and evolutionary analysis of gene
883   synteny and collinearity. *Nucleic Acids Res.* **40**, e49 (2012).

884   100.   C. Camacho, *et al.*, BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
885   (2009).

886   101.   R Foundation for Statistical Computing, Vienna, Austria., R: A language and environment
887   for statistical computing. (2024). Deposited 2024.

888   102.   J. Fox, S. Weisberg, *An R Companion to Applied Regression*, Third (Sage, 2019).

889   103.   R. V. Lenth, *et al.*, emmeans: Estimated Marginal Means, aka Least-Squares Means.
890   (2024). Deposited 23 January 2024.

891   104.   H. M. Kang, *et al.*, Variance component model to account for sample structure in
892   genome-wide association studies. *Nat. Genet.* **42**, 348–354 (2010).

893   105.   A. Alexa, J. Rahnenführer, Gene set enrichment analysis with topGO.

Figure 1. CNV Q$_{ST}$ values indicate regions of selection in the *A. artemisiifolia* genome. A. Q$_{ST}$ values of filtered coverage windows in European populations. Pink windows indicate those above 1% F$_{ST}$ threshold shown in B, blue windows indicate top 1% of Q$_{ST}$ values. C. Q$_{ST}$ values of filtered coverage windows in North American populations. Pink windows indicate those above 1% F$_{ST}$ threshold shown in D, blue windows indicate top 1% of Q$_{ST}$ values. Bars above manhattan plots indicate merged windows > 300 kbp.

Figure 2. Overlapping Q$_{ST}$ outliers in both ranges and their potential biological functions.

A. Distribution of Q$_{ST}$ values of all 10 kbp coverage windows in both North America and Europe. Overlapping windows exceeding the neutral F$_{ST}$ threshold of 1% in each respective range are colored in pink. B. Gene ontology enrichment plot of genes within overlapping outlier Q$_{ST}$ windows, with biological pathways only retained if represented by 2 or more genes (pink in A).

Figure 3. CNVr genotyping of cnv-chr4a. A. MDS coordinates from local PCA in 100kb windows across chromosome 4. Values diverging from zero indicate local structure compared with the rest of the chromosome. B. Distribution of merged window sizes along chromosome 4, with boxes indicating window size on the Y axis and chromosomal position on the X axis. C. Alignment of chromosome 4 Hap1 and Hap2 of the diploid reference genome. A clear gap is present corresponding to local PCA and merged window locations. D, E. Regional PC1 against regional read-depth corresponding to merged window identified in B. Clustering computed using k-means and manual annotation. D. represents modern samples and E. represents historic samples.

Figure 4. Spatio-temporal CNV frequency shifts. A, B. Logistic regression models for A) cnv-chr14a and B) cnv-chr-17a with error bands representing 95% CI of least-squares regressions of CNV frequency. Time binned into five categories, ranging from historic (purple) to modern (green). Model information found in Supplementary Table 6. C, D. Frequency of C) cnv-chr14a and D) cnv-chr17a in modern *A. artemisiifolia* populations.