Title: Integrated Analysis of Single-Cell and Bulk RNA-Seq Data reveals that

Ferroptosis-Related Genes Mediated the Tumor Microenvironment predicts Prognosis,

and guides Drug Selection in Triple-Negative Breast Cancer

We kindly request the opportunity to deposit our work as a preprint on bioRxiv or medRxiv, depending on its suitability.

Xuantong Gong[1#], Lishuang Gu[2#], Di Yang[1], Yu He[1], Qian Li[3*], Hao Qin[4*], Yong Wang[1,5*]

First authors: Xuantong Gong and Lishuang Gu contributed equally to this work.

Corresponding authors: Correspondence to Yong Wang, Hao Qin or Qian Li

[1]Department of Ultrasound, National Cancer Center/National Clinical Research Center for

Cancer/ Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical

College, Beijing, China

[2]Department of Ultrasound, Beijing Hospital of Traditional Chinese Medicine, Capital

Medical University, Beijing, China

[3]Department of Ultrasound, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou,

China

[4]Key Laboratory of Cancer and Microbiome, State Key Laboratory of Molecular Oncology,

National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital,

Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China.

[5]Department of Ultrasound, The First Affliated Hospital of China Medical University,

Shenyang, China

Email: Xuantong Gong: gongxuantong@163.com

Lishuang Gu: 13026300487@163.com

Di Yang: yangdi4134@163.com

Yu He: 1282302858@qq.com

*: Correspondence to Yong Wang, Hao Qin, Qian Li

Yong Wang

Department of Ultrasound, National Cancer Center/National Clinical Research Center for

Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical

College, Beijing 100021, China.

Department of Ultrasound, The First Affliated Hospital of China Medical University,

Shenyang, China

Email address: drwangyong77@163.com

Hao Qin

Key Laboratory of Cancer and Microbiome, State Key Laboratory of Molecular Oncology,

National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital,

Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China.

Email address: haoqin@cicams.ac.cn

Qian Li

Department of Ultrasound, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou,

China

Email address: 754427296@qq.com

Declarations

•Ethics approval and consent to participate: Not applicable

•Consent for publication: Not applicable

•Availability of datasets：The data used in this study are available in the public datasets. Nine

single-cell RNA-seq TNBC samples were obtained from the Gene Expression Omnibus (GEO)

database (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi). The data for external validation were

obtained from the GEO database. The data of 471 genes associated with ferroptosis were collected

from the FerrDb website (http://www.zhounan.org/ferrdb/current/).

•Competing Interest: The authors declare that they have no known competing financial

interests or personal relationships that could have appeared to influence the work reported in

this paper.

•Authors' contributions: YW ： Conceptualization, Methodology, Formal analysis,

Investigation, Writing-original draft preparation, Writing-review & editing, Project

administration and Funding acquisition. XTG: Conceptualization, Methodology, Formal

analysis, Investigation, Writing-original draft preparation, Writing-review & editing,

Resources and Data curation. LSG: Conceptualization, Methodology, Formal analysis,

Investigation, Writing-original draft preparation, Writing-review & editing, Resources and

Data curation. DY: Writing-original draft preparation and Writing-review & editing. YH:

Writing-original draft preparation and Writing-review & editing. QL: Conceptualization,

Methodology, Formal analysis, Investigation, Writing-original draft preparation, Writing-

review & editing,   Visualization and Supervision. HQ: Conceptualization, Methodology,

Formal analysis, Investigation, Writing-original draft preparation, Writing-review & editing,

Visualization and Supervision. All authors: reading and approval of the final manuscript.

•Impact Statement: Integrating single-cell and bulk RNA-seq data elucidates the role of

ferroptosis in TNBC, offering a prognostic model and personalized therapeutic insights.

Key words: RNA-seq; ferroptosis; tumor microenvironment; triple-negative breast cancer

Abstract

Background

Triple-negative breast cancer (TNBC) is aggressive, lacking methods to predict recurrence and

drug sensitivity. Ferroptotic heterogeneity varies in TNBC subtypes. However, the tumor

microenvironment (TME) mediated by ferroptosis genes is unclear. Our study aims to integrate

single-cell and bulk RNA sequencing (RNA-seq) data to reveal the ferroptosis-mediated TME in

TNBC, predicting prognosis and guiding treatment.

Methods

The single-cell and bulk RNA-seq data of TNBC were sourced from the Gene Expression

Omnibus (GEO) database. Using these data, a machine learning algorithm was employed to

integrate and analyze the characteristics of the TME mediated by ferroptosis-related genes in

TNBC. Prediction models for TNBC survival prognosis and drug treatment response were

established and then validated in an independent set.

Results

At the individual cell level, T cells were categorized into three distinct subpopulations, and local

macrophages into two subpopulations. The infiltration degree of these different cell

subpopulations was closely associated with prognosis and treatment outcomes. Based on this, the

risk score model we developed effectively predicted recurrence-free survival in TNBC patients,

with independently validated pooled predicted 3-, 4-, and 5-year Area Under the Curves (AUCs)

of 0.65, 0.67, and 0.71, respectively. Additionally, we found that patients in the high-risk group

may be more responsive to 27 drugs.

Conclusions

We have uncovered the tumor immune cell clusters in TNBC mediated by ferroptosis. A risk score

model was constructed to identify high-risk TNBC patients, which can assist physicians in disease

monitoring and precision therapy. The genes identified hold significant potential as therapeutic

targets for TNBC patients.

Funding

This project is funded by the National Natural Science Foundation of China (81974268,

82472000, 82304151), the Talent Incentive Program of Cancer Hospital Chinese, Academy of

Medical Sciences (801032247), the Cooperation Fund of CHCAMS (CFA202202023), and

the open project of Beijing Key Laboratory of Tumor Invasion and Metastasis Mechanism,

Capital Medical University(2023ZLKF03).

101 1. Background

Global cancer statistics show that breast cancer is the most common cancer in women and the

leading cause of cancer deaths [1]. Triple-negative breast cancer (TNBC) is a subtype of breast

cancer that lacks expression of estrogen receptor, progesterone receptor and human epidermal

growth factor receptor-2. Compared with other subtypes, TNBC is highly aggressive, with a poor

overall prognosis for patients and a median survival of less than 1 year after recurrent metastasis [2-

5]. Due to the lack of targets for endocrine therapy and targeted therapeutic agents, chemotherapy

and emerging immunotherapy are the main therapies. TNBC is highly heterogeneous and only

some patients benefit from treatment [6-8]. Unfortunately, there is no effective method that can

effectively predict the prognostic risk and drug sensitivity of TNBC, which is an urgent problem in

the clinic.

The tumor microenvironment (TME) is composed of various cell types such as cancer-associated

fibroblasts, tumor-associated macrophages, T cells, NK cells, B cells, endothelial cells, etc., which

interact with cancer cells and influence various aspects such as tumor progression, metastasis, and

response to therapy [9-11]. The composition and functional status of TME varies greatly among

different patients with breast cancer, and revealing the TME of each patient is crucial for selecting

a reasonable treatment and controlling tumor progression in the long term [12-13].

Ferroptosis is an iron-dependent form of non-apoptotic, oxidative form of regulated cell death

involving lipid hydroperoxides [14-15]. Studies have shown that drug-resistant cancer cells are more

sensitive to ferroptosis [16-18]. Therefore, ferroptosis is more recognized as a potential target for

cancer therapy. Ferroptotic heterogeneity exists in different subtypes of TNBC [19]. Since

ferroptosis is regulated by multiple metabolic pathways, the ferroptosis landscape of TNBC

remains unexplored and its relationship with patient prognosis and treatment response is uncertain

[20]. Therefore, exploring ferroptosis-related genes in TNBC and their correlation with the immune

microenvironment may provide a new treatment trend for TNBC.

RNA sequencing (RNA-seq) technology is a gene expression analysis method that can

qualitatively and quantitatively explore the transcriptome characteristics of biological samples at

the tissue and cellular levels [21-22]. Single-cell RNA-seq (scRNA-seq) has greatly enhanced our

understanding of transcriptional heterogeneity across cell types and states, and can be used to

explore the TME [23-24]. Bulk RNA-seq is used to analyze the average expression levels of RNA in

tissues or cell populations, and can be utilized to explore differences among individuals [25-26].

Machine learning is a branch of artificial intelligence that enables computer systems to learn from

training data and gain experience, and it is increasingly being used for the diagnosis, treatment,

and prognostic evaluation of cancer [27-28]. Literature reports that several models have been

constructed based on RNA-seq data for the prognostic evaluation of breast cancer patients, but the

reproducibility and interpretability of the results still deserve further exploration [29-32]. This study

intends to further explore the relationship among ferroptosis genes, prognosis, and drug sensitivity

from the perspective of ferroptotic heterogeneity in TNBC. In this work, we collected and

integrated data from single-cell and bulk RNA-seq to characterize the ferroptosis-related gene

mediated TME landscape of TNBC on multiple scales. furthermore, based on the findings in TME,

predictive models of patient prognosis and treatment response were constructed using machine

learning algorithms. We hope to provide a new way of individualized risk management for triple-

negative breast cancer patients and assist their precision treatment.

2. Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**

software, algorithm	R package Seurat	Microsoft	N/A	Version 4.1.1

148   2.1 Patient data collection and processing

149   The data used in this study were collected from public datasets. Nine single-cell RNA-seq TNBC

150   samples   were   obtained   from   the   Gene   Expression   Omnibus   (GEO)   database

151   (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi) under the accession number GSE176078 [33]

152   (Table1).

Table 1 Sample information for triple-negative breast cancer in GSE176078

Case ID	Gender	Age	T_Stage	Subtype by IHC
3946	Female	52	T2	TNBC
44041	Female	35	T2	TNBC
4465	Female	54	T2	TNBC
4495	Female	63	T1	TNBC
44971	Female	49	T2	TNBC
44991	Female	47	NA	TNBC
4513	Female	73	T3	TNBC
4515	Female	67	T1	TNBC
4523	Female	52	T2	TNBC

TNBC: Triple-negative breast cancer, IHC: Immunohistochemistry

To construct the model to predict prognosis and immunotherapy efficacy, bulk RNA-seq datasets

and clinicopathological information of TNBC were obtained from the GEO database under the

accession number GSE25066 [34]. The data for external validation were obtained from the GEO

database under the accession number GSE86166 [35].

2.2 Single-cell RNA-seq data processing

Data were processed using the R package Seurat (version 4.1.1) for quality control (QC). Based on

QC metrics suggested in Scanpy tutorial, outlier cells were removed based on relevant feature data

(nFeature_RNA, nCount_RNA, percent.mt), i.e., cells with less than 200 genes expressed or more

than 20% mitochondrial genes counts were filtered out. After quality control, the expression data

of each cell was normalized separately using the NormalizeData function and the top 2000 highly

variable genes (HVGs) of each cell were identified using the FindVariableFeatures function. To

remove batch effects among different samples, we utilized the FindIntegrationAnchors function to

find anchor genes and then the reclassified datasets were integrated utilized the IntegrateData

function. Data normalization was performed using the ScaleData function in Seurat. The RunPCA

function was utilized to reduce the dimension of principal component analysis (PCA) for the first

2000 HVGs screened above.

2.3 Genes associated with ferroptosis acquiring

The data of 471 genes (Supplementary Table 1) associated with ferroptosis were collected from

the FerrDb website (http://www.zhounan.org/ferrdb/current/). Then, from the above genes,

screened for genes with expression information in the single-cell matrix of TNBC patients. Firstly,

based on the clinical data of the samples, the expression values of ferroptosis-related genes in the

same type of samples were averaged and presented in a heatmap. Afterwards, the expression

values of ferroptosis-related genes were averaged across different cell types and also presented in

a heatmap.

2.4 Cell subtype identification associated with ferroptosis

Because information of cell type annotation was listed in the original literature of the GSE176078

dataset, the metadata of the original literature was used directly for the next analyses. The R

package Seurat (version 4.1.1) was used to identify cell types and presented results via t-SNE.

Non-negative matrix decomposition (NMF) is an algorithm based on high-throughput data to

identify and cluster out different molecular functional patterns [36]. In this work, subtypes of

macrophages and T cells based on genes associated with ferroptosis using the "NMF" R package

(v 0.26).

2.5 Cell-cell communication analysis

Cell-cell communication mediated by ligand-receptor complexes plays an important role in

multiple biological processes [37]. In this study, the "iTALK" R package was utilized to construct

the cellular

communication network. First, use the rawParse function to identify the top 50% highly expressed

genes for each cell type based on their expression means. Then, utilize the FindLR function to

identify ligands and receptors among the highly expressed genes. Finally, construct an interaction

network based on all the interaction relationships.

2.6 Pseudotime analysis

Pseudotime analysis of scRNA-seq snapshot data helps to provide an approximate landscape of

gene expression dynamics [38]. The "Monocle" R package (v 2.28.0) was applied for pseudotime

analysis to conduct cellular trajectory. The reduceDimensio function based on the DDRTree

algorithm is used to reduce the dimensions of the data. Based on the Progenitor Cell Biology

Consortium (PCBC) database (https://www.synapse.org), the stemness signature was identified via

the one-class logistic regression (OCLR) algorithm, then the stemness index of each TNBC cell

was calculated by scaling the Spearman correlation coefficients to be between 0 and 1. Eventually,

the order Cells function was utilized to sort cells and complete construction of trajectory.

2.7 Transcriptional factor analysis

The Single-cell Regulatory Network Inference and Clustering tool (SCENIC) enables

simultaneous gene regulatory network reconstruction and cell-state identification from scRNA-seq

data [39-40]. We utilized the "SCENIC" R package (v1.3.1) to establish the transcription factor (TF)

regulatory network. Specifically, first, coexpression modules are inferred using the

"GENIE3/GRNBOOST" function to identify gene set with co-expressed TFs. Next, the indirect

targets are pruned from these modules using cis-regulatory motif discovery (cisTarget). Finally,

the AUCell algorithm was utilized to evaluate the activity of regulons.

2.8 Hallmarks gene set enrichment analysis

Hallmarks gene set enrichment analysis was utilized the "irGSEA" R package (v2.1.5). The

ssGSEA algorithm was utilized to conduct differential pathway score between ferroptosis-related

immune cell subtypes.

2.9 Construction and validation of the prognostic model

First, based on the recurrence-free survival (RFS) data of 178 TNBC patients in the GSE25066

database, this study utilized the "survival" (v3.2-7) and "survminer" (v0.4.8) R package to

perform univariate Cox proportional hazards regression analyses on ferroptosis-related genes

signature of different immune cell subtypes.

The genes screened for significant association with RFS in the regression analysis (p<0.01) were

utilized to construct a risk factor-based model by the Least Absolute Shrinkage and Selection

Operator (LASSO) method implemented in the "glmnet" R package (v4.0-2). TNBC patients were

divided into low- and high-risk groups according to the median risk score, and Kaplan-Meier

survival curves were utilized to compare the RFS rates between the two groups, p value < 0.05

was considered to significance. To validate the performance of the model, receiver operating

characteristic (ROC) curves were demonstrated and the AUC values were calculated for

evaluating 3-year, 4-year, and 5-year RFS rates of TNBC patients in GSE25066 database. In

addition, the GSE86166 as well as the TCGA database were utilized as external validation sets to

further verify the robustness of the performance of the constructed model in this study.

2.10 Immune cells infiltration and drug sensitivity analysis

Immune cells infiltration is important to the anti-tumor response, and these cells are diverse

among patients [41-42]. The "CIBERSORT", "GSVA" and "TIMER" R package was utilized to

determine the distribution of different immune cell types between low- and high-risk groups. Next,

the 50% inhibitory concentration (IC50) of 138 chemotherapeutic drugs was calculated for each

patient using the "pRRophetic" R package (v 0.5), and drugs significantly associated with the risk

score were screened. Finally, we calculated the tumor immune dysfunction and exclusion (TIDE)

score (http://tide.dfci.harvard.edu.) for each patient and thus analyzed its difference between low-

and high-risk groups.

2.11 Statistical analysis

The R software (version 4.1.3) was utilized for all data analysis. Wilcoxon-rank sum test was

utilized to analyze associations of continuous variables. Log-rank test was utilized to analyze

differences in survival curves between groups. P value less than 0.05 was considered statistically

significant.

3. Results

3.1 The single-cell landscape of TNBC samples

In this study, a total of 9 TNBC single-cell samples were included. After QC, 38985 genes from

29733 cells per sample were finally selected for subsequent analysis (Supplementary Figure 1).

Based on original annotations of data, these cells were divided into 9 major cell clusters and 29

minor cell clusters (Figure 1a-c). Major cell clusters include B-cells, Cancer-associated Fibroblasts

(CAFs), Cancer Epithelial, Endothelial, Myeloid, Normal Epithelial, Plasmablasts, Perivascular-

like Cells (PVL), T-cells. Of these, Myeloid cluster is further divided into Cycling_Myeloid,

Dendritic Cells (DCs), Macrophage, and Monocyte clusters. As shown in Figure 1c, the immune

microenvironment of TNBC is characterized by a high proportion of T cells and macrophages. On

this basis, ligand-receptor interactions between different major cell clusters are frequent (Figure

1d).

Of the 471 ferroptosis-related genes from the FerrDb website, 391 were present in the single-cell

expression matrix of TNBC. We found that there were significant differences in the average

expression of these genes in different clinicopathological classifications, as shown in Figure 1e.

Moreover, there were significant differences in the expression of ferroptosis-related genes in 12

major cell clusters of TNBC (Figure 1f).

Fig. 1 Integration and clustering of scRNA-Seq data from triple-negative breast cancer. (a) t-SNE plot
of the 9 major cell clusters. (b) t-SNE plot of the 29 minor cell clusters. (c) t-SNE plot of the 12 major
cell clusters. (d) The number of ligand-receptor interactions of different major cell clusters in cell-cell
communication network, different colors represent different cell clusters and arrows represent ligand-
receptor orientation. (e) Heat map showing the average expression of ferroptosis-related genes in
different clinicopathological classifications. (f) Heat map showing the average expression of
ferroptosis-related genes in different major cell clusters.
280

3.2 Ferroptosis-related subpopulations of T cells in the TNBC

A total of 11,784 T cells in the scRNA-seq data of this study, including cycling T cells, NK cells,

NKT cells, T cells CD4+, and T cells CD8+ identified clusters (Figure 2a). Based on the NMF

algorithm (rank=3), each T cell cluster was further categorized into 3 ferroptosis-related

subpopulations (T_C1, T_C2, T_C3). Figure 2b shows that differentially expressed top 50

ferroptosis-related genes differed significantly among these 3 subpopulations. Among them, the

proportion of NK cells in the T_C2 subpopulation was significantly more than the other two

subpopulations (Figure 2c). According to pseudotime trajectories, we found that all three T cell

subpopulations were involved at different periods of differentiation (Figure 2d-f). Further, Figure

2g shows the average expression of signature genes associated with 8 functions in 15

subpopulations of T-cells, and we found that gene expression was significantly higher in the

Cycling T-cells. Moreover, different T-cell subpopulations play an important role in the anti-tumor

process (Figure 2h).

TFs are key regulators in cellular signal transduction [43]. As a result, RFX5, EOMES, TBX21,

CEBPB, RUNX3 and IKZF3 showed higher activities in NKT cells. Besides, IRF, NFATC2, STAT

and PRDM1 showed higher activities in CD8$^+$ T cells (Figure 2i). The HALLMARK analysis

results revealed a prominent enrichment in pathways such as pathways in DNA−Repair and

adipogenesis in cycling T cells, whereas in other clusters of T cells, pathways such as epithelial-

mesenchymal transition and KRAS signaling had higher enrichment. However, the difference in

enrichment scores between ferroptosis-related subpopulations of T cells was not significant.

Fig. 2 Ferroptosis-related subpopulations of T cells in triple-negative breast cancer. (a) t-SNE plot of
the 5 identified clusters of T cells. (b) Heat map showing the expression of ferroptosis-related genes
(Top 50) in 3 subpopulations of T cells. (c) Demonstration of the proportion of clusters of T cells in
ferroptosis-related subpopulations. (d) Pseudotime trajectories showing the developmental time course
of T cells. (e) Pseudotime trajectories of the 5 identified clusters of T cells. (f) Pseudotime trajectories
of ferroptosis-related subpopulations of T cells. (g) Heat map showing the average expression of
signature genes associated with 8 functions in 15 subpopulations of T-cells. (h) Network diagram of the
cell-cell communication between cancer epithelial cells and T cells. (i) Heat map for differential
analysis of transcription factors activity.

3.3 Ferroptosis-related subpopulations of macrophages in the TNBC

A total of 3671 macrophages in the scRNA-seq data of this study. Based on the NMF algorithm

(rank=4), 2 ferroptosis-related subpopulations of macrophages (M_C1, M_C2) were finally

obtained after clustering cell clusters with similar markers, the t-SNE plot was shown in Figure 3a.

The pseudotime trajectories showed that M_C2 cells belong to the early stage of macrophages and

then gradually develop into M_C1 cells (Figure 3b-c). Figure 3d shows that differentially

expressed top 50 ferroptosis-related genes differed significantly among subpopulations.

Further, we found that FOLR2, SEPP1, MRC1, LYVE1, SLC40A1, and CD163 were highly

expressed in M_C1 cells and named M_C1 as FOLR2+MAC. whereas, TREM2, FN1, CXCR4,

C3, S100A8, IFI6, and SPP1 were highly expressed in M_C2 and named M_C2 as TREM2+MAC.

Figure 3e-f shows the marker genes in each subpopulation of macrophages. Enrichment scores of

the 2 subpopulations in each cell are shown by t-SNE plots (Figure 3g-h). Moreover, the

enrichment score of TREM2+MAC and FOLR2+MAC were significantly negatively correlated

(r=-0.779) (Figure 3i). Communication between these two subpopulations and cancer epithelial

cells is vigorous (Figure 3j). In addition, detection of TFs activity in each macrophage cell

revealed that most of TFs had higher activity in FOLR2+ MAC cells (Supplementary Figure 2).

Fig. 3 Ferroptosis-related subpopulations of macrophages in triple-negative breast cancer. (a) t-SNE
plot of 2 ferroptosis-related subpopulations of macrophages. (b) Pseudotime trajectories showing the
developmental time course of macrophages. (c) Pseudotime trajectories of 2 ferroptosis-related
subpopulations of macrophages. (d) Heat map showing the average expression of ferroptosis-related
genes in 2 subpopulations of macrophages. (e) t-SNE plot showing the expression patterns of marker
genes in M_C2. (f) t-SNE plot showing the expression patterns of marker genes in M_C1. (g) t-SNE
plot showing the enrichment score of the TREM2+MAC in each cell. (h) t-SNE plot showing the
enrichment score of the FOLR2+MAC in each cell. (i) Bubble plot showing the correlation of
enrichment score between TREM2+MAC and FOLR2+MAC. (j) Network diagram of the cell-cell
communication between cancer epithelial cells and ferroptosis-related subpopulations of macrophages.

3.4 Construction and validation of predictive survival models based on ferroptosis-related genes

Based on the RFS data of TNBC in the GSE25066 database, univariate Cox regression analysis

was performed on 8371 specific markers of ferroptosis-related subpopulations, and 41 genes

significantly associated with RFS were screened (p < 0.01) (Supplementary Table 2,

Supplementary Figure 3). These genes were further analyzed by Least Absolute Shrinkage and

Selection Operator (LASSO) regression, and 23 genes were finally selected for the construction of

the risk model. We calculate the risk score using the following formula: risk score = TMEM160 *

(-0.4689) + EWSR1 * (-0.4237) + BCAT2 * (-0.2346) + PNKP * (-0.1592) + MLEC * (-0.1567) +

SAP30BP * (-0.1469) + TBL1XR1 * (-0.0894) + STAG1 * (-0.0695) + NR4A1 * (-0.0630) +

TNFRSF9 + (-0.0503) + PSD3 * (-0.0153) + BAD * (-0.0094) + MIIP * 0.0104 + HIST3H2A *

0.0294 + CDC25B * 0.0528 + TCEB1 * 0.0670 + HMGCS1 * 0.0988 + SPC25 * 0.1317 + TKT *

0.2234 + PTTG1 * 0.2960 + ADA * 0.3097 + AK1 * 0.4003 + AIMP2 * 0.4081 (Figure 4a-c). All

patients were categorized into low- and high-risk groups according to the median value of the risk

score (Figure 4d-f). Survival curves showed that patients in the high-risk group had a shorter RFS

compared to patients in the low-risk group (p<0.05, Figure 4g). In addition, the risk score had an

excellent predictive effect in predicting RFS in TNBC patients, its AUCs for 3-, 4-, and 5 years

RFS were 0.87, 0.88, and 0.88 (Figure 4h).

In external validation set (GSE86166), patients in the high-risk group also had a shorter RFS

compared to patients in the low-risk group (p<0.05). This risk model also performed well in

predicting RFS in TNBC patients, its AUCs for 3-, 4-, and 5 years RFS were 0.65, 0.67, and 0.71

(Figure 4i).

Fig. 4 Prognostic differences in triple-negative breast cancer with ferroptosis-related subpopulations.
(a) LASSO regression of 23 ferroptosis-related genes. (b) Cross-validation for optimizing the
parameter in LASSO regression. (c) Demonstration of regression coefficients corresponding to 23
genes. (d) Graph showing risk scores for all samples. (e) Scatterplot showing recurrence-free survival
for all samples. (f) Heat map showing the average expression of 23 genes in the low- and high-risk
groups. (g) Kaplan–Meier curves of survival analysis in the low- and high-risk groups. (h) Receiver
operating characteristic curves for predicting the recurrence-free survival at 3, 4 and 5 years in
training set. (i) Receiver operating characteristic curves for predicting the recurrence-free survival at 3,
4 and 5 years in external validation set.

3.5 Analysis of independent prognostic factors

Further, we performed univariate and multivariate Cox analyses to determine whether the risk

score could serve as an independent prognostic factor for TNBC patients compared to other

common clinicopathologic factors. In the GSE25066 database, Figure 5a shows that both the risk

factor and stage were significantly associated with TNBC patients' RFS and were independent

prognostic factors (p<0.05). In the external validation set, Figure 5b shows that only the risk factor

can be considered as independent prognostic factor in TNBC patients (p<0.05).

Fig. 5 Analysis of independent prognostic factors for triple-negative breast cancer patients. (a) The
forest plot showing the results of univariate and multivariate COX regression analysis of risk score,
age, grade, and stage in the GSE25066 database. (b) The forest plot showing the results of univariate
and multivariate COX regression analysis of risk score, grade, and stage in the GSE86166 database.

3.6 Correlation of the risk score with immune microenvironment

Next, we calculated the ImmuneScore, StromalScore, ESTIMATEScore, and TumorPurity for

samples in the low- and high-risk groups and compared the differences in scores between the two

groups. The results showed that these scores did not differ significantly between the low- and

high-risk groups (Figure 6a-f). Then, based on the CIBERSORT algorithm, we found that T cells

CD4 memory activated, NK cells resting, and Monocytes had significantly higher proportions in

the high-risk group, and the NK cells activated, and mast cells resting had significantly higher

proportions in the low-risk group (Figure 6g). Infiltration of these immune cells may play an

important role in the clinical course of TNBC patients.

Fig. 6 Analysis of immune microenvironment in low- and high-risk groups. (a-d) Correlation of the risk
score with the ImmuneScore, StromalScore, ESTIMATEScore, and TumorPurity. (e) Difference of
ImmuneScore, StromalScore, and ESTIMATEScore in low- and high-risk groups. (f) Difference of
TumorPurity in low- and high-risk groups. (g) Difference of immune infiltration score between low-
and high-risk groups calculated by CIBERSORT. * P < 0.05, ** P < 0.01, ns: p>0.05.

3.7 Analysis of clinical response to drugs with the risk score

Based on the "pRRophetic" R package, we explored the relationship between the risk score and

clinical response to 138 drugs and calculated IC50. As a result, 16 of 50 drugs whose clinical

response was significantly associated with the risk score were positively associated with the risk

score, and the remaining 34 were negatively associated with the risk score. Further, we found that

27 of drugs negatively associated with the risk score had significantly lower IC50 in the high-risk

group, suggesting that patients in the high-risk group may be more sensitive to these drugs,

favoring the choice of clinical medication (Figure 7a). And 13 of drugs positively associated with

the risk score had significantly higher IC50 in the high-risk group than in the low-risk group,

suggesting that patients in the high-risk group may be less sensitive to these drugs (Figure 7b).

Finally, we calculated the TIDE scores of patients in the low- and high-risk groups; unfortunately,

the TIDE scores were not significantly different between the two groups (Figure 7c).

Fig. 7 Analysis of immune microenvironment in low- and high-risk groups. (a) Box plot showing
differences of IC50 for drugs negatively associated with risk scores in different groups. (b) Box plot
showing differences of IC50 for drugs positively associated with risk scores in different groups. (c) Box
*plot showing differences of TIDE scores in different groups. * P < 0.05, ** P < 0.01, *** P < 0.001,*
***** P < 0.0001, ns: p>0.05.*

Discussion

TNBC is the most difficult subtype of breast cancer to treat, and survival and efficacy prediction

for each individual is a critical step in precision tumor therapy. Several studies have identified

ferroptosis-related genes as novel therapeutic targets to enhance treatment efficacy and improve

patient prognosis [44-47]. However, to the best of our knowledge, there is a lack of reports on the

association of ferroptosis-related genes with prognosis and clinical response to treatment in TNBC.

Here, firstly, we revealed ferroptosis-related immune cell clustering in TNBC patients at the

single-cell level, demonstrating the ferroptosis heterogeneity in TNBC, which was also supported

by other cohorts [19]. On the basis of the above results, we constructed an ferroptosis-related risk

score model based on bulk RNA-seq data, which successfully predicted the recurrence-free

survival of TNBC patients and validated the robustness of the results in various datasets. This is

important for the individualized management of TNBC patients. More importantly, the model can

also predict the sensitivity to drugs for TNBC patients with different risk stratification to guide

physicians in the selection of drugs for different patients, and it has significant clinical application

value.

Understanding the TME in TNBC is essential for improving prognosis and guiding treatment.

scRNA-seq enables transcriptomic analysis of individual cells to characterize the cellular diversity

in the TME in detail, thereby improving understanding of the disease [48-50]. In this study, our

analysis of scRNA-seq data from TNBC revealed that there are multiple immune cells infiltrating

in the TME and interacting frequently with each other, and they may impede or promote tumor

progression [51]. Since, we found that the average expression of ferroptosis-related genes was

significantly different in different T-stages of tumors, which might be strongly correlated with

patient prognosis. Further, based on ferroptosis-related genes, a higher proportion of T cells and

macrophages in the TME were categorized into different subpopulations to reveal the

heterogeneity of ferroptosis-related in TNBC. The T_C2 subpopulation was accompanied by a

greater infiltration of NK cells. NK cells have rapid and efficient anti-tumor immunity and their

activity is negatively correlated with breast cancer progression [52-54]. The relevant results were

validated in the dataset GSE25066. TFs are universal regulators of the transcription of many genes

and are closely associated with the progression of cancer [55]. The results indicate that TBX21,

which exhibits higher activity in NKT cells, is associated with the expression of SLC7A11, and

overexpression of SLC7A11 alters the expression of chemokines, resulting in the infiltration of

immune cells such as CD8$^+$ T cells and neutrophils [56]. It has been reported that RUNX3 induces

ferroptosis by activating the ING1/p53/SLC7A11 signaling pathway, thereby inhibiting the growth

of gallbladder cancer [57]. Furthermore, PRDM1 and IRF, which exhibit higher activity in CD8$^+$ T

cells, induce ferroptosis in cancer cells by inhibiting the transcription of GPX4 and driving the

expression of transferrin receptor, respectively, thereby improving the TME [58-59]. FOLR2, SEPP1,

MRC1, LYVE1, SLC40A1 and CD163 were highly expressed in the M_C1 subpopulation. Studies

have shown that these markers, which are normally present in mammary macrophages from

healthy humans and mice, correlate with a favorable prognosis in breast cancer [60-62]. Among them,

FOLR2 is positively correlated with antitumor immune players of CD8$^+$ T cells, DCs, B cells, and

tertiary lymphoid structures, and there is a strong correlation between FOLR2 expression and

various immune pathways including T-cell receptor and PD-1 signaling, as well as antigen

processing. FOLR2$^+$ macrophages are an important component in the initiation of antitumor

immunity [62]. In the M_C2 subpopulation, TREM2, FN1, C3, CXCR4, and SPP1 are highly

expressed. It has been shown that these markers are lowly expressed in macrophages of healthy

breast tissues and highly expressed in tumor tissues [62-64]. Specifically, the absence of TREM2

reshapes macrophage infiltration, fosters the enrichment and activation of T cells and NK cells,

and is correlated with an enhanced response to checkpoint blockade therapy [63]. In addition,

S100A8 and IFI6 were also highly expressed in the M_C2 subpopulation. Studies have reported

that a high percentage of S100A8[+] myeloid cell infiltration has been suggested as a potential

mechanism for worsening the prognosis of TNBC [65-66]. Furthermore, IFI6 is an interferon-

stimulated gene with anti-apoptotic and metastasis-promoting effects [67]. These findings

demonstrate the heterogeneity of the function of iron death-associated macrophage subpopulations,

which may play different roles in the anti-tumor process. In the original literature, Prabhakaran S

et al. [35] classified breast cancer into nine clusters (Ecotypes) based on single-cell characteristics,

which exhibited distinct cellular compositions and clinical outcomes. Among them, Ecotypes-2 is

primarily composed of LumA and Normal-like tumors, and patients in this group have the best

prognosis, whereas Ecotype-3 is enriched with Basal_SC, Cycling, Luminal_Progenitor, and a

Basal bulk PAM50 subtype, and patients in this group have the worst prognosis. In comparison,

this study focuses on TNBC, which has a poorer prognosis. The different distributions of T cell

and macrophage subtypes in different TNBC patients the study results revealed that ferroptosis-

related TME in TNBC patients may be valuable for patient survival and outcome prediction. In the

future, altering the degree of macrophage or T-cell infiltration in different subpopulations could be

potentially valuable in improving the prognosis of TNBC patients.

Intratumor heterogeneity is one of the most important factors contributing to poor clinical

outcomes in breast cancer, and ferroptosis is one of the major pathways of activation in highly

heterogeneous tumors [68-70]. In order to better address the problem of accurately predicting the

prognosis of TNBC, we firstly performed the correlation analysis of ferroptosis-related genes with

RFS and constructed a risk score model in the publicly available dataset of bulk RNA-seq. TNBC

patients were categorized into low- and high-risk groups based on the risk score. In the high-risk

group, TNBC patients had higher expression levels of the MIIP, HIST3H2A, CDC25B, TCEB1,

HMGCS1, SPC25, TKT, PTTG1, ADA, AK1, and AIMP2 genes. Among them, the cell division

cycle 25 (CDC25) family of proteins is dual-specific tyrosine phosphatases responsible including

three subtypes, CDC25A, CDC25B and CDC25C, which are used to regulate cell cycle transitions

[71-72]. CDC25B dephosphorylates and activates cyclin-dependent kinase/cyclin complex

(CDK1/Cyclin B), which have been shown to be overexpressed in breast cancer and promote cell

cycle progression [72-73]. Spindle component 25 (SPC25) is a key component of the nuclear

division cycle 80 (NDC80) complex, and its high expression promotes tumor cell proliferation by

inducing mitotic disorder [74]. Study shows that SPC25 promotes the proliferation of breast cancer

cells, and high levels of SPC25 mRNA levels are associated with high recurrence rates and low

survival rates in breast cancer patients [75]. Pituitary tumor transforming gene 1 (PTTG1) has been

demonstrated to promote proliferation, migration and invasion of cancer cells and is a gene that

promotes breast cancer development [76]. At the mRNA level, PTTG1 is more present in patients

with high histologic grade and with lymph node metastasis, is significantly associated with low

patient survival, and is a biomarker suggestive of poor prognosis in breast cancer, which may be

related to the regulation of the cell cycle by PTTG1 to promote the development of breast cancer

cells [77]. In the low-risk group, TNBC patients had higher expression levels of the TMEM160,

EWSR1, BCAT2, PNKP, MLEC, SAP30BP, TBL1XR1, STAG1, NR4A1, TNFRSF9, PSD3, and

BAD genes. These genes have different roles in regulating the TME in colorectal cancers, bladder

cancers, lymphomas, etc., but they have been rarely reported in TNBC [78-80]. In future studies,

these genes are valuable to investigate in the regulation of anti-TNBC.

In TNBC, we found that the risk score model we constructed had independent predictive power

for RFS and the predictive performance of this score was stable in an external validation set.

Considering the impact of the tumor immune microenvironment on the prognosis of TNBC

patients, we further explored the differences in immune cell infiltration in different subgroups of

patients. The results showed that T cells CD4 memory activated, NK cells resting, and Monocytes

had higher proportions in the high-risk group. This is in line with previous studies, such as that

CXCL7 expressed and released by monocytes can stimulate cancer cell migration, invasion and

metastasis [81]. While in the low-risk group, the proportion of NK cells activated and mast cells

resting was higher. It has been suggested that the effect of mast cells on tumor invasiveness may

vary among different breast cancer subtypes [82-83]. Our findings suggest that infiltration of mast

cells resting in the TME is more favorable to the prognosis of patients.

For high-risk TNBC patients, it is crucial to seek more effective drugs for adjuvant therapy in

clinical practice. In our study, we found that ABT-263 (navitoclax) and erlotinib exhibited the

greatest differences in drug sensitivity among different patient groups. ABT-263 is a mimetic of B-

cell lymphoma-2 (BCL-2) homology 3 (BH3) that can increase cellular reactive oxygen species

(ROS) levels and induce TNBC cell apoptosis by inhibiting the function of anti-apoptotic proteins

(BCL-2, BCL-XL, and BCL-W) [84-85]. ROS is a key regulatory factor in the occurrence of

ferroptosis, and excessive production of ROS can lead to the accumulation of lipid peroxides,

ultimately triggering cellular ferroptosis [86-87]. In addition, erlotinib is an epidermal growth factor

receptor (EGFR) tyrosine kinase inhibitor (TKI) that exerts its effects by competitively binding to

the ATP site within the catalytic domain and inhibiting the phosphorylation of EGFR, thereby

improving the progression-free survival of patients with EGFR mutations [88-89]. Studies have

shown that β-Elemene can enhance the sensitivity of EGFR-mutated non-small cell lung cancer to

erlotinib by upregulating lncRNA H19 and inducing ferroptosis, providing a new approach to

overcoming drug resistance in TNBC [90]. EGFR is a receptor tyrosine kinase that promotes the

proliferation and invasion of breast cancer by stimulating multiple oncogenic pathways such as

Ras-Raf-MEK-ERK, PI3K-AKT-mTOR, and Src-STAT3 [91]. The high sensitivity of high-risk

patients to erlotinib may be attributed to the higher expression of EGFR in their tumors. Currently,

the treatment of TNBC mainly relies on systemic chemotherapy, while small molecule inhibitors

and targeted drugs induce cell death through various pathways, holding great promise for anti-

tumor treatment in high-risk patient groups.

This study has some limitations. First, this study is based on the analysis of transcriptomic data,

and the integration of multi-omics data is needed to improve the predictive performance of the

model. Second, this study is a retrospective analysis of a public dataset, and future in vivo, in vitro

experiments as well as prospective clinical trials are needed to further validate the research

findings.

Conclusions

In conclusion, this study revealed the TNBC ferroptosis-mediated tumor immune cell clustering

based on scRNA-seq data, and the degree of infiltration of different subpopulations of cells had

different roles in the prognosis of TNBC patients. On this basis, combined with bulk RNA-seq

data, the survival prediction model for TNBC patients was established with excellent predictive

performance and stability. The high-risk TNBC patients screened by this prediction model and

their sensitive therapeutic drugs can help to guide physicians in disease monitoring and precise

treatment. The related genes screened also provide important value for TNBC patients to find

potential therapeutic targets to improve their prognosis.

Acknowledgements

Our thanks go to the Gene Expression Omnibus (GEO) database and FerrDb for providing

essential data resources. We extend our appreciation to the participating institutions and our

dedicated colleagues for their contributions and expertise. Special recognition is given to the

patients and clinical staff for their invaluable participation and assistance. We also thank the

reviewers for their insightful feedback.

List of abbreviations

Triple-negative breast cancer (TNBC); tumor microenvironment (TME); RNA sequencing (RNA-

seq); single-cell RNA-seq (scRNA-seq); Area Under the Curves(AUCs); Gene Expression

Omnibus (GEO); The Cancer Genome Atlas (TCGA); quality control (QC); highly variable genes

(HVGs); principal component analysis (PCA); t-distributed stochastic neighbor embedding (t-

SNE); Non-negative matrix decomposition (NMF); Progenitor Cell Biology Consortium (PCBC);

one-class logistic regression (OCLR); Single-cell Regulatory Network Inference and Clustering

tool (SCENIC); transcription factor (TF); recurrence-free survival (RFS); Least Absolute

Shrinkage and Selection Operator (LASSO); receiver operating characteristic (ROC); tumor

immune dysfunction and exclusion (TIDE); Cancer-associated Fibroblasts (CAFs); Perivascular-

like Cells (PVL); Dendritic Cells (DCs); Least Absolute Shrinkage and Selection

Operator(LASSO); cell division cycle 25 (CDC25); cyclin-dependent kinase/cyclin complex

(CDK1/Cyclin B); Spindle component 25 (SPC25); nuclear division cycle 80 (NDC80); Pituitary

tumor transforming gene 1 (PTTG1); B-cell lymphoma-2 (BCL-2); B-cell lymphoma-2 homology

3 (BH3); Reactive oxygen species (ROS); Epidermal growth factor receptor (EGFR); Tyrosine

kinase inhibitor (TKI).

References

1. Bray F, Laversanne M, Sung H, et al. Global cancer statistics 2022: GLOBOCAN estimates of

incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin.

2024;74(3):229-263.

2. Keenan TE, Tolaney SM. Role of Immunotherapy in Triple-Negative Breast Cancer. J Natl

Compr Canc Netw. 2020;18(4):479-489.

3. Yam C, Yen EY, Chang JT, et al. Immune Phenotype and Response to Neoadjuvant Therapy in

Triple-Negative Breast Cancer. Clin Cancer Res. 2021;27(19):5365-5375.

4. Howlader N, Cronin KA, Kurian AW, Andridge R. Differences in Breast Cancer Survival by

Molecular Subtypes in the United States. Cancer Epidemiol Biomarkers Prev.

2018;27(6):619-626.

5. Bardia A, Mayer IA, Vahdat LT, et al. Sacituzumab Govitecan-hziy in Refractory Metastatic

Triple-Negative Breast Cancer. N Engl J Med. 2019;380(8):741-751.

6. Cortazar P, Zhang L, Untch M, et al. Pathological complete response and long-term clinical

benefit in breast cancer: the CTNeoBC pooled analysis [published correction appears in

Lancet. 2019 Mar 9;393(10175):986]. Lancet. 2014;384(9938):164-172.

7. Shepherd JH, Ballman K, Polley MC, et al. CALGB 40603 (Alliance): Long-Term Outcomes

and Genomic Correlates of Response and Survival After Neoadjuvant Chemotherapy With or

Without Carboplatin and Bevacizumab in Triple-Negative Breast Cancer. J Clin Oncol.

2022;40(12):1323-1334.

8. Kim C, Gao R, Sei E, et al. Chemoresistance Evolution in Triple-Negative Breast Cancer

Delineated by Single-Cell Sequencing. Cell. 2018;173(4):879-893.e13.

9. Mayer S, Milo T, Isaacson A, et al. The tumor microenvironment shows a hierarchy of cell-

cell interactions dominated by fibroblasts. Nat Commun. 2023;14(1):5810.

10. Mao X, Xu J, Wang W, et al. Crosstalk between cancer-associated fibroblasts and immune

cells in the tumor microenvironment: new findings and future perspectives. Mol Cancer.

2021;20(1):131.

11. Binnewies M, Roberts EW, Kersten K, et al. Understanding the tumor immune

microenvironment (TIME) for effective therapy. Nat Med. 2018;24(5):541-550.

12. de Visser KE, Joyce JA. The evolving tumor microenvironment: From cancer initiation to

metastatic outgrowth. Cancer Cell. 2023;41(3):374-403.

13. Yang W, Xu M, Xu S, et al. Single-cell RNA reveals a tumorigenic microenvironment in the

interface zone of human breast tumors. Breast Cancer Res. 2023;25(1):100.

14. Fang X, Ardehali H, Min J, Wang F. The molecular and metabolic landscape of iron and

ferroptosis in cardiovascular disease. Nat Rev Cardiol. 2023;20(1):7-23.

15. Viswanathan VS, Ryan MJ, Dhruv HD, et al. Dependency of a therapy-resistant state of

cancer cells on a lipid peroxidase pathway. Nature. 2017;547(7664):453-457.

16. Kim R, Hashimoto A, Markosyan N, et al. Ferroptosis of tumour neutrophils causes immune

suppression in cancer. Nature. 2022;612(7939):338-346.

17. Hangauer MJ, Viswanathan VS, Ryan MJ, et al. Drug-tolerant persister cancer cells are

vulnerable to GPX4 inhibition. Nature. 2017;551(7679):247-250.

18. Tsoi J, Robert L, Paraiso K, et al. Multi-stage Differentiation Defines Melanoma Subtypes

with Differential Vulnerability to Drug-Induced Iron-Dependent Oxidative Stress. Cancer Cell.

2018;33(5):890-904.e5.

19. Yang F, Xiao Y, Ding JH, et al. Ferroptosis heterogeneity in triple-negative breast cancer

reveals an innovative immunotherapy combination strategy. Cell Metab. 2023;35(1):84-

100.e8.

20. Lu YJ, Gong Y, Li WJ, Zhao CY, Guo F. The prognostic significance of a novel ferroptosis-

related gene model in breast cancer. Ann Transl Med. 2022;10(4):184.

21. Hong M, Tao S, Zhang L, et al. RNA sequencing: new technologies and applications in cancer

research. J Hematol Oncol. 2020;13(1):166.

22. Kuksin M, Morel D, Aglave M, et al. Applications of single-cell and bulk RNA sequencing in

onco-immunology. Eur J Cancer. 2021;149:193-210.

23. Li F, Zhang H, Huang Y, et al. Single-cell transcriptome analysis reveals the association

between histone lactylation and cisplatin resistance in bladder cancer. Drug Resist Updat.

2024;73:101059.

24. Ma P, Amemiya HM, He LL, et al. Bacterial droplet-based single-cell RNA-seq reveals

antibiotic-associated heterogeneous cellular states. Cell. 2023;186(4):877-891.e14.

25. Chen J, Wang X, Ma A, et al. Deep transfer learning of cancer drug responses by integrating

bulk and single-cell RNA-seq data. Nat Commun. 2022;13(1):6494.

26. Janjic A, Wange LE, Bagnoli JW, et al. Prime-seq, efficient and powerful bulk RNA

sequencing. Genome Biol. 2022;23(1):88.

27. Guan C, Gong A, Zhao Y, et al. Interpretable machine learning model for new-onset atrial

fibrillation prediction in critically ill patients: a multi-center study. Crit Care. 2024;28(1):349.

28. Ching T, Zhu X, Garmire LX. Cox-nnet: An artificial neural network method for prognosis

prediction of high-throughput omics data. PLoS Comput Biol. 2018;14(4):e1006076.

29. Zhao H, Zhou X, Wang G, et al. Integrating Bulk and Single-cell RNA-seq to Construct a

Macrophage-related Prognostic Model for Prognostic Stratification in Triple-negative Breast

Cancer. J Cancer. 2024;15(18):6002-6015.

30. Pei S, Zhang P, Chen H, et al. Integrating single-cell RNA-seq and bulk RNA-seq to construct

prognostic signatures to explore the role of glutamine metabolism in breast cancer. Front

Endocrinol (Lausanne). 2023;14:1135297.

31. Lei H, Guo XA, Tao Y, et al. Semi-deconvolution of bulk and single-cell RNA-seq data with

application to metastatic progression in breast cancer. Bioinformatics. 2022;38(Suppl 1):i386-

i394.

32. Gao Y, Feng Z, Zhao H, et al. Integrating single-cell RNA-seq and bulk RNA-seq to explore

prognostic value and immune landscapes of methionine metabolism-related signature in breast

cancer. Front Genet. 2025;15:1521269.

33. Wu SZ, Al-Eryani G, Roden DL, et al. A single-cell and spatially resolved atlas of human

breast cancers. Nat Genet. 2021;53(9):1334-1347.

34. Hatzis C, Pusztai L, Valero V, et al. A genomic predictor of response and survival following

taxane-anthracycline chemotherapy for invasive breast cancer. JAMA. 2011;305(18):1873-

1881.

35. Prabhakaran S, Rizk VT, Ma Z, et al. Evaluation of invasive breast cancer samples using a 12-

chemokine gene expression score: correlation with clinical outcomes. Breast Cancer Res.

2017;19(1):71.

36. Ding X, Qin J, Huang F, Feng F, Luo L. The combination of machine learning and untargeted

metabolomics identifies the lipid metabolism -related gene CH25H as a potential biomarker in

asthma. Inflamm Res. 2023;72(5):1099-1119.

37. Efremova M, Vento-Tormo M, Teichmann SA, Vento-Tormo R. CellPhoneDB: inferring cell-

cell communication from combined expression of multi-subunit ligand-receptor complexes.

Nat Protoc. 2020;15(4):1484-1506.

38. Sugihara R, Kato Y, Mori T, Kawahara Y. Alignment of single-cell trajectory trees with

CAPITAL. Nat Commun. 2022;13(1):5972.

39. Aibar S, González-Blas CB, Moerman T, et al. SCENIC: single-cell regulatory network

inference and clustering. Nat Methods. 2017;14(11):1083-1086.

40. Van de Sande B, Flerin C, Davie K, et al. A scalable SCENIC workflow for single-cell gene

regulatory network analysis. Nat Protoc. 2020;15(7):2247-2276.

41. Iglesias-Escudero M, Arias-González N, Martínez-Cáceres E. Regulatory cells and the effect

of cancer immunotherapy. Mol Cancer. 2023;22(1):26. Published 2023 Feb 4.

42. Pérez-Romero K, Rodríguez RM, Amedei A, Barceló-Coblijn G, Lopez DH. Immune

Landscape in Tumor Microenvironment: Implications for Biomarker Development and

Immunotherapy. Int J Mol Sci. 2020;21(15):5521.

43. Gong TQ, Jiang YZ, Shao C, et al. Proteome-centric cross-omics characterization and

integrated network analyses of triple-negative breast cancer. Cell Rep. 2022;38(9):110460.

44. Li H, Yang P, Wang J, et al. HLF regulates ferroptosis, development and chemoresistance of

triple-negative breast cancer by activating tumor cell-macrophage crosstalk. J Hematol Oncol.

2022;15(1):2.

45. Zhang Z, Lu M, Chen C, et al. Holo-lactoferrin: the link between ferroptosis and radiotherapy

in triple-negative breast cancer. Theranostics. 2021;11(7):3167-3182. Published 2021 Jan 1.

46. Zhou TJ, Zhang MM, Liu DM, et al. Glutathione depletion and dihydroorotate dehydrogenase

inhibition actuated ferroptosis-augment to surmount triple-negative breast cancer.

Biomaterials.

47. Desterke C, Cosialls E, Xiang Y, et al. Adverse Crosstalk between Extracellular Matrix

Remodeling and Ferroptosis in Basal Breast Cancer. Cells. 2023;12(17):2176.

48. Li J, Hubisz MJ, Earlie EM, et al. Non-cell-autonomous cancer progression from

chromosomal instability. Nature. 2023;620(7976):1080-1088.

49. Van de Sande B, Lee JS, Mutasa-Gottgens E, et al. Applications of single-cell RNA

sequencing in drug discovery and development. Nat Rev Drug Discov. 2023;22(6):496-520.

50. Zhai Y, Zhang J, Huang Z, et al. Single-cell RNA sequencing integrated with bulk RNA

sequencing analysis reveals diagnostic and prognostic signatures and immunoinfiltration in

gastric cancer. Comput Biol Med. 2023;163:107239.

51. Deepak KGK, Vempati R, Nagaraju GP, et al. Tumor microenvironment: Challenges and

opportunities in targeting metastasis of triple negative breast cancer. Pharmacol Res.

2020;153:104683.

52. Gong Y, Chen W, Chen X, et al. An Injectable Epigenetic Autophagic Modulatory Hydrogel

for Boosting Umbilical Cord Blood NK Cell Therapy Prevents Postsurgical Relapse of Triple-

Negative Breast Cancer. Adv Sci (Weinh). 2022;9(23):e2201271.

53. Guillerey C, Huntington ND, Smyth MJ. Targeting natural killer cells in cancer

immunotherapy. Nat Immunol. 2016;17(9):1025-1036.

54. Bouzidi L, Triki H, Charfi S, et al. Prognostic Value of Natural Killer Cells Besides Tumor-

Infiltrating Lymphocytes in Breast Cancer Tissues. Clin Breast Cancer. 2021;21(6):e738-e747.

55. Wang S, Chen J, Li P, Chen Y. LINC01133 can induce acquired ferroptosis resistance by

enhancing the FSP1 mRNA stability through forming the LINC01133-FUS-FSP1 complex.

Cell Death Dis. 2023;14(11):767.

56. Cheng X, Wang Y, Liu L, Lv C, Liu C, Xu J. SLC7A11, a Potential Therapeutic Target

Through Induced Ferroptosis in Colon Adenocarcinoma. Front Mol Biosci. 2022;9:889688.

57. Cai C, Zhu Y, Mu J, et al. DNA methylation of RUNX3 promotes the progression of

gallbladder cancer through repressing SLC7A11-mediated ferroptosis. Cell Signal.

2023;108:110710.

58. Wu W, Ke W, Shi W, et al. PR/SET domain 1 targeting glutathione peroxidase 4 regulates

chronic hepatitis B liver fibrosis through ferroptosis. Cytojournal. 2024;21:78.

59. Song A, Wu L, Zhang BX, et al. Glutamine inhibition combined with CD47 blockade

enhances radiotherapy-induced ferroptosis in head and neck squamous cell carcinoma. Cancer

Lett. 2024;588:216727.

60. Jäppinen N, Félix I, Lokka E, et al. Fetal-derived macrophages dominate in adult mammary

glands. Nat Commun. 2019;10(1):281.

61. Wang Y, Chaffee TS, LaRue RS, et al. Tissue-resident macrophages promote extracellular

matrix homeostasis in the mammary gland stroma of nulliparous mice. Elife. 2020;9:e57438.

62. Nalio Ramos R, Missolo-Koussou Y, Gerber-Ferder Y, et al. Tissue-resident FOLR2+

macrophages associate with CD8+ T cell infiltration in human breast cancer. Cell.

2022;185(7):1189-1207.e25.

63. Molgora M, Esaulova E, Vermi W, et al. TREM2 Modulation Remodels the Tumor Myeloid

Landscape Enhancing Anti-PD-1 Immunotherapy. Cell. 2020;182(4):886-900.e17.

64. Müller A, Homey B, Soto H, et al. Involvement of chemokine receptors in breast cancer

metastasis. Nature. 2001;410(6824):50-56.

65. Drews-Elger K, Iorns E, Dias A, et al. Infiltrating S100A8+ myeloid cells promote metastatic

spread of human breast cancer and predict poor clinical outcome. Breast Cancer Res Treat.

2014;148(1):41-59.

66. Rigiracciolo DC, Nohata N, Lappano R, et al. Focal Adhesion Kinase (FAK)-Hippo/YAP

transduction signaling mediates the stimulatory effects exerted by S100A8/A9-RAGE system

in triple-negative breast cancer (TNBC). J Exp Clin Cancer Res. 2022;41(1):193.

67. Davenport AM, Morris M, Sabti F, et al. G1P3/IFI6, an interferon stimulated protein,

promotes the association of RAB5+ endosomes with mitochondria in breast cancer cells. Cell

Biol Int. 2023;47(11):1868-1879.

68. Zardavas D, Irrthum A, Swanton C, Piccart M. Clinical management of breast cancer

heterogeneity. Nat Rev Clin Oncol. 2015;12(7):381-394.

69. Su GH, Xiao Y, You C, et al. Radiogenomic-based multiomic analysis reveals imaging

intratumor heterogeneity phenotypes and therapeutic targets. Sci Adv. 2023;9(40):eadf0837.

70. Wu SZ, Al-Eryani G, Roden DL, et al. A single-cell and spatially resolved atlas of human

breast cancers. Nat Genet. 2021;53(9):1334-1347.

71. Boutros R, Lobjois V, Ducommun B. CDC25 phosphatases in cancer cells: key players? Good

targets?. Nat Rev Cancer. 2007;7(7):495-507.

72. Cairns J, Ly RC, Niu N, et al. CDC25B partners with PP2A to induce AMPK activation and

tumor suppression in triple negative breast cancer. NAR Cancer. 2020;2(4):zcaa039.

73. Guo YQ, Wang Q, Wang JG, et al. METTL3 modulates m6A modification of CDC25B and

promotes head and neck squamous cell carcinoma malignant progression. Exp Hematol Oncol.

2022;11(1):14.

74. Yang J, Huang Y, Song M, et al. SPC25 promotes proliferation and stemness of hepatocellular

carcinoma cells via the DNA-PK/AKT/Notch1 signaling pathway. Int J Biol Sci.

2022;18(14):5241-5259.

75. Wang Q, Zhu Y, Li Z, et al. Up-regulation of SPC25 promotes breast cancer. Aging (Albany

NY). 2019;11(15):5689-5704.

76. Qi L, Zhou B, Chen J, et al. Significant prognostic values of differentially expressed-

aberrantly methylated hub genes in breast cancer. J Cancer. 2019;10(26):6618-6634.

77. Meng C, Zou Y, Hong W, Bao C, Jia X. Estrogen-regulated PTTG1 promotes breast cancer

progression by regulating cyclin kinase expression. Mol Med. 2020;26(1):33.

78. Dai X, Wu Z, Ruan R, et al. TMEM160 promotes tumor immune evasion and radiotherapy resistance via PD-L1 binding in colorectal cancer. Cell Commun Signal. 2024;22(1):168.

79. Lei MZ, Li XX, Zhang Y, et al. Acetylation promotes BCAT2 degradation to suppress BCAA catabolism and pancreatic cancer growth. Signal Transduct Target Ther. 2020;5(1):70.

80. Venturutti L, Teater M, Zhai A, et al. TBL1XR1 Mutations Drive Extranodal Lymphoma by Inducing a Pro-tumorigenic Memory Fate. Cell. 2020;182(2):297-316.e27.

81. Wang YH, Shen CY, Lin SC, et al. Monocytes secrete CXCL7 to promote breast cancer progression. Cell Death Dis. 2021;12(12):1090. Published 2021 Nov 17.

82. Bense RD, Sotiriou C, Piccart-Gebhart MJ, et al. Relevance of Tumor-Infiltrating Immune Cell Composition and Functionality for Disease Outcome in Breast Cancer. J Natl Cancer Inst. 2016;109(1):djw192.

83. Majorini MT, Cancila V, Rigoni A, et al. Infiltrating Mast Cell-Mediated Stimulation of Estrogen Receptor Activity in Breast Cancer Cells Promotes the Luminal Phenotype. Cancer Res. 2020;80(11):2311-2324.

84. Oh Y, Jung HR, Min S, et al. Targeting antioxidant enzymes enhances the therapeutic efficacy of the BCL-XL inhibitor ABT-263 in KRAS-mutant colorectal cancers. Cancer Lett. 2021;497:123-136.

85. Lee A, Jin HO, Masudul Haque M, et al. Synergism of a novel MCL￿1 downregulator, acriflavine, with navitoclax (ABT￿263) in triple￿negative breast cancer, lung adenocarcinoma and glioblastoma multiforme. Int J Oncol. 2022;60(1):2.

86. Stockwell BR. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and

therapeutic applications. Cell. 2022;185(14):2401-2421.

87. Shi X, Xu T, Gao M, et al. Combined exposure of emamectin benzoate and microplastics

induces tight junction disorder, immune disorder and inflammation in carp midgut via

lysosome/ROS/ferroptosis pathway. Water Res. 2024;257:121660.

88. Yue D, Xu S, Wang Q, et al. Erlotinib versus vinorelbine plus cisplatin as adjuvant therapy in

Chinese patients with stage IIIA EGFR mutation-positive non-small-cell lung cancer (EVAN):

a randomised, open-label, phase 2 trial. Lancet Respir Med. 2018;6(11):863-873.

89. Roberts MS, Anstine LJ, Finke VS, et al. KLF4 defines the efficacy of the epidermal growth

factor receptor inhibitor, erlotinib, in triple-negative breast cancer cells by repressing the

EGFR gene. Breast Cancer Res. 2020;22(1):66.

90. Xu C, Jiang ZB, Shao L, et al. β-Elemene enhances erlotinib sensitivity through induction of

ferroptosis by upregulating lncRNA H19 in EGFR-mutant non-small cell lung cancer.

Pharmacol Res. 2023;191:106739.

91. Wee P, Wang Z. Epidermal Growth Factor Receptor Cell Proliferation Signaling Pathways.

Cancers (Basel). 2017;9(5):52.

• **Additional files File:** Appendix 1-figure 1.TIF. Appendix 1-figure 2.TIF. Appendix 1-figure

3_revised.TIF. Appendix 1-figure 4_revised.TIF. Appendix 1-figure 5.TIF. Appendix 1-figure

6.TIF. Appendix 1-figure 7.TIF. Appendix 1-Supplementary Figure 1.TIF. Appendix 1-

Supplementary Figure 2.TIF. Appendix 1-Supplementary Figure 3.TIF. Figure 1–figure

supplement.pdf. Figure 2–figure supplement.pdf. Figure 3–figure supplement.pdf. Figure 4–figure

supplement.pdf. Figure 5–figure supplement.pdf. Figure 6–figure supplement.pdf. Figure 7–figure

supplement.pdf. Supplementary Figure 1-figure supplement. Supplementary Figure 2-figure

supplement. Supplementary Figure 3-figure supplement. Appendix 2-ICMJE.zip. Appendix 3-

collaborators.xlsx. Supplementary Table 1. Supplementary Table 2.

a

b