# Hepatic Nrf1 (*Nfe2l1*) promotes VLDL dependent liver defense against sepsis

Michael J. Trites[1], Lei Li[1], May G. Akl[1], Aidan Hydomako[1], and Scott B. Widenmaier[1*]

[1] Department of Anatomy, Physiology, and Pharmacology, University of Saskatchewan, Saskatoon, SK, Canada.

* Corresponding author information
Scott B. Widenmaier
107 Wiggins Road, Health Sciences Building
Saskatoon, SK S7N 5E5, Canada
Phone: (306)966-8320
Email: scott.widenmaier@usask.ca

**Keywords:** Nrf1, Nfe2l1, liver, hepatocyte, disease tolerance, triglyceride, sepsis, stress defense

**Short title: Hepatic Nrf1 protects against sepsis**

# Abstract

Sepsis is a dysregulated inflammatory condition that causes mortality by triggering organ damage and dysfunction. Interest has emerged in stimulating disease tolerance to reduce organ damage and preserve organ function. Liver plays a role in disease tolerance by mediating metabolic adaptations that defend against sepsis, but sepsis-induced liver damage may limit these effects. Here, we investigated whether stress defending transcription factors nuclear factor erythroid 2 related factor-1 (Nrf1) and -2 (Nrf2) in hepatocytes protect liver defenses against sepsis. Using mice, we evaluate responses by hepatic Nrf1 and Nrf2 to sepsis as well as genetically altered hepatic Nrf1 and Nrf2 activity and then injected these mice with LPS or *Escherichia coli* to determine whether hepatic Nrf1 and Nrf2 protect against sepsis. Our results show hepatic Nrf1 and Nrf2 activity is reduced in severe sepsis and that hepatic Nrf1, but not Nrf2, deficiency predisposes for hypothermia and mortality. In stark contrast, enhancing hepatic Nrf1 activity protects against hypothermia and improves survival. These effects were unrelated to circulating glucose, ketones, bile acids, and cytokines. Instead, we show in sepsis that hepatic Nrf1 deficiency reduces VLDL secretion and enhancing hepatic Nrf1 activity increases VLDL secretion, and that inhibiting VLDL secretion blocks hepatic Nrf1-mediated protection against hypothermia and sepsis severity. Gene expression profiles suggest Nrf1 may promote this effect by increasing hepatic stress defense programming. Hence, we show mortality in sepsis may result from impaired stress defense and that hepatic Nrf1 can improve disease tolerance by promoting VLDL dependent liver defense against sepsis.

# Introduction

Sepsis is a life-threatening organ dysfunction caused by a dysregulated host response to infection[1]. In hospital mortality is greater than 10 % for sepsis and 40 % for septic shock, and survivors have poor long-term prognosis[1-5]. While current treatment strategies focus on eliminating the infection (e.g., antibiotics) and hemodynamic stabilization (e.g., administer fluid and vasopressor), there is a recognized need for adjunct therapy that can mitigate organ damage and restore organ function to improve patient outcome[2,6]. During infection, the host has two defense strategies: resistance and tolerance[7,8]. Resistance is mediated by the immune system and involves inflammatory processes tasked with detecting and eliminating the pathogen. However, this destructive and energetically costly process exerts collateral tissue damage that, in sepsis, leads to declining organ function and fitness[9]. In contrast, tolerance is a strategy in which host defenses reduce susceptibility to tissue damage, irrespective of pathogen burden, resulting in preserved organ function and fitness[7-9]. The mechanisms underlying disease tolerance and how they may be harnessed to improve the outcome of patients with sepsis is an emerging area of great interest[2,6-10].

Metabolic adaptations that protect against organ damage underlie disease tolerance[9-13]. In sepsis models using bacteria or endotoxin, reduced tissue damage and improved survival has been linked to adaptive shifts in glucose, ketone, heme, lipid, lipoprotein, and energy metabolism[14-23], which are similar to adaptive shifts during starvation and regulated by reactive oxygen species (ROS) and hormones such as glucocorticoid, fibroblast growth factor 21 (FGF21), growth and differentiation factor 15 (GDF15), and growth hormone[9,13,18,19,23-26]. Interestingly, liver plays a central role in these adaptations. Hepatocytes can promote tolerance via producing glucose[19,21-23], degrading heme[18,23], and responding to glucocorticoid[24] and GDF15[25] as well as by secreting acute phase proteins[14,27], FGF21[26], and triglyceride-rich very low density lipoprotein (VLDL)[15,25,26].

When managed well, the net effect is reduced organ damage, preserved heart function, and prevention of hypothermia[13,15,18,19,23,25,26,28]. Moreover, hepatocytes and other cells in the liver also coordinate to promote disease resistance by destroying the invading pathogen, altering systemic inflammation, and regulating lipoproteins that eliminate/clear the pathogen[14,16,17,20,27,29-34]. These defenses are important to avert life-threatening pathology, as patients with liver dysfunction due to cirrhosis are more prone to acquire sepsis and less able to survive sepsis[27,35,36]. Hence, liver may be a critical target organ to stimulate host defenses capable of improving sepsis outcomes.

Stress defense networks mediate damage control and promote metabolic adaptations that underlie disease tolerance[7,9-12]. Which networks in hepatocytes contribute to liver defenses against sepsis is not clear, but transcriptional programming is important for this process[37]. Previous works identified a role for the transcription factors glucocorticoid receptor, farnesoid X receptor, and peroxisome proliferator-activated receptor alpha[24,38-40]. In this study, we considered whether stress defending transcription factors nuclear factor erythroid 2 related factor-1 (Nrf1) and -2 (Nrf2) contribute to such programming. Nrf1 and Nrf2 can regulate genes that protect against ROS, proteotoxicity, impaired cell metabolism, and organelle dysfunction[41-46]. While a role for Nrf1 in sepsis has not yet been studied, whole body Nrf2 deficient mice were found to have greater mortality[47]. With respect to liver defenses, we have shown combined deletion of hepatic Nrf1 and Nrf2, by itself, results in predisposition to mortality and transcript profiles that resemble exposure to the sepsis-inducing endotoxin lipopolysaccharide (LPS)[48] as well as reduced glucose[49] and high density lipoprotein[50] output. Based on the role of liver and its adaptive role in disease tolerance, we reasoned these effects may influence sepsis outcomes. To investigate, here we modulate hepatic Nrf1 and Nrf2 level in mice and inject them with LPS or live *Escherichia coli* (*E. coli*) to determine whether the actions of hepatic Nrf1 and Nrf2 play a role in liver defenses against sepsis.

# Results

## Reduced hepatic Nrf1, but not Nrf2, activity impairs survival in LPS-induced sepsis

Nrf1 and Nrf2 can regulate genes that protect against ROS and other types of stress that emerge in liver during sepsis[41,42,44,45,47,48,51-57] . To determine the effect of sepsis on hepatic Nrf1 and Nrf2 activity, we examined liver and survival of mice injected with LPS from *E. coli* O111:B4, which has been extensively used for investigating sepsis[15,22,25,26,28,38,47]. During a 72-hour survival study, 1 mg/kg LPS resulted in 100 % survival, 10 mg/kg LPS resulted in 64 % survival, and 20 mg/kg LPS resulted in 10 % survival (Fig. 1A). These sub-lethal, low, and high LPS doses were used in further experiments. Using an established loss-of-function model[48-50], we verified the detection of Nrf1 and Nrf2 in liver nuclei (Fig. S1A) and found that LPS caused expected changes in liver, such as reduced growth hormone receptor[58,59] 8-hours post-injection with 20 mg/kg LPS (Fig. S1B) and increased mRNA of acute phase proteins serum amyloid A1, *Saa1*, and haptoglobin, *Hp*[14], 8- and 24-hours post-injection with 1 mg/kg LPS (Fig. S1C). Next, we examined surrogate factors for hepatic Nrf1 and Nrf2 activity. Compared to PBS, 8-hours post-injection with 20 mg/kg LPS reduced Nrf1 and Nrf2 in liver nuclei, while increasing NFκB (Fig. 1B,1C). This corresponded with reduced liver expression of the genes encoding Nrf1 (*Nfe2l1*) and Nrf2 (*Nfe2l2*) (Fig. S1D) and target genes of both[48], *Gclc*, *Gclm*, *Gsta3*, and *Gstm1* (Fig. 1D). 8- and 24-hours post-injection, the sub-lethal 1 mg/kg dose of LPS had little effect on liver mRNA for *Nfe2l1* and a trending reduction in *Nfe2l2* mRNA (Fig. S1E). But, re-analyzed RNA sequencing (RNA-seq) differential expression data comparing liver from mice 12-hours post-injection with 1 mg/kg LPS to untreated control, see Ganeshan et al.[15] and table S3, still support that LPS reduces expression of *Gclc*, *Gclm*, *Gsta3*, and *Gstm1* (Fig. S1F). Altogether, these findings indicate that hepatic Nrf1 and Nrf2 activity is acutely impaired in LPS-induced sepsis, particularly during exposure to a highly lethal dose.

The impact of ablating hepatic Nrf1 and Nrf2 activity days prior to sepsis onset on survival outcome was examined by utilizing mice with flox alleles for *Nfe2l1*, *Nfe2l2*, or both, which we described previously[48,53]. As before[48-50], *Nfe2l1^flox/flox* mice, *Nfe2l2^flox/flox* mice, and *Nfe2l1^flox/flox*; *Nfe2l2^flox/flox* mice were infected with adeno-associated virus (AAV) expressing Cre recombinase via thyroxine binding globulin promoter (AAV-CRE) to generate mice with hepatic deletion of Nrf1, Nrf2, or both, respectively. Controls received AAV expressing green fluorescent protein (AAV-GFP). As shown (Fig. 1E), mice received AAV infection 12 days prior to LPS injection, consistent with previous[48-50]. On day 12 (corresponding to ~5 days of gene deficiency), mice were injected with 10 mg/kg LPS, a dose found to result in 64 % survival in regular mice (see Fig. 1A). Survival was similar in controls, ranging from 50 % - 69 % between flox lines, and mice with hepatic Nrf2 deficiency were similar to control (survival: AAV-GFP = 50 % vs AAV-CRE = 45 %). In contrast, survival was reduced in mice with Nrf1 deficiency (survival: AAV-GFP = 69 % vs AAV-CRE = 26 %) and combined deficiency (survival: AAV-GFP = 58 % vs AAV-CRE = 23 %). Hence, we demonstrate that hepatic Nrf1, but not Nrf2, is required to protect against sepsis, and impairment to its activity, such as occurs in high LPS exposure, may predispose for lethality.

**Effects of hepatic Nrf1, Nrf2, and combined deficiency in LPS-induced sepsis**

We next sought to identify inflammatory and metabolic factors that may explain reduced survival in hepatic Nrf1 deficient mice exposed to LPS. Previous works show hepatic Nrf1 can counteract liver inflammation, promote proteostasis, and regulate lipid and glucose metabolism[45,46,48-54]. Here, compared to PBS, 10 mg/kg LPS robustly increased plasma levels of several inflammatory cytokines that underlie sepsis 8-hours post-injection, but hepatic deficiency for Nrf1, Nrf2, or both had no effect on these levels (Fig. 2A,S2A) or on LPS-induced changes to plasma glucose (Fig. 1B) and bile acids (Fig. S2B). In contrast, liver expression of the chemokine *Ccl2* was increased

and the transcription factor *Ppara* was decreased by hepatic Nrf1 and combined deficiency (Fig. S2C), irrespective of LPS exposure. Likewise, liver inflammation was increased in hepatic Nrf1 and combined deficiency, irrespective of LPS (Fig. S2D,S2E). Hepatic Nrf1 deficiency increased the unfolded protein response marker, phosphorylated eukaryotic initiation factor 2α (p-eIF2α), but had no effect on the protein translation activity marker, phosphorylated ribosomal protein S6 kinase (p-S6K), (Fig. S2F, S2G), suggesting these livers were under endoplasmic reticulum (ER) stress. Compared to PBS, LPS increased circulating triglyceride in control mice and in mice with hepatic Nrf2 deficiency, but this effect was blunted in the mice with hepatic Nrf1 deficiency and in those with combined deficiency (Fig. 2C). Altogether, we show that reduced survival in hepatic Nrf1 deficient mice is not due to altered systemic inflammation but does correspond with increased ER stress and inflammation in the liver as well as lower levels of circulating triglyceride.

**Enhancing hepatic Nrf1 activity increases survival and triglyceride in LPS-induced sepsis**

In complement to gene ablation (loss-of-function), we also investigated the effect of increasing hepatic Nrf1 activity (gain-of-function) prior to sepsis onset on survival outcome by comparing LPS responses of control mice infected with AAV-GFP to mice infected with an AAV expressing high levels of human NRF1 (AAV-hNRF1), which we described previously[48]. In this case, we used 20 mg/kg LPS, a dose we previously found to result in only 10 % survival in regular mice (see Fig. 1A). As shown (Fig. 3A), 12 days after AAV infection (corresponding to ~5 days of increased Nrf1 activity), mice were injected with LPS and a 72-hour survival study done. Compared to control, mice with hepatic Nrf1 gain-of-function had improved survival (survival: AAV-GFP = 11 % vs AAV-hNRF1 = 46 %). Hence, enhancing hepatic Nrf1 activity is protective against LPS-induced sepsis, which is the opposite effect of hepatic Nrf1 deficiency. Therefore, we show that hepatic Nrf1 is a critical factor involved in promoting liver defenses against sepsis.

Next, using *Nfe2l1^flox/flox* mice, we used complementary AAV-induced models for hepatic Nrf1 loss- (AAV-CRE) and gain-of-function (AAV-hNRF1) to identify sepsis-related parameters affected in the same opposing pattern as survival outcomes, compared to control mice (AAV-GFP) 8 hours post-injection with 20 mg/kg LPS or PBS (schematic in Fig. 3B). This was not the case for plasma glucose and ketones (Fig 3C,S3A). Moreover, hepatic Nrf1 loss-of-function decreased and gain-of-function increased plasma IGF1 in PBS injected but not in LPS injected mice (Fig. S3B) and the same pattern occurred for liver expression of growth hormone receptor in PBS and LPS injected mice (Fig. S3C). Liver expression of *Ccl2* was increased by hepatic Nrf1 loss-of-function, but unaffected by hepatic Nrf1 gain-of-function (Fig. S3D). Interestingly, in only the LPS injected mice, hepatic Nrf1 loss-of-function decreased and gain-of-function increased plasma triglyceride (Fig. 3D), whereas hepatic Nrf1 loss-of-function increased and gain-of-function decreased plasma free fatty acid (Fig. 3E). This shows the protective effect of hepatic Nrf1 activity against LPS-induced sepsis coincide with reduced free fatty acid and increased triglyceride in the bloodstream.

**Enhancing hepatic Nrf1 activity increases stress defense programming in LPS-induced sepsis**

Given current understanding[42,45,46,48,50-54,60,61], we reasoned that hepatic Nrf1 activity protects against LPS-induced sepsis via gene regulation. To investigate, RNA-seq was done on livers from control (AAV-GFP), hepatic Nrf1 loss-of-function (AAV-GFP), and hepatic Nrf1 gain-of-function (AAV-hNRF1) *Nfe2l1^flox/flox* mice 8 hours post-injection with 20 mg/kg LPS. Principle component analysis (PCA) comparing AAV-CRE to AAV-GFP and AAV-hNRF1 to AAV-GFP revealed hepatic Nrf1 loss- and gain-of-function did not cause major changes to transcript profiles (Fig. S4A,S4B). This was not surprising, given that LPS on its own drives broad and robust alterations to liver transcript profiles[15,37,62], and this may mask effects of altering hepatic Nrf1 activity. Interestingly,

in previous work on liver in less stressful conditions[48,51-53] hepatic Nrf1 loss-of-function has been shown to alter expression of thousands of genes, but there were no genes significantly affected by hepatic Nrf1 loss-of-function in this case (Fig. S4A) when liver is experiencing stress caused by LPS exposure and which, by itself, can impair hepatic Nrf1 activity (see Fig. 1B-1D). In contrast, compared to control using false discovery rate p-value adjusted DESeq2 analysis, expression of 330 genes were significantly different (69 down-regulated and 261 up-regulated) in liver of mice with hepatic Nrf1 gain-of-function (Fig. S4B). Hence, hepatic Nrf1 gain-of-function impacted hepatic transcriptional programming, and this may underlie coinciding protection against sepsis.

Ingenuity pathway analysis (IPA) was done to identify pathways most affected by hepatic Nrf1 gain-of-function. Genes undergoing IPA were selected using Wald statistics $\geq 3$ (391 genes) and $\leq$ -3 (76 genes), which amounted to 467 genes. Focusing on diseases-and-biofunctions, pathways most affected include "necrosis", "oxidative stress", "inflammation of body cavity", and "organismal death", and molecular activity prediction analysis suggest these pathways were less active in liver with hepatic Nrf1 gain-of-function (Fig. 4A). A heatmap of transformed RNA-seq values for genes corresponding to the "necrosis" pathway was also done (Fig. 4B), and show increased mRNA level of several genes that counteract cell stress and many of which are known to be regulated by Nrf1, such as *Gclm*, *Gstm1*, *Psma4*, *Psmb5*, and *Vcp*[42,45,48,51,52,54,60]. Altogether, this suggests enhancing hepatic Nrf1 activity increased hepatic resistance to sepsis-induced stress, and this may directly or indirectly promote liver defenses that protect against sepsis.

**Hepatic Nrf1 activity protects against *E. coli*-induced sepsis and hypothermia**

To ensure results using LPS translate to bacterial inflammation, we investigated whether hepatic Nrf1 loss- and gain-of-function *Nfe2l1^flox/flox^* mice had the same opposing effects on survival and sepsis-related parameters in mice infected with live *Escherichia coli* (*E. coli*). An *E. coli* dose

response was done (Fig. S5A), from which we determined that administering $4.0 \times 10^8$ colony forming units (CFU) *E. coli* in males and $3.2 \times 10^8$ CFU *E. coli* in females results in approximately 50 % survival during a 120-hour survival study. These doses were used for further investigation. Compared to PBS, *E. coli*-induced sepsis reduced liver expression of Nrf1 and Nrf2 target genes 24-hours post-infection (Fig. S5B), similar to the effect of high dose LPS (Fig. 1B-1D). In comparison to control, hepatic Nrf1 loss-of-function reduced survival and hepatic Nrf1 gain-of-function increased survival (Fig. 5A). Similar to the LPS model (Fig. 2B,3C), altering hepatic Nrf1 activity had no effect on sepsis-induced changes to plasma glucose (Fig. 5B) whereas plasma triglyceride was increased in mice with hepatic Nrf1 gain-of-function (Fig. 5C) 24 hours post-infection with *E. coli*. We also monitored body temperature prior to (i.e., time 0) and 24 hours post-infection, as hypothermia is a critical trigger for sepsis-induced lethality and the liver is capable of defending against this trigger by fueling thermogenic tissues with VLDL-triglyceride[15,25,26,63]. As with the survival outcomes (Fig. 5A), hepatic Nrf1 loss-of-function worsened hypothermia and gain-of-function improved hypothermia, compared to control mice, whereas there was no body temperature difference prior to infection (Fig. 5D). Thus, we show hepatic Nrf1 activity protects against *E. coli*-induced sepsis, with corresponding changes in circulating triglyceride, and also that the mechanism of this protection may involve liver defense against hypothermia.

**Hepatic Nrf1 activity protects against sepsis and hypothermia by promoting VLDL secretion**

Triglyceride rich VLDL promote survival in mice with sepsis by counteracting hypothermia[15,25,26]. So, we investigated whether reciprocal changes in circulating triglyceride that occurred in hepatic Nrf1 loss- and gain-of-function mice (Fig. 2,3,5) is causally linked to coinciding effects on liver defense against sepsis-induced hypothermia. First, we injected the sub-lethal dose of 1 mg/kg LPS (see Fig. 1A) to *Nfe2l1^flox/flox* mice infected 12 days prior with AAV-GFP (control), AAV-CRE (loss-

of-function), or AAV-hNRF1 (gain-of-function) and monitored body temperature, sepsis score, plasma triglyceride and other sepsis-related parameters 0-, 8-, and 24-hours post-injection (see Fig. 6A). There were changes to some plasma cytokines (Fig. S6A), liver inflammation (Fig. S6B,S6C), and plasma glucose (Fig. S6D), but this did not follow a consistent trend or pattern that may explain the outcomes. Similarly, hepatic Nrf1 gain-of-function increased liver expression of *Ghr* 0-, 8-, and 24-hours post-injection (Fig. S6E) but hepatic Nrf1 loss-of-function had no effect, whereas hepatic Nrf1 loss-of-function reduced plasma IGF1 0- and 24-hours post-injection (Fig. S6F), but hepatic Nrf1 gain-of-function had no effect. In contrast to these inconsistent patterns, hepatic Nrf1 loss-of-function worsened whereas gain-of-function improved hypothermia and the sepsis score (Fig. 6B,6C), which coincided with reduced plasma triglyceride in hepatic Nrf1 loss-of-function mice and increased plasma triglyceride in hepatic Nrf1 gain-of-function mice (Fig. 6D). Hence, we confirm there is a corresponding relationship between sepsis severity, body temperature, and circulating triglyceride with hepatic Nrf1 activity.

Liver-derived circulating triglyceride primarily originates from secreted VLDL, and this lipoprotein is capable of mediating disease tolerance[15,25,26,28]. So, we investigated whether hepatic Nrf1 promotes VLDL secretion to protect against sepsis severity and hypothermia in LPS exposed mice. First, we performed an *in vivo* VLDL secretion assay. 16-hours post-injection with 1 mg/kg LPS, mice were co-injected with lipoprotein lipase inhibitor poloxamer 407 and then triglyceride was monitored 0-, 2-, and 4-hours later. Compared to control, by hour 4 plasma triglyceride was reduced in mice with hepatic Nrf1 loss-of-function and increased in mice with hepatic Nrf1 gain-of-function (Fig. 6E), showing hepatic Nrf1 activity positively correlates with VLDL secretion. Second, we investigated whether increased VLDL secretion in hepatic Nrf1 gain-of-function mice is causally linked to protection against sepsis by employing use of the VLDL secretion inhibitor

lomitapide, which is a clinically used agent that inhibits triglyceride incorporation into VLDL[64].
As expected, lomitapide blunted plasma triglyceride levels in the AAV-GFP and AAV-hNRF1
infected mice (Fig. S6G). Control (AAV-GFP) and hepatic Nrf1 gain-of-function (AAV-hNRF1)
mice were injected with LPS + vehicle or LPS + lomitapide, whereas the hepatic Nrf1 loss-of-
function (AAV-CRE) mice were injected with only LPS + vehicle. Body temperature and sepsis
score was monitored 0-, 8-, and 24-hours post-injection (Fig. 6F,6G). As before, hepatic Nrf1 loss-
of-function mice had more severe hypothermia and sepsis score than LPS + vehicle treated control
mice. Conversely, hepatic Nrf1 gain-of-function mice treated with LPS + vehicle had improved
hypothermia and sepsis score than LPS + vehicle treated control mice. However, the beneficial
effect of hepatic Nrf1 gain-of-function was lost in mice treated with LPS + lomitapide, and this
was so severe that the body temperature and sepsis score in either control or hepatic Nrf1 gain-of-
function mice treated with LPS + lomitapide were now similar to that found in hepatic Nrf1 loss-
of-function mice treated with LPS + vehicle. Hence, we show that the secretion of triglyceride rich
VLDL is increased during sepsis by enhancing hepatic Nrf1 activity and that this effect is required
for hepatic Nrf1 activity to protect against sepsis.

# Discussion

Sepsis is a major health care challenge and there are limited interventional options[1-6]. One pressing need is a strategy for reducing tissue damage to preserve organ function and improve recovery[6]. Recent works show there are endogenous host processes capable of this feat, which is referred to as disease tolerance[7-10]. There is interest in stimulating mechanisms of disease tolerance to improve sepsis outcomes[2,6-10]. Mechanisms underlying disease tolerance are incompletely understood, but mediators of tissue damage control and their effect on systemic metabolism are considered critical components[7,9-11]. Here, we investigated whether transcription factors Nrf1 and Nrf2 in hepatocytes, which regulate genes involved in stress defense and damage control[41-46], protect against sepsis. We demonstrate that ablating hepatic Nrf1, but not Nrf2, predisposes for mortality whereas genetic-induction of hepatic Nrf1 activity improves survival in mice with LPS- and *E. coli*-induced sepsis. Thus, we identify hepatic Nrf1 as an important factor for liver defense against sepsis.

The liver plays a central role in defending against sepsis and hepatocytes contribute to this role, which includes adaptations in glucose, lipid, bile acid, ketone, and lipoprotein metabolism as well as influences on systemic inflammation[14-36,38-40]. We sought to determine whether effects of altered hepatic Nrf1 activity on survival in sepsis was linked to some of these parameters. Compared to control, hepatic Nrf1 loss- and gain-of-function did not alter systemic inflammation in mice with sepsis, and had no effect on sepsis-induced reductions in glycemia. Likewise, complementary alterations to hepatic Nrf1 activity had parallel, not opposing, effects on ketones and no effect on bile acids in *Nfe2l1^flox/flox* mice. Liver inflammation was increased in hepatic Nrf1 loss-of-function mice, and this was elevated 24-hours post-injection with sub-lethal sepsis, but the opposite did not occur in hepatic Nrf1 gain-of-function mice. Thus, the effect of altered hepatic Nrf1 activity on survival was apparently not linked to the effects of sepsis on glucose, ketone, and

bile acid metabolism as well as systemic inflammation. Since glucocorticoid receptor, farnesoid X receptor, and peroxisome proliferator-activated receptor has been shown to protect against sepsis by regulating these metabolic processes[24,38-40], hepatic Nrf1 may control an independent program that operates in parallel with these other factors. We did find a positive but inconsistent correlation for hepatic Nrf1 activity with liver growth hormone receptor and circulating IGF1, indicating there may be a link with the growth hormone/IGF1 axis. We previously found this link in a non-sepsis context[49] but are uncertain of the significance. In contrast, we consistently identified a positive correlation for hepatic Nrf1 activity with circulating triglyceride in sepsis, that hepatic Nrf1 loss-of-function mice have reduced whereas hepatic Nrf1 gain-of-function mice have increased VLDL secretion, and that inhibiting VLDL secretion blocked protective effects of enhancing hepatic Nrf1 activity against hypothermia and sepsis severity. Hence, our results show that hepatic Nrf1 against sepsis by promoting an increase in the secretion of triglyceride rich VLDL.

VLDL can protect against sepsis by fueling thermogenesis to avert hypothermia[15,25,26] and by facilitating pathogen elimination[30-32,34]. Here, we show Nrf1-regulated VLDL secretion protects against hypothermia, which underlies mortality[63]. While the mechanism by which hepatic Nrf1 controls VLDL secretion is not clearly understood from this study, we did examine the effect of hepatic Nrf1 on transcript profiles. RNA-seq results show hepatic Nrf1 gain-of-function liver has increased stress defense gene expression, consistent with previous[42,46,48,50-54], which may reduce susceptibility to liver damage. Indeed, hepatic Nrf1 loss-of-function liver had increased ER stress and inflammation. So, while we cannot draw a direct link, we speculate that liver with enhanced hepatic Nrf1 activity is better suited to manage the stress of sepsis and, in turn, to mediate defenses that promote survival. Consistent with this, we found that severe sepsis reduced Nrf1 and Nrf2 in liver nuclei, coinciding with reduced target gene expression, indicating susceptibility to mortality

in severe sepsis may in part result from impaired hepatic Nrf1 and Nrf2 activity. Indeed, we have shown that combined deletion of hepatic Nrf1 and Nrf2 in non-sepsis adult mice, by itself, results in a 36 % mortality rate just 28-days after inducing the gene deletion[48].

In this study, we identify an important role for hepatic Nrf1 in sepsis, which uncover four avenues for future research. First, we primarily use LPS here, due to its extensive use investigating sepsis and disease tolerance[15,22,25,26,28,38,47]. Though we confirmed key results using *E. coli*-induced sepsis, further work is needed to determine whether altering hepatic Nrf1 activity has similar effects in other types (e.g., polymicrobial, gram positive bacteria, virus, and fungal) and modes (e.g., intravenous versus peritonitis) of infection to ascertain general effectiveness in protecting against sepsis. Second, transcriptional programming is important for counteracting sepsis[24,37-40]. So, it is critical to delineate which genes Nrf1 regulates to elicit protective actions against sepsis, as this may reveal strategies for enhancing disease tolerance. Third, it is important to determine the mechanism by which Nrf1 is impaired in sepsis and whether reversing this effect can improve sepsis outcomes. This may also be the case for Nrf2. Though hepatic Nrf2 was not required to protect against sepsis, Nrf2 in other cell types and its induction has been shown to protect against sepsis[47,65,66] and may require hepatic Nrf2 to do so. Fourth, further insight is needed regarding the impact that hepatic Nrf1 has on the quantity and quality of VLDL and other sepsis counteracting lipoproteins such as high-density lipoprotein (HDL)[17,31,32,50]. Likewise, the mechanism by which VLDL and HDL protect against sepsis is unclear and likely a valuable area of future research.

In summary, we show hepatic Nrf1 protects against sepsis via promoting VLDL secretion, which prevents hypothermia to reduce sepsis severity and promote survival. Our finding underscores the value of investigating stress defense programming in liver and its impact on the mechanisms underlying tissue damage control in sepsis.

# Methods

## Animal used in study

Experiments were done on 20–30-week-old male and female mice on the C57BL/6J background, with the approval of the University of Saskatchewan's Animal Care Committee. Mice were group housed at 21 °C on a 12 h light/dark cycle and provided *ad libitum* access to chow from LabDiet (Prolab RMH 3000; catalog#: 5058) and water. Mice containing flox alleles in the genes for Nrf1 (*Nfe2l1^flox/flox*), Nrf2 (*Nfe2l2^flox/flox*), or both Nrf1 and Nrf2 (*Nfe2l1^flox/flox*; *Nfe2l2^flox/flox*) were bred in-house and described previously[48,53].

To induce hepatocyte-specific loss-of-function for Nrf1 activity, Nrf2 activity, or both, recombination of respective flox alleles was induced by retro-orbital infection of mice, while under anesthesia (3% isoflurane at oxygen flow rate of 1 L/minute), with a serotype 8 adeno-associated virus (AAV) that expresses Cre recombinase via the hepatocyte-specific thyroxin binding globulin promoter (AAV-CRE). To induce hepatocyte-specific gain-of-function for Nrf1 activity, mice were infected with AAV expressing human NRF1 (AAV-hNRF1). In each case, littermate control mice were infected with AAV expressing green fluorescent protein (AAV-GFP). Use of these AAVs were established by us previously[48-50]. In each experiment, males were infected with 3.0 x 10^11 AAV particles and females with 2.0 x 10^11 AAV particles, as this was found to result in more equivalent expression between sexes. All experiments involving AAVs were done within 17 days of infection. AAV-CRE (AAV8.TBG.PI.Cre.rBG) and AAV-GFP (AAV8.TBP.PI.eGFP.WPRE.bGH) were acquired from Addgene (#107787-AAV8 and #105535-AAV8). AAV-hNRF1 was produced by the University of Pennsylvania Vector Biocore and is described in our previous work[48].

**Sepsis induction and assessment of severity**

Sepsis was induced in mice via intraperitoneal (IP) injection with either lipopolysaccharide (LPS) from *E. coli* O111:B4 (Sigma, catalog #L2630) or with live DH5α *Escherichia coli* (*E. coli*; New England Biolabs, catalog #C2987H). Doses are described in the text, figures, and figure legends. In experiments that employed AAVs, sepsis was induced on day 12 after AAV administration. Mice were monitored for up to 72 hours when using LPS and 120 hours when using *E. coli*. Monitoring for symptoms and assignment of a murine sepsis score was done according to the guidelines by Shrum et al.[67]. Body temperature was measured using a rectal probe thermometer (BIOSEB Lab Instruments, catalog #BIO-TK8851). Mice were euthanized upon reaching a predefined humane endpoint and, at that point, determined to have not survived the sepsis challenge.

**Bacteria preparation for sepsis induction**

*E. coli* were heat shocked in a water bath at 42 °C for 30 seconds, placed on ice for 2 minutes, and grown to log phase in LB broth (Fisher, catalog #BP1426) at 37 °C in a shaking incubator. *E. coli* growth curve was quantified by correlating optical density at 600 nm to colony forming units (CFU), which was determined by spreading on LB agar plates (Fisher, catalog #BP1423). Prior to infecting mice, *E. coli* underwent centrifugation at 2000 g for 10 minutes at 4 °C, washed once with calcium and magnesium free phosphate buffered saline (PBS), and then resuspended in PBS.

**VLDL secretion assay and use of VLDL secretion inhibitor**

Mice underwent a very low-density lipoprotein (VLDL) secretion assay to examine the role of Nrf1 and hepatic triglyceride output in protecting against sepsis. For this, mice were injected with a sub-lethal 1 mg/kg dose of LPS, fasted overnight, and 16 hours later received IP injection with 1g/kg of lipoprotein lipase inhibiting agent, poloxamer 407 (Sigma, catalog #16758). Blood was

collected immediately prior to and up to 4 hours after receiving poloxamer 407 to measure plasma levels of triglyceride, as a surrogate readout for VLDL output. In addition to VLDL secretion assay, assessment of the effect of inhibiting VLDL secretion on body temperature and murine sepsis score post-LPS exposure was also done. In this case, prior to IP injection with LPS, mice were fasted overnight and received IP injection on the contralateral side with 2 mg/kg of lomitapide (Sigma, catalog #SML1385) in 10% DMSO/PBS, which suppresses VLDL secretion by inhibiting triglyceride loading into newly synthesizing VLDL particles in hepatocytes[64].

**Tissue collection**

Tissues were collected from mice anesthetized via 3 % isoflurane at oxygen flow rate of 1 L/minute and euthanized by exsanguination and cervical dislocation. Blood was collected using a 26-gauge syringe needle via cardiac puncture and placed into an ethylenediaminetetraacetic acid (EDTA) containing tube (5 mM final concentration). Plasma was separated via centrifugation at 8000 g and 4 °C for 10 minutes, placed into a fresh tube, and snap frozen on dry ice. Liver was collected by initially flushing the vasculature of euthanized mice with PBS. The liver was excised, and a piece was immediately processed for histological analysis, nuclear isolation, or snap frozen on dry ice, as described in text and figure legends. All tissues were stored at -80 °C until further use.

**Plasma analysis**

Plasma triglyceride levels were determined using Infinity Triglyceride reagent (ThermoFisher Scientific, catalog #TR22421) and a Synergy HT microplate reader. Plasma glucose was determined using an Accu-Check glucometer (Roche). Plasma cytokine levels were measured by Eve Technologies (https://www.evetechnologies.com/). Plasma IGF1 levels were determined by ELISA according to manufacturer's instructions (R&D Systems, catalog #DY791).

**Histological analysis**

A piece of the left liver lobe was fixed in 10 % neutral buffered formalin for 24 hours, subjected to three 24-hour washes in 70 % ethanol, embedded in wax, sectioned at 5 μm thickness, mounted on glass slides, and stained with hematoxylin and eosin (H&E). Images were capture using Aperio Scanscope CS image analysis System and Aperio Imagescope viewing software (version 12).

**Gene expression analysis using qPCR**

Ribonucleic acid (RNA) were extracted from frozen liver tissue homogenized in TRIzol Reagent (ThermoFisher Scientific, catalog #15596018). RNA clean-up was performed using a Qiagen RNeasy plus column (ThermoFisher Scientific, catalog #74034). RNA concentration and quality were determined using a NanoDrop One spectrophotometer (Thermo-Fisher Scientific). One μg of RNA were reverse transcribed into complementary deoxyribonucleic acid (cDNA) using a Maxima First strand cDNA synthesis kit, with dsDNase (ThermoFisher Scientific, catalog #K1672). Gene expression was determined using quantitative polymerase chain reaction (qPCR) on cDNA template with a Bio Rad CFX384-well Real-time PCR Detection System and PowerUp SYBR green Master Mix (ThermoFisher Scientific, catalog #A25742). A list of primer sequences for measuring gene expression are provided in table S1. Change in expression of target genes were normalized to the reference genes *36b4* and *Tbp* and all data presented were in good agreement. Gene expression data are presented as relative expression compared to the control group after normalization, as indicated in the figure legends.

**Gene expression analysis using RNA sequencing**

RNA sequencing (RNA-seq) was performed by the University of Saskatchewan Next Generation Sequencing Facility. As in previous[48], 500 ng of RNA per sample were used to prepare sequencing

libraries using TruSeq Stranded mRNA Library Prep Kit from Illumina (catalog #20020595 and 20022371). Libraries were quantified using the Qubit 4 fluorometer and dsDNA BR DNA assay (Invitrogen, catalog #Q32850). The fragment size of each library was determined using the DNA Screentape (D1000) assay on the TapeStation 4150 instrument. Libraries were pooled and sequenced using a NextSeq 500/550 High Output Kit v2.5 (75 cycles), which generated 38 bp paired end reads. Reads were extracted and adapter trimmed using Illumina fastp[68]. Reads were aligned using STAR version 2.7.9a[69], using the Gencode M32 mouse genome as reference. Generate FASTQ BaseSpace pipeline (version 1.1.0.64). There were > 25 million aligned reads per sample. Principal component analysis (PCA) was done to visualize sample clustering. Differential gene expression was determined using DESeq2 (version 1.22.2)[70].

**Nuclear fraction isolation**

Liver nuclei was isolated to assess Nrf1 and Nrf2 levels, as we did previously[48]. Approximately 250 mg of fresh liver from two mice were pooled (500 mg total), chopped into 1-3 mm pieces using a scalpel blade, and then homogenized with a Teflon homogenizer in STM buffer (250 mM sucrose, 50 mM Tris (pH 7.4), 5 mM $MgCl_2$) containing 1 mM phenylmethylsulfonyl fluoride and 1mM dithiothreitol. Samples underwent 800 g centrifugation for 15 minutes at 4 °C. The pellet was resuspended in 10 ml of ice-cold buffer A (10 mM HEPES (pH 7.9), 10 mM KCl, 1.5 mM MgCl2, 0.34 M sucrose, 10 % glycerol, 1 mM DTT, 0.1 % Triton X-100) and 1x protease inhibitor cocktail (Roche, catalog #11697498001) and incubated on ice for 10 minutes. Pellets were then collected and washed once with ice-cold buffer A. Samples were centrifuged at 4 °C at 800 g for 15 minutes, supernatants were removed, and nuclear lysates were prepared for protein analysis by immunoblot in 1 ml ice-cold RIPA buffer containing 1x protease inhibitor cocktail.

**Immunoblot**

Proteins were extracted from frozen liver pieces or nuclear fractions using RIPA buffer with fresh protease inhibitor cocktail. Concentrations were determined using a Pierce BCA protein assay kit (ThermoFisher Scientific, catalog #23250) and then heated at 95 °C for 5 minutes in loading buffer containing 1 % sodium dodecyl sulfate and then cooling on ice. 10 µg of protein was loaded onto a Novex 4-12 % Bis-Tris gel (ThermoFisher Scientific, catalog #NP0336), separated by size via electrophoresis in NuPAGE MOPS Running Buffer (ThermoFisher Scientific, catalog #NP0336), and transferred to a nitrocellulose membrane (Bio-Rad, catalog #1620115). Membranes were blocked in tris-buffer saline containing 0.1 % tween 20 (TBST) and 5 % powdered skim milk, and then incubated overnight at 4 °C with primary antibodies targeting the protein of interest. Antibodies used are described in table S2. Membranes were washed with TBST and incubated with horse radish peroxidase-conjugated secondary antibodies at room temperature for 60 minutes. Membranes were washed with TBST and levels of immuno-detected protein measured by incubating in Super Signal West Femto Maximum Sensitivity Substrate (ThermoFisher Scientific, catalog #34096) to generate chemiluminescence, which was detected using a BioRad ChemiDoc MP Imaging System. Level of signal was analyzed using ImageJ (https://imagej.net/ij/).

**Quantification and statistical analysis**

Male and female mice were used throughout the study. All between-sex results were in good agreement and so male and female samples were pooled throughout when performing statistical analysis. Statistical analysis of RNA-seq data was done using DESeq2[70]. All other data was analyzed using GraphPad PRISM (version 10.2.2). Significance was defined as $p < 0.05$, using the Mantel-Cox Log rank test, t-test, one-way, or two-way analysis of variance (ANOVA) and Dunnett's or Tukey's post-test, as indicated in figure legends.

# Resource availability, author contributions, and funding acknowledgment

**Supplemental Information.** Supplemental information is available for this manuscript. This includes the following: Supplemental Figures S1-S6 and Supplemental Tables S1-S3.

**Data Availability**. Upon acceptance for publication, data corresponding to figures will be publicly available to view in the Mendeley repository and raw sequence reads pertaining to RNA-seq data will be publicly available to view in the National Center for Biotechnology Information (NCBI) repository.

**Author Contributions**. Conceptualization, M.J.T and S.B.W.; methodology, M.J.T., L.L., M.G.A, A.H., and S.B.W.; formal analysis, M.J.T and S.B.W.; investigation, M.J.T and S.B.W.; data curation, M.J.T and S.B.W.; writing - original draft preparation, M.J.T. and S.B.W.; writing - review and editing, S.B.W.; visualization, M.J.T and S.B.W.; supervision, S.B.W.; funding acquisition, S.B.W..

**Declaration of Interests**. The authors declare there are no competing interests.

**Funding Sources.** Grants supporting this research were awarded to S.B.W. from the Canadian Institutes of Health Research (project #PJT-174988). M.J.T., M.G.A., and A.H. were supported by a scholarship from the University of Saskatchewan's College of Medicine. A.H. was also supported by a scholarship from the Natural Sciences and Engineering Research Council of Canada. S.B.W. was supported by a National New Investigator Award from the Heart and Stroke Foundation of Canada.

# References

Singer, M. *et al.* The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* **315**, 801-810 (2016). https://doi.org:10.1001/jama.2016.0287

Angus, D. C. & van der Poll, T. Severe sepsis and septic shock. *N Engl J Med* **369**, 840-851 (2013). https://doi.org:10.1056/NEJMra1208623

Fleischmann, C. *et al.* Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *Am J Respir Crit Care Med* **193**, 259-272 (2016). https://doi.org:10.1164/rccm.201504-0781OC

Rudd, K. E. *et al.* Global, regional, and national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study. *Lancet* **395**, 200-211 (2020). https://doi.org:10.1016/S0140-6736(19)32989-7

Shankar-Hari, M. *et al.* Evidence for a causal link between sepsis and long-term mortality: a systematic review of epidemiologic studies. *Crit Care* **20**, 101 (2016). https://doi.org:10.1186/s13054-016-1276-7

Lelubre, C. & Vincent, J. L. Mechanisms and treatment of organ failure in sepsis. *Nat Rev Nephrol* **14**, 417-427 (2018). https://doi.org:10.1038/s41581-018-0005-7

Medzhitov, R., Schneider, D. S. & Soares, M. P. Disease tolerance as a defense strategy. *Science* **335**, 936-941 (2012). https://doi.org:10.1126/science.1214935

Ayres, J. S. & Schneider, D. S. Tolerance of infections. *Annu Rev Immunol* **30**, 271-294 (2012). https://doi.org:10.1146/annurev-immunol-020711-075030

Wang, A., Luan, H. H. & Medzhitov, R. An evolutionary perspective on immunometabolism. *Science* **363** (2019). https://doi.org:10.1126/science.aar3932

Soares, M. P., Teixeira, L. & Moita, L. F. Disease tolerance and immunity in host protection against infection. *Nat Rev Immunol* **17**, 83-96 (2017). https://doi.org:10.1038/nri.2016.136

Soares, M. P., Gozzelino, R. & Weis, S. Tissue damage control in disease tolerance. *Trends Immunol* **35**, 483-494 (2014). https://doi.org:10.1016/j.it.2014.08.001

Kotas, M. E. & Medzhitov, R. Homeostasis, inflammation, and disease susceptibility. *Cell* **160**, 816-827 (2015). https://doi.org:10.1016/j.cell.2015.02.010

Willmann, K. & Moita, L. F. Physiologic disruption and metabolic reprogramming in infection and sepsis. *Cell Metab* **36**, 927-946 (2024). https://doi.org:10.1016/j.cmet.2024.02.013

Gabay, C. & Kushner, I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* **340**, 448-454 (1999). https://doi.org:10.1056/NEJM199902113400607

Ganeshan, K. *et al.* Energetic Trade-Offs and Hypometabolic States Promote Disease Tolerance. *Cell* **177**, 399-413 e312 (2019). https://doi.org:10.1016/j.cell.2019.01.050

Grin, P. M. *et al.* Low-density lipoprotein (LDL)-dependent uptake of Gram-positive lipoteichoic acid and Gram-negative lipopolysaccharide occurs through LDL receptor. *Sci Rep* **8**, 10496 (2018). https://doi.org:10.1038/s41598-018-28777-0

Khovidhunkit, W. *et al.* Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanisms and consequences to the host. *J Lipid Res* **45**, 1169-1196 (2004). https://doi.org:10.1194/jlr.R300019-JLR200

Larsen, R. *et al.* A central role for free heme in the pathogenesis of severe sepsis. *Sci Transl Med* **2**, 51ra71 (2010). https://doi.org:10.1126/scitranslmed.3001118

Ramos, S. *et al.* A hypometabolic defense strategy against malaria. *Cell Metab* **34**, 1183-1200 e1112 (2022). https://doi.org:10.1016/j.cmet.2022.06.011

Walley, K. R. Role of lipoproteins and proprotein convertase subtilisin/kexin type 9 in endotoxin clearance in sepsis. *Curr Opin Crit Care* **22**, 464-469 (2016). https://doi.org:10.1097/MCC.0000000000000351

Wang, A. *et al.* Glucose metabolism mediates disease tolerance in cerebral malaria. *Proc Natl Acad Sci U S A* **115**, 11042-11047 (2018). https://doi.org:10.1073/pnas.1806376115

Wang, A. *et al.* Opposing Effects of Fasting Metabolism on Tissue Tolerance in Bacterial and Viral Inflammation. *Cell* **166**, 1512-1525 e1512 (2016). https://doi.org:10.1016/j.cell.2016.07.026

Weis, S. *et al.* Metabolic Adaptation Establishes Disease Tolerance to Sepsis. *Cell* **169**, 1263-1275 e1214 (2017). https://doi.org:10.1016/j.cell.2017.05.031

Vandewalle, J. *et al.* Combined glucocorticoid resistance and hyperlactatemia contributes to lethal shock in sepsis. *Cell Metab* **33**, 1763-1776 e1765 (2021). https://doi.org:10.1016/j.cmet.2021.07.002

Luan, H. H. *et al.* GDF15 Is an Inflammation-Induced Central Mediator of Tissue Tolerance. *Cell* **178**, 1231-1244 e1211 (2019). https://doi.org:10.1016/j.cell.2019.07.033

Huen, S. C. *et al.* Hepatic FGF21 preserves thermoregulation and cardiovascular function during bacterial inflammation. *J Exp Med* **218** (2021). https://doi.org:10.1084/jem.20202151

Strnad, P., Tacke, F., Koch, A. & Trautwein, C. Liver - guardian, modifier and target of sepsis. *Nat Rev Gastroenterol Hepatol* **14**, 55-66 (2017). https://doi.org:10.1038/nrgastro.2016.168

Nikkanen, J. *et al.* An evolutionary trade-off between host immunity and metabolism drives fatty liver in male mice. *Science* **378**, 290-295 (2022). https://doi.org:10.1126/science.abn9886

Kubes, P. & Jenne, C. Immune Responses in the Liver. *Annu Rev Immunol* **36**, 247-277 (2018). https://doi.org:10.1146/annurev-immunol-051116-052415

Bartolome, N. *et al.* Biphasic adaptative responses in VLDL metabolism and lipoprotein homeostasis during Gram-negative endotoxemia. *Innate Immun* **18**, 89-99 (2012). https://doi.org:10.1177/1753425910390722

Harris, H. W., Grunfeld, C., Feingold, K. R. & Rapp, J. H. Human very low density lipoproteins and chylomicrons can protect against endotoxin-induced death in mice. *J Clin Invest* **86**, 696-702 (1990). https://doi.org:10.1172/JCI114765

Vreugdenhil, A. C., Snoek, A. M., van 't Veer, C., Greve, J. W. & Buurman, W. A. LPS-binding protein circulates in association with apoB-containing lipoproteins and enhances endotoxin-LDL/VLDL interaction. *J Clin Invest* **107**, 225-234 (2001). https://doi.org:10.1172/JCI10832

Wolfe, R. R., Shaw, J. H. & Durkot, M. J. Effect of sepsis on VLDL kinetics: responses in basal state and during glucose infusion. *Am J Physiol* **248**, E732-740 (1985). https://doi.org:10.1152/ajpendo.1985.248.6.E732

Read, T. E. *et al.* Triglyceride-rich lipoproteins prevent septic death in rats. *J Exp Med* **182**, 267-272 (1995). https://doi.org:10.1084/jem.182.1.267

Beyer, D. *et al.* The liver in sepsis: molecular mechanism of liver failure and their potential for clinical translation. *Mol Med* **28**, 84 (2022). https://doi.org:10.1186/s10020-022-00510-8

Arroyo, V., Moreau, R. & Jalan, R. Acute-on-Chronic Liver Failure. *N Engl J Med* **382**, 2137-2145 (2020). https://doi.org:10.1056/NEJMra1914900

Yoo, J. Y. & Desiderio, S. Innate and acquired immunity intersect in a global view of the acute-phase response. *Proc Natl Acad Sci U S A* **100**, 1157-1162 (2003). https://doi.org:10.1073/pnas.0336385100

Hao, H. *et al.* Farnesoid X Receptor Regulation of the NLRP3 Inflammasome Underlies Cholestasis-Associated Sepsis. *Cell Metab* **25**, 856-867 e855 (2017). https://doi.org:10.1016/j.cmet.2017.03.007

Paumelle, R. *et al.* Hepatic PPARalpha is critical in the metabolic adaptation to sepsis. *J Hepatol* **70**, 963-973 (2019). https://doi.org:10.1016/j.jhep.2018.12.037

Van Wyngene, L. *et al.* Hepatic PPARalpha function and lipid metabolic pathways are dysregulated in polymicrobial sepsis. *EMBO Mol Med* **12**, e11319 (2020). https://doi.org:10.15252/emmm.201911319

Yamamoto, M., Kensler, T. W. & Motohashi, H. The KEAP1-NRF2 System: a Thiol-Based Sensor-Effector Apparatus for Maintaining Redox Homeostasis. *Physiol Rev* **98**, 1169-1203 (2018). https://doi.org:10.1152/physrev.00023.2017

Zhang, Y. & Xiang, Y. Molecular and cellular basis for the unique functioning of Nrf1, an indispensable transcription factor for maintaining cell homoeostasis and organ integrity. *Biochem J* **473**, 961-1000 (2016). https://doi.org:10.1042/BJ20151182

Bartelt, A. & Widenmaier, S. B. Proteostasis in thermogenesis and obesity. *Biol Chem* **401**, 1019-1030 (2020). https://doi.org:10.1515/hsz-2019-0427

Kitteringham, N. R. *et al.* Proteomic analysis of Nrf2 deficient transgenic mice reveals cellular defence and lipid metabolism as primary Nrf2-dependent pathways in the liver. *J Proteomics* **73**, 1612-1631 (2010). https://doi.org:10.1016/j.jprot.2010.03.018

Lee, C. S., Ho, D. V. & Chan, J. Y. Nuclear factor-erythroid 2-related factor 1 regulates expression of proteasome genes in hepatocytes and protects against endoplasmic reticulum stress and steatosis in mice. *FEBS J* **280**, 3609-3620 (2013). https://doi.org:10.1111/febs.12350

Tsujita, T. *et al.* Transcription factor Nrf1 negatively regulates the cystine/glutamate transporter and lipid-metabolizing enzymes. *Mol Cell Biol* **34**, 3800-3816 (2014). https://doi.org:10.1128/MCB.00110-14

Thimmulappa, R. K. *et al.* Nrf2 is a critical regulator of the innate immune response and survival during experimental sepsis. *J Clin Invest* **116**, 984-995 (2006). https://doi.org:10.1172/JCI25790

Akl, M. G. *et al.* Complementary gene regulation by NRF1 and NRF2 protects against hepatic cholesterol overload. *Cell Rep* **42**, 112399 (2023). https://doi.org:10.1016/j.celrep.2023.112399

Akl, M. G., Baccetto, R., Stebbings, B. M., Li, L. & Widenmaier, S. B. Euglycemia is affected by stress defense factor hepatocyte NRF1, but not NRF2. *Biochem Biophys Res Commun* **668**, 96-103 (2023). https://doi.org:10.1016/j.bbrc.2023.05.082

Trites, M. J. *et al.* HDL functionality is dependent on hepatocyte stress defense factors Nrf1 and Nrf2. *Front Physiol* **14**, 1212785 (2023). https://doi.org:10.3389/fphys.2023.1212785

Hirotsu, Y., Hataya, N., Katsuoka, F. & Yamamoto, M. NF-E2-related factor 1 (Nrf1) serves as a novel regulator of hepatic lipid metabolism through regulation of the Lipin1

and PGC-1beta genes. *Mol Cell Biol* **32**, 2760-2770 (2012). https://doi.org:10.1128/MCB.06706-11

Ohtsuji, M. *et al.* Nrf1 and Nrf2 play distinct roles in activation of antioxidant response element-dependent genes. *J Biol Chem* **283**, 33554-33562 (2008). https://doi.org:10.1074/jbc.M804597200

Widenmaier, S. B. *et al.* NRF1 Is an ER Membrane Sensor that Is Central to Cholesterol Homeostasis. *Cell* **171**, 1094-1109 e1015 (2017). https://doi.org:10.1016/j.cell.2017.10.003

Xu, Z. *et al.* Liver-specific inactivation of the Nrf1 gene in adult mouse leads to nonalcoholic steatohepatitis and hepatic neoplasia. *Proc Natl Acad Sci U S A* **102**, 4120-4125 (2005). https://doi.org:10.1073/pnas.0500660102

Kamisako, T. *et al.* Role of Nrf2 in the alteration of cholesterol and bile acid metabolism-related gene expression by dietary cholesterol in high fat-fed mice. *J Clin Biochem Nutr* **54**, 90-94 (2014). https://doi.org:10.3164/jcbn.13-92

Okada, K. *et al.* Deletion of Nrf2 leads to rapid progression of steatohepatitis in mice fed atherogenic plus high-fat diet. *J Gastroenterol* **48**, 620-632 (2013). https://doi.org:10.1007/s00535-012-0659-z

Sugimoto, H. *et al.* Deletion of nuclear factor-E2-related factor-2 leads to rapid onset and progression of nutritional steatohepatitis in mice. *Am J Physiol Gastrointest Liver Physiol* **298**, G283-294 (2010). https://doi.org:10.1152/ajpgi.00296.2009

Defalque, D., Brandt, N., Ketelslegers, J. M. & Thissen, J. P. GH insensitivity induced by endotoxin injection is associated with decreased liver GH receptors. *Am J Physiol* **276**, E565-572 (1999). https://doi.org:10.1152/ajpendo.1999.276.3.E565

Wang, X. *et al.* Endotoxin-induced proteolytic reduction in hepatic growth hormone (GH) receptor: a novel mechanism for GH insensitivity. *Mol Endocrinol* **22**, 1427-1437 (2008). https://doi.org:10.1210/me.2007-0561

Radhakrishnan, S. K., den Besten, W. & Deshaies, R. J. p97-dependent retrotranslocation and proteolytic processing govern formation of active Nrf1 upon proteasome inhibition. *Elife* **3**, e01856 (2014). https://doi.org:10.7554/eLife.01856

Radhakrishnan, S. K. *et al.* Transcription factor Nrf1 mediates the proteasome recovery pathway after proteasome inhibition in mammalian cells. *Mol Cell* **38**, 17-28 (2010). https://doi.org:10.1016/j.molcel.2010.02.029

Sander, L. E. *et al.* Hepatic acute-phase proteins control innate immune responses during infection by promoting myeloid-derived suppressor cell function. *J Exp Med* **207**, 1453-1464 (2010). https://doi.org:10.1084/jem.20091474

Mai, S. H. C. *et al.* Body temperature and mouse scoring systems as surrogate markers of death in cecal ligation and puncture sepsis. *Intensive Care Med Exp* **6**, 20 (2018). https://doi.org:10.1186/s40635-018-0184-3

Cuchel, M. *et al.* Inhibition of microsomal triglyceride transfer protein in familial hypercholesterolemia. *N Engl J Med* **356**, 148-156 (2007). https://doi.org:10.1056/NEJMoa061189

Kong, X. *et al.* Enhancing Nrf2 pathway by disruption of Keap1 in myeloid leukocytes protects against sepsis. *Am J Respir Crit Care Med* **184**, 928-938 (2011). https://doi.org:10.1164/rccm.201102-0271OC

Thimmulappa, R. K. *et al.* Nrf2-dependent protection from LPS induced inflammatory response and mortality by CDDO-Imidazolide. *Biochem Biophys Res Commun* **351**, 883-889 (2006). https://doi.org:10.1016/j.bbrc.2006.10.102

Shrum, B. *et al.* A robust scoring system to evaluate sepsis severity in an animal model. *BMC Res Notes* **7**, 233 (2014). https://doi.org:10.1186/1756-0500-7-233

Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884-i890 (2018). https://doi.org:10.1093/bioinformatics/bty560

Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013). https://doi.org:10.1093/bioinformatics/bts635

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org:10.1186/s13059-014-0550-8

# Figure Legends

**Figure 1. Hepatic Nrf1 is important for surviving LPS-induced sepsis**. A) Survival plot of mice injected with indicated doses of LPS (n = 3-5 males; 3-6 females). B-C) Immunoblot (B) and its analysis (C), normalized by Lamin A/C, of liver nuclei isolated 8-hours post injection with 20 mg/kg LPS (n = 3 males; 3 females). D) Expression of Nrf1 and Nrf2 target genes in liver 8-hours post injection with 20 mg/kg LPS, normalized by *36b4* (n = 8 males; 8 females). E) Study design and survival plots of indicated flox mice infected as indicated in panel (n = 6-14 males; 6-13 females). p-value in C and D determined by t-test. p-value in E determined by Log rank (Mantel-Cox) test. In A and E, number of survivors in each group is shown in symbol legend. Data in C and D are mean ± standard error of the mean, with individual data points shown.

**Figure 2. Hepatic Nrf1 deficiency alters plasma triglyceride, but not cytokines or glycemia.** A-C) Plasma cytokines (A), glucose (B), and triglyceride (C) for indicated flox mice infected as indicated in panel (n = 2-7 males; 2-8 females). p-value in A determined by unpaired t-test, with adjustment for multiple comparisons. p-value in B and C determined by two-way ANOVA, with Tukey's post-test. Data in A-C are box and whisker plots, with individual data points shown.

**Figure 3. Increasing hepatic Nrf1 activity improves survival during sepsis and affects plasma triglyceride and free fatty acid in an inverse manner to hepatic Nrf1 deficiency**. A) Study design and survival plot of mice infected with AAV-GFP or AAV expressing human NRF1 in hepatocytes (AAV-hNRF1) and injected with 20 mg/kg LPS (n = 14 males; 14 females). Number of survivors in each group is shown in symbol legend. B) Study design comparing control to *Nfe2l1^{flox/flox}* mice with hepatic Nrf1 loss- or gain-of-function 8-hours post injection with PBS or

20 mg/kg LPS. C-E) Corresponds to study design in (B). Plasma glucose (C), triglyceride (D), and free fatty acid (E) in PBS and LPS injected *Nfe2ll^flox/flox* mice (n = 7-8 males; 8 females). p-value in A determined by Log rank (Mantel-Cox) test. p-value in C-E determined by two-way ANOVA, with Tukey's post-test. Data in C-E are box and whisker plots, with individual data points shown.

**Figure 4. Effect of increasing hepatic Nrf1 activity on liver gene expression during sepsis**. A) Compared to control mice (AAV-GFP), pathways enriched in liver of *Nfe2ll^flox/flox* mice with hepatic Nrf1 gain-of-function (AAV-hNRF1) 8-hours post injection with 20 mg/kg LPS, (n = 4 males; 4 females). p-value determined by Benjamin Hochberg analysis. B) Heatmap of relative expression for genes corresponding to pathway disease annotation: necrosis. Shown are log 2 values after transforming to Z-score.

**Figure 5. Hepatic Nrf1 is important for surviving *E. coli*-induced sepsis, corresponding with increasing plasma triglyceride and protection against hypothermia.** A) Study design and survival plot comparing control (AAV-GFP) to *Nfe2ll^flox/flox* mice with hepatic Nrf1 loss- (AAV-CRE) or gain-of-function (AAV-hNRF1) in response to 4.0 x 10^8 CFU *E. coli* (n = 9-10 males) or 3.2 x 10^8 CFU *E. coli* (n = 10 females). Number of survivors in each group is shown in symbol legend. B-D) Comparison of control to *Nfe2ll^flox/flox* mice with hepatic Nrf1 loss- or gain-of-function 24-hours post-injection of *E. coli*. Shown is glucose (B) in plasma (n = 10 males; 9-10 females), triglyceride (C) in plasma (n = 5 males; 5 females), and rectal temperatures (D) assessing hypothermia (n = 10 males; 9-10 females). p-value in A determined by Log rank (Mantel-Cox) test. p-value in B-D determined by two-way ANOVA with Tukey's post-test. Data in B-D are box and whisker plots, with individual data points shown.

**Figure 6. Hepatic Nrf1 promotes VLDL-dependent liver defenses that protect against sepsis severity and hypothermia.** A) Study design comparing control (AAV-GFP) to *Nfe2l1flox/flox* mice with hepatic Nrf1 loss- (AAV-CRE) or gain-of-function (AAV-hNRF1) 0-24 hours post-injection with 1 mg/kg LPS. B-D) Rectal temperature (B), sepsis score (C), and plasma triglyceride (D) 0-24 hours post-injection with LPS (n = 4-6 males; 4-6 females). E) VLDL secretion 16 hours post-injection with LPS. Plasma triglyceride was measured 0-4 hours after injecting 1 g/kg p-407, which inhibits metabolism of circulating triglyceride by lipoprotein lipase (n = 4 males; 4 females). F-G) Rectal temperature (F) and sepsis score (G) 0-24 hours post-injection with LPS ± co-injection with 2 mg/kg of hepatic VLDL secretion inhibitor, lomitapide (n = 4-6 males; 4-7 females). p-value in B-G determined by two-way ANOVA with Tukey's post-test. Data in B-G are box and whisker plots, with individual data points shown.

# Figure 1.

**A.** Survival response to lipopolysaccharide (LPS)

- PBS (3/3 males; 3/3 females)
- 1 mg/kg LPS (3/3 males; 3/3 females)
- 5 mg/kg LPS (2/3 males; 3/3 females)
- 10 mg/kg LPS (3/5 males; 4/6 females)
- 20 mg/kg LPS (1/4 males; 0/6 females)
- 30 mg/kg LPS (0/4 males; 0/3 females)

**B.** Immunoblot on liver nuclei 8 hours post-injection

**C.** Immunoblot quantification

**D.** Expression of Nrf1 and Nrf2 targets 8 hours post-injection with 20 mg/kg LPS or PBS

**E.**

in vivo systems
CRE/LoxP (flox) mouse lines:
1) *Nfe2l1^flox/flox* mice
2) *Nfe2l2^flox/flox* mice
3) *Nfe2l1^flox/flox*; *Nfe2l2^flox/flox* mice

experiment timeline
day 0: infect with AAV-GFP or AAV-CRE
day 12: inject 10 mg/kg LPS
day 15: complete survival assessment

10 mg/kg LPS injection to *Nfe2l1^flox/flox* mice
- AAV-GFP (8/13 males; 10/13 females)
- AAV-CRE (2/14 males; 5/13 females)

10 mg/kg LPS injection to *Nfe2l2^flox/flox* mice
- AAV-GFP (8/10 males; 2/10 females)
- AAV-CRE (4/10 males; 5/10 females)

10 mg/kg LPS injection to *Nfe2l1^flox/flox*; *Nfe2l2^flox/flox* mice
- AAV-GFP (3/6 males; 4/6 females)
- AAV-CRE (1/6 males; 2/7 females)

# Figure 2.

## A.

Plasma cytokines 8 hours post-injection with 10 mg/kg LPS or PBS

## B.

Plasma glucose 8 hours post-injection with 10 mg/kg LPS or PBS

## C.

Plasma triglyceride 8 hours post-injection with 10 mg/kg LPS or PBS

# Figure 3.

## A.

Experiment design: effect of Nrf1 gain-of-function on survival response to 20 mg/kg LPS

| in vivo system | AAV treatment | experiment timeline | | |
|---|---|---|---|---|
| C57bl/6J mice | 1. AAV-GFP (control)<br><br>2. AAV-hNRF1 (human NRF1 over-expression for gain-of-function effect) | day 0: infect with AAV-GFP or AAV-hNRF1 | day 12: inject 20 mg/kg LPS | day 15: survival assessment |

Survival response to 20 mg/kg LPS

— AAV-GFP (2/14 males; 1/14 females)
— AAV-hNRF1 (7/14 males; 6/14 females)

p<0.001

## B.

Experiment design: effect of Nrf1 loss- and gain-of-function on samples collected 8 hours post-LPS injection

| in vivo system | AAV treatment | experiment timeline | | |
|---|---|---|---|---|
| CRE/LoxP (flox) mouse line:<br><br>*Nfe2l1<sup>flox/flox</sup>* mice | 1. AAV-GFP (control)<br>2. AAV-CRE (loss-of-function)<br>3. AAV-hNRF1 (gain-of-function effect) | day 0: infect with AAV-GFP or AAV-hNRF1 | day 12: inject 20 mg/kg LPS | 8 hours later: collect samples for analysis |

## C.

Effect on plasma glucose

## D.

Effect on plasma triglyceride

## E.

Effect on plasma free fatty acid

# Figure 4.

**A.** Molecular activity prediction of ingenuity pathway analysis: AAV-hNRF1 compared to AAV-GFP

**B.** Gene expression heatmap on genes corresponding to the disease annotation 'necrosis': AAV-hNRF1 compared to AAV-GFP

# Figure 5.

**A.**  Experiment design: effect of Nrf1 gain-of-function on survival response to *E. coli* infection

| in vivo system | AAV treatment | experiment timeline | | |
|---|---|---|---|---|
| CRE/LoxP (flox) mouse line:<br><br>*Nfe2l1*<sup>flox/flox</sup> mice | 1. AAV-GFP (control)<br>2. AAV-CRE (loss-of-function)<br>3. AAV-hNRF1 (gain-of-function effect) | day 0: infect with AAV-GFP, AAV-CRE or AAV-hNRF1 | day 12: inject with *E. coli* | day 17: complete survival assessment |

Survival response to *E. coli* infection

AAV-GFP (5/10 males; 6/10 females)
AAV-CRE (1/10 males; 2/10 females)
AAV-hNRF1 (8/9 males; 8/10 females)

p<0.05
p<0.01

**B.**  Effect on plasma glucose

**C.**  Effect on plasma triglyceride

**D.**  Effect on hypothermia

# Figure 6.

**A.** Experiment design: impact of Nrf1 loss- and gain-of-function on sepsis related LPS effects

| in vivo system | AAV treatment | experiment timeline |
|---|---|---|
| CRE/LoxP (flox) mouse line:<br><br>*Nfe2l1^flox/flox* mice | 1. AAV-GFP (control)<br>2. AAV-CRE (loss-of-function)<br>3. AAV-hNRF1 (gain-of-function effect) | day 0: infect with AAV-GFP, AAV-CRE or AAV-hNRF1    day 12: inject 1 mg/kg LPS    up to 24 hours later:<br>- measure temperature<br>- collect sample |

**B.** Effect on hypothermia

**C.** Effect on sepsis severity

**D.** Effect on plasma triglyceride

**E.** Effect on VLDL secretion 16 hours post-LPS injection

**F.** Effect of hepatic VLDL secretion inhibitor, lomitapide, on hypothermia

**G.** Effect of hepatic VLDL secretion inhibitor, lomitapide, on sepsis severity