Hierarchical cortical plasticity in congenital sight impairment

Roni O. Maimon-Mor[1,2], Mahtab Farahbakhsh[1,2], Nicholas Hedger[3], Andrew T. Rider[1], Elaine J. Anderson[1,4,5], Geraint Rees[4,5], Tomas Knapen[6,7], Michel Michaelides[1,8], Tessa M. Dekker[1,2]

[1] UCL Institute of Ophthalmology, University College London, London EC1V 9EL, UK
[2] Experimental Psychology, University College London, London WC1H 0AP, UK
[3] Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading, RG6 6AL, UK
[4] UCL Institute of Cognitive Neuroscience, University College London, London WC1N 3AZ, UK
[5] The Wellcome Centre for Human Neuroimaging, University College London, London WC1N 3AR, UK
[6] Spinoza Centre for Neuroimaging, Royal Netherlands Academy of Sciences, 105 BK Amsterdam, The Netherlands
[7] Cognitive Psychology, Faculty of Behavioural and Movement Sciences, Vrije Universiteit, 1081 BT Amsterdam, The Netherlands
[8] Moorfields Eye Hospital, London EC1V 2PD, UK

Abstract

A robust learning system balances adaptability to new experiences with stability of its foundational architecture. To investigate how the human brain implements this we used a new approach to study plasticity and stability across hierarchical processing stages in visual cortex. We compare the rod system of individuals born with rod-only photoreceptor inputs (achromatopsia) to the typically developed rod system, allowing us to dissociate impacts of life-long versus transient responses to altered input. Cortical input stages (V1) exhibited high stability, with structural hallmarks of deprivation and no retinotopic reorganisation. However, plasticity manifested as reorganised read-out of these inputs by higher-order cortex, in a pattern that could compensate for the lower resolution of a rod-only system and its lack of high-density foveal input. We propose that these hierarchical dynamics robustly optimize processing of available input and could reflect a broader principle of brain organisation with important implications for emerging sight-recue therapies.

Keywords: achromatopsia; neural plasticity; fMRI; pRF modelling; connective field modelling; visual impairment;

Introduction

The need for neural systems to be robust on one hand but have plasticity to adapt to
experience on the other, places opposing pressures on the formation of the brain's neural
architecture. Historically, there have been two central contrasting views: early studies
suggested that the brain exhibits remarkable plasticity, with the capacity for significant
rewiring and reorganisation in response to new or altered experiences (Baker et al., 2005,
2008; Baseler et al., 2002; Calford et al., 2005; Dilks et al., 2009; Ferreira et al., 2016;
Schumacher et al., 2008). More recent work portrays the brain as a fundamentally stable
structure resistant to functional reorganization even when primary sensory inputs are lost
(Adams et al., 2007; Baseler et al., 2011; Goesaert et al., 2014; Molz et al., 2023; Ritter et
al., 2019). Our study reframes this debate by investigating how adaptations of higher-order
processing stages can compensate for stability of sensor-to-cortex connections, offering a
new perspective on neuroplasticity through the lens of hierarchical network dynamics.

In the visual domain, the focal point of the debate on plasticity and stability has hinged on
the extent to which retinal input deprivation can drive local reorganisation in early visual
cortex (Adams et al., 2007; Baker et al., 2005, 2008; Baseler et al., 2002, 2011; Calford et
al., 2005; Dilks et al., 2009; Dumoulin & Knapen, 2018; Ferreira et al., 2016; Goesaert et al.,
2014; Haak et al., 2015; Molz et al., 2023; Ritter et al., 2019; Schumacher et al., 2008) . In
reality visual impairment is a more global phenomenon with complex dynamics beyond
constricted local retinocortical projection zones (Carvalho et al., 2019). Investigating these
dynamics is crucial for understanding neural plasticity bottlenecks of regenerative treatments
under trial in human patients (Fischer et al., 2020; Georgiou et al., 2021; Michaelides et al.,
2023). In these treatments, signal is recovered at the level of the sensory organ, here the
eye. If tissue at any downstream processing stage has deteriorated due to inactivation or
has adopted new roles, it may be less responsive to its original signals. For ground-breaking
sight recovery treatments to reach their full potential, understanding these cascading
dynamics in the human brain is now paramount.

We revisit the debate on neuroplasticity and stability by investigating adaptations to altered
sensory input across hierarchical cortical processing stages, using the inherited retinal
disease achromatopsia as a model. Achromatopsia causes cone photoreceptors in the retina
to be inactive from birth (Aboshiha et al., 2014). This means visual development is purely
driven by rod photoreceptors. Rod-mediated vision causes low acuity, a blind spot (scotoma)
in the rod-free retina of the fovea, and no colour vision. In most studies on how the brain
adapts to sight loss, a major challenge is accurately matching the sensory input experienced
by patients in control individuals with typical sight (Brown et al., 2021; Jones et al., 2020;

Macnamara et al., 2021). This matching is critical for distinguishing neural adaptations to

long-term altered visual experience from normal transient responses to varying visual input

(Binda et al., 2013; Carvalho et al., 2021; Haak et al., 2012; Hummer et al., 2018). While

normally sighted individuals rely on cone vision most of their waking time, rod only vision can

also naturally occur or be induced experimentally (e.g. Barton & Brewer, 2015). This makes

it possible to directly compare lifelong rod-only vision in achromatopsia with transient rod-

only experiences in normal sight, distinguishing the long-term neural impacts of altered

sensory input from temporary changes within the typical system's range of flexibility.

It remains unclear how the brain adapts to altered retinal input in achromatopsia. Initially, a

functional MRI study showed that areas of the early visual cortex, typically receiving only

foveal cone input, might reorganise to process input from peripheral rod-containing regions

(Baseler et al., 2002). This suggests potential cortical plasticity. However, recent findings by

Molz et al. (2023) present a conflicting view, showing no reorganization in the deprived

primary visual cortex of achromats, and structural signs of deprivation (Molz et al., 2022).

This suggests functional stability. Here, we reasoned that both stability and plasticity in a

deprived visual system can co-exist across different cortical processing stages; A high

degree of stability in sensor-cortex connections (Haak et al., 2015; Molz et al., 2023) may be

offset by plasticity in downstream regions in the hierarchy, which may exhibit more scope to

flexibly adapt to optimise the use of available input.

To test this prediction, we drew on a uniquely large sample of achromats and controls. We

first characterised spatial tuning and remapping of V1 in a cone-input deprived visual system

to resolve ambiguity of earlier findings. We then used a connective field modelling approach

to assess sampling of V1 information by higher order visual cortex (V3). To truly distinguish

long-term adaptations from transient changes to altered input, we took a different approach

than prior studies. Rather than only manipulating luminance levels, we utilised the method of

silent-substitution (Estévez & Spekreijse, 1982) to create photoreceptor-specific population

receptive field (pRF) mapping that only stimulates rods. Crucially, this allowed us to obtain

well-matched, high-quality rod-driven maps from both achromats and controls, enabling

unbiased comparison of rod-mediated responses across the groups.

Using these specialised stimuli to characterise the impacts of purely rod-based vision on the

visual systems of achromats and normally-sighted controls, we first show that V1 retinotopy

presents the hallmarks of a highly robust system. There is no evident remapping of cortical

regions that are input-deprived under rod-only viewing circumstances, despite their life-long

altered input in achromatopsia. We found that the robustness to long-term altered input in

V1, is paired with reorganisation of retinotopic read-out by higher-order visual cortex (V3). A

comparison of the visual space organization of V1's connectivity with V3, reveals a shift in

the distribution of resources in achromats: more cortical resources in V3 are dedicated to

sampling the V1 cortical surface that encodes 1°-4° visual angles - the most foveal areas

accessible to rods. V3 also samples from fewer V1 vertices, effectively increasing its spatial

resolution. These changes in the rod-based visual system of achromats could act to

compensate for the absence of two features of the cone system that the rod system lacks: its

high-density foveal sampling and its overall higher resolution. This may reflect an

overarching broader organising principle where hierarchical optimisation processes allow the

brain to make more efficient use of the available (in this case atypical) input.

Results

113   In three results sections, we characterise the interplay between neural plasticity and stability

114   in achromatopsia across three complementary neural markers: brain structure, spatial tuning

115   characteristics of the functional MRI signal in V1, and the hierarchical retinotopic connectivity

116   patterns that indicate how higher-order visual cortex (V3) sample information from V1.

Structural analysis

As a proxy of structural integrity, we compared cortical thickness across 8 eccentricity

segments of V1 (0°-1°,1°-2°,2°-3°, etc.; see Figure 1B) in a large sample of 31 achromats

and 100 controls aged 6-35 years. Data were collated over two scanners. We expected

changes in achromats to be largest in the central 0°-2° eccentricities of the visual field,

encompassing the retinal 'rod-free' zone, an oval of approximately 1.25°x0.96° in size

(Curcio et al., 1990). V1 segments were defined using the Benson atlas (Benson & Winawer,

2018), which is fitted using cortical landmarks on individual brains. The atlas fit produces

predicted eccentricities to each vertex, this allows us to identify where the representation of

these eccentricities would have been given typical development and investigate what they

represent in atypical development.

Figure 1A shows an increase in cortical thickness in achromats compared to controls only in

the foveal projection zone (Controlling for scanner, overall cortical thickness, and age; 0°-1°

ANCOVA group effect: $F_{(1,126)=}9.05$, p_{unc}=0.003; 1°-2°: $F_{(1,126)=}7.79$, p_{unc}=0.006). This

increase was not present in more eccentric parts of V1 (p_{unc}>0.2 for all). This pattern is in

line with previously reported V1 foveal thickening in adult achromats (Molz et al., 2022), and

moreover suggests that this change is already present in late childhood. A further

comparison of the cortical thickness in the foveal projection zones (0°-2°) in V2 and V3

(Figure 1A right), revealed a trend toward significance in V2 ($F_{(1,26)}$=3.07, p=0.08) and a significant effect in V3 ($F_{(1,126)}$=6.32, p=0.01). As thicker cortex in early visual areas is a hallmark of visual deprivation from birth (Jiang et al., 2009; Park et al., 2009), this indicates that the foveal projection zones in achromats may lack retinal input. This provides evidence for stability, because if these cortical areas had substantially reorganised to receive more peripheral retinal input, we would not expect to see this marker of deprivation.

Figure 1 – **Structural properties of V1 in achromats and controls.** (**A**) Mean V1 cortical thickness of each eccentricity based V1 segment in millimetres (mm). Cortical thickness was increased in achromats (n=31) compared to controls (n=100) only for foveal projection segments 0°-2° and not in more eccentric segments 2°-8°. This pattern was trending in V2 and significant in V3. Error bars are 95% confidence intervals (**B**) Visualisation of regions of interest in one control participant. Segmentation of V1 to 1° eccentricity bins, and the foveal projection zone (0°-2°) of V2 and V3 was based on the Benson retinotopy atlas (Benson and Winawer,2018). The half circle on the right shows, in visual space, the area represented by each segment.

pRF properties of V1

We next test whether the visual-spatial tuning of V1 in a system that has developed using only rod input (i.e., achromats) remains stable or adapts, using population receptive field modelling. Our analyses focus on two types of plasticity. First, we ask what happens to the V1 cortex normally receiving only cone-mediated input from the foveal 'rod-free' zone, which shows structural abnormalities. Second, we test for changes in rod-selective visuospatial tuning across V1 (beyond the 0°-2° 'rod-free' fovea), in cortex which normally receives combinations of rod and cone input but develops with only rod inputs in achromats.

V1 receives retinotopically organised input, with adjacent neurons in the cortex receiving signals from adjacent neurons in the retina. Population receptive field modelling links retinotopic voxels in visual cortex to the locations in the visual field they represent (Dumoulin & Wandell, 2008). This is achieved by fitting population receptive fields estimates (pRFs; i.e.,

Gaussian tuning functions) to BOLD responses evoked by stimuli systematically traversing

the visual field. To compare rod-mediated cortex function in achromatopsia to normally

developed rod-responses under matched viewing circumstances, we used the silent

substitution method (Estévez & Spekreijse, 1982) and low luminance presentation which

allowed us to generate stimuli with a rod-isolating contrast ("rod-selective condition"). We

also included another stimulus condition that, in a typical retina, stimulates cones and the

rods equally ("non-selective condition"). This allowed us to assess how rod-mediated

measures in both groups compare to more typical pRF measures driven by cones in normal

sighted, and whether differences across achromats and controls remain consistent across

different visual stimulus properties.

Coverage and pRF properties of the foveal projection zone

We first compared retinotopic tuning in cortex containing the rod-free foveal projection zone

(atlas defined 0°-2°) across controls and achromats. In normally developed retinotopic rod

maps, the cortical projection of the foveal rod-free zone, appears as a central "hole" (Figure

2B; Barton & Brewer, 2015). To test if this normally silent area fills in with reorganised

retinotopic visual activity in achromats as has previously been reported (Baseler et al.,

2002), we compared the percentage of cortex that contains reliable pRFs (coverage;

meaning where pRFs could reliably be estimated from visually-driven signal. This is shown

in Figure 2 for pRF model fits of $R^2 > 0.1$, but effects were similar at $R^2 > 0.2$ and 0.3.

Figure 2 **– pRF properties of V1 in achromats and controls.** (**A**) Foveal projection zone

(0°-2°) coverage. Coverage is the fraction of vertices within the foveal confluence that

contain visual information (i.e., pRF R^2>0.1). Achromats (red markers) are not significantly

different from controls (black markers) in the rods only condition (square markers), and do

not show complete filling in even in the non-selective condition with higher contrast (triangle

markers). Triangle markers: group mean for the non-selective condition, square markers:

group mean for the rods only condition. Black markers: controls group mean, red markers:

achromat group means. Black/red vertical lines around the mean: 95% confidence interval,

grey shaded lines: 2.5%-97.5% percentile interval (only in controls). For achromats,

participants' dots are colour-coded to match across conditions and measures (**B**) Group

eccentricity plots on a flat surface for each condition (in standard fsaverage space),

presenting the group median value of each vertex. For visualisation purposes, vertices that

were absent in more than 30% of the group were left blank in each condition. Black lines

denote delineations of V1,2,3; white lines mark the foveal projection zone (0°-2°) in V1. As

shown in (A) we see no evidence of 'filling in' of the rod scotoma in achromats. (**C**) pRF

sizes along the atlas V1 segments. Achromats show a consistent increase in pRF size

compared to controls. (**D**) Eccentricity of modelled pRFs along the atlas V1 segments.

Achromats show an increase in eccentricity compared to controls, however that shift is not

consistent across the two stimuli types. Vertical lines in C&D represent 95% confidence

intervals.

In the rod-selective condition, the coverage in the 0°-2° region of interest was near-identical

across achromats and controls (on average 24% in controls, and 25% in achromats, $t_{(47)} =$

0.19, p=0.85, BF_{10}=0.31; see Figure 2A&B). For non-selective stimuli targeting both cones

and rods, controls exhibit greater visual coverage (75% on average) compared to

achromats, who are missing functional cones (43% on average) (Mann $U_{(52)}$=63, p<0.0001;

unequal variance reverted to non-parametric). Achromats show enhanced coverage under

non-selective compared to rod-selective conditions, likely due to increased luminance

enhancing the rod system's signal-to-noise ratio. We note that two achromats show potential

filling in with a foveal activation coverage beyond the 95% predictive interval of the normal

sighted rod gap. Their retinotopic profiles (see Supplement 1) suggest this may reflect

individual differences in retinal foveal photoreceptor distribution profiles rather than cortical

remapping. This highlights the importance of characterising variability in atypically

developing populations through larger samples rather than case studies.

Crucially, data quality was matched across groups, with no differences in pRF model fits

(see Supplement 2). It is therefore unlikely that any coverage inside the rod "hole" in

achromats was masked by factors such as eye-movement, visual responsiveness, or head-

movement. In line with this, we found no simple relationship between fixation stability and

coverage (Supplement 2). In addition, it is unlikely that these findings can be explained by

pRF fitting-related artefacts (Binda et al. 2013), as modelling the rod-scotoma in the pRF

fitting procedure gave similar results (see methods and Supplement 3).

Together these results provide strong evidence for stability in the face of life-long

deprivation, i.e., the conservation of the sensory topography. This is in line with the structural

changes in this cortical area presented above, and recent conclusions by an independent

group (Molz et al., 2023).

Altered rod representation in V1 - pRF size and eccentricity

We next tested for differences in spatial tuning across V1 regions that typically receive both rod and cone inputs but develop with only rod inputs in achromatopsia. Figure 2C shows that pRF sizes in achromats are systematically larger compared to controls across all eccentricities. This holds true for both the non-selective (V1 segments: 2°-3°: $t_{(52)}$=12.57, p_{unc}<0.001; 3°-4°: $t_{(52)}$=11.12, p_{unc}<0.001; 4°-5°: $t_{(52)}$=9.54, p_{unc}<0.001; 5°-6°: $t_{(52)}$=7.52, p_{unc}<0.001; 6°-7°: $t_{(52)}$=5.76, p_{unc}<0.001; 7°-8°: $t_{(52)}$=2.36, p_{unc}=0.022; all segments remain significant after FDR correction), and rod-selective stimulus conditions (V1 segments: 2°-3°: $t_{(48)}$=3.83, p_{unc}<0.001; 3°-4°: $t_{(48)}$=3.67, p_{unc} <0.001; 4°-5°: $t_{(48)}$=4.68, p_{unc} <0.001; 5°-6°: $t_{(48)}$=4.74, p_{unc} <0.001; 6°-7°: $t_{(48)}$=3.98, p_{unc} <0.001; 7°-8°: $t_{(48)}$=2.00, p_{unc} =0.051; all segments but the 7°-8° remain significant after FDR correction). In the non-selective condition, the smaller pRFs in controls are in line with the contribution the cone system's higher spatial resolution which is not active in the achromat group. However, for the rod-selective condition where both groups activate the same class of photoreceptors, the enlarged pRFs in achromats indicate altered rod-based visual processing.

Figure 2D shows that pRF estimates in achromats have higher eccentricities across V1 compared to controls, spanning all tested V1 segments (group comparisons of the rod-selective conditions: 2°-3°: $t_{(48)}$=3.32, p_{unc}=0.002; 3°-4°: $t_{(48)}$=4.42, p_{unc}<0.001; 4°-5°: $t_{(48)}$=3.59, p_{unc}=0.001; 5°-6°: $t_{(48)}$=3.17, p_{unc}=0.003; 6°-7°: $t_{(48)}$=3.15, p_{unc}=0.003; 7°-8°: $t_{(48)}$=2.23, p_{unc}=0.030; all segments but the 7°-8° remain significant after FDR correction). However, in the non-selective condition in achromats, even though still only rods are being stimulated, estimated eccentricities are lower than those observed under the rod-selective condition. In fact, the eccentricity estimates of the non-selective condition in achromats appear similar to those of the rod-selective condition in controls. This shows that it is possible, in the achromat system, to obtain eccentricity estimates in the normal rod range. We therefore do not consider the eccentricity shifts as evidence for plasticity or stability as it is unclear whether they are a property of the tissue or reflect other factors.

Together these analyses reveal subtle differences in the rod-mediated spatial tuning of V1 cortex beyond the foveal rod-free projection zone. Notably, achromats combine information across larger areas in visual space (larger pRFs). It is not clear whether larger pRFs can be purely attributed to coarser spatial sampling at the cortical level (i.e., a degree of cortical plasticity), because this could reflect changes photoreceptor properties at the retinal level, or blurring of visual information sampling at the ocular level due to nystagmus (eye-movement). The pRF size and eccentricity differences remain unchanged when accounting for the scotoma in the pRF fitting procedure (see Supplement 3).

Connective field modelling – hierarchical sampling

Connective field modelling – hierarchical sampling

We next used connective field modelling to test whether V3's sampling along the V1 cortical

surface in achromatopsia remains stable despite the drastically altered visual input from this

region since birth. We test for two hypothetical types of hierarchical plasticity, shifts in the

resource distribution across the V1 eccentricity map (CF centre), and shifts in the aperture of

the read-out area (CF size).

Higher-order visual regions receive retinotopic information from V1, with individual neurons

integrating information across multiple neighbouring V1 neurons. Connective field (CF)

modelling is a functional connectivity-based extension of pRF modelling which characterises

this process based on time-series correlations between brain regions. While pRF modelling

is 'stimulus referred', estimating a voxel's spatial tuning properties based on its response to

stimulus variations, connective field modelling is 'neural-referred', estimating a voxel's

'connective field' based on its correlations with aggregated voxels along the primary visual

cortex surface. Responses throughout cortex are modelled as deriving from Gaussian

shaped sampling profiles over the V1 cortical surface (connective fields), defined by a

position (centre of the Gaussian) and size (Gaussian spread in mm (σ)). CF modelling can

thus reveal the topographic profile of a brain region's neural connectivity with V1, including

the spatial extent of sampling. After estimating connective fields, the CF positions can be

combined with the retinotopic map of V1 to 'project' the retinotopy of V1 onto the rest of the

brain, thereby performing connectivity-derived retinotopic mapping. To create V3 connective-

field-based eccentricity maps (Figure 3B), each vertex was assigned the Benson atlas-based

eccentricity of its connective field centre in V1. Because CF modelling relies on inherent

signal fluctuations and involves no assumptions about visual input, it is more robust to

potential confounds such as poor fixation and low visual responsiveness (Tangtartharakul et

al., 2023). Here we focus on connective fields between V3 and V1. We omit V2, which

shares a border with V1, to avoid inaccuracies in CF estimation due to mis-delineation.

A

Distribution of sampled eccentricities

visual space

B

Controls

Tagging of V3 based on sampled CF centres in V1

Achromats

Tagging of V3 based on sampled CF centres in V1

*Figure 3 – **Connective field modelling (V3) in achromats and controls – sampled eccentricities.***

*(**A**) Right: The fraction of all vertices within V3 that sample from V1, binned by sampled V1 atlas*

eccentricity. In achromats there is a shift towards greater sampling from the most foveal area that

contains rods (1°-4°) of V1 relative to the normal-sighted rod and cone systems. Shaded areas:

*standard errors of the mean (s.e.m). All V1 eccentricities are derived from the Benson Atlas. **Left:** A*

schematic illustration of the sampled eccentricities in visual space. Dashed area represents the area

*where group differences in sampling were found, 1°-4°. (**B**) Visualisation of V3 sampled eccentricities*

of V1. From left to right: Half circle represents the visual field, where each colour represents 1 visual

degree. V1 is colour-coded into segments based on the atlas-assigned eccentricities. Each vertex in

V3 is then coloured based on the centre of Gaussian connective field within the V1 surface. Values in

V3 are based on group medians. To further highlight the group differences in the amount of cortical

surface devoted to the 1°-4° V1 segments we visualise V3 again on the right, colouring in all vertices

with a CF centre at the 1°-4° segments.

Shifts in V3 resource distribution along the V1 eccentricity map

When both rods and cones are stimulated in the normal sighted system, the largest proportion of V3 voxels is dedicated to sampling information from the V1 foveal projection (connective field centres in the 0°-1° degree range), with a gradual decrease in resources allocated to larger eccentricities (Figure 3A). When only rods are stimulated, foveal sampling is attenuated but the overall pattern of a gradual decrease in sampling from small to larger eccentricities remains. In achromats, the way in which V3 samples V1 is drastically different, with most V3 resources dedicated to sampling eccentricities at 1°-4° for both stimulation conditions (Figure 3B). When comparing the rod-selective condition in achromats to that of controls, we observe a shift towards greater V3 sampling from the V1 cortex encoding the (1°-4°) paracentral visual field (0°-1°: $t_{(47)}=1.18$, $p_{unc}=0.242$; 1°-2°: $t_{(47)}=4.03$, $p_{unc}<0.001$; 2°-3°: $t_{(47)}=5.32$, $p_{unc}<0.001$; 3°-4°: $t_{(47)}=3.70$, $p_{unc}=0.001$; 4°-5°: $t_{(47)}=2.32$, $p_{unc}=0.025$; 5°-6°: $t_{(47)}=0.84$, $p_{unc}=0.405$; 6°-7°: $t_{(47)}=-0.97$, $p_{unc}=0.337$; 7°-8°: $t_{(47)}=1.93$, $p_{unc}=0.060$; Segments including 1°-4° remain significant after FDR correction). It is unlikely that these differences result from differences in measurement quality, because the goodness of fit (CF R^2) was comparable across the two groups (Supplement 4). This shift in sampling could reflect increased neural resources dedicated to the most fovea-adjacent information that is available to the rod-only system. Surprisingly, even though we found no evidence of visually-driven signal in the region on V1's surface that normally takes foveal input (0°-1°) in achromats (see Figure 2), there is still a substantial portion of V3 vertices (15%) that sample from this region. This remaining foveal connectivity between V1 and V3 shows that there is also a degree of stability across retinotopic hierarchical sampling.

Figure 4 – **Connective field modelling (V3) in achromats and controls – connective**

field size and relationships with pRF size. (**A**) CF sizes in V3 in achromats and controls

across the rod-selective and non-selective condition. Controls show an expected increase in

CF size with sampled eccentricity while achromats show a flat pattern, with smaller CFs

between 4°-6°. In both groups results are consistent across the two stimulation conditions.

(**B**) Ratio of pRF size between achromats and controls in V1 (circles) and V3 (triangles). A

ratio of 1 indicates the mean pRF size of achromats is equivalent to that of controls. Across

both non-selective (left) and rod-selective (right) stimulus conditions, the ratio of pRF sizes in

V3 is closer to 1 than in V1, pointing towards a possible normalising effect of CF size on pRF

size. (**C**) Illustrations depicting the differences in pRF and CF sizes across the groups. (**D**) A

summary of the proposed model of how smaller CF sizes in V3 of achromats might operate

as a compensatory mechanism for increased pRF size in V1, resulting in normalized pRF

size at the level of V3.

Shifts in the V3 read-out aperture from V1

In normal sighted, we show that connective field size in V3 increases with eccentricity,

whether stimulating both rods and cones at the same time, or only rods. This replicates

findings by Haak et al., (2013) and Knapen (2021). In achromatopsia, however, V3

connective field sizes remain consistently small across all eccentricities, and are significantly

smaller than in the normal-sighted rod-system between 4°-6° eccentricity (comparing

achromats vs control rod-selective condition: 0°-1°: $t_{(44)}=1.07$, $p_{unc}=0.289$; 1°-2°: $t_{(45)}=-0.41$,

$p_{unc}=0.684$; 2°-3°: $t_{(45)}=-0.58$, $p_{unc}=0.568$; 3°-4°: $t_{(46)}=-1.70$, $p_{unc}=0.097$; 4°-5°: $t_{(47)}=-3.55$,

$p_{unc}=0.001$; 5°-6°: $t_{(47)}=-2.63$, $p_{unc}=0.011$; 6°-7°: $t_{(47)}=-0.81$, $p_{unc}=0.420$; 7°-8°: $t_{(45)}=-0.73$,

$p_{unc}=0.469$; degrees of freedom vary as not all participants have vertices sampling all V1

segments). These smaller connective field sizes in V3, mean that V3 voxels pool across a

smaller number of pRFs along the V1 cortical surface (4C, middle).

In V1, pRF sizes in achromats were larger than in controls both for rod-selective and non-

selective stimuli conditions (4C, left). Finer spatial sampling of the V1 cortex by V3 could

therefore effectively compensate for this coarser input, which should result in normalisation

of pRF size in V3 (4C, right). To further investigate this possibility, we plotted the ratio

between pRF sizes in achromats and pRF sizes in controls, with values above 1 indicating

enlarged pRFs in achromats (Figure 4B). As predicted by the smaller connective fields, the

ratio of pRF sizes is closer to 1 in V3 than in V1, indicating a normalisation of pRF size in the

higher-order region relative to the input stage (Figure 4D). This is consistently the case in

both stimulus conditions. This finer hierarchical sampling could reflect a higher-order

compensation for coarser visual inputs in achromats.

Discussion

Discussion

To test the effects of long-term altered visual input on neural plasticity across the cortical

hierarchy, we compared a unique group of individuals who have developed only with rod

photoreceptor inputs (achromatopsia) to normally sighted individuals who spend most of

their waking time relying on cone-dominated vision. As normal-sighted can also experience

rod-only vision under specific circumstances, this allowed us to dissociate the effects of life-

long altered experience from transient altered sensory input. By integrating markers of

plasticity and stability across structural, representational, and connectivity analyses we

reveal system-level dynamics of stability and adaptations to altered visual experience. The

results offer a new perspective on neural plasticity and stability, recasting these processes

as two interacting forces acting simultaneously on the cortex rather, than two separate

neural responses to change.

Specifically, in achromatopsia we found thickening of the cortex around the foveal projection

zone, which receives only cone-photoreceptor inputs and therefore would be signal-deprived

from birth in this group. In contrast to early reports (Baseler et al., 2002), we found no

evidence that visually-induced signalling inside this region had increased compared to the

typically developed rod system. We also observed subtle functional changes in V1

retinotopic tuning including increased population receptive field sizes, which may reflect

retinal or post-retinal changes. These data reveal that visual spatial representations in V1

are highly stable, integrating and resolving ambiguity of earlier findings (Baseler et al., 2002;

Mckyton et al., 2021; Molz et al., 2022, 2023). This high stability at the cortical input-stage

contrasted with reorganisation of the read-out of these inputs by higher-order visual cortex.

Notably, more cortical resources in V3 were dedicated to sampling input from the near-

fovea, and V3 neuron populations aggregated information across smaller areas along the V1

cortical surface. There was also evidence for stability of sampling in achromatopsia, because

V3 retained connectivity with the foveal confluence despite the life-long scotoma.

Together these findings show that while primary sensory areas such as V1 exhibit

considerable stability, higher-order areas such as V3 display marked plasticity that may

optimise use of the available information. Both increased sampling of the most fovea-

adjacent information available to rods, and finer-grained sampling of coarser inputs could

effectively counteract the limitations of developing with a fully rod-based system. These

findings add a new dimension to the ongoing debate on neural plasticity versus stability by

proposing a hierarchical network perspective where both processes interact across

processing stages to maintain the integrity of foundational architecture, whilst optimizing the

processing of available input.

Control measures and analyses presented above rule out several alternative explanations
for these results. Model fits were similar for both patients and controls, so group differences
are unlikely to reflect data quality discrepancies. Further, eye-tracking data, available for a
subset of participants, revealed no correlation between stable fixation and enhanced foveal
activity in achromatopsia (Supplement 2), suggesting that 'filling in' of the deprived V1 cortex
was not simply obscured by nystagmus. This is further supported by converging structural
evidence of cortical thickening of this region, a typical outcome of congenital visual
deprivation (Jiang et al., 2009; Park et al., 2009). Changes in V3 sampling of V1 are also
unlikely to be a by-product of eye-movement, as connective field modelling is locked to V1
inputs rather than screen locations, so insensitive to eye movement (Tangtartharakul et al.,
2023).
406
There are convincing arguments for the importance of a system that maintains highly stable
sensor-to-cortex connections at the first cortical input stage. Re-routing of cortical projections
in response to changing visual signals could lead to "coding catastrophe", because left
unchecked it could leave visual processing areas throughout the brain with mislabelled
retinotopic signals, corresponding to different regions of visual space than they would
normally represent (Dumoulin & Knapen, 2018; Haak et al., 2015; Wandell & Smirnakis,
2010). Higher-order regions aggregate more inputs across visual space and other sensory
domains, and display greater invariance to low-level stimulus properties (Lehky & Tanaka,
2016). Therefore, there is theoretically broader scope for experience-dependent reweighting
of inputs (Makin & Krakauer, 2023), to optimise use of inputs that are still available, more
reliable, or more relevant in the impaired system.
418
We propose a new hierarchical model of neural adaptation that hypothetically could act to
improve the limitations of a rod-only visual system (i.e., lacking foveal input and reduced
acuity). However, to confirm that the adjustments we observe are compensatory rather than
mere by-products of circuitry responses to visual deprivation, their functional benefits will
need to be demonstrated. The adaptations might suggest that individuals with
achromatopsia exhibit superior rod-based vision compared to typical systems. One study
supporting this, reported improved rod-mediated biological motion processing in some
achromatopsia patients, suggesting an enhanced rod system capacity (Burton et al., 2016).
Nonetheless, most studies typically find achromatopsia patients' rod vision to be similar or
inferior, potentially because retinal structure can also be impacted by the disease (Burton et
al., 2016; Hess & Nordby, 1986). It is likely that the functional outcome is shaped by a
mixture of impairments and compensatory mechanisms at various levels, presenting an ideal
case for examining such dynamics.

In light of breakthrough treatments set up to restore cone function in human patients with

achromatopsia (Farahbakhsh et al., 2022; Fischer et al., 2020; Michaelides et al., 2023;

Reichel et al., 2022), the new plasticity dynamics revealed by our work have significant

theoretical and practical implications. The possible dormancy of retinotopic maps in cortical

input stages as well as some preserved connectivity between V1 foveal areas and V3 bodes

well for these gene therapies, each indicating remaining infrastructure to process rescued

cone signals. However, our findings also show that the interpretation of signals from V1 by

higher-order networks is modified, and it remains to be seen how this effects the translation

of new sensory inputs into functional vision. Adaptations in neuronal tuning properties vary in

how stably ingrained they become in neural architecture. They can temporarily shift in

response to task-demands (transient modulation) (e.g., Klein et al., 2018), or unfold over

longer periods with extensive experience that can be acquired or reversed at any age

(perceptual learning) (Lu & Dosher, 2022). They can also involve more profound reweighting

of neural properties that do not easily revert after critical developmental windows

(developmental divergence) (Mitchell & Maurer, 2022). A previous study found that

connective fields stay stable with central field loss in adulthood (Haak et al., 2016), raising

the possibility of early windows of plasticity for these adaptations. Understanding whether

the adaptations in achromatopsia are within the normal range of flexibility for a typically

developed system or represent a fundamental reorganisation will be of high relevance to

treatment development, as it impacts the nature and extent of neural plasticity that can be

harnessed for therapeutic recovery at different ages.

With over 50 interventional studies for inherited retinal diseases initiated, the technology to

treat currently incurable heritable retinal diseases by restoring ocular cellular structure and

function is here. The next big hurdle will be to understand neural plasticity constraints on

integrating new signals from the eye into the existing neural infrastructure. Our research

provides a novel framework for understanding the interactions of neural stability and

plasticity as the human brain adapts to altered information, which is now of crucial

importance for harnessing the full potential of emerging visual restoration therapies.

Methods

Participants

All participants met MRI safety inclusion criteria and had no other known neurological disorders. Patients with achromatopsia had genetically confirmed disease (*CNGA3* or *CNGB3* bi-allelic disease-causing variants). The main dataset was collected at 1.5T. Additional structural data was collected at 3T.

Main 1.5T dataset: We obtained functional and structural 1.5T data from fourteen individuals with achromatopsia (age range = 8-27; mean±std = 13.7±5.7), and forty normally sighted controls (age range=6-34; mean±std=19.8±8.3). Additional structural 1.5T data was collected from 8 achromats (age range=8-35; mean±std=19.75±10.6) and 50 normally sighted controls (age range=6-29; mean±std=12.5±5.9).

Additional 3T dataset (previously published in Anderson et al. 2024): We obtained additional structural scans from ten normally sighted controls (age range=20-30; mean±std=23.8±3.7) and 9 individuals with achromatopsia (age range=16-33; mean±std=24.2±6.4).

For 4 normally-sighted individuals the rod-only pRF map was not collected. One achromat was excluded from functional analysis due to large head-movements. Data collection had ethics approval from the national ethics committee for individuals with achromatopsia (REC reference: 12/LO/1196, IRAS code: 106506, REC reference 11/LO/1229) and the UCL Research Ethics Committee for normal sighted control participants (4846/001).

MRI acquisition

Structural MRI

All 1.5T participants were scanned in a Siemens Avanto 1.5T MRI scanner with a 30-channel coil (a 32-channel coil customised to remove view obstructions). A high-resolution structural scan was acquired using a T1-weighted 3D MPRAGE (1 mm^3 voxel size, Bandwidth = 190 Hz/Px, 176 partitions, partition TR = 2730, TR = 8.4ms, TE = 3.57, effective T1 = 1000 ms, flip angle = 7 degrees).

All 3T participants were scanned in a Siemens 3T TIM-Trio scanner using a 32-channel head coil. A high-resolution structural scan was acquired using a T1-weighted 3D MDEFT, 1mm isotropic voxels, 176 sagittal slices, 256x240 matrix, TE 2.48ms, TR 7.92ms, TI 910ms.

Functional MRI

Functional T2*-weighted multiband 2D echo-planar images were collected using a multi-band sequence with 2.3 mm isotropic voxels (TR = 1000 ms, TE = 55 ms, volumes = 348, flip angle = 75°, MB acceleration factor = 4, Bandwidth = 1628 Hz/Px).

pRF stimuli and apparatus

Stimuli were presented on an MR-compatible LCD display (BOLDscreen 24, Cambridge Research Systems Ltd., UK; 51 x 32 cm; 1920 x 1200 pixels) viewed through a mirror in the scanner at 105 cm distance. Participants were lying supine in the scanner, with fixation stability recorded where possible via a mirror, with an Eyelink 1000 Plus (SR Research, Ottawa, ON) at the back of bore. Built-in gaze calibration was not possible in achromats due to nystagmus so the camera was calibrated in advance on a healthy eye. Before every other run, patients fixated on a 5-point custom calibration which allowed us to calibrate gaze measures post-hoc (Tailor et al., 2021) Stimuli were presented using MATLAB (R2016b, MathWorks, Natick, MA, USA), via the Psychophysics Toolbox 3 (Kleiner et al., 2007).

pRF mapping stimuli comprised a simultaneous rotating ring and contracting/expanding wedge embedded within a contrast-reversing checkerboard with 2Hz reversal rate (Figure 1B). Per run, the ring expanded/contracted for 6 cycles (48 secs/cycle) with logarithmic eccentricity scaling (van Dijk et al., 2016). The wedge (20° angle) rotated clockwise/anticlockwise for 8 cycles (36 sec/cycle). 20-second fixation baselines were embedded at the start, at the mid-point, and end of the run (total run duration 348 secs). The stimuli covered a maximum eccentricity of 8.6°, and moved to a new position each 1-second TR. The stimulus was overlaid with a small white central fixation dot (0.2° VA radius) and a black radial grid to encourage stable fixation. To keep participants engaged, they played a rewarded "kitten rescue mission game" in which they pressed a button when detecting a fixation target luminance change.

The rod-selective (stimulating only rods) and non-selective stimuli (stimulating rods and cones equally) differed in both the colour pair (chromatic pair) used to make up the checkerboard and luminance levels. For an in depth description of stimulus generation and validation see Farahbakhsh et al., 2022. In brief, a silent substitution approach was used to obtain a rod-selective colour pair (Estévez & Spekreijse, 1982). We generated a single chromatic pair that kept L and M cone contrast constant (i.e., invisible to L and M cones), whilst inducing a contrast response in rods. Since it is only possible to simultaneously silence two photoreceptor types with 3 colour channels, the resulting stimulus was not controlled for S-cone contribution. To eradicate the S-cone response, we presented these

stimuli at very low light level (max luminance: 0.02 cd/m^2). The non-selective stimuli

chromatic pair increased rod and L and M cone contrasts equivalently. This stimulus was

presented at mesopic luminance levels (max 0.5 cd/m2).

Prior to the rods only condition, which always appeared before the non-selective condition

participants laid in the scanner for a 15-minute dark adaptation. A group of 10 normal

sighted control adults had longer period of dark adaptation (45 minutes) test any impact on

the resultant pRF maps. However, since we did not find any differences between the two

groups they were combined and treated as a single control group. We note that excluding

these 10 participants did not change any of the reported results.

MRI pre-processing

536   Data were preprocessed using *fMRIPrep* 21.0.2 (Esteban, Blair, et al., 2018; Esteban,

537   Markiewicz, et al., 2018; RRID:SCR_016216), which is based on *Nipype* 1.6.1 (K.

538   Gorgolewski et al., 2011; K. J. Gorgolewski et al., 2018; RRID:SCR_002502).

fmriprep structural data pre-processing

The T1-weighted (T1w) image was corrected for intensity non-uniformity (INU) with

N4BiasFieldCorrection (Tustison et al., 2010), distributed with ANTs 2.3.3 (Avants et al.,

2008; RRID:SCR_004757), and used as T1w-reference throughout the workflow. The T1w-

reference was then skull-stripped with a *Nipype* implementation of the

antsBrainExtraction.sh workflow (from ANTs), using OASIS30ANTs as target template. Brain

tissue segmentation of cerebrospinal fluid (CSF), white-matter (WM) and gray-matter (GM)

was performed on the brain-extracted T1w using fast (FSL 6.0.5.1:57b01774,

RRID:SCR_002823, Zhang et al., 2001). Brain surfaces were reconstructed using recon-all

(FreeSurfer 6.0.1, RRID:SCR_001847, Dale et al., 1999), and the brain mask estimated

previously was refined with a custom variation of the method to reconcile ANTs-derived and

FreeSurfer-derived segmentations of the cortical gray-matter of Mindboggle

(RRID:SCR_002438, A. Klein et al., 2017). Volume-based spatial normalization to one

standard space (MNI152NLin2009cAsym) was performed through nonlinear registration with

antsRegistration (ANTs 2.3.3), using brain-extracted versions of both T1w reference and the

T1w template. The following template was selected for spatial normalization: *ICBM 152*

Nonlinear Asymmetrical template version 2009c [Fonov et al., 2009, RRID:SCR_008796;

TemplateFlow ID: MNI152NLin2009cAsym]. Many internal operations of **fMRIPrep** use

Nilearn 0.8.1 (Abraham et al., 2014, RRID:SCR_001362), mostly within the functional

processing workflow.

fmriprep functional data pre-processing

For each BOLD run found per subject (across all tasks), the following preprocessing was performed. First, a reference volume and its skull-stripped version were generated by aligning and averaging 1 single-band references (SBRefs). Head-motion parameters with respect to the BOLD reference (transformation matrices, and six corresponding rotation and translation parameters) are estimated before any spatiotemporal filtering using mcflirt (FSL 6.0.5.1:57b01774, Jenkinson et al., 2002). The BOLD time-series were resampled onto their original, native space by applying the transforms to correct for head-motion. These resampled BOLD time-series will be referred to as *preprocessed BOLD in original space*, or just *preprocessed BOLD*. The BOLD reference was then co-registered to the T1w reference using bbregister (FreeSurfer) which implements boundary-based registration (Greve & Fischl, 2009). Co-registration was configured with six degrees of freedom. First, a reference volume and its skull-stripped version were generated using a custom methodology of *fMRIPrep*. Several confounding time-series were calculated based on the *preprocessed BOLD*: framewise displacement (FD), DVARS and three region-wise global signals. FD was computed using two formulations following Power (absolute sum of relative motions, Power et al., 2014) and Jenkinson (relative root mean square displacement between affines, Jenkinson et al., 2002). FD and DVARS are calculated for each functional run, both using their implementations in *Nipype* (following the definitions by Power et al., 2014). The three global signals are extracted within the CSF, the WM, and the whole-brain masks. Additionally, a set of physiological regressors were extracted to allow for component-based noise correction (*CompCor*, Behzadi et al., 2007). Principal components are estimated after high-pass filtering the *preprocessed BOLD* time-series (using a discrete cosine filter with 128s cut-off) for the two *CompCor* variants: temporal (tCompCor) and anatomical (aCompCor). tCompCor components are then calculated from the top 2% variable voxels within the brain mask. For aCompCor, three probabilistic masks (CSF, WM and combined CSF+WM) are generated in anatomical space. The implementation differs from that of Behzadi et al. in that instead of eroding the masks by 2 pixels on BOLD space, the aCompCor masks are subtracted a mask of pixels that likely contain a volume fraction of GM. This mask is obtained by dilating a GM mask extracted from the FreeSurfer's *aseg* segmentation, and it ensures components are not extracted from voxels containing a minimal fraction of GM. Finally, these masks are resampled into BOLD space and binarized by thresholding at 0.99 (as in the original implementation). Components are also calculated separately within the WM and CSF masks. For each CompCor decomposition, the *k* components with the largest singular values are retained, such that the retained components' time series are sufficient to explain 50 percent of variance across the nuisance mask (CSF, WM, combined, or temporal). The remaining components are dropped from

consideration. The head-motion estimates calculated in the correction step were also placed

within the corresponding confounds file. The confound time series derived from head motion

estimates and global signals were expanded with the inclusion of temporal derivatives and

quadratic terms for each (Satterthwaite et al. 2013). Frames that exceeded a threshold of 0.9

mm FD or 1.5 standardised DVARS were annotated as motion outliers. The BOLD time-

series were resampled onto the following surfaces (FreeSurfer reconstruction

nomenclature): *fsnative*. All resamplings can be performed with *a single interpolation step* by

composing all the pertinent transformations (i.e., head-motion transform matrices,

susceptibility distortion correction when available, and co-registrations to anatomical and

output spaces). Surface resampling was performed using mri_vol2surf (FreeSurfer).

Additional pre-processing

The first 4 volumes of each run were discarded for signal stabilization. Bold data in fsnative

space was smoothed along the cortical surface (Gaussian kernel FWHM = 3 mm). For each

run, a denoising procedure was further applied on the data by inputting nuisance regressors

into a PCA and regressing out the top 5 components. The nuisance regressors include: Six

head motion regressors, FD and DVARS, CSF mean signal, white matter mean signal, the

first five anatomical CompCor regressors (csf+wm combined), and the first five temporal

CompCor regressors. Then data and regressors were temporally filtered with a Savitzky-

Golay filter (3rd order, 347 seconds in length) and zscored. The time series from the two

identical task runs were then averaged to increase signal to noise ratio.

MRI analysis

Region of interests

618 An anatomically defined retinotopy atlas was applied to the inflated cortical surface for each

619 participant in order to delineate the borders of V1, V2 and V3, and also each degree of

620 eccentricity up to 8° in V1 (Benson & Winawer, 2018). See Figure 1C for an example of an

621 eccentricity plot for one participant.

Structural data analysis

623 The 'mris_anatomical_stats' command was used to extract surface area and cortical

624 thickness properties for each participant, per hemisphere, for each region of interest. Overall

625 cortical thickness per participant was computed by weighting the cortical thickness of a

626 hemisphere by the number of vertices per hemisphere.

pRF analysis

pRF modelling was carried out with a custom Python package, prfpy (Aqil & Knapen, 2023).

To model population receptive fields, we used a isometric bivariate Gaussian model

(Dumoulin & Wandell, 2008), with *x,y* representing the preferred retinotopic position, spatial

standard deviation sigma (σ) representing pRF size as well as the baseline of the timeseries

and the signal amplitude beta (β). At every cortical location, the parameters of each model

were optimized by minimizing the residual sum of squares between the model prediction and

the measured BOLD signal. Fitting began with a grid search for the best model parameters

(pRF position and size) followed by an iterative optimization. In each participant, in each

condition, vertices with an $R^2 < 0.1$ (variance explained by the pRF model), a negative beta,

or a centre outside the stimulated radius (8.6°) were excluded from further analysis.

In addition, we ran a second model fit using a stimulus aperture file that incorporated a black

oval mask (1.25°x0.9°) over the area of the central rod-free scotoma (Curcio et al., 1990).

Previous research suggests that biases in pRF measures such as shifts in eccentricities and

size can occur when a scotoma is present and not explicitly modelled in the stimulus

apertures (Binda et al., 2013). Results from this analysis are reported in Supplement 3.

Connective field modelling

Connective field modelling was carried out with a dedicated Python package, prfpy (Aqil &

Knapen, 2023) based on the method described in Knapen (2021). In brief, gaussian

connective field profiles on the surface are defined for each vertex on the cortical manifold,

with a centre location of the connective field (v-centre), and σ (sigma) the Gaussian spread

in millimeters on the cortical surface.

To generate spatial Gaussian fields along the V1 surface, the diffusion of heat along the

cortical mesh was simulated (Crane et al., 2013) and used to infer geodesic distances

between all V1 vertices (V1 definition was based on the Benson atlas, see Regions of

Interest). As the two hemispheres are two separate surface meshes, this distance matrix

was calculated for each hemisphere separately. Grid-fit candidate CF sizes were: [0.5, 1, 2,

3, 4, 5, 7, 10, 15, 20] mm σ for the Gaussian CF. The predicted time course for each of the

resulting CF models was generated by taking the dot product between the CF's vertex profile

and the vertex by time-courses matrix in V1. The resulting CF model time courses were z

scored and correlated with the time courses throughout the brain. For each location in the

brain, the connective field resulting in the highest squared correlation was selected. Only

values within an $R^2 > 0.1$, beta > 0, and a centre within 8° of eccentricity (matching the

radius of the stimuli) were included in further analyses. To create connective-field-based

eccentricity maps, each vertex was assigned the Benson atlas-based eccentricities (Benson
& Winawer, 2018) of its CF V1 centre.

For visualisation of group maps all individual maps were transformed to fsaverage space.
Plotted values indicate the medians of each group and condition per vertex. A vertex was left
empty if no significant data was available for more than 30% of individuals within the group.

Statistical analysis

667 Jamovi R package (jmv) was used for Anovas and t-tests (The jamovi project, 2022). Jasp
668 (jsq) R package for Bayesian t-tests (JASP team, 2020). Where sphericity assumptions were
669 not met a greenhouse-geisser correction was applied. Uncorrected p-values are reported as
670 well as FDR multiple comparison corrections.

Acknowledgments

672 We thank Maria Molina Sanchez, Hunter Schone and Matan Mazor for providing comments
673 on an earlier version of this manuscript.

Funding

The research was supported by grants from the National Institute for Health Research
(NIHR) Biomedical Research Centre (BRC) at Moorfields Eye Hospital NHS Foundation
Trust and UCL Institute of Ophthalmology, the Economic and Social Research Council
(ESRC) of the UKRI (#ES/N000838/1); MeiraGTx, Retina UK, Moorfields Eye Charity
(R160035A, R190029A, R180004A), Foundation Fighting Blindness (USA) and The
Wellcome Trust (099173/Z/12/Z, 100227), Ardalan Family Scholarship, the Persia
Educational Foundation Maryam Mirzakhani Scholarship; and the Sir Richard Stapley
Educational Trust (#313812). The views expressed are those of the authors and not
necessarily those of the NHS, the NIHR, or the Department of Health.

Competing interests

685 Scanner time was funded by MeiraGTx via a research sponsorship awarded to UCL. M.M.
686 declares financial interest in MeiraGTx.

References

Aboshiha, J., Dubis, A. M., Cowing, J., Fahy, R. T. A., Sundaram, V., Bainbridge, J. W., Ali, R. R., Dubra, A., Nardini, M., Webster, A. R., Moore, A. T., Rubin, G., Carroll, J., & Michaelides, M. (2014). A prospective longitudinal study of retinal structure and function in achromatopsia. *Investigative Ophthalmology and Visual Science*, *55*(9), 5733–5743. https://doi.org/10.1167/iovs.14-14937

Abraham, A., Pedregosa, F., Eickenberg, M., Gervais, P., Mueller, A., Kossaifi, J., Gramfort, A., Thirion, B., & Varoquaux, G. (2014). Machine learning for neuroimaging with scikit-learn. *Frontiers in Neuroinformatics*, *8*. https://doi.org/10.3389/fninf.2014.00014

Adams, D. L., Sincich, L. C., & Horton, J. C. (2007). Complete Pattern of Ocular Dominance Columns in Human Primary Visual Cortex. *Journal of Neuroscience*, *27*(39), 10391–10403. https://doi.org/10.1523/JNEUROSCI.2923-07.2007

Aqil, M., & Knapen, T. (2023). *Prfpy: A python package to simulate and fit population receptive field models to time series data.* (v0.1.0-alpha) [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.10201022

Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis*, *12*(1), 26–41. https://doi.org/10.1016/j.media.2007.06.004

Baker, C. I., Dilks, D. D., Peli, E., & Kanwisher, N. (2008). Reorganization of visual processing in macular degeneration: Replication and clues about the role of foveal loss. *Vision Research*, *48*(18), 1910–1919. https://doi.org/10.1016/j.visres.2008.05.020

Baker, C. I., Peli, E., Knouf, N., & Kanwisher, N. G. (2005). Reorganization of Visual Processing in Macular Degeneration. *The Journal of Neuroscience*, *25*(3), 614–618. https://doi.org/10.1523/JNEUROSCI.3476-04.2005

Barton, B., & Brewer, A. A. (2015). fMRI of the rod scotoma elucidates cortical rod pathways

and implications for lesion measurements. *Proceedings of the National Academy of*

Sciences of the United States of America, *112*(16), 5201–5206.

https://doi.org/10.1073/PNAS.1423673112/-/DCSUPPLEMENTAL

Baseler, H. A., Brewer, A. A., Sharpe, L. T., Morland, A. B., Jaägle, H., & Wandell, B. A.

(2002). Reorganization of human cortical maps caused by inherited photoreceptor

abnormalities. *Nature Neuroscience 2002 5:4*, *5*(4), 364–370.

https://doi.org/10.1038/nn817

Baseler, H. A., Gouws, A., Haak, K. V., Racey, C., Crossland, M. D., Tufail, A., Rubin, G. S.,

Cornelissen, F. W., & Morland, A. B. (2011). Large-scale remapping of visual cortex

is absent in adult humans with macular degeneration. *Nature Neuroscience*, *14*(5),

649–657. https://doi.org/10.1038/nn.2793

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction

method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90–

101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Benson, N. C., & Winawer, J. (2018). Bayesian analysis of retinotopic maps. *eLife*, *7*.

https://doi.org/10.7554/ELIFE.40224

Binda, P., Thomas, J. M., Boynton, G. M., & Fine, I. (2013). Minimizing biases in estimating

the reorganization of human visual areas with BOLD retinotopic mapping. *Journal of*

Vision, *13*(7), 13. https://doi.org/10.1167/13.7.13

Brown, H. D. H., Gouws, A. D., Vernon, R. J. W., Lawrence, S. J. D., Donnelly, G., Gill, L.,

Gale, R. P., Baseler, H. A., & Morland, A. B. (2021). Assessing functional

reorganization in visual cortex with simulated retinal lesions. *Brain Structure and*

Function, *226*(9), 2855–2867. https://doi.org/10.1007/s00429-021-02366-w

Burton, E., Wattam-Bell*, J., S. Rubin, G., Aboshiha, J., Michaelides, M., Atkinson, J.,

Braddick, O., & Nardini, M. (2016). Dissociations in Coherence Sensitivity Reveal

Atypical Development of Cortical Visual Processing in Congenital Achromatopsia.

Investigative Ophthalmology & Visual Science, *57*(4), 2251–2259.

https://doi.org/10.1167/iovs.15-18414

Calford, M. B., Chino, Y. M., Das, A., Eysel, U. T., Gilbert, C. D., Heinen, S. J., Kaas, J. H., &

Ullman, S. (2005). Rewiring the adult brain. *Nature*, *438*(7065), E3–E3.

https://doi.org/10.1038/nature04359

Carvalho, J., Renken, R. J., & Cornelissen, F. W. (2019). Studying Cortical Plasticity in

Ophthalmic and Neurological Disorders: From Stimulus-Driven to Cortical Circuitry

Modeling Approaches. *Neural Plasticity*, *2019*, e2724101.

https://doi.org/10.1155/2019/2724101

Carvalho, J., Renken, R. J., & Cornelissen, F. W. (2021). Predictive masking of an artificial

scotoma is associated with a system-wide reconfiguration of neural populations in the

human visual cortex. *NeuroImage*, *245*, 118690.

https://doi.org/10.1016/j.neuroimage.2021.118690

Crane, K., Weischedel, C., & Wardetzky, M. (2013). Geodesics in heat: A new approach to

computing distance based on heat flow. *ACM Transactions on Graphics*, *32*(5),

152:1-152:11. https://doi.org/10.1145/2516971.2516977

Curcio, C. A., Sloan, K. R., Kalina, R. E., & Hendrickson, A. E. (1990). Human photoreceptor

topography. *Journal of Comparative Neurology*, *292*(4), 497–523.

https://doi.org/10.1002/cne.902920402

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical Surface-Based Analysis: I.

Segmentation and Surface Reconstruction. *NeuroImage*, *9*(2), 179–194.

https://doi.org/10.1006/nimg.1998.0395

Dilks, D. D., Baker, C. I., Peli, E., & Kanwisher, N. (2009). Reorganization of Visual

Processing in Macular Degeneration Is Not Specific to the "Preferred Retinal Locus".

The Journal of Neuroscience, *29*(9), 2768.

https://doi.org/10.1523/JNEUROSCI.5258-08.2009

Dumoulin, S. O., & Knapen, T. (2018). How Visual Cortical Organization Is Altered by

Ophthalmologic and Neurologic Disorders. *Annual Review of Vision Science*, *4*(1),

357–379. https://doi.org/10.1146/annurev-vision-091517-033948

Dumoulin, S. O., & Wandell, B. A. (2008). Population receptive field estimates in human

visual cortex. *NeuroImage*, *39*(2), 647–660.

https://doi.org/10.1016/j.neuroimage.2007.09.034

Esteban, O., Blair, R., Markiewicz, C. J., Berleant, S. L., Moodie, C., Ma, F., Isik, A. I.,

Erramuzpe, A., Kent, M., James D. andGoncalves, DuPre, E., Sitek, K. R., Gomez,

D. E. P., Lurie, D. J., Ye, Z., Poldrack, R. A., & Gorgolewski, K. J. (2018). fMRIPrep.

Software. https://doi.org/10.5281/zenodo.852659

Esteban, O., Markiewicz, C., Blair, R. W., Moodie, C., Isik, A. I., Erramuzpe Aliaga, A., Kent,

J., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S., Wright, J., Durnez, J.,

Poldrack, R., & Gorgolewski, K. J. (2018). fMRIPrep: A robust preprocessing pipeline

for functional MRI. *Nature Methods*. https://doi.org/10.1038/s41592-018-0235-4

Estévez, O., & Spekreijse, H. (1982). The 'silent substitution' method in visual research.

Vision Research, *22*(6), 681–691. https://doi.org/10.1016/0042-6989(82)90104-3

Farahbakhsh, M., Anderson, E. J., Maimon-Mor, R. O., Rider, A., Greenwood, J. A., Hirji, N.,

Zaman, S., Jones, P. R., Schwarzkopf, D. S., Rees, G., Michaelides, M., & Dekker,

T. M. (2022). A demonstration of cone function plasticity after gene therapy in

achromatopsia. *Brain*, *145*(11), 3803–3815. https://doi.org/10.1093/brain/awac226

Ferreira, S., Pereira, A. C., Quendera, B., Reis, A., Silva, E. D., & Castelo-Branco, M.

(2016). Primary visual cortical remapping in patients with inherited peripheral retinal

degeneration. *NeuroImage : Clinical*, *13*, 428–438.

https://doi.org/10.1016/j.nicl.2016.12.013

Fischer, M. D., Michalakis, S., Wilhelm, B., Zobor, D., Muehlfriedel, R., Kohl, S., Weisschuh,

N., Ochakovski, G. A., Klein, R., Schoen, C., Sothilingam, V., Garcia-Garrido, M.,

Kuehlewein, L., Kahle, N., Werner, A., Dauletbekov, D., Paquet-Durand, F., Tsang,

S., Martus, P., … Wissinger, B. (2020). Safety and Vision Outcomes of Subretinal

Gene Therapy Targeting Cone Photoreceptors in Achromatopsia: A Nonrandomized

Controlled Trial. *JAMA Ophthalmology*, *138*(6), 643–651.

https://doi.org/10.1001/jamaophthalmol.2020.1032

Fonov, V., Evans, A., McKinstry, R., Almli, C., & Collins, D. (2009). Unbiased nonlinear

average age-appropriate brain templates from birth to adulthood. *NeuroImage*, *47,*

Supplement 1, S102. https://doi.org/10.1016/S1053-8119(09)70884-5

Georgiou, M., Fujinami, K., & Michaelides, M. (2021). Inherited retinal diseases:

Therapeutics, clinical trials and end points—A review. *Clinical & Experimental*

Ophthalmology, *49*(3), 270–288. https://doi.org/10.1111/ceo.13917

Goesaert, E., Van Baelen, M., Spileers, W., Wagemans, J., & Op de Beeck, H. P. (2014).

Visual Space and Object Space in the Cerebral Cortex of Retinal Disease Patients.

PLoS ONE, *9*(2), e88248. https://doi.org/10.1371/journal.pone.0088248

Gorgolewski, K., Burns, C. D., Madison, C., Clark, D., Halchenko, Y. O., Waskom, M. L., &

Ghosh, S. (2011). Nipype: A flexible, lightweight and extensible neuroimaging data

processing framework in Python. *Frontiers in Neuroinformatics*, *5*, 13.

https://doi.org/10.3389/fninf.2011.00013

Gorgolewski, K. J., Esteban, O., Markiewicz, C. J., Ziegler, E., Ellis, D. G., Notter, M. P.,

Jarecka, D., Johnson, H., Burns, C., Manhães-Savio, A., Hamalainen, C., Yvernault,

B., Salo, T., Jordan, K., Goncalves, M., Waskom, M., Clark, D., Wong, J., Loney, F.,

… Ghosh, S. (2018). Nipype. *Software*. https://doi.org/10.5281/zenodo.596855

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using

boundary-based registration. *NeuroImage*, *48*(1), 63–72.

https://doi.org/10.1016/j.neuroimage.2009.06.060

Haak, K. V., Cornelissen, F. W., & Morland, A. B. (2012). Population Receptive Field

Dynamics in Human Visual Cortex. *PLoS ONE*, *7*(5), e37686.

https://doi.org/10.1371/journal.pone.0037686

Haak, K. V., Morland, A. B., & Engel, S. A. (2015). Plasticity, and its limits, in adult human

primary visual cortex. *Multisensory Research*, *28*(3–4), 297–307.

https://doi.org/10.1163/22134808-00002496

Haak, K. V., Winawer, J., Harvey, B. M., Renken, R., Dumoulin, S. O., Wandell, B. A., &

Cornelissen, F. W. (2013). Connective field modeling. *NeuroImage*, *66*, 376–384.

https://doi.org/10.1016/j.neuroimage.2012.10.037

Hess, R. F., & Nordby, K. (1986). Spatial and temporal limits of vision in the achromat. *The*

Journal of Physiology, 371(1), 365–385.

https://doi.org/10.1113/jphysiol.1986.sp015981

Hummer, A., Ritter, M., Woletz, M., Ledolter, A. A., Tik, M., Dumoulin, S. O., Holder, G. E.,

Schmidt-Erfurth, U., & Windischberger, C. (2018). Artificial scotoma estimation based

on population receptive field mapping. *NeuroImage*, *169*, 342–351.

https://doi.org/10.1016/J.NEUROIMAGE.2017.12.010

JASP team. (2020). *JASP (Version 0.18.3)* [Computer software]. https://jasp-stats.org/

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the

Robust and Accurate Linear Registration and Motion Correction of Brain Images.

NeuroImage, *17*(2), 825–841. https://doi.org/10.1006/nimg.2002.1132

Jiang, J., Zhu, W., Shi, F., Liu, Y., Li, J., Qin, W., Li, K., Yu, C., & Jiang, T. (2009). Thick

Visual Cortex in the Early Blind. *Journal of Neuroscience*, *29*(7), 2205–2211.

https://doi.org/10.1523/JNEUROSCI.5451-08.2009

Jones, P. R., Somoskeöy, T., Chow-Wing-Bom, H., & Crabb, D. P. (2020). Seeing other

perspectives: Evaluating the use of virtual and augmented reality to simulate visual

impairments (OpenVisSim). *Npj Digital Medicine*, *3*(1), 1–9.

https://doi.org/10.1038/s41746-020-0242-6

Klein, A., Ghosh, S. S., Bao, F. S., Giard, J., Häme, Y., Stavsky, E., Lee, N., Rossa, B.,

Reuter, M., Neto, E. C., & Keshavan, A. (2017). Mindboggling morphometry of

human brains. *PLOS Computational Biology*, *13*(2), e1005350.

https://doi.org/10.1371/journal.pcbi.1005350

Klein, B. P., Fracasso, A., van Dijk, J. A., Paffen, C. L. E., te Pas, S. F., & Dumoulin, S. O.

(2018). Cortical depth dependent population receptive field attraction by spatial

attention in human V1. *NeuroImage*, *176*, 301–312.

https://doi.org/10.1016/j.neuroimage.2018.04.055

Kleiner, M., Brainard, D. H., Pelli, D. G., Broussard, C., Wolf, T., & Niehorster, D. (2007).

What's new in Psychtoolbox-3? A free cross-platform toolkit for psychophysiscs with

Matlab and GNU/Octave. In *Cognitive and Computational Psychophysics* (Vol. 36).

Knapen, T. (2021). Topographic connectivity reveals task-dependent retinotopic processing

throughout the human brain. *Proceedings of the National Academy of Sciences*,

118(2), e2017032118. https://doi.org/10.1073/pnas.2017032118

Lehky, S. R., & Tanaka, K. (2016). Neural representation for object recognition in

inferotemporal cortex. *Current Opinion in Neurobiology*, *37*, 23–35.

https://doi.org/10.1016/j.conb.2015.12.001

Lu, Z.-L., & Dosher, B. A. (2022). Current directions in visual perceptual learning. *Nature*

Reviews Psychology, *1*(11), 654–668. https://doi.org/10.1038/s44159-022-00107-2

Macnamara, A., Chen, C., Schinazi, V. R., Saredakis, D., & Loetscher, T. (2021). Simulating

Macular Degeneration to Investigate Activities of Daily Living: A Systematic Review.

Frontiers in Neuroscience, *15*. https://doi.org/10.3389/fnins.2021.663062

Makin, T. R., & Krakauer, J. W. (2023). Against cortical reorganisation. *eLife*, *12*, e84716.

https://doi.org/10.7554/eLife.84716

Mckyton, A., Averbukh, E., Marks Ohana, D., Levin, N., & Banin, E. (2021). *Cortical Visual*

Mapping following Ocular Gene Augmentation Therapy for Achromatopsia.

https://doi.org/10.1523/JNEUROSCI.3222-20.2021

Michaelides, M., Hirji, N., Wong, S. C., Besirli, C. G., Zaman, S., Kumaran, N., Georgiadis,

A., Smith, A. J., Ripamonti, C., Gottlob, I., Robson, A. G., Thiadens, A., Henderson,

R. H., Fleck, P., Anglade, E., Dong, X., Capuano, G., Lu, W., Berry, P., …

Bainbridge, J. (2023). First-in-Human Gene Therapy Trial of AAV8-hCARp.hCNGB3

in Adults and Children With CNGB3-associated Achromatopsia. *American Journal of*

Ophthalmology, *253*, 243–251. https://doi.org/10.1016/j.ajo.2023.05.009

Molz, B., Herbik, A., Baseler, H. A., de Best, P. B., Vernon, R. W., Raz, N., Gouws, A. D.,

Ahmadi, K., Lowndes, R., McLean, R. J., Gottlob, I., Kohl, S., Choritz, L., Maguire, J.,

Kanowski, M., Käsmann-Kellner, B., Wieland, I., Banin, E., Levin, N., … Morland, A.

B. (2022). Structural changes to primary visual cortex in the congenital absence of

cone input in achromatopsia. *NeuroImage: Clinical*, *33*, 102925.

https://doi.org/10.1016/J.NICL.2021.102925

Molz, B., Herbik, A., Baseler, H. A., de Best, P., Raz, N., Gouws, A., Ahmadi, K., Lowndes,

R., McLean, R. J., Gottlob, I., Kohl, S., Choritz, L., Maguire, J., Kanowski, M.,

Käsmann-Kellner, B., Wieland, I., Banin, E., Levin, N., Morland, A. B., & Hoffmann,

M. B. (2023). Achromatopsia—Visual Cortex Stability and Plasticity in the Absence of

Functional Cones. *Investigative Ophthalmology & Visual Science*, *64*(13), 23.

https://doi.org/10.1167/iovs.64.13.23

Park, H.-J., Lee, J. D., Kim, E. Y., Park, B., Oh, M.-K., Lee, S., & Kim, J.-J. (2009).

Morphological alterations in the congenital blind based on the analysis of cortical

thickness and surface area. *NeuroImage*, *47*(1), 98–106.

https://doi.org/10.1016/j.neuroimage.2009.03.076

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.

(2014). Methods to detect, characterize, and remove motion artifact in resting state

fMRI. *NeuroImage*, *84*(Supplement C), 320–341.

https://doi.org/10.1016/j.neuroimage.2013.08.048

Reichel, F. F., Michalakis, S., Wilhelm, B., Zobor, D., Muehlfriedel, R., Kohl, S., Weisschuh,

N., Sothilingam, V., Kuehlewein, L., Kahle, N., Seitz, I., Paquet-Durand, F., Tsang, S.

H., Martus, P., Peters, T., Seeliger, M., Bartz-Schmidt, K. U., Ueffing, M., Zrenner, E.,

… Fischer, D. (2022). Three-year results of phase I retinal gene therapy trial for

CNGA3-mutated achromatopsia: Results of a non randomised controlled trial. *British*

Journal of Ophthalmology, *106*(11), 1567–1572. https://doi.org/10.1136/bjophthalmol-

2021-319067

Ritter, M., Hummer, A., Ledolter, A. A., Holder, G. E., Windischberger, C., & Schmidt-Erfurth,

U. M. (2019). Correspondence between retinotopic cortical mapping and

conventional functional and morphological assessment of retinal disease. *British*

Journal of Ophthalmology, *103*(2), 208–215. https://doi.org/10.1136/bjophthalmol-

2017-311443

Schumacher, E. H., Jacko, J. A., Primo, S. A., Main, K. L., Moloney, K. P., Kinzel, E. N., &

Ginn, J. (2008). Reorganization of visual processing is related to eccentric viewing in

patients with macular degeneration. *Restorative Neurology and Neuroscience*, *26*(4–

5), 391–402.

Tailor, V. K., Theodorou, M., Dahlmann-Noor, A. H., Dekker, T. M., & Greenwood, J. A.

(2021). Eye movements elevate crowding in idiopathic infantile nystagmus syndrome.

Journal of Vision, *21*(13), 9–9. https://doi.org/10.1167/JOV.21.13.9

Tangtartharakul, G., Morgan, C. A., Rushton, S. K., & Schwarzkopf, D. S. (2023).

Retinotopic connectivity maps of human visual cortex with unconstrained eye

movements. *Human Brain Mapping*, *44*(16), 5221–5237.

https://doi.org/10.1002/hbm.26446

The jamovi project. (2022). *Jamovi* (2.3) [Computer software]. https://www.jamovi.org

Tustison, N. J., Avants, B. B., Cook, P. A., Zheng, Y., Egan, A., Yushkevich, P. A., & Gee, J.

C. (2010). N4ITK: Improved N3 Bias Correction. *IEEE Transactions on Medical*

Imaging, *29*(6), 1310–1320. https://doi.org/10.1109/TMI.2010.2046908

van Dijk, J. A., de Haas, B., Moutsiana, C., & Schwarzkopf, D. S. (2016). Intersession

reliability of population receptive field estimates. *NeuroImage*, *143*, 293–303.

https://doi.org/10.1016/j.neuroimage.2016.09.013

Wandell, B. A., & Smirnakis, S. M. (2010). Plasticity and stability of visual field maps in adult

primary visual cortex. *Nature Reviews Neuroscience*, *10*, 873–884.

https://doi.org/10.1038/nrn2741

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a

hidden Markov random field model and the expectation-maximization algorithm. *IEEE*

Transactions on Medical Imaging, *20*(1), 45–57. https://doi.org/10.1109/42.906424