The functional roles of zebrafish *HoxA*- and *HoxD*-related clusters in the pectoral fin development

Mizuki Ishizaka[a], **Akiteru Maeno**[b], **Hidemichi Nakazawa**[a], **Renka Fujii**[a], **Sae Oikawa**[a], **Taisei Tani**[a], **Haruna Kanno**[a], **Rina Koita**[a], **and Akinori Kawamura**[a,*]

[a] Division of Life Science, Graduate School of Science and Engineering, Saitama University, Shimo-Okubo 255, Sakura-ku, Saitama 338-8570, Japan

[b] Cell Architecture Laboratory, National Institute of Genetics, Yata 1111, Mishima, Shizuoka 411-8540, Japan.

[*] Author to whom correspondence should be addressed:

Akinori Kawamura, Ph.D.

Division of Life Science, Graduate School of Science and Engineering, Saitama University, Saitama 338-8570, Japan

Phone: +81-48-858-3808

E-mail: akawamur@mail.saitama-u.ac.jp

Keywords: pectoral fin; *Hox* genes; zebrafish; X-ray CT scan

Abstract

The paralogs 9-13 *Hox* genes in mouse *HoxA* and *HoxD* clusters are critical for

limb development. When both *HoxA* and *HoxD* clusters are deleted in mice, significant

limb truncation is observed compared to the phenotypes of single and compound

mutants of *Hox9-13* genes in these clusters. In zebrafish, mutations in *hox13* genes in

HoxA- and *HoxD*-related clusters result in abnormal morphology of pectoral fins,

homologous to forelimbs. However, the effect of the simultaneous deletions of entire

HoxA- and *HoxD*-related clusters on pectoral fin development remains unknown. Here,

we generated mutants with several combinations of *hoxaa*, *hoxab*, and *hoxda* cluster

deletions and analyzed the pectoral fin development. In $hoxaa^{-/-};hoxab^{-/-};hoxda^{-/-}$ larvae,

the endoskeletal disc and the fin-fold are significantly shortened in developing pectoral

fins. In addition, we show that this anomaly is due to defects in the pectoral fin growth

after the fin bud formation. Furthermore, in the surviving adult mutants, micro-CT

scanning reveals defects in the posterior portion of the pectoral fin which is thought to

represent latent regions of the limb. Our results further support that the functional role

of *HoxA* and *HoxD* clusters is conserved in the paired appendage formation in bony

fishes.

Introduction

In the development of bilaterian animals, *Hox* genes, which encode homeodomain transcription factors, play a crucial role in determining the positional identities along the body axis. *Hox* genes are typically organized in clusters known as *Hox* clusters. The order of the *Hox* genes within a cluster corresponds to their collinear expression patterns in developing embryos along the body axis[1-3]. During the early stages of vertebrate evolution, a single primitive *Hox* cluster, consisting of 1–13 paralogous groups, underwent two rounds of whole-genome duplication, resulting in quadruplication[4,5]. As a result, mice now possess four *Hox* clusters known as *HoxA*, *HoxB*, *HoxC*, and *HoxD*, each located on different chromosomes.

Loss-of-function studies using knockout mice have revealed that *Hox* genes play important roles in vertebrate development[6,7]. Specifically, the 9-13 paralogs of *Hox* genes in *HoxA* and *HoxD* clusters, which all correspond to homologous to the *Drosophila Abd-B* homeotic gene, have been shown to cooperatively contribute to limb development[8-15]. In mouse and chick, it has been observed that *Hoxa9* to *Hoxa13* and *Hoxd9* to *Hoxd13* genes exhibit nested and collinear expression domains in the mesenchymal cells of developing limbs[1,6,16]. This occurs through the successive activation of these *Hox* genes according to their positional order in *HoxA* and *HoxD* clusters. Furthermore, knockout mice for *Hoxa* and *Hoxd* genes have been generated, providing insights into the roles of these *Hoxa* and *Hoxd* genes in providing the regional identities of each segment of the limbs[8-15]. For instance, mice lacking *Hoxa9* and *Hoxd9* exhibit abnormalities in the stylopod[12], which is the bone on the proximal side of the limb. Similarly, mice lacking *Hoxa13* and *Hoxd13* show defects in the autopod[13], which

is the bone on the distal side of the limb. Moreover, the simultaneous deletion of *HoxA*

and *HoxD* clusters leads to severe truncation of forelimbs, particularly distal elements,

in mice[14].

Zebrafish possess seven *hox* clusters that were further duplicated through

teleost-specific whole-genome duplication[17]. Currently, zebrafish have two clusters

derived from *HoxA*, namely *hoxaa* and *hoxab*. On the other hand, only *hoxda* cluster,

derived from *HoxD*, is present, as *hoxdb* cluster has been lost, except for a single

microRNA[17]. In the formation of the pectoral fin, which is homologous to the forelimb,

the 9-13 paralogous *hox* genes in zebrafish *hoxaa*, *hoxab*, and *hoxda* clusters exhibit

similar collinear expression patterns to those of mouse and chick[18-20]. Loss-of-function

study of *hoxa13a*, *hoxa13b*, and *hoxd13a* demonstrated the severe truncation of the

pectoral fin in adult zebrafish[19]. These findings suggest a deep conserved functional

homology of the posterior *HoxA*- and *HoxD*-related genes in the formation of the paired

appendages in bony fishes. However, other posterior *hox* genes besides these *hox13*

genes are also expressed during pectoral fin formation in zebrafish[18,20], and the function

of the entire *HoxA*- and *HoxD*-related clusters in pectoral fin development is yet to be

elucidated.

In a previous study, we used the CRISPR-Cas9 system to generate mutants

lacking each of the seven *hox* clusters in zebrafish[21]. Among the three *HoxA*- and

HoxD-related clusters, only *hoxab* cluster deletion mutants showed shortening of the

pectoral fin during embryogenesis[21], suggesting functional redundancy among these *hox*

clusters. In this study, we generated mutants with various combinations of deletions in

hoxaa, *hoxab*, and *hoxda* clusters and examined the role of these clusters in pectoral fin

development and skeletal structures of pectoral fins in adult fish.

Results

Previous studies have shown that mouse *Hox* genes in *HoxA* and *HoxD* clusters cooperate to play essential roles in limb patterning[8-15]. The absence of all *HoxA* and *HoxD* functions results in a significant truncation of distal limb elements[14]. To investigate the role of zebrafish *hoxaa*, *hoxab*, and *hoxda* clusters, which are homologous to mouse *HoxA* and *HoxD* clusters, in zebrafish development, we focused on the pectoral fin, a homologous to tetrapod forelimb in this study. To analyze the pectoral fin phenotypes, we examined larvae derived from intercrosses between triple hemizygous mutants for *hoxaa*, *hoxab*, and *hoxda* clusters (Fig. 1A-H). At 3 dpf, we observed that triple homozygous deletion mutants (hereafter referred to $hoxaa^{-/-};hoxab^{-/-};hoxda^{-/-}$) had present but severely shortened pectoral fins (Fig. 1H). The pectoral fin shortening in $hoxaa^{-/-};hoxab^{-/-};hoxda^{-/-}$ mutants was significantly greater than in any of the other *hox* cluster-deleted mutants. These results indicate that zebrafish *hoxaa*, *hoxab*, and *hoxda* clusters redundantly function in pectoral fin formation. The similar shortened pectoral fin/forelimbs phenotype in zebrafish and mouse *HoxA*- and *HoxD*-related mutants further suggest that the roles of *HoxA*- and *HoxD*-related clusters in paired appendage formation are conserved between zebrafish and mice.

The pectoral fin of zebrafish larvae consists mainly of the cartilaginous endoskeletal disc and the non-cartilaginous fin-fold (Fig. 2A)[22]. To examine the truncation of the pectoral fins in *hoxaa*, *hoxab*, and *hoxda* cluster mutant larvae in detail, we next compared the lengths of the endoskeletal disc and fin-fold, respectively (Fig. 2A-H). Cartilage-stained pectoral fins at 5 dpf showed that the lengths of the

endoskeletal disc along the anterior-posterior and proximal-distal axes were

significantly shorter in *hoxab$^{-/-}$;hoxda$^{-/-}$* and *hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$* larvae, whereas

the other combinations of *hoxaa*, *hoxab*, and *hoxda* cluster deletions did not appear to

affect the lengths of the endoskeletal disc (Fig. 2I, J). On the other hand, the effects of

hoxaa, *hoxab*, and *hoxda* cluster deletions were more pronounced with respect to the

length of fin-fold (Fig. 2K). The fin-fold in *hoxaa$^{-/-}$;hoxab$^{-/-}$* larvae was shortened

compared to that of wild-type larvae, whereas the differences in the length of the

endoskeletal disc were not observed. Among the three double *hox* cluster deletion

mutants, *hoxab$^{-/-}$;hoxda$^{-/-}$* larvae were the shortest in both the fin-fold and the

endoskeletal disc. Furthermore, the fin-fold length was shortest in

hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$ larvae, as was the endoskeletal disc. These results suggest that

the cooperative functions of *hoxaa*, *hoxab*, and *hoxda* clusters: *hoxab* cluster has the

highest contribution to pectoral fin formation in zebrafish, followed by *hoxda* cluster

and then *hoxaa* cluster.

To gain further insight into the shortening of pectoral fins in

hoxaa;hoxab;hoxda cluster mutants, we performed whole-mount *in situ* hybridization to

examine the expression patterns of genes essential for the pectoral fin formation.

Zebrafish *tbx5a* is essential for the initial formation of the pectoral fin buds, and loss of

tbx5a function leads to the absence of the pectoral fins[23,24]. Thus, we examined the

expression patterns of *tbx5a* in 30 hpf embryos derived from the intercrosses between

triple hemizygous mutants for *hoxaa*, *hoxab*, and *hoxda* clusters. We found no

significant differences in the expression of *tbx5a* in embryos obtained from this

intercross. Furthermore, we confirmed by genotyping of the stained embryos that

expression patterns of *tbx5a* in *hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$* larvae appeared

indistinguishable from that of sibling wild-type embryos (Fig. 3A, B). These results

indicate the normal establishment of the pectoral fin buds even in the absence of all the

functions of *hoxaa*, *hoxab*, and *hoxda* clusters. Therefore, we analyzed the expression of

sonic hedgehog a (*shha*) (Fig. 3C-J), which is subsequently required for the cell

proliferation of the developing pectoral fin[25]. In zebrafish, *shha* is expressed in the

posterior portion of the fin buds in 48 hpf embryos (Fig. 3C). Consistent with shortened

pectoral fins in zebrafish *HoxA-* and *HoxD*-related cluster mutants (Fig. 1, 2), reduced

expression of *shha* in the pectoral fin buds was observed. This was particularly evident

in *hoxab*$^{-/-}$;*hoxda*$^{-/-}$ and *hoxaa*$^{-/-}$;*hoxab*$^{-/-}$;*hoxda*$^{-/-}$ larvae, where *shha* expression in fin

buds was markedly down-regulated (Fig. 3I, J, Fig. S2). These results suggest that the

shortening of the pectoral fin observed in zebrafish *HoxA-* and *HoxD*-related cluster

mutants is due to retardation of pectoral fin growth after the normal formation of the fin

buds by *tbx5a* expression.

    Next, we investigated whether *HoxA-* and *HoxD*-related clusters contribute to

the skeletal formation of adult zebrafish pectoral fins. For this purpose, we raised larvae

to adult fish by multiple intercrosses between triple hemizygous mutants for *hoxaa*,

hoxab, and *hoxda* clusters. Our previous study demonstrated that homozygous deletion

of zebrafish *hoxab* cluster leads to lethality[21]. Consistently, fish with homozygous

deletion in *hoxab* cluster did not survive with any combination of deletions in *hoxaa*

and *hoxda* clusters (Table 1). Furthermore, fish with homozygous deletion of two or

three clusters in *hoxaa*, *hoxab*, and *hoxda* also did not survive. Therefore, we analyzed

the skeletal structures of the pectoral fins in *hoxaa*$^{-/-}$;*hoxab*$^{+/-}$;*hoxda*$^{+/-}$ and

hoxaa$^{+/-}$;*hoxab*$^{+/-}$;*hoxda*$^{-/-}$ fish, which were the surviving adult fish with the most of

these deletion mutations by using X-ray micro-CT scanning.

In our wild-type adult zebrafish, the number of fin rays in the pectoral fins is

10.8 on average (n=6, three specimens were examined for left and right pectoral fins;

Fig. 4A, J, Table 2). By contrast, statistical analysis showed that the number of fin rays

was slightly decreased to an average of 9.7 in the pectoral fins of

hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$ fish (n=6; Fig. 4B, J, Table 2), whereas that of

hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$ fish was comparable to that of wild-type fish (n=6; Fig. 4C, J,

Table 2). In addition, we observed that the anterior fin rays formed normally as in the

wild type, whereas one or two of the posterior fin rays were missing in

hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$ fish (Fig. 4B, Table 2; n=6/6, 100 % affected). Furthermore,

hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$ fish lost the pointed morphology of the proximal tips of the

posterior 7th and/or 8th fin rays (Table 2; n=3/6, 50 % affected).

hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$ fish retained the normal number of fin rays, but also lost the

pointed morphology of the proximal tips of the posterior four or five fin rays (Table 2;

n=3/6, 50 % affected). On the other hand, in the radial domains of endoskeletons

supporting the fin rays, the proximal radials are divided into four elements (pr1-pr4)

along the anteroposterior axis in wild-type fish [22] (Fig. 4A,D), although we noticed that

our wild-type zebrafish exhibited minor anomalies in the pectoral fins at low frequency

by micro-CT scanning (Table 2). However, *hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$* fish exhibited the

absence of the most posteriorly located pr4 and instead showed an enlargement of pr3

(n=5/6, 83 % affected; Fig. 4B, Table 2). Similarly, *hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$* fish also

showed the absence of pr4 (n=5/6, 83 % affected; Fig. 4C,F,I, Table 2). Furthermore, we

also observed a pectoral fin that lacked pr3 along with the absence of pr4 (n=3/6, 50 %

affected; Fig. 4C, Table 2). Previous study revealed that zebrafish *hoxa13a$^{-/-}$;a13b$^{-/-}$* and

hoxa13a$^{-/-}$;a13b$^{-/-}$;d13a$^{-/-}$ mosaic triple mutants showd the increased number of distal

radials[19]. To clarify whether the number of distal radials is also increased in

$hoxaa^{-/-};hoxab^{+/-};hoxda^{+/-}$ and $hoxaa^{+/-};hoxab^{+/-};hoxda^{-/-}$ fish, segmentation of micro-CT

scan images was performed to visualize the distal radials. In the wild-type, the number

of distal radials is 7.3 on average (Fig. 4G, K, Table 2). However, in

$hoxaa^{-/-};hoxab^{+/-};hoxda^{+/-}$ fish, the number of the distal radials averaged 11.2, and in

$hoxaa^{+/-};hoxab^{+/-};hoxda^{-/-}$ fish, the number increased to an average of 11.5 (Fig. 4H, I, K,

Table 2). Comparing the degree of abnormalities in the pectoral fins, slightly stronger

abnormalities were observed in $hoxaa^{+/-};hoxab^{+/-};hoxda^{-/-}$ fish than in

$hoxaa^{-/-};hoxab^{+/-};hoxda^{+/-}$ fish. This result may reflect the high contribution of the *hoxda*

cluster among the three *HoxA*- and *HoxD*-related clusters. These results suggest that in

zebrafish mutants with reduced function of *hoxaa*, *hoxab*, and *hoxda* clusters, a

pronounced structural defect was commonly observed in the posterior part of the

pectoral fins such as pr3, pr4, and the tip of the posterior fin rays. Interestingly, the

posterior part of the pectoral fin is thought to share molecular similarities with the limbs

of tetrapods[18,26-31]. Thus, the present loss-of-function study further supports the notion

that *Hox* genes in the *HoxA* and *HoxD* clusters already participated in patterning the

paired appendages before the divergence of the ray-finned and lobe-finned fishes.

Discussion

In this study, we generated compound mutants with different combinations of

zebrafish *hoxaa*, *hoxab*, and *hoxda* cluster deletions and analyzed their role in the

pectoral fin development. Combining these three *hox* cluster deletions increases early

pectoral fin development severity, suggesting that *hox* genes in *hoxaa*, *hoxab*, and

hoxda clusters function cooperatively as was observed in mouse *HoxA* and *HoxD*

clusters[8-15]. In a previous study, zebrafish *hoxa13a;hoxa13b* double mutants were

generated, showing that the endoskeletal disc is almost normal but the finfold is

shortened by about 30 % in the pectoral fin of larvae[19]. In this study, we showed the

pectoral fin phenotype of mutants lacking all *hoxaa*, *hoxab*, and *hoxda* clusters, and

both endoskeletal disc and finfold were found to be significantly shortened in

hoxaa$^{-/-}$*;hoxab*$^{-/-}$*;hoxda*$^{-/-}$ larvae. These abnormalities in our mutants are apparently

stronger than the pectoral fin abnormalities observed in *hoxa13a;hoxa13b* double

mutants[19]. The paralogous *hox* 9-13 genes of *HoxA*- and *HoxD*-related clusters are

expressed overlappingly in the developing pectoral fin in zebrafish[18,20], strongly

suggesting that these *hox9-hox13* genes contribute to pectoral fin formation in a

coordinated manner as was observed in mice.

When comparing the phenotypes of double *hox* cluster deletions in early

pectoral fin formation, the combination of *hoxab* and *hoxda* cluster deletions causes the

most severe phenotype (Fig. 1, 2). Detailed expression pattern analysis of the

developing pectoral fin demonstrated that zebrafish *hox9-13* genes in *hoxab* and *hoxda*

clusters are overlappingly expressed in mesenchymal cells in the fin buds from early

stages. In contrast, *hox* genes in *hoxaa* cluster are not expressed early, but are expressed

late in distal mesenchymal cells in the pectoral fins[18]. These similarities of the

expression patterns support that the most severe phenotype of *hoxab*$^{-/-}$*;hoxda*$^{-/-}$ larvae in

pectoral fin formation. It is also possible that the combination of deletions in *hoxab* and

hoxda clusters is the only combination of the three types of *hox* double deletions that

cannot be functionally complemented in part because the remaining *hoxaa* cluster has

no paralogous *hox10* and *hox12* (Fig. S1).

In the absence of 19 *hox* genes of zebrafish *hoxaa*, *hoxab*, and *hoxda* clusters,

we observed that the expression of *tbx5a* in the pectoral fin buds was not altered (Fig.

3A, B). These results suggest that the formation of the pectoral fin buds occurs normally

but subsequent cell differentiation or cell proliferation in the pectoral fin growth is

severely impaired in *hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$* larvae. Consistent with this observation,

after the formation of fin buds, we found that *shha* expression in the developing pectoral

fins is significantly down-regulated in mutants lacking zebrafish *HoxA-* and

HoxD-related clusters (Fig. 3C-J). Since zebrafish *shha/syu* mutants have been shown to

exhibit substantially reduced cell proliferation in pectoral fin buds which results in the

truncation of pectoral fins[25,32], the shortening of the pectoral fins in

hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$ larvae could be primarily due to reduced *shha* expression in fin

buds. In addition, these results are similar to previous results in mice that *Shh*

expression was not detectable in the severely truncated forelimbs of *HoxA* and *HoxD*

cluster double mutant mice[14]. Therefore, our loss-of-function results suggest not only

that the functions of *Hox* genes in *HoxA* and *HoxD* clusters are conserved in pectoral fin

and forelimb development, but also that the regulatory network comprising *Hox-Shh*

genes in appendage formation was already established in common ancestors of the

lobe-finned and ray-finned fishes.

In this study, we used micro-CT scans to examine the pectoral fins of

hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$ and *hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$* adult fish (Fig. 4). When

comparing our findings to previously reported *hoxa13a;a13b* mutants and

hoxa13a;a13b;d13a mosaic mutants (*hox13* mutants)[19], we noticed both similarities and

differences. One common feature is the increased number of distal radials, which is

consistent across all these mutants. Additionally, both *hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$* and

hoxaa$^{+/-}$*;hoxab*$^{+/-}$*;hoxda*$^{-/-}$ fish exhibit morphological abnormalities in the posterior

proximal radials, similar to *hox13* mutants[19]. Our multiple *hox* cluster deletion mutants

retain some *hox13* genes, suggesting that other posterior *hox* genes in *HoxA*- and

HoxD-related clusters may also play a role in the formation of proximal and distal

radials. On the other hand, unlike *hox13* mutants where the fin rays are lost[19], our

mutants only show a slight reduction in the number of fin rays while still retaining the

fin rays. This suggests that, unlike proximal and distal radials, a small amount of *hox13*

may be sufficient for the formation of fin rays in the pectoral fin.

We also detected common abnormalities in proximal radials and fin rays in the

posterior portion of the pectoral fins in *hoxaa*$^{-/-}$*;hoxab*$^{+/-}$*;hoxda*$^{+/-}$ and

hoxaa$^{+/-}$*;hoxab*$^{+/-}$*;hoxda*$^{-/-}$ fish, (Fig. 4). As no apparent abnormalities were observed in

these anterior skeletons, the reduced function of zebrafish *HoxA*- and *HoxD*-related

clusters seems to preferentially influence the posterior portion of the pectoral fins. In

zebrafish, the expression of posterior *HoxA*- and *HoxD*-related genes is progressively

restricted to the posterior portion of the developing pectoral fins, after the formation of

fin buds[18-20]. Furthermore, it has been shown that the phenotypes in which the posterior

skeletons of the pectoral fin are altered by specific mutations can be suppressed by

adding further null mutations in *hoxa11a*, *hoxa11b*, and *hoxd11a*[33]. Importantly, several

lines of evidence have indicated that the posterior structure of the pectoral fin has a gene

network similar to that of the limbs of tetrapods [18,26-31]. Our study demonstrates that

zebrafish *hox* clusters, homologous to *HoxA* and *HoxD* clusters, which are essential for

limb development in mice, also play an important role in the specific skeletal structures

of adult zebrafish pectoral fin which are considered to be potentially homologous to the

tetrapod limbs.

In this study, zebrafish three *HoxA*- and *HoxD*-related clusters were shown to

function together in zebrafish pectoral fin formation, similar to what has been observed

in mice[8-15]. While the focus of this study was on their role in pectoral fin development, it

is also possible that zebrafish *hoxaa*, *hoxab*, and *hoxda* clusters cooperate in other

developmental processes. Interestingly, we have shown that mutants with more than two

deletions of these clusters are lethal, although the exact cause of the lethality remains

unknown. Some of the finless mutants in zebrafish can survive [32,34,35], suggesting that

pectoral fin shortening does not necessarily result in lethality and may be linked to other

developmental events related to survival. In mice, it has been shown that cooperative

functions of *HoxA* and *HoxD* clusters are essential for urogenital development [10]. This

should be clarified in the future to elucidate the roles of *hoxaa*, *hoxab*, and *hoxda*

clusters during zebrafish development.

Materials and methods

Zebrafish husbandry

Zebrafish (Riken WT; RW), which were provided by the National BioResource Project (NBRP) Zebrafish in Japan, were maintained at 27°C with a 14-h light/10-h dark cycle. Embryos were obtained from natural spawning and the larvae were raised at 28.5°C. Developmental stages of the embryos were determined based on days post-fertilization (dpf) and morphological features as previously described[36]. Zebrafish mutants used in this study are *hoxaa* cluster-deficient*sud111*, *hoxab* cluster-deficient*sud112*, *hoxda* cluster-deficient*sud116*, which were previously generated[21].

Genotyping of mutants

For the genotyping of mutants, PCR was carried out. To partially dissect the caudal fins in juvenile and adult fish, the fish were anesthetized using the minimum required dose while checking the response with tricaine. Then, genomic DNA was extracted by NaOH method and used as a PCR template [37]. PCR was performed as described previously for the genotyping of *hox* cluster-deficient mutants[21]. After the reactions, the PCR products were separated by the electrophoresis in 2 % agarose gel in 0.5 x TBE buffer.

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization was performed for zebrafish embryos as previously described [38]. After staining, the tail region was dissected with a disposable scalpel (Feather). Genotyping was performed using genomic DNA extracted from the

315   dissected tail as described above. Stained embryos with the desired genotype were

316   selected and photographed under a stereomicroscope (Leica M205 FA) with a digital

317   camera (Leica DFC350FX).

319   **Alcian blue staining and the measurements of the endoskeletal disc and fin-fold**

320   Zebrafish larvae at 5 dpf were anesthetized in tricaine or MS-222 and then

fixed with 4 % paraformaldehyde in PBS. Alcian blue staining under strong acid

conditions makes it difficult to extract the genomic DNA required for genotyping after

staining. Thus, acid-free Alcian blue staining was performed as described previously[39].

After Alcian blue staining, genomic DNA was extracted from the posterior body, which

was manually detached under a stereomicroscope with forceps, and the genotype was

determined by PCR as previously described[21]. For the larvae with the desired genotype,

pectoral fins were manually dissected with forceps under a stereomicroscope, mounted

in 50 % glycerol on a glass slide (Matsunami), and photographed under a

stereomicroscope (Leica M205FA). From these images, the longest lengths of the

endoskeletal disc and fin-fold along the proximal-distal axis were measured using

ImageJ software.

X-ray micro-computed tomography (CT) scanning analysis

334   The pectoral fin skeletons of zebrafish were analyzed by X-ray micro-CT

scanning as essentially previously described[40]. Briefly, adult zebrafish were anesthetized

in 0.01% tricaine (w/v) and sacrificed in an ice water bath. Then specimens were fixed

with 4 % paraformaldehyde in PBS at 4 °C overnight and transferred to 70 % ethanol.

The regions surrounding the pectoral fins were manually dissected by forceps from the

zebrafish body under the stereomicroscope. Using an X-ray micro-CT

(ScanXmate-E090S105; Comscantechno), the dissected pectoral fins were scanned at a

tube peak voltage of 85 kV and a tube current of 90 µA. The micro-CT data were

constructed using coneCTexpress software (Comscantechno) and stored as a dataset

with an isotropic resolution of 3.1-3.4 µm. Three-dimensional image analysis was

performed using OsiriX MD software (Pixmeo). Finally, movies were edited using

Adobe Premiere Pro (Adobe) and DaVinci Resolve (Blackmagic Design).

Digital dissection of zebrafish pectoral fin skeletal structures

348 Based on the data of the skeletal structures of the pectoral fin obtained from

349 micro-CT scans, each zebrafish skeletal element was manually dissected using the

350 Imaris contour surface feature (in the Measurement Pro) and displayed in a 3D image

351 using Imaris v10.0 (Bitplane).

Ethics approval

354 All the experiments using live zebrafish were approved by the Committee for

355 Animal Care and Use of Saitama University (R2-A-1-6, R3-A-1-6, R4-A-1-7,

356 R5-A-1-7) and carried out according to the Animal Research Reporting of In Vivo

357 Experiments (ARRIVE) guidelines and to relevant guidelines and regulations.

Acknowledgments

We thank the NBRP zebrafish for providing the RW strain and for preserving

zebrafish *hox* cluster-deficient lines used in this study. We also thank Koji Tamura and

Gembu Abe for providing the zebrafish *tbx5a* plasmid for *in situ* hybridization. This

work was supported by KAKENHI Grants-in-Aid for Scientific Research from the

Ministry of Education, Culture, Sports, Science, and Technology, Japan (18K06177,

23K05790 to A.K.) and by the National Institute of Genetics under the Joint Research

and Research Meeting (NIG-JOINT) program (38A2019, 7A2020, 66A2021, 18A2022,

31A2023 to A.K.).

369 Data availability

370   The datasets used and/or analysed during the current study are available from

371   the corresponding author on reasonable request.

373 Author Contributions

374   A.K. conceived and designed the experiments, M.I., H.N., H.K., R.F., T.T.,

375   S.O., R.K., and A.K. performed the embryonic phenotype analysis, A.M. performed

376   X-ray micro-CT scan analysis and constructed the 3D-movie, A.K. wrote the manuscript.

377   All authors discussed the results and commented on the manuscript.

379 Conflict of interest

The authors declare no conflict of interest.

Figure legends

Figure 1. Shortening of pectoral fins in *hoxaa;hoxab;hoxda* cluster-deficient larvae.

(A-H) Shortening of the pectoral fins in *hoxaa*, *hoxab*, and *hoxda* cluster-deleted

mutants. Dorsal views of larvae at 3 dpf. Arrowhead indicates severely truncated

pectoral fins in *hoxaa*$^{-/-}$;*hoxab*$^{-/-}$;*hoxda*$^{-/-}$ larva. Embryos were obtained from the natural

intercrosses between *hoxaa*$^{+/-}$;*hoxab*$^{+/-}$;*hoxda*$^{+/-}$ triple hemizygous mutant fish and were

incubated at 28.5°C in water containing methylene blue as an anti-fungal agent (green

signal in the otic vesicle). After photography, PCR-based genotyping was performed. At

least three individuals (*n*>3) were identified for each genotype, and the representative

larvae are shown. Scale bar, 100 μm.

Figure 2. Shortening of the endoskeletal disc and fin-fold of the pectoral fins in

***hoxaa*$^{-/-}$;*hoxab*$^{-/-}$;*hoxda*$^{-/-}$ larvae.**

(A-H) Reduction of the endoskeletal disc (ed) and fin-fold (ff) in the pectoral fins of

hoxaa;hoxab;hoxda cluster-deleted mutants. Alcian blue staining was performed to

distinguish cartilaginous cells (endoskeletal disc) and non-cartilaginous fin-fold in

larvae at 5 dpf. After the staining, the pectoral fins were manually dissected and

mounted on a glass slide. The margin of the endoskeletal disc is emphasized with a

dashed line, and the margin of the fin-fold is highlighted with a line. Proximal to the left.

Scale bar, 100 μm. The number of dissected pectoral fins for which lengths were

measured is noted in the left bottom. (I-K) Comparisons of the length of the

endoskeletal disc and fin-fold in *hoxaa;hoxab;hoxda* cluster-deleted mutants. The

lengths of the endoskeletal disc and the fin-fold were measured by ImageJ. The

Tukey-Kramer test was performed with $*P < 0.05$, $**P < 0.01$, $***P < 0.001$,

and $****P < 0.0001$. Error bars represent the standard error of the mean.

Figure 3. Expression of *shha* in developing pectoral fin is significantly decreased in

***hoxaa;hoxab;hoxda* triple homozygous mutants.**

Whole-mount *in situ* hybridization was performed. Dorsal view. (A, B) Expression

patterns of *tbx5a* in wild-type siblings (n=5) and *hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$* embryos

(n=3) at 24 hpf. (C-J) Expression patterns of *shha* in the embryos at 48 hpf which were

derived from several intercrosses between *hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$* fish. Prior to

imaging, PCR-based genotyping was performed using genomic DNA extracted from the

tail of the stained embryos. At least three embryos were identified for each genotype to

confirm the reproducibility. Arrowhead indicates significantly decreased signal of *shha*

in the pectoral fin primordium of *hoxab$^{-/-}$;hoxda$^{-/-}$* and *hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$*

embryos. Scale bar, 100 µm.

Figure 4. Skeletal elements in the posterior regions of the pectoral fin are

preferentially affected in *hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$* and *hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$*

adult fish.

Micro-CT scan analysis of the skeletal structures in zebrafish pectoral fins. Both sides

of the pectoral fins in wild-type (n=3), *hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$* (n=3), and

hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$ (n=3) adult zebrafish were analyzed. The left pectoral fins are

shown. (A-C) The magnified views of pectoral fin endoskeletons from the lateral side.

(D-F) Magnified views of the pectoral fin endoskeletons from the opposite medial side

in the same specimens shown in A-C. Proximal radials (pr1-pr4) are indicated. Fin rays

are numbered from the anterior to the posterior. The asterisk indicates the absence of the

posterior proximal radials. The arrowhead indicates the loss of the proximal tip of the

fin rays. Anterior is to the top. The 3D movies were created using OsiriX MD ver13 and

are available in Movies S1-3. Scale bar, 500 μm. (G-I) Magnified images of proximal

radials (grey) and distal radials (red) as revealed by CT segmentation. The same

specimens as above were used for each genotype. Fin rays were removed by image

processing. The first distal radials are not shown due to their distance from the

remaining distal radials. 3D movies were created using Imaris ver10 (Movies S4-6). A

2D movie showing the distal radials and proximal radials was also created using Imaris

ver10 and is provided in Movie S7. Scale bar, 250 μm. (J,K) Comparisons of the

number of fin rays and distal radials between wild-type, *hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$* and

hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$ adult fish. The Tukey-Kramer test was performed with

P<0.05 and **P*<0.01.

Table 1.   Genotyping of juvenile fish derived from the intercrosses between

__hoxaa;hoxab;hoxda__ **triple hemizygous fish.**

Embryos were obtained by the natural spawning of intercrosses between triple

hemizygous *hoxaa;hoxab;hoxda* fish and reared to juveniles of 1-2 months of age. Their

genotype was then determined by PCR using genomic DNA extracted from the

dissected caudal fin. The expected percentage was calculated according to Mendel's law.

It appears that the nonviable mutants die after 7 days after fertilization.

Table 2. Summary of the abnormal morphology of the skeletal elements in

pectoral fins.

The left and right pectoral fins of three specimens each of wild-type,

hoxaa$^{-/-}$;hoxab$^{+/-}$;hoxda$^{+/-}$, and *hoxaa$^{+/-}$;hoxab$^{+/-}$;hoxda$^{-/-}$* mutants were analyzed by

micro-CT scanning and the abnormalities in the skeletal elements were summarized. In

the table, "-" indicates the normal morphology. For each genotype, #1 is shown in Fig.

4.

References

1. Dolle, P., Izpisua-Belmonte, J. C., Falkenstein, H., Renucci, A. & Duboule, D. Coordinate expression of the murine Hox-5 complex homoeobox-containing genes during limb pattern formation. *Nature*. **342**, 767-772 (1989).

2. Duboule, D. & Dolle, P. The structural and functional organization of the murine HOX gene family resembles that of Drosophila homeotic genes. *EMBO J*. **8**, 1497-1505 (1989).

3. Graham, A., Papalopulu, N. & Krumlauf, R. The murine and Drosophila homeobox gene complexes have common features of organization and expression. *Cell*. **57**, 367-378 (1989).

4. Dehal, P. & Boore, J. L. Two rounds of whole genome duplication in the ancestral vertebrate. *PLoS Biol*. **3**, e314 (2005).

5. Ohno, S. Evolution by gene duplication (1970).

6. Izpisua-Belmonte, J. C. & Duboule, D. Homeobox genes and pattern formation in the vertebrate limb. *Dev Biol*. **152**, 26-36 (1992).

7. Krumlauf, R. Hox genes in vertebrate development. *Cell*. **78**, 191-201 (1994).

8. Boulet, A. M. & Capecchi, M. R. Multiple roles of Hoxa11 and Hoxd11 in the formation of the mammalian forelimb zeugopod. *Development*. **131**, 299-309 (2004).

9. Davis, A. P. & Capecchi, M. R. Axial homeosis and appendicular skeleton defects in mice with a targeted disruption of hoxd-11. *Development*. **120**, 2187-2198 (1994).

10. Davis, A. P., Witte, D. P., Hsieh-Li, H. M., Potter, S. S. & Capecchi, M. R. Absence of radius and ulna in mice lacking hoxa-11 and hoxd-11. *Nature*. **375**, 791-795 (1995).

11. Dolle, P. *et al.* Disruption of the Hoxd-13 gene induces localized heterochrony leading to mice with neotenic limbs. *Cell*. **75**, 431-441 (1993).

12. Fromental-Ramain, C. *et al.* Specific and redundant functions of the paralogous Hoxa-9 and Hoxd-9 genes in forelimb and axial skeleton patterning. *Development*. **122**, 461-472 (1996).

13. Fromental-Ramain, C. *et al.* Hoxa-13 and Hoxd-13 play a crucial role in the patterning of the limb autopod. *Development*. **122**, 2997-3011 (1996).

14. Kmita, M. *et al.* Early developmental arrest of mammalian limbs lacking HoxA/HoxD gene function. *Nature*. **435**, 1113-1116 (2005).

15. Small, K. M. & Potter, S. S. Homeotic transformations and limb defects in Hox A11 mutant mice. *Genes Dev*. **7**, 2318-2328 (1993).

16. Nelson, C. E. *et al.* Analysis of Hox gene expression in the chick limb bud. *Development*. **122**, 1449-1466 (1996).

497   17.   Amores, A. *et al.* Zebrafish hox clusters and vertebrate genome evolution. *Science*. **282**,
498   1711-1714 (1998).

499   18.   Ahn, D. & Ho, R. K. Tri-phasic expression of posterior Hox genes during development
500   of pectoral fins in zebrafish: implications for the evolution of vertebrate paired
501   appendages. *Dev Biol*. **322**, 220-233 (2008).

502   19.   Nakamura, T., Gehrke, A. R., Lemberg, J., Szymaszek, J. & Shubin, N. H. Digits and
503   fin rays share common developmental histories. *Nature*. **537**, 225-228 (2016).

504   20.   Sordino, P., van der Hoeven, F. & Duboule, D. Hox gene expression in teleost fins and
505   the origin of vertebrate digits. *Nature*. **375**, 678-681 (1995).

506   21.   Yamada, K. *et al.* An atlas of seven zebrafish hox cluster mutants provides insights into
507   sub/neofunctionalization of vertebrate Hox clusters. *Development*. **148** (2021).

508   22.   Grandel, H. & Schulte-Merker, S. The development of the paired fins in the zebrafish
509   (Danio rerio). *Mech Dev*. **79**, 99-120 (1998).

510   23.   Ahn, D. G., Kourakis, M. J., Rohde, L. A., Silver, L. M. & Ho, R. K. T-box gene tbx5 is
511   essential for formation of the pectoral limb bud. *Nature*. **417**, 754-758 (2002).

512   24.   Garrity, D. M., Childs, S. & Fishman, M. C. The heartstrings mutation in zebrafish
513   causes heart/fin Tbx5 deficiency syndrome. *Development*. **129**, 4635-4645 (2002).

514   25.   Neumann, C. J., Grandel, H., Gaffield, W., Schulte-Merker, S. & Nusslein-Volhard, C.
515   Transient establishment of anteroposterior polarity in the zebrafish pectoral fin bud in
516   the absence of sonic hedgehog activity. *Development*. **126**, 4817-4826 (1999).

517   26.   Freitas, R., Gomez-Marin, C., Wilson, J. M., Casares, F. & Gomez-Skarmeta, J. L.
518   Hoxd13 contribution to the evolution of vertebrate appendages. *Dev Cell*. **23**,
519   1219-1229 (2012).

520   27.   Gehrke, A. R. *et al.* Deep conservation of wrist and digit enhancers in fish. *Proc Natl*
521   *Acad Sci U S A*. **112**, 803-808 (2015).

522   28.   Onimaru, K. *et al.* A shift in anterior-posterior positional information underlies the
523   fin-to-limb evolution. *Elife*. **4** (2015).

524   29.   Shubin, N., Tabin, C. & Carroll, S. Deep homology and the origins of evolutionary
525   novelty. *Nature*. **457**, 818-823 (2009).

526   30.   Tanaka, M. Fins into limbs: Autopod acquisition and anterior elements reduction by
527   modifying gene networks involving 5'Hox, Gli3, and Shh. *Dev Biol*. **413**, 1-7 (2016).

528   31.   Woltering, J. M., Noordermeer, D., Leleu, M. & Duboule, D. Conservation and
529   divergence of regulatory strategies at Hox Loci and the origin of tetrapod digits. *PLoS*
530   *Biol*. **12**, e1001773 (2014).

531   32.   van Eeden, F. J. *et al.* Genetic analysis of fin formation in the zebrafish, Danio rerio.
532   *Development*. **123**, 255-262 (1996).

33.   Hawkins, M. B., Henke, K. & Harris, M. P. Latent developmental potential to form
limb-like skeletal structures in zebrafish. *Cell*. **184**, 899-911 e813 (2021).

34.   Adachi, U. *et al.* Teleost Hox code defines regional identities competent for the
formation of dorsal and anal fins. *Proc Natl Acad Sci U S A*. **121**, e2403809121 (2024).

35.   Don, E. K. *et al.* Genetic basis of hindlimb loss in a naturally occurring vertebrate
model. *Biol Open*. **5**, 359-366 (2016).

36.   Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. & Schilling, T. F. Stages of
embryonic development of the zebrafish. *Developmental dynamics : an official*
publication of the American Association of Anatomists. **203**, 253-310 (1995).

37.   Meeker, N. D., Hutchinson, S. A., Ho, L. & Trede, N. S. Method for isolation of
PCR-ready genomic DNA from zebrafish tissues. *BioTechniques*. **43**, 610, 612, 614
(2007).

38.   Thisse, B. & Thisse, C. In situ hybridization on whole-mount zebrafish embryos and
young larvae. *Methods Mol Biol*. **1211**, 53-67 (2014).

39.   Walker, M. B. & Kimmel, C. B. A two-color acid-free cartilage and bone stain for
zebrafish larvae. *Biotech Histochem*. **82**, 23-28 (2007).

40.   Akama, K. *et al.* Role of somite patterning in the formation of Weberian apparatus and
pleural rib in zebrafish. *J Anat*. **236**, 622-629 (2020).

Figure 1. Shortening of pectoral fins in *hoxaa;hoxab;hoxda* cluster-deficient larvae.
(A-H) Shortening of the pectoral fins in *hoxaa*, *hoxab*, and *hoxda* cluster-deleted mutants. Dorsal views of larvae at 3 dpf. Arrowhead indicates severely truncated pectoral fins in *hoxaa⁻/⁻;hoxab⁻/⁻;hoxda⁻/⁻* larva. Embryos were obtained from the natural intercrosses between *hoxaa⁺/⁻;hoxab⁺/⁻;hoxda⁺/⁻* triple hemizygous mutant fish and were incubated at 28.5°C in water containing methylene blue as an anti-fungal agent (green signal in the otic vesicle). After photography, PCR-based genotyping was performed. At least three individuals (*n*>3) were identified for each genotype, and the representative larvae are shown. Scale bar, 100 μm.

Figure 2. Shortening of the endoskeletal disc and fin-fold of the pectoral fins in *hoxaa⁻/⁻;hoxab⁻/⁻;hoxda⁻/⁻* larvae.

(A-H) Reduction of the endoskeletal disc (ed) and fin-fold (ff) in the pectoral fins of *hoxaa;hoxab;hoxda* cluster-deleted mutants. Alcian blue staining was performed to distinguish cartilaginous cells (endoskeletal disc) and non-cartilaginous fin-fold in larvae at 5 dpf. After the staining, the pectoral fins were manually dissected and mounted on a glass slide. The margin of the endoskeletal disc is emphasized with a dashed line, and the margin of the fin-fold is highlighted with a line. Proximal to the left. Scale bar, 100 μm. The number of dissected pectoral fins for which lengths were measured is noted in the left bottom. (I-K) Comparisons of the length of the endoskeletal disc and fin-fold in *hoxaa;hoxab;hoxda* cluster-deleted mutants. The lengths of the endoskeletal disc and the fin-fold were measured by ImageJ. The Tukey-Kramer test was performed with $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. Error bars represent the standard error of the mean.

Figure 3. Expression of *shha* in developing pectoral fin is significantly decreased in *hoxaa;hoxab;hoxda* triple homozygous mutants.
Whole-mount *in situ* hybridization was performed. Dorsal view. (A, B) Expression patterns of *tbx5a* in wild-type siblings (*n*=5) and *hoxaa*⁻/⁻;*hoxab*⁻/⁻;*hoxda*⁻/⁻ embryos (*n*=3) at 24 hpf. (C-J) Expression patterns of *shha* in the embryos at 48 hpf which were derived from several intercrosses between *hoxaa*⁺/⁻;*hoxab*⁺/⁻;*hoxda*⁺/⁻ fish. Prior to imaging, PCR-based genotyping was performed using genomic DNA extracted from the tail of the stained embryos. At least three embryos were identified for each genotype to confirm the reproducibility. Arrowhead indicates significantly decreased signal of *shha* in the pectoral fin primordium of *hoxab*⁺/⁻;*hoxda*⁺/⁻ and *hoxaa*⁻/⁻;*hoxab*⁻/⁻;*hoxda*⁻/⁻ embryos. Scale bar, 100 μm.

Figure 4. Skeletal elements in the posterior regions of the pectoral fin are preferentially affected in *hoxaa⁻/⁻;hoxab⁺/⁻;hoxda⁺/⁻* **and** *hoxaa⁺/⁻;hoxab⁺/⁻;hoxda⁻/⁻* **adult fish.**
Micro-CT scan analysis of the skeletal structures in zebrafish pectoral fins. Both sides of the pectoral fins in wild-type (*n=3*), *hoxaa⁻/⁻;hoxab⁺/⁻;hoxda⁻/⁻* (*n=3*), and *hoxaa⁺/⁻;hoxab⁺/⁻;hoxda⁻/⁻* (*n=3*) adult zebrafish were analyzed. The left pectoral fins are shown. (A-C) The magnified views of pectoral fin endoskeletons from the lateral side. (D-F) The magnified views of the pectoral fin endoskeletons from the opposite medial side in the same specimens shown in A-C. Proximal radials (pr1-pr4) are indicated. Fin rays are numbered from the anterior to posterior. The asterisk indicates the absence of the posterior proximal radials. The arrowhead indicates the loss of the proximal tip of the fin rays. Anterior is to the top. The 3D movies are available in Movies S1-3. Scale bar, 500 μm. (G-I) Magnified images of proximal radials (grey) and distal radials (red) as revealed by CT segmentation. The same specimens as above were used for each genotype. Fin rays were removed by image processing. The first distal radials are not shown due to their distance from the remaining distal radials. For each genotype, 3D movie is available in Movies S4-6. 2D movie showing the distal radials and proximal radials is provided in Movie S7. Scale bar, 250 μm. (J,K) Comparisons of the number of fin rays and distal radials between wild-type, *hoxaa⁻/⁻;hoxab⁺/⁻;hoxda⁺/⁻* and *hoxaa⁻/⁻;hoxab⁺/⁻;hoxda⁻/⁻* adult fish. The Tukey-Kramer test was performed with *$P < 0.05$ and **$P < 0.01$.