Human microbiome-derived peptide affects the development of

experimental autoimmune encephalomyelitis via molecular

mimicry

Authors

Xin Ma[1, #], Jian Zhang[2, #], Qianling Jiang[1], Yong-Xin Li[2, *] and Guan Yang[1,3 *]

Affiliations

1. Department of Infectious Diseases and Public Health, Jockey Club College of

Veterinary Medicine and Life Sciences, City University of Hong Kong, Kowloon,

Hong Kong SAR, China.

2. Department of Chemistry and The Swire Institute of Marine Science, The

University of Hong Kong, Pokfulam Road, Hong Kong SAR, China.

3. Shenzhen Research Institute, City University of Hong Kong, Shenzhen, China

#equal contribution

*Corresponding Authors: Yong-Xin Li, Guan Yang.

Email: yxpli@hku.hk, gyang25@cityu.edu.hk

Abstract

Background: Gut commensal microbiota has been identified as a potential

environmental risk factor for multiple sclerosis (MS), and numerous studies have

linked the commensal microorganism with the onset of MS. However, little is known

about the mechanisms underlying the gut microbiome and host-immune system

interaction. **Results:** Here, we introduce the concept of molecular mimicry to

address this issue by mining human microbial-derived peptides based on their

similarity to the MHC II-TCR binding pattern of self-antigens. We analyzed 304,246

human microbiome genomes and 103 metagenomes collected from the MS cohort

and identified 731 nonredundant analogs of myelin oligodendrocyte glycoprotein

peptide 35-55 (MOG_{35-55}). Of note, half of these analogs could bind to MHC II and

interact with TCR through structural modeling of the interaction using fine-tuned

AlphaFold. Among the 8 selected peptides, the peptide (P3) derived from human gut

commensal *Akkermansia muciniphila* shows the ability to activate MOG_{35-55}-specific

CD4[+] T cells *in vitro* and exacerbate the development of experimental autoimmune

encephalomyelitis (EAE) in mice. Furthermore, dendritic cells could process and

present P3 to MOG-specific CD4[+] T cells and activate these cells. Collectively, our

data suggests the potential involvement of a MOG_{35-55}-mimic peptide derived from

the gut microbiota as a molecular trigger of EAE pathogenesis. **Conclusions:** Our

findings offer direct evidence of how microbes can initiate the development of EAE,

suggesting a potential microbiome-based therapeutic target for inhibiting the

progression of MS.

Keywords: multiple sclerosis, experimental autoimmune encephalomyelitis,

molecular mimicry, gut microbiota

Introduction

Multiple sclerosis (MS) is an autoimmune disease characterized by the involvement

of autoreactive T cells targeting myelin antigens, leading to the destruction of myelin

and damage to axons, resulting in neurologic syndromes and physical disability (1).

The global prevalence of MS stands at approximately 2.8 million individuals (2), with

an increasing incidence observed in developing countries and among children (3).

Although the etiology of MS remains unknown, both genetic and environmental

factors have been implicated in its development (1). Accumulating evidence

suggests that gut microorganisms play a pathogenic role in MS (4, 5). Research has

found notable differences in the composition of gut bacteria between MS

patients and healthy individuals (6, 7). Similar alterations in the gut microbiota,

including elevated levels of *Akkermansia muciniphila* (*A. muciniphila*), have also

been observed in experimental autoimmune encephalomyelitis (EAE), an animal

model employed to investigate MS (8). However, how the alterations in the gut

microbiota might affect MS/EAE development remains largely unknown.

Molecular mimicry is one of the leading mechanisms in linking the role of microbiota

in immune reactivity in autoimmune diseases and cancer (9-13). Reassuringly, it has

been reported that microbiota-derived peptides resembling myelin basic protein

(MBP) can activate autoreactive T cells in MS (14-16). MBP, along with proteolipid

protein (PLP) and myelin oligodendrocyte glycoprotein (MOG), are myelin antigens

targeted by HLA-DR-restricted T cells, leading to MS development (1). Of note,

MOG has been considered a primary target of cellular and humoral immune

responses in multiple diseases due to its special extracellular immunoglobulin

domain, which is directly accessible to binding partners including antibodies (17-19).

Furthermore, MOG 35-55 (MOG$_{35-55}$) peptide has been widely used to induce EAE.

Intrigued by these findings, we hypothesized that microbiome-derived peptides

resembling the MOG$_{35-55}$ epitope could trigger the autoimmune response

responsible for initiating EAE.

To address this hypothesis, we analyzed human microbiome genomes and

metagenome databases collected from MS cohort. Our aim was to identify microbial

peptides with MHC II-TCR recognition patterns resembling those of MOG$_{35-55}$, which

is presented by the MHC class II I-A^{b} and induces chronic EAE in C57BL/6J mice

(20). Subsequently, we identified, synthesized, and tested 8 of these peptides for

their ability to activate MOG$_{35-55}$-specific T cells. Our findings revealed that a

predicted peptide (P3) derived from the gut commensal organism *A. muciniphila* can

activate mouse MOG$_{35-55}$-specific T cells *in vitro*, albeit to a lesser extent compared

to the MOG$_{35-55}$ peptide. Additionally, P3 immunization can exacerbate the

development of EAE. We also observed some cross-reactivity between MOG_{35-55}

peptide and $CD4^+$ T cells obtained from mice immunized with P3. Furthermore,

dendritic cells (DCs) were able to process P3 and present it to $CD4^+$ T cells isolated

from 2D2 mice expressing a MOG-specific T cell receptor. Our findings suggest that

microbial peptide derived from the gut commensal, resembling MOG_{35-55}, may mimic

the native myelin peptide and potentially contribute to the EAE development.

Results

Well-defined peptide-MHC II-TCR binding motifs guided microbiome-derived

epitope discovery from the human microbiome

To uncover the microbiome-derived epitopes resembling the immunodominant

MOG_{35-55} peptide, we designed a genome mining workflow involving an analysis of

a vast dataset comprising 304,246 human microbiome genomes (21, 22) and 103

metagenomic samples from a MS study cohort (23) (Fig 1A). We first extracted

404,181 MOG_{35-55} analogs (Supplementary table 1) from these resources, followed

by a filtering process that retained 21,557 analogs (3,946 non-redundant sequences)

possessing well-defined binding residues at specific sequential spacing (Fig 1B,

Supplementary table 2).

To gain insight into the phylogenetic distribution of MOG_{35-55} analogs derived from

the human microbiome, we examined the 21,557 analogs with well-defined binding

motifs. Those analogs exhibited a wide distribution across 24 phyla, with the

occurrence of a specific motif per genome varying from 0 to 1 within each phylum

(Fig 1C). Of note, 10 phyla with fewer than 10 genomes and 19 unclassified

bacteria-derived contigs at the phylum level were grouped as "Others". Interestingly,

Verrucomicrobiota demonstrated a higher likelihood of encoding analogs featuring

the YRxxFxRVx motif. Besides, these analogs were predominantly encoded by

species such as *A. muciniphila* or *A. muciniphila.B*, which are enriched in patients

with MS (24).

Microbiome-derived peptide-MHC II-TCR complexes structure-prediction

Following the general model of antigen presentation (25, 26), we employed a fine-

tuned AlphaFold with a binding classifier (27) to discriminate ligands with mouse

MHC II molecules (H2-Ab) binding core peptides. Subsequently, the binding core

peptides were subjected to modeling the peptide-MHC II-TCR complex by TCRdock

(28). First, the performance of fine-tuned AlphaFold and TCRdock was assessed

using the MOG_{35-55} peptide as a test case (Fig 2). MOG_{35-55} peptide was further

chopped into 10 unique putative binding core peptide candidates (Fig 2A) for

subsequent analyses. This was necessary because the current state-of-the-art

machine learning-based structure predictors (27-30) for the peptide-MHC II-TCR

complex are designed to predict binding between a 9-mer binding core peptide and

its receptor (MHC II or TCR). Finally, the binding core peptide of MOG_{35-55}, with the

specific sequence "WYRPPFSRVVH", was predicted to be MHC II binders (Fig 2A),

and this binding core peptide exhibited reliable interactions with four TCRs that

contained different gene segments (Fig 2B). It is noteworthy that this binding core

peptide with the sequence "WYRPPFSRVVH" could interact with mouse MHC II and

TCR is consistent with the previous studies (20, 31). Inspired by these results, we

integrated these two methods to prioritize the microbiome-derived epitopes.

To mimic the process of proteolytical degradation of exogenous proteins before their

presentation to MHC II in antigen presentation (26), we truncated the corresponding

peptide windows (~ 21 residues) containing the binding motifs along these 3,946

analogs, resulting in 731 nonredundant putative ligand sequences (Fig 3A,

Supplementary table 2, 3). These ligands were further chopped into 5,783 unique

putative binding core peptide candidates (Supplementary table 3) for subsequent

analyses. Consequently, we identified 493 binding core peptides from 439 ligands

as MHC II binders (Fig 3B, Supplementary table 4). In addition, we successfully

modeled interactions between peptide-MHC II and 4 types of TCRs for 97.8% of the

MHC II binders using 4 peptide-MHC-TCR complexes templates (Supplementary

Fig 1 and Fig 2, Supplementary table 4). To determine the binding specificity,

binding motifs are generated from these binding core peptides and characterized by

anchor positions (Fig 3C and 3D). The motifs revealed that Y/F, R, F, R, and V

residues are enriched at well-defined anchor positions (loc_1, loc_2, loc_5, loc_7,

loc_8) in this study. Surprisingly, several noncanonical binding motifs starting with

"L", "A", "I", "S", "N", "H", "Q", "V", "T", "M", "W", and others were also observed (Fig

3C). These findings suggest that microbiome-derived epitopes could interact with

MHC II and TCRs with diverse binding characteristics.

Microbiome-derived peptide stimulates MOG$_{35-55}$-specific T cell

In order to investigate the potential modulatory effects of microbiome-derived

peptides resembling MOG$_{35-55}$ peptide sequence on the autoimmune response that

triggers EAE, a total of 8 peptides (Fig 3 and 4A) were selected from a pool of 439

ligands. These 8 peptides were confidently modeled in pMHC II-TCR interactions

(Fig 3), and their selection was based on four specific criteria. Firstly, ligands that

contained the "YRxxFxRVx" motif were considered. Secondly, ligands that could be

detected in the NBDC database were included (specifically, P1, P2, P4, P5, P6, and

P8). Thirdly, ligands collected from species related to MS were chosen (P3 and P5)

(24). Finally, ligands containing noncanonical binding motifs starting with "F" and "L"

were selected (P1, P3, P7, and P8), as these motifs were found to be most enriched

in the pool of 493 binding core peptides. To investigate the potential of microbe-

derived peptides to activate MOG$_{35-55}$-specific T cell, we stimulated splenocytes

isolated from MOG$_{35-55}$-induced EAE mice with MOG$_{35-55}$ peptide and 8 selected

peptides (Fig 4A). Of the tested 8 microbiome-derived peptides, P3 demonstrated

the capacity to stimulate the production of IL-17 and IFN-γ by MOG$_{35-55}$-specific T

cells (Fig 4B-D, supplementary Fig 3A). It is noteworthy that P3 was identified as

originating from *A. muciniphila*, a gut commensal that is significantly enriched in the

gut microbiota of individuals with MS (5).

P3 exacerbate EAE development in mice and shows bidirectional cross

reactivity with MOG$_{35-55}$ peptide *in vitro*

To further explore the impact of microbiome-derived peptide on EAE development,

C57BL/6J mice were immunized with a combination of P3 and a low dose of MOG$_{35-}$

$_{55}$. We found that P3 could exacerbate EAE development in mice (Fig 5A). At 17

days post-immunization, the spinal cord tissues were collected, and flow cytometry

as well as Hematoxylin and Eosin (H&E) or Luxol Fast Blue (LFB) staining were

performed to determine the impact of P3 on the severity of inflammation and

demyelination. As shown in Fig 5B, increased leukocyte infiltration was observed in

the MOG$_{35-55}$ + P3 group, implicating more severe immune infiltration in this group.

Moreover, more severe inflammation was observed in the white matter of spinal

cord from mice immunized with P3 and MOG$_{35-55}$ compared to the control mice (Fig

5C). Quantitative analysis revealed that the inflammatory scores (H&E score) in the

MOG$_{35-55}$ + P3 group were significantly higher than the MOG$_{35-55}$ + control group

(Fig. 5C). Additionally, the demyelination level (LFB score) showed an increasing

trend in mice immunized with MOG$_{35-55}$ + P3 compared to the control mice (Fig. 5D).

Splenocytes from P3 immunized mice (10-day post-immunization) were collected to

further test the cross-reactivity between P3 and MOG$_{35-55}$. Following an 18-hour

stimulation of P3-specific CD4$^+$ T cells either with P3 or MOG$_{35-55}$, the cells exhibited

a robust immune response to P3 and a relatively strong response to MOG_{35-55} as

indicated by the production of IFN-γ (Fig 6). These data suggest that the microbial

peptide P3 and MOG_{35-55} can stimulate a bidirectional cross-reactive immune

response.

DCs process microbiome-derived peptide P3 and further activate 2D2 CD4$^+$ T

cell proliferation

Given that P3 stimulated MOG_{35-55}-specific CD4$^+$ T cells *in vitro*, we investigated

whether DCs are capable of processing and presenting P3 to MOG_{35-55}-specific T

cells. Our pMHC II-TCR model predicted an interaction between the 11- residue

windows with the core 9-mer residues truncated from microbiome-derived peptide

and the MHC II-TCR complex (Fig 7A and 7B), leading us to hypothesize that this

peptide could be presented by DCs. To test this hypothesis, we performed an

antigen presentation assay using DCs isolated from the spleens of C57BL/6J mice.

The DCs were primed with either PBS, P3, or MOG_{35-55}, and then co-cultured with

sorted 2D2 CD4$^+$ T cells. The activation status of CD4$^+$ T cells was later assessed

by quantifying the expression of the early T cell activation marker CD69.

Furthermore, the proliferation of CD4$^+$ T cells was evaluated by monitoring the

changes in the population of CFSE-labeled cells. The results demonstrated that P3

could be presented by DCs and subsequently activate the proliferation of 2D2 CD4$^+$

T cells (Fig 7C). However, P3 induces a weaker CD4$^+$ T cell activation and

proliferation than MOG$_{35\text{-}55}$ (Fig 7D), which may explain the lower incidence and

clinical score observed in P3-induced EAE (Supplementary Fig 4).

Discussion

The gut microbiota has been implicated in the development of autoimmune

diseases. Investigations of MS patients and the EAE mouse model have revealed

that the gut microbiome is a crucial factor in disease progression and severity (4, 5).

Alterations in gut bacterial composition, including reductions in butyrate-producing

bacteria and increases in *Akkermansia* and *Clostridia* populations have been

consistently observed in MS patients (5, 7). Moreover, transfer of microbiota from

MS patients to animal models has been shown to increase disease incidence,

suggesting that alterations in the microbiome may actively contribute to the

development of MS (32). Germ-free and antibiotic-treated mice have shown

resistant to EAE, further supporting the necessity of the microbiome for disease

induction (33, 34).

Despite significant advances in understanding the role of the gut microbiota in MS

pathogenesis, the underlying mechanisms leading to disease development remain

poorly understood. Continuous sampling of gut microbial antigens by innate immune

cells enhances the probability of presenting specific antigens to antigen-specific T

cells. The presentation of these antigens is influenced by polymorphisms in MHC

genes, suggesting a potential role for host-commensal microorganisms cross-

reactivity in the development and persistence of autoimmunity among genetically

predisposed individuals. Molecular mimicry has been proposed as a possible

mechanism by which microorganisms can induce autoimmune diseases (15).

According to this hypothesis, T cells that respond to an epitope derived from an

infectious agent may cross-react with a self-antigen that share sequence or

structural similarities with the original microbial peptide (35). Evidence supporting

this hypothesis has been observed in numerous autoimmune diseases, such as

antiphospholipid syndrome and Type 1 Diabetes (36, 37). Given the diverse range

of microorganisms in the gut, peptides derived from gut flora could potentially trigger

the development of MS. The activation of autoreactive T cells is a critical component

of the early autoimmune response in individuals who ultimately develop MS. While

multiple potential antigens exist, MOG is a major EAE antigen in H-2b mice and a

possible autoantigen in MS (20). The binding pattern of the MOG-MHC II-

TCR complex has been characterized, revealing that tyrosine (Y) in position 40 of

the MOG$_{35-55}$ peptide is the dominant MHC II-binding residue, occupying the p1

pocket. Meanwhile, arginine (R) 41, phenylalanine (F) 44, R46, and valine (V) 47

are the major amino acids involved in TCR recognition (20). Research suggests that

low-affinity hotspot mimicry, which involves a subset of crucial residues within the

TCR-binding footprint, rather than high-affinity structural mimicry, may be a more

common contributor to the initiation of autoimmune diseases (14, 20). In our study,

we focused on identifying MOG$_{35-55}$ mimics based on the similarity of essential

residues within the TCR-binding footprint, rather than relying on the direct amino

acid similarity between microbial-derived peptides and self-antigens, as reported in

studies on Type 1 Diabetes (37).

By leveraging the expanding genome databases for human gut microbes, we

identified 8 microbial peptides that exhibit significant analogy in their antigen-MHC

II-TCR binding patterns with MOG_{35-55} peptide. Our findings indicated that

microbiome derived P3 can activate $CD4^+$ T cells specific to MOG_{35-55} peptide in

mice. Interestingly, our bioinformatic analysis revealed that this peptide was derived

from *A. muciniphila*, suggesting a potential impact of *A. muciniphila* on EAE

development. Previous studies have explored the link between *A. muciniphila* and

EAE or MS, such as the findings of Cekanaviciute *et al*., who observed that *A.*

muciniphila promotes Th1 differentiation in human peripheral blood mononuclear

cells (5). Moreover, previous studies have demonstrated that a peptide from *A.*

muciniphila shares a similar sequence with a MS associated self-antigen, guanosine

diphosphate (GDP)–L-fucose synthase. This shared peptide sequence has the

ability to stimulate the proliferation of $CD4^+$ cerebrospinal fluid-infiltrating T cells from

MS patients (16).

In this study, we optimized the process of screening self-antigen mimics from gut

microbiota by integrating the binding affinity to MHC II and TCR into the selection

criteria, thereby enhancing the accuracy of identifying self-antigen analogs in the gut

microbiome. Furthermore, our findings suggest that the enrichment of *A. muciniphila*

in the gut microbiota of EAE mice and MS patients may have pathological

implications as this bacterium has the capability of generating MOG_{35-55} analogs.

However, our method utilizing metagenomic data can only annotate genome to

species level, and further determination is needed to identify the specific *A.*

muciniphila strain capable of producing MOG_{35-55} analogs.

A. muciniphila, a next-generation probiotic, has shown positive effects on several

diseases including obesity and diabetes (38). However, recent studies have

revealed an enrichment of *A. muciniphila* in the gut microbiome of MS patients.

Furthermore, transplantation of fecal samples from MS patients into the intestines of

EAE mice leads to more severe EAE symptoms (5), indicating the need to assess

the potential beneficial or harmful effects of *A. muciniphila* from multiple

perspectives. Although multiple strains of *A. muciniphila* have been identified (39), it

remains unclear whether the observed effects in different disease contexts are

attributable to strain differences. The predominant use of metagenomics and 16S

sequencing in numerous studies exploring the relationship between various

diseases and *A. muciniphila* has impeded the identification of specific target strains,

as these methods primarily facilitate gene annotation at the species level,

presenting challenges in strain-level identification. Our study suggests that *A.*

muciniphila may contribute to the development of EAE due to its capacity to

generate MOG_{35-55}-resembling peptides. However, more evidence are needed to

determine the target strain that generates MOG_{35-55} analogs. Future studies will also

focus on elucidating the role of synergistic or antagonistic effect among different

microbial populations in autoimmune diseases. This study highlights the

necessity for further research on the correlation between gut microbiota and

autoimmune diseases in the growing trend of widespread probiotic consumption.

In summary, our findings prove a mechanism of molecular mimicry in which a

particular sequence within commensal gut microbe can imitate crucial residues in

the MOG-MHC II-TCR binding footprint, consequently initiating or modifying

the immune response associated with EAE development. While our study focused

only on MOG_{35-55} mimics, other myelin antigens, such as MBP and PLP, may also

serve as targets for autoreactive $CD4^+$ T cells (40). Given the enormous number of

microbial peptides generated by the gut microbiota, it is possible that other microbial

peptides may exist and can mimic these antigens and activate related autoantigen

reactive T cells. Meanwhile, it's important to note that the induction of autoimmune

diseases through molecular mimicry is complicated and a single factor acting alone

is unlikely to trigger the disease. Instead, accompanying host genetic susceptibility

is considered to be linked with an abnormal immune response triggered by one or

more microbial antigens or molecules. Nonetheless, this discovery may provide

a therapeutic target and an opportunity to impede the progression of MS.

Conclusion

We have investigated the role of molecular mimicry in the link between microbial

flora and EAE development. We have identified MOG_{35-55} mimics derived from gut

commensal cross-react with MOG_{35-55}-specific $CD4^+$ T cells and significantly

exacerbate the development of EAE in mice. Notably, antigen-presenting assays

confirmed that DCs can process and present this gut commensal derived peptide to

MOG_{35-55}-specific $CD4^+$ T cells. Our findings offer direct evidence of how microbes

can initiate the development of EAE, suggesting a potential microbiome-based

therapeutic target for inhibiting the progression of MS.

Material and Methods

Identify the human MOG_{35-55} peptide analog in the human microbiome

genomes and human-associated metagenomic samples

In brief, we first collected and curated human microbiome-derived peptide resources,

which included human microbiome genomes and human-associated metagenomics

samples. In detail, we collected 286,997 human microbiome genomes from Unified

Human Gastrointestinal Genome dataset (UHGG, v1.0) (41), and 17249 genomes

from CIBIO species-level genome bins (21). Taxonomic assignments were

performed for genomes from CIBIO by GTDB-tk (42). In addition, we downloaded

103 metagenomes from MS patients in National Bioscience Database Center

(NBDC) Human Database with the accession number of hum0197 (43). The data

obtained from the UHGG and NBDC were acquired in accordance with human

research ethics guidelines and were approved by the Human Ethics sub-Committee

of the City University of Hong Kong (jcc2223ay001). For these metagenomic

samples, we applied metagenomic assembly to obtain contigs using MEGAHIT (44)

for each sample separately. Taxonomic classification was further conducted for the

resulting contigs using CAT (v5.2.3) (45). Taxonomic assignments for genomes or

contigs based on the Genome Taxonomy Database (46). Finally, all open reading

frames were predicted in each genome or contig using Prodigal-short (47). Human

MOG$_{35-55}$ sequence analog searching was conducted by BLASTp with an e-value

threshold of 0.001 in a pool of proteins from the above Human microbiome-derived

peptide resources.

The structure-based prediction of peptide-MHC II-TCR binding

Identified analogs were first filtered based on whether they contained the motifs with

[YR..F.RV.], [YR..F.R..], [YR..F..V.], [YR....RV], and [Y...F.RV.] are involved in the

peptide-MHC-TCR interactions. Next, a special peptide region (~ 21 residues)

containing the above binding pattern was truncated from these filtered analogs. The

yielded nonredundant truncated peptides were subjected to further analyses.

The peptide-MHC II-TCR binding prediction was performed in two steps: the

peptide-MHC II binding specificity prediction and TCR: peptide-MHC II interactions

modeling. Because the advanced peptide-MHC II structure prediction tools often

predict the binding structure composed by the binding core of peptides (normally

with 9 residues) and receptors (MHC II or TCR) (27-30, 48), these nonredundant

truncated peptides were further chopped into a series of consecutive 11-residue

windows along peptide with the putative core nine-mer residues and their immediate

neighboring residues. The resulting 11-residue windows were subjected to a fine-

tuned AlphaFold to predict peptide-MHC II binding with the following parameters: --

model_names model_2_ptm_ft --ignore_identities (27). Peptide-MHC II templates

(4p23_MH2_H2ABa_H2ABb.pdb,6mng_MH2_H2ABa_H2ABb.pdb,1muj_MH2_H2A

Ba_H2ABb.pdb) in the PDB template set were used (27). Mean Inter-chain lowest

predicted aligned error (PAE) terms corresponding to the peptide-MHC interactions

≤ 4.34 and peptides' per-residue confidence score (pLDDT) corresponding to local

structural accuracy ≥ 90 were used to classify MHC II binder and non-binder. Next,

peptide-MHC II-TCR binding for the predicted binders was modeled by a specialized

version of the neural network predictor AlphaFold and TCR–pMHC complex

templates-based method, TCRDock with 'model_2_ptm' and '--benchmark'

parameter set (28). Based on TCRDock's workflow, three simulations were

conducted for each target peptide-MHC II-TCR complex, and finally chose the

model with the lowest PAE between the TCR and pMHC. Given the hypervariable

gene segments in TCR alpha and beta chain largely determine the structure

modeling (49), we took TCR:pMHC II structural templates (PDB IDs: 3c60, 3rdt,

3c5z, 6mnn), pairing with four types of mouse TCR (Supplementary table 5), to

model the peptide-MHC II-TCR structures. The predicted structures were visualized

by PyMoL.

Mice

Eight-week-old C57BL/6J mice were obtained from the Laboratory Animal Research

Unit (LARU) of City University of Hong Kong. The 2D2 TCR transgenic mice

C57BL/6-Tg (Tcra2D2, Tcrb2D2) 1Kuch/J were purchased from The Jackson

Laboratory and bred in LARU. Mice were randomly allocated to each treatment

group. All experimental procedures were approved by the Animal Ethics Committee

of the City University of Hong Kong and performed in accordance with LARU

Guidelines.

EAE induction

For induction of active EAE, mice (8-week-old) were immunized subcutaneously at

two sites on the back flank area with 200 µg of MOG_{35-55}

(MEVGWYRSPFSRVVHLYRNGK) peptide (Thermo Fisher) emulsified in 200 µl

incomplete Freund's adjuvant (BD bioscience) supplemented with 2 mg/ml heat-

killed *Mycobacterium tuberculosis* (BD bioscience). On days 0 and 2 after

immunization, mice received intraperitoneal 400 ng of pertussis toxin (Calbiochem).

The same immunization procedure was employed to detect the encephalitogenic

capacity of P3, using a combination of P3 and MOG_{35-55} (P3: 200 µg; MOG_{35-55}:

50µg;). The other group was administered same dose of a combination of irrelevant

peptide (set as a control of P3) and MOG_{35-55}. EAE severity was scored daily using

the standard scale: 0, no disease; 1, loss of tail tone; 2, hind limb weakness; 3, hind

limb paralysis; 4, hind limb paralysis and forelimb paralysis or weakness; and 5,

moribund/death (50). To detect the cross reactivity between $MOG_{35\text{-}55}$ and

synthesized peptides, mice were injected with peptide by following the similar

immunization procedure (DG peptides, China). The same immunization protocol

was employed to induce EAE using P3.

Antigen recall assay

T cell antigen recall response was assessed in cells harvested from the spleen of

EAE mice 10 days after the challenge. Mice were anesthetized by 3% isoflurane

inhalation and then euthanized by cervical dislocation. Spleens were then collected

from the euthanized mice and crushed through a 70-µm cell strainer and centrifuged

at 400 × g for 5 min at 4 °C, then incubated in red blood cell lysis buffer (Thermo

Fisher) for 5 min. After centrifuge, isolated cells were plated at a density of $1\text{x}10^{6}$

cells/well and stimulated with synthesized peptide or $MOG_{35\text{–}55}$ (20 µg/ml final

concentrations) in cell culture media (RPMI1640, 10% FBS, 2 mM L-glutamine, 10

mM Hepes, 50 µM 2-mercaptoethanol, 100 IU penicillin, and 100 µg/mL

streptomycin), in a total volume of 200 µL per well. Cells were stimulated for 18h at

37 °C and 5% CO_2 atmosphere, then analyzed by flow cytometry for cytokine (IFN-γ,

IL-17A) producing cells.

Histopathology

Inflammation and demyelination of spinal cords was assessed at day 17 after

immunization. The lumbar enlargement of the spinal cords was isolated from

animals and instantly fixed in 4% paraformaldehyde for a minimum of 48 h, then

embedded in paraffin and sectioned at 5-μm thickness. Sections were stained with

H&E (Solarbio) or LFB (Abcam) to evaluate inflammation and demyelination. The

sections were scanned using a NanoZoomer S60 Digital slide scanner

(Hamamatsu). The degrees of inflammation (H&E scores) were calculated as 0,

normal; 1, cellular infiltrates partially around meninges and blood vessels; 2, 1–10

lymphocyte infiltrates in the parenchyma; 3, 11–100 lymphocyte infiltrates in the

spinal cord; 4, 100-300 lymphocyte infiltrates in the spinal cord; 5, 300-1000

lymphocyte infiltrates in the spinal cord; and 6, over 1000 lymphocyte infiltrates in

the spinal cord (51). The severity of demyelination (LFB scores) was analyzed as 0,

normal; 1, myelin sheath injuries in rare areas; 2, a few areas of demyelination; 3,

confluent perivascular or subpial demyelination; 4, massive demyelination involving

one half of the spinal cord; and 5, extensive demyelination involving the whole

spinal cord (51).

Antigen presentation assays

To obtain stimulator DCs, single cell suspensions from the spleen collected from

C57BL/6J mice were prepared by mechanical disruption on a 70-μm cell strainer.

Cell suspensions were resuspended in MACS buffer (0.5% BSA, 2 mM EDTA in

PBS) after lysing red blood cells. CD11c$^+$ cells were then negatively selected

according to the manufacturer's instructions using the mouse Pan Dendritic Cell

Isolation Kit (Miltenyitec). Isolated DCs were pulsed with either PBS, P3, or MOG$_{35-}$

$_{55}$ at 20 µg/ml final concentrations, using 2 x 10^4 cells in 96-well plate for 1 h at

37 °C. Cells were then washed extensively and used for co-culture with responder

2D2 CD4$^+$ T cells. Responder MOG$_{35-55}$-specific CD4$^+$ T cells were isolated from

2D2 TCR transgenic mice spleen by using CD4$^+$ T cells Isolation Kit (Miltenyitec).

Sorted CD4$^+$ T cells were labeled with Carboxyfluorescein succinimidyl ester (CFSE,

Thermo Fisher) following the user guide. 10^5 CFSE labeled 2D2 CD4$^+$ T cells were

then seeded in triplicate in a round bottom 96-well plate with 2 x 10^4 stimulated DCs

(5:1 ratio). DCs and T cells were cultured together for 72 h at 37 °C and 5% CO$_2$

atmosphere, then cells were prepared for flow cytometry to detect T cell proliferation.

Flow cytometry

Single cell suspensions from the spleen were obtained by mechanical disruption on

a 70-µm cell strainer, then incubated in red blood cell lysis buffer for 5 min. For the

staining process, dead cells were stained with Propidium Iodide (Sigma-Aldrich) or

Live/Dead Ghost dye (Tonbo Biosciences). For staining of surface markers, cells

were incubated in fluorescently labeled antibodies (CD45, CD3, CD4, CD69, TCR

V$_\beta$11) for 30 min in FACS buffer (PBS containing 2% FBS) at 4 °C. For intracellular

staining, cells were washed with 2 mL wash buffer, then fixed and permeabilized by

using Cytofix/Cytoperm™ Fixation/Permeabilization Kit (BD Biosciences) as

manufacturer's instructions. Fixed cells were incubated in fluorescently labeled

antibodies (IFN-γ, IL-17A) for 30 min in permeabilization buffer at 4 °C. All data were

collected on a BD Celesta flow cytometer (BD Biosciences) and analyzed using

FlowJo software (v.10.2, BD Biosciences).

Statistical analysis

Data were analyzed using GraphPad Prism software and reported as mean ± SEM

unless otherwise noted. The two-tailed unpaired Student *t test* was used for

samples with equal variances, whereas Welch's *t test* was employed for samples

with unequal variances. Statistical significance was defined as a *p*-value <0.05.

Abbreviations

MS: Multiple sclerosis

MOG: Myelin oligodendrocyte glycoprotein

MOG_{35-55}: Myelin oligodendrocyte glycoprotein peptide 35-55

EAE: Experimental autoimmune encephalomyelitis

DCs: Dendritic cells

A. muciniphila: Akkermansia muciniphila

MBP: Myelin basic protein

PLP: Proteolipid protein

UHGG: Unified Human Gastrointestinal Genome dataset

NBDC: National Bioscience Database Center

PAE: Predicted aligned error

pLDDT: Peptides' per-residue confidence score

Hematoxylin and Eosin: H&E

Luxol Fast Blue: LFB

Declarations

Ethics approval and consent to participate: The data obtained from the UHGG

and NBDC were acquired in accordance with human research ethics guidelines and

were approved by the Human Ethics sub-Committee of the City University of Hong

Kong (jcc2223ay001). And all experimental procedures were approved by the

Animal Ethics Committee of the City University of Hong Kong and performed in

accordance with LARU Guidelines.

Consent for publication: Not applicable.

Availability of data and materials: All study data are included in the article and/or

Supplemantary files.

Competing interests: The authors declare no competing interest.

Funding: This work is supported by the National Natural Science Foundation of

China (82371350).

Authors' contributions: Conceptualization: GY, YXL; Methodology: XM, JZ, QLJ;

Investigation: XM, JZ; Visualization: XM, JZ; Supervision: GY, YXL; Writing: XM, JZ,

YXL, GY.

Acknowledgements: Not applicable.

Reference

1. Sospedra M, Martin R. Immunology of multiple sclerosis. Annu Rev Immunol.

2005;23:683-747.

2. Walton C, King R, Rechtman L, Kaye W, Leray E, Marrie RA, et al. Rising

prevalence of multiple sclerosis worldwide: Insights from the Atlas of MS. Multiple

Sclerosis Journal. 2020;26(14):1816-21.

3. Alroughani R, Boyko A. Pediatric multiple sclerosis: a review. BMC neurology.

2018;18:1-8.

4. Lee YK, Menezes JS, Umesaki Y, Mazmanian SK. Proinflammatory T-cell

responses to gut microbiota promote experimental autoimmune encephalomyelitis.

Proceedings of the National Academy of Sciences. 2011;108(supplement_1):4615-

22.

5. Cekanaviciute E, Yoo BB, Runia TF, Debelius JW, Singh S, Nelson CA, et al.

Gut bacteria from multiple sclerosis patients modulate human T cells and

exacerbate symptoms in mouse models. Proceedings of the National Academy of

Sciences. 2017;114(40):10713-8.

6. Chen J, Chia N, Kalari KR, Yao JZ, Novotna M, Paz Soldan MM, et al. Multiple

sclerosis patients have a distinct gut microbiota compared to healthy controls.

Scientific reports. 2016;6(1):28484.

7.   Jangi S, Gandhi R, Cox LM, Li N, Von Glehn F, Yan R, et al. Alterations of the

human gut microbiome in multiple sclerosis. Nature communications.

2016;7(1):12015.

8.   Johanson DM, Goertz JE, Marin IA, Costello J, Overall CC, Gaultier A.

Experimental autoimmune encephalomyelitis is associated with changes of the

microbiota composition in the gastrointestinal tract. Scientific reports.

2020;10(1):15183.

9.   Rojas M, Restrepo-Jiménez P, Monsalve DM, Pacheco Y, Acosta-Ampudia Y,

Ramírez-Santana C, et al. Molecular mimicry and autoimmunity. Journal of

autoimmunity. 2018;95:100-23.

10. Greiling TM, Dehner C, Chen X, Hughes K, Iñiguez AJ, Boccitto M, et al.

Commensal orthologs of the human autoantigen Ro60 as triggers of autoimmunity

in lupus. Science Translational Medicine. 2018;10(434):eaan2306.

11. Naghavian R, Faigle W, Oldrati P, Wang J, Toussaint NC, Qiu Y, et al. Microbial

peptides activate tumour-infiltrating lymphocytes in glioblastoma. Nature. 2023:1-11.

12. Kalaora S, Nagler A, Nejman D, Alon M, Barbolin C, Barnea E, et al.

Identification of bacteria-derived HLA-bound peptides in melanoma. Nature.

2021;592(7852):138-43.

13. Gil-Cruz C, Perez-Shibayama C, De Martin A, Ronchi F, Van Der Borght K,

Niederer R, et al. Microbiota-derived peptide mimics drive lethal inflammatory

cardiomyopathy. Science. 2019;366(6467):881-6.

14. Harkiolaki M, Holmes SL, Svendsen P, Gregersen JW, Jensen LT, McMahon R,

et al. T cell-mediated autoimmune disease due to low-affinity crossreactivity to

common microbial peptides. Immunity. 2009;30(3):348-57.

15. Wucherpfennig KW, Strominger JL. Molecular mimicry in T cell-mediated

autoimmunity: viral peptides activate human T cell clones specific for myelin basic

protein. Cell. 1995;80(5):695-705.

16. Planas R, Santos R, Tomas-Ojer P, Cruciani C, Lutterotti A, Faigle W, et al.

GDP-l-fucose synthase is a CD4+ T cell–specific autoantigen in DRB3* 02: 02

patients with multiple sclerosis. Science translational medicine.

2018;10(462):eaat4301.

17. Lerch M, Bauer A, Reindl M. The Potential pathogenicity of myelin

oligodendrocyte glycoprotein antibodies in the optic pathway. Journal of Neuro-

Ophthalmology. 2023;43(1):5.

18. Marignier R, Hacohen Y, Cobo-Calvo A, Pröbstel A-K, Aktas O, Alexopoulos H,

et al. Myelin-oligodendrocyte glycoprotein antibody-associated disease. The Lancet

Neurology. 2021;20(9):762-72.

19. Reindl M, Waters P. Myelin oligodendrocyte glycoprotein antibodies in

neurological disease. Nature Reviews Neurology. 2019;15(2):89-102.

20. Petersen TR, Bettelli E, Sidney J, Sette A, Kuchroo V, Backstrom BT.

Characterization of MHC- and TCR-binding residues of the myelin oligodendrocyte

glycoprotein 38-51 peptide. Eur J Immunol. 2004;34(1):165-73.

21. Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, et al. Extensive

Unexplored Human Microbiome Diversity Revealed by Over 150,000 Genomes from

Metagenomes Spanning Age, Geography, and Lifestyle. Cell. 2019;176(3):649-62

e20.

22. Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, et al. A

unified catalog of 204,938 reference genomes from the human gut microbiome.

Nature biotechnology. 2021;39(1):105-14.

23. Kishikawa T, Ogawa K, Motooka D, Hosokawa A, Kinoshita M, Suzuki K, et al. A

metagenome-wide association study of gut microbiome in patients with multiple

sclerosis revealed novel disease pathology. Frontiers in cellular and infection

microbiology. 2020;10:585973.

24. Zhou X, Baumann R, Gao X, Mendoza M, Singh S, Sand IK, et al. Gut

microbiome of multiple sclerosis patients and paired household healthy controls

reveal associations with disease risk and course. Cell. 2022;185(19):3467-86. e16.

25. Ishina IA, Zakharova MY, Kurbatskaia IN, Mamedov AE, Belogurov AA, Jr.,

Gabibov AG. MHC Class II Presentation in Autoimmunity. Cells. 2023;12(2).

26. Pishesha N, Harmand TJ, Ploegh HL. A guide to antigen processing and

presentation. Nat Rev Immunol. 2022;22(12):751-64.

27. Motmaen A, Dauparas J, Baek M, Abedi MH, Baker D, Bradley P. Peptide-

binding specificity prediction using fine-tuned protein structure prediction networks.

Proc Natl Acad Sci U S A. 2023;120(9):e2216697120.

28. Bradley P. Structure-based prediction of T cell receptor:peptide-MHC

interactions. Elife. 2023;12.

29. Marzella DF, Parizi FM, van Tilborg D, Renaud N, Sybrandi D, Buzatu R, et al.

PANDORA: A Fast, Anchor-Restrained Modelling Protocol for Peptide: MHC

Complexes. Front Immunol. 2022;13:878762.

30. Reynisson B, Barra C, Kaabinejadian S, Hildebrand WH, Peters B, Nielsen M.

Improved Prediction of MHC II Antigen Presentation through Integration and Motif

Deconvolution of Mass Spectrometry MHC Eluted Ligand Data. J Proteome Res.

2020;19(6):2304-15.

31. Lourbopoulos A, Deraos G, Matsoukas MT, Touloumi O, Giannakopoulou A,

Kalbacher H, et al. Cyclic MOG(35)(-)(55) ameliorates clinical and neuropathological

features of experimental autoimmune encephalomyelitis. Bioorg Med Chem.

2017;25(15):4163-74.

32. Berer K, Gerdes LA, Cekanaviciute E, Jia X, Xiao L, Xia Z, et al. Gut microbiota

from multiple sclerosis patients enables spontaneous autoimmune

encephalomyelitis in mice. Proceedings of the National Academy of Sciences.

2017;114(40):10719-24.

33. Berer K, Mues M, Koutrolos M, Rasbi ZA, Boziki M, Johner C, et al. Commensal

microbiota and myelin autoantigen cooperate to trigger autoimmune demyelination.

Nature. 2011;479(7374):538-41.

34. Ochoa-Repáraz J, Mielcarz DW, Ditrio LE, Burroughs AR, Foureau DM, Haque-

Begum S, et al. Role of gut commensal microflora in the development of

experimental autoimmune encephalomyelitis. The Journal of Immunology.

2009;183(10):6041-50.

35. Ochoa-Repáraz J, Kirby TO, Kasper LH. The gut microbiome and multiple

sclerosis. Cold Spring Harbor perspectives in medicine. 2018;8(6).

36. Ruff WE, Dehner C, Kim WJ, Pagovich O, Aguiar CL, Andrew TY, et al.

Pathogenic autoreactive T and B cells cross-react with mimotopes expressed by a

common human gut commensal to trigger autoimmunity. Cell host & microbe.

2019;26(1):100-13. e8.

37. Girdhar K, Huang Q, Chow I-T, Vatanen T, Brady C, Raisingani A, et al. A gut

microbial peptide and molecular mimicry in the pathogenesis of type 1 diabetes.

Proceedings of the National Academy of Sciences. 2022;119(31):e2120028119.

38. Cani PD, Depommier C, Derrien M, Everard A, de Vos WM. Akkermansia

muciniphila: paradigm for next-generation beneficial microorganisms. Nature

Reviews Gastroenterology & Hepatology. 2022;19(10):625-37.

39. Guo X, Zhang J, Wu F, Zhang M, Yi M, Peng Y. Different subtype strains of

Akkermansia muciniphila abundantly colonize in southern China. Journal of applied

microbiology. 2016;120(2):452-9.

40. Greer JM. Autoimmune T-cell reactivity to myelin proteolipids and glycolipids in

multiple sclerosis. Multiple sclerosis international. 2013;2013.

41. Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, et al. A

unified catalog of 204,938 reference genomes from the human gut microbiome. Nat

Biotechnol. 2021;39(1):105-14.

42. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk v2: memory

friendly classification with the genome taxonomy database. Bioinformatics.

2022;38(23):5315-6.

43. Kishikawa T, Ogawa K, Motooka D, Hosokawa A, Kinoshita M, Suzuki K, et al. A

Metagenome-Wide Association Study of Gut Microbiome in Patients With Multiple

Sclerosis Revealed Novel Disease Pathology. Front Cell Infect Microbiol.

2020;10:585973.

44. Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics. 2015;31(10):1674-6.

45. von Meijenfeldt FAB, Arkhipova K, Cambuy DD, Coutinho FH, Dutilh BE. Robust

taxonomic classification of uncharted microbial sequences and bins with CAT and

BAT. Genome Biol. 2019;20(1):217.

46. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P.

GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based

taxonomy. Nucleic Acids Res. 2022;50(D1):D785-D94.

47. Santos-Aberturas J, Chandra G, Frattaruolo L, Lacret R, Pham TH, Vior NM, et

al. Uncovering the unexplored diversity of thioamidated ribosomal peptides in

Actinobacteria using the RiPPER genome mining tool. Nucleic Acids Res.

2019;47(9):4624-37.

48. Mikhaylov V, Levine AJ. Accurate modeling of peptideMHC structures with

AlphaFold. 2023.

49. La Gruta NL, Gras S, Daley SR, Thomas PG, Rossjohn J. Understanding the

drivers of MHC restriction of T cell receptors. Nat Rev Immunol. 2018;18(7):467-78.

50. Shahi SK, Freedman SN, Dahl RA, Karandikar NJ, Mangalam AK. Scoring

disease in an animal model of multiple sclerosis using a novel infrared-based

automated activity-monitoring system. Scientific reports. 2019;9(1):19194.

51. Quan M-Y, Song X-J, Liu H-J, Deng X-H, Hou H-Q, Chen L-P, et al. Amlexanox

attenuates experimental autoimmune encephalomyelitis by inhibiting dendritic cell

maturation and reprogramming effector and regulatory T cell responses. Journal of

neuroinflammation. 2019;16:1-17.

Figure legends

Figure 1. Overview of MOG$_{35-55}$ analogs identified in the human microbiome.

(A) The workflow of human microbiome-derived peptides discovery. (B) The

proportion of MOG$_{35-55}$ analogs with well-defined binding motifs. (C) Chord diagram

showing the taxonomy distribution of prioritized analogs. The values in brackets

show the total number of analogs with corresponding binding motifs. per genome in

each phylum. The counts of analogs in each phylum were normalized by the

genome counts. The scale was the proportion of analogs in each motif (upper) or in

each phylum. Of note, the genomes less than 10 counts from *Firmicutes_I*,

Firmicutes_B, *Bacillota_B*, *Cyanobacteria*, *Desulfobacterota_I*, *Fusobacteriota*,

Pseudomonadota, *Euryarchaeota*, *Halobacterota*, *Patescibacteria*, and 19

unclassified contigs were merged into Others.

Figure 2. The confidence of hMOG$_{35\text{-}55}$ binding core peptide-MHC II and

peptide-MHC II-TCR complexes predicted TCRDock.

The scatter plots show the pLDDT value of a peptide versus an averaged inter-chain

PAE of peptide-MHC-II complexes (A) and peptide-MHC II-TCR complexes (B) for

binding core peptide candidates chopped from human MOG$_{35\text{-}55}$. The predicted

MHC II binding peptides (pLDDT ≥ 90, inter-PAE ≤ 4.34) are highlighted in red. The

sequence of binding core peptide candidates is listed on the right side of panel A.

The modeling of a peptide with pLDDT≥ 70 is considered as confident.

Figure 3. Polymorphism in 9-mer binding core peptides of the predicted

peptide-MHC II complex.

(A) The stacked bar plot shows the number of nonredundant ligands with 11 amino

acids residues truncated from analogs, colored with different motifs. (B) The scatter

plot shows the pLDDT value of a binding core peptide (corresponding to local

structural accuracy) versus an averaged MHC II-peptide complex's inter-chain PAE

score (corresponding to the peptide-MHC II interactions). The predicted binder is

highlighted in red. (C) The frequencies of amino acid residues (rows) at specific

positions (columns) among the predicted MHC II binding peptides. The loc_1 ~

loc_9 represent the binding core region among the binder, and loc_0 and loc_10

represent the flanking region of binders. (D) Representative sequence logos

illustrating peptide residue preference of MHC II. Each logo is labeled with the

starting amino acid residue of binding core peptides, and the number of binding core

peptides used to generate the logos is paired to the corresponding cells in panel [B].

The logos are colored by the chemical polarity of the amino acid residue. In panels

C and D, sequence/residue numbers are 0-indexed: they start at 0 and end with N-1,

where N is the length of the sequence.

Figure 4. Microbiome-derived peptide can stimulate MOG$_{35-55}$-specific T cells

from EAE mice.

(A) The origin and sequence of microbial peptides tested in this study. The amino

acids highlighted in green are the dominant MHC II-binding residue, while major

TCR contact residues are colored in red. The amino acids colored in blue represent

different residue between human MOG$_{35-55}$ and mouse MOG$_{35-55}$. The asterisk

highlights that P5 is a peptide predicted in both *Alistipes putredinis* and *A.*

muciniphila. (B, C and D) MOG$_{35-55}$–specific CD4$^+$ T cells collected from the spleen

of EAE mice were treated with each of the selected 8 microbial peptides. PBS and

MOG$_{35-55}$ served as negative control and positive control; Irrelevant peptide was set

as scrambled peptide control. Flow cytometry was used to determine IFN-γ and IL-

17 production (n = 3). Representative FACS plots (gated on CD3$^+$ and CD4$^+$ cells)

for the population of IFN-γ- or IL-17-secreted CD4$^+$ T cells that were stimulated with

mimic peptides or MOG$_{35-55}$ (20 µg/ml final concentration). Only data from P3, which

induced the highest cytokine-producing cell population was shown (B), along with

the frequency (C) and cell counts (D) of each population among CD4$^+$ T cells (Red

triangle specifically point P3 treatment group). Data shown are the average ± SEM.

Figure 5. P3 immunization exacerbates EAE development *in vivo*.

(A) Mean EAE clinical scores and incidence rate from each group of mice. (B)

Spinal cord samples were collected at day 17 post-immunization and the leukocyte

population within the CNS was assessed by flow cytometry. The cell count of

leukocytes were significantly higher in MOG$_{35-55}$+P3 group than MOG$_{35-55}$+control

group (n = 8). (C, D) Spinal cord tissue sections were obtained at day 17 post-

immunization and stained with H&E (C) or LFB (D). The inflammatory level (H&E

scores) and demyelination level (LFB scores) were quantified (n=4-5). Data shown

are the average ± SEM. *$p<0.05$*.

Figure 6. P3 shows bidirectional cross-reactivity with MOG$_{35-55}$.

(A, B and C) Splenocytes from P3 immunized mice (n=5) were stimulated with either

P3 (positive control), MOG$_{35-55}$, or PBS (negative control) in a concentration of 40

µg/ml to detect the cross-reactivity between P3 and MOG$_{35-55}$, and activation of

CD4$^+$ T cells was assessed by measuring the IL-17- and IFN-γ-producing population

using flow cytometry. Representative FACS plots (gated on CD3$^+$ and CD4$^+$ cells)

demonstrate the population of activated CD4$^+$ T cells (A), along with the frequency

(B) and cell counts (C) of IL-17$^+$ and INF-γ$^+$ population among CD4$^+$ T cells. Data

shown are the average ± SEM. *$p<0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 7. P3 stimulates 2D2 CD4$^+$ T cell proliferation through DCs presentation

in vitro.

(A) Predicted structure of peptide-MHC II-TCR complex with binding core peptide

from P3. TCR alleles same from complex template 3c60 (PDB ID). Surface

representation is displayed in lighter shades. (B) Close-up view of peptide-TCR

interactions (upper) and peptide-MHC II (bottom) in panel D, colored amino acid

residues represent the dominant MHC II-binding residue (green) and major TCR

recognition residue (red). Peptide residues and TCR alpha CDR3 (orange) and TCR

beta CDR3 (hot pink) residues were shown in sticks, and residues from MHC II

interacting with peptide were shown in slate blue. Hydrogen bonds are highlighted in

black dashed lines. (C, D) The histogram FACS plots (C) and quantification (D) of

CFSE labeled 2D2 CD4$^+$ T cells proliferation, as well as the frequency of CD69$^+$

cells within the CD4$^+$ T cells population, following co-culture with DCs pulsed with

PBS, P3, or MOG$_{35\text{-}55}$. Dash line in histogram represents the proliferation level of

PBS control. Data shown are the average ± SEM. **$p < 0.01$, ****$p < 0.0001$.

Figure 1

Figure 2

Figure 3

Figure 4

A

No	Taxonomy	Sequence
No	*Homo sapiens* MOG$_{35-55}$	MEVGWYRPPFSRVVHLYRNGK
	Mus musculus MOG$_{35-55}$	MEVGWYRSPFSRVVHLYRNGK
P1	*Bacteroides* MBL fold metallo-hydrolase	PIIYFYRSAFDRVMDGVKNSK
P2	*Alistipes sp902388705*	MDVNTYRPVFLRVPMDMILK
P3	*Akkermansia muciniphila*	TTLSFYRPPFLRVRRPFYIIF
P4	*Acutalibacteraceae_UMGS1623 sp934647945*	TFVQDYRYPFLRVISLLALAA
P5	*Alistipes putredinis, Akkermansia muciniphila*	AIDWFYRRPFSRVISETVFRY*
P6	*Bacteroides caccae*	LLLLWYRKIFVRVVWLGRTRH
P7	*Angelakisella*	EELEQYRPLFGRVIIRIRTPS
P8	*Pararuminococcus gallinarum*	KLKIKYREGFFRVVHAIRDFF

B

C

D

Figure 5

Figure 6

Figure 7

A

TCR beta
TCR alpha
Peptide 3
MHCII

B

Peptide-TCR

TCR alpha CDR3
TCR beta CDR3
Arg
Pro
Peptide 3

P3: TTLSFYRPPFLRVRRPFYIIF

Peptide-MHC II

MHCII
Thr
Leu
Ser
Phe
Tyr
Arg
Pro
Pro
Phe
Leu
Arg
Peptide 3

C

PBS   MOG₃₅₋₅₅   P3

Relative cells number

5.89   94.0   29.9

CFSE

D

% Proliferation

****
****

PBS   MOG₃₅₋₅₅   P3

% of CD69⁺ cells in CD4⁺ T cells

**
**

PBS   MOG₃₅₋₅₅   P3