Title: Novel quantal analysis method reveals conservation of injected charge by excitatory synapses

across cortical cells of different sizes

Author Names and Affiliations: Martynas Dervinis[1], Guy Major[1]

[1]: Cardiff University, Cardiff, Wales, CF10 3AT, UK

Corresponding author email address: martynas.dervinis@gmail.com

Number of pages: 39

Number of figures: 9

Number of words

Abstract: 252

Body text excluding Materials and Methods: 3061

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: Cardiff University (GM, MD), The Medical Research Council of the UK (MD

PhD). Neural activity simulations were performed on the Cardiff School of Biosciences' Biocomputing

Hub HPC/Cloud infrastructure.

Abstract

In chemical synapses of the central nervous system (CNS), information is transmitted via the presynaptic release of a vesicle (or 'quantum') of neurotransmitter, which elicits a postsynaptic electrical response with an amplitude termed the 'quantal size'. This key determinant of neural computation is hard to measure reliably due to its small size and the multiple sources of noise within neurons and electrophysiological recordings. Measuring amplitudes of miniature postsynaptic currents (mPSCs) or potentials (mPSPs) at the cell soma potentially offers a technically straightforward way to estimate quantal sizes, as each of these miniature responses (or 'minis') is elicited by the spontaneous release of a single neurotransmitter vesicle. However, a somatically recorded mini is typically massively attenuated compared with at its input site, and a significant fraction are indistinguishable from (or cancelled out by) background noise fluctuations. Here we describe a novel quantal analysis method that estimates electrical size of the synapse by combining separate somatic recordings of background physiological noise and minis with simulations. With the help of a genetic algorithm, simulations are successfully used to infer the combined amplitude and rise time distribution of minis that would otherwise be inaccessible due to low signal to noise ratio. The estimated distributions reveal a striking inverse dependence of mean minis' amplitudes on cell's total capacitance (a proxy for the size of cells) that firmly supports the conservation of the electrical size of excitatory cortical synapses in rats. We incorporate the novel quantal analysis method into our patch clamp data analysis software called 'minis'.

Abstract

Abbreviations

ABF – Axon Binary File

aCSF – artificial cerebrospinal fluid

AMPA – α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid

ANN – artificial neural network

AP – action potential

CPP – (*RS*)-3-(2-carboxypiperazin-4-yl)-propyl-1-phosphonic acid

EPSC – excitatory postsynaptic current

EPSP – excitatory postsynaptic potential

GA – genetic algorithm

GABA – gamma-aminobutyric acid

IPSC – inhibitory post-synaptic current

IPSP – inhibitory post-synaptic potential

MAD – maximal absolute deviation

mEPSC– miniature excitatory post-synaptic current

mEPSP– miniature excitatory post-synaptic potential

mIPSP– miniature inhibitory post-synaptic potential

mPSC – miniature post-synaptic current

mPSP – miniature post-synaptic potential

NBQX – 2,3-dioxo-6-nitro-1,2,3,4-tetrahydrobenzo[*f*]quinoxaline-7-sulfonamide

NMDA – N-methyl-D-aspartate

SAD – sum of absolute deviations

SD – standard deviation

smEPSP – simulated mEPSP

TTX – tetrodotoxin

V_m – membrane potential

Introduction

Synaptic transmission in the CNS occurs in 'quanta' (Forti et al., 1997; Hardingham et al., 2010; Isaacson and Walmsley, 1995; Jonas et al., 1993; Kraszewski and Grantyn, 1992; Kullmann and Nicoll, 1992; Kuno, 1964; Larkman et al., 1991; Sahara and Takahashi, 2001; Stern et al., 1992; Wall and Usowicz, 1998) usually thought to correspond to the release of a single neurotransmitter vesicle (Heuser et al., 1979). The amplitude of the postsynaptic response to a quantum of transmitter, the quantal size, is a key parameter defining the efficacy of synaptic transmission (Katz, 1969; Vere-Jones, 1966) and, therefore, an important determinant of neural computation.

Quantal size measurements in the CNS are difficult due to multi-synaptic contacts, limited accessibility, significant quantal variance, temporal non-stationarity of synaptic properties, and low signal-to-noise ratios of somatic electrophysiological recordings of quantal events (Bekkers, 1994; Korn and Faber, 1991). For these reasons classical quantal analysis as done at the neuromuscular junction has run into problems in central synapses. More recent methods such as noise deconvolution, Variance-Mean analysis, and optical quantal analysis have not fared much better either (Biró et al., 2005; Clamann et al., 1991; Clements, 2003; Clements and Silver, 2000; Edwards et al., 1976; Enoki et al., 2009; Jack et al., 1981; Liu et al., 2022; Oertner et al., 2002; Peled and Isacoff, 2011; Redman, 1990; Reid and Clements, 1999; Silver, 2003; Yuste et al., 1999).

A complementary method of assessing synaptic function is to measure properties of mPSCs. Miniature PSC amplitude measurements are often taken as quantal size estimates across the population of active synapses onto a neuron, as the mPSC (under voltage clamp) or mPSP (under voltage recording) likely correspond to the response to a spontaneous release of a single neurotransmitter vesicle (Brown et al., 1979; del Castillo and Katz, 1954; Fatt and Katz, 1952; Isaacson and Walmsley, 1995; Wall and Usowicz, 1998). Measurements of mPSC amplitudes are

routine in the literature. However, they are likely overestimates due to small amplitude events being

difficult to detect and largely being missed.

The use of voltage clamp can be a major reason why mPSCs "go missing". Theoretical studies (Major,

1993; Rall and Segev, 1985; Spruston et al., 1993) predicted and experimental findings (Williams and

Mitchell, 2008) confirmed that voltage clamp fails to clamp extended dendrites, and to capture the

current flowing in at electrically remote synapses. Amplitude attenuation can also be caused if series

resistance is not compensated (Williams and Mitchell, 2008). These issues can be somewhat

sidestepped by switching to voltage recording.

However, most mPSCs go undetected because of the electrophysiology of the neuron itself. Distal

minis leak charge via the dendritic membrane, but an even more significant amplitude attenuation,

40-fold or more, is inflicted by charging the membrane capacitance of dendritic trees (Larkum et al.,

2009; Major et al., 2013a; Nevian et al., 2007; Stuart and Spruston, 1998; Williams and Mitchell,

2008; Williams and Stuart, 2002). Such amplitudes, at the soma, are often indistinguishable from

noise fluctuations. Recording from thin dendrites close to synaptic input sites is technically

challenging (Nevian et al. 2007). Therefore, a reliable yet widely accessible quantal analysis method,

based on somatic whole-cell patch recording, would require taking account of the background noise

component in addition to measuring putative mPSPs (or mPSCs).

Here we describe such a method based on separate recordings of excitatory mPSPs (mEPSPs) in the

presence of background physiological noise ('noise + minis' recording) and of noise alone after

blocking mEPSPs pharmacologically ('noise-alone' recording). To detect mEPSP-like events in these

recordings we used a newly developed algorithm called 'minis' which is described in our companion

article (Dervinis and Major, 2023). Because of larger minis overshadowing smaller potentials, events

detected in the 'noise-alone' recording cannot be used to account for the noise component in the

'noise + minis' recording. Instead, simulated minis must be used by superimposing them on the

'noise-alone' recording until the 'noise + simulated minis' voltage trace closely matches the 'noise +

real minis' recording. Therefore, we augmented 'minis' software with a genetic optimisation

algorithm (GA) that automatically adjusts amplitude and rise times of a simulated minis. The novel

quantal analysis method produced an unbiased quantal size estimate for excitatory synapses in the

CNS which was approximately equal to 28.53 pC and was conserved across pyramidal cells of all sizes

in cortical layers 2/3 and 5. 'Minis' thus provides a straightforward quantal analysis method of

detecting mPSPs and mPSCs in somatic recordings and separating smaller of these events from

background noise fluctuations.

Materials and Methods

Animals and electrophysiology

All experimental procedures were carried out in accordance with the UK Animals (Scientific

Procedures) Act 1986 at the Cardiff University under Home Office personal and project licenses. The

full description of electrophysiology procedures is provided in our companion article (Dervinis and

Major, 2023). Briefly, coronal somatosensory cortical slices obtained from 19- to 27-day old Wistar

rats (RRID:RGD_13508588; n = 11) of either sex were used to obtain whole-cell recordings (n = 14) of

mEPSPs in the background of physiological noise ('noise + minis' condition) by applying artificial

cerebrospinal fluid (aCSF) solution added with an action potential (AP) and inhibitory postsynaptic

potential (IPSP) blockers. Following this phase, the recordings were continued further after adding

excitatory postsynaptic potential (EPSP) blockers ('noise-alone' condition).

Passive membrane time constant, total capacitance, and cell's input resistance (the bridge balance

error) were estimated by injecting current pulses as described in (Major G et al., 1993).

Experimental design

The goal of the study was to design a quantal analysis method based on somatic recordings of minis.

Due to low signal to noise ratio, simulations of minis were used to infer the signal component and to

eliminate the noise component in these recordings.

The experimental design of this study was similar to the one described in our companion article

(Dervinis and Major, 2023). As described in the previous section, two types of whole-cell recordings

were obtained where mEPSPs were recorded in the background of physiological noise or where

mEPSPs were pharmacologically blocked and only noise was recorded. The noise recordings were

then superimposed with simulated mEPSPs (smEPSPs) and 'minis' algorithm was applied to this

'noise + simulated minis' voltage trace. The resulting distribution of detected events in terms of their

amplitudes and 10-90% rise times was compared to the equivalent distribution based on 'noise +

real minis' recording. Discrepancy between the two distributions was used to inform the GA which

would then adjust the distribution of smEPSPs and generate new 'noise + simulated minis' voltage

traces and compare them again to the equivalent distribution based on 'noise + real minis'

recording. This procedure was repeated multiple times until the two distributions were closely

matched. At the end of this process the minis' distribution used for simulating a closely matched

recording was deemed to be the inference of the true minis' distribution underlying the 'noise + real

minis' recording.

Detection of miniature postsynaptic potentials

Detection of real and simulated mEPSPs was carried out using 'minis' software. The detailed

description of this detection algorithm is provided in our companion article (Dervinis and Major,

2023). We used the same detection algorithm parameters as described in that article except for the

amplitude range which was set to 0.02-10 mV.

173 **Selection of recordings for fitting distributions of detected real minis-like events**

174 Prior to carrying out any estimation of minis' distribution, segments of 'noise + real minis' and

175 'noise-alone' recordings were carefully selected. Segments containing unusual amount of noise or V_m

176 glitches were rejected. Parts of recording showing instability or drift of recording parameters were

177 also rejected to ensure that only good quality data was used. We called this a preprocessing stage

178 and data analyses we performed at this stage were inbuilt into 'minis' software.

179

180 As part of the preprocessing stage 'minis' produced 7 graphs showing the evolution of various

181 recording parameters across the entire duration of a single recording. An example is provided in

182 Figure 1 showing these graphs as individual panels of the figure for the longest obtained recording.

183 Figure 1A shows the evolution of V_m from the very beginning of the recording obtained in the regular

184 aCSF with APs present to the very end of the recording where APs as well as mEPSPs and mIPSPs

185 were blocked for some time ('noise-alone' recording phase). The data is shown after removing V_m

186 segments containing pulses with the total duration of 8519.9 s. Throughout the entirety of this

187 recording the baseline V_m varied between −70 and −67 mV and, thus remained relatively stable.

188

189 There are 3 time markers that extend across all figure panels marking the time when the aSCF

190 containing TTX and gabazine was infused to block APs and IPSPs (the first marker), the time when

191 APs were observed to cease (the second marker), and the time when the aCSF containing NBQX and

192 CPP was infused to block remaining EPSPs (the third marker). The recording segment between the

193 second and the third markers constituted the 'noise + minis' recording, while the segment following

194 the third marker when mEPSPs were eventually blocked constituted the 'noise-alone' recording. The

195 shaded regions that also extend across all panels of Figure 1 mark files that were selected for

196 subsequent minis' distribution fitting analyses. In this example files 27 to 88 and files 100 to 104

197 were selected for further analyses (10 recording sweeps per each file).

198

The remaining panels of Figure 1 were used to guide the data selection for subsequent analyses. The

standard deviation (SD) of the V_m (Figure 1B) was useful in determining the borders of different

recording types. Stable prolonged decreases in SD reliably indicated when APs or minis were

effectively blocked by corresponding pharmacological agents. Sudden brief increases in SD flagged

problematic recording segments containing higher levels of noise or V_m glitches. Gradual increases in

SD suggested the ensuing gradual deterioration of the recording quality.

Similar information was also provided by the evolution of amplitudes of the top 10% (amplitude-

wise) of all detected minis-like events in a single sweep (Figure 1C). Prolonged drop in the

amplitudes marked borders between recording types, while sudden jumps were indicative of V_m

instabilities. Finally, effective decay time constants of the same top 10% (amplitude-wise) of

detected events (Figure 1D) could be used as an indicator for the presence/absence of minis.

Blocking minis resulted in time constants becoming large indicating that only noise fluctuations were

being detected that did not possess typical decay dynamics common to PSPs.

Further insights into the recording quality over time were provided by impulse based measures.

Passive membrane time constant measurements over time (Figure 1E) were expected to remain

stable. Deviations from this expectation could signal a change in passive membrane properties and

the degradation in the recording quality. Total membrane capacitance (Figure 1F) was also expected

to be stable over time. Change in the capacitance could indicate tear of the membrane. Finally, input

resistance (the bridge balance error; Figure 1G) was not permitted to deviate by more than twice its

estimate obtained at the beginning of the recording. Violation of this rule of thumb indicated a

deterioration in the recording quality potentially resulting from debris accumulating in the recording

pipette or other reasons.

We obtained the above-discussed 7 longitudinal recording state/quality indicators for all our

recordings. Multipaneled graphs showing these indicators for each individual recording separately

are provided in Supplementary Figures 1-13.

Simulations

Simulating distributions of miniature excitatory postsynaptic potentials with a range of amplitudes

and rise times

All simulations were carried out using 'minis' software developed in Matlab (Mathworks;

RRID:SCR_001622) and performed on Cardiff School of Biosciences' Biocomputing Hub HPC/Cloud

infrastructure. Simulation of minis-like events was based on an analytical solution to the passive

cable equation for a transient current (Rall, 1977) using 100 lumped terms with a double exponential

synaptic current with τ_1= 0.15-3 ms (optimised) and τ_2 = 2 ms (Major et al., 1993). The detailed

description of how arbitrary shape PSPs were obtained is provided in our companion article (Dervinis

and Major, 2023).

A pool of simulated events was generated combining 4 bivariate (amplitude and 10-90% rise time)

normal distributions. 6 parameters described each distribution: amplitude mean μ_X and standard

deviation σ_X, 10-90% rise time mean μ_Y and standard deviation σ_Y, a correlation factor ρ, and a scale

factor n:

243
$$f(x,y) = \frac{n}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} e^{-\frac{1}{2(1-\rho^2)}\left(\left(\frac{x-\mu_X}{\sigma_X}\right)^2 - 2\rho\left(\frac{x-\mu_X}{\sigma_X}\right)\left(\frac{y-\mu_Y}{\sigma_Y}\right) + \left(\frac{y-\mu_Y}{\sigma_Y}\right)^2\right)}.$$

All distribution parameters were controlled by the GA.

Fitting distributions of detected real minis-like events

Matlab's ga function was used to control 24 parameters describing 4 bivariate normal distributions

(6 paprameters per distribution). Additional controlled parameters were the synaptic rise time

constant τ_1 which ranged between 0.15 and 3 ms and the passive membrane time constant τ_m. The

range for the latter constant was taken to be between 0.95 times and 1.05 times the initial estimate

based on the average waveform of all minis-like events detected in the 'noise + real minis' recording.

Thus, there were 26 parameters in total controlled by the GA.

The GA would start by generating multiple distributions smEPSPs (i.e., 'individuals'; we used 260

individuals per 'generation'). Events would be drawn pseudo-randomly from a distribution and then

simulated events of the right shape (in terms of amplitude and 10-90% rise time) would be

superimposed on the 'noise-alone' recording at time points that were chosen pseudo-randomly

(Figure 2). For generating the initial set of individuals ('population'), the GA chose parameters

pseudo-randomly within prescribed ranges. For each of the four bivariate normal distributions the

parameter ranges were as follows: μ_X (-0.1-1 mV), σ_X (0.01-1 mV), μ_Y (-2.5-10 ms), σ_Y (0.25-10 ms), ρ

(-1 to 1), and n (0-10000 for the first distribution and –10000 to 10000 for the remaining three

distributions). Any negative count bins within the resulting final distribution were zeroed out.

In the next step the GA assessed the fitness of each individual in the generation by using the cost

function. The top 5% of the most fit individuals survived to the next generation without change while

individual crossover, genetic migration, and other GA parameters were set to default values of the

Matlab's ga function except for the mutation function type which was set to

'mutationadaptfeasible'. A new evolved generation of individuals was then created. Which then

served as the basis for yet another generation. This process was repeated until an individual with the

lowest cost (highest fitness value) possible was created, there was no improvement in the fitness for

100 consecutive generations, or the number of generations reached 1000.

In the case when the GA terminated without attaining the highest possible fitness value, distribution

fitting procedure was reinitiated using the most fit individual as a seed. The population genetic

diversity was reset by setting the GA-controlled parameter ranges to ±5% of the original ranges

relative to the seed parameters. The rerunning of distribution fitting was repeated until the GA

generated the most fit individual possible, an acceptable level of fitness was reached (defined in the

next subsection), or there was no improvement in fitness compared to the previous run.

If the most fit individual was within acceptable levels of fitness values, the lower threshold on

smEPSP amplitude was increased by 0.005 mV and the entire fitting procedure rerun from the

beginning till the end. This procedure was repeated until the GA stopped converging on a solution

with acceptable fitness values. The reason for pushing the lower limit of smEPSPs' amplitudes is that

GA-generated solutions are not unique. Matching 'noise + real minis' recording can be achieved by

using a distribution of fewer relatively large amplitude smEPSPs or more numerous smaller

amplitude smEPSPs. There is an infinite number of such distributions with ever decreasing

amplitudes of smEPSPs. Therefore, GA has no guaranteed way of arriving at the proven true

distribution of minis, only at an upper amplitude limit of this distribution by the above-described

thresholding procedure.

Cost function

The goal of the minis' distribution fitting procedure was to match 'noise + simulated minis' and

'noise + real minis' recordings as close as possible in terms of the distribution of amplitudes and 10-

90% rise times of detected minis-like events in the two types of V_m recording traces ('simulated' vs

'real' distributions). This was achieved by minimising the cost function using the GA. The cost

function was composed of three types of components: shape constraints on the simulated minis

distribution, sums of absolute deviations (SADs) between 'simulated' and 'real' distributions, and

maximal absolute deviations (MADs) between 'simulated' and 'real' distributions.

We calculated the typical ranges of SADs and MADs for each of the 14 recordings by taking the

individual 'noise + real minis' recording and dividing it into smaller files typically consisting of 5

sweeps each (to match the size of the segment of 'noise-alone' recording used in the distribution

fitting procedure). Each file in this set was subjected to minis detection procedure resulting in a set

of 'real' distributions. They were then subtracted from each other, and corresponding SAD and MAD

scores were calculated for each possible $n*(n-1)$ file combinations. 6 different SAD and MAD scores

were calculated based on: distributions of (1) amplitudes of all detected events, (2) amplitudes of

top 50% largest detected events, (3) amplitudes of top 10% largest detected events, (4) amplitudes

of top 2% largest detected events, (5) distribution of 10-90% rise times, and (6) a joint amplitude and

10-90% rise time distribution. The smallest percentile that guaranteed to include half of SAD or MAD

scores of all file combinations simultaneously for all 6 types of scores ('6-score combined 50th

centile') was used to determine which SAD or MAD values should be used for all 6 types of scores to

construct the cost function.

The cost function was constructed of 6 types of SAD scores and 6 types of MAD scores estimated

using the '6-score combined 50th centile' procedure. Additional 3 SAD scores were added to penalise

'simulated' amplitude distributions that had lower counts of high amplitude events (top 50%, 10%,

and 2%) compared to 'real' distributions. These 15 scores were used to evaluate how close the

'simulated' distribution was to a single selected 'real' distribution that was deemed to be of the best

quality by a human observer. Additional corresponding 15 'deviation' scores (same as SADs and

MADs but comparing to all distributions at once) were calculated assessing how much each 'real' or

'simulated' distribution deviated from the full set consisting of all 'real' distributions and the single

'simulated' distribution. The 'simulated' distribution was expected to have a lower deviation score

than the worst performing 'real' distribution. 'Simulated' distributions were further penalised if their

underlying 'simulated minis' distribution violated the 10-90% rise time skewness constraint which

stipulated that the mean 10-90% rise time of simulated minis should exceed the median. Finally,

'simulated' distributions were also penalised if their underlying 'simulated minis' amplitude

distribution had a sharp cliff at its lower end. Hence, there were 32 fitness components in total (15 +

15 + 1 + 1 = 32).

If all amplitude SAD, MAD, and deviation scores for comparing the 'simulated' distribution to the

closest 'real' distribution were satisfied at the end of the fitting procedure, the fit was deemed to be

'good enough'. Alternatively, the fit could also be deemed good enough if all amplitude and 10-90%

rise time SAD and MAD scores were satisfied (deviation scores excluded).

Statistical analyses

All inferential statistics were based on the assumption that distributions of all measures were

normal. All mean distribution values are stated/depicted with 95% confidence limits.

Data accessibility

All data analysed in this study are publicly available (GIN DOI will appear in version 2 of the

manuscript). The available data include 'noise + minis' and 'noise-alone' whole-cell patch membrane

potential recordings that were used in computer simulations and minis' distribution fitting. All

electrophysiological recordings were stored in Axon Binary File (ABF) format.

Code accessibility

All analyses were carried out in Matlab and the analysis code is publicly available on GitHub (DOI will

appear in version 2 of the manuscript). The code is complete with instructions on how to reproduce

all figures reported in this study.

Software accessibility

The present study reported the use of a novel quantal analysis method that is part of 'minis'

software available on GitHub (Dervinis, 2022).

Results

Estimating quantal size by direct subtraction of noise component

We have subjected 'noise + minis' recordings and 'noise-alone' recordings to the 'minis' detection

algorithm. Detected mEPSP-like events had a particular shape amplitude and 10-90% rise time

distributions that resembled log-normal or gamma distributions (Figure 3). In the amplitude domain

the lower end of the distribution contains a mixture of real minis and noise fluctuations that were

misidentified as real events.

We argued that if the noise component could be reliably estimated based on the 'noise-alone'

recording where synaptic activity has been pharmacologically blocked, it could be directly subtracted

from the 'noise + minis' distribution yielding an estimate of the pure minis' component. Figure 3A

shows both 'noise + minis' and 'noise-alone' detected event amplitude distributions superimposed

over each other. After subtracting the 'noise-alone' distribution from the 'noise + minis' distribution

there appear negative low amplitude bins in the resulting distribution (indicated by an arrowhead in

Figure 3A). These missing low amplitude events are caused by larger minis overshadowing smaller

minis and noise fluctuations in the 'noise + minis' recording (we have discussed this phenomenon in

great length in our companion article (Dervinis and Major, 2023)). Therefore, the direct subtraction

approach can only provide a high-end upper limit on the mean of the minis' amplitude distribution.

Nevertheless, this over-estimate is more reliable than a highly subjective approach of simply

estimating the mean amplitude based on events detected using a high amplitude threshold. Figure 4

shows the distribution of mean minis' amplitudes obtained using the direct noise subtraction

method. When mean amplitude estimates are compared to capacitance estimates (or

1/capacitance), there is a strong inverse correlation between the mean amplitude and cell's

capacitance (the proxy measure for cell's size; r = 0.754, p = 0.00275, R^2 = 0.568). That is, the larger

the cell, the smaller the amplitude of its somatically detected minis and vice-versa. This relationship

is predicted by the theory of capacitors (C = Q/V, where C stands for capacitance, Q stands for

charge, and V stands for voltage across the cell's membrane), and it is a direct demonstration

attenuating effect of cell's capacitance on the amplitude of somatically detected minis. If a direct

line was drawn through the data and the origin of the axes in Figure 4, the slope of this line would

provide a quantal size estimate for excitatory synapses in the CNS in terms of charge (Q = ΔV/

Δ(1/C)) which is approximately equal to 34.33 fC in this case.

Estimating quantal size by simulating excitatory postsynaptic potential source distribution

An alternative and more reliable way of estimating the quantal size in our recordings is to estimate

the minis component directly using simulations. This relies on 'minis' optimisation algorithm (GA)

being able to closely match a 'noise + simulated minis' voltage trace (Figure 2) with 'noise + real

minis' recordings in terms of amplitude and 10-90% rise time distributions of minis-like events

detected within these traces. We were able to successfully do that for all our recordings (see Figure

5 and Supplementary Figures 14-26) with one particular example shown in Figure 5. All simulations

produced 'good enough' matches showing amplitude and 10-90% rise time distributions of detected

minis-like events being within the range of amplitude and 10-90% rise time count values measured

in sets of 'real event' distributions. All resulting minis source distributions are displayed in Figure 6

and Supplementary Figures 27-39. An example in Figure 6 shows that the 'real minis' source

distribution likely has a log-normal or gamma-like shape in both amplitude and rise time domains.

Incidence rates estimated based on simulated minis source distributions ranged roughly between 10

and 100 minis/s (Figure 7). This is estimate is an order of magnitude higher than the one based on

the literature and significantly higher than the one based on the direct subtraction method (t(13) =

4.33, p = 0.0008) and also higher than the estimate based on amplitudes higher than a chosen 0.8

mV threshold (t(13) = 4.89, p = 0.0003). It is precisely within the range predicted in our companion

article (Dervinis and Major, 2023). However, this range is still somewhat of an underestimate as

minis source distributions were often generated using a lower limit on simulated amplitudes to

constrain the distribution non-uniqueness as discussed in the methods subsection on the

distribution fitting.

Mean minis' amplitudes derived using simulations ranged between 0.053 and 0.14 mV (Figure 8A).

Just like amplitudes derived by the noise subtraction method in Figure 4, amplitudes derived using

simulations had a strong association with the size of the cell represented by cell's capacitance

(Figure 8A). The effect size of this relationship was even stronger for simulated amplitudes

explaining 63% of the total variance (r = 0.793, p = 0.00115). The slope of the line fitted through the

data and passing the origin of axes in Figure 8A provided the estimate of quantal size in the cortical

excitatory synapses that was approximately equal to 28.53 fC and, as expected, was smaller than

that derived by the noise subtraction method. The two types of mean amplitude estimates derived

either by noise subtraction or by minis source distribution simulation were mutually correlated (r =

0.596, p = 0.0275, R^2 = 0.355, Figure 8B). The equation of the line (1.01x + 0.005) fitted to this data

provided a way to approximate the mean amplitude of fitted minis' source distribution by adding the

intercept term to the means derived by noise subtraction (no multiplication term as the slope of the

line was approximately equal to 1).

Generic miniature excitatory postsynaptic potential distribution in neocortex

To construct a generic mEPSP distribution, we normalised amplitudes of simulated minis' source

distributions shown in Figure 6 and Supplementary Figures 27-39 by dividing them by their

respective means and taking the total average. The resulting combined (amplitude and 10-90% rise

time) distribution is shown in Figure 9A and its smoothed version is shown in Figure 9B. The

corresponding marginal amplitude and 10-90% rise time distributions are shown in Figures 9C and E,

respectively, with their smoothed versions shown in Figures 9D and F. Both marginal distributions

have a log-normal or gamma-like distributions.

Discussion

In this article we reported two major scientific advances. First, we described a novel quantal analysis

method that is based on separate somatic recordings of mEPSPs in the background of physiological

noise and recordings of noise alone after pharmacological blockade of mEPSPs. The latter type of

recording provides a means to estimate the properties of the noise component which can then be

used to account for noise fluctuations in the combined minis and noise recording. However, this was

done not by a direct subtraction of the noise component but rather by the simulation of the

underlying minis component and repeatedly superimposing simulated minis shapes onto the noise

trace until the 'noise + simulated minis' voltage trace closely matched the 'noise + real minis'

recording. The key tool to achieving this match was the GA that optimised the parameters of the

simulated minis distribution in terms of amplitudes and 10-90% rise times.

The second major advance was the use case for the novel quantal analysis method that successfully

measured the electrical size of excitatory synapses of cortical pyramidal cells which was shown to be

approximately equal to 28.53 fC. This number was inferred from the slope of the line fitting minis'

mean amplitude and cell's total capacitance data in accordance with the theory of capacitors. The

fact that this line had an intercept at the origin of the axes indicated that there was no active

compensation for minis' amplitude attenuation due to cell's size (i.e., total capacitance). Indeed,

cell's capacitance played a major role affecting the amplitude of mEPSPs resulting in a large

proportion of mEPSPs being otherwise indistinguishable from noise. The number of these so-called

'missing' minis was so high that the estimated minis' incidence rate was in the order of 10-100

minis/s: A number that is larger by an order of magnitude than the one typically encountered in the

synaptic physiology literature. Finally, both the amplitude and 10-90% rise time distributions of minis

had a log-normal- or gamma-like shapes.

These findings have a few major implications. First, the method, as used for estimating the mEPSPs'

distributions in cortical layers 2/3 and 5 of cells of various size and then using these distributions to

arrive at the electrical size of excitatory synapses serves as a proof of principle that it can be

successfully used to estimate PSP distributions of various types in the CNS. The next logical step of

this method is to extend its application to the estimation of amplitude and rise time distributions of

mIPSPs generated by different types of inhibitory synapses and, as a result, the electrical sizes of

these different inhibitory synapses. Moreover, the ability to get relatively accurate quantal size

estimates unobscured by physiological noise is important for constraining synaptic parameters in

physiologically realistic simulations of neural activity. For example, now we should be able to

estimate how many active synapses are needed to generate an NMDA spike within a single dendritic

segment (Major et al., 2013b). The importance of these number estimates is not restricted to the

domain of physiologically realistic simulations but also extends to more abstract computations that

use NMDA spike-like mechanisms as coincidence detectors in latest cortically inspired models of

artificial intelligence (Hawkins et al., 2019). It is thus imperative to arrive at accurate estimates of

synaptic transmission parameters.

These findings also draw attention to methodological issues in the field of synaptic physiology. As

discussed in our companion article (Dervinis and Major, 2023), the commonly reported incidence

rates for mEPSCs are well below 10 minis/s. Even if one is in doubt of the incidence rates that we

arrived with the help of simulations that lie in the range of 10-100 minis/s, the range of incidence

rates given by direct subtraction of the distribution of events detected in the 'noise-alone' recording

from the distribution based on the 'noise + minis' recording is still within 5.9 and 26.7 minis/s which

we know to be an underestimate. Therefore, reported minis' incidence rates below 10 minis/s are

certainly underestimates. As discussed earlier, these estimates could be improved by switching to

voltage recording instead of using the voltage clamp (Major, 1993; Rall and Segev, 1985; Spruston et

al., 1993; Williams and Mitchell, 2008). But more importantly, a strong association between cell's

capacitance (or its size) and how much somatically recorded minis' amplitudes are attenuated

indicates that special care must be taken to account for the effect of background noise fluctuations

on the amplitude measurements as we have done here. This association is yet another evidence

supporting experimental observations of massive minis' amplitude attenuation as the PSP spreads

from its dendritic source location to the soma (Larkum et al., 2009; Major et al., 2013a; Nevian et al.,

2007; Stuart and Spruston, 1998; Williams and Mitchell, 2008; Williams and Stuart, 2002).

It is also important to note that understanding the link between cell's capacitance and somatically

recorded minis' average amplitudes as well as knowing actual shapes of minis' distributions in terms

of their amplitudes and rise times allows us to generate artificial data where ground truth data is

known. Such labelled data could be used to train artificial neural networks (ANNs) which could then

be used develop the next generation powerful quantal analysis methods allowing detection of minis

directly in mixed 'noise + minis' recordings. Such methods would not only do away with the need to

obtain relatively long voltage recordings with consecutive pharmacological phases but could

potentially fully automate the minis' detection process. We are not the first to suggest the

application of ANNs to this issue (see (Pircher et al., 2022) and (Wang et al., 2024)). However, we are

not aware of attempts to use realistic minis' distributions for this purpose.

Even though the novel quantal analysis method we present here is an important advancement, it

nevertheless has limitations. One of them is the need for long duration recordings using different

pharmacological agents at various stages of recording. However, the somatic recording set up is not

as complicated as an alternative multisite dendritic recording set up would be. Moreover, the

distribution fitting procedure involving the GA can be lengthy and computationally intensive.

However, 'minis' software is parallelised at the PSP detection level, as well as at the GA optimisation

level making it particularly suitable to be executed on multi-core computing cluster. Although 'minis'

software combined with the two-stage recording set up offers an unprecedented control potential

over the background physiological noise, we understand that many researchers may not be

persuaded to adopt it themselves. Therefore, the adoption of ANNs for the next generation of

'minis' software based on current findings and future use-cases offers a way to simplify the current

approach.

Some researchers may be critical of a method that does not rely on a direct measurement of minis

but rather on an approximation via indirect means using simulations. This is a valid criticism that

could be addressed in the future by using a method of repeatedly subjecting a voltage recording to

minis detection procedure with increasingly lower threshold while subtracting detected events from

the voltage recording. This approach potentially could recover most of the smaller minis that are

overshadowed by the larger ones (the so called 'negative bins' problem illustrated in Figure 3A).

Furthermore, other researchers may question the ability to pharmacologically isolate the

background physiological noise. Our 'noise-alone' recordings may contain random openings of

individual AMPA channels or may fail to fully eliminate channel openings in the presence of quantal

release of neurotransmitter. Physiological noise itself may be different in the two conditions with

the presence or absence of mEPSP pharmacological blockers. Although there may be some reason to

suspect that may be the case, these factors are unlikely to have affected the results significantly due

to small amplitude nature of these membrane potential variations.

References

Bekkers JM. 1994. Quantal analysis of synaptic transmission in the central nervous system. *Curr Opin*

Neurobiol **4**:360–365. doi:https://doi.org/10.1016/0959-4388(94)90097-3

Biró ÁA, Holderith NB, Nusser Z. 2005. Quantal Size Is Independent of the Release Probability at

Hippocampal Excitatory Synapses. *The Journal of Neuroscience* **25**:223–232.

doi:10.1523/JNEUROSCI.3688-04.2005

Brown TH, Wong RKS, Prince DA. 1979. Spontaneous miniature synaptic potentials in hippocampal

neurons. *Brain Res* **177**:194–199. doi:https://doi.org/10.1016/0006-8993(79)90931-4

Clamann HP, Rioult-Pedotti MS, Luscher HR. 1991. The influence of noise on quantal EPSP size

obtained by deconvolution in spinal motoneurons in the cat. *J Neurophysiol* **65**:67–75.

doi:10.1152/jn.1991.65.1.67

Clements JD. 2003. Variance–mean analysis: a simple and reliable approach for investigating

synaptic transmission and modulation. *J Neurosci Methods* **130**:115–125.

doi:https://doi.org/10.1016/j.jneumeth.2003.09.019

Clements JD, Silver RA. 2000. Unveiling synaptic plasticity: a new graphical and analytical approach.

Trends Neurosci **23**:105–113. doi:10.1016/S0166-2236(99)01520-9

del Castillo J, Katz B. 1954. Quantal components of the end-plate potential. *J Physiol* **124**:560–573.

doi:https://doi.org/10.1113/jphysiol.1954.sp005129

Dervinis M, Major G. 2023. A novel method for reliably measuring miniature and spontaneous

postsynaptic potentials/currents in whole-cell patch clamp recordings in the central nervous

system. *bioRxiv*. doi:10.1101/2022.03.20.485046

Edwards FR, Redman SJ, Walmsley B. 1976. Statistical fluctuations in charge transfer at Ia synapses

on spinal motoneurones. *J Physiol* **259**:665–688.

doi:https://doi.org/10.1113/jphysiol.1976.sp011488

Enoki R, Hu Y, Hamilton D, Fine A. 2009. Expression of Long-Term Plasticity at Individual Synapses in

Hippocampus Is Graded, Bidirectional, and Mainly Presynaptic: Optical Quantal Analysis.

Neuron **62**:242–253. doi:10.1016/j.neuron.2009.02.026

Fatt P, Katz B. 1952. Spontaneous subthreshold activity at motor nerve endings. *J Physiol* **117**:109–

128.

Forti L, Bossi M, Bergamaschi A, Villa A, Malgaroli A. 1997. Loose-patch recordings of single quanta at

individual hippocampal synapses. *Nature* **388**:874–878. doi:10.1038/42251

Hardingham NR, Read JCA, Trevelyan AJ, Nelson JC, Jack JJB, Bannister NJ. 2010. Quantal Analysis

Reveals a Functional Correlation between Presynaptic and Postsynaptic Efficacy in Excitatory

Connections from Rat Neocortex. *The Journal of Neuroscience* **30**:1441–1451.

doi:10.1523/JNEUROSCI.3244-09.2010

Hawkins J, Lewis M, Klukas M, Purdy S, Ahmad S. 2019. A Framework for Intelligence and Cortical

Function Based on Grid Cells in the Neocortex. *Front Neural Circuits* **12**.

doi:10.3389/fncir.2018.00121

Heuser JE, Reese TS, Dennis MJ, Jan Y, Jan L, Evans L. 1979. Synaptic vesicle exocytosis captured by

quick freezing and correlated with quantal transmitter release. *Journal of Cell Biology* **81**:275–

300. doi:10.1083/jcb.81.2.275

Isaacson JS, Walmsley B. 1995. Counting quanta: Direct measurements of transmitter release at a

central synapse. *Neuron* **15**:875–884. doi:10.1016/0896-6273(95)90178-7

Jack JJ, Redman SJ, Wong K. 1981. The components of synaptic potentials evoked in cat spinal

motoneurones by impulses in single group Ia afferents. *J Physiol* **321**:65–96.

doi:https://doi.org/10.1113/jphysiol.1981.sp013972

Jonas P, Major G, Sakmann B. 1993. Quantal components of unitary EPSCs at the mossy fibre

synapse on CA3 pyramidal cells of rat hippocampus. *J Physiol* **472**:615–663.

doi:https://doi.org/10.1113/jphysiol.1993.sp019965

Katz B. 1969. The Release of Neural Transmitter Substances. Liverpool: Liverpool Univ. Press.

Korn H, Faber DS. 1991. Quantal analysis and synaptic efficacy in the CNS. *Trends Neurosci* **14**:439–

445. doi:https://doi.org/10.1016/0166-2236(91)90042-S

Kraszewski K, Grantyn R. 1992. Unitary, quantal and miniature gaba-activated synaptic chloride

currents in cultured neurons from the rat superior colliculus. *Neuroscience* **47**:555–570.

doi:https://doi.org/10.1016/0306-4522(92)90165-X

Kullmann DM, Nicoll RA. 1992. Long-term potentiation is associated with increases in quantal

content and quantal amplitude. *Nature* **357**:240–244. doi:10.1038/357240a0

Kuno M. 1964. Quantal components of excitatory synaptic potentials in spinal motoneurones. *J

Physiol* **175**:81–99. doi:https://doi.org/10.1113/jphysiol.1964.sp007504

Larkman A, Stratford K, Jack J. 1991. Quantal analysis of excitatory synaptic action and depression in

hippocampal slices. *Nature* **350**:344–347. doi:10.1038/350344a0

Larkum ME, Nevian T, Sandler M, Polsky A, Schiller J. 2009. Synaptic Integration in Tuft Dendrites of

Layer 5 Pyramidal Neurons: A New Unifying Principle. *Science (1979)* **325**:756–760.

doi:10.1126/science.1171958

Liu Z, Lu X, Villette V, Gou Y, Colbert KL, Lai S, Guan S, Land MA, Lee J, Assefa T, Zollinger DR,

Korympidou MM, Vlasits AL, Pang MM, Su S, Cai C, Froudarakis E, Zhou N, Patel SS, Smith CL,

Ayon A, Bizouard P, Bradley J, Franke K, Clandinin TR, Giovannucci A, Tolias AS, Reimer J,

Dieudonné S, St-Pierre F. 2022. Sustained deep-tissue voltage recording using a fast indicator

evolved for two-photon microscopy. *Cell* **185**:3408-3425.e29. doi:10.1016/J.CELL.2022.07.013

Major G. 1993. Solutions for transients in arbitrarily branching cables: III. Voltage clamp problems.

Biophys J **65**:469–491. doi:10.1016/S0006-3495(93)81039-7

Major G, Evans J D, Jack J J. 1993. Solutions for transients in arbitrarily branching cables: I. Voltage

recording with a somatic shunt. *Biophys J* **65**:423–449.

Major G, Evans JD, Jack JJ. 1993. Solutions for transients in arbitrarily branching cables: I. Voltage

recording with a somatic shunt. *Biophys J* **65**:423–449. doi:10.1016/S0006-3495(93)81037-3

Major G, Larkum ME, Schiller J. 2013a. Active Properties of Neocortical Pyramidal Neuron Dendrites.

Annu Rev Neurosci **36**:1–24. doi:10.1146/annurev-neuro-062111-150343

Major G, Larkum ME, Schiller J. 2013b. Active Properties of Neocortical Pyramidal Neuron Dendrites.

Annu Rev Neurosci **36**:1–24. doi:10.1146/annurev-neuro-062111-150343

Nevian T, Larkum ME, Polsky A, Schiller J. 2007. Properties of basal dendrites of layer 5 pyramidal

neurons: a direct patch-clamp recording study. *Nat Neurosci* **10**:206–214. doi:10.1038/nn1826

Oertner TG, Sabatini BL, Nimchinsky EA, Svoboda K. 2002. Facilitation at single synapses probed with

optical quantal analysis. *Nat Neurosci* **5**:657–664. doi:10.1038/nn867

Peled ES, Isacoff EY. 2011. Optical quantal analysis of synaptic transmission in wild-type and rab3-

mutant Drosophila motor axons. *Nat Neurosci* **14**:519–526. doi:10.1038/nn.2767

Pircher T, Pircher B, Feigenspan A. 2022. A novel machine learning-based approach for the detection

and analysis of spontaneous synaptic currents. *PLoS One* **17**:1–16.

doi:10.1371/journal.pone.0273501

Rall W. 1977. Core Conductor Theory and Cable Properties of Neurons In: Kandel ER, Brookhardt JM,

Mountcastle VM, editors. Handbook of Physiology, Cellular Biology of Neurons. Bethesda, MD:

American Physiological Society. pp. 39–97.

Rall W, Segev I. 1985. Space-clamp problems when voltage clamping branched neurons with

intracellular microelectrodes In: T. G. Smith Jr, Lecar H, Redman SJ, Gage P, editors. Voltage and

Patch Clamping with Microelectrodes. Bethesda, MD: American Physiological Society. pp. 191–

215.

Redman S. 1990. Quantal analysis of synaptic potentials in neurons of the central nervous system.

Physiol Rev **70**:165–198. doi:10.1152/physrev.1990.70.1.165

Reid CA, Clements JD. 1999. Postsynaptic expression of long-term potentiation in the rat dentate

gyrus demonstrated by variance-mean analysis. *J Physiol* **518**:121–130.

doi:https://doi.org/10.1111/j.1469-7793.1999.0121r.x

Sahara Y, Takahashi T. 2001. Quantal components of the excitatory postsynaptic currents at a rat

central auditory synapse. *J Physiol* **536**:189–197. doi:https://doi.org/10.1111/j.1469-

7793.2001.00189.x

Silver RA. 2003. Estimation of nonuniform quantal parameters with multiple-probability fluctuation

analysis: theory, application and limitations. *J Neurosci Methods* **130**:127–141.

doi:https://doi.org/10.1016/j.jneumeth.2003.09.030

Spruston N, Jaffe DB, Williams SH, Johnston D. 1993. Voltage- and space-clamp errors associated

with the measurement of electrotonically remote synaptic events. *J Neurophysiol* **70**:781–802.

doi:10.1152/jn.1993.70.2.781

Stern P, Edwards FA, Sakmann B. 1992. Fast and slow components of unitary EPSCs on stellate cells

elicited by focal stimulation in slices of rat visual cortex. *J Physiol* **449**:247–278.

doi:https://doi.org/10.1113/jphysiol.1992.sp019085

Stuart G, Spruston N. 1998. Determinants of Voltage Attenuation in Neocortical Pyramidal Neuron

Dendrites. *The Journal of Neuroscience* **18**:3501–3510. doi:10.1523/JNEUROSCI.18-10-

03501.1998

Vere-Jones D. 1966. SIMPLE STOCHASTIC MODELS FOR THE RELEASE OF QUANTA OF TRANSMITTER

FROM A NERVE TERMINAL. *Australian Journal of Statistics* **8**:53–63.

doi:https://doi.org/10.1111/j.1467-842X.1966.tb00164.x

Wall MJ, Usowicz MM. 1998. Development of the quantal properties of evoked and spontaneous

synaptic currents at a brain synapse. *Nat Neurosci* **1**:675–682. doi:10.1038/3677

Wang N-S, Marino M, Malinow R. 2024. Detecting unitary synaptic events with machine learning.

Proceedings of the National Academy of Sciences **121**:e2315804121.

doi:10.1073/pnas.2315804121

Williams SR, Mitchell SJ. 2008. Direct measurement of somatic voltage clamp errors in central

neurons. *Nat Neurosci* **11**:790–798. doi:10.1038/nn.2137

Williams SR, Stuart GJ. 2002. Dependence of EPSP Efficacy on Synapse Location in Neocortical

Pyramidal Neurons. *Science (1979)* **295**:1907–1910. doi:10.1126/science.1067903

Yuste R, Majewska A, Cash SS, Denk W. 1999. Mechanisms of Calcium Influx into Hippocampal

Spines: Heterogeneity among Spines, Coincidence Detection by NMDA Receptors, and Optical

Quantal Analysis. *The Journal of Neuroscience* **19**:1976–1987. doi:10.1523/JNEUROSCI.19-06-

01976.1999

Figures

Figure 1: Recording quality measures used during the data preprocessing stage to select 'noise +

minis' and 'noise-alone' recording sweeps for further data analysis for recording p103a (layer 5).

(A) Baseline V_m across the entire recording session. The raw trace is very similar to the smoothed

one and, therefore, is occluded by the latter. Stimulation pulses were excluded from the display. The

total recording duration (excluding pulse periods) was 8520 s. The first red vertical line marks the

time when pharmacological blockade of APs was initiated. The second red vertical line marks the

time when no more APs were observing in the recording. The third red vertical line marks the time

when pharmacological blockade of mEPSPs was initiated. Bright yellow periods mark 'noise + minis'

recording segment selected as the main comparison file for the distribution fitting. Light yellow

marks all the remaining good quality 'noise + minis' recording sweeps that were also included in the

distribution fitting procedure for estimating target SAD, MAD, and 'deviation' scores.

(B) V_m standard deviation measured over consecutive 15 ms windows over the same recording

session. The meaning of symbols is the same as in panel (A).

(C) Evolution of mean and median amplitudes of the top 10% (amplitude-wise) of all detected minis-

like events averaged over single sweeps across the entire recording session.

(D) Effective decay (drop to 1/e of the peak value) time constants of the same top 10% (amplitude-

wise) of detected events as in panel (C) averaged over consecutive 20-second-long segments across

the same session.

(E) Evolution of passive membrane time constant over the entire recording session. Measurements

were averaged over consecutive 20-second-long segments. Both measurements based on

exponential decay curve fit and on effective V_m drop to 1/e of the initial value were calculated and

displayed.

(F) Evolution of total cell's membrane capacitance across the recording session. Measurements were

averaged over consecutive 20-second-long segments. Both measurements based on exponential

decay curve fit and on effective V_m drop to 1/e of the initial value were calculated and displayed.

(G) Evolution of cell's input resistance (the bridge balance error) across the recording session.

Measurements were averaged over consecutive 20-second-long segments.

Figure 2: Constructing 'noise + simulated minis' voltage trace.

(A) A voltage trace of a physiological noise recording superimposed with simulated mEPSPs.

(B) A voltage trace showing simulated mEPSPs only that were used to produce the voltage trace in

(A).

(C) A segment of 'noise-alone' recording that was used to produce the voltage trace in (A). The noise

recording was obtained from cell p131a in the cortical layer 2/3.

Figure 3: Amplitude and rise time distributions of detected minis-like events in 'noise + minis' and

'noise-alone' recordings.

(A) Amplitude distributions of detected minis-like events in 'noise + minis' and 'noise-alone'

recordings (p108b cell in the cortical layer 5). The legend is provided in panel (B). The read

arrowhead indicates negative bins after subtracting 'noise-alone' distribution from 'noise + minis'

distribution.

(B) 10-90% rise time distributions of detected minis-like events in 'noise + minis' and 'noise-alone'

recordings.

Figure 4: Relationship between the inverse of cell's total capacitance and the mean amplitude of

minis recorded at its soma after subtracting the amplitude distribution based on the 'noise-alone'

recording from the amplitude distribution based on the 'minis + noise' distribution. The blue line is

the least square fit to the data while the black line is the fit to the data that passes through the

origin of the axes.

Figure 5: Distributions of minis-like events detected in a 'noise + simulated minis' voltage trace that

closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p106b of the cortical layer 5. Interval marked by green indicates the range of values

across multiple 'noise + real minis' recording files. The yellow shaded interval indicates the triple size

range.

(A) Amplitude distribution of minis-like events detected in a 'noise + simulated minis' voltage trace

(the blue curve).

(B) 10-90% rise time distribution of minis-like events detected in a 'noise + simulated minis' voltage

trace (the blue curve).

Figure 6: An example of simulated minis distribution that was used to produce a close match

between distributions of detected minis-like events in a 'noise + simulated minis' voltage trace and

in a 'noise + real minis' recording for a cell p131c of cortical layer 2/3.

(A) A joint amplitude and 10-90% rise time distribution of simulated minis.

(B) The same distribution but viewed from a different angle.

(C) Marginal amplitude distribution of the same distribution as in (A).

(D) Marginal 10-90% rise time distribution of the same distribution as in (A).

Figure 7: Estimated minis incidence rates using various methods.

'Thresholded' method: 0.8 mV detection threshold was applied to all 14 'noise + real minis'

recordings and only events with larger amplitudes were included in the estimate.

'Subtracted' method: Distribution of minis-like events in the 'noise-alone' recording was subtraceted

from the distribution of minis-like events in the 'noise + real minis' recording and the incidence rate

estimate was based on the remaining distribution.

'Fitted' method: Simulated minis distribution was used to estimate the minis' incidence rate.

Black markers and black vertical lines correspond to means and 95% confidence intervals. Red

dotted lines mark the range of incidence rates typically reported in the literature and discussed in

the companion article (Dervinis and Major, 2023).

Figure 8: Relationship between cell's inverse total capacitance and estimated mean minis amplitude

using simulations of minis and minis' distribution fitting.

(A) Correlation between mean minis amplitudes as derived using simulated minis' disitrbution fitting

method and the inverse of cell's total capacitance. The blue line is the least square fit to the data

while the black line is the fit to the data that passes through the origin of the axes.

(B) Correlation between the 'subtracted' mean minis amplitude as shown in Figure 3 and the 'fitted'

mean minis amplitude.

Figure 9: Generic minis distribution.

Distributions resulting from all 14 recordings were averaged and the resulting distribution was

smoothed. All minis' amplitudes were normalised by means of their respective source distributions.

(A) A joint amplitude and 10-90% rise time distribution of simulated minis.

(B) The same distribution but viewed from a different angle.

(C) Marginal amplitude distribution of the same distribution as in (A).

(D) Marginal 10-90% rise time distribution of the same distribution as in (A).

 Supplementary Figures

Supplementary Figure 1: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p106b (layer 5).

(A) Baseline V_m across the entire recording session. The raw trace is very similar to the smoothed

one and, therefore, is occluded by the latter. Stimulation pulses were excluded from the display. The

total recording duration (excluding pulse periods) was 4960 s. The first red vertical line marks the

time when pharmacological blockade of APs was initiated. The second red vertical line marks the

time when no more APs were observing in the recording. The third red vertical line marks the time

when pharmacological blockade of mEPSPs was initiated. Bright yellow periods mark 'noise + minis'

recording segment selected as the main comparison file for the distribution fitting. Light yellow

marks all the remaining good quality 'noise + minis' recording sweeps that were also included in the

distribution fitting procedure for estimating target SAD, MAD, and 'deviation' scores.

(B) V_m standard deviation measured over consecutive 15 ms windows over the same recording

session. The meaning of symbols is the same as in panel (A).

(C) Evolution of mean and median amplitudes of the top 10% (amplitude-wise) of all detected minis-

like events averaged over single sweeps across the entire recording session.

(D) Effective decay (drop to 1/e of the peak value) time constants of the same top 10% (amplitude-

wise) of detected events as in panel (C) averaged over consecutive 20-second-long segments across

the same session.

(E) Evolution of passive membrane time constant over the entire recording session. Measurements

were averaged over consecutive 20-second-long segments. Both measurements based on

exponential decay curve fit and on effective V_m drop to 1/e of the initial value were calculated and

displayed.

(F) Evolution of total cell's membrane capacitance across the recording session. Measurements were

averaged over consecutive 20-second-long segments. Both measurements based on exponential

decay curve fit and on effective V_m drop to 1/e of the initial value were calculated and displayed.

(G) Evolution of cell's input resistance (the bridge balance error) across the recording session.

Measurements were averaged over consecutive 20-second-long segments.

Supplementary Figure 2: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p108a (layer 5).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 5140 s.

Supplementary Figure 3: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p108b (layer 5).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 2400 s.

Supplementary Figure 4: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p108c (layer 5).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 2000 s.

Supplementary Figure 5: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p120b (layer 2/3).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 4000 s.

Supplementary Figure 6: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p122a (layer 2/3).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 4200 s.

Supplementary Figure 7: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p124b (layer 5).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 2400 s.

Supplementary Figure 8: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p125a (layer 2/3).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 3800 s.

Supplementary Figure 9: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p127c (layer 2/3).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 3200 s.

Supplementary Figure 11: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p129a (layer 5).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 4000 s.

Supplementary Figure 12: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p131a (layer 2/3).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 2400 s.

Supplementary Figure 13: Recording quality measures used during the data preprocessing stage to

select 'noise + minis' and 'noise-alone' recording sweeps for further data analysis for recording

p131c (layer 2/3).

The meaning of panels A-G is explained in Supplementary figure 1 except for data being averaged

over 100-second-long windows in panels D-G. The figure shows only part of the recording following

the blockade of APs with only a single red vertical time marker visible indicating the time when

pharmacological EPSP blockers were infused. The total duration of the recording shown in the figure

was 2200 s.

Supplementary figure 14: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p103a of the cortical layer 5. Interval marked by green indicates the range of values

across multiple 'noise + real minis' recording files. The yellow shaded interval indicates the triple size

range.

(A) Amplitude distribution of minis-like events detected in a 'noise + simulated minis' voltage trace

(the blue curve).

(B) 10-90% rise time distribution of minis-like events detected in a 'noise + simulated minis' voltage

trace (the blue curve).

Supplementary figure 15: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p108a of the cortical layer 5. The meaning of the two panels and different colours is

explained in Supplementary figure 14.

Supplementary figure 16: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p108b of the cortical layer 5. The meaning of the two panels and different colours is

explained in Supplementary figure 14.

Supplementary figure 17: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p108c of the cortical layer 5. The meaning of the two panels and different colours is

explained in Supplementary figure 14.

Supplementary figure 18: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p120b of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

Supplementary figure 19: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p122a of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

Supplementary figure 20: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p124b of the cortical layer 5. The meaning of the two panels and different colours is

explained in Supplementary figure 14.

Supplementary figure 21: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p125a of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

Supplementary figure 22: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p127c of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

Supplementary figure 23: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p128c of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

Supplementary figure 24: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p129a of the cortical layer 5. The meaning of the two panels and different colours is

explained in Supplementary figure 14.

Supplementary figure 25: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p131a of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

Supplementary figure 26: Distributions of minis-like events detected in a 'noise + simulated minis'

voltage trace that closely matched 'noise + real minis' recordings.

Parameters of simulated minis were controlled by the GA, whereas the 'noise-alone' recording was

obtained in cell p131c of the cortical layer 2/3. The meaning of the two panels and different colours

is explained in Supplementary figure 14.

1003

1004

Supplementary figure 27: An example of simulated minis distribution that was used to produce a

close match between distributions of detected minis-like events in a 'noise + simulated minis'

voltage trace and in a 'noise + real minis' recording for a cell p103a of cortical layer 5.

(A) A joint amplitude and 10-90% rise time distribution of simulated minis.

(B) The same distribution but viewed from a different angle.

(C) Marginal amplitude distribution of the same distribution as in (A).

(D) Marginal 10-90% rise time distribution of the same distribution as in (A).

1012

1013

1014   **Supplementary figure 28:** An example of simulated minis distribution that was used to produce a

1015   close match between distributions of detected minis-like events in a 'noise + simulated minis'

1016   voltage trace and in a 'noise + real minis' recording for a cell p106b of cortical layer 5.

1017   The meaning of different figure panels is explained in Supplementary figure 27.

1018

1019

1020 **Supplementary figure 29:** An example of simulated minis distribution that was used to produce a

1021 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1022 voltage trace and in a 'noise + real minis' recording for a cell p108a of cortical layer 5.

1023 The meaning of different figure panels is explained in Supplementary figure 27.

1024

1025

1026 **Supplementary figure 30:** An example of simulated minis distribution that was used to produce a

1027 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1028 voltage trace and in a 'noise + real minis' recording for a cell p108b of cortical layer 5.

1029 The meaning of different figure panels is explained in Supplementary figure 27.

1030

1031

1032   **Supplementary figure 31:** An example of simulated minis distribution that was used to produce a

1033   close match between distributions of detected minis-like events in a 'noise + simulated minis'

1034   voltage trace and in a 'noise + real minis' recording for a cell p108c of cortical layer 5.

1035   The meaning of different figure panels is explained in Supplementary figure 27.

1036

1037

1038 **Supplementary figure 32:** An example of simulated minis distribution that was used to produce a

1039 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1040 voltage trace and in a 'noise + real minis' recording for a cell p120b of cortical layer 2/3.

1041 The meaning of different figure panels is explained in Supplementary figure 27.

1042

1043

1044 **Supplementary figure 33:** An example of simulated minis distribution that was used to produce a

1045 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1046 voltage trace and in a 'noise + real minis' recording for a cell p122a of cortical layer 2/3.

1047 The meaning of different figure panels is explained in Supplementary figure 27.

1048

1049

1050 **Supplementary figure 34:** An example of simulated minis distribution that was used to produce a

1051 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1052 voltage trace and in a 'noise + real minis' recording for a cell p124b of cortical layer 5.

1053 The meaning of different figure panels is explained in Supplementary figure 27.

1054

1055

1056 **Supplementary figure 35:** An example of simulated minis distribution that was used to produce a

1057 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1058 voltage trace and in a 'noise + real minis' recording for a cell p125a of cortical layer 2/3.

1059 The meaning of different figure panels is explained in Supplementary figure 27.

1060

1061

1062 **Supplementary figure 36:** An example of simulated minis distribution that was used to produce a

1063 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1064 voltage trace and in a 'noise + real minis' recording for a cell p127c of cortical layer 2/3.

1065 The meaning of different figure panels is explained in Supplementary figure 27.

1066

1067

1068 **Supplementary figure 37:** An example of simulated minis distribution that was used to produce a

1069 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1070 voltage trace and in a 'noise + real minis' recording for a cell p128c of cortical layer 2/3.

1071 The meaning of different figure panels is explained in Supplementary figure 27.

1072

1073

1074 **Supplementary figure 38:** An example of simulated minis distribution that was used to produce a

1075 close match between distributions of detected minis-like events in a 'noise + simulated minis'

1076 voltage trace and in a 'noise + real minis' recording for a cell p129a of cortical layer 5.

1077 The meaning of different figure panels is explained in Supplementary figure 27.

1078

1079

1080   **Supplementary figure 39:** An example of simulated minis distribution that was used to produce a

1081   close match between distributions of detected minis-like events in a 'noise + simulated minis'

1082   voltage trace and in a 'noise + real minis' recording for a cell p131a of cortical layer 2/3.

1083   The meaning of different figure panels is explained in Supplementary figure 27.