Ultra-resolution Deconvolution of Spatial Transcriptomics Unveils Spatiotemporal

Cellular Dynamics in Complex Microenvironments

Authors: Yin Xu[1,2]†, Zurui Huang[1,2]†, Yawei Zhang[1,2]†, Zhenghang Wang[3]†, Peijin Guo[4],

Feifan Zhang[1,2], Wenxuan Gong[1,2], Guanghao Liang[1,2], Boyuan Mei[4], Lihui Dong[1,2],

Renbao Chang[1,2], Minghui Gong[1,2], Yu Xia[4], Haochen Ni[1,2], Jing Yang[1,2], Yuan Gao[1,2], Zhaoqi

Liu[1,2], Lin Shen[3], Jian Li[3], Meng Michelle Xu[4]*, and Dali Han[1,2]*

Affiliations:

[1] Department of Computational Biology, Key Laboratory of RNA Science and Engineering,

China National Center for Bioinformation, Beijing Institute of Genomics, Chinese Academy

of Sciences; Beijing 100101, China.

[2]College of Future Technology, Sino-Danish College, University of Chinese Academy of

Sciences, Beijing 100049, China.

[3]Department of Gastrointestinal Oncology, Key Laboratory of Carcinogenesis and

Translational Research (Ministry of Education), Peking University Cancer Hospital &

Institute, Beijing 100142, China.

[4]State Key Laboratory of Molecular Oncology, School of Basic Medical Sciences, Institute

for Immunology, Tsinghua University; Beijing 100871, China.

† These authors contributed equally.

*Corresponding author. Email: michellexu@mail.tsinghua.edu.cn (M.M.X.);

20   handl@big.ac.cn (D. H.)

Abstract:

The current challenge in analyzing sequencing-based spatial transcriptomics (ST) is to precisely

resolve the spatial distribution of cell subpopulations with low abundance and detect context-

dependent variation in cell states within complex tissue microenvironments. Here, we introduce

an ultra-resolution ST deconvolution algorithm (UCASpatial) that improves the resolution of

mapping cell subpopulations to spatial locations by leveraging the contribution of genes

indicative of cell identity through entropy-based weighting. Using both in silico and real ST

datasets, we demonstrate that UCASpatial improves the robustness and accuracy in identifying

low-abundant cell subpopulations and distinguishing transcriptionally heterogeneous cell
subpopulations. Applying UCASpatial to murine wound healing, we recapitulate known
spatiotemporal dynamics of multiple cell subpopulations involved in wound healing and reveal
the spatial segregation of distinct macrophage subpopulations embedded within different
cellular communities at the wound center. In human colorectal cancer (CRC), we link genomic
alterations in individual cancer clones to multi-cellular characteristics of the tumor immune
microenvironment (TIME) and reveal the co-evolution of tumor cells and TIME within the
same tumor. We show that the copy number gain on chromosome 20q (chr20q-gain) in tumor
cells orchestrates a T cell-excluded TIME, indicative of resistance to immunotherapy in CRC,
and is associated with tumor-intrinsic endogenous retrovirus silencing and impaired type I
interferon response. Our findings present UCASpatial as a versatile tool for deciphering ultra-
resolution cellular landscapes in ST and exploring intercellular mechanisms in complex and
dynamic microenvironments.

Introduction
The ability to elucidate the spatial organization of cells within the microenvironment is crucial
for understanding cellular functions and exploring intercellular mechanisms that govern
development, physiology, and pathogenesis[1,2]. Emerging spatial transcriptomics technologies
offer promising avenues for achieving this; however, current spot-based techniques face a
significant limitation in lacking single-cell resolution. Methods like Visium[3], Tomo-Seq[4], Slide-
seq[5], and HDST[6] amalgamate RNA-seq measurements from multiple cells or cell fractions at
each spot. As a result, these techniques struggle with the precise identification of individual cell
types within the microenvironment. Consequently, annotating cell types at each spot becomes
imperative for accurately capturing spatial cell organization[7,8].

To address this challenge, various computational approaches have been proposed to
deconvolute spatial transcriptomic profiles into cellular compositions based on gene expression
profiles of each cell type identified through single-cell RNA sequencing (scRNA-seq), such as
SPOTlight[9], RCTD[10], cell2location[11], CARD[12], and stereoscope[13]. These methods effectively
delineate significant differences in cell type compositions across diverse anatomical or
pathological layers in well-structured tissues such as the brain, heart, and lymphatic structures,
even at fine-grained resolutions[11,12,14-16].

In recent years, a growing body of research has emphasized the importance of
characterizing the complex microenvironment, particularly the influx of immune cells that
orchestrate dynamic spatial patterns, varying between tissue homeostasis and non-homeostasis

conditions. This complexity is evident in scenarios such as tissue damage[17,18] and the tumor

microenvironment (TME)[19-23]. However, several immune cells, particularly those infiltrating

into tumors, are known for their sparse presence and low transcriptional activity[24-27]. Their

discontinuous spatial distributions in these microenvironments limit the effectiveness of

algorithms relying on spot-neighbor information[27,28]. Additionally, the low abundance and the

context-dependent phenotypic variation of heterogeneous immune cells, particularly T cells,

pose challenges for conventional deconvolution algorithms[24]. Furthermore, their scarcity makes

them difficult to capture accurately using histological images, thus constraining the

applicability of image-based algorithms[29,30]. Consequently, the precise characterization of

complex and dynamic microenvironments remains a significant challenge.

To address the aforementioned challenge, we developed UCASpatial, an ultra-resolution

cellular deconvolution algorithm in spatial transcriptomics based on entropy-weighted

regression. A unique advancement of UCASpatial is its formulation of the Shannon entropy-

based weights to quantify the ability of each gene for indicating cell identity and incorporate

this information in the deconvolution process. This approach improves the robustness and

accuracy in identifying low-abundant cell subpopulations and distinguishing transcriptionally

heterogeneous cell subpopulations in ST, thus allowing for precise and unbiased deconvolution

of the cellular landscape. Through simulations and applications to murine wound healing and

human CRC ST datasets, we illustrate UCASpatial's improvements in deconvolution and

demonstrate its utility in revealing spatiotemporal transitions and multicellular communities

among cell subpopulations.

Result

UCASpatial: an entropy-weighted regression model for spatial deconvolution of cellular

landscapes.

UCASpatial deconvolutes ST data into cellular compositions by leveraging cell identity-

specific gene expression profiles (cGEPs) derived from either paired or unpaired scRNA-seq

data (Fig. 1). First, UCASpatial extracts cGEPs from the referenced scRNA-seq dataset with a

set of user-provided cell-subpopulation labels. It strategically selects cells meeting a meta-

purity threshold to enhance discrimination of referenced cell identities, and then extracts cGEPs

through optimized Non-negative Matrix Factorization (NMF). Second, UCASpatial formulates

Shannon entropy-based weights for each gene within the extracted cGEPs to enhance genes

indicative of cell identities (Methods). Third, UCASpatial employs Weighted Non-Negative

Least Square (WNNLS) to deconvolute the mRNA counts of each spot in the ST data using the

weighted cGEPs, thereby estimating the cellular composition. Additionally, UCASpatial offers

a suite of downstream analysis tools. These include identifying tumor clones and their

corresponding TIME, as well as spatially resolved quantification of transposable elements

(TEs). To ensure versatility in applications, UCASpatial is seamlessly integrated with the Seurat

framework.

Benchmarking the performance of UCASpatial in the simulated ST dataset

To assess UCASpatial's performance, we generated a simulated ST dataset comprising 750 ST

spots designed to mimic the real cellular composition of the TME. Each spot was constituted

by the transcripts of 5 to 20 cells sampled from scRNA-seq data of human CRC, encompassing

20 reference cell subpopulations (Supplementary Fig. 1a-c, and Methods). The cellular

composition of simulated spots reflects the actual cellular composition identified in typical

tumor architectures within the TME[16,31] (Supplementary Fig. 1d, and Supplementary Table. 1).

To assess the performance of UCASpatial on the simulated ST dataset with varying

complexity, we ranked the complexity of these simulated spots based on their cell-type

composition entropy. Subsequently, we categorized the simulated spots into three groups (low,

medium, and high complexity), where higher entropy denotes increased complexity (Fig. 2a,

and Supplementary Fig. 1d). We compared UCASpatial with several recently established

methods, including SPOTlight, RCTD, cell2location, CARD, and stereoscope, using Root

Mean Square Error (RMSE) to first quantify the deviation between estimations and ground truth

at the spot level. UCASpatial exhibited superior performance across all complexity levels

(median RMSE = 0.051) compared to the aforementioned methods, with 10%, 21%, 31%, 36%,

and 35% improvement in terms of RMSE compared to RCTD (0.058), SPOTlight (0.066),

CARD (0.075), cell2location (0.081), and stereoscope (0.080), respectively (Fig. 2b).

Additionally, UCASpatial demonstrated a comparable Pearson Correlation Coefficient (PCC)

to cell2location and outperformed other methods, when comparing estimations with the ground

truth (Supplementary Fig. 2b). Moreover, we also assessed the ability for identifying the

existence of cell subpopulations within spots using the F1 score. UCASpatial consistently

outperformed other methods across all complexity levels with the highest F1 score (with a 6-

25% improvement) (Fig. 2b, and Supplementary Fig. 2c).

We next evaluated the precision and robustness of UCASpatial in deconvoluting different

cell subpopulations. Examination of PCC between estimations and ground truth for each

method unveiled UCASpatial's superior performance, achieving the highest PCC in 15 out of

20 cell subpopulations (Fig. 2c), particularly notable for NK cell and CD8[+] T cell

subpopulations (with a 39-187% improvement) (Fig. 2d). By categorizing the PCC values of

each cell subpopulation into four groups, we noticed UCASpatial exhibited PCC values

surpassing the medium-high correlation level (PCC > 0.5) for 19 out of 20 cell subpopulations,

with the sole exception $CD4^+$ Tfh (PCC = 0.47) while remained the best compared to alternative

methods (Fig. 2c, e). In contrast, other methods yielded PCC values below the medium-high

correlation level for at least 30% of cell subpopulations, with some even falling below 0.25 (Fig.

2e), underscoring UCASpatial's capability in discriminating cell subpopulations, even with

subtle transcriptomic differences (Supplementary Fig. 3a, b). Furthermore, assessments using

the PCC and F1 score reaffirmed UCASpatial's precise and robust decomposition across each

cell subpopulation (Supplementary Fig. 3c, d).

Considering the challenge of ST analysis in precisely mapping the spatial location of low-

abundant cell subpopulations, we next investigated UCASpatial's sensitivity in annotating cell

subpopulations with low frequencies within spots. We categorized the proportion of each cell

subpopulation within simulated spots into ten groups, forming a gradient from 0-10% to 0-100%

(Fig. 2f). Subsequently, we calculated the RMSE across various proportions to evaluate the

deviation of cell-subpopulation proportions. We observed that UCASpatial exhibited the lowest

overall RMSE (averaged by each cell subpopulation, medium RMSE = 0.051) across different

proportions with a 7-31% improvement over the other methods (Fig. 2g). Moreover, we

specifically examined the RMSE of different cell subpopulations, revealing that UCASpatial

notably outperformed other methods for cell subpopulations with subtle transcriptomic

differences such as $CD8^+$ Tex and $CD8^+$ Tem (with 25-63% and 13-26% improvements,

respectively) (Fig. 2g and Supplementary Fig. 4). Consistent results were observed in

assessments using the PCC and F1 score (Supplementary Figs. 5-6), highlighting UCASpatial's

superiority in accurately annotating cell subpopulations with low frequencies within spots, even

with subtle transcriptomic differences.

Furthermore, we illustrated how UCASpatial's entropy-based weighting and meta-purity

filtering designs enhance the performance of cellular deconvolution (Supplementary Fig. 7).

Specifically, the entropy-based weighting design showed improvements of 22%, 50%, and 17%

in terms of RMSE, PCC, and F1 score, respectively, compared to UCASpatial without

weighting. The meta-purity filtering design exhibited improvements across different parameters

with 4-11%, 1-8%, and 1-3% in terms of RMSE, PCC, and F1 score. In addition, the same

analyses as before showed that UCASpatial, without entropy-weighting and meta-purity

filtering, lost the previously mentioned improvements (Supplementary Figs. 2-6). Collectively,

our in-silico benchmarking demonstrates that UCASpatial enables precise and sensitive spatial

deconvolution, excelling in discerning cell subpopulations with subtle transcriptomic
differences, even when they exist in low proportions.

UCASpatial accurately annotates cell subpopulations in the murine wound healing ST
dataset

To demonstrate UCASpatial's utility, we applied it to real ST datasets. We primarily
investigated murine wound healing, a well-organized multicellular process characterized by
high complexity and spatiotemporal dynamics[32-34]. Specifically, we employed a 2 mm ear punch
wound model in MRL/MpJ mice (hereafter denoted as MRL), a strain with super-healing
capacity that completely regenerates the ear tissue post-wounding[35-37]. Tissue samples were
collected from MRL mice at various stages of wound healing: unwounded (day 0),
inflammatory stage (day 3), return-to-unwounded stage (day 7), and regeneration stage (day
15)[33] (Fig. 3a). The ST dataset was generated using the 10x Genomics Visium platform. For
reference, we generated scRNA-seq data from day 7 ear tissue, annotated with 20 cell
subpopulations (Fig. 3b, c, and Supplementary Fig. 8a). We then utilized UCASpatial for the
deconvolution of these 20 cell subpopulations within the murine wound healing ST dataset
(Supplementary Fig. 8b).

To delineate the spatial distribution of cell subpopulations in different regions, we divided
the sections into three subdomains: the wound center (2 mm center diameter), wound proximal
(0-1 mm from the wound edge), and wound distal (1-2 mm from the wound edge)
(Supplementary Fig. 8c). We firstly examined the infiltrating cell subpopulations within the
wound center. We noticed the structural cell types, including keratinocytes, endothelial cells,
and fibroblasts, revealing alignment with their respective anatomical positions (Fig. 3c, and
Supplementary Figs. 8b and 10). Notably, keratinocytes demonstrated enrichment in regions of
regenerative epithelium from the return-to-unwounded stage, signifying successful re-
epithelialization[38] (Fig. 3c, and Supplementary Figs. 8b and 10). Additionally, endothelial cells
were observed proximal to blood vessels, while 4 fibroblast subpopulations were enriched at
the wound center during the return-to-unwounded and regeneration stages (Fig. 3c, and
Supplementary Figs. 8b and 10). For example, $Lrrc15^+$ fibroblasts, which exhibited the
characteristics of mature myofibroblasts, were accumulated at the wound center at the return-
to-unwounded stage. Notably, $Lrrc15^+$ fibroblasts comprised over 33% of all cell
subpopulations at that location, consistent with their function in promoting wound contraction[39]
(Fig. 3b, and Supplementary Figs. 8a, d, and 10). The above observations recapitulate the spatial
distribution and dynamics of structural cells during tissue repair[33,38,39].

We also investigated whether UCASpatial could capture the spatial dynamics of wound-
associated immune cells over time. We observed the enrichment of myeloid cells, including
neutrophils and macrophages, at the wound center during the inflammatory stage, in line with
previous studies demonstrating their pivotal role in orchestrating the early immune response to
tissue injury[40,41] (Supplementary Figs. 8b, d, and 10). Interestingly, while neutrophils decreased
during the return-to-unwounded stage, macrophages persisted in high abundance within the
repaired tissues, consistent with previous findings[33,40,42] (Supplementary Figs. 8d and 10).
Similar to neutrophils, basophils were enriched at the wound center in the inflammatory stage
and decreased during the return-to-unwounded stage, indicating that they are the first
responders to injury[43] (Supplementary Figs. 8d and 10).

Subsequently, we conducted a comparative analysis of UCASpatial alongside other
methods in murine wound healing. We calculated the PCC between the spatial distribution of
cell-subpopulation proportion and marker-gene score (Supplementary Table. 2 and Methods),
which was used to quantify the performance of deconvolution without any ground truth[15].
UCASpatial, along with cell2location, SPOTlight, and RCTD, demonstrated high PCC between
the spatial distribution of cell-subpopulation proportions and marker gene expression, while
UCASpatial still showed an improvement of 5%-20% in comparison (Supplementary Fig. 8e,
f). While all considered methods performed well at annotating structural cell subpopulations,
some methods failed to identify several immune cell subpopulations. For example, RCTD and
stereoscope overlooked the spatiotemporal dynamics of neutrophils, which was supposed to be
enriched at the inflammatory stage and declined at the return-to-unwounded stage
(Supplementary Figs. 8g and 10-15). Additionally, RCTD and Cell2location exhibited limited
capability in detecting macrophages on day 0, with proportions less than or equal to 1%
(Supplementary Figs. 8h and 10-15). We also noticed that UCASpatial without entropy
weighting and meta-purity filtering exhibited lower PCC compared with UCASpatial
(Supplementary Figs. 8e and 16). These results demonstrate the superior performance of
UCASpatial in delineating the spatiotemporal dynamics of cell subpopulations.

UCASpatial reconstructs spatiotemporal dynamics of macrophage subpopulations during
murine wound healing
Recent studies suggest that macrophages infiltrate healing tissues and respond to various
microenvironmental signals, leading to their differentiation into specialized subpopulations
with distinct functions and locations[33]. To examine whether UCASpatial could detect the
spatiotemporal dynamics of macrophage subpopulations including *Cd163*[+] macrophages, *Spp1*[+]

macrophages, *MHCII*[+] macrophages (encoded by *H2-Aa* and *H2-Ab1*), and *Il1b*[+] macrophages,

we focused on the wound center at the return-to-unwounded stage (day 7) for further analysis

(Supplementary Fig. 9a, b). We subdivided the wound center into two layers: the epidermis

layer and the dermis layer, based on hematoxylin and eosin (H&E) staining (Fig. 3c).

All four macrophage subpopulations annotated by UCASpatial exhibited distinct spatial

distributions at the wound center, which were consistent with previous reports[33,44] (Fig. 3d).

Specifically, *Cd163*[+] and *Il1b*[+] macrophages were predominantly found around the epidermis

layer, whereas *Spp1*[+] and *MHCII*[+] macrophages were confined to the dermis layer, aligning

with their marker-expression patterns (Supplementary Fig. 9c). Notably, while *Cd163*[+]

macrophages were evenly distributed across the entire wound center, *Il1b*[+] macrophages

exhibited spatial exclusivity, preferentially clustering at the migrating epithelial tongue (Fig.

3d). Functional enrichment analysis indicated that *Cd163*[+] macrophages may play roles in

mesenchymal cell differentiation, endocytosis, and efferocytosis, while *Il1b*[+] macrophages

were enriched for functions related to leukocyte activation and epithelial cell differentiation

(Supplementary Fig. 9d). Intriguingly, multicellular community analysis also confirmed that

Il1b[+] macrophages and keratinocytes co-occurred in the same community, indicating the

function of *Il1b*[+] macrophages in re-epithelialization[44] (Supplementary Fig. 9e).

Although all considered methods displayed exceptional proficiency in elucidating *Cd163*[+]

macrophages, UCASpatial achieved higher accuracy for annotating other macrophage

subpopulations. Particularly, CARD, cell2location, and RCTD failed to identify *Il1b*[+]

macrophages in the migrating epithelial tongue, while SPOTlight inaccurately annotated the

unique spatial distribution patterns of *Spp1*[+] macrophages in the dermis (Fig. 3d).

Subsequently, we examined the spatiotemporal dynamics of macrophages throughout the

wound-healing process. We observed that *Il1b*[+] macrophages appeared in the wound center at

the inflammatory stage but gradually decreased in abundance at the return-to-unwounded stage

(Fig. 3e). In contrast, the other three macrophage subpopulations were predominantly located

on the wound proximal and distal, and had distinct temporal dynamics. While *MHCII*[+]

macrophages were present at all measured time points, *Cd163*[+] macrophages were detected at

the inflammatory stage and were subsequently replaced by *Spp1*[+] macrophages, which peaked

in abundance at the return-to-unwounded stage and declined at the regeneration stage. These

spatial segregations and temporal dynamics of macrophage subpopulations annotated by

UCASpatial indicate their role in coordinating tissue repair towards an unwounded state[40] (Fig.

3e). Collectively, UCASpatial reconstructs an ultra-resolution spatiotemporal cellular

landscape in murine wound healing and reveals the unique spatial segregation and temporal

dynamics of macrophage subpopulations that served in different steps of wound healing.

UCASpatial accurately annotates cell subpopulations in the human CRC ST dataset

Spatial omics has provided insights into the intricate interplay between tumor alterations and

the co-evolved immune microenvironment surrounding tumors[21,45,46]. However, due to the low

abundance and transcriptional output of tumor-infiltrating immune cells[24,26,27], characterization

of immune cell compositions in the vicinity of tumor clones remains a challenge[21,47]. To

investigate whether UCASpatial can precisely resolve the tumor immune microenvironment,

we curated an ST dataset from eight tumor sections of seven patients with human CRC using

the 10x Genomics Visium platform[48] (Fig. 4a, Supplementary Fig. 17a, and Supplementary

Table. 3). We leveraged an integrated scRNA-seq dataset encompassing 34 cell subpopulations

as a reference[48] (Fig. 4b, Supplementary Fig. 17b, c, and Methods).

We initially annotated the spatial distribution of 34 cell subpopulations in the human CRC

ST dataset using UCASpatial (Supplementary Figs. 18-25). Notably, stromal cell

subpopulations showed consistent distribution with anatomical locations as expected from prior

literature, such as smooth muscle cells of the muscular layer and endothelial cells around blood

vessels, along with pericytes[49,50]. Similarly, epithelial cell subpopulations displayed enrichment

in the epithelial region, consistent with corresponding H&E staining and pathological

annotations (Supplementary Fig. 17a). Upon further examination of immune subpopulations,

we noticed B cell and various T cell and dendritic cell (DC) subpopulations, including CD4$^+$

Tn, CD4$^+$ Tfh, CD8$^+$ Tex, CD8$^+$ Tem, cDC2, and cDC3, formed a multicellular community

resembling a tertiary lymphoid structure (TLS)[51,52] (Supplementary Figs. 18-25, see

Supplementary Table. 4 for abbreviation information). Additionally, we noticed the spatial

exclusivity of *C1QC*$^+$ macrophages and *PDPN*$^+$ macrophages, indicating their distinct functions

in TME (Supplementary Figs. 18-25).

For comparative analysis, we also annotated this ST dataset using other methods

(Supplementary Figs. 26-65). Remarkably, UCASpatial achieved the highest average PCC with

an improvement of 17-104% over other methods between the cell-subpopulation proportions

and signature expressions, especially in T cell subpopulations (with at least a 26% improvement)

(Supplementary Fig. 74a, Supplementary Table. 5, and Methods). In alignment with in-silico

benchmarking results, all evaluated methods correctly annotated structural cell subpopulations,

for example, *FAP*$^+$ myofibroblast, yet UCASpatial showed a higher sensitivity in annotating

immune cell subpopulations (Fig. 4c and Supplementary Fig. 74a, b). For instance, UCASpatial

showed superior performance over other methods in identifying the spatial distribution of T cell

subpopulations in almost all five TLS regions (Supplementary Fig. 74c). While UCASpatial

depicted 4-6 T cell subpopulations consistent with their marker gene expressions

(Supplementary Fig. 74c, d), all compared methods only identified 0-2 T cell subpopulations.

Moreover, only UCASpatial captured distinct spatial distribution patterns of three CD8$^+$ T cell

subpopulations infiltrating tumors: CD8$^+$ Tex and CD8$^+$ Trm were proximal to the tumor,

whereas CD8$^+$ Tem was distal (Fig. 4c). Additionally, we noticed UCASpatial without entropy-

weighting and meta-purity filtering were unable to either identify T cell subpopulations in TLS

regions or discriminate the spatial distribution patterns of three CD8$^+$ T cell subpopulations,

exhibiting a comparable overall PCC with cell2location and SPOTlight (Supplementary Figs.

66-74). Collectively, these results demonstrate that UCASpatial can accurately resolve cell

subpopulations in the complex tumor immune microenvironment.

UCASpatial spatially links the tumor genomics with diverse TIME at a clonal resolution.

Tumor-intrinsic genomic alterations have collateral effects that modify the spatial organization

of the immune microenvironment[53-55]. To achieve the spatial linking of tumor genomics with

diverse immune microenvironments at a detailed clonal resolution, we generated an integrated

immunogenomics platform termed 'UCASpatial-clonalTIME'. This platform enables the

identification of tumor clones within tissue sections by inferring spatial copy number variations

(CNVs)[46] and concurrently estimating the composition of the TIME for each clone (clonalTIME)

using UCASpatial (Fig. 4c, Supplementary Fig. 75a, and Methods).

By assessing the proportions of CD4$^+$ and CD8$^+$ T cell subpopulations, we categorized the

clonalTIME into two phenotypes: T cell-rich and T cell-excluded (Supplementary Fig. 75b, c,

and Methods). We detected three out of thirteen tumor clones displaying the T cell-rich

clonalTIME phenotype (Fig. 4d, and Supplementary Fig. 75c, d). Within these T cell-rich

clonalTIMEs, all three subpopulations of CD8$^+$ T cells-CD8$^+$ Tem, CD8$^+$ Trm, and CD8$^+$ Tex-

were notably enriched (Fig. 4d). Analysis of the CD4$^+$ T cell compartment indicated significant

enrichment of CD4$^+$ Tm and Th1/Th17 within the T cell-rich clonalTIMEs (Fig. 4d).

Additionally, CD4$^+$ Tn and CD4$^+$ Tfh showed enrichment within these T cell-rich clonalTIMEs,

while CD4$^+$ Treg did not exhibit a significant enrichment pattern between these two

clonalTIME phenotypes in our ST dataset (Fig. 4d).

We next explored local cues that may influence the infiltration of cytotoxic T cells and the

transition of clonalTIMEs. Given that antigen-presenting cells (APCs) such as DCs and

macrophages can instruct T cell response[56-60], we assumed that these APC subpopulations with

functional complexity might exhibit spatial specifications that either support T cell infiltration

or hinder it. To this end, we identified potential macrophage-T multicellular communities. By

analyzing the PCC between various cell subpopulations across thirteen tumor clones, we

identified a community (community3) comprising $C1QC^+$ macrophages, CD8$^+$ Tem, and CD8$^+$

Tex, which displayed a significant enrichment in T cell-rich clonalTIMEs (Fig. 4e, f, and

Supplementary Fig. 76a). This enrichment pattern of community3 was observed during the

progression of intra-tumoral clonalTIME transitions, indicating the spatiotemporal dynamics of

this community during tumor evolution (Fig. 4g).

Intriguingly, $C1QC^+$ macrophages exhibited upregulated expression of genes encoding

migration-inducing chemokines (*CCL8*, *CCL13*, and *CXCL12*), major histocompatibility

complex (MHC) molecules (*HLA-DPA1*, *HLA-DPB1*, *HLA-DQA1*, *HLA-DQB1*, and *HLA-*

DQB2), and complement components (*C1QA*, *C1QB*, and *C1QC*) (Supplementary Fig. 76b, c).

Furthermore, co-expression module analysis also suggested that $C1QC^+$ macrophages support

T cell activation in CRC[57,61,62] (Supplementary Fig. 65d-f). However, employing alternative

methods for ultra-resolution characterization of the clonalTIME revealed significant limitations.

For instance, cell2location, CARD, SPOTlight, and RCTD failed to capture the intra-tumor

heterogeneity across the distribution of T cell subpopulations at a clonal resolution

(Supplementary Fig. 77). Moreover, stereoscope, SPOTlight, and CARD failed to detect the co-

occurrence pattern between $C1QC^+$ macrophages and CD8$^+$ Tem or CD8$^+$ Tex (Supplementary

Figs. 26-33, 50-57, 78). Together, these data show that UCASpatial links the tumor genomics

with diverse immune microenvironments at a clonal resolution, and reveals the spatiotemporal

dynamics of multicellular communities during clonal evolution in CRC.

UCASpatial identifies chr20q-gain as a tumor genomic feature associated with the

formation of T cell-excluded clonalTIME in CRC

To investigate whether genetic alterations are associated with immune phenotypic transitions

in tumor clones, we conducted a comparison of CNVs across clones using our 'UCASpatial-

clonalTIME' platform (Supplementary Fig. 79a, and Supplementary Table. 6). We found that T

cell-excluded clones displayed specific CNVs compared to T cell-rich clones (Fig. 4d, and Fig.

5a). Notably, we observed a frequent copy number gain on chromosomes 20q and 20p (chr20q-

gain and chr20p-gain) in T cell-excluded clones. We hypothesized that these copy number gains

on chr20q and 20p emerged during clonal evolution and were co-opted with clonalTIME

transitions.

To further investigate this, we focused on a particular section (CRC4) containing three

clones with sequential clonal events. In this section, clone1, with chr1p-loss-2 as a common

ancestor, gave rise to clone2 with chr8p-gain and chr8q-gain, alongside clone3 with chr20q-

gain, chr20p-gain, and three additional CNVs (Fig. 5b, c). Interestingly, as clone1 progressed

to clone2, the clonalTIME remained T cell-rich (Fig. 4d, Fig. 5a). However, upon progression

from clone1 to clone3, we observed a significant decrease in CD8$^+$ T cell infiltration, especially

CD8$^+$ Tex, and myeloid infiltration, especially $C1QC^+$ macrophages within their clonalTIMEs

(Fig. 4d, e).

For further validation, we conducted an analysis of the PCC between the CNVs detected

in CRC4 and the signature expression of distinct cell subpopulations using the TCGA-COAD

dataset (Supplementary Fig. 79b). Our examination revealed significant correlations between

clone3-specific CNVs, encompassing chr20q-gain, chr20p-gain, chr7p-gain, chr7q-gain, and

chr17p-loss, and reduced signature expression of CD8$^+$ Tex, CD8$^+$ Tem, CD8$^+$ Trm, CD4$^+$ Tm,

CD4$^+$ Tfh, and CD4$^+$ Treg cells (Fig. 5d). Notably, these five CNVs exhibited a concurrent

occurrence pattern within the TCGA-COAD dataset (Fig. 5e). To ascertain which CNV

primarily contributes to diminished T cell infiltration, we isolated the individual impacts of each

CNV in the TCGA-COAD dataset based on the status of these five CNVs (Methods). Our

analysis revealed that only the group harboring chr20q-gain, prevalent in 57.9% of CRC

patients, exhibited a significant reduction in the expression levels of T cell marker genes (*CD3G*,

CD8A, and *CD4*) compared to the wild-type (WT) group (Fig. 5f and Supplementary Fig. 79c).

These findings suggest that the emergence of chr20q-gain may contribute to the transition of

the clonalTIME phenotype from T cell-rich to T cell-excluded.

Molecular characteristics of clones with T cell-excluded clonalTIME

We proceeded to investigate the spatial molecular characteristics of these T cell-excluded tumor

clones. Functional enrichment analysis unveiled an enrichment for pathways related to

translation machinery among the upregulated genes in clone3 as compared to clone1 and clone2.

Conversely, pathways related to type I interferon (IFN) production were notably enriched

among the downregulated genes (Fig. 5g, and Supplementary Fig. 80a-d). Notably, chemokine-

coding genes responsible for cytotoxic CD8$^+$ T cell recruitment, such as *CXCL9* and *CXCL10*,

and genes associated with antigen presentation, including *TAP1*, *TAP2*, *HLA-A*, *HLA-B*, and

B2M, exhibited reduced expression[63,64] (Fig. 5g, and Supplementary Fig. 80a-c). This

observation potentially elucidates the T cell-excluded phenotype of clone3.

Research has demonstrated that the induction of the type I IFN response in tumors can be

attributed to the innate sensing of cytosolic RNA derived from TEs[65]. Considering that TEs

serve as internal adjuvants eliciting T cell responses against tumors, we hypothesized that differences in TE levels might underlie the immune phenotypic transition from T cell-rich to T cell-excluded. To examine this hypothesis, we quantified the spatially-resolved expression of TEs[66] (Fig. 5h and Methods). Our analysis revealed that two CRC-specifically upregulated TEs-the internal sequence of human endogenous retrovirus subfamily H (*HERV-H*) and its associated long terminal repeat *LTR7Y*-were downregulated in clone3 with chr20q-gain compared to clone1 and clone2[67,68] (Fig. 5i, j, Supplementary Fig. 80d, and Methods). These findings suggest that *HERV-H* downregulation could be employed by tumor cells as a mechanism to evade immune recognition in CRC. Correspondingly, we also noted decreased expression levels of *HERV-H* and attenuated signaling activity in viral transcription and type I IFN production in tumor samples with chr20q-gain relative to chr20q-WT in TCGA-COAD (Fig. 5k, l, Supplementary Fig. 80e, and Methods).

Considering the association of chr20q-gain with T cell excluded clonalTIME in CRC, and the potential impact of T cells on clinical response to immune checkpoint blockade (ICB) treatment, we hypothesized that chr20q-gain might influence the response to ICB treatment in CRC patients. To examine this, we evaluated survival post-ICB treatment by categorizing patients from Memorial Sloan Kettering Cancer Center (MSKCC) IMPACT (MSK-IMPACT) cohorts (comprising 71 patients) into two groups based on their chr20q status[69,70]. Notably, we observed a correlation between the presence of chr20q-gain and poorer survival in patients undergoing ICB, which was not evident in CRC patients who did not receive ICB treatment (TCGA-COAD dataset) (Fig. 5m, and Supplementary Fig. 80f, g). To rule out the potential impact of tumor mutation burden (TMB), we categorized patients based on both CNV status and TMB. Our analysis indicated that the correlation between chr20q-gain and diminished survival in the CRC ICB cohort remained irrespective of TMB levels (Supplementary Fig. 69h). Taken together, we identified chr20q-gain, a prevalent genomic defect in CRC, as a genomic marker for T cell exclusion, suggesting its utility as an immunogenomic parameter for predicting responsiveness to ICB. Furthermore, we propose a potential model wherein tumor cells orchestrate a T cell-excluded clonalTIME by suppressing *HERV-H* expression during clonal evolution.

Discussion

Most current spatial deconvolution methodologies excel in decomposing well-structured tissues like the mouse brain[9-13,16]. However, unlike these structured tissues, sublayers within complex microenvironments typically exhibit subtle differences, often with low abundance and high

heterogeneity. Here, we introduce UCASpatial, a robust, sensitive, and versatile algorithm designed to deconvolute ultra-resolution spatiotemporal cellular composition. Additionally, we introduce clonalTIME, a downstream function tailored to analyze UCASpatial results within the TIME. These tools are seamlessly integrated into the Seurat framework, facilitating a broad range of applications.

UCASpatial shares conceptual similarities with existing methodologies. For instance, SPOTlight utilizes a seeded topic model and a modified non-negative least squares (NNLS) approach to estimate cell type compositions[9], while stereoscope employs a model-based probabilistic method to estimate both cell type signatures and relative cell abundance across spots. RCTD[10] and cell2location[11] utilize statistical models (Poisson distribution and Bayesian model, respectively) to estimate cell abundance across spots, while CARD[12] incorporates a conditional autoregressive (CAR) assumption into the NMF model, accommodating spatial correlation in cell-type composition across spots. However, specific meta-purity filtering and entropy-based weighting mechanisms in UCASpatial offer distinct advantages, as evidenced by both our in-silico benchmarking results and analysis of real ST datasets (Supplementary Figs. 2-7, 16, 66-74). The meta-purity filtering design in UCASpatial filters out intermediate cells with transcriptomes similar to other cell types, thereby enhancing the representativeness of each cell subpopulation[71-73]. Furthermore, through entropy-based weighting, UCASpatial enhances the discrimination of fine cell subpopulations with subtle transcriptomic differences by quantifying the ability of each gene to indicate cell identity and incorporating this information in the deconvolution process.

Moreover, UCASpatial demonstrates versatility across platforms for spatial transcriptomic sequencing. To illustrate, we applied UCASpatial to mouse hippocampus data from the Slide-seqV2 platform[12,74], exemplifying its adaptability. As anticipated, UCASpatial accurately delineated the three anatomical structures of the hippocampus: the cornu ammonis 1 (CA1)/CA2 region, the CA3 region, and the dentate gyrus (Supplementary Fig. 81). Specifically, CA1 cells predominantly localized in the CA1 region, CA3 cells exhibited high enrichment in CA3, while dentate cells were situated in the dentate gyrus forming a C-shaped ring. Additionally, ependymal cells, characterized by their irregular and columnar shape[75], lined the brain ventricles, whereas choroid cells resided beneath the ependymal cells and were found in the choroid plexus[76] along with Cajal-Retzius cells[77] (Supplementary Fig. 81).

We notice that UCASpatial does not incorporate neighboring or image information, a strategy commonly employed previously to enhance performance[12,29,30]. However, integrating such information might not necessarily enhance the annotation of infiltrated immune cell

subpopulations with low proportions in complex microenvironments. This is due to the discrete distribution of these subpopulations resulting from recruitment dynamics. Moreover, the efficacy of UCASpatial is contingent upon the quality of the referenced scRNA-seq data. By leveraging unpaired scRNA-seq data as a reference, this limitation could potentially be mitigated. Additionally, establishing a high-quality scRNA-seq database encompassing various common tissues could address this concern.

Spatially inferred CNVs within ST provide insights into the tumor genomics landscapes at a clonal resolution. However, linking tumor genomics and diverse immune landscapes at such detailed clonal resolution remains constrained by the deciphering of the TIME. UCASpatial enables ultra-resolution analyses of TIMEs and provides an immunogenetic platform for addressing this challenge. By applying UCASpatial in human CRC, we identified a prevalent CNV, chr20q-gain, correlating with a T cell-excluded immune phenotype and poor responsiveness to ICB, which had not been previously reported. Additionally, considering the significant correlation of the chr20q-gain with the p53 mutation in CRC, and the potential effect of the p53 mutation in TE activation[78-80], we hypothesize that patients with a concurrent p53 mutation and chr20q-WT might have better responsiveness to ICB in CRC. Furthermore, extending UCASpatial to other cancer types could help identify novel CNV markers that affect clonalTIMEs or predict the efficacy of specific immunotherapies.

In future iterations, enhancing UCASpatial could involve integrating whole exome sequencing or whole genome sequencing to offer a more comprehensive elucidation of the co-option between tumor mutations and spatially-resolved clonalTIMEs. Additionally, employing spatial cellular deconvolution alongside cell interaction analysis holds promise for unraveling the spatiotemporal dynamics across multiple biological processes and understanding how they are modulated by external cues. Similarly, gaining insights into the perturbations of the microenvironment in response to disease can deepen our comprehension of pathological mechanisms and aid in the development of therapeutic strategies.

In conclusion, we anticipate that UCASpatial will find myriad applications in deciphering complex microenvironments across various developmental stages, states of health, and disease conditions within a spatial framework.

Methods

UCASpatial model.

In brief, UCASpatial is an entropy-weighted model that annotates the cell-identity (equal to cell-type) compositions on each spatial location by deconvoluting the spatial expression count

matrix into a predefined set of reference cell types. The model requires both spatial
transcriptomics data and paired or unpaired scRNA-seq data from a given tissue as input. By
default, the model takes an untransformed spatial-expression count matrix $\mathbf{P} \in \mathbb{R}^{S \times G}$ of genes
$g = \{1,..,G\}$ at spots $s = \{1,..,S\}$ as input, as, for example, obtained from the 10x Space
Ranger software (10x Visium data). The scRNA-seq data $\mathbf{B} \in \mathbb{R}^{G \times N}$ serve as a reference and
consist of N cells with the same set of G genes with $\mathbf{P}$. All the N cells will be classified
into K cell types which need to be provided by the users.

We denote $\mathbf{X} \in \mathbb{R}^{S \times K}$ as the cell-type composition matrix, where each row x_s represents
the proportions of the K cell types on spot s. By default, this matrix is estimated using an
entropy-weighted non-negative least square model, which allows for accurate and robust
annotation in complex situations. (see below).

Meta-purity filtering

The existence of intermediate cells blurs cell-type borders due to their mixed identities from
multiple cell types, which will disturb the recognition of cell-identity specific gene expression
profiles (cGEPs). To remove the effect brought by intermediate cells in the reference scRNA-
seq data, we designed the meta-purity filtering strategy as an optional preprocessing.

Based on the input scRNA-seq reference matrix $\mathbf{B}$, we construct meta cells using the
function 'FindClusters' from R package 'Seurat' with a default resolution of 100. We
recommend the resolution parameter by an expectation of an average of 100 cells per meta-cell,
which serves as a parameter selection criterion. Using pre-existing cell-type labels, we define
the purity of meta-cells by $pu(MC_i) = \max_{j \le K}(\dfrac{n_{ct_j}}{n_{MC_i}})$, where MC_i denotes the ith meta cell,
$pu(MC_i)$ denotes the purity of MC_i, n_{ct_j} denotes the cell number of cell type j in MC_i, and
n_{MC_i} denotes the cell number of MC_i. The cells contained in meta cells with purity below the
given threshold (default 0.95) will be filtered out.

Down sampling

529 Another optional preprocessing for referenced scRNA-seq is down sampling. By default, we
530 will randomly downsample the scRNA-seq reference matrix $\mathbf{B}$ to 100 cells for each cell type
531 based on a given random seed to improve the speed for downstream analysis. Thus, the
532 processed scRNA-seq expression matrix $\widetilde{\mathbf{B}} \in \mathbb{R}^{G \times \widetilde{N}}$ has been established, in which
533 $\widetilde{N} = 100 \times K$ by default.

Optimized non-negative matrix factorization (NMF).

535 In order to efficiently extract the cGEP matrix $\mathbf{E} \in \mathbb{R}^{G \times K}$, we utilize non-negative matrix
536 factorization (NMF) on the processed scRNA-seq expression matrix $\widetilde{\mathbf{B}}$. Based on the provided

cell-type labels, we initialize $\mathbf{H} \in \mathbb{N}^{K \times \tilde{N}}$ as a one-hot matrix representing the identity of each cell, thus guiding it towards a biologically relevant result. This initialization also aims at reducing variability and improving the consistency between runs. We iterated through NMF to estimate the cGEP matrix $\mathbf{E}$:

$$\tilde{\mathbf{B}} \sim \mathbf{E} \times \mathbf{H} \tag{1}$$

where each element in the matrix $\mathbf{E}$ and $\mathbf{H}$ is constrained to be non-negative.

Entropy-based cGEPs weighting

Considering that the indicative capacity for cell types of each gene is unequal, quantifying the indicative capacity of each gene and taking it into account in the decomposition step would benefit from distinguishing the heterogenous cell states with subtle transcriptome differences. Shannon entropy for the expression percentage of each gene across different cell types can be a great quantification for the indicative capacity of each gene. Thus, we formulate Shannon entropy-based weights for each gene g of the extracted cGEP matrix $\mathbf{E}$ to highlight genes indicative of cell identities.

UCASpatial calculates the expression percentage matrix $\mathbf{PCT} \in \mathbb{R}^{K \times G}$ based on the processed scRNA-seq expression matrix $\tilde{\mathbf{B}}$:

$$\mathbf{PCT}_{k,g} = \frac{\text{num}(\tilde{\mathbf{B}}_{g,n \in k} > 0)}{\text{num}(\tilde{\mathbf{B}}_{g,n \in k})} \tag{2}$$

where $\mathbf{PCT}_{k,g}$ denotes the element of $\mathbf{PCT}$ representing cell type k for gene g, $\text{num}(\tilde{\mathbf{B}}_{g,n \in k})$ represents the number of cells in cell type k in $\tilde{\mathbf{B}}$, $\text{num}(\tilde{\mathbf{B}}_{g,n \in k} > 0)$ represents the number of cells in cell type k with gene g expression over 0. The expression percentage matrix $\mathbf{PCT}$ will be further normalized into $\widetilde{\mathbf{PCT}}$ via each element $\mathbf{PCT}$ being divided by the sum of the corresponding row:

$$\widetilde{\mathbf{PCT}}_{k,g} = \frac{\mathbf{PCT}_{k,g}}{\sum_{i=1}^{K} \mathbf{PCT}_{i,g}} \tag{3}$$

Based on the $\widetilde{\mathbf{PCT}}$ matrix, UCASpatial further calculates the Shannon entropy for each gene:

$$\text{Entropy}(g) = -\sum_{i=1}^{K} \widetilde{\mathbf{PCT}}_{i,g} \times \log(\widetilde{\mathbf{PCT}}_{i,g}) \tag{4}$$

For better understanding, UCASpatial provides an adjusted entropy using the maximum entropy minus the entropy for each gene g_0:

$$\text{adj.ent}(g) = Ent.maxm - \text{Entropy}(g) \tag{5}$$

where $Ent.maxm = \sup(\text{Entropy}(g)) = -\sum_1^K \frac{1}{K} \times \log(\frac{1}{K})$ represents the supremum of

$\text{Entropy}(g)$. The maximum entropy $Ent.maxm$ is only related to the number of cell types K.

The adjusted entropy could excellently represent the expression specificity for genes, with a

higher adjusted entropy representing a higher indicative capacity for cell types of a gene.

Considering that the adjusted entropy can only evaluate the expression specificity of each

gene, we thus took not only the adjusted entropy but also the expression percentage and

expression level into account, designing the entropy-based weight of each gene as:

573
$$\omega(g) = \max_{i \in \{1,2,\dots,K\}} (\frac{2 \times \text{adj.ent}(g)}{Ent.maxm} \times 2 \times \widetilde{\mathbf{PCT}}_{i,g} \times \text{avg\_logFC}(i,g)) \tag{6}$$

574
$$\omega(g) = \begin{cases} \omega(g), & \omega(g) \le 1 \\ 1, & \omega(g) > 1 \end{cases} \tag{7}$$

where $\text{avg\_logFC}(i,g)$ denotes the average of $\log_2$ fold-change expression for gene g in cell

type i, $\omega(g)$ denotes the entropy-based weight of gene g which is constrained to be non-

negative. UCASpatial also combined common marker finding methods including the

FindAllMarkers function in Seurat (version 3.1.4) and cosine similarity method[81] (version

0.9.0), for optional gene filtering in cGEPs, which can further improve the indicative capacity

of genes in cGEPs.

Entropy-weighted non-negative least square model.

582 Taking the untransformed spatial expression count matrix $\mathbf{P}$ as input, this step aims to

583 decompose the RNA counts of each spot into referenced cell type composition $\mathbf{X} \in \mathbb{R}^{S \times K}$

584 based on $\mathbf{E}$. Unit variance normalization by gene is performed in $\mathbf{P}$ in order to standardize

585 discretized gene expression levels[5,82]. Commonly, this step will utilize the non-negative least

586 square (NNLS) model for each spot s:

587
$$\mathbf{P}_s = \mathbf{X}_s \times \mathbf{E} + \varepsilon_s \tag{8}$$

588 where $\mathbf{P}_s$ denotes the spatial expression count vector of spot s, $\varepsilon_s \in \mathbb{R}^{S \times 1}$ represents the

589 residual error of spot s which is restricted to being non-negative. By minimizing the loss

590 function for each spot s:

591
$$\min L(\varepsilon) = \min \|\varepsilon_s\|_2^2 = \min_{\mathbf{X}_{s,k}} \sum_{k=1}^K \sum_{g=1}^G (\mathbf{P}_{s,g} - \mathbf{X}_{s,k}\mathbf{E}_{g,k})^2 \tag{9}$$

where $\mathbf{X}_{s,k}$ denotes the cell type composition of spot s, with each element restricted to be non-

negative; $\mathbf{P}_{s,g}$ denotes the spatial expression count of gene j in spot s; ε_s denotes the residual

error between the deconvoluted expression vector $\mathbf{X}_{s,k}\mathbf{E}_{g,k}$ and true spatial expression $\mathbf{P}_{s,g}$

where $j \in \{1, 2, \ldots, G\}$. By minimizing the loss function $L(\varepsilon) = \|\varepsilon_s\|_2^2$, we can obtain the $\mathbf{X}_{s,i}$

which can best mimic the real cell type composition of spot s under the given cell identities.

To further take the entropy-based weight into account, we extended the common NNLS

into an Entropy-weighted Non-Negative Least Squares (ENNLS) model. We first extend the

weight vector $\omega(g)$ to a diagonal matrix $\mathbf{\Omega} \in \mathbb{R}^{G \times G}$:

600
$$\mathbf{\Omega} = \begin{pmatrix} \omega_1 & 0 & \cdots & 0 \\ 0 & \omega_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \omega_G \end{pmatrix} \tag{10}$$

601   the corresponding ENNLS model has been extended into:

602
$$\mathbf{P}_s \sqrt{\mathbf{\Omega}} - \mathbf{X}_s \mathbf{E} \sqrt{\mathbf{\Omega}} \tag{11}$$

603   The loss function of the ENNLS model weighted by the diagonal matrix $\mathbf{\Omega}$ is:

604
$$\min \widetilde{L}(\varepsilon) = \min \left\| \widetilde{\varepsilon}_s \right\|_2^2 = \min_{\mathbf{X}_{s,k}} \left\| \mathbf{P}_s \sqrt{\mathbf{\Omega}} - \mathbf{X}_s \mathbf{E} \sqrt{\mathbf{\Omega}} \right\|_2^2$$
$$= \min_{\mathbf{X}_{s,k}} \sum_{k=1}^{K} \sum_{g=1}^{G} (\mathbf{P}_{s,g} \sqrt{\mathbf{\Omega}_{g,g}} - \mathbf{X}_{s,k} \mathbf{E}_{g,k} \sqrt{\mathbf{\Omega}_{g,g}})^2 \tag{12}$$

where each element in $\mathbf{X}_{s,i}$ is restrained to be non-negative. Each row of the cell-type

composition spectrum $\mathbf{X}$ is normalized into $[0,1]$ by $\mathbf{X}_{s,k} = \dfrac{\mathbf{X}_{s,k}}{\sum_{i=1}^{K} \mathbf{X}_{s,i}}$.

By minimizing the weighted loss function $\min \widetilde{L}(\varepsilon) = \min \left\| \widetilde{\varepsilon}_s \right\|_2^2$, we can highlight the

contribution of genes indicative of cell identities, and finally calculate the cell-type composition

spectrum $\mathbf{X}$ in a more biological way.

Referring to SPOTlight, by selecting all cells in $\widetilde{\mathbf{B}}$ from the same cell type k, and

computing the median of each column of $\mathbf{E}$ for a consensus cell type-specific topic signature,

we use $\mathbf{H}$ to further correct $\mathbf{E}$.

Additionally, while UCASpatial is implemented in R, its deconvolution algorithm NMF is

implemented with an efficient C++ code that is linked back to the main functions of UCASpatial

through Rcpp, ensuring scalable computation.

Finally, considering that the number of cells in each spot is generally within a certain range,

for example, 5-20 cells for 10x Visium, we believe that a proportion less than the threshold will

be noise in the calculation process. By default, we use a less stringent filter of 0.01 to remove

potential noise in the cell-type composition $\mathbf{X}$.

620	
621	**Simulation strategy and evaluation metrics.**
622	**Simulation strategy**
623	We generated simulated spatial transcriptomic data sets from a reference CRC scRNA-seq data
624	set (Supplementary Figs. 1a-c and 4, see in the *Human specimens (in-house cohort)* part). To
625	simulate different situations of tumor microenvironment, we generated three different regions
626	including tumor core (TC), tumor margin (TM), and tumor stroma (TS)[16,54]. Each region has
627	five replicates with each replicate containing N_{spot} = 50 spots, generated based on random seeds
628	(19980205, 19990911, 19960330, 20000502, 20230509) The number of cells in each spot is
629	restricted to less than 20. To mimic the true situation as much as possible, we also imposed
630	restrictions on the types of cells present in different regions, so that spots from different regions
631	have different cell type co-localization patterns. Detailed restrictions for regional simulation are
632	listed in Supplementary Table 1. To better mimic the real capture, if the synthetic spot had over
633	25,000 UMI counts, it would be randomly down-sampled into 20,000 UMI counts.
634	We applied different methods (see in the *Compared methods* part) to deconvolute the above
635	datasets. In each analysis, we provided identical simulated spatial data and reference scRNA-
636	seq data as input for all methods. To ensure an equitable assessment of algorithm performance,
637	we adopted the identical threshold of 0.05 (established on spots containing up to 20 cells) as
638	UCASpatial for algorithms lacking default filtering thresholds, aiming to eliminate noise in
639	accessing the ability to determine whether individual cell types exist at a spot (i.e., spots with
640	non-zero cell abundance or not).
641	**Construct complexity hierarchy**
642	To evaluate UCASpatial in microenvironments with different complexity, we rank the
643	complexity hierarchy of these simulated spots by calculating their cell-type composition
644	entropy: $\text{Entropy}(s) = -\sum_{i=1}^{K} \mathbf{X}_{s,i} \cdot \log(\mathbf{X}_{s,i})$, where $\mathbf{X}_{s,i}$ denotes the proportion of cell type i in
645	spot s. We classify them into three groups (low, medium, and high), with N_{spot} =250 in each
646	group, where higher entropy indicates a higher level of complexity.
647	**Evaluation metrics**
648	In each simulation replicate, we calculated the true cell-subpopulation proportions on each
649	spatial location as the number of cells in each cell subpopulation divided by the total number
650	of cells in the location. $T_{s,k}$ and $P_{s,k}$ represent the true and predicted proportion of cell
651	subpopulation k, respectively, in the sth spot through the number of total cell subpopulations K.
652	To evaluate the performance of the methods comprehensively, the main benchmark metrics

653 used were PCC, RMSE, and F1 score. The definitions of these metrics are provided below.

654 *PCC.* The cell subpopulation-level PCC value was calculated between the predicted proportion

655 of the specific cell subpopulation P_k and the ground truth proportion of the specific cell

656 subpopulation T_k using the following equation:

$$657 \quad \text{PCC}_{\text{cell}}(P_k, T_k) = \frac{\mathbb{E}[P_k, T_k] - \mathbb{E}[P_k]\mathbb{E}[T_k]}{\sqrt{\mathbb{E}[P_k^2] - (\mathbb{E}[P_k])^2}\sqrt{\mathbb{E}[T_k^2] - (\mathbb{E}[T_k])^2}} \tag{13}$$

658 The spot level PCC value was calculated between the predicted spot composition of P_s

659 and the ground truth spot composition T_s using the following equation:

$$660 \quad \text{PCC}_{\text{spot}}(P_s, T_s) = \frac{\mathbb{E}[P_s, T_s] - \mathbb{E}[P_s]\mathbb{E}[T_s]}{\sqrt{\mathbb{E}[P_s^2] - (\mathbb{E}[P_s])^2}\sqrt{\mathbb{E}[T_s^2] - (\mathbb{E}[T_s])^2}} \tag{14}$$

661 *RMSE.* The cell subpopulation-level RMSE was calculated between the predicted proportion of

662 the specific cell subpopulation P_k and the ground truth proportion of the specific cell

663 subpopulation T_k using the following equation:

$$664 \quad \text{RMSE}_{\text{cell}}(P_k, T_k) = \sqrt{\frac{1}{S}\sum_{s=1}^{S}(T_{s,k} - P_{s,k})^2} \tag{15}$$

665 The spot level RMSE value was calculated between the predicted spot composition of P_i

666 and the ground truth spot composition T_i using the following equation:

$$667 \quad \text{RMSE}_{\text{spot}}(P_s, T_s) = \sqrt{\frac{1}{K}\sum_{k=1}^{K}(T_{s,k} - P_{s,k})^2} \tag{16}$$

F1 score. F1 score was selected to evaluate the ability of alternative methods to determine the

presence of individual cell types at specific locations, i.e., with non-zero cell abundances, based

on the confusion matrix. A confusion matrix is a table used in the field of machine learning and

statistics to assess the performance of a classification model. It summarizes the performance of

a classification algorithm by showing the counts of true positive (TP), true negative (TN), false

positive (FP), and false negative (FN) predictions for each class in a dataset.

$$674 \quad \text{F1 score} = \frac{2 \cdot \frac{TP}{TP + FP} \cdot \frac{TP}{TP + FN}}{\frac{TP}{TP + FP} + \frac{TP}{TP + FN}} \tag{17}$$

Compared methods.

We compared UCASpatial with six deconvolution methods: (1) SPOTlight (version 0.1.7), (2)

RCTD (version 1.2.0) (3) cell2location (version 0.8a0), (4) CARD (version 1.0) and (5)

stereoscope (v 0.3.1). Results of the stereoscope were computed with parameters "-sce 25000

-n 5000 -ste 25000 -stb 500 -scb 500 -gp". For other methods, we followed the tutorial on the

corresponding GitHub pages and used the recommended default parameter settings for

deconvolution analysis.

Experimental and Analysis Approach to Mouse Ear

Ear wound creation

All mice utilized in the wounding experiments were aged between 8-10 weeks. Mice were

subjected to short-term anesthesia induced by isoflurane prior to wound generation. The dorsal

and ventral aspects of mouse ears were sanitized with cotton balls saturated in 75% alcohol.

Mouse ears were delicately flattened using forceps, and a tissue-penetrating ear puncher (2 mm)

was used to create perforating injuries within the vascularized regions of the ear. Hemostatic

powder was applied to the wounds using disposable cotton swabs. All instruments were

sterilized under high pressure before use.

Single-cell RNA-seq

Single-cell suspensions were obtained from MRL mouse ear tissue at day 7 post-wounding.

Specifically, mouse ear tissues were mechanically minced and then incubated at 37°C for 5-10

minutes with 40 µL of 50 mg/mL collagenase II and 40 µL of 50 mg/mL collagenase IV.

Following a brief low-speed centrifugation to precipitate undigested tissue fragments, the

supernatant was transferred into a stop solution composed of fetal bovine serum, EDTA, and

D-HBSS. Additional digestion enzyme was added to the partially digested tissue fragments, and

the digestion process was repeated until complete tissue digestion was achieved. The digested

mixture was filtered through a 70 µm membrane and centrifuged at 4°C at 1200 rpm for 10

minutes. After discarding the supernatant, the cell pellet was resuspended in FACS buffer for

cell sorting. Calcein and 7-AAD were freshly added before cell sorting. Then FITC$^+$ 7-AAD$^-$

single cells were sorted for library construction of scRNA-seq. The libraries were performed

using Chromium Next GEM Single Cell 3' GEM, Library & Gel Bead Kit V3 (PN-1000075)

purchased from 10x Genomics according to the protocols provided by manufacturers. The

libraries were sequenced on an Illumina Novaseq 6000 platform.

Spatial gene expression assay

Fresh mouse ear tissues were embedded in the Optimal Cutting Temperature Compound (OCT,

Sakura Tissue-TEK), and stored at -80°C. The 10-µm section was placed on the pre-chilled

Optimization slides (Visium, 10x Genomics, PN-1000193). After collecting histological

morphological information, the tissue was processed within the optimal permeabilization time

of 18 minutes to digest and release mRNA from tissue slides. The released mRNA was

subsequently captured by proximal oligonucleotides within the tissue, culminating in the

generation of the respective libraries. The libraries were sequenced on an Illumina Hiseq X Ten

platform.

Single-cell RNA-seq data processing

10x Genomics scRNA-seq data were aligned and quantified using the Cell Ranger Single-Cell

Software Suit (version 3.0.2) against the mm10 reference genome. Genes with less detected cell

numbers (less than 3) and cells with less detected genes (less than 200) were removed. To

further filter out low-quality data, the following criteria were applied to filter cells: (1) the

number of expressed genes greater than 500 or less than 5,000; (2) mitochondrial proportion

less than 20%. To cluster single cells by their expression, the unsupervised graph-based

clustering algorithm implemented in the Seurat R package (version 3.1.4) was used. The top

2,000 variable genes identified by the vst method were used to perform principal component

analysis. The RunUMAP and RunTSNE functions were employed to generate UMAP and t-

SNE dimensionality reduction results, while the FindNeighbors function was used to construct

a shared nearest neighbor graph. DoubletFinder was utilized to predict potential doublets and

filter them out.

Spatial transcriptomics data processing

Using the Space Ranger software (https://support.10xgenomics.com/spatial-gene-

expression/software/pipelines/latest/tutorials), the 10x Visium spatial transcriptomics data was

aligned and quantified against the mm10 version of the mouse reference genome. Downstream

analysis was conducted using the R package Seurat. Integration of unpaired scRNA-seq data

with spatial transcriptomics sequencing data was performed for deconvolution analysis to

obtain the proportion of each cell subtype within each spot of the spatial transcriptome.

Parameters for UCASpatial: downsample_n = 1000, meta.resolution = 200, meta.filter = 0.95,

cos.filter.threshold = 0.1, the others using default. Parameters for other methods: Default. Using

the threshold of 3% to filter the noise after decomposition by each method (Supplementary Figs.

10-16).

Spatial transcriptomics data visualization

The original data placed mouse ear tissues on two slides: tissues from day 0, day 7, and day 15

were on the first slide, and tissues from day 3 were on the second slide. To better visualize the

spatial transcriptomics data of mouse ears, we merged all the tissues into one slide and aligned

them based on the time point post-wounding. Specifically, we first extracted the coordinates of

each spot in all 4 tissues. We then transformed the coordinates of spots from day 3, with each

row replaced by the col, and each column replaced by the row. We next transformed the

coordinates of each spot's cols abiding by the time points. ggplot2 was used for visualizing

spatial transcriptomics data after coordinate transformation.

Experimental and Analysis Approach to Human CRC

Tissue dissociation of patient samples

Fresh human tumor specimens and patient-matched adjacent normal tissues were washed twice with DPBS (gibco) and then mechanically disrupted into approximately 1 mm^3 pieces in the RPMI-1640 (gibco) and further enzymatically digested with Human Tumor Dissociation Kit (Miltenyi Biotec, 130-095-929) in C-tube in combination with gentleMACS™ Dissociator (Miltenyi Biotec, 130-093-235) presetting program h_tumor_01, h_tumor_02 and h_tumor_03 sequentially according to the manufacturer's instruction. Between each program, samples were incubated for 30 minutes at 37°C under continuous rotation using a rotator. The dissociated cells were subsequently passed through a 70-μm cell strainer (Biologix, 15-1070) and centrifuged at 300g for 7 min. After the supernatant was removed, the cell pellet was washed with DPBS. The pelleted cells were resuspended in a cell cryopreservation medium (FBS with 10% DMSO) at a concentration of $1×10^6$ cells/ml and further cryopreserved in liquid nitrogen for long-term storage.

Single-cell RNA-seq for patient samples

On the day of scRNA-seq droplet capture, cells were thawed rapidly in a 37°C-water bath and washed off DMSO in DPBS. Palleted cells were resuspended in a sorting buffer (PBS with 2% FBS and 2 mM EDTA). Cells were stained with CD45-FITC (Biolegend, 368508) for 20 minutes at 4°C in the dark. 7-AAD was freshly added prior to FACS sorting performed on the BD FACS Aria III instrument. Based on FACS analysis, 7-AAD$^-$CD45$^+$ cells and 7-AAD$^-$CD45$^-$ cells were sorted into 1.5ml tubes (ExCell Bio, CS015-0041) and counted manually under the microscope (Leica), respectively. Libraries were generated using Chromium Next GEM Single Cell 5' Library & Gel Bead Kit v1.1 (10x Genomics, PN-1000165) according to the manufacturer's protocol. Purified libraries were sequenced by the Illumina NovaSeq 6000 platform with 150-bp paired-end reads.

10x Visium spatial transcriptomics

CRC samples were frozen and embedded in OCT (Sakura Tissue-Tek). All tissue blocks were sectioned using the Leica CM1950 cryostat and were cut at 10 μm. Samples were selected on the basis of morphology, orientation (based on H&E staining), and RNA integrity number (RIN) that was obtained using Agilent 2100 Bioanalyzer. The RIN of selected samples ranged from 4.5 to 8.3. The sections were treated as recommended by 10x Genomics and the optimization procedure showed the optimal permeabilization time determined as 12 min for CRC samples using the Visium Spatial Tissue Optimization Kit (10x Genomics, PN-1000193). Spatial gene

expression slides were equilibrated inside the cryostat at -20℃ before sectioning and all of the

reagents used for spatial transcriptomics were from the Visium Spatial Gene Expression Kit

(10x Genomics, PN-1000187). According to the manufacturer's instruction, the tissue sections

were fixed in chilled methanol at -20℃ for 30 min and stained with H&E. Brightfield

histological images were performed on the Olympus Microscopes BX53 at ×20 magnification

and then taken by cellSens Dimension (version 1.18) software. Images were exported as tiled

tiffs for analysis. After imaging, tissue sections were permeabilized at 37℃ for optimal

permeabilization time of release of mRNA as mentioned above. Following permeabilization,

mRNA was reverse-transcribed and underwent second-strand synthesis, denaturation, cDNA

amplification, and cleanup by SPRIselect beads sequentially (Visium, 10x Genomics, User

Guide CG000239). cDNA amplification cycles were determined by quantitative PCR (qPCR).

Library quality was evaluated using TapeStation (Agilent Technologies). Sequencing was

performed on the Illumina Hiseq X Ten platform.

Human specimens (in-house cohort)

One male (CRC3) and two female (CRC1 and CRC2) patients were enrolled and pathologically

diagnosed with colon cancer at Peking University Cancer Hospital & Institute (Supplementary

Table. 3). All patients provided written informed consent. This study was approved by the

Institutional Ethics Committee of Beijing Cancer Hospital (2023YJZ16). Among these patients,

CRC1 was diagnosed as stage II, and CRC3 and CRC2 were diagnosed as stage IV according

to the guidance of AJCC version 8. Fresh tumor tissues and patient-matched normal adjacent

colon tissues were collected from surgery resection, which was never treated with

chemotherapy or radiation therapy prior to the study. The resected samples were stored in RPMI

1640 (gibco) added with Amphotericin B (Shelleck, S1636) and Gentamicin Sulfate (gibco,

15710-064) and transported at 4 ˚C immediately. A small part of each tissue was frozen and

embedded in OCT (Sakura Tissue-Tek) and the remainder of the same tissue was dissociated to

a single-cell suspension.

Single-cell RNA-seq data processing and annotation

In-house single-cell RNA-seq data from 10x Genomics were processed using Cell Ranger

(version 6.0.1), aligning to the hg38 reference genome. Low-quality cells were filtered out if

they expressed fewer than 500 or more than 7,000 genes. Cells with mitochondrial gene

expression exceeding 20% were also excluded. Batch effects of in-house and public scRNA-

seq datasets were corrected using the integration workflow in the Seurat R package[83] (version

3.1.4). The top 2,000 variable features were used to identify anchors using the

FindIntegrationAnchors function, and data from in-house and public scRNA-seq datasets were

integrated using the IntegrateData function.

To identify major cell types, the top 50 PCs were calculated using the RunPCA function, and

the top 20 PCs were used as input to the uniform manifold approximation and projection

(UMAP) algorithm (RunUMAP). Nearest neighbor graphs were constructed using the

FindNeighbors function, and clusters were identified using the Louvain function (resolution =

0.6). Cell types were annotated based on well-known marker genes.

A second round of clustering was performed to further characterize subpopulations within

each cell type. Due to the variability in cell number and features among cell types, T&NK,

myeloid, and epithelial cells were re-integrated using the top 2,000 variable features and re-

clustering using different parameters, respectively. NK and T cells were clustering (resolution

= 0.4) on the basis of the top 15 PCs. Myeloid cells were clustering (resolution = 0.4) on the

basis of the top 30 PCs. Epithelial cells were clustering (resolution = 0.05) on the basis of the

top 30 PCs. One T cell and myeloid cluster with few specific markers but exhibiting high

expression of cell cycle-related genes was removed, as these genes could not be fully regressed

out after cell cycle score scaling. In total, we defined 34 cell types in CRC, and this CRC

scRNA-seq reference was used throughout the manuscript to maintain consistency in presenting

spatial mapping results (Fig. 4b, Supplementary Fig. 17b, c).

Spatial transcriptomics data processing

Raw sequencing reads of spatial transcriptomics were quality-checked and mapped by Space

Ranger v1.3.0. The gene-spot matrices generated after ST data processing from ST and Visium

samples were analyzed with the Seurat package in R. The processed data were used for

downstream analysis of UCASpatial (meta.resolution = 300, meta.purity = 0.9, other

parameters using default) and other methods (using default parameters). Using the threshold of

3% to filter the noise after decomposition by each method (Supplementary Figs. 18-73).

Gene signature score calculating in ST

Select the top 20 genes of each cell subpopulation provided in Supplementary Table. 2 or

Supplementary Table. 5 as the signature gene list for murine wound healing ST dataset and

human CRC ST dataset, respectively. Use the "AddModuleScore" method in the Seurat R

package to calculate the signature score of each cell subpopulation in the ST dataset using the

signature gene list.

Gene co-expression module analysis

To analyze the gene program pattern in macrophage subpopulations, the correlations of the top

2000 variance genes obtained from single-cell clustering step by the Seurat R package were

calculated with the Pearson method. To identify the gene co-expression module, a hierarchical

clustering method with complete distance was used for the correlation matrix of top variance

genes. Seven co-expression modules were obtained (Supplementary Fig. 76d-f). Gene

expression data were normalized by the "LogNormalize" method in the Seurat R package. The

z-score transformed normalization data were then used to calculate the correlations of top

variance genes. The score was calculated by averaging the normalized expression of genes in

each set.

Tumor clonal inference and clonalTIME quantification using UCASpatial-clonalTIME

To infer tumor clonal diversity within each spatial transcriptomics section, CNVs were

evaluated by ordering genes according to their chromosomal location and applying a moving

average to the relative expression values. Genes were arranged according to their genomic

localization using the SpatialInferCNV R package[46]. CNVs in high epithelial spots (overall

epithelial proportion > 40%) were estimated using reference spots containing few or no

malignant cells selected from UCASpatial annotated remaining spots (overall epithelial

proportion < 40%). Considering the non-dispersed spatial distribution of epithelial cells,

scattered high epithelial spots (neighboring same-type spots < 1, step = 1) were considered

noise and defined as remaining regions. The spot-resolution CNV profiles for each spatial

transcriptomics section were then estimated using the SpatialInferCNV workflow (cutoff = 0.1,

cluster_by_groups = FALSE, denoise = TRUE, HMM = FALSE), excluding chrM, chrX and

chrY genes. To further define tumor regions and infer tumor clonal diversity based on CNV

data, we constructed dendrogram trees to determine clonal groupings of each spot. Dendrogram

nodes with at least one high-probability CNV region were defined as tumor clones and assigned

a specific identity. Nodes without high-probability CNVs were assigned to normal epithelial or

remaining regions based on their histological structure. The average cell subpopulation

composition across all spots within a clonal region was defined as the cell subpopulation

composition of that clonalTIME (Fig. 4c). To perform a comprehensive comparison of

individual CNVs across clones, we initially identified the CNV locations and statuses of each

clone using an HMM model in SpatialInferCNV (Fig. 5a-c, and Supplementary Fig. 79a). CNV

regions locating on the same chromosome with similar variation patterns in the TCGA-COAD

dataset were merged if their PCC ($\log_2$FC_CNV) > 0.8, where $\log_2$FC_CNV denotes the log2

fold change of the copy number of the tumor over the copy number of normal. These CNVs

were then separated based on the location of the centromere (Supplementary Table. 6).

ClonalTIME phenotyping and identification of multi-cellular community

To classify the clonalTIME phenotype of each tumor clone, the scaled proportion of the T cell

subpopulation of each clone was clustered using the K-means function (centers = 2).

ClonalTIME were classified as T cell-riched or T cell-excluded based on the proportion of T

cell subpopulation within each clustering group (Fig. 4d, and Supplementary Fig. 75b-d). To

determine whether a given cell subpopulation was enriched in a specific clonalTIME phenotype,

we calculated the ratio of observed cell number to expected cell number (Ro/e) for T cell-rich

clonalTIME, where the expected proportion of cell subpopulation was determined from the chi-

squared test. We assumed that a subpopulation was enriched in T cell-rich clonalTIME if Ro/e >

1.2 and enriched in T cell-excluded clonalTIME if Ro/e < 0.8. To identify the cell subpopulation

co-change pattern at a clonal resolution, the correlations of the average proportion of cell

subpopulation in each clone were calculated with the Pearson method. To identify the multi-

cellular community, a hierarchical clustering method with complete distance was used for the

correlation matrix. Four cellular communities were obtained (Fig. 4e, and Supplementary Fig.

76a).

Spatial quantification of TE expression

We quantify the expression of transposable elements (TEs) using spatial transcriptomic data

based on a previously reported algorithm[66]. This algorithm allocates TE reads to TE metagenes

based on the TE type-specific sequence (Fig. 5h). In brief, the genes/TEs and cell count matrix

were generated according to default parameters and normalized using the NormalizeData

function. Clonal-specific TE expression programs were identified by comparing each tumor

clone region using the FindAllMarkers function. The list of CRC-specifically upregulated TEs

was downloaded from previous literature[67]. The *HERV-H* expression level was calculated as the

average expression of previously reported *HERV-H* elements (*HERVH-int*, *LTR7*, *LTR7B*, and

LTR7Y) (Fig. 5j).

Differential expression analysis using spatial transcriptomics

Differentially expressed genes across each clone were analyzed using the Seurat R package

(version 4.0.5). Space Ranger output files were imported, and the data were normalized and

scaled using the default NormalizeData and ScaleData functions. Differential gene expression

analyses were performed comparing groups using the FindMarkers function, with the following

parameters: $\log_2$(fold change) ≥ 0.1 and adjusted P-value ≤ 0.05 (Fig. 5g). The list of CHIP-

seq-based STAT1 target genes was downloaded from previous literature[84].

Functional enrichment analysis

The signatures of macrophage subpopulations in murine wound healing and human CRC were

obtained in scRNA-seq data by FindAllMarkers function in the Seurat package with min.pct =

0.25, min.diff.pct = 0.25, logfc.threshold = 0.25 and other default parameters. Metascape was

used to perform functional enrichment analysis for each macrophage subpopulation[85]

(Supplementary Fig. 76f). For differentially expressed genes in the CRC4 section, Gene

Ontogeny (GO) enrichment analysis was performed by clusterProfiler (v4.6.0)[86] (Fig. 5g).

TCGA-COAD data analysis

Gene expression and clinical data for Colon Adenocarcinoma (COAD) were downloaded from

The Cancer Genome Atlas (TCGA). Average copy number data were downloaded from the

UCSC Xena browser (https://xena.ucsc.edu). For CNV analysis, only the samples with CNV

information were obtained. The average $\log_2$FC_CNV > 0.35 was defined as copy number gain,

while the average $\log_2$FC_CNV < -0.35 was defined as copy number loss (Fig. 5f). The

signatures of each cell subpopulation were obtained in CRC single-cell data by the

FindAllMarkers function in the Seurat package with default parameters. The normalized

expression of the top 10 signatures in each cell subpopulation was used to calculate the average

signature score. The reads per million (RPM) of known human endogenous retroviruses (HERV)

in TCGA-COAD were downloaded from a previous report[87,88]. For HERV analysis, samples

without HERV expression information were excluded. Overall *HERV-H* expression was

calculated as the average expression of known transcribed *HERV-H* loci (*HERV-H1-7*)[87] (Fig.

5k). The pathway activity of each sample was calculated using GSVA with default parameter

settings[89] (Fig. 5l). Survival analyses were then performed with the survival R package (version

3.4-0) (Fig. 5m, and Supplementary Fig. 80g).

MSK-IMPACT data analysis

The genomic and clinical data of 71 CRC patients with Immune Checkpoint Blockade (ICB)

therapy were downloaded from published studies[69,70]. The copy number status of each loci was

determined using the GISTIC v2.0 function in GenePattern platform[90,91]. The average

$\log_2$FC_CNV > 0.35 was defined as copy number gain, while the average $\log_2$(copy number/2)

< -0.35 was defined as copy number loss. Survival analyses were then performed with the

survival R package (Fig. 5m, and Supplementary Fig. 80f, h).

Analysis of mouse hippocampus Slide-seq V2 data.

To test UCASpatial's ability in Slide-seq V2 platform, we used UCASpatial (cos.filter.threshold

= 0.1, ent.filter.threshold = 0.2, logfc.threshold = 0.1,min.diff.pct = 0.05, meta.assay = 'RNA')

to spatially annotate cell subpopulations in Slide-seq V2 data of the mouse hippocampus using

published scRNA-seq datasets of mouse hippocampus[10,92] (Supplementary Fig. 81).

Data availability

The sequencing data of both scRNA-seq and spatial transcriptomics of murine wound healing

Code availability

The UCASpatial package and code to reproduce the results in this study are available in the
GitHub repositories at https://github.com/BIGHanLab/UCASpatial.

Acknowledgments

This work was funded by the National Natural Science Foundation of China (NSFC) (32121001
to D.H.), Strategic Priority Research Program of the Chinese Academy of Science
(XDB0570101 to D.H.), Beijing Natural Science Foundation (Z200023 to D.H.)

Author contributions

D.H. and M.M.X. conceived the study and provided overall supervision of the study. D.H.,
M.M.X., and Yin Xu designed the study. Yin Xu designed and developed UCASpatial. Yin Xu
conducted benchmark analyses with assistance from F.Z.. R.C. conducted library construction
for murine scRNA-seq. M.G. and R.C. conducted library construction for murine spatial
transcriptomics. Y.Z. performed bioinformatics analyses for the murine dataset. Yu Xia and
M.G. performed IHC staining of murine ears. Z.W., L.S., and J.L. provided human specimens.
J.Y. organized and summarized the patient information. P.G. and L.D. conducted library
construction for human scRNA-seq and spatial transcriptomics. G.L. preprocessed the human
scRNA-seq data. Z.H. and Yin Xu performed bioinformatics analyses for the human colorectal
cancer dataset. Z.H. performed bioinformatics analyses of TEs. D.H., M.M.X., Yin Xu, Y.Z.,
and Z.H. wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

Main Figures

Figure 1

Fig. 1 | Schematic overview of UCASpatial. UCASpatial takes both referenced single-cell RNA sequencing (scRNA-seq) data with cluster information and spatial transcriptomic data as input. After three-step processing involving cell type-specific GEPs (cGEPs) extracting, entropy-based cGEPs weighting, and deconvolution, UCASpatial decomposes spatially resolved, multi-cell RNA count matrices (i.e., spatial transcriptomic data) into a referenced cell-type composition. GEP, gene expression profile. NMF, non-negative matrix factorization. e-weight, Shannon entropy-based weight. NNLS, non-negative least square. Figure adapted from images created with BioRender.com.

Figure 2

Fig. 2 | Validation of UCASpatial using simulated data. a, Graphic model showing the classification of 750 simulated spots into three groups, each representing a different complexity level - low, medium, and high, respectively. The classification is based on the entropy of cell-type composition for each spot, with each group containing 250 spots. **b**, Box plot showing spot-level root mean square error (RMSE) (left) and F1 score (right) for each method across three complexity levels. Each box plot ranges from the third and first quartiles with the median as the horizontal line, while whiskers represent 1.5 times the interquartile range from the lower and upper bounds of the box. **c**, Line charts showing cell-subpopulation-level Pearson correlation coefficient (PCC) for each method across 20 cell subpopulations. 'avg' and 'maxm' represent the average and maximum PCC, respectively, of the five methods excluding UCASpatial. **d**, Left, line charts representing cell-subpopulation-level PCC of CD8$^+$ effector memory T cells (CD8$^+$ Tem), CD8$^+$ exhausted T cells (CD8$^+$ Tex), CD8$^+$ tissue-resident memory T cells (CD8$^+$ Trm), and natural killer cells (NK) for each method. Right, stacked

barplot showing the accumulated count of false predictions of these cell subpopulations for each

method. Colors represent different cell subpopulations. FP, false positive; FN, false negative. **e**,

The cell-subpopulation-level PCC values for each method were binned into four confidence

groups: 0-0.25 (low), 0.25-0.5 (medium-low), 0.5-0.75 (medium-high), and 0.75-1 (high).

Stacked barplot showing the ratio of four confidence groups for each method. **f**, Graphic model

representing the classification based on the proportion range of a specific cell subpopulation

within the simulated spots. All simulated spots were classified into ten groups from proportion

0-10% to proportion 0-100% for each cell subpopulation. **g**, Left, line charts representing the

average RMSE of 20 cell subpopulations across various proportions for each method. Middle,

line charts showing an example of the RMSE for CD8$^+$ Tex across different proportions. Right,

line charts showing an example of the RMSE for CD8$^+$ Tem across different proportions.

Figures (**a** and **f**) adapted from images created with BioRender.com.

Figure 3

Fig. 3 | UCASpatial reconstructs spatiotemporal micro-anatomy dynamic of diverse macrophage subpopulations during murine wound healing. a, Graphical overview of experimental design and analysis workflow. scRNA-seq was performed using total cells from MRL mice ear tissue on Day 7 post-wounding. Spatial transcriptomics data were generated from MRL mice ear tissue on Day 0 (unwounded), Day 3, Day 7, and Day 15 post-wounding. Figure adapted from images created with BioRender.com. **b**, Uniform Manifold Approximation and Projection (UMAP) plot showing 20 cell subpopulations of all cells from scRNA-seq. **c**,

Left, hematoxylin and eosin (H&E) staining of mouse ear tissues. Scale bar, 500 μm. Middle,

micro-anatomic annotation of wound center on Day 7, the green dashed line represents the edge

of the dermis-epidermis, and the red dashed line represents the blood vessel. Right, Spatial

visualization of keratinocytes, endothelial cells, and fibroblasts at the wound center on Day 7.

Colors represent the proportion of these three cell types. The proportion of fibroblast was

calculated by the sum of 4 fibroblast subpopulations' proportion. **d**, Spatial visualization of the

distribution of indicated macrophage subpopulations at the wound center on Day 7 based on

UCASpatial, CARD, Cell2location, RCTD, SPOTlight, and stereoscope. Colors represent the

proportion of each macrophage subpopulation. **e**, Left, schematic representation of the spot

grouping criteria. Spots were divided into three subdomains: wound center (1 mm radius),

wound proximal (0-1 mm from the wound center), and wound distal (1-2 mm from the wound

center). Right, Heatmap showing the spatiotemporal micro-anatomy dynamics of macrophage

subpopulations of three subdomains. The numbers in the heatmap represent the average

proportion of macrophage subpopulations at specific subdomains and time points. Colors

represent the average proportion for each macrophage subpopulation. The number of UW

(above the heatmap) represents the average proportion of each macrophage subpopulation on

Day 0, respectively. UW, unwounded.

Figure 4

1042

Fig. 4 | Characteristics of clonal-resolution landscape with two TIME phenotypes in human CRC. a, Left, experimental design of in-house colorectal cancer (CRC) dataset. Right, an overview of the CRC dataset used for UCASpatial analysis, with samples organized by

resource. The red, orange, and blue dots represent the presence of spatial transcriptomics,

scRNA-seq, or pathological annotation, respectively. The number in red dots indicates the count

of spatial transcriptomic sections per patient. **b**, UMAP plot displaying 34 cell subpopulations

identified from scRNA-seq reference. **c**, Left, the pathological annotations and indicating

markers' expression (*CD8A*, *GZMK*, *CXCL13*, *XCL1*, and *FAP*) in the CRC3b section. Colors

represent the expression level of each gene. Right, Spatial visualization of the distribution of

indicated CD8$^+$ T cell subpopulations at the CRC3b section based on UCASpatial, CARD,

cell2location, RCTD, SPOTlight, and stereoscope. Colors represent the proportion of each

CD8$^+$ T cell subpopulation. **d**, Schematic representation of UCASpatial-clonalTIME functions.

From left to right: identification of epithelial regions via UCASpatial; delineation of tumor

clone regions and normal epithelial regions using spatialCNV. clonalTIME, clonal tumor

microenvironment. **e**, Heatmap showing the distribution of cell subpopulations across clones

grouped by clonalTIME phenotypes. Colors reflect scaled cell subpopulation proportions

estimated by UCASpatial. Ro/e scores represent the enrichment pattern for each cell

subpopulation between 2 clonalTIME phenotypes. Columns in the heatmap were annotated by

cloneID, sectionID, and clonalTIME phenotype. Ro/e, the ratio of observed over expected cell

subpopulation proportions. **f**, Heatmap showing the PCC between the proportion of each two

cell subpopulations across 13 tumor clones. Communities were identified using hierarchical

clustering. **g**, Boxplot showing the proportion of community3 between T cell-excluded and T

cell-rich clonalTIMEs. Boxplot elements: center line, median; box limits, upper and lower

quartiles; whiskers, 1%–99%. Statistical analysis was performed using a two-tailed unpaired *t*-

test. *$0.01 \leq p < 0.05$. **h**, Left, the spatial distribution of each clone in the CRC4 section. Spots

filled with green, blue, and red represent the tumor region of clone1, clone2, and clone3,

respectively. Dashed lines mark the general contour of each tumor clone region. Middle, barplot

showing the proportion of community3 in each tumor clone of CRC4. Right, spatial

visualization of the proportions for 3 cell subpopulations (CD8$^+$ Tex, CD8$^+$ Tem, and *C1QC*$^+$

macrophage) in community3 estimated by UCASpatial. Figures (**a** and **c**) adapted from images

created with BioRender.com.

Figure 5

1074

associated endogenous retrovirus silencing. a, Heatmap showing copy number variation

(CNV) landscape of each clone grouped by clonalTIME phenotypes. The color indicates the

CNV status (Orange indicates CNV gain, and blue indicates CNV loss). Columns in the

heatmap were annotated by cloneID, sectionID, and clonalTIME phenotype. Forest plots

showing the odds ratio scores associated with clonalTIME phenotypes of each CNV. Dots

indicate the calculated odds ratio and whiskers indicate 95% confidence intervals (CIs). Only

CNVs occurring in at least 2 clones were shown in the heatmap. **b**, Heatmap showing the CNV

profile inferred from spatial transcriptomics data from the CRC4 section. Clonal groupings of

spots were determined by hierarchical clustering. **c**, Phylogenetic tree of three clones in the

CRC4 section. The corresponding clonalTIME phenotype of each clone is listed alongside. The

CNV status of each clone is listed (summarized by the chromosome number where the event

was located, p/q arm, and gain/loss; full genomic locations of each CNV are provided in

Supplementary Table 6). Gray dots represent benign cells, while red dots signify tumor clones.

The intensity of red corresponds to the number of CNVs. **d**, Left, bar plot showing the frequency

of each CNV status in the TCGA-COAD dataset. Right, heatmap showing the PCC between

the copy number of indicated CNV status and signature scores of each cell subpopulation in the

TCGA-COAD dataset. Colors represent the PCC value. Blank squares indicate non-significant

associations (PCC < 0.2). The color of CNV names represents associated clonalTIME

phenotypes in the CRC4 section. **e**, Heatmap displaying the PCC between the copy numbers of

different CNVs status within the TCGA-COAD dataset. Colors represent the PCC value.

Statistical analysis was performed using a two-tailed unpaired *t*-test. **f**, Boxplot showing *CD8A*

expression in samples with indicated CNV status in the TCGA-COAD dataset. **g**, Gene

expression difference between clone3 and other clones in CRC4 section. Left, volcano plot

displaying differentially expressed genes in clone3 versus other clones in CRC4 section.

Differentially expressed genes were defined by adjusted p-value < 0.01 and |log$_2$(fold change)|

$\geq$ 0.1. P-values were calculated using the Wilcoxon rank-sum test and adjusted by the

Bonferroni correction method. Right, functional enrichment analysis of upregulated (n=615)

and downregulated genes (n=1148) of clone3. **h**, Schematic representation of transposable element (TE) quantification in spatial transcriptomics. The resulting assignments are condensed into a matrix of read counts for each cell, versus each TE. Figure adapted from images created with BioRender.com. **i**, Volcano plot showing the differentially expressed TEs between clone3 and other clones in CRC4 section. Only TEs previously identified as being specifically expressed in CRC tumors were considered in this study. Differentially expressed TEs of clone3 were defined by adjusted p-value < 0.01 and a $|\log_2(\text{fold change})| \geq 0.25$. P-values were calculated using the Wilcoxon rank-sum test and adjusted by the Bonferroni correction method. **j**, *HERV-H* expression of each clone in CRC4 section. Left, spatial visualization of *HERV-H* expression in CRC4 section. Dashed lines mark the general contour of each tumor clone region. Right, violin plot displaying the *HERV-H* expression across 3 clones in CRC4 section. Statistical analysis was performed using a two-tailed unpaired *t*-test. **k**, Boxplot representing the *HERV-H* expression across normal tissue, chr20q-WT, and chr20q-gain tumor samples in the TCGA-COAD dataset. **l**, Boxplot displaying the distribution of GSVA activity of the indicated pathways (antigen processing and presentation, type-I IFN production, viral transcription) between chr20q-WT and chr20q-gain tumor samples in the TCGA-COAD dataset. **m**, Prognostic significance of chr20q-gain in ICB-treated CRC patients. Left, overall survival (Kaplan-Meier curves) of CRC patients post ICB in the MSK-IMPACT cohort stratified by the levels of chr20q CNV status. Right, overall survival (Kaplan-Meier curves) of CRC patients without ICB treatment in the TCGA-COAD dataset, stratified by the levels of chr20q CNV status. Log-Rank p-values are shown. Boxplot elements: center line, median; box limits, upper and lower quartiles; whiskers, 1%–99%. Statistical analysis was performed using a two-tailed unpaired *t*-test. $*0.01 \leq p < 0.05$ $**p < 0.01$ $***p < 0.001$ $****p < 0.0001$.

Reference

1126

1   Brbic, M. *et al.* Annotation of spatially resolved single-cell data with STELLAR. *Nat Methods* **19**,

1411-1418, doi:10.1038/s41592-022-01651-8 (2022).

2   Vandereyken, K., Sifrim, A., Thienpont, B. & Voet, T. Methods and applications for single-cell

and spatial multi-omics. *Nat Rev Genet* **24**, 494-515, doi:10.1038/s41576-023-00580-2 (2023).

3   Stahl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by spatial

transcriptomics. *Science* **353**, 78-82, doi:10.1126/science.aaf2403 (2016).

4   Junker, J. P. *et al.* Genome-wide RNA Tomography in the zebrafish embryo. *Cell* **159**, 662-675,

doi:10.1016/j.cell.2014.09.038 (2014).

5   Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide expression

at high spatial resolution. *Science* **363**, 1463-1467, doi:10.1126/science.aaw1219 (2019).

6   Vickovic, S. *et al.* High-definition spatial transcriptomics for in situ tissue profiling. *Nat Methods*

16, 987-990, doi:10.1038/s41592-019-0548-y (2019).

7   Liao, J., Lu, X., Shao, X., Zhu, L. & Fan, X. Uncovering an Organ's Molecular Architecture at

Single-Cell Resolution by Spatially Resolved Transcriptomics. *Trends Biotechnol* **39**, 43-58,

doi:10.1016/j.tibtech.2020.05.006 (2021).

8   Rao, A., Barkley, D., Franca, G. S. & Yanai, I. Exploring tissue architecture using spatial

transcriptomics. *Nature* **596**, 211-220, doi:10.1038/s41586-021-03634-9 (2021).

9   Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF regression

to deconvolute spatial transcriptomics spots with single-cell transcriptomes. *Nucleic Acids Res*

49, e50, doi:10.1093/nar/gkab043 (2021).

10   Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial transcriptomics. *Nat*

Biotechnol **40**, 517-526, doi:10.1038/s41587-021-00830-w (2022).

11   Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial transcriptomics.

Nat Biotechnol **40**, 661-671, doi:10.1038/s41587-021-01139-4 (2022).

12   Ma, Y. & Zhou, X. Spatially informed cell-type deconvolution for spatial transcriptomics. *Nat*

Biotechnol **40**, 1349-1359, doi:10.1038/s41587-022-01273-7 (2022).

13   Andersson, A. *et al.* Single-cell and spatial transcriptomics enables probabilistic inference of cell

type topography. *Commun Biol* **3**, 565, doi:10.1038/s42003-020-01247-y (2020).

14   Li, B. *et al.* Benchmarking spatial and single-cell transcriptomics integration methods for

transcript distribution prediction and cell type deconvolution. *Nat Methods* **19**, 662-670,

doi:10.1038/s41592-022-01480-9 (2022).

15   Li, H. *et al.* A comprehensive benchmarking with practical guidelines for cellular deconvolution

of spatial transcriptomics. *Nat Commun* **14**, 1548, doi:10.1038/s41467-023-37168-7 (2023).

16   Sun, D., Liu, Z., Li, T., Wu, Q. & Wang, C. STRIDE: accurately decomposing and integrating
spatial transcriptomics using single-cell RNA sequencing. *Nucleic Acids Res* **50**, e42,
doi:10.1093/nar/gkac150 (2022).

17   Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *Nature* **608**, 766-777,
doi:10.1038/s41586-022-05060-x (2022).

18   Ackerman, J. E. *et al.* Defining the spatial-molecular map of fibrotic tendon healing and the
drivers of Scleraxis-lineage cell fate and function. *Cell Rep* **41**, 111706,
doi:10.1016/j.celrep.2022.111706 (2022).

19   Anderson, A. C. *et al.* Spatial transcriptomics. *Cancer Cell* **40**, 895-900,
doi:10.1016/j.ccell.2022.08.021 (2022).

20   Foster, D. S. *et al.* Multiomic analysis reveals conservation of cancer-associated fibroblast
phenotypes across species and tissue of origin. *Cancer Cell* **40**, 1392-1406 e1397,
doi:10.1016/j.ccell.2022.09.015 (2022).

21   Heiser, C. N. *et al.* Molecular cartography uncovers evolutionary and microenvironmental
dynamics in sporadic colorectal tumors. *Cell* **186**, 5620-5637 e5616,
doi:10.1016/j.cell.2023.11.006 (2023).

22   Matusiak, M. *et al.* Spatially Segregated Macrophage Populations Predict Distinct Outcomes In
Colon Cancer. *Cancer Discov*, doi:10.1158/2159-8290.CD-23-1300 (2024).

23   Kersten, K. *et al.* Spatiotemporal co-dependency between macrophages and exhausted CD8(+)
T cells in cancer. *Cancer Cell* **40**, 624-638 e629, doi:10.1016/j.ccell.2022.05.004 (2022).

24   van der Leun, A. M., Thommen, D. S. & Schumacher, T. N. CD8(+) T cell states in human cancer:
insights from single-cell analysis. *Nat Rev Cancer* **20**, 218-232, doi:10.1038/s41568-019-0235-
4 (2020).

25   Kirschenbaum, D. *et al.* Time-resolved single-cell transcriptomics defines immune trajectories
in glioblastoma. *Cell* **187**, 149-165 e123, doi:10.1016/j.cell.2023.11.032 (2024).

26   Benotmane, J. K. *et al.* High-sensitive spatially resolved T cell receptor sequencing with SPTCR-
seq. *Nat Commun* **14**, 7432, doi:10.1038/s41467-023-43201-6 (2023).

27   Liu, S. *et al.* Spatial maps of T cell receptors and transcriptomes reveal distinct immune niches
and interactions in the adaptive immune response. *Immunity* **55**, 1940-1952 e1945,
doi:10.1016/j.immuni.2022.09.002 (2022).

28   Delfini, M., Stakenborg, N., Viola, M. F. & Boeckxstaens, G. Macrophages in the gut: Masters in
multitasking. *Immunity* **55**, 1530-1548, doi:10.1016/j.immuni.2022.08.005 (2022).

29   Xun, Z. *et al.* Reconstruction of the tumor spatial microenvironment along the malignant-
boundary-nonmalignant axis. *Nat Commun* **14**, 933, doi:10.1038/s41467-023-36560-7 (2023).

30 Zhang, D. *et al.* Inferring super-resolution tissue architecture by integrating spatial
transcriptomics with histology. *Nat Biotechnol*, doi:10.1038/s41587-023-02019-9 (2024).

31 Fu, T. *et al.* Spatial architecture of the immune microenvironment orchestrates tumor immunity
and therapeutic response. *J Hematol Oncol* **14**, 98, doi:10.1186/s13045-021-01103-4 (2021).

32 Eming, S. A., Martin, P. & Tomic-Canic, M. Wound repair and regeneration: mechanisms,
signaling, and translation. *Sci Transl Med* **6**, 265sr266, doi:10.1126/scitranslmed.3009337
(2014).

33 Hu, K. H. *et al.* Transcriptional space-time mapping identifies concerted immune and stromal
cell patterns and gene programs in wound healing and cancer. *Cell Stem Cell* **30**, 885-903 e810,
doi:10.1016/j.stem.2023.05.001 (2023).

34 Mascharak, S. *et al.* Multi-omic analysis reveals divergent molecular events in scarring and
regenerative wound healing. *Cell Stem Cell* **29**, 315-327 e316, doi:10.1016/j.stem.2021.12.011
(2022).

35 McBrearty, B. A., Clark, L. D., Zhang, X. M., Blankenhorn, E. P. & Heber-Katz, E. Genetic
analysis of a mammalian wound-healing trait. *Proc Natl Acad Sci U S A* **95**, 11792-11797,
doi:10.1073/pnas.95.20.11792 (1998).

36 Clark, L. D., Clark, R. K. & Heber-Katz, E. A new murine model for mammalian wound repair
and regeneration. *Clin Immunol Immunopathol* **88**, 35-45, doi:10.1006/clin.1998.4519 (1998).

37 Heydemann, A. The super super-healing MRL mouse strain. *Front Biol (Beijing)* **7**, 522-538,
doi:10.1007/s11515-012-1192-4 (2012).

38 Pastar, I. *et al.* Epithelialization in Wound Healing: A Comprehensive Review. *Adv Wound Care*
(New Rochelle) **3**, 445-464, doi:10.1089/wound.2013.0473 (2014).

39 desJardins-Park, H. E., Foster, D. S. & Longaker, M. T. Fibroblasts and wound healing: an
update. *Regen Med* **13**, 491-495, doi:10.2217/rme-2018-0073 (2018).

40 Wynn, T. A. & Vannella, K. M. Macrophages in Tissue Repair, Regeneration, and Fibrosis.
Immunity **44**, 450-462, doi:10.1016/j.immuni.2016.02.015 (2016).

41 Kim, M. H. *et al.* Dynamics of neutrophil infiltration during cutaneous wound healing and
infection using fluorescence imaging. *J Invest Dermatol* **128**, 1812-1820,
doi:10.1038/sj.jid.5701223 (2008).

42 Phillipson, M. & Kubes, P. The Healing Power of Neutrophils. *Trends Immunol* **40**, 635-647,
doi:10.1016/j.it.2019.05.001 (2019).

43 Raziyeva, K. *et al.* Immunology of Acute and Chronic Wound Healing. *Biomolecules* **11**,
doi:10.3390/biom11050700 (2021).

44 Liu, S. *et al.* A tissue injury sensing and repair pathway distinct from host pathogen defense.

Cell **186**, 2127-2143 e2122, doi:10.1016/j.cell.2023.03.031 (2023).

45   Seferbekova, Z., Lomakin, A., Yates, L. R. & Gerstung, M. Spatial biology of cancer evolution.

Nat Rev Genet **24**, 295-313, doi:10.1038/s41576-022-00553-x (2023).

46   Erickson, A. *et al.* Spatially resolved clonal copy number alterations in benign and malignant

tissue. *Nature* **608**, 360-367, doi:10.1038/s41586-022-05023-2 (2022).

47   Lomakin, A. *et al.* Spatial genomics maps the structure, nature and evolution of cancer clones.

Nature **611**, 594-602, doi:10.1038/s41586-022-05425-2 (2022).

48   Qi, J. *et al.* Single-cell and spatial analysis reveal interaction of FAP(+) fibroblasts and SPP1(+)

macrophages in colorectal cancer. *Nat Commun* **13**, 1742, doi:10.1038/s41467-022-29366-6

(2022).

49   Trimm, E. & Red-Horse, K. Vascular endothelial cell development and diversity. *Nat Rev Cardiol*

20, 197-210, doi:10.1038/s41569-022-00770-1 (2023).

50   Li, G., Gao, J., Ding, P. & Gao, Y. The role of endothelial cell-pericyte interactions in

vascularization and diseases. *J Adv Res*, doi:10.1016/j.jare.2024.01.016 (2024).

51   Fridman, W. H. *et al.* B cells and tertiary lymphoid structures as determinants of tumour immune

contexture and clinical outcome. *Nat Rev Clin Oncol* **19**, 441-457, doi:10.1038/s41571-022-

00619-z (2022).

52   Munoz-Erazo, L., Rhodes, J. L., Marion, V. C. & Kemp, R. A. Tertiary lymphoid structures in

cancer - considerations for patient prognosis. *Cell Mol Immunol* **17**, 570-575,

doi:10.1038/s41423-020-0457-0 (2020).

53   Danenberg, E. *et al.* Breast tumor microenvironment structures are associated with genomic

features and clinical outcome. *Nat Genet* **54**, 660-669, doi:10.1038/s41588-022-01041-y (2022).

54   Dhainaut, M. *et al.* Spatial CRISPR genomics identifies regulators of the tumor

microenvironment. *Cell* **185**, 1223-1239 e1220, doi:10.1016/j.cell.2022.02.015 (2022).

55   Ramirez, C. F. A. *et al.* Cancer cell genetics shaping of the tumor microenvironment reveals

myeloid cell-centric exploitable vulnerabilities in hepatocellular carcinoma. *Nat Commun* **15**,

2581, doi:10.1038/s41467-024-46835-2 (2024).

56   DeNardo, D. G. & Ruffell, B. Macrophages as regulators of tumour immunity and immunotherapy.

Nat Rev Immunol **19**, 369-382, doi:10.1038/s41577-019-0127-6 (2019).

57   Dong, L. *et al.* The loss of RNA N(6)-adenosine methyltransferase Mettl14 in tumor-associated

macrophages promotes CD8(+) T cell dysfunction and tumor growth. *Cancer Cell* **39**, 945-957

e910, doi:10.1016/j.ccell.2021.04.016 (2021).

58   Cassetta, L. *et al.* Human Tumor-Associated Macrophage and Monocyte Transcriptional

Landscapes Reveal Cancer-Specific Reprogramming, Biomarkers, and Therapeutic Targets.

Cancer Cell **35**, 588-602 e510, doi:10.1016/j.ccell.2019.02.009 (2019).

Han, D. *et al.* Anti-tumour immunity controlled through mRNA m(6)A methylation and YTHDF1 in dendritic cells. *Nature* **566**, 270-274, doi:10.1038/s41586-019-0916-x (2019).

Kloosterman, D. J. & Akkari, L. Macrophages at the interface of the co-evolving cancer ecosystem. *Cell* **186**, 1627-1651, doi:10.1016/j.cell.2023.02.020 (2023).

Zhang, L. *et al.* Single-Cell Analyses Inform Mechanisms of Myeloid-Targeted Therapies in Colon Cancer. *Cell* **181**, 442-459 e429, doi:10.1016/j.cell.2020.03.048 (2020).

Li, N. *et al.* Mapping and modeling human colorectal carcinoma interactions with the tumor microenvironment. *Nat Commun* **14**, 7915, doi:10.1038/s41467-023-43746-6 (2023).

Honda, K. *et al.* IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* **434**, 772-777, doi:10.1038/nature03464 (2005).

Liu, S. Y., Sanchez, D. J., Aliyari, R., Lu, S. & Cheng, G. Systematic identification of type I and type II interferon-induced antiviral factors. *Proc Natl Acad Sci U S A* **109**, 4239-4244, doi:10.1073/pnas.1114981109 (2012).

Chiappinelli, K. B. *et al.* Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* **162**, 974-986, doi:10.1016/j.cell.2015.07.011 (2015).

He, J. *et al.* Identifying transposable element expression dynamics and heterogeneity during development at the single-cell level with a processing pipeline scTE. *Nat Commun* **12**, 1456, doi:10.1038/s41467-021-21808-x (2021).

Kong, Y. *et al.* Transposable element expression in tumors is associated with immune infiltration and increased antigenicity. *Nat Commun* **10**, 5228, doi:10.1038/s41467-019-13035-2 (2019).

Sexton, C. E., Tillett, R. L. & Han, M. V. The essential but enigmatic regulatory role of HERVH in pluripotency. *Trends Genet* **38**, 12-21, doi:10.1016/j.tig.2021.07.007 (2022).

Samstein, R. M. *et al.* Tumor mutational load predicts survival after immunotherapy across multiple cancer types. *Nat Genet* **51**, 202-206, doi:10.1038/s41588-018-0312-8 (2019).

Zehir, A. *et al.* Mutational landscape of metastatic cancer revealed from prospective clinical sequencing of 10,000 patients. *Nat Med* **23**, 703-713, doi:10.1038/nm.4333 (2017).

Haghverdi, L., Buttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. Diffusion pseudotime robustly reconstructs lineage branching. *Nat Methods* **13**, 845-848, doi:10.1038/nmeth.3971 (2016).

MacLean, A. L., Hong, T. & Nie, Q. Exploring intermediate cell states through the lens of single cells. *Curr Opin Syst Biol* **9**, 32-41, doi:10.1016/j.coisb.2018.02.009 (2018).

Zhou, P., Wang, S., Li, T. & Nie, Q. Dissecting transition cells from single-cell transcriptome data through multiscale stochastic dynamics. *Nat Commun* **12**, 5609, doi:10.1038/s41467-021-

1296   25548-w (2021).

74   Stickels, R. R. *et al.* Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-
seqV2. *Nat Biotechnol* **39**, 313-319, doi:10.1038/s41587-020-0739-1 (2021).

75   Del Bigio, M. R. Ependymal cells: biology and pathology. *Acta Neuropathol* **119**, 55-73,
doi:10.1007/s00401-009-0624-y (2010).

76   Ramachandran, V. S. *Encyclopedia of the Human Brain*. (Elsevier, 2002).

77   Meyer, G. Building a human cortex: the evolutionary differentiation of Cajal-Retzius cells and
the cortical hem. *J Anat* **217**, 334-343, doi:10.1111/j.1469-7580.2010.01266.x (2010).

78   Ptashkin, R. N. *et al.* Chromosome 20q Amplification Defines a Subtype of Microsatellite Stable,
Left-Sided Colon Cancers with Wild-type RAS/RAF and Better Overall Survival. *Mol Cancer Res*
15, 708-713, doi:10.1158/1541-7786.MCR-16-0352 (2017).

79   Shah, N. M. *et al.* Pan-cancer analysis identifies tumor-specific antigens derived from
transposable elements. *Nat Genet* **55**, 631-639, doi:10.1038/s41588-023-01349-3 (2023).

80   Liang, Y., Qu, X., Shah, N. M. & Wang, T. Towards targeting transposable elements for cancer
therapy. *Nat Rev Cancer* **24**, 123-140, doi:10.1038/s41568-023-00653-8 (2024).

81   Dai, M., Pei, X. & Wang, X. J. Accurate and fast cell marker gene identification with COSG. *Brief*
Bioinform **23**, doi:10.1093/bib/bbab579 (2022).

82   Kotliar, D. *et al.* Identifying gene expression programs of cell-type identity and cellular activity
with single-cell RNA-Seq. *Elife* **8**, doi:10.7554/eLife.43803 (2019).

83   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420,
doi:10.1038/nbt.4096 (2018).

84   Satoh, J. & Tabunoki, H. A Comprehensive Profile of ChIP-Seq-Based STAT1 Target Genes
Suggests the Complexity of STAT1-Mediated Gene Regulatory Mechanisms. *Gene Regul Syst*
Bio **7**, 41-56, doi:10.4137/GRSB.S11433 (2013).

85   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-
level datasets. *Nat Commun* **10**, 1523, doi:10.1038/s41467-019-09234-6 (2019).

86   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing biological
themes among gene clusters. *OMICS* **16**, 284-287, doi:10.1089/omi.2011.0118 (2012).

87   Mayer, J., Blomberg, J. & Seal, R. L. A revised nomenclature for transcribed human endogenous
retroviral loci. *Mob DNA* **2**, 7, doi:10.1186/1759-8753-2-7 (2011).

88   Rooney, M. S., Shukla, S. A., Wu, C. J., Getz, G. & Hacohen, N. Molecular and genetic
properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48-61,
doi:10.1016/j.cell.2014.12.033 (2015).

89   Hanzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray
and RNA-seq data. *BMC Bioinformatics* **14**, 7, doi:10.1186/1471-2105-14-7 (2013).

90   Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the targets of
focal somatic copy-number alteration in human cancers. *Genome Biol* **12**, R41, doi:10.1186/gb-
2011-12-4-r41 (2011).

91   Reich, M. *et al.* GenePattern 2.0. *Nat Genet* **38**, 500-501, doi:10.1038/ng0506-500 (2006).

92   Saunders, A. *et al.* Molecular Diversity and Specializations among the Cells of the Adult Mouse
Brain. *Cell* **174**, 1015-1030 e1016, doi:10.1016/j.cell.2018.07.028 (2018).

1338