Hypoxia inducible factor HIF1α elevates expression of mRNA capping enzyme during cobalt
chloride-induced hypoxia

Safirul Islam and Chandrama Mukherjee*

Institute of Health Sciences, Presidency University, Plot No. DG/02/02, Premises No. 14-0358,
Action Area 1D, Kolkata 700156, India

*Corresponding author, email: chandrama.dbs@presiuniv.ac.in

Email and ORCID IDs for all authors

Safirul Islam: safirul.rs@presiuniv.ac.in, ORCID: 0000-0001-5859-1331

Chandrama Mukherjee: chandrama.dbs@presiuniv.ac.in, ORCID: 0000-0002-4749-1295

Abstract

In response to hypoxia, hypoxia-inducible factors (HIFs) control the transcriptomic output to mitigate
the hypoxic stress. Long noncoding RNAs (lncRNA) are found to be very crucial in regulating
hypoxia. Like mRNAs, lncRNAs are protected by 5' caps that are added by mRNA capping enzyme
(CE) in the nucleus. The previous concept that capping takes place in the nucleus was changed by the
recognition of a cytoplasmic pool of capping enzyme (cCE). cCE has been shown to recap its
substrate uncapped mRNAs or long noncoding RNAs (lncRNAs) present in the cytoplasm, preventing
their degradation, even during arsenite-induced oxidative stress. In this study, we examined the effect
of $CoCl_2$ induced hypoxia on cCE and its function in regulating the substrate lncRNAs.

Here, we show that $CoCl_2$ induced hypoxia elevates the expressions of nuclear and cytoplasmic CE in
HIF1α dependent manner as evidenced by Chromatin immunoprecipitation and HIF1α inhibitor
experiments. Furthermore, we found cCE post-transcriptionally controls the stability of its target
lncRNAs amidst $CoCl_2$ induced hypoxia. These results suggest that cCE, upregulated by HIF1α, may
act as a posttranscriptional modulator for a few cCE-targeted lncRNAs.

Keywords. mRNA capping enzyme/*RNGTT*, cytoplasmic capping, $CoCl_2$ induced hypoxia, HIF1α,
long noncoding RNA

List of Abbreviations:

CE, mRNA capping enzyme

cCE, cytoplasmic mRNA capping enzyme

HIFs, hypoxia inducible factors

HRR, hypoxia responsive RNA

CoCl$_2$, Cobalt chloride

ChIP, chromatin immunoprecipitation

LncRNA, long noncoding RNA

miRNA, microRNA

circRNA, circular RNA

RNGTT, RNA Guanylyltransferase And 5'-Phosphatase

RNMT, RNA guanine 7-methyl transferase

RAMAC, RNA guanine 7-methyl transferase activating subunit

K294A, dominant negative mutant of cCE

bHLH, basic-helix–loop–helix

LUAD, lung adenocarcinoma

RCC, renal cell carcinoma

1. Introduction

Hypoxia is one of the contributing factors for generating a solid tumor microenvironment and plays a key role in the progression of tumor metastasis. In 1953 *Gray et. al.* showed the responsiveness of hypoxic tumor cells toward radiotherapy is poor as compared to well-oxygenated tumor cells (1). Additional studies revealed that hypoxia has a profound effect on tumor cell biology which includes its metabolism, angiogenic potential, and sensitivity to chemotherapy or immunotherapy (2). Oxygen-sensitive hypoxia-inducible factor 1α (HIF1α) is the master regulator of hypoxia which reprograms the cellular transcriptomic output in concert with constitutively expressed β subunit so that cells can withstand hypoxic stress (3–5). In addition to HIF1α, several noncoding RNAs like microRNAs (miRNA), long noncoding RNAs (lncRNA), circular RNAs (circRNA)) and epigenetic factors also regulate the cellular response to hypoxia (6).

Cellular stresses are categorized into two groups type 1 and type 2. Hypoxia, heat shock, and arsenite treatment belong to type 1 stress whereas genotoxic drugs belong to type 2 stress. Cells initiate specific cell stressor mechanisms for their survival upon confronting any environmental stress (7). Studies have indicated that hypoxia markedly reduces *de novo* RNA synthesis mediated by modulating chromatin architecture (8,9). In addition, cellular response to hypoxia is also achieved at the post-transcriptional level which includes mRNA splicing, stability, etc. Hypoxia-induced transcriptional regulation is well studied whereas its effect on RNA stability at the post-transcriptional level is still not comprehensive. The 5' cap is one of the crucial factors that determines the stability of an RNA transcript and how this structure is affected during hypoxic stress, remains elusive.

The majority of eukaryotic RNAs that are transcribed by RNA pol II must have a 5' cap structure which consists of an N^7 methylated guanosine residue that is attached covalently to the first nucleotide of the transcripts by an enzyme called RNA guanylyltransferase and 5'-phosphatase (RNGTT) or mRNA capping enzyme (CE) (10,11). Canonical capping takes place inside the nucleus where CE in complex with RNA guanine 7-methyl transferase (RNMT) and RNA guanine 7-methyl transferase activating subunit (RAMAC) act on the 5'end of the nascent RNA transcripts to add a cap (12–14). Later, a pool of cytosolic capping enzymes (cCE) was identified which can recap the uncapped transcripts in the cytoplasm resulting in their stabilization (15,16). cCE forms a complex with the cytoplasmic pool of RNMT, RAMAC along with an unknown 5'RNA kinase and an adaptor protein Nck1 that provides a platform for the assembly of these components to make active cCE that regulates the stability and translability of certain mRNAs and thus has been speculated in maintaining 'cap homeostasis' to regulate gene expression (16–18). Inhibiting cytoplasmic capping by overexpressing a dominant negative mutant of cCE (K294A) increased a pool of uncapped RNAs (16,19). In agreement with these findings, studies across several labs revealed the presence of uncapped RNAs in various cells pointing to the possibilities of cytoplasmic recapping. Indeed, with the advancement of imaging

techniques, mRNAs have been shown to undergo cyclical decapping and recapping (20). Later,

recapping was revealed to act as a nexus of post-transcriptional control as cCE not only regulates

mRNAs but also a subset of lncRNAs (21).

Besides these canonical functions of CE/cCE either in the protection and maturation of RNA

pol II transcribed transcripts or recapping cytosolic transcripts, recent studies have indicated its

potential role in various biological processes ranging from modulating the Hedgehog signaling

pathway during *Drosophila* development to its involvement in certain cancers (22,23). A current study

demonstrates CE governs biliary-derived liver regeneration initiation by transcriptional regulation of

mTORC1 and Dnmt1 in Zebrafish (24). Additionally, one recent study uncovers that METTL3

regulates the expression and activity of CE in non-small cell lung cancer (NSCLC) cell lines (25).

Recently, a study revealed the role of cCE in protecting the cap status of its target mRNAs during

arsenite-induced oxidative stress (26). However, there is no study regarding the effect of hypoxic

stress on CE/cCE.

In this study, we examine the effect of Cobalt chloride ($CoCl_2$) induced hypoxia on the expression of

CE or cCE for the first time. $CoCl_2$ is a well-known hypoxia-mimetic chemical compound that

triggers hypoxia-like conditions in vitro (27). Using chemically induced hypoxia on osteosarcoma cell

lines (U2OS and MG63) as our model systems, we found $CoCl_2$ induced hypoxia elevates the

expression of CE/cCE. Furthermore, we examine the steady state level of previously validated cCE-

targeted lncRNAs (21) in $CoCl_2$ induced hypoxic conditions in cells expressing a dominant negative

mutant of cCE (K294A) (21). Our study shows some of these lncRNA are upregulated in $CoCl_2$

induced hypoxia and cCE regulates their stability indicating its role in post-transcriptional regulation

of a subset of hypoxia-responsive lncRNAs.

2. Materials and methods

2.1 Cloning of K294A in pLVX-TetOne-Puro vector:

Mouse cytoplasmic restricted catalytically inactive capping enzyme (K294A) was amplified from

pcDNA4/TO-bio-myc-NES-mCEΔNLS (K294A) construct (21) using Q5® High-Fidelity DNA

Polymerase (NEB, cat no# M0491L). The pLVX-TetOne-Puro-myc-NES-mCEΔNLS(K294A)

construct was prepared by subcloning myc-NES-mCEΔNLS(K294A) into an empty pLVX-TetOne-

Puro vector (Takara Bio, cat. No# 631847) using In-Fusion HD Cloning Kit (Takara Bio, cat no#

638909). The construct was further confirmed by Sanger sequencing and is available upon request.

The sequences of the cloning primers are listed in Table S1.

2.2 Cell culture, treatments, and generation of stable cell lines:

Human osteosarcoma cell line U2OS was purchased from the American Type Culture Collection

(ATCC), and MG63 from the National Centre for Cell Science (NCCS, Pune). Both the cell lines were

routinely cultured in 1X DMEM (Gibco, cat no# 11965092) supplemented with 10% fetal bovine

serum (Himedia, cat no# RM1112) and 1% penicillin-streptomycin (Gibco, cat no# 15140-122) in the

presence of 5% CO_2 at 37 □C in a humidified incubator (Eppendorf). Mycoplasma contamination was

tested to confirm contamination-free cell culture using a PCR-detection kit (Sigma-Aldrich, cat no#

MP0035).

For hypoxia induction *in vitro*, cells were treated with different concentrations (100 µM, 200 µM, and

300 µM) of $CoCl_2$ (Sigma Aldrich, cat no# C8661) solution that was made by dissolving in DPBS

(Gibco, cat no# 14040182). After 24 hours of treatment, cells were harvested for RNA/protein

extraction (28).

HIF1α inhibitor PX478 was procured from the MedChemExpress (cat #HY-10231). 10mM stock of

PX478 was prepared by dissolving it in deionized water. Cells were treated with PX478 to a final

concentration of 40µM to inhibit HIF1α activity (29).

To generate inducible stable cell lines expressing bio-myc K294A, U2OS cells were transfected with

pLVX-TetOne-Puro bio-myc K294A. After 48 hours of transfection, Puromycin (50µg/ml, Sigma

Aldrich, cat# P9620) was added to multiple dilutions of transfected cells and transfectants were

incubated for clonal selection of the stable cells according to the manufacturer's protocols till the

isolation of the single colonies. After growing a few selected colonies, Doxycyclin (Dox) (1ug/ml)

(Sigma Aldrich, cat no# D5207) was added to the selected clones for inducible expression of K294A.

The stable cell line expressing the highest bio myc K294A protein (Fig: S3C-D) was selected for

subsequent experiments.

2.3 XTT cell viability assay:

The viability of osteosarcoma cells upon treatment with different concentrations of $CoCl_2$ solution

was assessed by using an XTT cell viability assay kit (Cell Signalling Technology, cat no#9095)

following the manufacturer's protocol. Briefly, 2000 cells of U2OS/MG63 were seeded in 96 well

plates and treated with different concentrations of $CoCl_2$ solutions. 24 hours post-treatment, 50 µl of

XTT reagent was added to each well and incubated for 3 h in 5% CO_2 at 37 □C. The absorbance was

measured at 450 nm using a BioTek Synergy microplate reader.

2.4 RNA interference and treatment:

For Xrn1 knockdown, K294A stable cells uninduced or induced were transfected with Xrn1 siRNA

(Ambion, assay ID# s29016) targeting the coding region of *XRN1* and scramble siRNA at 10nM final

concentration using Lipofectamine RNAiMAX reagent following manufacturers protocol (Invitrogen,

cat no# 13778-075). 24 hrs post-transfection, transfection mix containing media was replenished with

fresh growth medium. Total of 48 hrs post-transfection half of the cultured cells were treated with

Dox (1µg/ml) to induce K294A expression and maintained in culture for another 24 hrs. Cells were

harvested for protein and RNA isolation. The knockdown efficiency was affirmed by western blotting

using rabbit anti-Xrn1 antibody (Invitrogen, cat no# PA5-57110).

2.5 Cytoplasmic and nuclear fractionations:

To isolate cytoplasmic fractions, U2OS cell pellets were resuspended in cytoplasmic lysis buffer (20

mM Tris-Cl of pH 7.5, 10 mM NaCl, 10 mM MgCl$_2$, 10 mM KCl, 0.2% NP40, 1mM PMSF,

RNAseOUT and 1X protease inhibitor cocktail) and incubated on ice for 10 min with intermittent

gentle agitation. Nuclei were removed by centrifugation at 1,000 xg for 10 minutes at 4°C.

To isolate nuclear fractions, the nuclear pellet was resuspended in nuclear lysis buffer (150 mM NaCl,

1% Triton X-100, 0.5% Sodium Deoxycholate, 1% SDS, and 25 mM Tris-Cl of pH 7.5) for 40

minutes with occasional agitation. The nuclear extract was prepared by centrifugation at 10,000 xg for

10 minutes at 4 □C.

2.6 RNA extraction and poly(A) selection:

Total RNA was extracted by adding TRIzol to the harvested cells (Invitrogen cat# 15596026)

according to the manufacturer's protocol. To isolate cytoplasmic RNA TRIzol reagent was added to

cytoplasmic lysate (2.5) following by manufacturer's protocol. RNAs were then treated with DNase I

(Thermo Scientific, cat no# EN0521) and extracted using phenol: chloroform: isoamyl alcohol (Sisco

Research Laboratories Pvt Ltd, cat no# 69031). 2μg of DNA-free RNA was used for poly(A) selection

using Dynabeads mRNA DIRECT Kit (Invitrogen, cat no# 61011) according to the manufacturer's

instruction.

2.7 Quantitative real-time PCR analysis:

2 μg of either total RNA or cytoplasmic RNA was reverse transcribed using Super Script III First

Strand cDNA Synthesis kit (Invitrogen cat# 18080051) according to the manufacturer's instruction. A

1:1 cDNA dilution was done to perform real-time PCR using SsoAdvanced Universal SYBR Green

Supermix (Bio-Rad cat no#172-5271) in CFX connect Real-Time instrument (Bio-Rad). The PCR

cycle was carried out using the following thermal conditions initial denaturation at 95 □C for 30 s

followed by 40 cycles of 95 □C for 10 s and 60 □C for 15 s and a melting curve was, generated by

steadily increasing the temperature from 65 □C to 95 □C in 0.5 □C increments with a duration of 2 s/

step. *Rplpo* was used as an internal control. The relative fold change of gene expression was

calculated using the comparative CT method (ΔΔCt). Unless specified otherwise, at least three

different biological replicates were used. The complete list of primers is listed in Table S1.

2.8 Western blotting analysis:

Cells were harvested in ice-cold DPBS and the cell pellet was collected by centrifuging at 1000 g for

5 min at 4 □C. For Whole cell lysis, harvested cells were resuspended in whole cell lysis buffer (50

mM Tris-Cl pH 8.0, 150 mM NaCl, and 1% NP-40) supplemented with 1mM PMSF and protease

cocktail inhibitor (Roche, cat no# 4693159001) and kept in ice for 30 minutes with intermittent

vortex. Following incubation, the whole cell lysate was collected by centrifuging the suspension at

12000 xg for 10 minutes at 4 □C. A Rapid Gold BCA Protein Assay kit (Pierce, cat no# A53226) was

used to quantify protein samples. 30-60 µg protein extracts were fractionated using 8-10% SDS-

PAGE under reducing conditions and transferred onto Immobilon-FL PVDF (Millipore, cat no#

IPFL00010) membranes. The membranes were blocked in 3% skimmed milk in phosphate-buffered

saline (PBS) for 30 min at room temperature followed by incubation in respective primary antibody

dilution. The dilutions are as follows 1:1000 rabbit anti-HIF1α (Cell Signaling Technology, cat no#

36169); 1:1000 mouse anti-beta actin (Santa Cruz Biotechnology, cat no# SC-517582), 1:1000 rabbit

anti-CE (Novus Biologicals, cat no# NPB1-49973); 1:2000 mouse anti-GAPDH (Novus Biologicals,

cat#2D4A7); 1:2000 rabbit anti-CA9 (Cloud-clone Crop, cat# PAD076Hu01); 1:2000 mouse anti-

lamin (DHSB, cat no# MANLAC1(4A7), 1:1000 rabbit anti-Xrn1 antibodyat 4 □C overnight with

constant shaking. Membranes were washed thrice in PBS-T after primary antibody incubation and

incubated with secondary antibodies like donkey anti-MouseDyLight 800 (Cell Signaling Technology,

cat no#5257) and donkey anti-Rabbit680 (Invitrogen, cat no#A10043) at 1: 10000 dilutions in PBS-T.

Blots were scanned in OdysseyCLx Imaging System (LiCor Inc.) and band intensities were calculated

using ImageJ software (NIH, USA).

2.9 Immunofluorescence:

Cells were cultured on coverslips and fixed in freshly prepared fixing solution [4%PFA(EMS)+0.2%

Triton-100 in 1X PBS] for 10 min at room temperature. Coverslips were washed thrice in washing

solution (0.5mM MgCl$_2$, 0.05% Triton-100 in 1X PBS) and blocked in blocking buffer (5% FBS,

0.2M glycine, 0.1% Triton-100 in 1X PBS) for 30 min at room temperature. Coverslips were

incubated with primary antibodies (1:100, rabbit anti-HIF1α; 1:100 rabbit anti-CE) overnight at 4 □C.

Following washing coverslips were incubated with secondary antibodies of donkey IgG conjugated

with either Alexa Fluor 488 (Invitrogen cat no# A21206) or 568 (Invitrogen, cat no# A10037) at a

dilution of 1:1000 and Hoechst 33358 (Sigma, cat no# 94403) diluted in blocking buffer for 1 hr at

room temperature. Coverslips were mounted on glass slides with a mounting medium (Invitrogen, cat

no# S36967). Images were acquired using a Confocal laser scanning microscope (Leica) fitted with a

63X Plan Apo oil immersion objective (NA 1.4). Images were analyzed, and the fluorescence intensity

measurements were performed using the Zeiss Zen 3.3 software package.

2.10 Chromatin Immunoprecipitation (ChIP) assay:

HIF1α binding sites on CE promoter were predicted by using JASPAR software

(http://www.jaspar.genereg.net). The ChIP assay was performed using a Simple Chip Enzymatic

Chromatin IP Kit (Cell Signaling Technology, cat# 9003) as per the manufacturer's

instructions. Briefly, the treated/ untreated cells were fixed in 1% formaldehyde for 10 min at room
temperature to cross-link the DNA-protein complex. Glycine was added to the cell to quench the
unreacted formaldehyde. Following washing with ice-cold PBS containing 1mM PMSF, cells were
harvested by centrifugation at 2000 g for 5 min at 4 □C. Cell pellet was lysed in lysis buffer (1×
buffer A, 0.5 mM DTT, 1× Protease Inhibitor Cocktail, 1 mM PMSF) and nuclei were pellet down at
2000 g for 5 min at 4 □C. The nuclei pellet was resuspended in buffer B supplemented with 0.5 mM
DTT and treated with micrococcal nuclease for 20 min at 37 □C. Following centrifugation at 16000 g
for 1 min at 4 □C. Nuclei pellet was resuspended in ChIP buffer (1× ChIP buffer, 1× Protease
Inhibitor Cocktail, 1 mM PMSF), followed by sonication to break the nuclei membrane and lysate
was clarified by centrifugation at 9400 g for 10 min at 4 □C. 10 μg of chromatin DNA was incubated
with rabbit anti-HIF1α (Cell Signaling Technology, cat no# 36169) or normal rabbit IgG (Cell
Signaling Technology) overnight at 4°C.To the IP samples protein G magnetic beads were added and
incubated 2 hrs at 4°C with rotation. The bead-chromatin-antibody complexes were washed three
times in a low-salt wash buffer and one time in a high-salt buffer and eluted in elution buffer heating
at 65 °C for 30 min. The proteins were digested by treating them with proteinase K at 65 °C for 2 hrs
followed by DNA isolation using a spin column. The immunoprecipitated and input DNA were
analyzed by Agarose gel electrophoresis as well as subjected to real-time PCR using SsoAdvanced
Universal SYBR Green Supermix (Bio-Rad cat#172-5271) in CFX connect Real-Time instrument
(Bio-Rad). The primers for ChIP qPCR are listed in Table S1. Relative fold enrichment was calculated
using the comparative CT method (ΔΔCt) described previously where IgG was used as the negative
control.

2.11 *In Vitro* Xrn1 susceptibility assay:

30 ng poly(A) selected cytoplasmic RNAs of different experimental conditions from three
independent biological replicates were heat denatured at 65 □C for 5 minutes and set up for digestion
by adding 0.5 units of Xrn1 enzyme (NEB, cat no# M0338S). Prior to digestion, the cytoplasmic
poly(A)RNAs were spiked with synthetic capped Renilla luciferase mRNA as negative control for the
experiments respectively. The RNA samples were treated with Xrn1 incubated for 1 hr at 37 □C.
After incubation, the digested RNAs were recovered using phenol: chloroform extraction followed by
ethanol precipitation. cDNA was prepared using iscript cDNA synthesis kit (Bio-Rad cat no#
1708890). RT-qPCR was performed using gene-specific primer pairs against the 5'-ends of the
candidate transcripts as done in earlier studies. The C_t values for each cCE targeted transcripts and
non-targeted controls were normalized against the C_t value of capped Renilla luciferase obtained from
each condition. The loss of 5' ends from each experimental condition was obtained by calculating the
differences in normalized C_t values of each transcript before and after Xrn1 digestion. ΔX or change
in Xrn1 susceptibility in cells was measured by $\Delta X_{K294A}/\Delta X_{Control}$ ratio, where $\Delta X_{Control}$ is the relative

loss of transcript 5' ends in control, and ΔX_{K294A} is their relative loss in K294A-expressing cells. The

plot showed the relative change in Xrn1 susceptibility between $CoCl_2$ treated and untreated cells.

2.12 Statistical analysis:

257   The statistical analysis was calculated by performing Two-tail Student's *t*-tests for two groups and

258   one-way ANOVA for multiple groups using GraphPad Prism 9 Software (GraphPad Software, Inc).

259   All the data were represented as ±SD of at least three independent biological replicates (n=3). Data

260   with $p<0.05$ were considered statistically significant.

3. Result and Discussion

3.1 Hypoxia mimetic $CoCl_2$ augments HIF1α expression in U2OS and MG63 osteosarcoma cells.

The characterization and activity of cCE have been well elucidated in U2OS osteosarcoma cells (15–

17,21). Other cell lines like NIH3T3 mouse fibroblasts or human RPE1 (diploid) or primary cells like

mouse astrocytes or cardiomyocytes express variable amounts of cCE (26). To determine the effect of

hypoxia on the CE/cCE, an *in vitro* chemically induced hypoxic cellular model was established using

two osteosarcoma cell lines U2OS and MG63 along with the diploid cell line hTERT RPE-1 (Fig.1,

S1). $CoCl_2$ is one of the most commonly used hypoxia mimetic reagents for the induction of chemical

hypoxia in cellular models and the presence of HIF1α in the cellular extract is the indicator of

successful hypoxia induction (28,30,31). Both U2OS and MG63 cells were exposed to different

concentrations (100 µM, 200 µM, and 300 µM) of $CoCl_2$ solution for 24 h and the induction of $CoCl_2$

triggered hypoxia was confirmed by determining the presence of hypoxia marker HIF1α at the protein

level by immunoblotting and immunofluorescence staining (Fig. 1). The gradual increase of HIF1α

protein expression with the increasing dose of $CoCl_2$ at 24 h was observed in both cell lines (Figs.

1A,1B, 1D, & 1E) which affirms the establishment of hypoxia. The toxicity of $CoCl_2$ in the cell lines

was examined by performing a XTT cell viability assay (32). Concentration-dependent cytotoxicity of

$CoCl_2$ was observed in both cell lines where 300 µM showed the highest toxicity (Figs. 1C and 1F).

Treatment of cells with 200 µM $CoCl_2$ concentration did not show any toxicity to both the cell lines.

So, considering the expression of hypoxic marker HIF1α and the minimal cytotoxicity of $CoCl_2$, 200

µM was selected for further experiments with both cell lines. The immunofluorescence images of

HIF1α also corroborate with the immunoblotting results for $CoCl_2$ induced hypoxia in U2OS and

MG63 cell lines (Figs. 1G and 1H). RPE-1 cells also showed elevated expression of HIF-1α after

being treated with 200 µM $CoCl_2$ for 24 h (Figs. S1A and B) suggesting a similar mechanism exists.

3.2 $CoCl_2$-induced hypoxia elevates mRNA-capping enzyme expression in osteosarcoma cells

CE protects mRNAs and lncRNAs by adding a 5' cap structure and besides this protection of 5' ends

of these RNAs it also helps the RNA transcripts for their efficient export from the nucleus to the

cytoplasm, co-transcriptional processing, and decay (33,34). However, there are a few studies that

report the regulation of expression of mammalian CE itself or its recruitment to target genes. A recent
study shows catalytically active CE can bind to RNA polymerase II and its target genes in the
presence of c-Myc (35). Earlier, we have shown that CE can shuttle between the nucleus and the
cytoplasm and localize to stress granules during arsenite-induced stress (26). However, the impact of
other stressors like hypoxia, a major contributing factor to the development of solid tumors, on CE is
yet to be known. This led us to examine the effect of $CoCl_2$ mimetic hypoxia on CE. To understand
this, both U2OS and MG63 cells were incubated with $CoCl_2$ for 24 h, followed by staining using CE
antibody, and images were captured by a confocal microscope. The immunofluorescence results
showed the enhanced expression of total CE, particularly cytoplasmic CE as evidenced by increased
staining intensity in the cytoplasm in $CoCl_2$ treated cells, compared to untreated cells in both cell lines
(Figs. 2A & 2C). We further quantified the mean intensity of CE and our data revealed significant
upregulation of CE in $CoCl_2$ induced hypoxic cells compared to untreated cells (Figs.2B & 2D). To
determine if this increased level of CE was due to the change in the level of the total protein or change
in the cytoplasmic-nuclear distribution, total protein levels were measured from untreated and $CoCl_2$
treated cells and an increase in the level of CE was observed in both cell lines (Figs: 4B and 4D,
discussed later) supporting the immunofluorescence data (Figs. 2A-D). To examine the distribution of
nuclear and cytoplasmic CE upon $CoCl_2$ treatment, U2OS cells were fractionated in nuclear and
cytoplasmic fractions using a standard method (21) and the efficiency of separations were measured
by western hybridization with antibodies against Lamin A/C (nuclear protein) and GAPDH
(cytoplasmic protein) (Fig: 2E). The chemical induced hypoxic condition was confirmed by the
cytosolic expression of CA9, in $CoCl_2$ treated cells since it was a positive marker for $CoCl_2$ induced
hypoxia (36) (Fig: 2E). An increased level of CE was observed in both nuclear and cytoplasmic
fractions after $CoCl_2$ treatment (Fig: 2E). However, in agreement with the previous
immunofluorescence results (Figs. 2A, 2C), cytosolic CE was expressed more in $CoCl_2$ induced
hypoxia as evidenced by the quantitation (Fig: 2F). A similar increase in the expression of the total
population of CE is also observed in RPE1 cells (Fig. S1C-D). Collectively, these data suggest $CoCl_2$
induced hypoxia increased expression of CE/cCE in osteosarcoma and human diploid cell lines
indicating possible regulation of CE by HIF1α.

3.3 Mammalian *RNGTT* promoter harbours hypoxia-responsive elements (HRE)

Previous studies showed hypoxia or $CoCl_2$ induced hypoxia stimulates the expression of HIF1α (37).
Similarly, in our $CoCl_2$ induced hypoxia model, we observed an accumulation of HIF1α protein
during hypoxia (Fig: 1). HIF1α is a member of basic-helix–loop–helix (bHLH)-PAS domain-
containing protein family (38) and known to induce the transcription of gene expression by binding to
the HRE (R/CGTG) elements present upstream of the transcription start sites in the promoter region
(39,40). This led us to speculate that HIF1α might control *RNGTT* expression. *In silico* promoter
analysis revealed putative HRE sequences in the promoter of *RNGTT* as represented schematically in

Fig. 3A. The *in-silico* data was validated in CoCl$_2$ treated or untreated U2OS cells by performing ChIP

assay with anti-HIF1α antibody, followed by PCR analysis with different sets of primers targeted to

two putative HIF1α binding sites (BS1 & BS2) within 5 kb upstream of the transcription start sites.

PCR results showed the direct binding of HIF1α, but not the IgG control to the *RNGTT* promoter in

both the binding sites in CoCl$_2$ treated U2OS cells but not in untreated cells (Fig. 3B). The absence of

binding of HIF1α in the untreated cells may be due to the degradation of HIF1α in the untreated cells

as it is unstable in presence of oxygen (41). Further qPCR analysis revealed that in CoCl$_2$ induced

hypoxic U2OS cells, HIF1α binds to *RNGTT* promoter to BS1 and BS2 with 12-fold and 15-fold

enrichment respectively, as compared to untreated cells (Fig. 3C).

3.4 Depletion of HIF1α reduces expression of CE in osteosarcoma cells

To corroborate the ChIP assay results which showed binding of HIF1α to *RNGTT* promoter, HIF1α

was depleted using pharmacological inhibitor PX478. PX478 is a well-known inhibitor of HIF1α (29).

It affects its expression both at transcriptional, and translational levels. Moreover, PX478 prevents de-

ubiquitination of ubiquitin-tagged HIF1α molecules (42). Osteosarcoma cells were treated with CoCl$_2$

and PX478 was added to these cells only. To test if the binding of HIF1α to *RNGTT* promoter was

direct or not, we performed ChIP assay as before (3.3) with PX478 treated or untreated CoCl$_2$ induced

hypoxic cells. The binding of HIF1α to BS1 and BS2 sites were markedly reduced in PX478 treated

cells compared to untreated cells (Fig: S2A). Later, qPCR analysis showed that PX478 treatment led

to ~7 and 13-fold reduced binding of HIF1α to BS1 and BS2 sites respectively on *RNGTT* promoter

as compared to PX478 untreated CoCl$_2$ induced hypoxic cells. (Fig: S2B) thus confirming the direct

binding of HIF1α to the *RNGTT* promoter.

Based on this result, we speculated the compromised binding of HIF1α to *RNGTT* promoter would

result reduced expression of CE in PX478 treated cells. To test this, we treated U2OS cells and MG63

cells with CoCl$_2$ or kept untreated. PX478 was added to CoCl$_2$ induced hypoxic cells and total RNA

and protein levels were measured from different treatment groups. We observed a reduction in the

level of *RNGTT* mRNA in HIF1α inhibited, CoCl$_2$ induced hypoxic samples compared to HIF1α non-

inhibited CoCl$_2$ induced hypoxic samples for both cell lines (Fig. 4A). Western blot analysis showed

increased expression of HIF1α in CoCl$_2$ induced hypoxic samples but decreased expression of HIF1α

in PX478 treated, CoCl$_2$ induced hypoxic samples in osteosarcoma cells (Figs. 4B-4E).

Next, we investigated the effect of HIF1α inhibition on the expression of CE. We observed CE

expression also reduced in HIF1α inhibited samples as compared to respective controls in

osteosarcoma cells (Figs. 4B-4E). Therefore, our study showed the elevated expression of *RNGTT*

mRNA or the encoded protein during CoCl$_2$ induced hypoxia. This was due to the direct binding of

the HIF1α transcription factor on the *RNGTT* promoter which was abolished upon inhibition of
HIF1α.

3.5 cCE stabilizes its targeted lncRNAs post-transcriptionally during hypoxia

The hypoxia microenvironment is one of the common features of solid tumors that regulates tumor
cells both genotypically and phenotypically (43). Hypoxia, in turn, modulates the expression of
various non-coding RNAs to make phenotypic changes in tumor cells, and lncRNAs are one of them
(44,45). Previous findings from our group suggested a few lncRNAs could be targeted post-
transcriptionally by cCE (21). Considering this fact, we are intrigued to understand whether cCE has
any role in regulating its target lncRNAs during chemical induced hypoxia. For that, we have chosen
ASH1L-AS1, GAS5-215, ENTPD3-AS1, NFYC-AS1, and H19-009 which are related to any solid
tumors and a few of them established as substrates of cCE (21). H19-009 and GAS5 lncRNAs are
elevated in various tumors (46,47). The other candidate lncRNAs are also involved in developing
cancer occurrence and development; such as participation of ASH1L-AS1 in gastric cancer (GC),
ENTPD3-AS1 in lung adenocarcinoma (LUAD), and renal cell carcinoma (RCC), NFYC-AS1 in
LUAD (48–51).

To analyze the effect of hypoxia on these cCE-targeted lncRNAs, $CoCl_2$ treated or untreated U2OS
cells were separated into nuclear and cytoplasmic fractions efficiently as judged by the detection of
Lamin A/C in the nuclear fractions and GAPDH in the cytoplasmic fractions (Fig. S3A). RNA was
extracted from cytoplasmic fractions and relative expressions of the candidate lncRNAs were
measured using RT-qPCR. We found that out of five lncRNAs, four of them (ASH1L-AS1, GAS5-
215, ENTPD3-AS1, and H19-009) were significantly upregulated, whereas the expression of NFYC-
AS1 was significantly reduced in $CoCl_2$ treated cells (Fig: 5A). As the stability of some of these
lncRNAs was regulated by cCE (20), we wondered if the upregulated expressions of these HRLs were
contributed by cCE.

To examine this, we made Tetracycline-inducible stable U2OS cell lines expressing cytoplasmic
restricted, catalytically inactive dominant negative form CE (referred to here as K294A), as
schematically represented in Fig S3B. In this cell line, the nuclear localization signal (NLS) was
deleted and the nuclear export signal (NES) was added to the CE construct to make it cytoplasm-
restricted, as evident from the immunofluorescence image (Fig. 5B). These cells were treated with
Doxycycline (Dox) to overexpress K294A (Fig: S3D). The same cell line was used as Control for
subsequent assays where no overexpression was observed in absence of Dox (Fig. S3D). Control and
K294A cells were treated with or without $CoCl_2$ for 24 hrs as described earlier. Following incubation,
all the cells were fractionated into nuclear and cytoplasmic fractions. The efficiency of fractions was
assessed by western hybridization using antibodies against Lamin A/C and GAPDH as done earlier
(Fig. 5C). Our data showed no cross-contamination among the fractions. The expression of CA9, was

also checked to confirm the $CoCl_2$ induced hypoxic condition (Fig. 5C). The steady-state expressions

of the candidate lncRNAs in cytoplasmic fractions isolated from Control and K294A cells in normal

condition, as well as $CoCl_2$ induced hypoxic conditions, were measured using RT-qPCR (Fig. 5D).

In agreement with our previous study (21), no significant changes were observed in the expression of

ENTPD3-AS1 whereas ASH1L-AS1 showed significant elevation in its expression in K294A-

expressing cells (Fig. 5D). However, in contrast to our earlier observation with transiently transfected

cells (21), we noticed significantly reduced expressions of GAS5-215, NFYC-AS1, and H19-009 in

K294A cells as compared to Control cells under similar conditions (Fig. 5D). This difference may be

attributed to the usage of clonal stable cell lines in the current study expressing the similar level of

K294A in contrast to a mixed population of cells expressing different levels of K294A along with

untransfected cells generated from transient transfection experiments performed in the previous study

(21). A similar trend was also observed for mRNA substrates of cCE, as reported in previous studies

where decreased expressions of cCE targeted mRNAs were noticed in U2OS cell line stably

expressing K294A (18,52). We hypothesized since K294A cells express inactive cCE, the cCE-

targeted transcripts like GAS5-215, NFYC-AS1, and H19-009 present in the cytoplasm could not be

recapped and would be susceptible to exoribonuclease-mediated degradation resulting in reduced

expression in K294A cells. To address this, we depleted Xrn1, the major exoribonuclease from K294A

and Control cells using specific siRNA targeting Xrn1 as shown in the schematic Fig: 5E.

Cytoplasmic fractions were isolated to collect RNA and protein from these treatment groups. Our data

showed more than 80% depletion of Xrn1 (Figs: 5F-G). Further qPCR analysis of cytoplasmic RNAs

from the above treatment groups revealed elevated expressions of GAS5-215, NFYC-AS1 and H19-

009 in K294A cells compared to Control cells when Xrn1 was depleted (Fig. 5H) supporting our

hypothesis. On the other hand, ASH1L-AS1 and ENTPD3-AS1 might be protected by some other

mechanism that yet to be studied.

Interestingly, we observed the expressions of ASH1L-AS1, GAS5-215 and H19-009, which were

significantly upregulated in $CoCl_2$ induced hypoxic control cells, were reduced in $CoCl_2$ induced

hypoxic K294A cells (Fig. 5D). In addition, the expression of NFYC-AS1 that did not elevate in

response to CoCl2 treatment, was reduced markedly in presence of K294A (Fig. 5D). The possible

reason could be during hypoxic stress, these transcripts might have lost its 5' cap structure and in the

absence of active cCE, they could not be recapped, subjected to exoribonucleases. Hence, their

expressions were reduced in $CoCl_2$ mimetic hypoxic K294A cells (Fig. 5D). To examine this

possibility, we performed Xrn1 susceptibility assay that can distinguish uncapped RNA from capped

RNAs (16). Cytoplasmic poly(A) RNAs from $CoCl_2$ treated and untreated cells mixed with capped

Rennila luciferase mRNA were subjected to *in vitro* Xrn1 digestion followed by RT-qPCR for 5

candidate lncRNAs, *STRN4* and *STAT3*, which were negative and positive controls of cCE

respectively. The C_t values obtained for each transcript in $\pm$ CoCl$_2$ treated cells were normalized
against that of the spike luciferase mRNA, which remained unchanged by Xrn1 treatment.

Xrn1 susceptibility or loss of 5' ends are represented as the difference in relative Ct values before and
after Xrn1 digestion of the same transcript in $\pm$K294A cells as described in the methods (Fig: 5H).
We observed in absence of active cCE, GAS5-215, H19-009 and NFYC-AS1 transcripts lost their
caps during CoCl$_2$ mimetic hypoxic conditions as evidenced by high susceptibility to Xrn1 similar to
STAT3 transcript suggesting role of cCE protecting these transcripts during CoCl$_2$ treatment. In
contrast, expression of ASH1L-AS1 transcript that was reduced in CoCl$_2$ treated K294A expressing
cells, remained unaffected by Xrn1 digestion, indicating retaining cap structure under same condition
(Fig: 5H). The similar notion was observed for expression of ENTPD3-AS1 transcript too that
increased in K294A expressing cells, retained its cap structures under the same condition like STRN4
(Fig: 5H). These data suggest that cCE plays an important role in regulating the stability of GAS5-
215, H19-009 and NFYC-AS1 transcripts in CoCl$_2$ mimetic hypoxic conditions post-transcriptionally.

4. Conclusion

Nuclear capping is a co-transcriptional event whereas cytoplasmic capping is a post-transcriptional
event. In our previous study, we demonstrated that a subset of lncRNAs is post-transcriptionally
stabilized by cCE which recapped the uncapped transcripts in the cytoplasm. In this study, we showed
for the first time that CE/cCE is upregulated by HIF1α due to the binding of this transcription factor to
HRE-containing sites in the *RNGTT* promoter. We also examined some tumor-associated lncRNAs
and found they are differentially expressed during CoCl$_2$ induced hypoxia. Importantly, a few of them
(H19-009, GAS5-215, and NFYC-AS1) are post-transcriptionally regulated by cCE as evidenced by
their decreased expressions in K294A cells under normal conditions. Additionally, we observed
reduced expressions of these transcripts in CoCl$_2$ induced hypoxic K294A cells which suggests cCE
post-transcriptionally regulates the stability of these transcripts. Further, we showed cCE protects it's
target transcripts, H19-009, GAS5-215, and NFYC-AS1 during hypoxia when they lost their caps.
However, no significant change in Xrn1 susceptibility was seen for ASH1L-AS1 under this condition.
Perhaps, other mechanisms may be responsible for its down regulation which needs to be studied in
the future.

In this present study, we have examined the effect of hypoxia on the total pool of CE using CoCl$_2$ as a
hypoxia mimetic agent on osteosarcoma cells and hTERT RPE-1 as cellular models. Mechanistically,
the enhanced expression of CE/cCE under this chemical induced hypoxia is executed by HIF1α which
is a known transcription factor and the master regulator of hypoxia. CE is known to maintain the
stability of most RNA pol II transcripts that include both mRNAs and lncRNAs. Further investigation
should be carried out to identify all hypoxia-responsive RNAs regulated by cCE. Since the present
study relies upon only on CoCl$_2$ that stimulated HIF1α-mediated responses via its inhibition of prolyl

hydroxylase (39), and this treatment might have some off target effects. It would be better for future experiments to be conducted under physical hypoxic condition that mimics more closely the tumour microenvironment to examine the effect of hypoxic stress on CE/cCE and the plausible post transcriptional regulation of the transcriptome by CE/cCE.

Moreover, earlier study showed the protective function of cCE on its target transcripts during recovery from a brief arsenite induced oxidative stress through recapping the decapped transcripts. Further studies should examine if cCE play similar protective role to its target transcripts during recovery from hypoxic conditions. Taken together, our data reveals HIF1α mediated upregulation of CE/cCE and a novel function of cCE in post-transcriptional regulation of target transcripts during $CoCl_2$ induced hypoxia.

Acknowledgments

This work was supported by extramural grants CRG/2019/006427 and CRG/2023/006384 from the Department of Science and Technology, Government of India to Dr. Chandrama Mukherjee. The authors also thank UGC, Government of India for a UGC-SRF fellowship to Mr. Safirul Islam and the Department of Biotechnology, India for a Ramlingaswami Re-entry fellowship to Dr. Chandrama Mukherjee. The authors thank the Institute of Health Sciences, Presidency University for the departmental instrument facility and Presidency University for necessary infrastructural support, Dr. Piyali Mukherjee, Presidency University for providing Lamin antibody, Dr. Shubhra Majumder, Presidency University for providing access to Zeiss Microscope and Dr. Abhik Saha, Presidency University for providing access to Licor imager.

Author contributions: CRediT

Safirul Islam: Conceptualization, Methodology, Validation, Investigation, Formal Analysis, Visualization, Writing-original draft, Writing - Review & Editing; **Chandrama Mukherjee:** Conceptualization, Resources, Writing - Original Draft, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

References

1. Gray LH, Conger AD, Ebert M, Hornsey S, Scott OCA. The Concentration of Oxygen Dissolved in Tissues at the Time of Irradiation as a Factor in Radiotherapy. Br J Radiol [Internet]. 1953 Dec;26(312):638–48. Available from: https://academic.oup.com/bjr/article/26/312/638-648/7299137

2. Chen Z, Han F, Du Y, Shi H, Zhou W. Hypoxic microenvironment in cancer: molecular mechanisms and therapeutic interventions. Signal Transduct Target Ther [Internet]. 2023 Feb 17;8(1):70. Available from: http://www.ncbi.nlm.nih.gov/pubmed/36797231

3. Parodi M, Raggi F, Cangelosi D, Manzini C, Balsamo M, Blengio F, et al. Hypoxia Modifies the Transcriptome of Human NK Cells, Modulates Their Immunoregulatory Profile, and Influences NK Cell Subset Migration. Front Immunol [Internet]. 2018;9(OCT):2358.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/30459756

4.   Abou Khouzam R, Sharda M, Rao SP, Kyerewah-Kersi SM, Zeinelabdin NA, Mahmood AS,
et al. Chronic hypoxia is associated with transcriptomic reprogramming and increased genomic
instability in cancer cells. Front cell Dev Biol [Internet]. 2023 Mar 9;11(March):1095419.
Available from: https://www.frontiersin.org/articles/10.3389/fcell.2023.1095419/full

5.   Reheman A, Wu Q, Xu J, He J, Qi M, Li K, et al. Transcriptomic analysis of the hypoxia-
inducible factor 1α impact on the gene expression profile of chicken fibroblasts under hypoxia.
Poult Sci [Internet]. 2024;103(3):103410. Available from:
https://doi.org/10.1016/j.psj.2023.103410

6.   Islam S, Mukherjee C. Molecular regulation of hypoxia through the lenses of noncoding RNAs
and epitranscriptome. Wiley Interdiscip Rev RNA [Internet]. 2023 Mar;14(2):e1750. Available
from: http://www.ncbi.nlm.nih.gov/pubmed/35785444

7.   Arimoto K, Fukuda H, Imajoh-Ohmi S, Saito H, Takekawa M. Formation of stress granules
inhibits apoptosis by suppressing stress-responsive MAPK pathways. Nat Cell Biol [Internet].
2008 Nov;10(11):1324–32. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18836437

8.   Johnson AB, Barton MC. Hypoxia-induced and stress-specific changes in chromatin structure
and function. Mutat Res. 2007 May 1;618(1–2):149–62.

9.   Johnson AB, Denko N, Barton MC. Hypoxia induces a novel signature of chromatin
modifications and global repression of transcription. Mutat Res Mol Mech Mutagen [Internet].
2008 Apr 2;640(1–2):174–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18294659

10.   Shatkin AJ. Capping of eucaryotic mRNAs. Cell. 1976 Dec;9(4 PT 2):645–53.

11.   Ensinger MJ, Moss B. Modification of the 5' terminus of mRNA by an RNA (guanine-7-)-
methyltransferase from HeLa cells. J Biol Chem. 1976 Sep 10;251(17):5283–91.

12.   Ramanathan A, Robb GB, Chan S-H. mRNA capping: biological functions and applications.
Nucleic Acids Res. 2016 Sep 19;44(16):7511–26.

13.   Pillutla RC, Yue Z, Maldonado E, Shatkin AJ. Recombinant human mRNA cap
methyltransferase binds capping enzyme/RNA polymerase IIo complexes. J Biol Chem. 1998
Aug 21;273(34):21443–6.

14.   Varshney D, Lombardi O, Schweikert G, Dunn S, Suska O, Cowling VH. mRNA Cap
Methyltransferase, RNMT-RAM, Promotes RNA Pol II-Dependent Transcription. Cell Rep.
2018 May 1;23(5):1530–42.

15.   Otsuka Y, Kedersha NL, Schoenberg DR. Identification of a cytoplasmic complex that adds a
cap onto 5'-monophosphate RNA. Mol Cell Biol [Internet]. 2009 Apr;29(8):2155–67.
Available from: http://www.ncbi.nlm.nih.gov/pubmed/19223470

16.   Mukherjee C, Patil DP, Kennedy BA, Bakthavachalu B, Bundschuh R, Schoenberg DR.
Identification of Cytoplasmic Capping Targets Reveals a Role for Cap Homeostasis in
Translation and mRNA Stability. Cell Rep [Internet]. 2012 Sep 27;2(3):674–84. Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf

17.   Mukherjee C, Bakthavachalu B, Schoenberg DR. The cytoplasmic capping complex assembles
on adapter protein nck1 bound to the proline-rich C-terminus of Mammalian capping enzyme.
PLoS Biol [Internet]. 2014 Aug;12(8):e1001933. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/25137142

18.   Trotman JB, Giltmier AJ, Mukherjee C, Schoenberg DR. RNA guanine-7 methyltransferase
catalyzes the methylation of cytoplasmically recapped RNAs. Nucleic Acids Res. 2017 Oct
13;45(18):10726–39.

19.   Berger MR, Alvarado R, Kiss DL. mRNA 5′ ends targeted by cytoplasmic recapping cluster at
CAGE tags and select transcripts are alternatively spliced. FEBS Lett. 2019;593(7).

20.   Iwasaki S, Ingolia NT. PROTEIN TRANSLATION. Seeing translation. Science [Internet].
2016 Jun 17;352(6292):1391–2. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/27313023

21.   Mukherjee A, Islam S, Kieser RE, Kiss DL, Mukherjee C. Long noncoding RNAs are
substrates for cytoplasmic capping enzyme. FEBS Lett. 2023;597(7):947–61.

22.   Chen P, Zhou Z, Yao X, Pang S, Liu M, Jiang W, et al. Capping Enzyme mRNA-cap/RNGTT
Regulates Hedgehog Pathway Activity by Antagonizing Protein Kinase A. Sci Rep. 2017 Jun
6;7(1):2891.

23.   Borden K, Culjkovic-Kraljacic B, Cowling VH. To cap it all off, again: dynamic capping and
recapping of coding and non-coding RNAs to control transcript fate and biological activity.
Cell Cycle. 2021 Jul;20(14):1347–60.

24.   Ma J, Yang Z, Huang Z, Li L, Huang J, Chen J, et al. Rngtt governs biliary-derived liver
regeneration initiation by transcriptional regulation of mTORC1 and Dnmt1 in zebrafish.
Hepatology [Internet]. 2023 Jul 1;78(1):167–78. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/36724876

25.   Valle-morales D, Romano G, Le P, Saviana M, Brown R, Li H, et al. METTL3 alters capping
enzyme expression and its activity on ribosomal proteins . 2023;3.

26.   Gayen A, Mukherjee A, Kumar K, Majumder S, Chakrabarti S, Mukherjee C. The mRNA-
capping enzyme localizes to stress granules in the cytoplasm and maintains cap homeostasis of
target mRNAs. J Cell Sci [Internet]. 2024 Jun 1;137(11). Available from:
https://www.biorxiv.org/content/10.1101/2023.03.28.534620v1%0Ahttps://www.biorxiv.org/c
ontent/10.1101/2023.03.28.534620v1.abstract

27.   Hirsilä M, Koivunen P, Xu L, Seeley T, Kivirikko KI, Myllyharju J. Effect of desferrioxamine
and metals on the hydroxylases in the oxygen sensing pathway. FASEB J [Internet]. 2005 Aug
7;19(10):1308–10. Available from: https://onlinelibrary.wiley.com/doi/10.1096/fj.04-3399fje

28.   Rana NK, Singh P, Koch B. CoCl2 simulated hypoxia induce cell proliferation and alter the
expression pattern of hypoxia associated genes involved in angiogenesis and apoptosis. Biol
Res [Internet]. 2019 Dec 15;52(1):12. Available from:
https://biolres.biomedcentral.com/articles/10.1186/s40659-019-0221-z

29.   Tiwari A, Tashiro K, Dixit A, Soni A, Vogel K, Hall B, et al. Loss of HIF1A From Pancreatic
Cancer Cells Increases Expression of PPP1R1B and Degradation of p53 to Promote Invasion
and Metastasis. Gastroenterology. 2020;159(5):1882-1897.e5.

30.   Muñoz-Sánchez J, Chánez-Cárdenas ME. The use of cobalt chloride as a chemical hypoxia
model. J Appl Toxicol [Internet]. 2019 Apr 28;39(4):556–70. Available from:
https://analyticalsciencejournals.onlinelibrary.wiley.com/doi/10.1002/jat.3749

31.   Bae S, Jeong HJ, Cha HJ, Kim K, Choi YM, An IS, et al. The hypoxia-mimetic agent cobalt
chloride induces cell cycle arrest and alters gene expression in U266 multiple myeloma cells.
Int J Mol Med. 2012;30(5).

32.   Brosin A, Wolf V, Mattheus A, Heise H. Use of XTT-assay to assess the cytotoxicity of
different surfactants and metal salts in human keratinocytes (HaCaT). A feasible method for in
vitro testing of skin irritants. Acta Derm Venereol [Internet]. 1997 Jan;77(1):26–8. Available
from: http://www.ncbi.nlm.nih.gov/pubmed/9059672

33.   Topisirovic I, Svitkin Y V., Sonenberg N, Shatkin AJ. Cap and cap-binding proteins in the
control of gene expression. Wiley Interdiscip Rev RNA [Internet]. 2011 Mar 28;2(2):277–98.

Available from: https://wires.onlinelibrary.wiley.com/doi/10.1002/wrna.52

34.   Schoenberg DR, Maquat LE. Regulation of cytoplasmic mRNA decay. Nat Rev Genet
[Internet]. 2012 Mar 6;13(4):246–59. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/22392217

35.   Lombardi O, Varshney D, Phillips NM, Cowling VH. c-Myc deregulation induces mRNA
capping enzyme dependency. Oncotarget [Internet]. 2016 Dec 13;7(50):82273–88. Available
from: http://www.ncbi.nlm.nih.gov/pubmed/27756891

36.   Chu CY, Jin YT, Zhang W, Yu J, Yang HP, Wang HY, et al. CA IX is upregulated in CoCl2-
induced hypoxia and associated with cell invasive potential and a poor prognosis of breast
cancer. Int J Oncol. 2016;48(1).

37.   Semenza GL. HIF-1 and mechanisms of hypoxia sensing. Curr Opin Cell Biol [Internet]. 2001
Apr;13(2):167–71. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0955067400001940

38.   Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-
helix-PAS heterodimer regulated by cellular O2 tension. Proc Natl Acad Sci U S A.
1995;92(12):5510–4.

39.   Semenza GL, Agani F, Booth G, Forsythe J, Iyer N, Jiang B-H, et al. Structural and functional
analysis of hypoxia-inducible factor 1. Kidney Int [Internet]. 1997 Feb;51(2):553–5. Available
from: https://linkinghub.elsevier.com/retrieve/pii/S0085253815599171

40.   Weidemann A, Johnson RS. Biology of HIF-1α. Cell Death Differ [Internet]. 2008 Apr
15;15(4):621–7. Available from: https://www.nature.com/articles/cdd200812

41.   Jaśkiewicz M, Moszyńska A, Króliczewski J, Cabaj A, Bartoszewska S, Charzyńska A, et al.
The transition from HIF-1 to HIF-2 during prolonged hypoxia results from reactivation of
PHDs and HIF1A mRNA instability. Cell Mol Biol Lett. 2022;27(1).

42.   Koh MY, Spivak-Kroizman T, Venturini S, Welsh S, Williams RR, Kirkpatrick DL, et al.
Molecular mechanisms for the activity of PX-478, an antitumor inhibitor of the hypoxia-
inducible factor-1alpha. Mol Cancer Ther [Internet]. 2008 Jan;7(1):90–100. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/18202012

43.   Macharia LW, Muriithi W, Heming CP, Nyaga DK, Aran V, Mureithi MW, et al. The
genotypic and phenotypic impact of hypoxia microenvironment on glioblastoma cell lines.
BMC Cancer [Internet]. 2021 Nov 19;21(1):1248. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/34798868

44.   Barth DA, Prinz F, Teppan J, Jonas K, Klec C, Pichler M. Long-Noncoding RNA (lncRNA) in
the Regulation of Hypoxia-Inducible Factor (HIF) in Cancer. Non-coding RNA [Internet].
2020 Jul 6;6(3):27. Available from: https://www.mdpi.com/2311-553X/6/3/27

45.   Peng P-H, Hsu K-W, Chieh-Yu Lai J, Wu K-J. The role of hypoxia-induced long noncoding
RNAs (lncRNAs) in tumorigenesis and metastasis. Biomed J [Internet]. 2021 Oct;44(5):521–
33. Available from: https://linkinghub.elsevier.com/retrieve/pii/S2319417021000305

46.   Liu F, Pan H, Xia G, Qiu C, Zhu Z. Prognostic and clinicopathological significance of long
noncoding RNA H19 overexpression in human solid tumors: evidence from a meta-analysis.
Oncotarget [Internet]. 2016 Dec 13;7(50):83177–86. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/27825121

47.   Ni W, Yao S, Zhou Y, Liu Y, Huang P, Zhou A, et al. Long noncoding RNA GAS5 inhibits
progression of colorectal cancer by interacting with and triggering YAP phosphorylation and
degradation and is negatively regulated by the m6A reader YTHDF3. Mol Cancer [Internet].
2019 Oct 16;18(1):143. Available from: http://www.ncbi.nlm.nih.gov/pubmed/31619268

48. Xie M, Zhang L, Han L, Huang L, Huang Y, Yang M, et al. The ASH1L-AS1-ASH1L axis
controls NME1-mediated activation of the RAS signaling in gastric cancer. Oncogene
[Internet]. 2023 Nov;42(46):3435–45. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/37805663

49. Chiang H, Ong C, Chang C, Wu K-L, Wu Y, Lai J, et al. Downregulated antisense lncRNA
ENTPD3-AS1 contributes to the development of lung adenocarcinoma. Am J Cancer Res
[Internet]. 2024;14(2):854–68. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/38455397

50. Pandini C, Pagani G, Tassinari M, Vitale E, Bezzecchi E, Saadeldin MK, et al. The pancancer
overexpressed NFYC Antisense 1 controls cell cycle mitotic progression through in cis and in
trans modes of action. Cell Death Dis [Internet]. 2024 Mar 11;15(3):206. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/38467619

51. Song Y, Du J, Lu P, Zou Q, Zeng S, Liu M, et al. LncRNA NFYC-AS1 promotes the
development of lung adenocarcinomas through autophagy, apoptosis, and MET/c-Myc
oncogenic proteins. Ann Transl Med [Internet]. 2021 Nov;9(21):1621. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/34926665

52. Del Valle Morales D, Trotman JB, Bundschuh R, Schoenberg DR. Inhibition of cytoplasmic
cap methylation identifies 5 top mrnas as recapping targets and reveals recapping sites
downstream of native 5 ends. Nucleic Acids Res. 2020;48(7).

Figure Legends

Figure 1: Hypoxia mimetic CoCl$_2$ augments HIF1α expression in U2OS and MG63 osteosarcoma
cells. A & D. Western blot analysis of hypoxic stress marker HIF1α, showing gradual increase upon
increasing concentrations of CoCl$_2$ for 24 hrs treatment in U2OS and MG63 osteosarcoma cells
respectively. B & E. Quantification of HIF1α protein by ImageJ software using three independent
biological replicates. β-actin was used as the loading control. C & F. XTT cell viability assay showing
CoCl$_2$ dose-dependent loss of cellular viability in U2OS and MG63 cells. G & H. Immunostaining
analysis showing an elevated expression of HIF1α (green) during hypoxia in osteosarcoma cells. DNA
is represented by DAPI. The scale bar represents 50 μm. Statistical significance was calculated by
unpaired two-tailed Student's t-test represented as ± SD from three biological replicates. *ns, P*: non-
significant, *$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$, ****$P < 0.0001$; n≥3.

Figure 2: CoCl$_2$-induced hypoxia elevates expression of CE/cCE in osteosarcoma cells. A & C.
Immunostaining analysis showing enhanced expression of nuclear (CE) and cytoplasmic mRNA
capping enzyme (cCE) in CoCl$_2$ treated cells as compared to untreated cells. The scale bar represents
50 μm. B&D. Quantification of mean CE intensity using Zeiss Zen 3.3 software package, n≥20 cells. E.
Western blot analysis of CE in cytoplasmic and nuclear fractions of U2OS cells treated or untreated
with CoCl$_2$. GAPDH and Lamin A/C were used as cytoplasmic and nuclear markers respectively
whereas CA9 was used as a hypoxic marker. F. Quantification of CE bands intensity from E showing
the amount of CE in the nucleus and cytoplasm in U2OS cells treated or untreated with CoCl$_2$.
Normalization of cytoplasmic and nuclear CE was done using GAPDH and Lamin A/C respectively.
Statistical significance was calculated by unpaired two-tailed Student's t-test represented as ± SD
from three biological replicates. *ns, P*: non-significant, *$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$, ****$P <$
0.0001; n≥3.

Figure 3: The promotor of mammalian CE (*RNGTT*) harbors hypoxia-responsive elements (HRE). A.
Human RNGTT gene contains two HIF1α binding motif (R/CGTG) in its promoter region as shown in

the schematic. B. Chromatin immunoprecipitation (ChIP) analysis showed direct binding of HIF1α in
the *RNGTT* promoter regions (BS1 and BS2). 2% input samples were loaded. The molecular weight of
the DNA ladder is shown on the left. C. ChIP assay followed by quantitative RT–PCR (RT–qPCR)
showed ~12 and 15-fold enrichment of HIF1α binding to the BS1 and BS2 of RNGTT promoter region
in $CoCl_2$ treated cells respectively. Statistical significance was calculated by unpaired two-tailed
Student's t-test represented as ± SD from three biological replicates. ns: non-significant, *P< 0.05,
**P< 0.005, ***P< 0.0005, ****P< 0.0001; n≥3.

Figure 4: Depletion of HIF1α reduces expression of CE in osteosarcoma cells. A. Quantitative RT–
PCR (RT–qPCR) analysis revealed treatment of osteosarcoma cells (U2OS and MG63) with 40µM
HIF1α inhibitor PX478 in $CoCl_2$ treated conditions resulting in a significant reduction in *RNGTT* mRNA
expression. B & E. Western blot analysis shows an increased expressions of both HIF1α and CE
protein upon $CoCl_2$ treatment which were decreased significantly upon treatment with PX478 in
presence of $CoCl_2$. C & E. Quantification of HIF1α and CE protein by ImageJ software using three
independent biological replicates. β-actin was used as the loading control. Values are represented as
± SD from three biological replicates. Statistical significance was drawn by performing unpaired two-
tailed Student's t-test, ns, P: non-significant, *P< 0.05, **P< 0.005, ***P< 0.0005, ****P< 0.0001;
n≥3.

Figure 5: cCE stabilizes its targeted lncRNAs posttranscriptionally during hypoxia. A. RT-qPCR
analysis of cCE-targeted lncRNAs shows differential expression patterns in cytoplasmic extracts of
$CoCl_2$ treated U2OS cells. The C_t values of each lncRNA was normalized to the internal control, *Rplp0*.
The relative fold change of the transcripts was plotted after normalizing to the control, untreated
samples. $CoCl_2$ treatment significantly elevates the expression of ASH1L-AS1, GAS5-215, ENTPD3-AS1,
and H19-009, whereas NFYC-AS1 expression was significantly reduced. A paired Two-tailed Student's
t-test was performed for statistical analysis. B. Confocal image of Doxycycline inducible stable clonal
cell line C1 which expressed Myc tagged K294A in the cytoplasm only. C. Nuclear and cytoplasmic
fractionation of U2OS cells expressing K294A in the presence or absence of $CoCl_2$. Western blot of
Myc and CA9 confirm K294A expression and $CoCl_2$ induced hypoxia, respectively. The quality of
fractionation was further substantiated by western blotting of nuclear marker (Lamin A/C) and
cytoplasmic marker (GAPDH). D. The steady-state expressions of the candidate transcripts in K294A
stable cells in ± $CoCl_2$ conditions. GAS5-215, NFYC-AS1, and H19-009 were significantly decreased in
K294A-expressing $CoCl_2$ untreated cells. But ASH1L-AS1 was significantly increased and ENTPD3-AS1
did not show any significant change. Only GAS5-215 and H19-009 showed significantly reduced
expression in $CoCl_2$ treated K294A-expressing cells. The expression of each transcript was normalized
to *Rplp0* and the relative fold change was plotted after comparing to control cells. E. Schematic of
Xrn1 knockdown experiment. F. Western blot analysis of Xrn1 using cell lysate of K294A
expressing/non-expressing cells transfected with either scramble or Xrn1 siRNA. G. Densitometric
quantification of protein bands. β-actin was used as a loading control. H. RT-qPCR analysis of
candidate transcripts using cytoplasmic RNAs of Xrn1 knockdown samples. *STRN4* is a control
transcript whereas *STAT3* is a recapping target transcript respectively. The C_t values of each transcript
was calculated by normalizing to the C_t values of the internal control, *Rplp0*. The relative fold change
was calculated by normalizing to the control cells. Similar to STAT3, the steady-state levels of GAS5-
215, NFYC-AS1, and H19-009 showed significant enrichment upon Xrn1 knockdown. Statistical
significance was calculated using One way ANOVA to compare scr vs scr+K294A and si RNA Xrn1 vs si
RNA Xrn1+ K294A. I. Xrn1 susceptibility assay was performed using poly (A) selected cytoplasmic
RNAs from different conditions as described before (5D) and treated with ± Xrn1 followed by RNA
extraction and RT-qPCR using gene-specific primers. Renila luciferase spike in control that was
unchanged during ± Xrn1 was used as an internal control for normalization. The Xrn1 susceptibility is

measured by differences in normalized C_t values between Xrn1 treated and non-treated sample. The
Xrn1 susceptibility of cells expressing K294A and the non-expressing cells were plotted according to
the previous study (Mukherjee et al., 2012). Unpaired Student's t-test was used for statistical
analysis. Values are represented as ± SD from three biological replicates. ns, P: non-significant, *P<
0.05, **P< 0.005, ***P< 0.0005, ****P< 0.0001; n≥3.