Title page

Dynamics and Regulatory Roles of RNA m^{6}A Methylation in Unbalanced

Genomes

Shuai Zhang,[1,2,†] Xinyu Liu,[1,2,†] Ruixue Wang,[1,2,†] Junhan Wang,[1,2] Ludan Zhang,[1,2] and Lin Sun[1,2,*]

[1] Beijing Key Laboratory of Gene Resource and Molecular Development, College of Life Sciences,

Beijing Normal University, Beijing 100875, China

[2] Key Laboratory of Cell Proliferation and Regulation Biology of Ministry of Education, College of

Life Sciences, Beijing Normal University, Beijing 100875, China

† These authors contributed equally to this work.

* For correspondence: sunlin@bnu.edu.cn (Lin Sun)

Abstract

N^6-methyladenosine (m^{6}A) in eukaryotic RNA is an epigenetic modification that is critical for RNA

metabolism, gene expression regulation, and the development of organisms. Aberrant expression

of m^{6}A components appears in a variety of human diseases. RNA m^{6}A modification in *Drosophila*

has proven to be involved in sex determination regulated by *Sxl* and may affect X chromosome

expression through the MSL complex. The dosage-related effects under the condition of genomic

imbalance (i.e., aneuploidy) are related to various epigenetic regulatory mechanisms. Here, we

investigated the roles of RNA m^{6}A modification in unbalanced genomes using aneuploid

Drosophila. The results showed that the expression of m^{6}A components changed significantly

under genomic imbalance, and affected the abundance and genome-wide distribution of m^{6}A,

which may be related to the developmental abnormalities of aneuploids. The relationships

between methylation status and classical dosage effect, dosage compensation, and inverse

dosage effect were also studied. In addition, we demonstrated that RNA m^{6}A methylation may

affect dosage-dependent gene regulation through dosage-sensitive modifiers, alternative splicing,

the MSL complex, and other processes. More interestingly, there seems to be a closely relationship between MSL complex and RNA m^{6}A modification. It is found that ectopically overexpressed MSL complex, especially the levels of H4K16Ac through MOF could influence the expression levels of m^{6}A modification and genomic imbalance may be involved in this interaction. We found that m^{6}A could affect the levels of H4K16Ac through MOF, a component of the MSL complex, and that genomic imbalance may be involved in this interaction. Altogether, our work reveals the dynamic and regulatory role of RNA m^{6}A modification in unbalanced genomes, and may shed new light on the mechanisms of aneuploidy-related developmental abnormalities and diseases.

Introduction

Epigenetic modifications regulate gene expression in response to environmental changes and play important roles in the development of organisms and a variety of human diseases (Jaenisch and Bird, 2003; Lence et al., 2017). In addition to DNA and chromatin modifications, which are well studied, more than one hundred RNA chemical modifications have been identified in cells to date (Lee et al., 2014; Roundtree et al., 2017a). As a marker of post-transcriptional regulation, RNA modifications participate in almost all aspects of RNA metabolism (Roundtree et al., 2017a). N^6-methyladenosine (m^{6}A) is the most prevalent internal modification in many eukaryotic messenger RNAs (mRNAs) and long noncoding RNAs (lncRNAs) (Lee et al., 2014; Roundtree et al., 2017a; Yang et al., 2018), which widely affects RNA alternative splicing (Dominissini et al., 2012), export (Roundtree et al., 2017b), stability (Wang X et al., 2014), and translation (Meyer et al., 2015). RNA m^{6}A modifications have been found to be enriched on the transcripts of genes that regulate development and cell fate specification (Dominissini et al., 2012; Meyer et al., 2012; Geula et al., 2015), and some m^{6}A sites are regulated in a tissue- or disease-specific manner (Zaccara et al., 2019). In addition, the abnormal expression of m^{6}A components is related to the tumorigenesis, proliferation, and metastasis of many types of cancers (Pinello et al., 2018; Ma et al., 2019). Therefore, it is of great significance to study RNA m^{6}A methylation for revealing the mechanisms of gene expression regulation and human diseases.

At present, there are few studies on RNA m^{6}A modification in *Drosophila*, possibly due to its

relatively low abundance (m^{6}A/A < 0.2%) (Haussmann et al., 2016), and the mutation of some

m^{6}A component genes will affect their viability and fertility (Hongay and Orr-Weaver, 2011;

Haussmann et al., 2016; Kan et al., 2017). However, as a model organism with specific genetic

and developmental advantages, *Drosophila* remains an excellent tool for studying the roles of

epigenetic modifications in gene regulation, individual development, and disease process. Several

components of the *Drosophila* m^{6}A methyltransferase complex (*Ime4*, *dMettl14*, and *fl(2)d*

constitute the core complex, with *vir* and *nito* acting as cofactors) and a m^{6}A reader protein

(*Ythdc1*) have been identified, all of which have homologues in mammals (Lence et al., 2017).

Deletion of the major methyltransferase gene *Ime4* or the reader *Ythdc1* causes locomotion

defects, and the splicing of sex-determining factor *Sxl* is affected (Haussmann et al., 2016; Lence

et al., 2016; Kan et al., 2017). However, homozygous mutations in *fl(2)d*, *vir*, and *nito* were lethal,

suggesting that these subunits have important functions other than methylation (Penn et al.,

2008; Yan et al., 2015; Kan et al., 2017; Lence et al., 2017). RNA m^{6}A modification has an obvious

sexual dimorphism in *Drosophila*, and reduced m^{6}A levels severely decreased the survival of

females. It is thought to be due to the derepression of *msl-2* caused by aberrantly spliced *Sxl*,

which forms the Male Specific Lethal (MSL) complex that associates with the X chromosome

(Haussmann et al., 2016).

Dosage compensation is a widespread phenomenon in unbalanced genomes (Lucchesi, 2018;

Birchler and Veitia, 2021). The deletion or duplication of some chromosomes rather than the

whole chromosome set leads to genomic imbalance, that is, aneuploidy (Orr et al., 2015).

Aneuploid variation is usually detrimental to organisms (Birchler and Veitia, 2012; Orr et al., 2015;

Birchler and Veitia, 2021), and is associated with developmental abnormalities, mental

retardation, and various congenital defects (Williams et al., 2008; Huang et al., 2021;

Sanchez-Pavon et al., 2021), possibly due to disorders in their gene expression systems (Prestel et

al., 2010; Letourneau et al., 2014). Studies across species have pointed out that there is

genome-wide *trans* modulation in aneuploidy, and the genes on the varied chromosomes are

compensated to a certain extent, while genes located on the rest of the genome are mainly

regulated in the opposite direction to the changes of chromosome numbers, which is known as

the inverse dosage effect (Birchler and Veitia, 2012, 2021; Sun et al., 2013c; Hou et al., 2018; Shi

et al., 2021). Histone modification (Zhang et al., 2021a), chromatin remodeling (Birchler, 2016),

lncRNAs (Zhang et al., 2023) and microRNAs (Shi et al., 2022) have all been shown to play a role

in genomic imbalance.

Because of the haploinsufficiency for X-linked genes, heterogametic individuals in organisms

with XY sex determination systems could be regarded as analogous to aneuploidy (Disteche,

2016), including humans and *Drosophila*. Some studies have linked histone H4 lysine 16

acetylation (H4K16ac) and non-coding *roX* RNAs to the dosage compensation of *Drosophila* (Park

et al., 2010; Conrad et al., 2012). On the other hand, the compensation of X chromosome in

human is thought to be regulated by lncRNA X-inactive specific transcript (*XIST*) (Jordan et al.,

2019). Interestingly, in *Drosophila*, RNA m^6A modification indirectly affects gene expression

through *Sxl* and *msl-2*; while in human, m^6A methylation of the key lncRNA *XIST* is necessary for

the silencing of gene transcription on one of the female X chromosomes (Patil et al., 2016).

Moreover, most tumor cells have genomic instability and high levels of aneuploidy (Ben-David

and Amon, 2020; Chiarle, 2021), and at the same time, m^6A component genes are often

aberrantly expressed in various cancers (Pinello et al., 2018; Ma et al., 2019).

An increasing number of studies have found that dosage-related effects in aneuploidy may be

the integration of multiple modulations rather than through a single mechanism (Prestel et al.,

2010; Birchler, 2016). The effects of genomic imbalance are complicated, and the model of

dosage compensation and global gene regulation in *Drosophila* has been extended to autosomal

aneuploidies and sex chromosome aneuploid metafemales where MSL complexes are not

assembled (Sun et al., 2013b,c; Zhang et al., 2021b). In addition, genomic imbalance and *trans*

regulatory mechanisms in other species such as maize, *Arabidopsis*, and humans have also been

investigated (Hou et al., 2018; Raznahan et al., 2018; Shi et al., 2021; Yang et al., 2021; San

Roman et al., 2023). To reveal the role of RNA m^6A modification in unbalanced genomes, we

studied the dynamic changes and regulatory functions of m^6A methylation under genetic

imbalance conditions using autosomal and sex chromosome aneuploid *Drosophila* maintained in

our laboratory. Meanwhile, dosage-sensitive modifiers, differential alternative splicing events,

the MSL complex and other factors that may mediate the relationships between RNA m^6A

modification and the dosage-related effects of aneuploidy were also investigated. In summary,

we provided a comprehensive picture of RNA m^6A methylation in unbalanced genomes.

Results

The responses of m^{6}A components under genomic imbalance

RNA m^{6}A methylation is a reversible epigenetic modification, and its dynamic process is mediated by m^{6}A methyltransferases (writers), demethylases (erasers), and m^{6}A recognition proteins (readers) (Lee et al., 2014; Yang et al., 2018; Zaccara et al., 2019) (Figure 1A). We first detected the expression of m^{6}A components in *Drosophila* larvae with karyotypes of normal diploids and trisomies that have an additional chromosome arm using RT-qPCR to determine whether RNA m^{6}A dynamics are affected by correlative enzymes in unbalanced genomes (Figure 1B; Figure 1—figure supplement 1A,B). It was found that the transcription levels of most m^{6}A writers and the major m^{6}A reader are down-regulated in aneuploids compared with their respective sex-corresponding controls (Figure 1B). Unlike the other components, the cofactor *vir* of methyltransferase complex is up-regulated in trisomy 2L females (Haussmann et al., 2016; Lence et al., 2016; Kan et al., 2017). Because RNA m^{6}A modification is enriched in the nervous system of *Drosophila*, the brains of aneuploid third instar larvae were also used to detect the expression of m^{6}A components. As expected, a decreased expression trend of m^{6}A components similar to that of the whole larvae was observed (Figure 1—figure supplement 1C).

Previous studies have pointed out that the m^{6}A methylomes have temporal and spatial specificity in the development process of organisms, especially for embryonic development and cell differentiation (Meyer and Jaffrey, 2014; Wang Y et al., 2014). Aberrant expression of m^{6}A components may cause defects in embryogenesis and even early embryonic lethality (Zhong et al., 2008; Wang Y et al., 2014; Geula et al., 2015; Zhao et al., 2017). Therefore, we designed probes of m^{6}A components (Figure 1—figure supplement 1D,E) to examine the mRNA expression and localization patterns during early embryonic development of aneuploid *Drosophila* using high-resolution TSA-FISH (Lecuyer et al., 2007; Jandura et al., 2017). The results showed that the subembryonic distribution patterns of the five components of m^{6}A methyltransferase complex and one m^{6}A reading protein are similar in wildtype and aneuploidies (Figure 1C,D; Figure

1—figure supplement 2A-F). At the blastoderm stage (stage 1-5), the probe signals of m^{6}A

components are widely distributed in the surface cell layer, yolk plasma, yolk cortex, and show a

pattern of basal enrichment (Figure 1—figure supplement 2A-F). For the gastrulae at stage 6-11,

the transcripts of m^{6}A components are mainly located in head, amnioproctodeal invagination,

germ band, and as in previous studies (Lence et al., 2016), and show an enrichment in

neuroectoderm (Figure 1C; Figure 1—figure supplement 2A-F). At later stages of embryonic

development (stages 12-13 and 14-17), the probes are distributed in brain, ventral nerve cord,

midgut, and salivary gland (Figure 1C; Figure 1—figure supplement 2A-F).

By observing the subcellular localization of the probe signals of m^{6}A components at higher

magnification, it can be found that most of the mRNAs of these genes have nuclei-associated

localization patterns (Figure 1D; Figure 1—figure supplement 2A'-F'). In the early and late stages

of embryonic development, the probe signals of *Ime4*, *dMettl14*, *fl(2)d*, *vir*, and *Ythdc1* form

dense small foci near the nucleus, which is a perinuclear distribution (Figure 1D; Figure 1—figure

supplement 2A'-F'). For *nito*, in addition to the perinuclear signals, there is also an obvious signal

of intranuclear localization during early embryogenesis, which is manifested as one or two small

foci in the blastoderm nucleus (Figure 1D; Figure 1—figure supplement 2E').

Although there seems to be no difference in the localization of the transcripts of m^{6}A

components that we detected in aneuploidy and wildtype embryos, the expression levels of

these genes, as determined by relative fluorescence intensity, are significantly changed (Figure

1E,F; Figure 1—figure supplement 2A"-F"). Except for the irregular fluctuations at early

embryonic stages, the expression levels of most m^{6}A components in aneuploids are lower than

those in wildtype at more mature stages (Figure 1E,F; Figure 1—figure supplement 2A"-F"), which

is similar to the trend we detected in third instar larvae (Figure 1B; Figure 1—figure supplement

1C). The transcripts of m^{6}A methyltransferase and reader protein have specific subembryonic and

subcellular localization patterns in the development of *Drosophila* embryos, which may be a

mechanism to regulate cellular functions, and ensure appropriate cell growth and differentiation

(Lecuyer et al., 2007).

Genome-wide mapping of RNA m^{6}A methylation in aneuploid *Drosophila*

Subsequently, we detected the global levels of RNA m^{6}A methylation in wildtype and aneuploid

Drosophila larvae to determine whether m^{6}A abundance is altered under the condition of

genomic imbalance due to modulation by transmethylases. The overall abundance of m6A was

represented by the m^{6}A/A ratio in total RNA (Figure 2A). It was found that the m^{6}A abundance of

wildtype males is higher than that of females, but both are lower than 0.2%, which is consistent

with the finding that m^{6}A levels in *Drosophila* are relatively low (Haussmann et al., 2016; Lence et

al., 2016). Females with triple chromosome 2 left arms (2L) and metafemales with triple X

chromosomes have significantly higher m^{6}A abundances than diploid females, whereas there is

no significant difference between trisomy 2L males and wildtype males (Figure 2A). Therefore,

genomic imbalance can affect the m^{6}A methylation status to some extent, and this epigenetic

modification is different between males and females. However, the m^{6}A abundance in

aneuploidies did not follow the expression levels of transmethylases.

To obtain m^{6}A mapping across the transcriptome of aneuploid *Drosophila*, samples with

different karyotypes were used for m^{6}A methylated RNA immunoprecipitation sequencing

(MeRIP-Seq). By identifying consistent peaks in two biological replicates, approximately ten

thousand m^{6}A peaks were found in wildtype females and males, whereas there were less than

half the number of methylation sites in the three kinds of aneuploidies (Figure 2B). When these

m^{6}A sites were annotated according to the nearest genes, the changes in the number of

m^{6}A-marked genes in samples of different genotypes were in accordance with that of m^{6}A peaks,

in which there are about 2,500 m^{6}A-marked genes in wildtype and about 1,000 m^{6}A-modified

genes in trisomies (Figure 2C). Both the number of m^{6}A peaks and the number of m^{6}A-modified

genes are the least in trisomy 2L females (Figure 2B,C). The changes in the number of RNA

methylation sites in aneuploids did not coincide with the changes of overall m^{6}A abundance, but

instead matched the expression of m^{6}A components. We speculate that this may be caused by

the nonuniformity and heterogeneity of RNA m^{6}A modification, including the tissue specificity,

the developmental specificity, the different numbers of m^{6}A sites in one transcript, the different

proportions of methylated transcripts, et cetera (Meyer et al., 2012; Meyer and Jaffrey, 2014;

Zaccara et al., 2019). Analysis of m^{6}A-modified reads in wildtype and aneuploidy MeRIP samples

showed that the expression levels at m^{6}A sites in aneuploidies are significantly higher than that in

wildtype (Mann Whitney U test p-values < 0.001; Figure 2D), which confirmed our hypothesis.

These results suggest that RNA m^{6}A modification exhibits greater heterogeneity in unbalanced

genomes.

We next studied the overall characteristics of m^{6}A methylation in aneuploid and control

Drosophila. m^{6}A peaks are localized along all autosomes and sex chromosomes, and the numbers

of peaks are correlated with chromosome length (Figure 2—figure supplement 1A). Methylation

sites are widely distributed on 5'UTR (30-50%), 3'UTR (10-15%), promoter (15-20%), and internal

exons (20-35%), and the aneuploidies appeared to have a higher proportion of 5'UTR peaks and a

lower proportion of exon peaks than the wildtype (Figure 2E). The highest values of density

distributions of the length of exons containing m^{6}A peaks are around 250 bp, and approximately

90% of the exons are longer than the typical 140 bp (Figure 2—figure supplement 1B). *De novo*

motif analysis of m^{6}A methylation peaks in each sample showed that the top-ranked motif is

consistent with the conserved m^{6}A motif DRACH (D=G/A/U, R=G/A, H=U/A/C; Figure 2F). In

addition, enrichment analysis of the known motif DRACH showed that it is significantly enriched

in all samples, and more than 97% of the m^{6}A sites have this sequence (Figure 2—figure

supplement 1C).

There are 370 common genes to which the m^{6}A peaks were annotated in all genotypes

(Figure 2G). We classified the m^{6}A-modified genes in each sample and found that they were

mostly protein-coding genes, whereas only about 7% were ncRNAs (Figure 2H). As described in

previous studies (Meyer et al., 2012; Zaccara et al., 2019), most methylated RNAs have one m^{6}A

peak (Figure 2—figure supplement 1D-H). However, there are still some genes whose transcripts

can be highly methylated and contain more than five m^{6}A sites (Figure 2—figure supplement

1D-I). It was found that transcription factors were significantly enriched in 70 genes with more

than 5 m^{6}A peaks in both wildtype and aneuploidy (Fisher's exact test p-value = 7.7e-4; Figure 2I;

Figure 2—figure supplement 1I). Thus, specific types of genes may be preferentially targeted by

m^{6}A modification, such as transcriptional regulators (Kan et al., 2017). Analysis of the functions of

m^{6}A-modified genes in each sample revealed that they were enriched for a large number of

common functions, among which the most significant were those related to morphogenesis,

development, and growth (Figure 2—figure supplement 2A). The 141 GO terms shared by the

five genotypes were summarized and found to include functions about metabolism, regulation,

cellular processes, signaling pathways, behavior, and immune response (Figure 2—figure

supplement 2B,C). Through pathway enrichment analysis of m^{6}A-modified genes, 9 terms were

found to be consistently enriched in all samples, including MAPK signaling pathway, Hippo

signaling pathway, TGF-β pathway, etc. (Figure 2—figure supplement 2D-F).

Differentially methylated peaks (DMPs) and their associated genes

We then searched for the differential methylation peaks (DMPs) in autosomal and sex

chromosome aneuploidies compared with wildtype *Drosophila* of their corresponding sex. The

number of DMPs and the changing trend of DMP methylation status are shown in the Figures

(Figure 3A-D). In trisomy 2L females, the number of upregulated DMPs (1,131) is less than that of

downregulated DMPs (1,602). In trisomy 2L males, the number of upregulated DMPs (1,521) is

slightly higher than that of downregulated DMPs (1,259). In metafemales, the number of

upregulated DMPs (1,393) is higher than that of downregulated DMPs (926) (Figure 3A-D). More

DMP-associated genes were found in autosomal aneuploidies than in sex chromosome

aneuploidy (trisomy 2L female = 2,148, trisomy 2L male = 2,055, metafemale = 1,771; Figure 3E).

Also, the total number of peaks after the differential analysis would be different from the number

of m^{6}A peaks identified in the previous section, because the analysis software merged and

recalculated the peaks (Stark and Brown, 2013).

DMPs are distributed on all chromosomes (Figure 3—figure supplement 1A). Specifically, the

MeRIP-Seq signal intensity at the location of DMPs showed that the m^{6}A-marked reads in

trisomies are denser than those of wildtypes, especially on the triple chromosomes (Figure 3F-H).

The density of reads on chromosome 2L of trisomy 2L *Drosophila* is higher than that on other

chromosomes, while the density of reads on chromosome X of metafemales is significantly higher

than that of autosomes (Figure 3F-H). Previous studies on aneuploid *Drosophila* showed that the

expression levels of genes on chromosomes with increased copy numbers were mostly similar to

those in diploids, and the expression of genes on unvaried chromosomes were widely inverse

regulated (Sun et al., 2013c; Birchler, 2016; Zhang et al., 2023). It can be concluded that the

changes in the density of reads at DMP sites are not due to a direct gene dosage effect. Similar

results were obtained by counting the methylation status of cis and trans transcripts (Figure 3I-K).

Overall, the transcripts of genes located on the varied chromosomes in three trisomies have

more up-regulated DMPs than down-regulated DMPs, which is more obvious in metafemales

(Figure 3I-K). In trisomy 2L females and males, genes localized on other autosomes have a higher

proportion of down-regulated DMPs, while the X-linked *trans* genes show sexual dimorphism and

X chromosome-specific response to genomic imbalance. (Figure 3I,J). For metafemales, the

transcripts of *trans* genes have similar numbers of up-regulated and down-regulated DMPs

(Figure 3K). Therefore, *cis* genes in trisomy generally possessed a higher proportion of

up-regulated DMP-associated transcripts, whereas the methylation states of *trans* genes are

variable, depending on the identities of varied chromosomes and their genomic locations. The

above results give us reason to speculate that the changes of m^{6}A methylation may affect dosage

compensation of *cis*-genes and inverse dosage modulation of *trans*-genes in aneuploidy.

The distribution of DMPs along gene features includes 5'UTR (30-35%), 3'UTR (20-25%),

promoter (10-25%), and internal exon (25-35%), among which the proportion of 3'UTR is higher

than that of all m^{6}A sites (Figure 3—figure supplement 1B). The length distributions of exons with

DMPs are similar to that of all m^{6}A-modified exons (Figure 3—figure supplement 1C). *De novo*

motif analysis of the sequences where DMPs are located or enrichment analysis of the known

motif DRACH both showed that DMP sites are enriched for the conserved motif of m^{6}A (Figure

3—figure supplement 1D,E). There are also similarities between the types of DMPs-associated

genes and m^{6}A-marked genes (Figure 3—figure supplement 1F). The DMP-associated genes of

the three aneuploidies have complex overlapping relationships (Figure 3—figure supplement 1G).

Most DMP-associated genes have one differentially methylated site, and only a few genes

contained more than three DMPs (Figure 3—figure supplement 1H-J). There are 1,042 common

DMP-associated genes in all trisomies, of which 71 were transcription factors and were

significantly enriched (Fisher's exact test p-value = 6.9e-5; Figure 4A,B). We speculate that these

differentially methylated transcription factors may play an important role in the regulation of

gene expression and development of aneuploidy.

Functional enrichment analysis showed that genes with up-regulated DMPs in three trisomies

were enriched for 153 common functions, while the genes with down-regulated DMPs shared

only 6 functions (Figure 4—figure supplement 1A,B). Similarly, genes with up-regulated DMPs

were enriched for more consistent pathways in all aneuploids (Figure 4—figure supplement 1C,D).

Most of the functions of DMP-associated genes are the same as those of m^{6}A-marked genes, and

these genes are also enriched for cell fate commitment, dorsal/ventral pattern formation, sex

differentiation, post-transcriptional regulation of gene expression, and other additional GO terms.

We found that many DMP-associated genes have functions related to gene expression regulation

and dosage compensation, including some components of MSL complex (*msl-1*, *msl-3*,

lncRNA:roX1) and the major sex-determining factor *Sxl* (Figure 4C). We also noticed that m^{6}A

component genes *fl(2)d*, *nito*, and *Ythdc1* themselves are differentially methylated in aneuploidy

(Figure 4C), for example, there was a common significantly upregulated m^{6}A peak in the 5'UTR of

the transcripts of *fl(2)d* in three aneuploidies (Figure 4D). These data suggest that m^{6}A dynamics

in aneuploids may be involved in the expression regulation of unbalanced genomes through

various genes with regulatory functions.

304   Relationships between m^{6}A and dosage-related modulation of gene expression in

305   aneuploidy

To explore the relationships between RNA m^{6}A modification and gene expression regulation

under genomic imbalance, we combined the results of MeRIP-Seq and RNA-seq, and performed a

comprehensive analysis. The results showed that about 12%-15% of genes differentially

expressed in aneuploid *Drosophila* also belong to DMPs-associated genes (Figure 5A-C). However,

except for trisomy 2L males, the DEGs of the other two trisomies did not show enrichment of

differential m^{6}A methylation (Fisher's exact test p-values: trisomy 2L male = 0.012, trisomy 2L

female and metafemale > 0.05). There are 55 genes that are both differentially expressed and

differentially methylated in all aneuploidies (Figure 5D). By analyzing the functions of genes that

are simultaneously differentially expressed and differentially methylated, we obtained a number

of consistent biological functions, including post-embryonic animal morphogenesis, cell

communication, signal transduction, response to stimulation, and so on (Figure 5E), possibly

reflecting the important roles of m^{6}A-regulated expression in cell signaling and development of

organisms.

By analyzing the crossover of different groups of m^{6}A-modified genes and genes with

canonical dosage effect, dosage compensation, and inverse dosage effect, we further revealed

the relationships between m^{6}A-modified genes and dosage-related effects in aneuploidy (Figure

5F-H). The results showed that in trisomy 2L females, *cis* dosage compensation genes, *trans*

autosomal dosage effect genes, and *trans* unchanged genes are significantly enriched in m^{6}A

group with methylation in both trisomy and control; meanwhile, *cis* dosage effect genes, other

autosomal dosage effect genes, and X-linked dosage effect genes are enriched in the group

without methylation in both trisomy and control (Figure 5F). On the contrary, in trisomy 2L males,

cis dosage effect genes, *trans* dosage effect genes, and *trans* unchanged genes are significantly

enriched in m^{6}A group whose genes are methylated in both trisomy and control; while the group

of genes that are not methylated at all is enriched for *cis* dosage effect genes, *cis* dosage

compensation genes, *trans* inverse dosage effect genes, and other autosomal dosage effect genes

(Figure 5G). Metafemales performed similarly to trisomy 2L female, with significant enrichment

of *cis* dosage compensation genes, autosomal *trans* dosage effect genes, and *trans* unchanged

genes in m^{6}A group where both trisomy and control genes are methylated; and the group

without methylation is enriched for *cis* dosage effect genes and *trans* dosage effect genes (Figure

5H). These results indicated that m^{6}A-modified genes in aneuploid *Drosophila* females are mainly

related to dosage compensation and inverse dosage effect, while genes not modified by m^{6}A are

mainly related to direct dosage effect. Male aneuploids did not follow this trend.

Furthermore, the distributions of m^{6}A modification sites along gene features in genes with

classical dosage-related effects were also studied (Figure 5I-K). We found that in wildtype females,

trisomy 2L females, and metafemales, a higher proportion of m^{6}A sites on genes with canonical

dosage compensation and inverse dosage effect are distributed in the 5'UTR than genes with

dosage effects (Figure 5I,K). At the same time, this proportion is higher in trisomy than in

wildtype. In contrast to females, trisomy 2L males have a higher proportion of 5'UTR RNA m^{6}A

modification on dosage effect genes (Figure 5J). The above study again demonstrates the sexual

dimorphism of RNA m^{6}A modification in response to aneuploidy. In addition, it has been

suggested that m^{6}A residues in the 5'UTR region may have unique regulatory functions (Luo et al.,

2014; Meyer and Jaffrey, 2014). Therefore, we hypothesized that the high level of 5'UTR m^{6}A

modification on genes with dosage compensation and inverse dosage effect might regulate the

down-regulation of these genes.

Dozens of dosage-sensitive modifiers have been identified in *Drosophila*, whose dosage

changes can negatively or positively regulate the gene expression across the whole genome like

aneuploidy (Birchler et al., 2001; Birchler and Veitia, 2007). Protein-protein interaction (PPI)

networks between dosage-sensitive modifiers and differentially expressed genes (DEGs) were

constructed to investigate whether RNA m^{6}A modification could participate in the gene

regulatory networks in unbalanced genomes through these regulators (Figure 5—figure

supplement 1A-C). In the three types of aneuploidies, there are more dosage-sensitive modifiers

with up-regulated DMPs than with down-regulated DMPs. The regulators *ox* and *Vha55* without

DMPs interact with a large number of significantly up-regulated DEGs in trisomy 2L females and

metafemales, whereas these two regulators interact with a small number of down-regulated

DEGs in trisomy 2L males. Up-regulated DMP-associated regulators *wg*, *Uba1*, *ap*, *Atg1*, *osa*, *rdx*,

and *sd* in trisomy 2L females mainly interacted with significantly down-regulated DEGs (Figure

5—figure supplement 1A); and up-regulated DMP-associated regulators *Uba1*, *osa*, and *sd* in

metafemales also interacted with down-regulated DEGs (Figure 5—figure supplement 1C).

Up-regulated DMP-associated regulators *wg*, *Kr-h1*, and *Trl* mainly interacted with up-regulated

DEGs in trisomy 2L males (Figure 5—figure supplement 1B). In addition, a preponderance of

significantly down-regulated DEGs interacted with dosage-sensitive modifiers is observed on

trans chromosomes of all *Drosophila* trisomies (Figure 5—figure supplement 1A-C).

Alterations of m^{6}A may be involved in differential alternative splicing in imbalanced

genomes

Next, we analyzed the differential alternative splicing events in aneuploid *Drosophila*. More than

1,000 differential splicing events have been identified in different aneuploids, and about

one-third of them are of the type of skipped exon (SE) (Figure 6—figure supplement 1A-C). The

biological functions of differentially spliced transcripts are mainly involved in macromolecular

fiber organization, locomotion, and growth (Figure 6—figure supplement 1D), and these genes

are enriched in heterogeneous pathways in different aneuploidies (Figure 6—figure supplement

1E). Notably, we found that genes with DMPs are significantly enriched for differential alternative

splicing events in three kinds of aneuploid *Drosophila* (Fisher's exact test p-values < 0.05; Figure

6A-C). Among the genes whose transcripts are differentially spliced, 27%-32% are also

differentially methylated (Figure 6A-C). There are 67 genes with both differential alternative

splicing and differential m^{6}A methylation in all aneuploidies (Figure 6D). The functions of these

genes are similar to those of all differentially spliced genes (Figure 6E), but more consistent KEGG

pathways are enriched, including endocytosis, mTOR signaling pathway, Hedgehog signaling

pathway, and so on (Figure 6F).

In all trisomy *Drosophila*, 10 genes are shown to be differentially expressed, with their

transcripts also being differentially spliced and differentially methylated. Three of them are

illustrated below [*su(Hw)*, *Ppn*, and *CG13124*]. *su(Hw)* is a component of the gypsy chromatin

insulator complex, which is a regulatory element that establish independent domains of

transcriptional activity (Roseman et al., 1993). Due to the close relationship between *su(Hw)* and

second-site modifiers (Rabinow and Birchler, 1989), and *BEAF-32*, which is also an insulator

DNA-binding protein, has been proposed as a possible inverse dosage regulator (Gurudatta et al.,

2012; Zhang et al., 2021b), We speculated that the transcription factor *su(Hw)* may also be a

dosage-sensitive regulator. The data showed that the transcription levels of *su(Hw)* are

up-regulated in all three aneuploidies, and there is a consistent m^{6}A modification site with

significantly down-regulated methylation (Figure 6G). At the same time, the transcripts of this

gene have a common alternative 5' splice site (A5SS), and its inclusion levels in trisomies are

down-regulated, that is, more short transcript isoforms are generated (Figure 6G). *Ppn* gene

encodes an essential extracellular matrix protein that influences cell rearrangements. The

expression level of *Ppn* and its m^{6}A methylation in 5'UTR region are both significantly

up-regulated in trisomies (Figure 6—figure supplement 1F). In addition, an alternatively spliced

exon is significantly less frequently skipped in all aneuploid *Drosophila* (Figure 6—figure

supplement 1G). The third gene, *CG13124*, which may be involved in involved in regulation of

translational initiation, is up-regulated in aneuploids. Two m^{6}A sites in its 5'UTR region are

methylated at higher levels in all aneuploid *Drosophila* (Figure 6—figure supplement 1H). Its

transcripts also have a common significantly different exon-skipping event (Figure 6—figure

supplement 1I).

These results indicate that there are complicated relationships among RNA m^{6}A modification,

gene expression, and alternative splicing under the condition of genome imbalance. RNA splicing

seems to be more closely related to m^{6}A methylation than gene transcription. The m^{6}A sites

located in 5'UTR show remarkable changes in methylation levels in aneuploid *Drosophila*, and

may be involved in the regulation of some differential alternative splicing events, such as exon

skipping.

Interactions between m^{6}A and _Drosophila_ Male Specific Lethal (MSL) complex

Previous studies have shown that m^{6}A components are involved in regulating the alternative

splicing of sex-determining gene _Sxl_ in _Drosophila_, and the deficiency of m^{6}A writers or readers

will lead to the reduction of female-specific isoforms of _Sxl_ (Haussmann et al., 2016; Lence et al.,

2016; Kan et al., 2017). _Sxl_ is also a direct target of RNA m^{6}A modification (Kan et al., 2017). We

found that _Sxl_ transcripts are both differentially methylated and differentially spliced in three

kinds of aneuploid _Drosophila_ (Figure 7A,B). For trisomy 2L females and metafemales, two

common m^{6}A peaks are significantly up-regulated in the 5'UTR region of _Sxl_. However, trisomy 2L

males have a significantly down-regulated m^{6}A peak in the 5'UTR (Figure 7A). Meanwhile,

multiple junctions of _Sxl_ transcripts undergo complicated alternative splicing in aneuploid

Drosophila, including SE, A5SS, alternative 3' splice site (A3SS), and mutually exclusive exons

(MXE) types (Figure 7A). By checking the distributions of RNA sequencing reads near the

male-specific exon (namely the third exon), it can be observed that there are almost no mapped

reads on the third exon in wildtype females, trisomy 2L females, and metafemales, while the

reads mapped to the second and fourth exons are highly prevalent (Figure 7B). On the contrary,

wildtype males and trisomy 2L males have a substantial number of reads on the third exon,

accompanied by a smaller number of reads on the adjacent two exons (Figure 7B). Notably, we

found that a small number of RNA-seq reads aligned to the male-specific exon appeared in

trisomy 2L females, which was identified as an SE-type differential alternative splicing event (FDR

= 7.4e-7; Figure 7B). This variation may be related to the abnormal m^{6}A methylation levels under

the condition of genomic imbalance.

The abnormal expression of m^{6}A components reduces the survival of female _Drosophila_,

which is thought to be probably caused by the expression of downstream MSL complex

(Haussmann et al., 2016). To investigate the interplay between the MSL complex and m^{6}A

modification, we examined the responses of m^{6}A regulators in transgenic _Drosophila_ strains with

MSL2 mutation or overexpression (Figure 7—figure supplement 1A-E; Figure 7C,D). The results

revealed significant changes in the expression profiles of m^{6}A regulators in MSL2 transgenic

strains, especially the m^{6}A reader protein Ythdc1, which increased tens of fold in MSL2

knockdown and overexpressed *Drosophila* (Figure 7C,D; Figure 7—figure supplement 1E). We

also observed that the trends of Ime4 expression in females and males were the same in

MSL2-overexpressed *Drosophila*, whereas there was obvious sexual dimorphism in

MSL2-knockdown samples (Figure 7C,D), which may be due to the ectopic assembly of MSL

complex in MSL2-overexpressed females (Zhang et al., 2021a). In addition, we also examined the

expression levels of MSL2 when *Ythdc1* was knocked down, and found that MSL2 was also

significantly increased in females (Figure 7—figure supplement 1F). All these results strongly

suggest a potential relationship between MSL2 and Ythdc1. Next, we further compared the

overall abundance of m^{6}A in MSL2 transgenic and wildtype *Drosophila* (Figure 7E,F). The results

showed that m^{6}A levels were significantly decreased with MSL2 knockdown and significantly

increased with MSL2 overexpression in females (Figure 7E). However, overexpression of MSL2

failed to exert a discernible effect on m^{6}A abundance in males, although m^{6}A level decreased

with MSL2 depletion (Fig.7F). This difference between the sexes may be due to the endogenous

MSL2 in males, but not in females.

In the next, we investigated the expression of m^{6}A regulators in aneuploid *Drosophila*

overexpressing MSL2, according the results that the MSL complex could be regulated directly or

indirectly by m^{6}A modification, and the sexual dimorphism of RNA m^{6}A modification in response

to aneuploidy. It is found that the transcription levels of m^{6}A regulators are significantly

up-regulated in the brains of most aneuploid larvae that overexpressed MSL2 (Figure 7G; Figure

7—figure supplement 1G). These results are not completely consistent with the quantitative

results when MSL2 was overexpressed in diploids, which may be related to the effect of

unbalanced genomes. We also used TSA-FISH to detect the expression and distribution of m^{6}A

components during embryogenesis in aneuploidies overexpressing MSL2 (Figure 7H; Figure

1—figure supplement 2). The subembryonic and subcellular distributions of mRNAs for m^{6}A

methyltransferases and reading protein did not appear to be affected by ectopic expression of

MSL2 in aneuploid *Drosophila* embryos (Figure 1—figure supplement 2A-F; Figure 1—figure

supplement 2A'-F'). But the relative expression of m^{6}A components showed diverse dynamics,

among which the levels of the most important methyltransferase Ime4 are significantly

up-regulated in autosomal trisomy and sex chromosome trisomy with MSL2 overexpression

(Figure 7H; Figure 1—figure supplement 2A"-F"). Considering that unbalanced genomes can

affect the expression of MSL complex subunits, and the above results show that there is a close

relationship between MSL complex and m^{6}A modification, we speculate that unbalanced

genomes may influence the expression of m^{6}A regulators, possibly through the MSL complex.

476 Relationship of H4K16Ac with m^{6}A modification

As an important component of the MSL complex, MOF is a histone acetyltransferase that

specifically acetylates histone H4 at lysine 16 (H4K16Ac), thereby affecting chromatin structure

and functions, and activating transcription (Kind et al., 2008; Conrad et al., 2012). Previous

studies have elucidated the regulatory role of m^{6}A in histone modifications, and histone 3 lysine

36 trimethylation (H3K36me3) also plays a role in recruiting methyltransferase complexes to

deposit m^{6}A markers on RNA (Wang et al., 2018; Huang et al., 2019; Li et al., 2020). To investigate

the functions of MOF-mediated H4K16Ac on RNA m^{6}A modification, we analyzed RNA-seq data

from *Drosophila* strains overexpressing MOF (Figure 8A; Figure 7—figure supplement 1H-J). We

found obvious changes in the expression of m^{6}A regulators in strains overexpressing MOF, with

the levels of almost all m^{6}A regulators being elevated (Figure 8A). It was also observed that after

overexpression of MOF, the increasing trends of m^{6}A regulators in females and males were not

exactly the same, mostly showing more pronounced increases in females, except for Mettl3

(Ime4) and dMettl14. According to the results of RT-qPCR, the expression of Ythdc1 was

decreased in MOF knockdown strains and increased in MOF overexpression strains (Figure 8B;

Figure 7—figure supplement 1J); meanwhile, the expression levels of MOF were significantly

reduced in *Ythdc1* knockdown strains (Figure 8C). These results were further verified by polytene

chromosome immunofluorescence experiments (Figure 8D,E). In males, the expression level of

MOF was significantly decreased in *Ythdc1* knockdown *Drosophila*; while in *Ythdc1* knockdown

females, the expression level of MOF was not significantly changed compared with wildtype

females (Figure 8D,E). These results suggest that MOF-mediated acetylation modification can

have a certain effect on RNA m^{6}A methylation in a sexually dimorphic manner, which may be due

to the endogenous MSL complex and the imbalance of X chromosome dosage in males.

Next, we investigated whether the m^{6}A reader protein Ythdc1 would also have an effect on

the levels of H4K16Ac in *Drosophila*. Quantitative analysis of polytene chromosome

immunofluorescence results in *Ythdc1* knockdown *Drosophila* showed that the changes of

H4K16Ac levels also showed sexual dimorphism (Figure 8F,G). In males, knockdown of *Ythdc1* led

to a decrease in H4K16Ac level; whereas in females, knockdown of *Ythdc1* did not affect the level

of H4K16Ac (Figure 8F,G). These observed changes in H4K16Ac were consistent with the pattern

of changes in MOF protein in *Ythdc1* knockdown *Drosophila* strains, suggesting that m^{6}A

modification may affect H4K16Ac levels through the mediation of MOF. To further study whether

unbalanced genomes are involved in the interaction between m^{6}A and histone acetylation

modification, we also analyzed RNA sequencing data from trisomy 2L *Drosophila* strains

overexpressing MOF. The data showed that overexpression of MOF in trisomy 2L resulted in

significant changes in the expression of m^{6}A regulators, with a trend different from that observed

in diploids (Figure 8A). These results suggest that genomic imbalance might affect the interaction

between MOF and m^{6}A regulators to some extent, and the potential mechanisms require further

investigation.

Discussion

As an emerging epigenetic modification, RNA m^{6}A methylation has been found to be involved in

almost all aspects of RNA fate and metabolism (Gilbert et al., 2016; Yang et al., 2018). RNA m^{6}A

modification is also closely related to the development of organisms and a variety of human

diseases (Barbieri et al., 2017; Zhao et al., 2017; Pinello et al., 2018; Ma et al., 2019; Liu et al.,

2021; Shafik et al., 2021). However, the roles of m^{6}A methylation in development and gene

expression of aneuploidy have not been studied yet. Aneuploid variation is usually more

detrimental than changes of the entire chromosome set due to genomic imbalance (Birchler and

Veitia, 2007, 2012). The global changes of gene regulatory networks in unbalanced genomes

involves various epigenetic mechanisms, including histone modification, chromatin remodeling,

lncRNAs, microRNAs, etc. (Birchler, 2016; Zhang et al., 2021a, 2023; Shi et al., 2022). This study

demonstrated that the expression of m^{6}A components was altered under genomic imbalance,

leading to dynamic changes in the entire methylome. Potential intermediaries by which m^{6}A

modification could affect *trans* regulation and achieve dosage compensation in aneuploid

Drosophila were also investigated, such as dosage-sensitive modifiers, alternative splicing events,

and the MSL complex.

Our experiments show that the expression levels of most m^{6}A component genes are

significantly down-regulated in aneuploid *Drosophila* larvae (Figure 1A; Figure 1—figure

supplement 1C). Depletion of m^{6}A components interferes with the development of animals and

plants, especially leading to impaired self-renewal and differentiation of embryonic stem cells,

defects in embryonic development, and even early embryonic lethality (Granadino et al., 1990;

Horiuchi et al., 2006; Raffel et al., 2007; Luo et al., 2014; Wang Y et al., 2014; Geula et al., 2015).

We demonstrated by TSA-FISH that appropriate temporal and spatial specific distributions of the

transcripts for m^{6}A components are vital during *Drosophila* embryogenesis (Figure 1C-F; Figure

1—figure supplement 2). The abnormal expression of m^{6}A-related genes may affect the

development of aneuploid embryos.

In previous studies, the abundance of m^{6}A modification is usually positively correlated with

the number of m^{6}A peaks and m^{6}A-marked genes (Luo et al., 2014; Zhu et al., 2023). However,

our results obviously did not conform to this rule, with higher m^{6}A abundance and fewer

MeRIP-Seq peaks in aneuploids (Figure 2A-C). This reflects the complexity and heterogeneity of

m^{6}A modification. We suspect that in aneuploidy many RNAs may be lost that are methylated at

a low level in wildtype, and possess a higher proportion of highly methylated RNAs. Analysis of

the expression levels at each m^{6}A site confirmed our hypothesis (Figure 2D). Thus, this

phenomenon represents an imbalance in m^{6}A methylation caused by aneuploidy.

The distribution of m^{6}A sites on gene features, m^{6}A consensus motifs, biotypes of

m^{6}A-marked genes, enriched functions and pathways of methylated genes in aneuploidies are

similar to those in wildtype. We also found that genes highly methylated in all genotypes are

enriched for transcription factors (Figure 2I). It is consistent with previous studies suggesting that

transcription regulatory genes may be preferentially targeted by m^{6}A (Kan et al., 2017).

We also analyzed the characteristics of DMPs and DMP-associated genes in trisomies. By

observing the distributions of MeRIP-Seq reads around DMP sites, it can be found that the

densities of m^{6}A-marked reads on chromosome 2L are relatively higher in trisomy 2L females and

males (Figure 3F,G). Meanwhile, metafemales with triple X have a higher density of m^{6}A-marked

reads on chromosome X (Figure 3H). Previous studies have found that there is dosage

compensation for *cis*-genes in autosomal and sex chromosome aneuploid *Drosophila*, and the

expression of most genes on varied chromosomes approaches diploid levels (Sun et al., 2013b,c).

Therefore, the up-regulation of m^{6}A levels is not directly caused by the increased number of

chromosomes. A recent study found that the selective enrichment of m^{6}A methylation may play a

role as a transcript degradation signal in dosage compensation in mammals (Rücklé et al., 2023).

Therefore, we speculate that changes in m^{6}A levels in aneuploid *Drosophila* may affect its dosage

compensation and inverse dosage effect by regulating the stability of the transcripts. In addition,

more up-regulated DMPs are detected in trisomy 2L males and metafemales, while trisomy 2L

females have more down-regulated DMPs (Figure 3A-D). Combined with the fact that the survival

rate of trisomy 2L female larvae is relatively lower than that of the other two trisomies, it can be

speculated that the regulation of DMPs may affect the survival and development of aneuploid

Drosophila.

Gene expression in unbalanced genomes is extensively modulated (Sun et al., 2013b,c; Hou et

al., 2018; Raznahan et al., 2018; Shi et al., 2021; Yang et al., 2021; Zhang et al., 2021b; San Roman

et al., 2023). Previous studies have analyzed the transcriptome data of autosomal and sex

chromosome trisomic *Drosophila*, and found that most of the genes on the triple chromosomes

were compensated, and the ratios of their expression levels to wildtype were approximately 1;

meanwhile, the expression of genes on other chromosomes was close to two-thirds of that of the

wildtype, which was called an inverse dosage effect (Sun et al., 2013b,c; Zhang et al., 2021b). We

investigated the relationships between genes with different m^{6}A methylation status and genes

modulated by classical dosage-related effects (Figure 5F-H). The results showed that for

aneuploid females, genes methylated in both trisomy and control are significantly enriched in

dosage compensated *cis* genes and inverse dosage effect *trans* genes, and genes not methylated

at all are enriched in dosage effect genes. However, in aneuploid males, methylated genes are

associated with gene dosage effect (Figure 5F-H). We also found that the proportion of 5'UTR

m^{6}A peaks on dosage compensation and inverse dosage effect genes are increased in aneuploid

females (Figure 5I-K). These results provide evidence of sexual dimorphism in the relationships

between RNA m^{6}A modification and dosage-related effects. m^{6}A located in the 5'UTR generally

shows higher tissue specificity and richer dynamic changes, and may have unique regulatory

functions (Dominissini et al., 2012; Meyer and Jaffrey, 2014; Gilbert et al., 2016). The increased

proportion of m^{6}A in the 5'UTR of genes with classical dosage-related effects in aneuploidies

suggests that m^{6}A modification may be involved in the regulation of dosage-dependent genes.

Dosage-related effects of aneuploidy are thought to be caused by dosage-sensitive genes (Shi

et al., 2021; Yang et al., 2021). Some dosage-sensitive regulators have been identified in

Drosophila, and changes in the dosage of individual regulatory genes can mimic the effects of

aneuploidy in the whole genome (Birchler et al., 2001; Xie and Birchler, 2012; Zhang et al., 2021b).

Most dosage-sensitive regulators are transcription factors, signal transduction components, and

chromatin proteins, which have in common being members of macromolecular complexes or

having multicomponent interactions (Birchler et al., 2001; Birchler and Veitia, 2007). We studied

the m^{6}A modification of dosage-dependent genes and their protein-protein interaction networks

in aneuploid *Drosophila* (Figure 5—figure supplement 1). The results showed that there are

complex interactions between dosage-sensitive regulators and differentially expressed genes.

Among them, most of the DEGs located on the unvaried chromosomes are down-regulated,

indicating that the dosage-sensitive regulators mainly have negative effects on *trans* target genes.

There are more regulatory genes with up-regulated DMPs than down-regulated DMPs in all three

aneuploidies, and many of the regulators with up-regulated methylation are connected with

interactors that have down-regulated expression.

We comprehensively analyzed the relationships among RNA m^{6}A modification, gene

expression levels, and alternative splicing. The data showed that differential m^{6}A methylation

under genomic imbalance appeared to be more closely associated with differential alternative

splicing (Figure 5A-C; Figure 6A-C). This phenomenon is reasonable because the mutation or

knockout of *Ime4*, *dMettl14*, *fl(2)d*, and *Ythdc1* has been found to affect a large number of

alternative splicing events in *Drosophila*, and *fl(2)d* itself is thought to encode a splicing factor

(Penn et al., 2008; Haussmann et al., 2016; Lence et al., 2016). A small set of transcripts are

simultaneously differentially m^{6}A methylated, differentially expressed, and differentially

alternative spliced in aneuploidies, including the transcription factor *su(Hw)*, which may be a

dosage-sensitive regulator (Figure 6G-H). Besides, m^{6}A modification in the 5'UTR regions may

play a special role in some differential alternative splicing events (Figure 6—figure supplement

1F-I).

RNA m^{6}A modification in *Drosophila* has been shown to be involved in the alternative splicing

of *Sxl* (Lence et al., 2017). The deletion of some m^{6}A components in females will result in a

reduction of female-specific splicing of *Sxl* and an increase of the inclusion of the third exon,

along with phenotypic sexual transformation (Haussmann et al., 2016; Lence et al., 2016; Kan et

al., 2017). We found that *Sxl* transcripts of aneuploid *Drosophila* are both differentially m^{6}A

methylated and differentially spliced (Figure 7A-B). Among multiple alternative splicing events,

trisomy 2L females have a higher level of the third exon inclusion compared with wildtype

females (Figure 7B), which may be related to the differential methylation at 5'UTR of the *Sxl*

transcripts.

Some studies proposed that the deletion of RNA m^{6}A methyltransferase Ime4 harms the

survival of female *Drosophila* because of the insufficient inhibition of *msl-2* caused by decreased

Sxl levels, which in turn leads to up-regulation of X-linked genes (Haussmann et al., 2016).

However, other studies have found that the expression of genes on the X chromosome in females

with ectopic expression of MSL2 does not increase twofold, that is, the MSL complex does not

directly mediate dosage compensation, and the global inverse dosage effect caused by the

imbalance of sex chromosomes is the basis for dosage compensation (Birchler, 1981, 2016; Sun et

al., 2013a). Therefore, the lethality of the lack of m^{6}A in female *Drosophila* may not be directly

caused by up-regulation of X-linked genes through ectopic assembly of MSL complex, and the

specific reasons need to be further studied.

The functions of MSL complex in dynamic regulation of global gene expression in aneuploid

genomes has been described (Zhang et al., 2021a). Here, we found that MSL2, a structure

component of MSL complex, not only affects the expression of m^{6}A regulators, but also

influences the overall abundance of m^{6}A in *Drosophila* (Figure 7C-F), proving a close relationship

between the MSL complex and RNA m^{6}A modification. In addition, under the condition of

genomic imbalance, the relative expression of m^{6}A components was also changed in larvae and

embryos of MSL2-overexpressed trisomies (Figure 7G,H). Another component of the MSL

complex, MOF, which mediates histone acetylation and transcriptional activation (Kind et al.,

2008; Conrad et al., 2012), is also closely associated with the m^{6}A reader Ythdc1. Overexpression

of MOF increased the expression of Ythdc1 (Figure 8A,B), and in turn, knockdown of *Ythdc1*

reduced the expression of MOF and the level of H4K16Ac catalyzed by it (Figure 8C-G). These

results demonstrate complicated interactions between RNA m^{6}A methylation and the MSL

complex in unbalanced genomes, which may affect the gene expression, sexual dimorphism,

development and survival of aneuploid *Drosophila*.

Materials and methods

655   **Key resources table**

656   Softwares used in this study:

Resource	Designation	Source or reference
software, algorithm	R v4.2.2	http://www.R-project.org
software, algorithm	trim-galore v0.6.10	https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
software, algorithm	FastQC v0.12.1	https://www.bioinformatics.babraham.ac.uk/projects/fastqc/
software, algorithm	HISAT2 v2.2.1	Kim et al., 2019
software, algorithm	StringTie v2.1.5	Pertea et al., 2015
software, algorithm	exomePeak2 v1.10.0	Meng et al., 2013
software, algorithm	HOMER v4.11	Heinz et al., 2010
software, algorithm	DiffBind v3.8.4	Stark and Brown, 2013
software, algorithm	ChIPseeker v1.34.1	Yu et al., 2015
software, algorithm	deepTools v3.5.4	Ramírez et al., 2014
	rMATS v4.2.0	Shen et al., 2014
software, algorithm	ggplot2 v3.4.0	Villanueva and Chen, 2019
software, algorithm	clusterProfiler v4.6.0	Wu et al., 2021
software, algorithm	org.Dm.eg.db v3.16.0	https://www.bioconductor.org/
software, algorithm	DESeq2 v1.38.2	Love et al., 2014
software, algorithm	Integrative Genomics Viewer v2.12.0	Thorvaldsdóttir et al., 2013
software, algorithm	Fiji v1.53c	Schindelin et al., 2012
software, algorithm	AnimalTFDB	Hu et al., 2019

658   Antibodies used in this study:

Antibody	Manufacturer	Product #	RRID #
Mouse anti-SXL	Developmental Studies Hybridoma	M18-s	AB_528464

	Bank		
Goat anti-MSL2	Santa Cruz	sc-32459	AB_672213
Goat anti-MOF	Santa Cruz	sc-22351	AB_670132
Rabbit anti-H4K16Ac	EMD Millipore	07-329	AB_310525
Alexa Fluor 488/594	Jackson Immuno Research	N/A	N/A
Biotin-Conjugated Mouse Monoclonal Anti-Digoxigenin antibody	Jackson Immuno Research	200-062-156	AB_2339017
Streptavidin-HRP	Invitrogen	S991	N/A
Cyanine 3 Tyramide	Akoya Biosciences	FP1046	N/A

Drosophila stocks and genetic crosses

The *Drosophila* strains mentioned in this study were all maintained and crossed in our laboratory.

The crossing methods have been described previously (Zhang et al., 2021a). Trisomy

chromosome 2 left arm (2L) female and male third instar larvae were obtained from the cross of

y; C(2L)dp; F(2R) bw females and Canton S males. The metafemale larvae were obtained from the

cross of *C(1)DX, ywf/winscy* females and Canton S males. Aneuploidies overexpressing MSL2 were

generated from the crosses of *Drosophila* with compound chromosomes and MSL2 homozygotes.

The MSL2 transgene strain was constructed and validated in a previous study (Sun et al., 2013a).

All *Drosophila* strains were cultured on cornmeal dextrose medium at 25°C. Genes and

chromosomal balancers are described in Flybase (https://flybase.org/).

RNA m^{6}A methylation quantification

Total RNA was extracted from *Drosophila* larvae using TRIzol Reagent (Invitrogen) to detect the

abundance of m^{6}A. The relative quantification of m^{6}A methylation was performed using EpiQuik

m^{6}A RNA Methylation Quantification Kit (Colorimetric) (Epigentek, NY, USA, Cat # P-9005).

Specifically, 80 μl of Binding Solution was first added to each well of the plate, and 200 ng of RNA

samples, 2 μl of Negative Control, or 2 μl of diluted Positive Control were added to the designed

wells. Subsequently, the plate was incubated at 37°C for 90 min. After washing with Wash Buffer,

50 μl of Capture Antibody, 50 μl of Detection Antibody, and 50 μl of Enhancer Solution were

added to each well in order, and wells were emptied before adding a new solution each time.

Finally, 100 µl of Developer Solution and Stop Solution were added to each well away from light,

and the absorbance was read on a microplate reader at a wavelength of 450 nm.

m^{6}A methylated RNA immunoprecipitation sequencing (MeRIP-Seq)

The MeRIP-Seq service was provided by Cloudseq Biotech Inc. (Shanghai, China), and this

technology was developed on the basis of published experimental methods (Meyer et al., 2012).

In brief, *Drosophila* larvae of five genotypes, each with two biological replicates, were collected

for sequencing. Ribosomal RNAs were removed from total RNA using Ribo-Zero rRNA Removal

Kits (Illumina, USA). Immunoprecipitation of m^{6}A RNA was performed using GenSeq® m^{6}A RNA IP

kit (GenSeq, Shanghai, China). The NEBNext® Ultra II Directional RNA Library Prep kit (New

England Biolabs, USA) was used for RNA sequencing library construction. High-throughput

sequencing was performed using Illumina NovaSeq 6000 sequencers with the paired-end 150 bp

protocol.

Analysis of MeRIP-Seq data

The raw sequencing data was first filtered by Trim Galore (version 0.6.10)

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) to remove adapters and

low-quality reads. The quality of the data was then assessed by FastQC (version 0.12.1)

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Subsequently, clean reads were

aligned to the *Drosophila* reference genome (Drosophila_melanogaster.BDGP6.32.dna.toplevel.fa,

downloaded from the Ensembl database) using HISAT2 (version 2.2.1) (Kim et al., 2019). Next,

The R package exomePeak2 (version 1.10.0) (Meng et al., 2013) was used for m^{6}A peak calling

(the screening criteria were log2FC ≥ 1, RPM.IP ≥ 0.5, and score ≥ 5). Motif analysis was

performed using HOMER (version 4.11) (Heinz et al., 2010). Differentially methylated peaks were

analyzed by the R package DiffBind (version 3.8.4) (Stark and Brown, 2013), which employs the

DESeq2 (Love et al., 2014) method. m^{6}A peaks with a p-value of 0.1 or less were considered as

differentially methylated peaks (DMPs). ChIPseeker (version 1.34.1) (Yu et al., 2015) was used to

annotate the peaks, and plotted some of the figures. The profiles and heatmaps of MeRIP-Seq

reads around DMPs were generated by deepTools (version 3.5.4) (Ramírez et al., 2014). The signal

distribution on the genome was visualized using the Integrative Genomics Viewer (IGV) software

(version 2.12.0) (Thorvaldsdóttir et al., 2013).

RNA sequencing data and analysis

The RNA-seq data of aneuploid *Drosophila* used in this article were generated in a previous study

(Zhang et al., 2023) and can be downloaded from the Gene Expression Omnibus (GEO) database

(GSE233534). Genome mapping of sequencing data and gene expression quantification were

carried out through HISAT2-StringTie pipeline (Pertea et al., 2015; Kim et al., 2019). Differential

alternative splicing events were identified by rMATS (version 4.2.0) program (Shen et al., 2014)

(with the parameters of -t paired --readLength 150 --cstat 0.0001 --libType fr-firststrand). The

method of generating the ratio distributions of gene expression changes was as described

previously (Zhang et al., 2022). The plots were generated using ggplot2 (version 3.4.0) (Villanueva

and Chen, 2019) in the R program (version 4.2.2). Differential expression analysis was performed

by DESeq2 (version 1.38.2) (Love et al., 2014), and the threshold was set to adjusted p-value ≤

0.05. The functional enrichment analysis was performed by ClusterProfiler (version 4.6.0) (Wu et

al., 2021) based on org.Dm.eg.db (version 3.16.0) from Bioconductor

(https://www.bioconductor.org/). The Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathway data was obtained from the network (https://www.kegg.jp/). The list of transcription

factors of *Drosophila* was downloaded from AnimalTFDB (Hu et al., 2019). Protein-protein

interaction (PPI) relationships were obtained from the STRING database (Szklarczyk et al., 2019).

Real-time quantitative PCR

Total RNA of *Drosophila* whole larvae, larval brains, or adult heads was extracted using TRIzol

Reagent (Invitrogen), and reverse transcription was done with *TransScript* one-step gDNA

Removal and cDNA Synthesis SuperMix (TransGen Biotech). The sequences of primers designed

for RT-qPCR were listed in Figure 1—figure supplement 1A. The usability of these primers was

verified by agarose gel electrophoresis (Figure 1—figure supplement 1B). The real-time PCR was

performed with *TransStart* Tip Green qPCR SuperMix (+ Dye II) (TransGen Biotech) using ABI

QuantStudio 6 Flex Real-Time PCR System. Relative quantification of gene expression was

determined using the 2$^{-\Delta\Delta Ct}$ method.

Fluorescence *in situ* hybridization (FISH) of *Drosophila* embryos

Collection of *Drosophila* embryos and fluorescence *in situ* hybridization were performed as

previously described (Jandura et al., 2017; Zhang et al., 2021a). The probe primers containing

flanking T7 promoter elements were listed in Figure 1—figure supplement 1D. After PCR

amplification and *in vitro* transcription, their products were examined by agarose gel

electrophoresis (Figure 1—figure supplement 1E). Digoxygenin (DIG)-labelled antisense RNA

probes were hybridized to transcripts of interest, and then detected using a succession of

anti-DIG antibody conjugated to biotin, streptavidin conjugated to horseradish peroxidase (HRP)

and fluorescently conjugated tyramide. This hierarchical process can greatly enhance the probe

signals and is referred to as a tyramide signal amplification (TSA) system. All images were

acquired with a Zeiss LSM880 laser confocal fluorescence microscope using ZEN software. The

same probes for relative fluorescence intensity analysis were photographed using the same

parameters. Fluorescence images were processed and analyzed using Fiji (version 1.53c)

(Schindelin et al., 2012).

Polytene chromosomes immunostaining

Salivary gland chromosomes immunostaining was performed as previously described (Zhang et

al., 2021a). Briefly speaking, the salivary glands from third-instar larvae were first fixed in 3.7%

formaldehyde for 1 min, and then dissociated with 50% acetic acid for 5 min. Polytene

chromosomes were treated with the following primary antibodies at a dilution of 1:100: anti-SXL

(Developmental Studies Hybridoma Bank, M18-s), anti-MSL2 (Santa Cruz, sc-32459), anti-MOF

(Santa Cruz, sc-22351), and anti-H4K16Ac (EMD Millipore, 07-329). Finally,

fluorescence-conjugated secondary antibodies (Alexa Fluor 488 and Alexa Fluor 594, Jackson

Immuno Research) were used for detection.

Acknowledgements

This work was supported by National Natural Science Foundation of China (Grant No. 31871243 and No. 32070566) to L.S. We thank James A.Birchler for revising the manuscript.

Additional information

Funding

Funder	Grant reference number	Author
National Natural Science Foundation of China	No. 31871243	Lin Sun
National Natural Science Foundation of China	No. 32070566	Lin Sun

Author contributions

Shuai Zhang: Methodology, Investigation, Data Curation, Formal analysis, Writing - Original Draft, Writing - Review & Editing, Visualization. Xinyu Liu, Methodology, Investigation, Validation, Writing - Original Draft, Writing - Review & Editing, Visualization. Ruixue Wang, Methodology, Investigation, Validation, Writing - Original Draft, Writing - Review & Editing, Visualization. Junhan Wang: Methodology, Investigation, Formal analysis. Ludan Zhang: Methodology, Investigation, Validation. Lin Sun: Conceptualization, Writing - Original Draft, Writing - Review & Editing, Supervision, Project administration, Funding acquisition. All authors have read and approved the final manuscript.

Additional files

Supplementary files

Supplementary File 1

Data availability

All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE253401. The previously published RNA-seq data used in this study are available at the GEO repository under the accession number GSE233534.

The following datasets were generated:

Author(s)	Year	Dataset title	Dataset URL	Database and Identifier
Zhang S, Liu X, Wang R, Wang J, Zhang L, Sun L	2024	Dynamics and regulatory roles of RNA m^{6}A methylation in unbalanced genomes	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE253401	NCBI Gene Expression Omnibus, GSE253401

The following previously published datasets were used:

Author(s)	Year	Dataset title	Dataset URL	Database and Identifier
Zhang S, Wang R, Zhu X, Zhang L, Liu X, Sun L	2024	Characteristics and expression of lncRNA in *Drosophila* aneuploidy	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE233534	NCBI Gene Expression Omnibus, GSE233534

References

Barbieri, I., Tzelepis, K., Pandolfini, L., Shi, J., Millan-Zambrano, G., Robson, S. C., Aspris, D., Migliori, V., Bannister, A. J., Han, N., De Braekeleer, E., Ponstingl, H., Hendrick, A., Vakoc, C. R., Vassiliou, G. S., & Kouzarides, T. (2017). Promoter-bound METTL3 maintains myeloid leukaemia by m(6)A-dependent translation control. *Nature*, *552*(7683), 126-131. https://doi.org/10.1038/nature24678

Ben-David, U., & Amon, A. (2020). Context is everything: aneuploidy in cancer. *Nature Reviews Genetics*, *21*(1), 44-62. https://doi.org/10.1038/s41576-019-0171-x

Birchler, J. A. (1981). The genetic basis of dosage compensation of alcohol dehydrogenase-1 in maize. *Genetics*, *97*(3-4), 625-637. https://www.ncbi.nlm.nih.gov/pubmed/17249082

Birchler, J. A. (2016). Parallel Universes for Models of X Chromosome Dosage Compensation in

Drosophila: A   Review.   *Cytogenetic   and   Genome   Research*,   *148*(1),   52-67.

https://doi.org/10.1159/000445924

Birchler, J. A., Bhadra, U., Bhadra, M. P., & Auger, D. L. (2001). Dosage-dependent gene regulation

in multicellular eukaryotes: implications for dosage compensation, aneuploid syndromes, and

quantitative   traits.   *Developmental   Biology*,   *234*(2),   275-288.

https://doi.org/10.1006/dbio.2001.0262

Birchler, J. A., & Veitia, R. A. (2007). The gene balance hypothesis: from classical genetics to

modern genomics. *Plant Cell*, *19*(2), 395-402. https://doi.org/10.1105/tpc.106.049338

Birchler, J. A., & Veitia, R. A. (2012). Gene balance hypothesis: connecting issues of dosage

sensitivity across biological disciplines. *Proceedings of the National Academy of Sciences of*

the   United   States   of   America,   *109*(37),   14746-14753.

https://doi.org/10.1073/pnas.1207726109

Birchler, J. A., & Veitia, R. A. (2021). One Hundred Years of Gene Balance: How Stoichiometric

Issues Affect Gene Expression, Genome Evolution, and Quantitative Traits. *Cytogenetic and*

Genome Research, *161*(10-11), 529-550. https://doi.org/10.1159/000519592

Chiarle, R. (2021). Solving the chromosome puzzle of aneuploidy in cancer. *Genes Dev*, *35*(15-16),

1073-1075. https://doi.org/10.1101/gad.348773.121

Conrad, T., Cavalli, F. M. G., Holz, H., Hallacli, E., Kind, J., Ilik, I., Vaquerizas, J. M., Luscombe, N. M.,

& Akhtar, A. (2012). The MOF Chromobarrel Domain Controls Genome-wide H4K16

Acetylation and Spreading of the MSL Complex. *Developmental Cell*, *22*(3), 610-624.

https://doi.org/10.1016/j.devcel.2011.12.016

Disteche, C. M. (2016). Dosage compensation of the sex chromosomes and autosomes. *Semin*

Cell Dev Biol, *56*, 9-18. https://doi.org/10.1016/j.semcdb.2016.04.013

Dominissini, D., Moshitch-Moshkovitz, S., Schwartz, S., Salmon-Divon, M., Ungar, L., Osenberg, S.,

Cesarkas, K., Jacob-Hirsch, J., Amariglio, N., Kupiec, M., Sorek, R., & Rechavi, G. (2012).

Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature*,

485(7397), 201-206. https://doi.org/10.1038/nature11112

Geula, S., Moshitch-Moshkovitz, S., Dominissini, D., Mansour, A. A., Kol, N., Salmon-Divon, M.,

Hershkovitz, V., Peer, E., Mor, N., Manor, Y. S., Ben-Haim, M. S., Eyal, E., Yunger, S., Pinto, Y.,

Jaitin, D. A., Viukov, S., Rais, Y., Krupalnik, V., Chomsky, E., . . . Hanna, J. H. (2015). Stem cells.

m6A mRNA methylation facilitates resolution of naive pluripotency toward differentiation.

Science, *347*(6225), 1002-1006. https://doi.org/10.1126/science.1261417

Gilbert, W. V., Bell, T. A., & Schaening, C. (2016). Messenger RNA modifications: Form, distribution,

and function. *Science*, *352*(6292), 1408-1412. https://doi.org/10.1126/science.aad8711

Granadino, B., Campuzano, S., & Sanchez, L. (1990). The *Drosophila melanogaster fl(2)d* gene is

needed for the female-specific splicing of *Sex-lethal* RNA. *EMBO Journal*, *9*(8), 2597-2602.

https://doi.org/DOI 10.1002/j.1460-2075.1990.tb07441.x

Gurudatta, B. V., Ramos, E., & Corces, V. G. (2012). The BEAF insulator regulates genes involved in

cell polarity and neoplastic growth. *Developmental Biology*, *369*(1), 124-132.

https://doi.org/10.1016/j.ydbio.2012.06.013

Haussmann, I. U., Bodi, Z., Sanchez-Moran, E., Mongan, N. P., Archer, N., Fray, R. G., & Soller, M.

(2016). m(6)A potentiates *Sxl* alternative pre-mRNA splicing for robust *Drosophila* sex

determination. *Nature*, *540*(7632), 301-304. https://doi.org/10.1038/nature20577

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H.,

& Glass, C. K. (2010). Simple Combinations of Lineage-Determining Transcription Factors

Prime *cis*-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell*,

38(4), 576-589. https://doi.org/10.1016/j.molcel.2010.05.004

Hongay, C. F., & Orr-Weaver, T. L. (2011). *Drosophila* Inducer of MEiosis 4 (*IME4*) is required for

Notch signaling during oogenesis. *Proceedings of the National Academy of Sciences of the*

United States of America, *108*(36), 14855-14860. https://doi.org/10.1073/pnas.1111577108

Horiuchi, K., Umetani, M., Minami, T., Okayama, H., Takada, S., Yamamoto, M., Aburatani, H., Reid,

P. C., Housman, D. E., Hamakubo, T., & Kodama, T. (2006). Wilms' tumor 1-associating protein

regulates G2/M transition through stabilization of cyclin A2 mRNA. *Proceedings of the*

National Academy of Sciences of the United States of America, *103*(46), 17278-17283.

https://doi.org/10.1073/pnas.0608357103

Hou, J., Shi, X., Chen, C., Islam, M. S., Johnson, A. F., Kanno, T., Huettel, B., Yen, M. R., Hsu, F. M., Ji,

T., Chen, P. Y., Matzke, M., Matzke, A. J. M., Cheng, J., & Birchler, J. A. (2018). Global impacts

of chromosomal imbalance on gene expression in *Arabidopsis* and other taxa. *Proceedings of*

the National Academy of Sciences of the United States of America, *115*(48), E11321-E11330.

https://doi.org/10.1073/pnas.1807796115

Hu, H., Miao, Y. R., Jia, L. H., Yu, Q. Y., Zhang, Q., & Guo, A. Y. (2019). AnimalTFDB 3.0: a

comprehensive resource for annotation and prediction of animal transcription factors. *Nucleic*

Acids Research, *47*(D1), D33-D38. https://doi.org/10.1093/nar/gky822

Huang, A. C., Olson, S. B., & Maslen, C. L. (2021). A Review of Recent Developments in Turner

Syndrome   Research.   *J   Cardiovasc   Dev   Dis*,   *8*(11),   475-491.

https://doi.org/10.3390/jcdd8110138

Huang, H., Weng, H., Zhou, K., Wu, T., Zhao, B. S., Sun, M., Chen, Z., Deng, X., Xiao, G., Auer, F.,

Klemm, L., Wu, H., Zuo, Z., Qin, X., Dong, Y., Zhou, Y., Qin, H., Tao, S., Du, J., . . . Chen, J. (2019).

Histone H3 trimethylation at lysine 36 guides m(6)A RNA modification co-transcriptionally.

Nature, *567*(7748), 414-419. https://doi.org/10.1038/s41586-019-1016-7

Jaenisch, R., & Bird, A. (2003). Epigenetic regulation of gene expression: how the genome

integrates intrinsic and environmental signals. *Nature Genetics*, *33 Suppl*, 245-254.

https://doi.org/10.1038/ng1089

Jandura, A., Hu, J., Wilk, R., & Krause, H. M. (2017). High Resolution Fluorescent *In Situ*

Hybridization in Drosophila Embryos and Tissues Using Tyramide Signal Amplification. *J Vis*

Exp, *128*(128), 56281. https://doi.org/10.3791/56281

Jordan, W., Rieder, L. E., & Larschan, E. (2019). Diverse Genome Topologies Characterize Dosage

Compensation   across   Species.   *Trends   in   Genetics*,   *35*(4),   308-315.

https://doi.org/10.1016/j.tig.2019.02.001

Kan, L., Grozhik, A. V., Vedanayagam, J., Patil, D. P., Pang, N., Lim, K. S., Huang, Y. C., Joseph, B., Lin,

C. J., Despic, V., Guo, J., Yan, D., Kondo, S., Deng, W. M., Dedon, P. C., Jaffrey, S. R., & Lai, E. C.

(2017). The m(6)A pathway facilitates sex determination in *Drosophila*. *Nat Commun*, *8*,

15737. https://doi.org/10.1038/ncomms15737

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8),

907-915. https://doi.org/10.1038/s41587-019-0201-4

Kind, J., Vaquerizas, J. M., Gebhardt, P., Gentzel, M., Luscombe, N. M., Bertone, P., & Akhtar, A.

(2008). Genome-wide analysis reveals MOF as a key regulator of dosage compensation and

gene   expression   in   *Drosophila*.   *Cell*,   *133*(5),   813-828.

https://doi.org/10.1016/j.cell.2008.04.036

Lecuyer, E., Yoshida, H., Parthasarathy, N., Alm, C., Babak, T., Cerovina, T., Hughes, T. R., Tomancak, P., & Krause, H. M. (2007). Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell*, *131*(1), 174-187. https://doi.org/10.1016/j.cell.2007.08.003

Lee, M., Kim, B., & Kim, V. N. (2014). Emerging roles of RNA modification: m(6)A and U-tail. *Cell*, *158*(5), 980-987. https://doi.org/10.1016/j.cell.2014.08.005

Lence, T., Akhtar, J., Bayer, M., Schmid, K., Spindler, L., Ho, C. H., Kreim, N., Andrade-Navarro, M. A., Poeck, B., Helm, M., & Roignant, J. Y. (2016). m(6)A modulates neuronal functions and sex determination in *Drosophila*. *Nature*, *540*(7632), 242-247. https://doi.org/10.1038/nature20568

Lence, T., Soller, M., & Roignant, J. Y. (2017). A fly view on the roles and mechanisms of the m(6)A mRNA modification and its players. *RNA Biology*, *14*(9), 1232-1240. https://doi.org/10.1080/15476286.2017.1307484

Letourneau, A., Santoni, F. A., Bonilla, X., Sailani, M. R., Gonzalez, D., Kind, J., Chevalier, C., Thurman, R., Sandstrom, R. S., Hibaoui, Y., Garieri, M., Popadin, K., Falconnet, E., Gagnebin, M., Gehrig, C., Vannier, A., Guipponi, M., Farinelli, L., Robyr, D., . . . Antonarakis, S. E. (2014). Domains of genome-wide gene expression dysregulation in Down's syndrome. *Nature*, *508*(7496), 345-350. https://doi.org/10.1038/nature13200

Li, Y., Xia, L., Tan, K., Ye, X., Zuo, Z., Li, M., Xiao, R., Wang, Z., Liu, X., Deng, M., Cui, J., Yang, M., Luo, Q., Liu, S., Cao, X., Zhu, H., Liu, T., Hu, J., Shi, J., . . . Xia, L. (2020). *N*6-Methyladenosine co-transcriptionally directs the demethylation of histone H3K9me2. *Nature Genetics*. https://doi.org/10.1038/s41588-020-0677-3

Liu, J., Gao, M., He, J., Wu, K., Lin, S., Jin, L., Chen, Y., Liu, H., Shi, J., Wang, X., Chang, L., Lin, Y., Zhao, Y. L., Zhang, X., Zhang, M., Luo, G. Z., Wu, G., Pei, D., Wang, J., . . . Chen, J. (2021). The RNA m(6)A reader YTHDC1 silences retrotransposons and guards ES cell identity. *Nature*, *591*(7849), 322-326. https://doi.org/10.1038/s41586-021-03313-9

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Lucchesi, J. C. (2018). Transcriptional modulation of entire chromosomes: dosage compensation.

Journal of Genetics, *97*(2), 357-364. https://doi.org/10.1007/s12041-018-0919-7

Luo, G. Z., MacQueen, A., Zheng, G., Duan, H., Dore, L. C., Lu, Z., Liu, J., Chen, K., Jia, G., Bergelson,

J., & He, C. (2014). Unique features of the m6A methylome in *Arabidopsis thaliana*. *Nat*

Commun, *5*, 5630. https://doi.org/10.1038/ncomms6630

Ma, S., Chen, C., Ji, X., Liu, J., Zhou, Q., Wang, G., Yuan, W., Kan, Q., & Sun, Z. (2019). The interplay

between m6A RNA methylation and noncoding RNA in cancer. *Journal of Hematology &*

Oncology, *12*(1), 121. https://doi.org/10.1186/s13045-019-0805-7

Meng, J., Cui, X. D., Rao, M. K., Chen, Y. D., & Huang, Y. F. (2013). Exome-based analysis for RNA

epigenome   sequencing   data.   *Bioinformatics*,   *29*(12),   1565-1567.

https://doi.org/10.1093/bioinformatics/btt171

Meyer, K. D., & Jaffrey, S. R. (2014). The dynamic epitranscriptome: *N*6-methyladenosine and

gene expression control. *Nature Reviews: Molecular Cell Biology*, *15*(5), 313-326.

https://doi.org/10.1038/nrm3785

Meyer, K. D., Patil, D. P., Zhou, J., Zinoviev, A., Skabkin, M. A., Elemento, O., Pestova, T. V., Qian, S.

B., & Jaffrey, S. R. (2015). 5' UTR m6A Promotes Cap-Independent Translation. *Cell*, *163*(4),

999-1010. https://doi.org/10.1016/j.cell.2015.10.012

Meyer, K. D., Saletore, Y., Zumbo, P., Elemento, O., Mason, C. E., & Jaffrey, S. R. (2012).

Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop

codons. *Cell*, *149*(7), 1635-1646. https://doi.org/10.1016/j.cell.2012.05.003

Orr, B., Godek, K. M., & Compton, D. (2015). Aneuploidy. *Current Biology*, *25*(13), R538-R542.

https://doi.org/10.1016/j.cub.2015.05.010

Park, S. W., Oh, H., Lin, Y. R., & Park, Y. (2010). MSL *cis*-spreading from *roX* gene up-regulates the

neighboring   genes.   *Biochem   Biophys   Res   Commun*,   *399*(2),   227-231.

https://doi.org/10.1016/j.bbrc.2010.07.059

Patil, D. P., Chen, C. K., Pickering, B. F., Chow, A., Jackson, C., Guttman, M., & Jaffrey, S. R. (2016).

m(6)A RNA methylation promotes *XIST*-mediated transcriptional repression. *Nature*,

537(7620), 369-373. https://doi.org/10.1038/nature19342

Penn, J. K., Graham, P., Deshpande, G., Calhoun, G., Chaouki, A. S., Salz, H. K., & Schedl, P. (2008).

Functioning of the *Drosophila* Wilms'-tumor-1-associated protein homolog, *Fl(2)d*, in

Sex-lethal-dependent alternative splicing. *Genetics*, *178*(2), 737-748. https://doi.org/10.1534/genetics.107.081679

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T. C., Mendell, J. T., & Salzberg, S. L. (2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology*, *33*(3), 290-295. https://doi.org/10.1038/nbt.3122

Pinello, N., Sun, S., & Wong, J. J. (2018). Aberrant expression of enzymes regulating m(6)A mRNA methylation: implication in cancer. *Cancer Biol Med*, *15*(4), 323-334. https://doi.org/10.20892/j.issn.2095-3941.2018.0365

Prestel, M., Feller, C., & Becker, P. B. (2010). Dosage compensation and the global re-balancing of aneuploid genomes. *Genome Biology*, *11*(8), 216. https://doi.org/10.1186/gb-2010-11-8-216

Rabinow, L., & Birchler, J. A. (1989). A dosage-sensitive modifier of retrotransposon-induced alleles of the *Drosophila white* locus. *EMBO Journal*, *8*(3), 879-889. https://www.ncbi.nlm.nih.gov/pubmed/2542025

Raffel, G. D., Mercher, T., Shigematsu, H., Williams, I. R., Cullen, D. E., Akashi, K., Bernard, O. A., & Gilliland, D. G. (2007). *Ott1*(*Rbm15*) has pleiotropic roles in hematopoietic development. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(14), 6001-6006. https://doi.org/10.1073/pnas.0609041104

Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A., & Manke, T. (2014). deepTools: a flexible platform for exploring deep-sequencing data. *Nucleic Acids Research*, *42*(W1), W187-W191. https://doi.org/10.1093/nar/gku365

Raznahan, A., Parikshak, N. N., Chandran, V., Blumenthal, J. D., Clasen, L. S., Alexander-Bloch, A. F., Zinn, A. R., Wangsa, D., Wise, J., Murphy, D. G. M., Bolton, P. F., Ried, T., Ross, J., Giedd, J. N., & Geschwind, D. H. (2018). Sex-chromosome dosage effects on gene expression in humans. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(28), 7398-7403. https://doi.org/10.1073/pnas.1802889115

Roseman, R. R., Pirrotta, V., & Geyer, P. K. (1993). The *su(Hw)* protein insulates expression of the *Drosophila melanogaster white* gene from chromosomal position-effects. *EMBO Journal*, *12*(2), 435-442. https://doi.org/DOI 10.1002/j.1460-2075.1993.tb05675.x

Roundtree, I. A., Evans, M. E., Pan, T., & He, C. (2017a). Dynamic RNA Modifications in Gene Expression Regulation. *Cell*, *169*(7), 1187-1200. https://doi.org/10.1016/j.cell.2017.05.045

Roundtree, I. A., Luo, G. Z., Zhang, Z. J., Wang, X., Zhou, T., Cui, Y. Q., Sha, J. H., Huang, X. X.,

Guerrero, L., Xie, P., He, E., Shen, B., & He, C. (2017b). YTHDC1 mediates nuclear export of

*N*6-methyladenosine methylated mRNAs. *Elife*, *6*, e31311. https://doi.org/ARTN e31311

Rücklé, C., Körtel, N., Basilicata, M. F., Busch, A., Zhou, Y., Hoch-Kraft, P., Tretow, K., Kielisch, F.,

Bertin, M., Pradhan, M., Musheev, M., Schweiger, S., Niehrs, C., Rausch, O., Zarnack, K.,

Valsecchi, C. I. K., & König, J. (2023). RNA stability controlled by m(6)A methylation

contributes to X-to-autosome dosage compensation in mammals. *Nature Structural &*

Molecular Biology, *30*(8). https://doi.org/10.1038/s41594-023-00997-7

San Roman, A. K., Godfrey, A. K., Skaletsky, H., Bellott, D. W., Groff, A. F., Harris, H. L., Blanton, L.

V., Hughes, J. F., Brown, L., Phou, S., Buscetta, A., Kruszka, P., Banks, N., Dutra, A., Pak, E.,

Lasutschinkow, P. C., Keen, C., Davis, S. M., Tartaglia, N. R., . . . Page, D. C. (2023). The human

inactive X chromosome modulates expression of the active X chromosome. *Cell Genom*, *3*(2),

100259. https://doi.org/10.1016/j.xgen.2023.100259

Sanchez-Pavon, E., Mendoza, H., & Garcia-Ferreyra, J. (2021). Trisomy 21 and Assisted

Reproductive Technologies: A review. *JBRA Assist Reprod*, *26*(1), 129-141.

https://doi.org/10.5935/1518-0557.20210047

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,

Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K.,

Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis.

Nature Methods, *9*(7), 676-682. https://doi.org/10.1038/Nmeth.2019

Shafik, A. M., Zhang, F., Guo, Z., Dai, Q., Pajdzik, K., Li, Y., Kang, Y., Yao, B., Wu, H., He, C., Allen, E.

G., Duan, R., & Jin, P. (2021). *N*6-methyladenosine dynamics in neurodevelopment and aging,

and its potential role in Alzheimer's disease. *Genome Biology*, *22*(1), 17.

https://doi.org/10.1186/s13059-020-02249-z

Shen, S. H., Park, J. W., Lu, Z. X., Lin, L., Henry, M. D., Wu, Y. N., Zhou, Q., & Xing, Y. (2014). rMATS:

Robust and flexible detection of differential alternative splicing from replicate RNA-Seq data.

Proceedings of the National Academy of Sciences of the United States of America, *111*(51),

E5593-E5601. https://doi.org/10.1073/pnas.1419161111

Shi, X., Yang, H., Chen, C., Hou, J., Hanson, K. M., Albert, P. S., Ji, T., Cheng, J., & Birchler, J. A.

(2021). Genomic imbalance determines positive and negative modulation of gene expression

in diploid maize. *Plant Cell*, *33*(4), 917-939. https://doi.org/10.1093/plcell/koab030

Shi, X., Yang, H., Chen, C., Hou, J., Ji, T., Cheng, J., & Birchler, J. A. (2022). Dosage-sensitive miRNAs

trigger modulation of gene expression during genomic imbalance in maize. *Nat Commun*,

13(1), 3014. https://doi.org/10.1038/s41467-022-30704-x

Stark, R., Brown, G. (2013). DiffBind: differential binding analysis of ChIP-seq peak data.

Bioconductor package. http://www.bioconductor.org. DOI: 10.18129/B9.bioc.DiffBind.

Sun, L., Fernandez, H. R., Donohue, R. C., Li, J., Cheng, J., & Birchler, J. A. (2013a). Male-specific

lethal complex in *Drosophila* counteracts histone acetylation and does not mediate dosage

compensation. *Proceedings of the National Academy of Sciences of the United States of*

America, *110*(9), E808-E817. https://doi.org/10.1073/pnas.1222542110

Sun, L., Johnson, A. F., Donohue, R. C., Li, J., Cheng, J., & Birchler, J. A. (2013b). Dosage

compensation and inverse effects in triple X metafemales of *Drosophila*. *Proceedings of the*

National Academy of Sciences of the United States of America, *110*(18), 7383-7388.

https://doi.org/10.1073/pnas.1305638110

Sun, L., Johnson, A. F., Li, J., Lambdin, A. S., Cheng, J., & Birchler, J. A. (2013c). Differential effect of

aneuploidy on the X chromosome and genes with sex-biased expression in *Drosophila*.

Proceedings of the National Academy of Sciences of the United States of America, *110*(41),

16514-16519. https://doi.org/10.1073/pnas.1316041110

Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,

Doncheva, N. T., Morris, J. H., Bork, P., Jensen, L. J., & Mering, C. (2019). STRING v11:

protein-protein association networks with increased coverage, supporting functional

discovery in genome-wide experimental datasets. *Nucleic Acids Research*, *47*(D1), D607-D613.

https://doi.org/10.1093/nar/gky1131

Thorvaldsdóttir, H., Robinson, J. T., & Mesirov, J. P. (2013). Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. *Briefings in Bioinformatics*,

14(2), 178-192. https://doi.org/10.1093/bib/bbs017

Villanueva, R. A. M., & Chen, Z. J. (2019). ggplot2: Elegant Graphics for Data Analysis, 2nd edition.

Measurement-Interdisciplinary Research and Perspectives, *17*(3), 160-167.

https://doi.org/10.1080/15366367.2019.1565254

Wang, X., Lu, Z. K., Gomez, A., Hon, G. C., Yue, Y. N., Han, D. L., Fu, Y., Parisien, M., Dai, Q., Jia, G.

F., Ren, B., Pan, T., & He, C. (2014). *N6*-methyladenosine-dependent regulation of messenger

RNA stability. *Nature*, *505*(7481), 117-120. https://doi.org/10.1038/nature12730

Wang, Y., Li, Y., Toth, J. I., Petroski, M. D., Zhang, Z., & Zhao, J. C. (2014). *N6*-methyladenosine

modification destabilizes developmental regulators in embryonic stem cells. *Nature Cell*

Biology, *16*(2), 191-198. https://doi.org/10.1038/ncb2902

Wang, Y., Li, Y., Yue, M. H., Wang, J., Kumar, S., Wechsler-Reya, R. J., Zhang, Z. L., Ogawa, Y., Kellis,

M., Duester, G., & Zhao, J. C. (2018). N(6)-methyladenosine RNA modification regulates

embryonic neural stem cell self-renewal through histone modifications. *Nature Neuroscience*,

21(8), 1139-1139. https://doi.org/10.1038/s41593-018-0169-2

Williams, B. R., Prabhu, V. R., Hunter, K. E., Glazier, C. M., Whittaker, C. A., Housman, D. E., &

Amon, A. (2008). Aneuploidy affects proliferation and spontaneous immortalization in

mammalian cells. *Science*, *322*(5902), 703-709. https://doi.org/10.1126/science.1160058

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S.,

Bo, X., & Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics

data. *Innovation (N Y)*, *2*(3), 100141. https://doi.org/10.1016/j.xinn.2021.100141

Xie, W., & Birchler, J. A. (2012). Identification of Inverse Regulator-a (*Inr-a*) as Synonymous with

Pre-mRNA Cleavage Complex II Protein (*Pcf11*) in *Drosophila*. *G3 (Bethesda)*, *2*(6), 701-706.

https://doi.org/10.1534/g3.112.002071

Yan, D., & Perrimon, N. (2015). *spenito* is required for sex determination in *Drosophila*

melanogaster. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(37), 11606-11611. https://doi.org/10.1073/pnas.1515891112

Yang, H., Shi, X., Chen, C., Hou, J., Ji, T., Cheng, J., & Birchler, J. A. (2021). Predominantly inverse

modulation of gene expression in genomically unbalanced disomic haploid maize. *Plant Cell*,

33(4), 901-916. https://doi.org/10.1093/plcell/koab029

Yang, Y., Hsu, P. J., Chen, Y. S., & Yang, Y. G. (2018). Dynamic transcriptomic m(6)A decoration:

writers, erasers, readers and functions in RNA metabolism. *Cell Research*, *28*(6), 616-624.

https://doi.org/10.1038/s41422-018-0040-8

Yu, G. C., Wang, L. G., & He, Q. Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak

annotation, comparison and visualization. *Bioinformatics*, *31*(14), 2382-2383.

https://doi.org/10.1093/bioinformatics/btv145

Zaccara, S., Ries, R. J., & Jaffrey, S. R. (2019). Reading, writing and erasing mRNA methylation.

Nature Reviews: Molecular Cell Biology, *20*(10), 608-624.

https://doi.org/10.1038/s41580-019-0168-5

Zhang, L. D., Zhang, S., Wang, R. X., & Sun, L. (2022). Genome-Wide Identification of Long

Noncoding RNA and Their Potential Interactors in *ISWI* Mutants. *International Journal of*

Molecular Sciences, *23*(11), 6247. https://doi.org/10.3390/ijms23116247

Zhang, S., Qi, H., Huang, C., Yuan, L., Zhang, L., Wang, R., Tian, Y., & Sun, L. (2021a). Interaction of

Male Specific Lethal complex and genomic imbalance on global gene expression in *Drosophila*.

Scientific Reports, *11*(1), 19679. https://doi.org/10.1038/s41598-021-99268-y

Zhang, S., Wang, R., Huang, C., Zhang, L., & Sun, L. (2021b). Modulation of Global Gene

Expression by Aneuploidy and CNV of Dosage Sensitive Regulatory Genes. *Genes (Basel)*,

12(10), 1606. https://doi.org/10.3390/genes12101606

Zhang, S., Wang, R., Zhu, X., Zhang, L., Liu, X., & Sun, L. (2023). Characteristics and expression of

lncRNA and transposable elements in *Drosophila* aneuploidy. *iScience*, *26*(12), 108494.

https://doi.org/10.1016/j.isci.2023.108494

Zhao, B. S., Wang, X., Beadell, A. V., Lu, Z., Shi, H., Kuuspalu, A., Ho, R. K., & He, C. (2017).

m(6)A-dependent maternal mRNA clearance facilitates zebrafish maternal-to-zygotic

transition. *Nature*, *542*(7642), 475-478. https://doi.org/10.1038/nature21355

Zhong, S. L., Li, H. Y., Bodi, Z., Button, J., Vespa, L., Herzog, M., & Fray, R. G. (2008). MTA is an

messenger RNA adenosine methylase and interacts with a homolog of a sex-specific splicing

factor. *Plant Cell*, *20*(5), 1278-1288. https://doi.org/10.1105/tpc.108.058883

Zhu, C., Zhang, S., Zhou, C., Tian, C., Shi, B., Xu, K., Huang, L., Sun, Y., Lin, Y., Lai, Z., & Guo, Y.

(2023). RNA Methylome Reveals the m(6)A-mediated Regulation of Flavor Metabolites in Tea

Leaves under Solar-withering. *Genomics Proteomics Bioinformatics*, *21*(4), 769-787.

https://doi.org/10.1016/j.gpb.2023.02.003

1105

1106

1107 **Figures**

1108

1109

Figure 1. The responses of m^{6}A methyltransferases and reader protein under the condition of genomic imbalance. (**A**) Schematic diagram of m^{6}A components in *Drosophila*. (**B**) RT-qPCR analysis of mRNA levels of m^{6}A methyltransferases and reader protein in third instar larvae of wildtype and trisomy *Drosophila*. CF, wildtype female control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale; 2L, chromosome 2 left arm. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001. (**C**) Subembryonic distribution patterns of the transcripts of m^{6}A components in wildtype and trisomy 2L *Drosophila*. The names of the genes were shown in the left of the pictures; the genotypes and stages were shown above. Red, probes; green, DAPI. Scale bar, 100 μm. (**D**) Subcellular localization of probe signals. Probe name was written in the corner of each picture. Red, probe; green, DAPI. Arrowheads indicate the foci of probe signals. Scale bars, 10 μm. (**E,F**) The expression levels of m^{6}A component genes in stage 6-11 (**E**) and stage 14-17 (**F**) represented by relative fluorescence intensity of probes compared with DAPI signals. The expression of wildtype embryos was set as one. Sample size = 10. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 2. Overview of RNA m⁶A methylation in aneuploid *Drosophila*. (**A**) Global m⁶A abundance in third instar larvae of wildtype and aneuploid *Drosophila*. Data represent the mean of two independent experiments, each containing three or four biological replicates. Error bar indicates the standard error of the means (SEM). Student's *t* test *p < 0.05, **p < 0.01. (**B**) The number of m⁶A peaks identified by MeRIP-Seq. (**C**) The number of m⁶A-modified genes obtained by annotating the peaks. (**D**) Expression levels of m⁶A modification sites in IP samples expressed as log10-transformed reads per million (RPM). The number on the boxplot indicates the median RPM of each sample. Mann-Whitney U test *p < 0.05, **p < 0.01, ***p < 0.001. (**E**) Percentages of peaks localized on different gene features. (**F**) The most enriched motifs obtained by *de novo*

motif analysis of the m⁶A peaks. (**G**) Venn diagram showing the intersection of m⁶A-modified

genes in each sample. (**H**) Gene biotypes of m⁶A-modified genes. (**I**) Venn diagram showing the

intersection of transcription factors and m⁶A-modified genes with more than five m⁶A peaks in all

samples. Seventy genes with more than five m⁶A peaks were listed on the right, with

transcription factors in red and others in blue. P-value indicates one-tailed Fisher's exact test. CF,

wildtype female control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male;

XXX, metafemale; 5'UTR, 5' untranslated region; 3'UTR, 3' untranslated region; MeRIP-Seq, m⁶A

methylated RNA immunoprecipitation sequencing.

1143

1144

1145   **Figure 3.** Differential m⁶A methylome analysis of aneuploid *Drosophila*. (**A**-**C**) Heatmaps of

differentially methylated peaks (DMPs) in trisomy 2L females (**A**), trisomy 2L males (**B**) and

metafemales (**C**), and their corresponding control groups. The threshold of significance was

p-value ≤ 0.1. (**D**) The number of DMPs. The threshold of significance was set to p-value ≤ 0.1.

The numbers above the horizontal dashed lines indicate peaks with upregulated methylation

levels, and the numbers below indicate peaks with downregulated methylation levels. (**E**) The

number of DMP-associated genes. (**F**-**H**) Profiles and heatmaps illustrating the density of

m^6A-modified reads at the DMP positions in trisomy 2L females (**F**), trisomy 2L males (**G**),

metafemales (**H**), and their corresponding controls. The DMPs were divided into five groups

according to the chromosomes they located. (**I**-**K**) Scatter plots showing the concentration of

reads at methylation sites on different chromosomes in trisomy 2L females (**I**), trisomy 2L males

(**J**), and metafemales (**K**). Red points indicate significantly up-regulated m^6A peaks, blue points

indicate significantly down-regulated m^6A peaks, and gray points indicate m^6A peaks without

significant changes. The percentages of DMPs on *cis* and *trans* chromosomes were indicated in

the corners of the plots. CF, wildtype female control; CM, wildtype male control; 2LF, trisomy 2L

female; 2LM, trisomy 2L male; XXX, metafemale.

Figure 4. Differentially methylated peak (DMP) associated genes and their functions. (A) Venn diagram showing the number of common DMP-associated genes in three types of aneuploidies compared with wildtypes. (B) Venn diagram showing the intersection of transcription factors and the common DMP-associated genes in all comparisons. Transcription factors with DMPs were listed on the right. P-value indicates one-tailed Fisher's exact test. (C) Network showing the functions related to expression regulation and dosage compensation enriched by DMP-associated genes. The color of the nodes indicates the comparison, and the size of the function nodes represents the number of DMP-associated genes connected with them. (D) Genome browser example of *fl(2)d* for indicated MeRIP-seq data. Steelblue color represents input reads, while other colors represent IP reads. Signals were displayed as the mean CPM of two biological replicates. The gene architectures were shown at the bottom. The magenta rectangles at above represent differentially methylated peak. CF, wildtype female control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale.

Figure 5. Relationships between RNA m^{6}A methylation and gene expression in aneuploidy. (**A**-**C**) Venn diagrams showing the intersections of differentially expressed genes (DEGs) and differentially methylated peak (DMP) associated genes in trisomy 2L females (**A**), trisomy 2L males (**B**), and metafemales (**C**) compared with their corresponding controls. P-values indicate one-tailed Fisher's exact tests. (**D**) The common differentially expressed and differentially methylated genes in all groups. (**E**) Functional enrichment analysis of simultaneously differentially expressed and differentially methylated genes. Top 10 enriched GO terms (Biological Process) with p-value < 0.1 in each comparison were shown. (**F**-**H**) Sankey diagrams showing the

relationships between genes with different m^{6}A-modified states and genes with canonical dosage

effect (DE), dosage compensation (DC), and inverse dosage effect (IDE) in trisomy 2L females (**F**),

trisomy 2L males (**G**), and metafemales (**H**). Enrichment analysis was performed on each two

groups of genes, and deep color lines indicate significant connection relationships (Fisher's exact

test p-value < 0.05). (**I**-**K**) Gene feature distributions for m^{6}A peaks on genes with canonical DE,

DC, and IDE in trisomy 2L females (**I**), trisomy 2L males (**J**), metafemales (**K**), and their

corresponding controls. CF, wildtype female control; CM, wildtype male control; 2LF, trisomy 2L

female; 2LM, trisomy 2L male; XXX, metafemale; DEG, differentially expressed gene; DMP,

differentially methylated peak; DE, dosage effect; DC, dosage compensation; IDE, inverse dosage

effect. Canonical DE refers to ratio > 1.25, DC stands for 0.8 < ratio < 1.25, IDE stands for 0.5 <

ratio < 0.8, and unchanged refers to 0.8 < ratio < 1.25.

1197

Figure 6. Combined analysis of differential alternative splicing and differential methylation. (**A**-**C**) Venn diagrams showing the intersections of differential alternative splicing (DAS) genes and differentially methylated peak (DMP) associated genes in trisomy 2L females (**A**), trisomy 2L males (**B**), and metafemales (**C**) compared with their corresponding controls. P-values indicate one-tailed Fisher's exact tests. (**D**) The common differentially alternatively spliced and differentially methylated genes in all groups. (**E**) Functional enrichment analysis of simultaneously differentially alternatively spliced and differentially methylated genes. Top 10 enriched GO terms (Biological Process) with p-value < 0.05 in each comparison were shown. (**F**) KEGG pathway enrichment analysis of simultaneously differentially alternatively spliced and differentially methylated genes. Top 10 enriched pathways with p-value < 0.1 in each comparison were shown. (**G**) Genome browser example of *su(Hw)* for indicated MeRIP-Seq data. Steelblue color represents

input reads, while other colors represent IP reads. Signals were displayed as the mean CPM of two biological replicates. The gene architecture was shown at the bottom (only two representative transcript isoforms were shown). The magenta rectangle at above represents DMP. The magenta arrowhead indicates the position of differential alternative splicing. (**H**) Sashimi plot depicting RNA sequencing reads and exon junction reads at the position where the differential splicing events occurs on *su(Hw)*. The gene model was shown below. One of the biological replicates was chosen for representation. CF, wildtype female control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale; DAS, differential alternative splicing; DMP, differentially methylated peak; CPM, Counts per million; RPKM, Reads per kilobase per million mapped reads.

Figure 7. Interactions between m^{6}A and *Drosophila* Male Specific Lethal (MSL) complex. (**A**)

Genome browser example of *Sxl* for indicated MeRIP-Seq data. Steelblue color represents input

reads, while other colors represent IP reads. Signals were displayed as the mean CPM of two

biological replicates. The gene architecture was shown at the bottom (only four representative

transcript isoforms were shown). The rectangles at above represent m^{6}A peaks, where red

indicates up-regulated DMPs, green indicates down-regulated DMP, and gray indicates no

significant changes. The magenta arrowheads indicate the positions of alternative splicing. (**B**)

Sashimi plot depicting RNA sequencing reads and exon junction reads at the position where the

differential splicing events occurs on *Sxl*. The gene model was shown below, with the third exon

indicated in red. One of the biological replicates was chosen for representation. CF, wildtype

female control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX,

metafemale. (**C,D**) RT-qPCR analysis of mRNA levels of m^{6}A components in the heads of MSL2

transgenic female (**C**) and male (**D**) *Drosophila* adults. (**E,F**) Abundance of m^{6}A modification in the

heads of MSL2 transgenic females (**E**) and males (**F**). Student's *t* test *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001. MSL2 KDF, MSL2 neural-knockdown female; MSL2 KDM, MSL2 neural-knockdown male;

MSL2 OEF, MSL2-overexpressed female; MSL2 OEM, MSL2-overexpressed male. (**G**) RT-qPCR

analysis of mRNA levels of m^{6}A regulators in the brains of trisomy and MSL2-overexpressed

trisomy *Drosophila* larvae. Student's *t* test *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. 2LF, trisomy 2L

female; 2LM, trisomy 2L male; XXX, metafemale; M2LF, MSL2-overexpressed trisomy 2L female;

M2LM, MSL2-overexpressed trisomy 2L male; MXXX, MSL2-overexpressed metafemale. (**H**) The

expression levels of Ime4 in trisomy and MSL2-overexpressed trisomy embryos represented by

relative fluorescence intensity of probes. The expression of wildtype embryos was set as one.

Sample size = 10. Student's *t* test *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. 2L, trisomy 2L; M2L,

MSL2-overexpressed trisomy 2L; XXX, metafemale; MXXX, MSL2-overexpressed metafemale.

1246

1247

Figure 8. RNA m^{6}A modification may regulate H4K16Ac through histone acetyltransferase MOF.

(**A**) Heatmap of the expression changes of m^{6}A regulators in MOF overexpressing *Drosophila* larvae. The color of the heatmap represents log2(ratio). CF, wildtype female control; CM, wildtype male control; MOF-F, MOF-overexpressed female; MOF-M, MOF-overexpressed male; 2LF, trisomy 2L female; 2LM, trisomy 2L male; MOF-2LF, MOF-overexpressed trisomy 2L female; MOF-2LM, MOF-overexpressed trisomy 2L male. (**B**) RT-qPCR analysis of mRNA levels of Ythdc1 in the heads of MSL2 transgenic *Drosophila.* (**C**) RT-qPCR analysis of mRNA levels of MOF in the heads of *Ythdc1* knockdown *Drosophila*. (**D**) Immunofluorescence of polytene chromosomes showed the signals of SXL (red channel) and MOF (green channel) in the salivary glands of

Drosophila larvae of different genotypes. (**E**) The levels of MOF in the salivary glands of *Ythdc1*

knockdown *Drosophila* quantified by immunofluorescence. (**F**) Immunofluorescence of polytene

chromosomes showed the signals of SXL (red channel) and H4K16Ac (green channel) in the

salivary glands of *Drosophila* larvae of different genotypes. (**G**) The levels of H4K16Ac in the

salivary glands of *Ythdc1* knockdown *Drosophila* quantified by immunofluorescence. Scale bars,

10 μm. Student's *t* test *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. CF, wildtype female control; CM,

wildtype male control; MOF KDF, MOF neural-knockdown female; MOF KDM, MOF

neural-knockdown male; MOF OEF, MOF-overexpressed female; MOF OEM, MOF-overexpressed

male; Ythdc1 KDF, Ythdc1-knockdown female; Ythdc1 KDM, Ythdc1-knockdown male; Ythdc1 OEF,

Ythdc1-overexpressed female; Ythdc1 OEM, Ythdc1-overexpressed male.

1267

1268

1269 **Figure supplements**

1270

1271

Figure 1—figure supplement 1. Primers of m⁶A methyltransferases and reader protein for

RT-qPCR and TSA-FISH. (**A**) The sequences of RT-qPCR primers. (**B**) Agarose gel electrophoresis

validation of qPCR primers. The left lane of each gene is negative control with primers only and

no DNA substrate. (**C**) RT-qPCR analysis of mRNA levels of m⁶A methyltransferases and reader

protein in the brains of wildtype and trisomy *Drosophila* larvae. CF, wildtype female control; CM,

wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale. Student's

t test *p < 0.05, **p < 0.01, ***p < 0.001. (**D**) The sequences of FISH primers. (**E**) The sizes of the

products of the FISH primers after PCR amplification and in vitro transcription were tested by

agarose gel electrophoresis. 2L, chromosome 2 left arm; RT-qPCR, Real-time quantitative PCR;

TSA-FISH, Tyramide signal amplification-based fluorescence *in situ* hybridization.

1282

1283

Figure 1—figure supplement 2. Embryo TSA-FISH of m^6A components in wildtype, trisomy, and MSL2-overexpressed trisomy *Drosophila*. The images represent m^6A component *Ime4* (**A-A"**), *dMettl14* (**B-B"**), *fl(2)d* (**C-C"**), *vir* (**D-D"**), *nito* (**E-E"**), and *Ythdc1* (**F-F"**), respectively. (**A-F**) Gene expression patterns in entire embryos. The genotypes of the samples were shown above, and the development stages (St.) were shown in the left of the pictures. The red pseudocolor represents signal from the probes, and green is the signal of nucleus. Arrowheads indicate regions where the probe signal are enriched. Scale bars, 100 μm. (**A'-F'**) RNA subcellular location patterns of m^6A components. Arrowheads indicate probe signals. Scale bars, 10 μm. (**A"-F"**) The expression levels of m^6A component genes represented as relative fluorescence intensity of the probes compared with DAPI. The expression of wildtype embryos was set as base line. WT, wildtype; 2L, trisomy 2L; M2L, MSL2-overexpressed trisomy 2L. Sample size = 10. Student's *t* test *$p < 0.05$, **$p < 0.01$,

1295   ***p < 0.001.

1296

1297

Figure 2—figure supplement 1. Characteristics of m⁶A-modified genes in wildtype and aneuploid

Drosophila. (**A**) Distribution of m⁶A peaks along each chromosome. (**B**) Kernel density plot

showing the distribution of exon lengths of m⁶A-modified genes. CF, wildtype female control; CM,

wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale. (**C**)

Enrichment analysis performed on the known m⁶A consensus motif DRACH (D = G/A/U, R = G/A,

H = U/A/C). (**D**-**H**) Statistics of the number of m⁶A peaks on m⁶A-modified genes in wildtype

females (**D**), wildtype males (**E**), trisomy 2L females (**F**), trisomy 2L males (**G**), and metafemales

(**H**). (**I**) Venn diagram showing the number of m⁶A-modified genes with more than five m⁶A

peaks.

1307

1308

Figure 2—figure supplement 2. Functional and pathway analysis of m⁶A-modified genes in

wildtype and aneuploid *Drosophila*. (**A**) Functional enrichment analysis of m⁶A-modified genes.

Top 10 enriched GO terms (Biological Process) with adjusted p-value < 0.05 in each sample were

shown. (**B**) UpSet plot showing the shared sets of enriched GO terms of m⁶A-modified genes in

different groups. (**C**) Summary of the common functions enriched by m⁶A-modified genes in all

genotypes. (**D**) KEGG pathway enrichment analysis of m⁶A-modified genes. Top 10 enriched

pathways with p-value < 0.05 in each sample were shown. (**E**) UpSet plot showing the shared sets

of enriched KEGG pathways of m⁶A-modified genes in different groups. (**F**) The pathways

enriched in all genotypes were listed. CF, wildtype female control; CM, wildtype male control; 2LF,

trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale; GO, Gene Ontology; BP, Biological

Process; KEGG, Kyoto Encyclopedia of Genes and Genomes.

1320

1321

Figure 3—figure supplement 1. Characteristics of differentially methylated peak (DMP) associated genes. (**A**) Distribution of differentially methylated peaks (DMPs) along each chromosome. (**B**) Percentages of DMPs localized on different gene features. (**C**) Kernel density plot showing the distribution of exon lengths of DMP-associated genes. (**D**) The top enriched motifs with *de novo* motif analysis of DMPs in each comparison. (**E**) Enrichment analysis performed on the known m^6A consensus motif DRACH (D = G/A/U, R = G/A, H = U/A/C). (**F**) Gene biotypes of DMP-associated genes. (**G**) Venn diagram showing the intersection of DMP-associated genes in each comparison. (**H**-**J**) Statistics of the number of DMPs on DMP-associated genes in

1330 trisomy 2L females compared with wildtype females (**H**), trisomy 2L males compared with

1331 wildtype males (**I**), and metafemales compared with wildtype females (**J**). CF, wildtype female

1332 control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX,

1333 metafemale; 5'UTR, 5' untranslated region; 3'UTR, 3' untranslated region.

1334

1335

Figure 4—figure supplement 1. Functional and pathway analysis of DMP-associated genes. (**A**)

Functional enrichment analysis of DMP-associated genes. Top 10 enriched GO terms (Biological

Process) with adjusted p-value < 0.05 in each comparison were shown. (**B**) UpSet plot showing

the shared sets of enriched GO terms of DMP-associated genes in different aneuploidies. (**C**)

KEGG pathway enrichment analysis of DMP-associated genes. Top 10 enriched pathways with

p-value < 0.1 in each comparison were shown. (**D**) UpSet plot showing the shared sets of

enriched KEGG pathways of DMP-associated genes in different aneuploidies. CF, wildtype female

control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX,

metafemale.

1345

Figure 5—figure supplement 1. Dosage-sensitive modifiers and their interactors in aneuploid *Drosophila*. (**A**-**C**) Networks showing the methylation status of known dosage-sensitive modifiers and the expression changes of genes interact with these modifiers in trisomy 2L females (**A**), trisomy 2L males (**B**), and metafemales (**C**), respectively. Protein-protein interaction (PPI) relationships were obtained from the STRING database. The colors of the heatmaps on chromosomes represent the gene density. Dosage-sensitive modifiers were arranged according to their localizations, with red indicating genes with upregulated DMPs, green indicating genes with

1354 downregulated DMPs, yellow indicating genes with both up- and down-regulated DMPs, and grey

1355 indicating genes without DMPs. The interactors of these modifiers were also separated by

1356 chromosomes, and the colors of the nodes and edges indicate the log2(fold changes) of gene

1357 expression in trisomies. The width of the edges represents the combined score, and only

1358 interactions with combined score ≥ 700 were selected for presentation.

1359

1360

Figure 6—figure supplement 1. Alternative splicing analysis in aneuploidies. (**A**-**C**) Numbers of

differential alternative splicing events in trisomy 2L females (**A**), trisomy 2L males (**B**), and

metafemales (**C**). SE, skipped exon; A5SS, alternative 5' splice site; A3SS, alternative 3' splice site;

MXE, mutually exclusive exons; RI, retained intron. (**D**) Functional enrichment analysis of

differentially alternatively spliced genes. Top 10 enriched GO terms (Biological Process) with

p-value < 0.05 in each comparison were shown. (**E**) KEGG pathway enrichment analysis of differentially alternatively spliced genes. Top 10 enriched pathways with p-value < 0.05 in each comparison were shown. (**F,H**) Genome browser example of *Ppn* (**F**) and *CG13124* (**H**) for indicated MeRIP-seq data. Steelblue color represents input reads, while other colors represent IP reads. Signals were displayed as the mean CPM of two biological replicates. The gene architectures were shown at the bottom (only two representative transcript isoforms were shown). The magenta rectangles at above represent DMPs. The magenta arrowheads indicate the positions of differential alternative splicing. (**G,I**) Sashimi plots depicting RNA sequencing reads and exon junction reads at the positions where the differential splicing events occurs on *Ppn* (**G**) and *CG13124* (**I**). The gene models were shown below. One of the biological replicates was chosen for representation. CF, wildtype female control; CM, wildtype male control; 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale; CPM, Counts per million; RPKM, Reads per kilobase per million mapped reads.

Figure 7—figure supplement 1. The relationships between m⁶A methylation and MSL complex. (**A,C**) Immunofluorescence of polytene chromosomes in MSL2-overexpressed female (**A**) and

male (**C**) larvae. The red channel is the signal from SXL and the green channel is the signal from antibodies of MSL complex components. DNA is stained with DAPI in blue. Scale bars, 10 μm. OE, overexpressed. (**B,D**) RT-qPCR analysis of mRNA levels of MSL2 in the heads of MSL2 transgenic female (**B**) and male (**D**) *Drosophila* adults. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001. OE, overexpressed. (**E**) RT-qPCR analysis of mRNA levels of m^6A regulators in the heads of MSL2 transgenic *Drosophila*. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001. CF, wildtype female control; CM, wildtype male control; OEF, overexpressed female; OEM, overexpressed male; KDF, knockdown female; KDM, knockdown male. (**F**) RT-qPCR analysis of mRNA levels of MSL2 in the heads of m^6A regulators knockdown *Drosophila*. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001. CF, wildtype female control; CM, wildtype male control; OEF, overexpressed female; OEM, overexpressed male; KDF, knockdown female; KDM, knockdown male. (**G**) RT-qPCR analysis of mRNA levels of m^6A regulators in the brains of trisomy and MSL2-overexpressed trisomy *Drosophila* larvae. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001. 2LF, trisomy 2L female; 2LM, trisomy 2L male; XXX, metafemale; M2LF, MSL2-overexpressed trisomy 2L female; M2LM, MSL2-overexpressed trisomy 2L male; MXXX, MSL2-overexpressed metafemale. (**H,I**) Immunofluorescence of polytene chromosomes in MOF-overexpressed female (**H**) and male (**I**) larvae. The red channel is the signal from SXL and the green channel is the signal from MOF or H4K16Ac. DNA is stained with DAPI in blue. Scale bars, 10 μm. OE, overexpressed. (**J**) RT-qPCR analysis of mRNA levels of MOF in the heads of MOF transgenic *Drosophila*. Student's *t* test *p < 0.05, **p < 0.01, ***p < 0.001. CF, wildtype female control; CM, wildtype male control; OEF, overexpressed female; OEM, overexpressed male; KDF, knockdown female; KDM, knockdown male.