Main Manuscript for

N-acetyl-transferases required for iron uptake and aminoglycoside resistance promote
virulence lipid production in *M. marinum*

Bradley S. Jones[1,2], Vikram Pareek[1], Daniel D. Hu[3], Simon D. Weaver[3], Camille
Syska[1,4], Grace Galfano[1], Matthew M. Champion[2,3], Patricia A. Champion[1,2*]

[1]Department of Biological Sciences, [2]Eck Institute for Global Health, [3]Department of
Chemistry and Biochemistry, University of Notre Dame, Notre Dame IN, 46556

[4]Current address: CNRS – CRBM, Montpellier, France

*Patricia A Champion.

Email: pchampio@nd.edu

Author Contributions: Paste the author contributions here.

Competing Interest Statement: None.

Keywords: Mycobacterium, PDIM, GNAT, Acetylation, MbtK, Eis

This PDF file includes:

Main Text
Figures 1 to 5
Table 1

Abstract

Phagosomal lysis is a key aspect of mycobacterial infection of host macrophages. Acety-

lation is a protein modification mediated enzymatically by N-acetyltransferases (NATs)

that impacts bacterial pathogenesis and physiology. To identify NATs required for lytic

activity, we leveraged *Mycobacterium marinum,* a nontubercular pathogen and an estab-

lished model for *M. tuberculosis. M. marinum* hemolysis is a proxy for phagolytic activity.

We generated *M. marinum* strains with deletions in conserved NAT genes and screened

for hemolytic activity. Several conserved lysine acetyltransferases (KATs) contributed to

hemolysis. Hemolysis is mediated by the ESX-1 secretion system and by phthiocerol

dimycocerosate (PDIM), a virulence lipid. For several strains, the hemolytic activity was

restored by the addition of second copy of the ESX-1 locus. Using thin-layer chromatog-

raphy (TLC), we found a single NAT required for PDIM and phenolic glycolipid (PGL)

production. MbtK is a conserved KAT required for mycobactin siderophore synthesis and

virulence. Mycobactin J exogenously complemented PDIM/PGL production in the Δ*mbtK*

strain. The Δ*mbtK M. marinum* strain was attenuated in macrophage and *Galleria*

mellonella infection models. Constitutive expression of either *eis* or *papA5,* which en-

code a KAT required for aminoglycoside resistance and a PDIM/PGL biosynthetic en-

zyme, rescued PDIM/PGL production and virulence of the Δ*mbtK* strain. Eis N-terminally

acetylated PapA5 *in vitro,* supporting a mechanism for restored lipid production. Overall,

our study establishes connections between the MbtK and Eis NATs, and between iron

uptake and PDIM and PGL synthesis in *M. marinum.* Our findings underscore the multi-

functional nature of mycobacterial NATs and their connection to key virulence pathways.

Significance Statement

Acetylation is a modification of protein N-termini, lysine residues, antibiotics and lipids.

Many of the enzymes that promote acetylation belong to the GNAT family of proteins. *M.*

marinum is a well-established as a model to understand how *M. tuberculosis* causes tu-

berculosis. In this study we sought to identify conserved GNAT proteins required for ear-

ly stages of mycobacterial infection. Using *M. marinum,* we determined that several

GNAT proteins are required for the lytic activity of *M. marinum.* We uncovered previously

unknown connections between acetyl-transferases required for iron uptake and antimi-

crobial resistance, and the production of the unique mycobacterial lipids, PDIM and

PGLOur data support that acetyl-transferases from the GNAT family are interconnected,

and have activities beyond those previously reported.

53
54
55 **Main Text**
56
57 **Introduction**
58
Acetylation is the movement of an acetyl group from an acetyl donor to an amine

including N^{α}-amino groups (N-terminal acetylation), the ε-amine of lysine (N^{ε}-acetylation)

or primary amines of nucleotides, peptides, and small molecules (1). Acetylation can be

facilitated enzymatically by N-acetyltransferases (NATs). N-terminal protein acetylation

is considered irreversible because N-terminal deacetylases have not been identified. Ly-

sine acetylation is reversible by lysine deacetylases (2-4) . Protein acetylation regulates

protein folding, stability, localization, and protein-protein interactions (5-11).

Mycobacterium tuberculosis is the cause of the human disease tuberculosis (12).

In 2022, *M. tuberculosis* infected 10.6 million individuals, resulting in an estimated 1.3

million deaths worldwide (13). *M. marinum* causes a tuberculosis-like disease in poikilo-

thermic fish, and is an occasional human pathogen.

M. marinum serves as a model for several aspects of *M. tuberculosis* pathogene-

sis (14, 15). We previously reported the N-terminal acetylomes of *M. tuberculosis* and *M.*

marinum (16) . We found that 15% of *M. tuberculosis* proteins and 11% of *M. marinum*

proteins are acetylated at their N-termini (16). Approximately 16% of the mycobacterial

proteome is acetylated at lysine residues (17)**.** Acetylation mapped to metabolism, anti-

biotic resistance, iron uptake, and lipid synthesis pathways critical to pathogenesis (17-

20)**.**

The majority of predicted and established NATs belong to the GCN-5 Related N-

acetyltransferase (GNAT) superfamily of enzymes (1). *M. tuberculosis* and *M. marinum*

share 23 highly conserved acetyltransferases (16), several of which are capable of acet-

ylating more than one type of target (21-31). Substrates of NATs cannot be predicted by

protein sequence alone, requiring that the substrates be determined empirically (*3*).

Early following infection, mycobacterial pathogens are taken up by professional

phagocytes (32, 33). Following adaptation to the phagosome (34, 35), pathogenic myco-

bacteria use the ESAT-6-system-1 (ESX-1) Type VII secretion system to lyse the

phagosomal membrane and promote interaction with the macrophage cytoplasm (36-

40). Several mycobacterial lipids, including phthiocerol dimycocerosate (PDIM) and phe-

nolic glycolipid (PGL), promote optimal phagosomal lysis (41-43), likely because these

lipids are required for the wild-type levels of ESX-1 effector proteins (44). Several ESX-1

proteins (EsxA, EspA, EspB, EspC, EspL, $EccA_1$, $EccB_1$, $EccCb_1$, $EccE_1$, EsxH, and

EspB), and PDIM/PGL biosynthesis proteins (Mas, PpsA, PpsE, Pks15/1, PapA5 and

DrrA) are N-terminally or lysine acetylated (16, 17, 45). We recently identified the Emp1

(MMAR_1839) NAT as the sole enzyme that promotes N-terminal acetylation of EsxA in

M. marinum in vivo (46). Here, we sought to identify highly conserved GNAT proteins

between *M. marinum* and *M. tuberculosis* that promote virulence through the ESX-1 and

PDIM pathways, leveraging genetics, biochemistry, proteomics, functional assays and

models of infection.

97
98   **Results**
99
Several GNAT proteins contribute to *M. marinum* hemolytic activity. We generated

M. marinum strains lacking individual genes with predicted or established NAT activity

[Table 1, (46)], and the corresponding complementation strains. We confirmed each

strain using PCR (Figure S1) followed by targeted DNA sequencing. ESX-1 and

PDIM/PGL are required for optimal phagolytic activity of *Mycobacterium* (42, 47, 48). We

tested the requirement of 19 conserved NATs for *M. marinum* hemolytic activity, which

can serve as a proxy for phagolytic activity. Six NATs significantly impacted *M. marinum*

hemolytic activity (Fig. 1A), while the remaining 13 NATs did not significantly impact he-

molysis (Fig. S2A). WT *M. marinum* lysed red blood cells (RBCs), while the $\Delta eccCb_1$

strain, which lacks a functional ESX-1 secretion system (37, 49), failed to lyse RBCs

(*P*<0.0001 relative to the WT strain) similar to the cell-free control (PBS). None of the *M.*

marinum strains lacking individual NATs phenocopied the $\Delta eccCb_1$ strain (Figs. 1A and

S2A). However, the deletion of several individual NAT genes significantly reduced he-

molysis compared to the WT strain, including *argA, riml*, *pat, MMAR_3205, MMAR_0332*

and *mbtK* (all *P*<0.0001 vs the WT strain, Fig. 1A). The *Δeis* strain was significantly

more hemolytic than the WT strain (*P*=0.0032). For all strains, except Δ*pat,* expression

of the NAT gene of interest from the mycobacterial optimal promoter significantly in-

creased hemolysis compared to the deletion strain. From these data, we conclude that

several putative and established NATs are required for wild-type hemolytic activity in *M.*

marinum.

Several of the NATs were either established or predicted lysine acetyl transfer-

ases [KATs in Fig 1A indicated by open bars, (21, 24, 26, 50-56)], and are the focus of

this study. To test if the predicted NAT impacted ESX-1 activity, we added an additional

copy of the ESX-1 system to the NAT deletion strains and tested for restoration of he-

molysis. We introduced the pRD1-2F9 cosmid containing the extended RD1 locus from

M. tuberculosis, which includes the majority of the ESX-1 component and substrate

genes (57), into the *M. marinum* strains lacking individual KAT genes, as well as the WT

and $\Delta eccCb_1$ strains. The addition of a second copy of the ESX-1 locus to the WT strain

did not significantly impact hemolysis (Fig. S2B). Because the $eccCb_{1MT}$ (MT = *M. tuber-*

culosis) gene is included on the pRD1-2F9 cosmid, the pRD1-2R9 plasmid complement-

ed the hemolytic activity of the $\Delta eccCb_1$ strain. Interestingly, the pRD1-2F9 cosmid sig-

nificantly increased the hemolytic activity of the Δpat and $\Delta 3205$ strains (P<0.0001, rela-

tive to the parent strains), supporting that Pat and MMAR_3205 impact hemolytic activity

through the ESX-1 system. The pRD1-2F9 cosmid did not significantly impact the hemo-

lytic activity of the $\Delta riml,$ Δeis or $\Delta mbtK$ strains, suggesting a specific connection be-

tween Pat and MMAR_3205.

We next tested if the deletion strains had altered production of PDIM, phenolic

glycolipids (PGLs) and tri-acyl glycerides (TAG). We extracted total lipids from the *M.*

marinum strains lacking individual KAT genes and their respective complementation

strains and measured PGLs, PDIM and TAG using TLC. The WT strain, but not the

Δmas strain produced PDIM, TAG (Fig. 1B, top TLC) and PGL [bottom TLC, lane 1 and

2, (44, 46, 58-63)]. Both PDIM and PGL were reduced and TAG levels were elevated in

total lipids extracted from the $\Delta mbtK$ strain (lane 3). Expression of the *mbtK* gene in the

$\Delta mbtK$ strain increased PDIM and PGL, but did not reduce TAG levels compared to the

$\Delta mbtK$ strain (lane 4). Deletion or complementation of the other potential KAT genes did

not noticeably impact the lipids tested under these conditions (lanes 5-14). From these

data, we conclude that MbtK is required for the production of WT levels of PDIM and

PGLs.

Mycobactin deficiency is linked to reduced growth and PDIM and PGL production

in *M. marinum.*   MbtK in is required for synthesizing mycobactin, a secreted

siderophore required for iron scavenging (55). In the absence of MbtK, *M. tuberculosis*

growth *in vitro* was severely impacted due to the loss of mycobactin production (55).

Likewise, we found that the $\Delta mbtK$ strain exhibited significantly reduced growth com-

pared to the WT and complemented strains (Fig. S3A). We next tested if the reduced

growth and PDIM production in the Δ*mbtK M. marinum* strain could be rescued by sup-

plementing the Δ*mbtK* strain with ferric Mycobactin J. As shown in Figure S3C, addition

of mycobactin to the growth media restored growth of the Δ*mbtK* strain, similar to the

complemented and WT strains. We grew the *M. marinum* strains in the presence of a

range of mycobactin concentrations (from 2 µg/ml to 0.0002 µg/ml or in the absence of

mycobactin) and extracted total lipids. Using TLC, we found that PDIM and PGL produc-

tion was restored to WT levels in the presence of 2 µg/ml or 0.02 µg/ml mycobactin (Fig.

S3C). 0.0002 µg/ml did not restore PDIM or PGL production to the Δ*mbtK* strain From

these data we conclude that the addition of mycobactin J to the grown media trans-

complemented the growth defect and PDIM/PGL production of the Δ*mbtK* strain. These

data support the idea that PDIM and PGL production was reduced in the Δ*mbtK* strain

due to a deficiency in iron scavenging.

MbtK, and its role in mycobactin synthesis, was previously linked to phospholipid

homeostasis (55). However, MbtK has not previously been linked to PDIM and PGL pro-

duction. To gain insight into the mechanisms underlying the phenotypes associated with

the loss of MbtK we used proteomics to measure the changes in protein levels. We gen-

erated cell-associated and secreted protein fractions from the WT, Δ*mbtK,* and the

Δ*mbtK/*p*mbtK* strains (Dataset S1) and measured protein levels in each fraction using

nUHPLC MS-MS/MS quantitative bottom-up proteomics (44, 64). The deletion of *mbtK*

resulted in a significant increase in 15 proteins (Figure S4A), the majority of which are

associated with three chromosomal regions associated with iron uptake, including ESX-3

(EccB3 and MycP3), MMAR_2874, MMAR_2875, cyp278A1 and MMAR_2878, which

are colocalized on the chromosome, and MbtF, MbtE and MbtD, which are colocalized

with MbtK on the chromosome. The Cyp278A1 protein is annotated as being involved in

heme uptake (65). The additional genes are conserved hypothetical proteins. Expression

of *mbtK* in the Δ*mbtK* strain restored protein levels to those in the WT strain. We also

measured the protein levels in the secreted fractions. The levels of the Mbt proteins

were increased in the secreted fraction from the Δ*mbtK* strain as compared to the strains

expressing *mbtK*. There was no significant change in the secretion of any ESX proteins

to explain the reduced hemolytic activity in Fig. 1A. These data did not allow us to con-

nect the loss of MbtK to the production of PDIM and PGL.

Overexpression of *eis* restores virulence lipids to the Δ*mbtK* strain. To understand

the requirement for MbtK in the lipid biosynthesis, we asked if constitutive expression of

the other putative KAT genes could restore wild-type PDIM, PGL and TAG levels to the

Δ*mbtK* strain. We introduced each KAT gene into the Δ*mbtK* strain on an integrating

plasmid, expressed from the MOP promoter, and confirmed the presence of the plasmid

in each strain by PCR (Fig. S5). The expression of the *riml, pat* or *MMAR_3205* genes

did not detectably impact the levels of PDIM, TAG or PGLs in the Δ*mbtK* strain (Fig. 2A,

lanes 5-7). However, constitutive expression of the *eis* gene specifically restored PDIM,

TAG and PGL levels in the Δ*mbtK* strain (lane 8) to those in the WT and complemented

strains.

Eis is a N-acetyltransferase that is clinically linked to aminoglycoside resistance

in *M. tuberculosis* (28, 66-71). To our knowledge, there are no known connections be-

tween the Eis and MbtK acetyltransferases. We first considered if Eis restored lipid pro-

duction simply because it was the most similar NAT to MbtK. Based on structural predic-

tions, MbtK and Eis proteins do not share domains beyond the conserved GNAT do-

mains (AlphaFold, Fig 2B). The RimI, Pat or MMAR_3205 proteins that did not function-

ally complement the Δ*mbtK* also share the GNAT domain. At the amino acid level, Eis,

RimI and MMAR_3205 are all ~20% identical to MbtK (Fig. 2C), and both MbtK1 and

MMAR_3205 are more phylogenetically related KAT to MbtK the Eis (Fig. 2D). There-

fore, we hypothesized that Eis specifically restored PDIM/PGL production to the Δ*mbtK*

strain. Our proteomics analysis shows that expression of *eis* in the Δ*mbtK* strain restored

the levels of 12 of the 15 previously discussed proteins, with the exception of

MMAR_2874, MMAR_2875 and Cyp278A1, which are at the same genetic locus (Fig.

S4). These data suggest that *eis* partially complements the Δ*mbtK* strain.

M. tuberculosis required supplementation with iron during laboratory culture

when the *mbtK$_{MT}$* gene was deleted (55). The Δ*mbtK* (*MMAR_3692*) *M. marinum* strain

exhibited reduced growth without the addition of supplemental iron compared to the WT

strain (Fig. S3A). An orthologous *mbtK* gene, *MMAR_1587*, in *M. marinum* genome

which may allow growth of the Δ*mbtK* strain without additional iron. MMAR_1587 is

28.5% identical at the amino acid level, and is predicted to be structurally similar to MbtK

(Fig. S6A) MMAR_1587 is 54.73% identical to the MbtK$_{MT}$ protein at the amino acid lev-

el, and is annotated as the MbtK ortholog on Mycobrowser (15). Although the

MMAR_3692 protein is 26% identical to MbtK$_{MT}$, the *MMAR_3692* gene is syntenic with

the *mbtK$_{MT}$*. As such, we refer to *MMAR_3692* as *mbtK,* and *MMAR_1587* as *mbtK1.*

(Fig. 2D). *M. marinum* has an orthologous *eis* gene, *MMAR_0151,* which is annotated

as *eis1* (65). The Eis and Eis1 proteins are 31.44% identical, and are predicted to be

structurally similar (Fig. S6A). To further probe the interaction between MbtK and Eis, we

constitutively expressed *mbtK1* and *eis1* in the Δ*mbtK* strain. Constitutive expression of

the *mbtK1* or *eis1* genes did not completely rescue PDIM or PGL levels in the Δ*mbtK*

strain as visualized by TLC (Fig. S6B). These data suggest that additional copies of

MbtK or Eis are functionally redundant, and that Eis likely specifically rescues PDIM,

PGL and TAG production in the Δ*mbtK* strain.

Eis N-terminally acetylates PapA5 *in vitro*. The proteomics data did not provide an

obvious mechanism for how Eis restored PDIM and PGL production to the Δ*mbtK* strain.

We demonstrated that several proteins required for PDIM and/ or PGL production (PpsA,

PapA5, Mas and PpsE, Pks15/1) were N-terminally acetylated [Fig S7, Pink, (*1*)]. Alt-

hough neither MbtK nor Eis have reported to N-terminally acetylate proteins, we tested if

MbtK and/or Eis could N-terminally acetylate one or more biosynthetic proteins required

for PDIM and PGL production.

It was previously established that the first six amino acids of a substrate protein

were sufficient for N-terminal acetylation *in vitro* (30, 72). We had N-terminal six AA pep-

tides synthesized based on the N-terminal peptides we previously measured using pro-

teomics [Fig. 3A, (16)]. Only the Mas peptide contained a "K" residue, suggesting that

we could measure N-terminal acetylation of these peptides. The N-terminal peptides

were used as substrates for *in vitro* acetylation assays with the WT or inactive versions

of MbtK (MbtKE161A, active site mutation as predicted by UniProt) and Eis [EisY128A,

proton donor active site mutant based on (29)] that we expressed and purified from *E.*

coli (Fig. S8). Transferring the acetyl group from Acetyl-CoA to a peptide substrate re-

sults in a free thiol group on the resulting CoA molecule. The reaction of this free thiol

with 5, 5'-dithiobis-(2-nitrobenzoic acid) DTNB results in a colorimetric change that can

be measured at 412nm. The resulting product has 1:1 stoichiometry with the acetylated

peptide and can be used to quantify protein acetylation (29, 73). The RimI NAT from

Salmonella with the S18 substrate [(72, 74), Fig. S8] was a positive control for the reac-

tion. Incubation of the PpsE (*P*=0.0147) or PapA5 (*P*<0.0001) peptides with the WT

MbtK resulted in significantly higher DTNB production than incubation with the

MbtKE161A protein (Fig. 3B). Incubation of the PpsA, Mas or Pks15/1 peptides resulted

in product formation that was not significantly different between the active and inactive

enzymes. These data support that MbtK may acetylate the N-terminal amino acid of

PpsE or PapA5.

257   We conducted the same experiment using Eis or EisY128A, a proton donor ac-

258   tive site mutant (29). Incubation of the PapA5 peptide with the WT Eis protein resulted in

259   significantly higher DTNB production than incubation with the Eis Y128A protein

260   *(P*=0.0096, Fig. 3C). None of the other peptides tested exhibited significantly reduced

261   product formation between the WT and dead enzymes. These data are consistent with

262   the possibility that Eis may N-terminally acetylate PapA5.

263   To directly test the acetylation of PpsE and PapA5 by MbtK and Eis, we used FIA

mass spectrometry. As shown in Figure 3D, MbtK did not N-terminally acetylate the

PpsE peptide. Although we detected the unacetylated peptide (left), we did not detect

the acetylated peptide. We likewise tested if Eis or MbtK could N-terminally acetylate the

PapA5 peptide. The unacetylated PapA5 peptide was detected when mixed with either

Eis (Fig. 3E, top left) or MbtK (bottom left). The acetylated PapA5 peptide was detected

in the presence of Eis, but not MbtK (Fig. 3E, right). We conclude that Eis, but not MbtK,

N-terminally acetylates PapA5 *in vitro*. These data suggest that *eis* expression in the

Δ*mbtK* strain restores PDIM production by N-terminal acetylation of the PapA5 protein

(Fig. 3F). We attempted to measure N-terminal acetylation in the Δ*mbtK* strain, and in

the corresponding complementation and *eis* expression strain (Dataset S1). We were

unable to measure the N-terminal peptide of PapA5 in these strains. We did not identify

N-terminal peptides whose acetylation in the Δ*mbtK* strain was significantly decreased

and was restored by both the *mbtK* and *eis* expression, or by expression of the *eis* gene.

Potential candidates for targets that did not reach significance are in (Fig. S4B). These

data did not allow us to determine if the N-terminal acetylation of PapA5 was promoted

by *eis* expression in *M. marinum*.

We tested if *papA5* could restore PDIM/PGL production or growth to the Δ*mbtK*

strain by collecting total lipids and visualizing PDIM and PGL on TLCs. Expression of

papA5 did not increase PDIM or PGL Levels (Fig. 3F, lanes 2 and 3). Expression of the

papA5 gene in the Δ*mbtK* strain increased the levels of PDIM and PGLs in this strain,

but not to WT levels (lanes 4 and 5, and additional replicates in Fig S9). Together these

data support the conclusion that although PapA5 partially rescued the reduced

PDIM/PGL levels in the Δ*mbtK* strain, it did not rescue growth under the conditions test-

ed.

The requirement of MbtK for mycobacterial pathogenesis can be sup-

pressed by Eis or PapA5. Because the Δ*mbtK$_{MT}$* was attenuated for virulence in a

mouse model of infection (55), we tested if the Δ*mbtK* strain was attenuated in two mod-

els of mycobacterial infection. First, we tested if *mbtK* was required for macrophage cy-

tolysis using Ethidium homodimer (Eth-D1), which is a membrane impermeable nucleic

acid stain (*16*). If *M. marinum* infection perforates the macrophage membrane, the cells

stain red. We infected RAW 264.7 cells with *M. marinum* at an M OI of 5, and counted

the presence of red cells (Fig. 4A). Infection of RAW 264.7 cells with wild-type *M.*

marinum (*P*<0.0001)*,* but not the Δ*eccCb$_1$* strain, resulted in significant levels of cytolysis

compared to the uninfected cells. The Δ*mas* strain and the Δ*mbtK* strains resulted in

significantly less cytolysis than the WT strain ($P<0.0001$). The complementation strains

significantly increased cytolysis compared to the deletion strains ($P<0.0001$) but were

still significantly lower than the WT strain ($P<0.0001$ and $P=0.0090$, respectively). The

Δ*mbtK* strain expressing the *mbtKE161A* gene did not significantly affect cytolysis com-

pared to the Δ*mbtK* strain. From these data we conclude that the NAT activity of MbtK is

required for macrophage cytolysis.

We tested macrophage cytolysis following infection with the Δ*mbtK* strain consti-

tutively expressing the *eis* or the *eisY128A* gene (Fig. 4B). Expression of the *eis* or

eisY128A gene in the Δ*mbtK* strain significantly increased cytolysis compared to the

Δ*mbtK* strain ($P<0.0001$), similar to the WT strain. Infection with the Δ*mbtK* strain ex-

pressing the *eisY128A* gene resulted in significantly reduced cytolysis compared to the

eis gene ($P=0.0001$). We conclude that constitutive expression of *eis* suppresses the

loss of MbtK during macrophage infection, and that this suppression requires the NAT

activity of Eis.

We tested if constitutive expression of *papA5* would likewise restore cytolysis to

the Δ*mbtK* strain. Infection of RAW 264.7 cells with the Δ*mbtK* strain constitutively ex-

pressing *papA5* resulted in significantly more cytolysis than the WT ($P=0.0218$) or

Δ*mbtK* strains ($P<0.0001$, Fig. 4A). To test if suppression of cytolysis by *papA5* expres-

sion was specific we added two controls. First, constitutive expression of *papA5* in the

Δ*mas* strain failed to restore cytolysis to the Δ*mas* strain. Second, constitutive expres-

sion of the *rimI* NAT in the Δ*mbtK* strain failed to restore cytolysis to the Δ*mbtK* strain.

Collectively, our data support that acetylation of PapA5 by Eis bypasses the loss of

MbtK.

We tested if the orthologous *mbtK1* and *eis1* genes were essential for macro-

phage cytolysis. Expression of the *mbtK1* gene partially restored cytolysis to the Δ*mbtK*

strain (Fig. S7B) Expression of the *eis1* gene in the Δ*mbtK* strain resulted in cytolysis

levels significantly higher than the WT strain (*P*<0.0001). Deletion of the *mbtK1* or *eis*

genes significantly reduced macrophage cytolysis compared to the WT strain

(*P*<0.0001), but cytolysis was not restored upon expression of *mbtK1* or *eis* in these

strains. Deletion of the *eis1* gene resulted in a significant reduction of macrophage cytol-

ysis compared to the WT strain, and this was significantly increased upon expression of

the *eis1* gene, though not to WT levels.

To complement the cellular model of infection, we tested the ability of *M.*

marinum to kill the waxworm, *Galleria mellonella* larvae*. G. mellonella* was previously

established as a model for mycobacterial infection (75-79). For each experiment, we in-

fected 30 worms, each injected with 1x10^7 bacteria in a 5 µL volume. Experiments were

performed in biological triplicate. As shown in Figure 4B, WT *M. marinum* killed all of the

infected *G. mellonella* by 7 days post-infection. In contrast, infection with either the

Δ*eccCb₁* or Δ*mbtK* strains resulted in less killing, similar to the uninfected worms. Dele-

tion of *eis* did not significantly impact the killing of *G. mellonella.* Expression of *mbtK* re-

stored the ability of the Δ*mbtK* strain to kill *G. mellonella,* similar to the WT strain. Ex-

pression of either *eis* or *papA5* in the Δ*mbtK* strain partially restored the ability of *M.*

marinum to kill *G. mellonella.* Introduction of the pRD1-2F9 plasmid to the Δ*mbtK* strain

or growing the Δ*mbtK* strain in ferric mycobactin did not restore *G. mellonella* killing to

the ΔmbtK strain (Fig. S3B). Together, these data support that the expression of Eis or

PapA5 suppresses the loss of *mbtK* in *M. marinum.*

345
346   **Discussion**

347
We sought to identify conserved GNAT containing proteins important for the ESX-1 and

PDIM virulence pathways. We found that the majority of GNAT proteins previously anno-

tated as lysine acetyltransferases were required for wild-type levels of hemolysis by *M.*

marinum. We identified several potential NATs required for hemolysis that could be res-

cued by the addition of a second ESX-1 locus. These GNAT proteins will be followed up

in future studies. Our data support a model where mycobactin-mediated iron uptake

promotes the PapA5-mediated biosynthesis of PDIM and PGLs (80). In the absence of

MbtK, the production of PDIM and PGLs is reduced compared to the WT *M. marinum*

strain. PDIM and PGL production was complemented by expression of either *mbtK* or *in*

trans by the addition of Mycobactin J to the growth media. Complementation was de-

pendent upon the acetyltransferase activity of MbtK, as expression of the *mbtKE161A*

allele, which does not have NAT activity, did not restore PDIM and PGL production. We

showed that the expression of *papA5* or the *eis* NAT in the Δ*mbtK* strain specifically

suppressed the requirement of MbtK in PDIM and PGL production. The simplest mecha-

nism, which is supported by our *in vitro* assays, is that Eis N-terminally acetylates Pa-

pA5, promoting the production of PDIM and PGL in the absence of MbtK. Overall, our

studies support a previously undiscovered link between MbtK, iron uptake and the final

step of PDIM and PGL biosynthesis (Fig. 5).

Our model is supported by several lines of evidence previously published in the

literature. It was previously established by Madigan et al that MbtK is required broadly

for lipid and mycobactin production (55). MbtK is required for the synthesis of the lipid

tail of the mycobactin siderophore, but more broadly, the loss of MbtK and iron uptake in

M. tuberculosis resulted in a depletion of phospholipids during iron starvation (55). Alt-

hough Madigan et al connected lipid production to MbtK mediated siderophore iron up-

take in *M. tuberculosis,* the PDIM and PGL lipids were not readily measured using

lipidomics approaches in this study. As such, the connection between mycobactin de-

pendent iron uptake and PDIM/PGL biosynthesis has not been previously investigated.

The connection between iron and PDIM/PGL biosynthesis could be direct. For example,

the PapA5 mediated synthesis step in PDIM biosynthesis could require iron. In support

of this, *papA5* overexpression or the addition of mycobactin restored PDIM/PGL biosyn-

thesis in the Δ*mbtK* strain. Alternatively, the connection between iron and PDIM/PGL

biosynthesis could be indirect. For example, it is possible that iron levels are a signal

that regulate the production of PDIM and PGL. Further experimentation is required to

differentiate between these two possibilities.

The mechanisms connecting MbtK to mycobacterial pathogenesis are unclear.

Yet, it was surprising that constitutive expression of the *papA5* or *eis* genes fully restored

macrophage cytolysis to the Δ*mbtK* strain. Infection of *G. mellonella* revealed that in-

creasing PapA5 or Eis restored virulence to the Δ*mbtK* strain, but the kinetics of killing

differed from that of the WT strain. These data support that perhaps expression of these

genes restored PDIM/PGL production, but did not fully complement the Δ*mbtK* strain.

Further work is required to determine if constitutive expression of *eis* or *papA5* restore

more that the PDIM/PGL deficiency to the Δ*mbtK* strain.

In this study, we uncovered a link between Eis and PapA5. Eis is a bifunctional

N-acetyl-transferase that acetylates aminoglycoside antibiotics to inactivate them. Muta-

tions that lead to *eis* overexpression naturally occur in clinical isolates of *M. tuberculosis,*

leading to kanamycin resistance (28, 66-71). Numerous studies have reported inhibitors

of Eis as a potential path to prevent aminoglycoside resistance during *M. tuberculosis*

infection (81-90). Eis reversibly acetylates lysine residues in the HupB protein, affecting

chromosomal condensation in *M. tuberculosis* (26)*.* We showed that Eis N-terminally

acetylates PapA5 *in vitro*, adding another potential activity to this enzyme. We were un-

able to detect the N-terminal peptide of PapA5 in our proteomics experiment. However,

expression of *papA5* suppressed several phenotypes of the Δ*mbtK* strain supporting this

connection (Fig. 5). If Eis acetylates the N-terminus of PapA5 *in vivo*, this modification

must promote PapA5 activity, allowing suppression of the PDIM/PGL defect in the

Δ*mbtK* strain. Our data further support an emerging literature that NATs, including Eis,

are multifunctional (91).

Knobloch et al previously suggested that the orphan *mbtK* gene (*MMAR_1587*)

was the *M. marinum* ortholog of *M. tuberculosis* MbtK (92). Despite being located at the

mbt locus, *MMAR_3692* was not mentioned in this publication. In our study, we found

that deletion of the *MMAR_3692* gene resulted in reduced growth during standard cultur-

ing conditions. The growth defect was significant, but unlike in *M. tuberculosis,* the dele-

tion of *mbtK* did not abrogate growth. Similar to *M. tuberculosis,* supplementation with

mycobactin restored growth of the Δ*mbtK* strain (55). Deletion of the *mbtK* gene, but not

the *MMAR_1587* gene resulted in a measurable PDIM/PGL defect. Likewise, expression

of *MMAR_1587* in the Δ*mbtK* strain did not restore PDIM levels. From our studies, we

propose that the *MMAR_3692* gene is the *M. marinum* ortholog of the *M. tuberculosis*

mbtK. Our data also suggests that the two genes in *M. marinum* are functionally distinct.

In non-tubercular mycobacterial species, there are two functionally distinct eis

genes. In *M. abscessus,* these are annotated as Eis1Mab and Eis2mab (93). Overex-

pression of *eis2mab,* but not *eis1mab,* resulted in resistance to Kanamycin. We likewise

found that there are two potential *eis* genes in *M. marinum.* Deletion of either *eis* gene

attenuated *M. marinum* in a macrophage model of infection (Fig. S6C). Overexpression

of *eis,* but not *eis1* in *M. marinum* resulted in low level Kan resistance (Fig S10), support-

ing that robust literature linking Eis levels to aminoglycoside resistance, and that the two

eis genes are functionally distinct. Our macrophage infection data supports that overex-

pression of *eis1* restored cytolysis of the Δ*mbtK* strain to greater than WT levels (Fig.

S6). Yet, expression of *eis1* did not restore PDIM/PGL production to the Δ*mbtK* strain.

The mechanisms connecting Eis1 to MbtK need further experimentation.

Overall, this study revealed a new connection between iron scavenging and

PDIM and PGL biosynthesis that requires further investigation. We uncovered that the

Eis NAT, which has been widely linked to drug resistance in *Mycobacterium,* also has

the potential to N-terminally acetylate proteins, including those required for the produc-

tion of the PDIM and PGL lipids. Our data support that NATs cannot be characterized by

the targets they modify, and likely play unexpected, interconnected and as of yet unde-

fined roles in mycobacterial pathogenesis and physiology.

433
434 **Materials and Methods**
435
A summary of the methods are found here. Where necessary, detailed methods are pro-

vided in the Supplementary Information. *Mycobacterium marinum* strains were derived

from the *Mycobacterium marinum* M strain (ATCC BAA-535), and maintained as de-

scribed previously. In frame deletion strains were generated using allelic exchange, and

can be found in TableS1. Complementation plasmids, found in Table S2, were con-

structed using the primers in Table S3 using FastCloning. Site directed mutagenesis was

performed as described previously (94). Hemolysis assays were performed as de-

scribed. RAW 264.7 cells were maintained as described and infected with *M. marinum* at

an MOI of 5. Macrophage cytotoxicity were measured by measuring Ethidium

homodimer uptake as described. *G. mellonella* larval infections were performed by in-

jecting *M. marinum* at a concentration1x10^7/5μL. Thin-layer chromatography experi-

ments was performed as described in (46). Expression and purification of 6xHis-tagged

proteins were expressed in *E. coli,* and purified using Ni-NTA resin, followed by removal

of the epitope tag using TEV. Proteins were used to test activity in vitro against N-

terminal peptides (Genscript) using 5,5-dithio-bis-(2-nitrobenzoic acid) (DTNB) assays

as (73) with minor modifications. Peptide acetylation following the DTNB assay was con-

firmed using MALDI. Bottom-up proteomics was performed after labelling samples with

deuterated acetyl groups as (16) to quantify protein levels and N-terminal acetylation.

Label-free quantification (LFQ) was used for differential protein expression analysis, and

data were processed using R (v4.3.2). The code can be found on github

(https://github.com/Champion-Lab/3692). Rawdata is available through MassIVE and

The Proteome Exchange at ftp://MSV000095200@massive.ucsd.edu with Identifier

MSV000095200 PXD053528.

459
460 **Acknowledgments**
461
We would like to thank Bill Boggess at the Notre Dame Mass Spectrometry and Prote-

omics Facility for continued assistance. Research reported in this publication was sup-

ported by NIAID of the National Institutes of Health under award number AI106872. BSJ

was supported by a Graduate Fellowship from the Eck Institute for Global Health at the

University of Notre Dame. We thank Dr. Micah Ferrell, who constructed several NAT de-

letion strains used in Figure S1.

References

1.   L. Favrot, J. S. Blanchard, O. Vergnolle, Bacterial GCN5-Related N-
Acetyltransferases: From Resistance to Regulation. *Biochemistry* **55**, 989-1002
(2016).

2.   J. Gu *et al.*, Cloning and characterization of NAD-dependent protein deacetylase
(Rv1151c) from Mycobacterium tuberculosis. *Biochemistry (Mosc)* **74**, 743-748
(2009).
3.   J. S. Smith *et al.*, A phylogenetically conserved NAD+-dependent protein
deacetylase activity in the Sir2 protein family. *Proc Natl Acad Sci U S A* **97**, 6658-
6663 (2000).
4.   J. Landry *et al.*, The silencing protein SIR2 and its homologs are NAD-dependent
protein deacetylases. *Proc Natl Acad Sci U S A* **97**, 5807-5811 (2000).
5.   C. S. Hwang, A. Shemorry, A. Varshavsky, N-terminal acetylation of cellular
proteins creates specific degradation signals. *Science* **327**, 973-977 (2010).
6.   A. Drazic, L. M. Myklebust, R. Ree, T. Arnesen, The world of protein acetylation.
Biochim Biophys Acta **1864**, 1372-1401 (2016).
7.   H. Nagai *et al.*, A C-terminal translocation signal required for Dot/Icm-dependent
delivery of the Legionella RalF protein to host cells. *Proc Natl Acad Sci U S A*
102, 826-831 (2005).
8.   W. M. Holmes, B. K. Mannakee, R. N. Gutenkunst, T. R. Serio, Loss of amino-
terminal acetylation suppresses a prion phenotype by modulating global protein
folding. *Nat Commun* **5**, 4383 (2014).
9.   D. C. Scott *et al.*, Blocking an N-terminal acetylation-dependent protein
interaction inhibits an E3 ligase. *Nat Chem Biol* **13**, 850-857 (2017).
10.   M. Ventura *et al.*, Nuclear translocation of glyceraldehyde-3-phosphate
dehydrogenase is regulated by acetylation. *Int J Biochem Cell Biol* **42**, 1672-
1680 (2010).
11.   R. Ree, S. Varland, T. Arnesen, Spotlight on protein N-terminal acetylation. *Exp*
Mol Med **50**, 1-13 (2018).
12.   R. Koch, [Tuberculosis etiology]. *Dtsch Gesundheitsw* **7**, 457-465 (1884).
13.   WHO, Global Tuberculosis Report 2023. (2024).
14.   T. C. Pozos, L. Ramakrishnan, New models for the study of Mycobacterium-host
interactions. *Curr Opin Immunol* **16**, 499-505 (2004).
15.   T. P. Stinear *et al.*, Insights from the complete genome sequence of
Mycobacterium marinum on the evolution of Mycobacterium tuberculosis.
Genome Res **18**, 729-741 (2008).
16.   C. R. Thompson, M. M. Champion, P. A. Champion, Quantitative N-Terminal
Footprinting of Pathogenic Mycobacteria Reveals Differential Protein Acetylation.
J Proteome Res **17**, 3246-3258 (2018).
17.   L. Xie *et al.*, Proteome-wide lysine acetylation profiling of the human pathogen
Mycobacterium tuberculosis. *Int J Biochem Cell Biol* **59**, 193-202 (2015).
18.   F. Liu *et al.*, Acetylome analysis reveals diverse functions of lysine acetylation in
Mycobacterium tuberculosis. *Mol Cell Proteomics* **13**, 3352-3366 (2014).
19.   A. Singhal *et al.*, Systematic Analysis of Mycobacterial Acylation Reveals First
Example of Acylation-mediated Regulation of Enzyme Activity of a Bacterial
Phosphatase. *J Biol Chem* **290**, 26218-26234 (2015).
20.   X. Wu *et al.*, Differential lysine acetylation profiles of Erwinia amylovora strains
revealed by proteomics. *J Proteomics* **79**, 60-71 (2013).
21.   W. Lee, B. C. VanderVen, S. Walker, D. G. Russell, Novel protein
acetyltransferase, Rv2170, modulates carbon and energy metabolism in
Mycobacterium tuberculosis. *Scientific reports* **7**, 72 (2017).

22.   K. B. Arun *et al.*, Acetylation of Isoniazid Is a Novel Mechanism of Isoniazid
521   Resistance in Mycobacterium tuberculosis. *Antimicrob Agents Chemother* **65**
522   (2020).
523   23.   S. Nambi, N. Basu, S. S. Visweswariah, cAMP-regulated protein lysine
524   acetylases in mycobacteria. *J Biol Chem* **285**, 24313-24323 (2010).
525   24.   H. Yang *et al.*, Lysine acetylation of DosR regulates the hypoxia response of
526   Mycobacterium tuberculosis. *Emerg Microbes Infect* **7**, 34 (2018).
527   25.   S. Nambi *et al.*, Cyclic AMP-dependent protein lysine acylation in mycobacteria
528   regulates fatty acid and propionate metabolism. *J Biol Chem* **288**, 14114-14124
529   (2013).
530   26.   K. D. Green *et al.*, Acetylation by Eis and Deacetylation by Rv1151c of
531   Mycobacterium tuberculosis HupB: Biochemical and Structural Insight.
532   *Biochemistry* **57**, 781-790 (2018).
533   27.   K. H. Kim *et al.*, Mycobacterium tuberculosis Eis protein initiates suppression of
534   host immune responses by acetylation of DUSP16/MKP-7. *Proc Natl Acad Sci U*
535   *S A* **109**, 7729-7734 (2012).
536   28.   M. A. Zaunbrecher, R. D. Sikes, Jr., B. Metchock, T. M. Shinnick, J. E. Posey,
537   Overexpression of the chromosomally encoded aminoglycoside acetyltransferase
538   eis confers kanamycin resistance in Mycobacterium tuberculosis. *Proc Natl Acad*
539   *Sci U S A* **106**, 20004-20009 (2009).
540   29.   W. Chen, T. Biswas, V. R. Porter, O. V. Tsodikov, S. Garneau-Tsodikova,
541   Unusual regioversatility of acetyltransferase Eis, a cause of drug resistance in
542   XDR-TB. *Proc Natl Acad Sci U S A* **108**, 9804-9808 (2011).
543   30.   D. Pathak, A. H. Bhat, V. Sapehia, J. Rai, A. Rao, Biochemical evidence for
544   relaxed substrate specificity of Nalpha-acetyltransferase (Rv3420c/rimI) of
545   Mycobacterium tuberculosis. *Scientific reports* **6**, 28892 (2016).
546   31.   P. I. Pletnev *et al.*, Ribosomal protein S18 acetyltransferase RimI is responsible
547   for the acetylation of elongation factor Tu. *J Biol Chem* **298**, 101914 (2022).
548   32.   J. A. Armstrong, P. D. Hart, Response of cultured macrophages to
549   Mycobacterium tuberculosis, with observations on fusion of lysosomes with
550   phagosomes. *J Exp Med* **134**, 713-740 (1971).
551   33.   J. A. Armstrong, P. D. Hart, Phagosome-lysosome interactions in cultured
552   macrophages infected with virulent tubercle bacilli. Reversal of the usual
553   nonfusion pattern and observations on bacterial survival. *J Exp Med* **142**, 1-16
554   (1975).
555   34.   M. A. Fisher, B. B. Plikaytis, T. M. Shinnick, Microarray analysis of the
556   Mycobacterium tuberculosis transcriptional response to the acidic conditions
557   found in phagosomes. *J Bacteriol* **184**, 4025-4032 (2002).
558   35.   K. H. Rohde, R. B. Abramovitch, D. G. Russell, Mycobacterium tuberculosis
559   invasion of macrophages: linking bacterial gene expression to environmental
560   cues. *Cell host & microbe* **2**, 352-364 (2007).
561   36.   T. Hsu *et al.*, The primary mechanism of attenuation of bacillus Calmette-Guerin
562   is a loss of secreted lytic function required for invasion of lung interstitial tissue.
563   *Proc Natl Acad Sci U S A* **100**, 12420-12425 (2003).
564   37.   S. A. Stanley, S. Raghavan, W. W. Hwang, J. S. Cox, Acute infection and
565   macrophage subversion by Mycobacterium tuberculosis require a specialized
566   secretion system. *Proc Natl Acad Sci U S A* **100**, 13001-13006 (2003).

38. P. S. Manzanillo, M. U. Shiloh, D. A. Portnoy, J. S. Cox, Mycobacterium
Tuberculosis Activates the DNA-Dependent Cytosolic Surveillance Pathway
within Macrophages. *Cell host & microbe* **11**, 469-480 (2012).
39. N. van der Wel *et al.*, M. tuberculosis and M. leprae translocate from the
phagolysosome to the cytosol in myeloid cells. *Cell* **129**, 1287-1298 (2007).
40. R. Simeone *et al.*, Phagosomal rupture by Mycobacterium tuberculosis results in
toxicity and host cell death. *PLoS Pathog* **8**, e1002507 (2012).
41. J. Quigley *et al.*, The Cell Wall Lipid PDIM Contributes to Phagosomal Escape
and Host Cell Exit of Mycobacterium tuberculosis. *MBio* **8** (2017).
42. M. M. Osman, A. J. Pagan, J. K. Shanahan, L. Ramakrishnan, Mycobacterium
marinum phthiocerol dimycocerosates enhance macrophage phagosomal
permeabilization and membrane damage. *PloS one* **15**, e0233252 (2020).
43. J. Augenstreich, V. Briken, Host Cell Targets of Released Lipid and Secreted
Protein Effectors of Mycobacterium tuberculosis. *Frontiers in cellular and
infection microbiology* **10**, 595029 (2020).
44. B. S. Jones *et al.*, The loss of the PDIM/PGL virulence lipids causes differential
secretion of ESX-1 substrates in Mycobacterium marinum. *mSphere* **9**, e0000524
(2024).
45. A. G. Birhanu *et al.*, N(epsilon)- and O-Acetylation in Mycobacterium tuberculosis
Lineage 7 and Lineage 4 Strains: Proteins Involved in Bioenergetics, Virulence,
and Antimicrobial Resistance Are Acetylated. *J Proteome Res* **16**, 4045-4059
(2017).
46. O. A. Collars *et al.*, An N-acetyltransferase required for ESAT-6 N-terminal
acetylation and virulence in Mycobacterium marinum. *mBio* **14**, e0098723 (2023).
47. J. Augenstreich *et al.*, ESX-1 and phthiocerol dimycocerosates of Mycobacterium
tuberculosis act in concert to cause phagosomal rupture and host cell apoptosis.
Cell Microbiol 10.1111/cmi.12726 (2017).
48. L. Y. Gao *et al.*, A mycobacterial virulence gene cluster extending RD1 is
required for cytolysis, bacterial spreading and ESAT-6 secretion. *Mol Microbiol*
53, 1677-1693 (2004).
49. K. M. Guinn *et al.*, Individual RD1-region genes are required for export of ESAT-
6/CFP-10 and for virulence of Mycobacterium tuberculosis. *Mol Microbiol* **51**,
359-370 (2004).
50. D. G. Christensen *et al.*, Identification of Novel Protein Lysine Acetyltransferases
in Escherichia coli. *MBio* **9** (2018).
51. H. Xu, S. S. Hegde, J. S. Blanchard, Reversible acetylation and inactivation of
Mycobacterium tuberculosis acetyl-CoA synthetase is dependent on cAMP.
Biochemistry **50**, 5883-5892 (2011).
52. H. J. Lee, P. T. Lang, S. M. Fortune, C. M. Sassetti, T. Alber, Cyclic AMP
regulation of protein lysine acetylation in Mycobacterium tuberculosis. *Nat Struct
Mol Biol* **19**, 811-818 (2012).
53. S. Ghosh, B. Padmanabhan, C. Anand, V. Nagaraja, Lysine acetylation of the
Mycobacterium tuberculosis HU protein modulates its DNA binding and genome
organization. *Mol Microbiol* **100**, 577-588 (2016).
54. O. Vergnolle, H. Xu, J. S. Blanchard, Mechanism and regulation of mycobactin
fatty acyl-AMP ligase FadD33. *J Biol Chem* **288**, 28116-28125 (2013).
55. C. A. Madigan *et al.*, Lipidomic analysis links mycobactin synthase K to iron
uptake and virulence in M. tuberculosis. *PLoS Pathog* **11**, e1004792 (2015).

56. R. Krithika *et al.*, A genetic locus required for iron acquisition in Mycobacterium
tuberculosis. *Proc Natl Acad Sci U S A* **103**, 2069-2074 (2006).
57. P. Brodin *et al.*, Dissection of ESAT-6 system 1 of Mycobacterium tuberculosis
and impact on immunogenicity and virulence. *Infect Immun* **74**, 88-98 (2006).
58. J. Yu *et al.*, Both phthiocerol dimycocerosates and phenolic glycolipids are
required for virulence of Mycobacterium marinum. *Infect Immun* **80**, 1381-1389
(2012).
59. A. K. Azad, T. D. Sirakova, L. M. Rogers, P. E. Kolattukudy, Targeted
replacement of the mycocerosic acid synthase gene in Mycobacterium bovis
BCG produces a mutant that lacks mycosides. *Proc Natl Acad Sci U S A* **93**,
4787-4792 (1996).
60. A. K. Azad, T. D. Sirakova, N. D. Fernandes, P. E. Kolattukudy, Gene knockout
reveals a novel gene cluster for the synthesis of a class of cell wall lipids unique
to pathogenic mycobacteria. *J Biol Chem* **272**, 16741-16745 (1997).
61. O. A. Trivedi *et al.*, Dissecting the mechanism and assembly of a complex
virulence mycobacterial lipid. *Mol Cell* **17**, 631-643 (2005).
62. L. R. Camacho *et al.*, Analysis of the phthiocerol dimycocerosate locus of
Mycobacterium tuberculosis. Evidence that this lipid is involved in the cell wall
permeability barrier. *J Biol Chem* **276**, 19845-19854 (2001).
63. M. Mathur, P. E. Kolattukudy, Molecular cloning and sequencing of the gene for
mycocerosic acid synthase, a novel fatty acid elongating multifunctional enzyme,
from Mycobacterium tuberculosis var. bovis Bacillus Calmette-Guerin. *J Biol
Chem* **267**, 19388-19395 (1992).
64. D. D. Hu, S. D. Weaver, O. A. Collars, P. A. Champion, M. M. Champion, N-
terminal proteomics of Mycobacterium marinum using bottom-up label-free
quantitative analysis in data-dependent acquisition mode on a timsTOF Pro mass
spectrometer. *Microbiol Resour Announc* **13**, e0126323 (2024).
65. A. Kapopoulou, J. M. Lew, S. T. Cole, The MycoBrowser portal: a comprehensive
and manually annotated resource for mycobacterial genomes. *Tuberculosis
(Edinburgh, Scotland)* **91**, 8-13 (2011).
66. S. D. Ley *et al.*, Melting the eis: Nondetection of Kanamycin Resistance Markers
by Routine Diagnostic Tests and Identification of New eis Promoter Variants.
Antimicrob Agents Chemother **65**, e0250220 (2021).
67. M. B. Gikalo, E. Y. Nosova, L. Y. Krylova, A. M. Moroz, The role of eis mutations
in the development of kanamycin resistance in Mycobacterium tuberculosis
isolates from the Moscow region. *J Antimicrob Chemother* **67**, 2107-2109 (2012).
68. A. Sowajassatakul, T. Prammananan, A. Chaiprasert, S. Phunpruch,
Overexpression of eis without a mutation in promoter region of amikacin- and
kanamycin-resistant Mycobacterium tuberculosis clinical strain. *Ann Clin
Microbiol Antimicrob* **17**, 33 (2018).
69. A. Sowajassatakul, T. Prammananan, A. Chaiprasert, S. Phunpruch, Molecular
characterization of amikacin, kanamycin and capreomycin resistance in M/XDR-
TB strains isolated in Thailand. *BMC Microbiol* **14**, 165 (2014).
70. P. Kambli *et al.*, Correlating rrs and eis promoter mutations in clinical isolates of
Mycobacterium tuberculosis with phenotypic susceptibility levels to the second-
line injectables. *Int J Mycobacteriol* **5**, 1-6 (2016).
71. J. Carter, C. R. Consortium, Quantitative measurement of antibiotic resistance in
Mycobacterium tuberculosis reveals genetic determinants of resistance and

susceptibility in a target gene approach. *Res Sq* 10.21203/rs.3.rs-3378915/v1
(2023).
72.   M. W. Vetting, D. C. Bareich, M. Yu, J. S. Blanchard, Crystal structure of RimI
from Salmonella typhimurium LT2, the GNAT responsible for N(alpha)-
acetylation of ribosomal protein S18. *Protein Sci* **17**, 1781-1790 (2008).
73.   H. Foyn, P. R. Thompson, T. Arnesen, DTNB-Based Quantification of In Vitro
Enzymatic N-Terminal Acetyltransferase Activity. *Methods in molecular biology*
(Clifton, N.J **1574**, 9-15 (2017).
74.   A. Yoshikawa, S. Isono, A. Sheback, K. Isono, Cloning and nucleotide
sequencing of the genes rimI and rimJ which encode enzymes acetylating
ribosomal proteins S18 and S5 of Escherichia coli K12. *Mol Gen Genet* **209**, 481-
488 (1987).
75.   E. Rwegasila, E. B. Mubofu, S. S. Nyandoro, P. Erasto, J. J. Munissi,
Preparation, Characterization and in Vivo Antimycobacterial Studies of
Panchovillin-Chitosan Nanocomposites. *Int J Mol Sci* **17** (2016).
76.   M. Meir, T. Grosfeld, D. Barkan, Establishment and Validation of Galleria
mellonella as a Novel Model Organism To Study Mycobacterium abscessus
Infection, Pathogenesis, and Treatment. *Antimicrob Agents Chemother* **62**
(2018).
77.   Y. Li *et al.*, Galleria mellonella - a novel infection model for the Mycobacterium
tuberculosis complex. *Virulence* **9**, 1126-1137 (2018).
78.   F. M. Entwistle, P. J. Coote, Evaluation of greater wax moth larvae, Galleria
mellonella, as a novel in vivo model for non-tuberculosis Mycobacteria infections
and antibiotic treatments. *J Med Microbiol* **67**, 585-597 (2018).
79.   M. Asai *et al.*, Galleria mellonella as an infection model for the virulent
Mycobacterium tuberculosis H37Rv. *Virulence* **13**, 1543-1557 (2022).
80.   S. S. Chavadi *et al.*, The mycobacterial acyltransferase PapA5 is required for
biosynthesis of cell wall-associated phenolic glycolipids. *Microbiology (Reading)*
158, 1379-1387 (2012).
81.   P. Santos-Junior *et al.*, A consensus reverse docking approach for identification
of a competitive inhibitor of acetyltransferase enhanced intracellular survival
protein from Mycobacterium tuberculosis. *Bioorg Med Chem* **108**, 117774 (2024).
82.   L. Radhakrishnan, R. Dani, I. Navabshan, S. Jamal, N. Ahmed, Targeting
Aminoglycoside Acetyltransferase Activity of Mycobacterium tuberculosis
(H37Rv) Derived Eis (Enhanced Intracellular Survival) Protein with Quercetin.
Protein J **43**, 12-23 (2024).
83.   A. H. Pang, K. D. Green, O. V. Tsodikov, S. Garneau-Tsodikova, Discovery and
development of inhibitors of acetyltransferase Eis to combat Mycobacterium
tuberculosis. *Methods Enzymol* **690**, 369-396 (2023).
84.   S. P. Swain *et al.*, In silico studies of alkaloids and their derivatives against N-
acetyltransferase EIS protein from Mycobacterium tuberculosis. *J Biomol Struct*
Dyn 10.1080/07391102.2023.2259487, 1-15 (2023).
85.   A. H. Pang *et al.*, Discovery and Mechanistic Analysis of Structurally Diverse
Inhibitors of Acetyltransferase Eis among FDA-Approved Drugs. *Biochemistry* **62**,
710-721 (2023).
86.   A. H. Pang *et al.*, Discovery of substituted benzyloxy-benzylamine inhibitors of
acetyltransferase Eis and their anti-mycobacterial activity. *Eur J Med Chem* **242**,
114698 (2022).

87. A. Punetha *et al.*, Structure-based design of haloperidol analogues as inhibitors
of acetyltransferase Eis from Mycobacterium tuberculosis to overcome
kanamycin resistance. *RSC Med Chem* **12**, 1894-1909 (2021).
88. A. Punetha *et al.*, Structure-Guided Optimization of Inhibitors of Acetyltransferase
Eis from Mycobacterium tuberculosis. *ACS chemical biology* **15**, 1581-1594
(2020).
89. K. D. Green, A. Punetha, C. Hou, S. Garneau-Tsodikova, O. V. Tsodikov,
Probing the Robustness of Inhibitors of Tuberculosis Aminoglycoside Resistance
Enzyme Eis by Mutagenesis. *ACS Infect Dis* **5**, 1772-1778 (2019).
90. H. X. Ngo *et al.*, Potent 1,2,4-Triazino[5,6 b]indole-3-thioether Inhibitors of the
Kanamycin Resistance Enzyme Eis from Mycobacterium tuberculosis. *ACS
Infect Dis* **4**, 1030-1040 (2018).
91. F. Sanz-Garcia *et al.*, Mycobacterial Aminoglycoside Acetyltransferases: A Little
of Drug Resistance, and a Lot of Other Roles. *Frontiers in microbiology* **10**, 46
(2019).
92. P. Knobloch *et al.*, Mycobacterium marinum produces distinct mycobactin and
carboxymycobactin siderophores to promote growth in broth and phagocytes.
Cell Microbiol **22**, e13163 (2020).
93. K. L. Ung, L. Kremer, M. Blaise, Structural analysis of the N-acetyltransferase
Eis1 from Mycobacterium abscessus reveals the molecular determinants of its
incapacity to modify aminoglycosides. *Proteins* **89**, 94-106 (2021).
94. P. A. Champion, S. A. Stanley, M. M. Champion, E. J. Brown, J. S. Cox, C-
terminal signal sequence promotes virulence factor secretion in Mycobacterium
tuberculosis. *Science* **313**, 1632-1636 (2006).
95. F. Sievers *et al.*, Fast, scalable generation of high-quality protein multiple
sequence alignments using Clustal Omega. *Molecular systems biology* **7**, 539
(2011).
.

Figures and Tables

741
742

Figure 1. At least four Lysine acetyltransferases are required for WT levels of hemolytic activity in *M. marinum*. A. Hemolytic activity of *M. marinum* strains lacking genes encoding predicted NATs. The putative/ known KATs are not filled. -, + refers to absence or presence of complementation plasmid, which gene of interest expressed from the mycobacterial optimal promoter. Significance relative to the WT strain; Brackets report significance between the deletion and complemented strains. **** *P*<0.0001, *** *P*=0.0004, 0.0002, * *P*=0.0250, 0.0298, ** *P*=0.0032 **B.** TLC analysis of total lipids from the indicated *M. marinum* strains. 6 µl of total lipids were analyzed in each experiment. The solvents used to visualize the indicated lipid species are indicated. The results are representative of at least three biological replicates.

753
754

755
756
Figure 2. Overexpression of Eis functionally restores PDIM production in the
Δ*mbtK* strain A. TLC analysis of total lipids from the indicated *M. marinum* strains. The
NAT genes in lanes 4-8 were constitutively expressed in the ΔmbtK strain using the my-
cobacterial optimal strain. 6 µl of total lipids was separated for each strain. The indicated
solvents and lipids visualized are indicated. The TLC is representative of at least three
biological replicates. **B.** Overlay of the predicted structures for MbtK and Eis NATs. The
PDB files used were: https://alphafold.ebi.ac.uk/files/AF-B2HMD6-F1-model_v3.pdb and
https://alphafold.ebi.ac.uk/files/AF-B2HM88-F1-model_v3.pdb. The structures were overlaid
using PyMOL v.2.4.0. **C**. % amino acid identify of the NATs from panel A. Protein se-
quences were obtained from Mycobrowser (65), and aligned using Clustal Omega (95).
D. Clustering by phylogenetic neighbor joining tree without distance corrections, gener-
ated based on protein alignments generated by Clustal Omega.

769
770

771
772
Figure 3. PapA5 is N-terminally acetylated by Eis *in vitro.* A. Table of peptides from
proteins in the PDIM/PGL pathways. B. DTNB assay measuring the production of TNB
(412nm) upon the acetylation of the PpsA, PpsE, Mas, Pks15/1 and PapA5 peptides by
B. WT or inactive MbtK and **C.** the RimI protein from *Salmonella typhimurium* with the
S18 N-terminal peptide (ARYFRR). DTNB assays were performed in technical triplicate.
The data represent the mean product formation from all three experiments. Significance
was determined using a Student's t-test between the active and inactive forms of each
NAT protein. B.* P=0.0417, **** P<0.0001 C. ** P=0.0096. **D.** Flow injection analysis of
the N-terminal peptide of PpsE with purified MbtK protein. Left panel: peaks correspond-
ing to the unacetylated N-terminal peptide of PpsE. Right panel: mass range corre-

sponding to the acetylated version. **E.** Flow injection of the N-terminal peptide of PapA5
incubated with either purified Eis protein or purified MbtK protein. Top left panel: peaks
corresponding to unacetylated PapA5. Top right: the acetylated version of PapA5 when
incubated with Eis. Bottom left panel: peaks corresponding to unacetylated PapA5, Bot-
tom right panel: the same mass range corresponding to the acetylated version of PapA5.
F. TLC on total lipids from *M. marinum.* 6µl total lipid loaded into each lane, PDIM run 3x
consecutively in the solvent indicated, PGLs run once in the solvent indicated. PDIM li-
pids visualized by phosphomolybdic acid and subsequent charring, PGL lipids visualized
by 1% a-naphthol and subsequent charring. The TLC is representative of 3 biological
replicates, the remaining 2 are shown in Fig. S9.

794
795

A

B

796
797

**Figure 4. The requirement of MbtK for mycobacterial pathogenesis can be sup-
pressed by constitutive expression of Eis or PapA5. A**. RAW 264.7 cells infected
with *M. marinum* strains at an MOI of 5. The data is the average of at least 3 independ-
ent infections, each performed in biological triplicate. **** $P<0.0001$, *** $P=0.0001$, **
$P=0.0218$, #### $P<0.0001$. **B.** *Galleria mellonella* infected with the *M. marinum* strains
indicated. Larvae were injected with 1×10^{7} bacteria into the right hindmost proleg. Larvae
were infected in 3 groups of 10, for a total of 30 larvae per strain per infection. Data is
representative of at least 3 biological replicates, each performed in technical triplicate.
Statistical analysis was performed as in (79). Survival curves were compared using a
log-rank Mantel-Cox test, $P<0.0001$.

810   .
811

Figure 5. Model of Interactions between MbtK, Eis and PDIM biosynthesis. This
study revealed that MbtK, likely through siderophore production, is required for the pro-
duction of PDIM and PGL in *M. marinum.* This requirement was complemented in trans
by the addition of Mycobactin J. Constitutive expression of either Eis, or PapA5 specifi-
cally suppressed the loss of PDIM/PGL production in the absence of MbtK, suggesting
that the N-terminal acetylation of PapA5 could promote the restoration of PDIM/PGL
synthesis in the Δ*mbtK* strain.

Table 1. Table of the *M. marinum* genes with their orthologs from *M. tuberculosis*
and their commonly used names.

M. marinum gene	*M. tuberculosis* gene	Name
MMAR_1067	*Rv0730*	
MMAR_1968	*Rv2747*	ArgA
MMAR_4519	*Rv0995*	RimJ
MMAR_1839	*Rv2867c*	Emp1
MMAR_1882	*Rv2851c*	ElaA
MMAR_2463	*Rv1653*	ArgJ
MMAR_0522	*Rv0262c*	Aac
MMAR_1123	*Rv3420c*	RimI
MMAR_4496	*Rv0998*	mPat
MMAR_4889	*Rv0802c*	
MMAR_3205	*Rv2170*	
MMAR_0332	*Rv0133*	
MMAR_1341	*Rv3216*	
MMAR_3740	*Rv2416c*	Eis
MMAR_2836	*Rv3225c*	
MMAR_2615		
MMAR_0151		Eis1
MMAR_3692*	*Rv1347c*	MbtK
MMAR_2744		