Title: N-acetyl-transferases required for iron uptake and aminoglycoside resistance promote viru-

lence lipid production in *M. marinum*

Authors:

Bradley S. Jones[1,2], Vikram Pareek[1], Daniel D. Hu[3], Simon D. Weaver[3], Camille Syska[1,4], Grace

Galfano[1], Matthew M. Champion[2,3], Patricia A. Champion[1,2]

[1]Department of Biological Sciences, [2]Eck Institute for Global Health, [3]Department of Chemistry

and Biochemistry, University of Notre Dame, Notre Dame IN, 46556

[4]Current address: CNRS – CRBM, Montpellier, France

Abstract

Mycobacterial pathogens, including *M. tuberculosis,* the cause of human tuberculosis, and the

non-tubercular species, *M. marinum* exhibit lytic activity in the host phagosome that is essential

for disease. Acetylation is a conserved protein modification that is mediated enzymatically by N-

acetyltransferases (NATs). NATs acetylate protein N^{α}-termini, lysine residues, and small mole-

cules like antibiotics to alter bacterial physiology and pathogenesis. We sought to define con-

served mycobacterial NATs required for lytic activity. We generated and characterized a library

of *M. marinum* strains with deletions in predicted conserved NAT genes*. M. marinum* lyses red

blood cells in a contact dependent manner. We screened our strains for hemolytic activity. We

found that nearly all of the conserved lysine acetyltransferases (KATs) contributed to the hemo-

lytic activity of *M. marinum*. Hemolysis is mediated by both the ESX-1 secretion system and by

phthiocerol dimycocerosate (PDIM), and outer lipid required for virulence. We found that the he-

molytic activity in several KAT deletion strains was restored by the addition of second copy of the

ESX-1 locus. Using thin-layer chromatography (TLC), we found that MbtK was essential for PDIM

and phenolic glycolipid (PGL) production. MbtK is a conserved KAT required for mycobactin si-

derophore synthesis and virulence. We found that the Δ*mbtK M. marinum* strain was attenuated

in macrophage and *Galleria mellonella* infection models. We show that the addition of Mycobactin

J exogenously complements PDIM/PGL production in the *ΔmbtK* strain, linking production of

these lipids to iron uptake. Constitutive expression of either *eis* or *papA5,* which encode a known

KAT and a PDIM/PGL biosynthetic enzyme, rescued PDIM/PGL production and virulence of the

ΔmbtK strain. Eis is an N-acetyltransferase linked to aminoglycoside resistance. We found that

Eis N-terminally acetylated PapA5 *in vitro*, suggesting that Eis modification of PapA5 in the ab-

sence of MbtK restored lipid production. Overall, our study defines new connections between the

MbtK and Eis NATs, and between iron uptake and PDIM and PGL synthesis in *M. marinum*.

Moreover, out findings underscore the multifunctional nature of mycobacterial NATs and their

connection to key virulence pathways.

Importance: Acetylation is a ubiquitous modification of protein N-termini, lysine residues, antibi-

otics and lipids. Many of the enzymes that promote acetylation belong to the GNAT family of

proteins. *M. tuberculosis* is the causative agent of Tuberculosis. *M. marinum* is a related pathogen

that is well established as a model to understand aspects of how *M. tuberculosis* causes disease.

In this study we sought to identify conserved GNAT proteins required for early stages of myco-

bacterial infection. Using *M. marinum,* we determined that several GNAT proteins are required for

the lytic activity of *M. marinum.* We uncovered previously unknown connections between acetyl-

transferases required for iron uptake and antimicrobial resistance, and the production of the

unique mycobacterial lipids, PDIM and PGL. Importantly, our data support that acetyl-transferases

from the GNAT family are interconnected, and have activities beyond those previously reported

in the literature.

Introduction:

Acetylation is the movement of an acetyl group from an acetyl donor to an amine including

N^α-amino groups (N-terminal acetylation), the ε-amine of lysine (N^ε-acetylation) or primary amines

of nucleotides, peptides, and small molecules (1). Acetylation can be facilitated enzymatically by

N-acetyltransferases (NATs). N-terminal protein acetylation is considered irreversible because N-

terminal deacetylases have not been identified. Lysine acetylation is reversible by lysine deacety-

lases (2-4) . Protein acetylation regulates protein folding, stability, localization, and protein-protein

interactions (5-11).

Mycobacterium tuberculosis is the cause of the human disease tuberculosis (12). In 2022,

M. tuberculosis infected 10.6 million individuals, resulting in an estimated 1.3 million deaths world-

wide (13). *M. marinum* causes a tuberculosis-like disease in poikilothermic fish, and is an occa-

sional human pathogen.

M. marinum serves as a model for several aspects of *M. tuberculosis* pathogenesis (14,

15). We previously reported the N-terminal acetylomes of *M. tuberculosis* and *M. marinum (16)* .

We found that 15% of *M. tuberculosis* proteins and 11% of *M. marinum* proteins are acetylated

at their N-termini (16). Approximately 16% of the mycobacterial proteome is acetylated at lysine

residues (17)**.** Acetylation mapped to metabolism, antibiotic resistance, iron uptake, and lipid syn-

thesis pathways critical to pathogenesis (17-20)**.**

The majority of predicted and established NATs belong to the GCN-5 Related N-acetyl-

transferase (GNAT) superfamily of enzymes (1). *M. tuberculosis* and *M. marinum* share 23 highly

conserved acetyltransferases (16), several of which are capable of acetylating more than one

type of target (21-31). Substrates of NATs cannot be predicted by protein sequence alone, requir-

ing that the substrates be determined empirically (*3*).

Early following infection, mycobacterial pathogens are taken up by professional phago-

cytes (32, 33). Following adaptation to the phagosome (34, 35), pathogenic mycobacteria use the

ESAT-6-system-1 (ESX-1) Type VII secretion system to lyse the phagosomal membrane and

promote interaction with the macrophage cytoplasm (36-40). Several mycobacterial lipids, includ-

ing phthiocerol dimycocerosate (PDIM) and phenolic glycolipid (PGL), promote optimal phagoso-

mal lysis (41-43), likely because these lipids are required for the wild-type levels of ESX-1 effector

proteins (44). Several ESX-1 proteins (EsxA, EspA, EspB, EspC, EspL, EccA$_1$, EccB$_1$, EccCb$_1$,

EccE$_1$, EsxH, and EspB), and PDIM/PGL biosynthesis proteins (Mas, PpsA, PpsE, Pks15/1,

PapA5 and DrrA) are N-terminally or lysine acetylated (16, 17, 45). We recently identified the

Emp1 (MMAR_1839) NAT as the sole enzyme that promotes N-terminal acetylation of EsxA in

M. marinum in vivo (46). Here, we sought to identify highly conserved GNAT proteins between *M.*

marinum and *M. tuberculosis* that promote virulence through the ESX-1 and PDIM pathways,

leveraging genetics, biochemistry, proteomics, functional assays and models of infection.

86
Results:

Several GNAT proteins contribute to *M. marinum* hemolytic activity. We generated *M. mari-*

num strains lacking individual genes with predicted or established NAT activity [Table 1, (46)],

and the corresponding complementation strains. We confirmed each strain using PCR (Figure

S1) followed by targeted DNA sequencing. ESX-1 and PDIM/PGL are required for optimal phag-

olytic activity of *Mycobacterium* (42, 47, 48). We tested the requirement of 19 conserved NATs

for *M. marinum* hemolytic activity, which can serve as a proxy for phagolytic activity. Six NATs

significantly impacted *M. marinum* hemolytic activity (Fig. 1A), while the remaining 13 NATs did

not significantly impact hemolysis (Fig. S2A). WT *M. marinum* lysed red blood cells (RBCs), while

the Δ*eccCb$_1$* strain, which lacks a functional ESX-1 secretion system (37, 49), failed to lyse RBCs

($P<0.0001$ relative to the WT strain) similar to the cell-free control (PBS). None of the *M. marinum*

strains lacking individual NATs phenocopied the Δ*eccCb$_1$* strain (Figs. 1A and S2A). However,

the deletion of several individual NAT genes significantly reduced hemolysis compared to the WT

strain, including *argA, rimI*, *pat, MMAR_3205, MMAR_0332* and *mbtK* (all $P<0.0001$ vs the WT

strain*,* Fig. 1A). The Δ*eis* strain was significantly more hemolytic than the WT strain ($P=0.0032$).

For all strains, except Δ*pat,* expression of the NAT gene of interest from the mycobacterial optimal

promoter significantly increased hemolysis compared to the deletion strain. From these data, we

conclude that several putative and established NATs are required for wild-type hemolytic activity

in *M. marinum.*

Several of the NATs were either established or predicted lysine acetyl transferases [KATs

in Fig 1A indicated by open bars, (21, 24, 26, 50-56)], and are the focus of this study. To test if

the predicted NAT impacted ESX-1 activity, we added an additional copy of the ESX-1 system to

the NAT deletion strains and tested for restoration of hemolysis. We introduced the pRD1-2F9

cosmid containing the extended RD1 locus from *M. tuberculosis*, which includes the majority of

the ESX-1 component and substrate genes (57), into the *M. marinum* strains lacking individual

KAT genes, as well as the WT and $\Delta eccCb_1$ strains. The addition of a second copy of the ESX-1

locus to the WT strain did not significantly impact hemolysis (Fig. S2B). Because the $eccCb_{1MT}$

(MT = *M. tuberculosis*) gene is included on the pRD1-2F9 cosmid, the pRD1-2R9 plasmid com-

plemented the hemolytic activity of the $\Delta eccCb_1$ strain. Interestingly, the pRD1-2F9 cosmid sig-

nificantly increased the hemolytic activity of the Δpat and $\Delta 3205$ strains ($P<0.0001$, relative to the

parent strains), supporting that Pat and MMAR_3205 impact hemolytic activity through the ESX-

1 system. The pRD1-2F9 cosmid did not significantly impact the hemolytic activity of the $\Delta rimI$,

Δeis or $\Delta mbtK$ strains, suggesting a specific connection between Pat and MMAR_3205.

We next tested if the deletion strains had altered production of PDIM, phenolic glycolipids

(PGLs) and tri-acyl glycerides (TAG). We extracted total lipids from the *M. marinum* strains lacking

individual KAT genes and their respective complementation strains and measured PGLs, PDIM

and TAG using TLC. The WT strain, but not the Δmas strain produced PDIM, TAG (Fig. 1B, top

TLC) and PGL [bottom TLC, lane 1 and 2, (44, 46, 58-63)]. Both PDIM and PGL were reduced

and TAG levels were elevated in total lipids extracted from the $\Delta mbtK$ strain (lane 3). Expression

of the *mbtK* gene in the $\Delta mbtK$ strain increased PDIM and PGL, but did not reduce TAG levels

compared to the $\Delta mbtK$ strain (lane 4). Deletion or complementation of the other potential KAT

genes did not noticeably impact the lipids tested under these conditions (lanes 5-14). From these

data, we conclude that MbtK is required for the production of WT levels of PDIM and PGLs.

Mycobactin deficiency is linked to reduced growth and PDIM and PGL production in *M. marinum.* MbtK in is required for synthesizing mycobactin, a secreted siderophore required for iron scavenging (55). In the absence of MbtK, *M. tuberculosis* growth *in vitro* was severely impacted due to the loss of mycobactin production (55). Likewise, we found that the Δ*mbtK* strain exhibited significantly reduced growth compared to the WT and complemented strains (Fig. S3A). We next tested if the reduced growth and PDIM production in the Δ*mbtK M. marinum* strain could be rescued by supplementing the Δ*mbtK* strain with ferric Mycobactin J. As shown in Figure S3C, addition of mycobactin to the growth media restored growth of the Δ*mbtK* strain, similar to the complemented and WT strains. We grew the *M. marinum* strains in the presence of a range of mycobactin concentrations (from 2 µg/ml to 0.0002 µg/ml or in the absence of mycobactin) and extracted total lipids. Using TLC, we found that PDIM and PGL production was restored to WT levels in the presence of 2 µg/ml or 0.02 µg/ml mycobactin (Fig. S3C). 0.0002 µg/ml did not restore PDIM or PGL production to the Δ*mbtK* strain From these data we conclude that the addition of mycobactin J to the grown media trans-complemented the growth defect and PDIM/PGL production of the Δ*mbtK* strain. These data support the idea that PDIM and PGL production was reduced in the Δ*mbtK* strain due to a deficiency in iron scavenging.

MbtK, and its role in mycobactin synthesis, was previously linked to phospholipid homeostasis (55). However, MbtK has not previously been linked to PDIM and PGL production. To gain insight into the mechanisms underlying the phenotypes associated with the loss of MbtK we used proteomics to measure the changes in protein levels. We generated cell-associated and secreted protein fractions from the WT, Δ*mbtK,* and the Δ*mbtK*/p*mbtK* strains (Dataset S1) and measured protein levels in each fraction using nUHPLC MS-MS/MS quantitative bottom-up proteomics (44, 64). The deletion of *mbtK* resulted in a significant increase in 15 proteins (Figure S4A), the majority of which are associated with three chromosomal regions associated with iron uptake, including ESX-3 (EccB3 and MycP3), MMAR_2874, MMAR_2875, cyp278A1 and MMAR_2878, which

are colocalized on the chromosome, and MbtF, MbtE and MbtD, which are colocalized with MbtK

on the chromosome. The Cyp278A1 protein is annotated as being involved in heme uptake (65).

The additional genes are conserved hypothetical proteins. Expression of *mbtK* in the Δ*mbtK* strain

restored protein levels to those in the WT strain. We also measured the protein levels in the

secreted fractions. The levels of the Mbt proteins were increased in the secreted fraction from the

Δ*mbtK* strain as compared to the strains expressing *mbtK*. There was no significant change in

the secretion of any ESX proteins to explain the reduced hemolytic activity in Fig. 1A. These data

did not allow us to connect the loss of MbtK to the production of PDIM and PGL.

Overexpression of *eis* restores virulence lipids to the Δ*mbtK* strain. To understand the re-

quirement for MbtK in the lipid biosynthesis, we asked if constitutive expression of the other pu-

tative KAT genes could restore wild-type PDIM, PGL and TAG levels to the Δ*mbtK* strain. We

introduced each KAT gene into the Δ*mbtK* strain on an integrating plasmid, expressed from the

MOP promoter, and confirmed the presence of the plasmid in each strain by PCR (Fig. S5). The

expression of the *rimI, pat* or *MMAR_3205* genes did not detectably impact the levels of PDIM,

TAG or PGLs in the Δ*mbtK* strain (Fig. 2A, lanes 5-7). However, constitutive expression of the

eis gene specifically restored PDIM, TAG and PGL levels in the Δ*mbtK* strain (lane 8) to those in

the WT and complemented strains.

Eis is a N-acetyltransferase that is clinically linked to aminoglycoside resistance in *M. tu-*

berculosis (28, 66-71)*.* To our knowledge, there are no known connections between the Eis and

MbtK acetyltransferases. We first considered if Eis restored lipid production simply because it was

the most similar NAT to MbtK. Based on structural predictions, MbtK and Eis proteins do not share

domains beyond the conserved GNAT domains (AlphaFold, Fig 2B). The RimI*,* Pat or

MMAR_3205 proteins that did not functionally complement the Δ*mbtK* also share the GNAT do-

main*.* At the amino acid level, Eis, RimI and MMAR_3205 are all ~20% identical to MbtK (Fig.

2C), and both MbtK1 and MMAR_3205 are more phylogenetically related KAT to MbtK the Eis

(Fig. 2D). Therefore, we hypothesized that Eis specifically restored PDIM/PGL production to the

Δ*mbtK* strain. Our proteomics analysis shows that expression of *eis* in the Δ*mbtK* strain restored

the levels of 12 of the 15 previously discussed proteins, with the exception of MMAR_2874,

MMAR_2875 and Cyp278A1, which are at the same genetic locus (Fig. S4). These data suggest

that *eis* partially complements the Δ*mbtK* strain.

M. tuberculosis required supplementation with iron during laboratory culture when the

mbtK$_{MT}$ gene was deleted (55). The Δ*mbtK* (*MMAR_3692*) *M. marinum* strain exhibited reduced

growth without the addition of supplemental iron compared to the WT strain (Fig. S3A). An orthol-

ogous *mbtK* gene, *MMAR_1587*, in *M. marinum* genome which may allow growth of the Δ*mbtK*

strain without additional iron. MMAR_1587 is 28.5% identical at the amino acid level, and is pre-

dicted to be structurally similar to MbtK (Fig. S6A) MMAR_1587 is 54.73% identical to the MbtK$_{MT}$

protein at the amino acid level, and is annotated as the MbtK ortholog on Mycobrowser (15).

Although the MMAR_3692 protein is 26% identical to MbtK$_{MT}$, the *MMAR_3692* gene is syntenic

with the *mbtK$_{MT}$*. As such, we refer to *MMAR_3692* as *mbtK,* and *MMAR_1587* as *mbtK1.* (Fig.

2D). *M. marinum* has an orthologous *eis* gene, *MMAR_0151,* which is annotated as *eis1* (65).

The Eis and Eis1 proteins are 31.44% identical, and are predicted to be structurally similar (Fig.

S6A). To further probe the interaction between MbtK and Eis, we constitutively expressed *mbtK1*

and *eis1* in the Δ*mbtK* strain. Constitutive expression of the *mbtK1* or *eis1* genes did not com-

pletely rescue PDIM or PGL levels in the Δ*mbtK* strain as visualized by TLC (Fig. S6B). These

data suggest that additional copies of MbtK or Eis are functionally redundant*,* and that Eis likely

specifically rescues PDIM, PGL and TAG production in the Δ*mbtK* strain.

Eis N-terminally acetylates PapA5 *in vitro*. The proteomics data did not provide an obvious

mechanism for how Eis restored PDIM and PGL production to the Δ*mbtK* strain. We demonstrated

that several proteins required for PDIM and/ or PGL production (PpsA, PapA5, Mas and PpsE,

Pks15/1) were N-terminally acetylated [Fig S7, Pink, (*1*)]. Although neither MbtK nor Eis have

reported to N-terminally acetylate proteins, we tested if MbtK and/or Eis could N-terminally acet-

ylate one or more biosynthetic proteins required for PDIM and PGL production.

It was previously established that the first six amino acids of a substrate protein were

sufficient for N-terminal acetylation *in vitro* (30, 72). We had N-terminal six AA peptides synthe-

sized based on the N-terminal peptides we previously measured using proteomics [Fig. 3A, (16)].

Only the Mas peptide contained a "K" residue, suggesting that we could measure N-terminal acet-

ylation of these peptides. The N-terminal peptides were used as substrates for *in vitro* acetylation

assays with the WT or inactive versions of MbtK (MbtKE161A, active site mutation as predicted

by UniProt) and Eis [EisY128A, proton donor active site mutant based on (29)] that we expressed

and purified from *E. coli* (Fig. S8). Transferring the acetyl group from Acetyl-CoA to a peptide

substrate results in a free thiol group on the resulting CoA molecule. The reaction of this free thiol

with 5, 5'-dithiobis-(2-nitrobenzoic acid) DTNB results in a colorimetric change that can be meas-

ured at 412nm. The resulting product has 1:1 stoichiometry with the acetylated peptide and can

be used to quantify protein acetylation (29, 73). The RimI NAT from *Salmonella* with the S18

substrate [(72, 74), Fig. S8] was a positive control for the reaction. Incubation of the PpsE

(P=0.0147) or PapA5 (P<0.0001) peptides with the WT MbtK resulted in significantly higher DTNB

production than incubation with the MbtKE161A protein (Fig. 3B). Incubation of the PpsA, Mas or

Pks15/1 peptides resulted in product formation that was not significantly different between the

active and inactive enzymes. These data support that MbtK may acetylate the N-terminal amino

acid of PpsE or PapA5.

We conducted the same experiment using Eis or EisY128A, a proton donor active site

mutant (29). Incubation of the PapA5 peptide with the WT Eis protein resulted in significantly

higher DTNB production than incubation with the Eis Y128A protein *(P=0.0096, Fig. 3C)*. None of

the other peptides tested exhibited significantly reduced product formation between the WT and

dead enzymes. These data are consistent with the possibility that Eis may N-terminally acetylate

PapA5.

To directly test the acetylation of PpsE and PapA5 by MbtK and Eis, we used FIA mass

spectrometry. As shown in Figure 3D, MbtK did not N-terminally acetylate the PpsE peptide. Alt-

hough we detected the unacetylated peptide (left), we did not detect the acetylated peptide. We

likewise tested if Eis or MbtK could N-terminally acetylate the PapA5 peptide. The unacetylated

PapA5 peptide was detected when mixed with either Eis (Fig. 3E, top left) or MbtK (bottom left).

The acetylated PapA5 peptide was detected in the presence of Eis, but not MbtK (Fig. 3E, right).

We conclude that Eis, but not MbtK, N-terminally acetylates PapA5 *in vitro*. These data suggest

that *eis* expression in the Δ*mbtK* strain restores PDIM production by N-terminal acetylation of the

PapA5 protein (Fig. 3F). We attempted to measure N-terminal acetylation in the Δ*mbtK* strain,

and in the corresponding complementation and *eis* expression strain (Dataset S1). We were un-

able to measure the N-terminal peptide of PapA5 in these strains. We did not identify N-terminal

peptides whose acetylation in the Δ*mbtK* strain was significantly decreased and was restored by

both the *mbtK* and *eis* expression, or by expression of the *eis* gene. Potential candidates for

targets that did not reach significance are in (Fig. S4B). These data did not allow us to determine

if the N-terminal acetylation of PapA5 was promoted by *eis* expression in *M. marinum*.

We tested if *papA5* could restore PDIM/PGL production or growth to the Δ*mbtK* strain by

collecting total lipids and visualizing PDIM and PGL on TLCs. Expression of *papA5* did not in-

crease PDIM or PGL Levels (Fig. 3F, lanes 2 and 3). Expression of the *papA5* gene in the Δ*mbtK*

strain increased the levels of PDIM and PGLs in this strain, but not to WT levels (lanes 4 and 5,

and additional replicates in Fig S9). Together these data support the conclusion that although

PapA5 partially rescued the reduced PDIM/PGL levels in the Δ*mbtK* strain, it did not rescue

growth under the conditions tested.

The requirement of MbtK for mycobacterial pathogenesis can be suppressed by Eis

or PapA5. Because the Δ*mbtK$_{MT}$* was attenuated for virulence in a mouse model of infection (55),

we tested if the Δ*mbtK* strain was attenuated in two models of mycobacterial infection. First, we

tested if *mbtK* was required for macrophage cytolysis using Ethidium homodimer (Eth-D1), which

is a membrane impermeable nucleic acid stain (*16*). If *M. marinum* infection perforates the mac-

rophage membrane, the cells stain red. We infected RAW 264.7 cells with *M. marinum* at an M

OI of 5, and counted the presence of red cells (Fig. 4A). Infection of RAW 264.7 cells with wild-

type *M. marinum* ($P<0.0001$)*,* but not the $\Delta eccCb_1$ strain, resulted in significant levels of cytolysis

compared to the uninfected cells. The Δmas strain and the $\Delta mbtK$ strains resulted in significantly

less cytolysis than the WT strain ($P<0.0001$). The complementation strains significantly increased

cytolysis compared to the deletion strains ($P<0.0001$) but were still significantly lower than the

WT strain ($P<0.0001$ and $P=0.0090$, respectively). The $\Delta mbtK$ strain expressing the *mbtKE161A*

gene did not significantly affect cytolysis compared to the $\Delta mbtK$ strain. From these data we con-

clude that the NAT activity of MbtK is required for macrophage cytolysis.

We tested macrophage cytolysis following infection with the $\Delta mbtK$ strain constitutively

expressing the *eis* or the *eisY128A* gene (Fig. 4B). Expression of the *eis* or *eisY128A* gene in the

$\Delta mbtK$ strain significantly increased cytolysis compared to the $\Delta mbtK$ strain ($P<0.0001$), similar

to the WT strain. Infection with the $\Delta mbtK$ strain expressing the *eisY128A* gene resulted in signif-

icantly reduced cytolysis compared to the *eis* gene ($P=0.0001$). We conclude that constitutive

expression of *eis* suppresses the loss of MbtK during macrophage infection, and that this sup-

pression requires the NAT activity of Eis.

We tested if constitutive expression of *papA5* would likewise restore cytolysis to the $\Delta mbtK$

strain. Infection of RAW 264.7 cells with the $\Delta mbtK$ strain constitutively expressing *papA5* resulted

in significantly more cytolysis than the WT ($P=0.0218$) or $\Delta mbtK$ strains ($P<0.0001$, Fig. 4A). To

test if suppression of cytolysis by *papA5* expression was specific we added two controls. First,

constitutive expression of *papA5* in the Δmas strain failed to restore cytolysis to the Δmas strain.

Second, constitutive expression of the *rimI* NAT in the $\Delta mbtK$ strain failed to restore cytolysis to

the $\Delta mbtK$ strain. Collectively, our data support that acetylation of PapA5 by Eis bypasses the

loss of MbtK.

We tested if the orthologous *mbtK1* and *eis1* genes were essential for macrophage cytol-

ysis. Expression of the *mbtK1* gene partially restored cytolysis to the Δ*mbtK* strain (Fig. S7B)

Expression of the *eis1* gene in the Δ*mbtK* strain resulted in cytolysis levels significantly higher

than the WT strain (*P*<0.0001). Deletion of the *mbtK1* or *eis* genes significantly reduced macro-

phage cytolysis compared to the WT strain (*P*<0.0001), but cytolysis was not restored upon ex-

pression of *mbtK1* or *eis* in these strains. Deletion of the *eis1* gene resulted in a significant reduc-

tion of macrophage cytolysis compared to the WT strain, and this was significantly increased upon

expression of the *eis1* gene, though not to WT levels.

To complement the cellular model of infection, we tested the ability of *M. marinum* to kill

the waxworm, *Galleria mellonella* larvae. *G. mellonella* was previously established as a model for

mycobacterial infection (75-79). For each experiment, we infected 30 worms, each injected with

1×10^7 bacteria in a 5 µL volume. Experiments were performed in biological triplicate. As shown in

Figure 4B, WT *M. marinum* killed all of the infected *G. mellonella* by 7 days post-infection. In

contrast, infection with either the Δ*eccCb_1* or Δ*mbtK* strains resulted in less killing, similar to the

uninfected worms. Deletion of *eis* did not significantly impact the killing of *G. mellonella.* Expres-

sion of *mbtK* restored the ability of the Δ*mbtK* strain to kill *G. mellonella,* similar to the WT strain.

Expression of either *eis* or *papA5* in the Δ*mbtK* strain partially restored the ability of *M. marinum*

to kill *G. mellonella.* Introduction of the pRD1-2F9 plasmid to the Δ*mbtK* strain or growing the

Δ*mbtK* strain in ferric mycobactin did not restore *G. mellonella* killing to the ΔmbtK strain (Fig.

S3B). Together, these data support that the expression of Eis or PapA5 suppresses the loss of

mbtK in *M. marinum.*

Discussion:

We sought to identify conserved GNAT containing proteins important for the ESX-1 and PDIM

virulence pathways. We found that the majority of GNAT proteins previously annotated as lysine

acetyltransferases were required for wild-type levels of hemolysis by *M. marinum.* We identified

several potential NATs required for hemolysis that could be rescued by the addition of a second

ESX-1 locus. These GNAT proteins will be followed up in future studies. Our data support a model

where mycobactin-mediated iron uptake promotes the PapA5-mediated biosynthesis of PDIM and

PGLs (80). In the absence of MbtK, the production of PDIM and PGLs is reduced compared to

the WT *M. marinum* strain. PDIM and PGL production was complemented by expression of either

mbtK or *in trans* by the addition of Mycobactin J to the growth media. Complementation was

dependent upon the acetyltransferase activity of MbtK, as expression of the *mbtKE161A* allele,

which does not have NAT activity, did not restore PDIM and PGL production. We showed that the

expression of *papA5* or the *eis* NAT in the ΔmbtK strain specifically suppressed the requirement

of MbtK in PDIM and PGL production. The simplest mechanism, which is supported by our *in vitro*

assays, is that Eis N-terminally acetylates PapA5, promoting the production of PDIM and PGL in

the absence of MbtK. Overall, our studies support a previously undiscovered link between MbtK,

iron uptake and the final step of PDIM and PGL biosynthesis (Fig. 5).

Our model is supported by several lines of evidence previously published in the literature.

It was previously established by Madigan et al that MbtK is required broadly for lipid and myco-

bactin production (55). MbtK is required for the synthesis of the lipid tail of the mycobactin sider-

ophore, but more broadly, the loss of MbtK and iron uptake in *M. tuberculosis* resulted in a deple-

tion of phospholipids during iron starvation (55). Although Madigan et al connected lipid produc-

tion to MbtK mediated siderophore iron uptake in *M. tuberculosis,* the PDIM and PGL lipids were

not readily measured using lipidomics approaches in this study. As such, the connection between

mycobactin dependent iron uptake and PDIM/PGL biosynthesis has not been previously investi-

gated. The connection between iron and PDIM/PGL biosynthesis could be direct. For example,

the PapA5 mediated synthesis step in PDIM biosynthesis could require iron. In support of this,

papA5 overexpression or the addition of mycobactin restored PDIM/PGL biosynthesis in the

ΔmbtK strain. Alternatively, the connection between iron and PDIM/PGL biosynthesis could be

indirect. For example, it is possible that iron levels are a signal that regulate the production of

PDIM and PGL. Further experimentation is required to differentiate between these two possibili-

ties.

The mechanisms connecting MbtK to mycobacterial pathogenesis are unclear. Yet, it was

surprising that constitutive expression of the *papA5* or *eis* genes fully restored macrophage cytol-

ysis to the Δ*mbtK* strain. Infection of *G. mellonella* revealed that increasing PapA5 or Eis restored

virulence to the Δ*mbtK* strain, but the kinetics of killing differed from that of the WT strain. These

data support that perhaps expression of these genes restored PDIM/PGL production, but did not

fully complement the Δ*mbtK* strain. Further work is required to determine if constitutive expression

of *eis* or *papA5* restore more that the PDIM/PGL deficiency to the Δ*mbtK* strain.

In this study, we uncovered a link between Eis and PapA5. Eis is a bifunctional N-acetyl-

transferase that acetylates aminoglycoside antibiotics to inactivate them. Mutations that lead to

eis overexpression naturally occur in clinical isolates of *M. tuberculosis,* leading to kanamycin

resistance (28, 66-71). Numerous studies have reported inhibitors of Eis as a potential path to

prevent aminoglycoside resistance during *M. tuberculosis* infection (81-90). Eis reversibly acety-

lates lysine residues in the HupB protein, affecting chromosomal condensation in *M. tuberculosis*

(26). We showed that Eis N-terminally acetylates PapA5 *in vitro*, adding another potential activity

to this enzyme. We were unable to detect the N-terminal peptide of PapA5 in our proteomics

experiment. However, expression of *papA5* suppressed several phenotypes of the Δ*mbtK* strain

supporting this connection (Fig. 5). If Eis acetylates the N-terminus of PapA5 *in vivo*, this modifi-

cation must promote PapA5 activity, allowing suppression of the PDIM/PGL defect in the Δ*mbtK*

strain. Our data further support an emerging literature that NATs, including Eis, are multifunctional

(91).

Knobloch et al previously suggested that the orphan *mbtK* gene (*MMAR_1587*) was the

M. marinum ortholog of *M. tuberculosis* MbtK (92). Despite being located at the *mbt* locus,

MMAR_3692 was not mentioned in this publication. In our study, we found that deletion of the

MMAR_3692 gene resulted in reduced growth during standard culturing conditions. The growth

defect was significant, but unlike in *M. tuberculosis,* the deletion of *mbtK* did not abrogate growth.

Similar to *M. tuberculosis,* supplementation with mycobactin restored growth of the Δ*mbtK* strain

(55). Deletion of the *mbtK* gene, but not the *MMAR_1587* gene resulted in a measurable

PDIM/PGL defect. Likewise, expression of *MMAR_1587* in the Δ*mbtK* strain did not restore PDIM

levels. From our studies, we propose that the *MMAR_3692* gene is the *M. marinum* ortholog of

the *M. tuberculosis mbtK*. Our data also suggests that the two genes in *M. marinum* are function-

ally distinct.

In non-tubercular mycobacterial species, there are two functionally distinct eis genes. In

M. abscessus, these are annotated as Eis1Mab and Eis2mab (93). Overexpression of *eis2mab,*

but not *eis1mab,* resulted in resistance to Kanamycin. We likewise found that there are two po-

tential *eis* genes in *M. marinum.* Deletion of either *eis* gene attenuated *M. marinum* in a macro-

phage model of infection (Fig. S6C). Overexpression of *eis,* but not *eis1* in *M. marinum* resulted

in low level Kan resistance (Fig S10), supporting that robust literature linking Eis levels to amino-

glycoside resistance, and that the two *eis* genes are functionally distinct. Our macrophage infec-

tion data supports that overexpression of *eis1* restored cytolysis of the Δ*mbtK* strain to greater

than WT levels (Fig. S6). Yet, expression of *eis1* did not restore PDIM/PGL production to the

Δ*mbtK* strain. The mechanisms connecting Eis1 to MbtK need further experimentation.

Overall, this study revealed a new connection between iron scavenging and PDIM and

PGL biosynthesis that requires further investigation. We uncovered that the Eis NAT, which has

been widely linked to drug resistance in *Mycobacterium,* also has the potential to N-terminally

acetylate proteins, including those required for the production of the PDIM and PGL lipids. Our

data support that NATs cannot be characterized by the targets they modify, and likely play unex-

pected, interconnected and as of yet undefined roles in mycobacterial pathogenesis and physiol-

ogy.

Methods: A summary of the methods are found here. Where necessary, detailed methods are provided in the Supplementary Information. *Mycobacterium marinum* strains were derived from the *Mycobacterium marinum* M strain (ATCC BAA-535), and maintained as described previously. In frame deletion strains were generated using allelic exchange, and can be found in TableS1. Complementation plasmids, found in Table S2, were constructed using the primers in Table S3 using FastCloning. Site directed mutagenesis was performed as described previously (94). Hemolysis assays were performed as described. RAW 264.7 cells were maintained as described and infected with *M. marinum* at an MOI of 5. Macrophage cytotoxicity were measured by measuring Ethidium homodimer uptake as described. *G. mellonella* larval infections were performed by injecting *M. marinum* at a concentration $1\times10^7/5\mu L$. Thin-layer chromatography experiments was performed as described in (46). Expression and purification of 6xHis-tagged proteins were expressed in *E. coli,* and purified using Ni-NTA resin, followed by removal of the epitope tag using TEV. Proteins were used to test activity in vitro against N-terminal peptides (Genscript) using 5,5-dithio-bis-(2-nitrobenzoic acid) (DTNB) assays as (73) with minor modifications. Peptide acetylation following the DTNB assay was confirmed using MALDI. Bottom-up proteomics was performed after labelling samples with deuterated acetyl groups as (16) to quantify protein levels and N-terminal acetylation. Label-free quantification (LFQ) was used for differential protein expression analysis, and data were processed using R (v4.3.2). The code can be found on github (https://github.com/Champion-Lab/3692). Rawdata is available through MassIVE and The Proteome Exchange at ftp://MSV000095200@massive.ucsd.edu with Identifier MSV000095200 PXD053528.

Acknowledgements We would like to thank Bill Boggess at the Notre Dame Mass Spectrometry and Proteomics Facility for continued assistance. Research reported in this publication was supported by NIAID of the National Institutes of Health under award number AI106872. BSJ was supported by a Graduate Fellowship from the Eck Institute for Global Health at the University of

414   Notre Dame. We thank Dr. Micah Ferrell, who constructed several NAT deletion strains used in

415   Figure S1, during his training at Notre Dame.

416
Figure Legends

Figure 1. At least four Lysine acetyltransferases are required for WT levels of hemolytic

activity in *M. marinum.* A. Hemolytic activity of *M. marinum* strains lacking genes encoding pre-

dicted NATs. The putative/ known KATs are not filled. -, + refers to absence or presence of com-

plementation plasmid, which gene of interest expressed from the mycobacterial optimal promoter.

Significance relative to the WT strain; Brackets report significance between the deletion and com-

plemented strains. **** $P<0.0001$, *** $P=0.0004$, 0.0002, * $P=0.0250$, 0.0298, ** $P=0.0032$ **B.** TLC

analysis of total lipids from the indicated *M. marinum* strains. 6 µl of total lipids were analyzed in

each experiment. The solvents used to visualize the indicated lipid species are indicated. The

results are representative of at least three biological replicates.

Figure 2. Overexpression of Eis functionally restores PDIM production in the Δ*mbtK* strain

A. TLC analysis of total lipids from the indicated *M. marinum* strains. The NAT genes in lanes 4-

8 were constitutively expressed in the ΔmbtK strain using the mycobacterial optimal strain. 6 µl

of total lipids was separated for each strain. The indicated solvents and lipids visualized are indi-

cated. The TLC is representative of at least three biological replicates. **B.** Overlay of the predicted

structures for MbtK and Eis NATs. The PDB files used were: https://alphafold.ebi.ac.uk/files/AF-

B2HMD6-F1-model_v3.pdb and https://alphafold.ebi.ac.uk/files/AF-B2HM88-F1-model_v3.pdb.

The structures were overlaid using PyMOL v.2.4.0. **C**. % amino acid identify of the NATs from

panel A. Protein sequences were obtained from Mycobrowser (65), and aligned using Clustal

Omega (95). **D.** Clustering by phylogenetic neighbor joining tree without distance corrections,

generated based on protein alignments generated by Clustal Omega.

Figure 3. PapA5 is N-terminally acetylated by Eis *in vitro.* A. Table of peptides from proteins in the PDIM/PGL pathways. B. DTNB assay measuring the production of TNB (412nm) upon the acetylation of the PpsA, PpsE, Mas, Pks15/1 and PapA5 peptides by **B.** WT or inactive MbtK and **C.** the RimI protein from *Salmonella typhimurium* with the S18 N-terminal peptide (ARYFRR). DTNB assays were performed in technical triplicate. The data represent the mean product formation from all three experiments. Significance was determined using a Student's t-test between the active and inactive forms of each NAT protein. B.* P=0.0417, **** P<0.0001 C. ** P=0.0096**. D.** Flow injection analysis of the N-terminal peptide of PpsE with purified MbtK protein. Left panel: peaks corresponding to the unacetylated N-terminal peptide of PpsE. Right panel: mass range corresponding to the acetylated version. **E.** Flow injection of the N-terminal peptide of PapA5 incubated with either purified Eis protein or purified MbtK protein. Top left panel: peaks corresponding to unacetylated PapA5. Top right: the acetylated version of PapA5 when incubated with Eis. Bottom left panel: peaks corresponding to unacetylated PapA5, Bottom right panel: the same mass range corresponding to the acetylated version of PapA5. **F.** TLC on total lipids from *M. marinum.* 6µl total lipid loaded into each lane, PDIM run 3x consecutively in the solvent indicated, PGLs run once in the solvent indicated. PDIM lipids visualized by phosphomolybdic acid and subsequent charring, PGL lipids visualized by 1% a-naphthol and subsequent charring. The TLC is representative of 3 biological replicates, the remaining 2 are shown in Fig. S9.

Figure 4. The requirement of MbtK for mycobacterial pathogenesis can be suppressed by constitutive expression of Eis or PapA5. A. RAW 264.7 cells infected with *M. marinum* strains at an MOI of 5. The data is the average of at least 3 independent infections, each performed in biological triplicate. **** P<0.0001, *** P=0.0001, ** P=0.0218, #### P<0.0001. **B.** *Galleria mellonella* infected with the *M. marinum* strains indicated. Larvae were injected with 1x10[7] bacteria into the right hindmost proleg. Larvae were infected in 3 groups of 10, for a total of 30 larvae per strain per infection. Data is representative of at least 3 biological replicates, each performed

in technical triplicate. Statistical analysis was performed as in (79). Survival curves were com-

pared using a log-rank Mantel-Cox test, $P<0.0001$.

Figure 5. Model of Interactions between MbtK, Eis and PDIM biosynthesis. This study re-

vealed that MbtK, likely through siderophore production, is required for the production of PDIM

and PGL in *M. marinum.* This requirement was complemented in trans by the addition of Myco-

bactin J. Constitutive expression of either Eis, or PapA5 specifically suppressed the loss of

PDIM/PGL production in the absence of MbtK, suggesting that the N-terminal acetylation of

PapA5 could promote the restoration of PDIM/PGL synthesis in the Δ*mbtK* strain.

Table Legend

Table 1. Table of the *M. marinum* genes with their orthologs from *M. tuberculosis* and their

commonly used names.

M. marinum gene	*M. tuberculosis* gene	Name
MMAR_1067	Rv0730	
MMAR_1968	Rv2747	ArgA
MMAR_4519	Rv0995	RimJ
MMAR_1839	Rv2867c	Emp1
MMAR_1882	Rv2851c	ElaA
MMAR_2463	Rv1653	ArgJ
MMAR_0522	Rv0262c	Aac
MMAR_1123	Rv3420c	RimI
MMAR_4496	Rv0998	mPat
MMAR_4889	Rv0802c	
MMAR_3205	Rv2170	
MMAR_0332	Rv0133	
MMAR_1341	Rv3216	
MMAR_3740	Rv2416c	Eis
MMAR_2836	Rv3225c	
MMAR_2615		
MMAR_0151		Eis1
MMAR_3692*	Rv1347c	MbtK
MMAR_2744		

References

1. L. Favrot, J. S. Blanchard, O. Vergnolle, Bacterial GCN5-Related N-Acetyltransferases: From Resistance to Regulation. *Biochemistry* **55**, 989-1002 (2016).
2. J. Gu *et al.*, Cloning and characterization of NAD-dependent protein deacetylase (Rv1151c) from Mycobacterium tuberculosis. *Biochemistry (Mosc)* **74**, 743-748 (2009).
3. J. S. Smith *et al.*, A phylogenetically conserved NAD+-dependent protein deacetylase activity in the Sir2 protein family. *Proc Natl Acad Sci U S A* **97**, 6658-6663 (2000).
4. J. Landry *et al.*, The silencing protein SIR2 and its homologs are NAD-dependent protein deacetylases. *Proc Natl Acad Sci U S A* **97**, 5807-5811 (2000).
5. C. S. Hwang, A. Shemorry, A. Varshavsky, N-terminal acetylation of cellular proteins creates specific degradation signals. *Science* **327**, 973-977 (2010).
6. A. Drazic, L. M. Myklebust, R. Ree, T. Arnesen, The world of protein acetylation. *Biochim Biophys Acta* **1864**, 1372-1401 (2016).
7. H. Nagai *et al.*, A C-terminal translocation signal required for Dot/Icm-dependent delivery of the Legionella RalF protein to host cells. *Proc Natl Acad Sci U S A* **102**, 826-831 (2005).
8. W. M. Holmes, B. K. Mannakee, R. N. Gutenkunst, T. R. Serio, Loss of amino-terminal acetylation suppresses a prion phenotype by modulating global protein folding. *Nat Commun* **5**, 4383 (2014).
9. D. C. Scott *et al.*, Blocking an N-terminal acetylation-dependent protein interaction inhibits an E3 ligase. *Nat Chem Biol* **13**, 850-857 (2017).
10. M. Ventura *et al.*, Nuclear translocation of glyceraldehyde-3-phosphate dehydrogenase is regulated by acetylation. *Int J Biochem Cell Biol* **42**, 1672-1680 (2010).
11. R. Ree, S. Varland, T. Arnesen, Spotlight on protein N-terminal acetylation. *Exp Mol Med* **50**, 1-13 (2018).
12. R. Koch, [Tuberculosis etiology]. *Dtsch Gesundheitsw* **7**, 457-465 (1884).
13. WHO, Global Tuberculosis Report 2023. (2024).
14. T. C. Pozos, L. Ramakrishnan, New models for the study of Mycobacterium-host interactions. *Curr Opin Immunol* **16**, 499-505 (2004).
15. T. P. Stinear *et al.*, Insights from the complete genome sequence of Mycobacterium marinum on the evolution of Mycobacterium tuberculosis. *Genome Res* **18**, 729-741 (2008).
16. C. R. Thompson, M. M. Champion, P. A. Champion, Quantitative N-Terminal Footprinting of Pathogenic Mycobacteria Reveals Differential Protein Acetylation. *J Proteome Res* **17**, 3246-3258 (2018).
17. L. Xie *et al.*, Proteome-wide lysine acetylation profiling of the human pathogen Mycobacterium tuberculosis. *Int J Biochem Cell Biol* **59**, 193-202 (2015).
18. F. Liu *et al.*, Acetylome analysis reveals diverse functions of lysine acetylation in Mycobacterium tuberculosis. *Mol Cell Proteomics* **13**, 3352-3366 (2014).
19. A. Singhal *et al.*, Systematic Analysis of Mycobacterial Acylation Reveals First Example of Acylation-mediated Regulation of Enzyme Activity of a Bacterial Phosphatase. *J Biol Chem* **290**, 26218-26234 (2015).
20. X. Wu *et al.*, Differential lysine acetylation profiles of Erwinia amylovora strains revealed by proteomics. *J Proteomics* **79**, 60-71 (2013).
21. W. Lee, B. C. VanderVen, S. Walker, D. G. Russell, Novel protein acetyltransferase, Rv2170, modulates carbon and energy metabolism in Mycobacterium tuberculosis. *Scientific reports* **7**, 72 (2017).
22. K. B. Arun *et al.*, Acetylation of Isoniazid Is a Novel Mechanism of Isoniazid Resistance in Mycobacterium tuberculosis. *Antimicrob Agents Chemother* **65** (2020).
23. S. Nambi, N. Basu, S. S. Visweswariah, cAMP-regulated protein lysine acetylases in mycobacteria. *J Biol Chem* **285**, 24313-24323 (2010).

24. H. Yang *et al.*, Lysine acetylation of DosR regulates the hypoxia response of
Mycobacterium tuberculosis. *Emerg Microbes Infect* **7**, 34 (2018).
25. S. Nambi *et al.*, Cyclic AMP-dependent protein lysine acylation in mycobacteria regulates
fatty acid and propionate metabolism. *J Biol Chem* **288**, 14114-14124 (2013).
26. K. D. Green *et al.*, Acetylation by Eis and Deacetylation by Rv1151c of Mycobacterium
tuberculosis HupB: Biochemical and Structural Insight. *Biochemistry* **57**, 781-790 (2018).
27. K. H. Kim *et al.*, Mycobacterium tuberculosis Eis protein initiates suppression of host
immune responses by acetylation of DUSP16/MKP-7. *Proc Natl Acad Sci U S A* **109**,
7729-7734 (2012).
28. M. A. Zaunbrecher, R. D. Sikes, Jr., B. Metchock, T. M. Shinnick, J. E. Posey,
Overexpression of the chromosomally encoded aminoglycoside acetyltransferase eis
confers kanamycin resistance in Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A*
106, 20004-20009 (2009).
29. W. Chen, T. Biswas, V. R. Porter, O. V. Tsodikov, S. Garneau-Tsodikova, Unusual
regioversatility of acetyltransferase Eis, a cause of drug resistance in XDR-TB. *Proc Natl*
Acad Sci U S A **108**, 9804-9808 (2011).
30. D. Pathak, A. H. Bhat, V. Sapehia, J. Rai, A. Rao, Biochemical evidence for relaxed
substrate specificity of Nalpha-acetyltransferase (Rv3420c/rimI) of Mycobacterium
tuberculosis. *Scientific reports* **6**, 28892 (2016).
31. P. I. Pletnev *et al.*, Ribosomal protein S18 acetyltransferase RimI is responsible for the
acetylation of elongation factor Tu. *J Biol Chem* **298**, 101914 (2022).
32. J. A. Armstrong, P. D. Hart, Response of cultured macrophages to Mycobacterium
tuberculosis, with observations on fusion of lysosomes with phagosomes. *J Exp Med* **134**,
713-740 (1971).
33. J. A. Armstrong, P. D. Hart, Phagosome-lysosome interactions in cultured macrophages
infected with virulent tubercle bacilli. Reversal of the usual nonfusion pattern and
observations on bacterial survival. *J Exp Med* **142**, 1-16 (1975).
34. M. A. Fisher, B. B. Plikaytis, T. M. Shinnick, Microarray analysis of the Mycobacterium
tuberculosis transcriptional response to the acidic conditions found in phagosomes. *J*
Bacteriol **184**, 4025-4032 (2002).
35. K. H. Rohde, R. B. Abramovitch, D. G. Russell, Mycobacterium tuberculosis invasion of
macrophages: linking bacterial gene expression to environmental cues. *Cell host &*
microbe **2**, 352-364 (2007).
36. T. Hsu *et al.*, The primary mechanism of attenuation of bacillus Calmette-Guerin is a loss
of secreted lytic function required for invasion of lung interstitial tissue. *Proc Natl Acad Sci*
U S A **100**, 12420-12425 (2003).
37. S. A. Stanley, S. Raghavan, W. W. Hwang, J. S. Cox, Acute infection and macrophage
subversion by Mycobacterium tuberculosis require a specialized secretion system. *Proc*
Natl Acad Sci U S A **100**, 13001-13006 (2003).
38. P. S. Manzanillo, M. U. Shiloh, D. A. Portnoy, J. S. Cox, Mycobacterium Tuberculosis
Activates the DNA-Dependent Cytosolic Surveillance Pathway within Macrophages. *Cell*
host & microbe **11**, 469-480 (2012).
39. N. van der Wel *et al.*, M. tuberculosis and M. leprae translocate from the phagolysosome
to the cytosol in myeloid cells. *Cell* **129**, 1287-1298 (2007).
40. R. Simeone *et al.*, Phagosomal rupture by Mycobacterium tuberculosis results in toxicity
and host cell death. *PLoS Pathog* **8**, e1002507 (2012).
41. J. Quigley *et al.*, The Cell Wall Lipid PDIM Contributes to Phagosomal Escape and Host
Cell Exit of Mycobacterium tuberculosis. *MBio* **8** (2017).
42. M. M. Osman, A. J. Pagan, J. K. Shanahan, L. Ramakrishnan, Mycobacterium marinum
phthiocerol dimycocerosates enhance macrophage phagosomal permeabilization and
membrane damage. *PLoS one* **15**, e0233252 (2020).

43. J. Augenstreich, V. Briken, Host Cell Targets of Released Lipid and Secreted Protein
Effectors of Mycobacterium tuberculosis. *Frontiers in cellular and infection microbiology*
10, 595029 (2020).
44. B. S. Jones *et al.*, The loss of the PDIM/PGL virulence lipids causes differential secretion
of ESX-1 substrates in Mycobacterium marinum. *mSphere* **9**, e0000524 (2024).
45. A. G. Birhanu *et al.*, N(epsilon)- and O-Acetylation in Mycobacterium tuberculosis Lineage
7 and Lineage 4 Strains: Proteins Involved in Bioenergetics, Virulence, and Antimicrobial
Resistance Are Acetylated. *J Proteome Res* **16**, 4045-4059 (2017).
46. O. A. Collars *et al.*, An N-acetyltransferase required for ESAT-6 N-terminal acetylation and
virulence in Mycobacterium marinum. *mBio* **14**, e0098723 (2023).
47. J. Augenstreich *et al.*, ESX-1 and phthiocerol dimycocerosates of Mycobacterium
tuberculosis act in concert to cause phagosomal rupture and host cell apoptosis. *Cell*
Microbiol 10.1111/cmi.12726 (2017).
48. L. Y. Gao *et al.*, A mycobacterial virulence gene cluster extending RD1 is required for
cytolysis, bacterial spreading and ESAT-6 secretion. *Mol Microbiol* **53**, 1677-1693 (2004).
49. K. M. Guinn *et al.*, Individual RD1-region genes are required for export of ESAT-6/CFP-
10 and for virulence of Mycobacterium tuberculosis. *Mol Microbiol* **51**, 359-370 (2004).
50. D. G. Christensen *et al.*, Identification of Novel Protein Lysine Acetyltransferases in
Escherichia coli. *MBio* **9** (2018).
51. H. Xu, S. S. Hegde, J. S. Blanchard, Reversible acetylation and inactivation of
Mycobacterium tuberculosis acetyl-CoA synthetase is dependent on cAMP. *Biochemistry*
50, 5883-5892 (2011).
52. H. J. Lee, P. T. Lang, S. M. Fortune, C. M. Sassetti, T. Alber, Cyclic AMP regulation of
protein lysine acetylation in Mycobacterium tuberculosis. *Nat Struct Mol Biol* **19**, 811-818
(2012).
53. S. Ghosh, B. Padmanabhan, C. Anand, V. Nagaraja, Lysine acetylation of the
Mycobacterium tuberculosis HU protein modulates its DNA binding and genome
organization. *Mol Microbiol* **100**, 577-588 (2016).
54. O. Vergnolle, H. Xu, J. S. Blanchard, Mechanism and regulation of mycobactin fatty acyl-
AMP ligase FadD33. *J Biol Chem* **288**, 28116-28125 (2013).
55. C. A. Madigan *et al.*, Lipidomic analysis links mycobactin synthase K to iron uptake and
virulence in M. tuberculosis. *PLoS Pathog* **11**, e1004792 (2015).
56. R. Krithika *et al.*, A genetic locus required for iron acquisition in Mycobacterium
tuberculosis. *Proc Natl Acad Sci U S A* **103**, 2069-2074 (2006).
57. P. Brodin *et al.*, Dissection of ESAT-6 system 1 of Mycobacterium tuberculosis and impact
on immunogenicity and virulence. *Infect Immun* **74**, 88-98 (2006).
58. J. Yu *et al.*, Both phthiocerol dimycocerosates and phenolic glycolipids are required for
virulence of Mycobacterium marinum. *Infect Immun* **80**, 1381-1389 (2012).
59. A. K. Azad, T. D. Sirakova, L. M. Rogers, P. E. Kolattukudy, Targeted replacement of the
mycocerosic acid synthase gene in Mycobacterium bovis BCG produces a mutant that
lacks mycosides. *Proc Natl Acad Sci U S A* **93**, 4787-4792 (1996).
60. A. K. Azad, T. D. Sirakova, N. D. Fernandes, P. E. Kolattukudy, Gene knockout reveals a
novel gene cluster for the synthesis of a class of cell wall lipids unique to pathogenic
mycobacteria. *J Biol Chem* **272**, 16741-16745 (1997).
61. O. A. Trivedi *et al.*, Dissecting the mechanism and assembly of a complex virulence
mycobacterial lipid. *Mol Cell* **17**, 631-643 (2005).
62. L. R. Camacho *et al.*, Analysis of the phthiocerol dimycocerosate locus of Mycobacterium
tuberculosis. Evidence that this lipid is involved in the cell wall permeability barrier. *J Biol*
Chem **276**, 19845-19854 (2001).
63. M. Mathur, P. E. Kolattukudy, Molecular cloning and sequencing of the gene for
mycocerosic acid synthase, a novel fatty acid elongating multifunctional enzyme, from

Mycobacterium tuberculosis var. bovis Bacillus Calmette-Guerin. *J Biol Chem* **267**, 19388-19395 (1992).

64. D. D. Hu, S. D. Weaver, O. A. Collars, P. A. Champion, M. M. Champion, N-terminal proteomics of Mycobacterium marinum using bottom-up label-free quantitative analysis in data-dependent acquisition mode on a timsTOF Pro mass spectrometer. *Microbiol Resour Announc* **13**, e0126323 (2024).

65. A. Kapopoulou, J. M. Lew, S. T. Cole, The MycoBrowser portal: a comprehensive and manually annotated resource for mycobacterial genomes. *Tuberculosis (Edinburgh, Scotland)* **91**, 8-13 (2011).

66. S. D. Ley *et al.*, Melting the eis: Nondetection of Kanamycin Resistance Markers by Routine Diagnostic Tests and Identification of New eis Promoter Variants. *Antimicrob Agents Chemother* **65**, e0250220 (2021).

67. M. B. Gikalo, E. Y. Nosova, L. Y. Krylova, A. M. Moroz, The role of eis mutations in the development of kanamycin resistance in Mycobacterium tuberculosis isolates from the Moscow region. *J Antimicrob Chemother* **67**, 2107-2109 (2012).

68. A. Sowajassatakul, T. Prammananan, A. Chaiprasert, S. Phunpruch, Overexpression of eis without a mutation in promoter region of amikacin- and kanamycin-resistant Mycobacterium tuberculosis clinical strain. *Ann Clin Microbiol Antimicrob* **17**, 33 (2018).

69. A. Sowajassatakul, T. Prammananan, A. Chaiprasert, S. Phunpruch, Molecular characterization of amikacin, kanamycin and capreomycin resistance in M/XDR-TB strains isolated in Thailand. *BMC Microbiol* **14**, 165 (2014).

70. P. Kambli *et al.*, Correlating rrs and eis promoter mutations in clinical isolates of Mycobacterium tuberculosis with phenotypic susceptibility levels to the second-line injectables. *Int J Mycobacteriol* **5**, 1-6 (2016).

71. J. Carter, C. R. Consortium, Quantitative measurement of antibiotic resistance in Mycobacterium tuberculosis reveals genetic determinants of resistance and susceptibility in a target gene approach. *Res Sq* 10.21203/rs.3.rs-3378915/v1 (2023).

72. M. W. Vetting, D. C. Bareich, M. Yu, J. S. Blanchard, Crystal structure of RimI from Salmonella typhimurium LT2, the GNAT responsible for N(alpha)-acetylation of ribosomal protein S18. *Protein Sci* **17**, 1781-1790 (2008).

73. H. Foyn, P. R. Thompson, T. Arnesen, DTNB-Based Quantification of In Vitro Enzymatic N-Terminal Acetyltransferase Activity. *Methods in molecular biology (Clifton, N.J* **1574**, 9-15 (2017).

74. A. Yoshikawa, S. Isono, A. Sheback, K. Isono, Cloning and nucleotide sequencing of the genes rimI and rimJ which encode enzymes acetylating ribosomal proteins S18 and S5 of Escherichia coli K12. *Mol Gen Genet* **209**, 481-488 (1987).

75. E. Rwegasila, E. B. Mubofu, S. S. Nyandoro, P. Erasto, J. J. Munissi, Preparation, Characterization and in Vivo Antimycobacterial Studies of Panchovillin-Chitosan Nanocomposites. *Int J Mol Sci* **17** (2016).

76. M. Meir, T. Grosfeld, D. Barkan, Establishment and Validation of Galleria mellonella as a Novel Model Organism To Study Mycobacterium abscessus Infection, Pathogenesis, and Treatment. *Antimicrob Agents Chemother* **62** (2018).

77. Y. Li *et al.*, Galleria mellonella - a novel infection model for the Mycobacterium tuberculosis complex. *Virulence* **9**, 1126-1137 (2018).

78. F. M. Entwistle, P. J. Coote, Evaluation of greater wax moth larvae, Galleria mellonella, as a novel in vivo model for non-tuberculosis Mycobacteria infections and antibiotic treatments. *J Med Microbiol* **67**, 585-597 (2018).

79. M. Asai *et al.*, Galleria mellonella as an infection model for the virulent Mycobacterium tuberculosis H37Rv. *Virulence* **13**, 1543-1557 (2022).

80. S. S. Chavadi *et al.*, The mycobacterial acyltransferase PapA5 is required for biosynthesis
of cell wall-associated phenolic glycolipids. *Microbiology (Reading)* **158**, 1379-1387
(2012).
81. P. Santos-Junior *et al.*, A consensus reverse docking approach for identification of a
competitive inhibitor of acetyltransferase enhanced intracellular survival protein from
Mycobacterium tuberculosis. *Bioorg Med Chem* **108**, 117774 (2024).
82. L. Radhakrishnan, R. Dani, I. Navabshan, S. Jamal, N. Ahmed, Targeting Aminoglycoside
Acetyltransferase Activity of Mycobacterium tuberculosis (H37Rv) Derived Eis (Enhanced
Intracellular Survival) Protein with Quercetin. *Protein J* **43**, 12-23 (2024).
83. A. H. Pang, K. D. Green, O. V. Tsodikov, S. Garneau-Tsodikova, Discovery and
development of inhibitors of acetyltransferase Eis to combat Mycobacterium tuberculosis.
Methods Enzymol **690**, 369-396 (2023).
84. S. P. Swain *et al.*, In silico studies of alkaloids and their derivatives against N-
acetyltransferase EIS protein from Mycobacterium tuberculosis. *J Biomol Struct Dyn*
10.1080/07391102.2023.2259487, 1-15 (2023).
85. A. H. Pang *et al.*, Discovery and Mechanistic Analysis of Structurally Diverse Inhibitors of
Acetyltransferase Eis among FDA-Approved Drugs. *Biochemistry* **62**, 710-721 (2023).
86. A. H. Pang *et al.*, Discovery of substituted benzyloxy-benzylamine inhibitors of
acetyltransferase Eis and their anti-mycobacterial activity. *Eur J Med Chem* **242**, 114698
(2022).
87. A. Punetha *et al.*, Structure-based design of haloperidol analogues as inhibitors of
acetyltransferase Eis from Mycobacterium tuberculosis to overcome kanamycin
resistance. *RSC Med Chem* **12**, 1894-1909 (2021).
88. A. Punetha *et al.*, Structure-Guided Optimization of Inhibitors of Acetyltransferase Eis from
Mycobacterium tuberculosis. *ACS chemical biology* **15**, 1581-1594 (2020).
89. K. D. Green, A. Punetha, C. Hou, S. Garneau-Tsodikova, O. V. Tsodikov, Probing the
Robustness of Inhibitors of Tuberculosis Aminoglycoside Resistance Enzyme Eis by
Mutagenesis. *ACS Infect Dis* **5**, 1772-1778 (2019).
90. H. X. Ngo *et al.*, Potent 1,2,4-Triazino[5,6 b]indole-3-thioether Inhibitors of the Kanamycin
Resistance Enzyme Eis from Mycobacterium tuberculosis. *ACS Infect Dis* **4**, 1030-1040
(2018).
91. F. Sanz-Garcia *et al.*, Mycobacterial Aminoglycoside Acetyltransferases: A Little of Drug
Resistance, and a Lot of Other Roles. *Frontiers in microbiology* **10**, 46 (2019).
92. P. Knobloch *et al.*, Mycobacterium marinum produces distinct mycobactin and
carboxymycobactin siderophores to promote growth in broth and phagocytes. *Cell
Microbiol* **22**, e13163 (2020).
93. K. L. Ung, L. Kremer, M. Blaise, Structural analysis of the N-acetyltransferase Eis1 from
Mycobacterium abscessus reveals the molecular determinants of its incapacity to modify
aminoglycosides. *Proteins* **89**, 94-106 (2021).
94. P. A. Champion, S. A. Stanley, M. M. Champion, E. J. Brown, J. S. Cox, C-terminal signal
sequence promotes virulence factor secretion in Mycobacterium tuberculosis. *Science*
313, 1632-1636 (2006).
95. F. Sievers *et al.*, Fast, scalable generation of high-quality protein multiple sequence
alignments using Clustal Omega. *Molecular systems biology* **7**, 539 (2011).

732   Figure 1.

733
734
735

736   Figure 2.

739   Figure 3.

740
741

742   Figure 4.

A

B

743
744
745

746   Figure 5.
747

748