Natural variation in starvation sensitivity maps to a point mutation in
phospholipase IPLA2-VIA in *Drosophila melanogaster*

Shawn B. Williams [*†], Brian Franklin[†‡], Faye A. Lemieux[†], David M Rand[†]

[*]Department of Molecular Biology, Cell Biology and Biochemistry, Brown University,
Providence, RI 02906, USA

[†]Department of Ecology and Evolutionary Biology, Brown University, Providence, RI
02906, USA

[‡]Current address: University of Pennsylvania Perelman School of Medicine,

Philadelphia, PA 19104, USA

Correspondence:

David M Rand, PhD

Brown University

Box G-W

80 Waterman Street

Providence, RI 02912

Phone: (401) 863-2890

Email: david_rand@brown.edu

Abstract

Resistance to starvation is a classic complex trait, where genetic and environmental variables can greatly modify an animal's ability to survive without nutrients. In this study, we investigate the genetic basis of starvation resistance using complementary quantitative and classical genetic mapping in *Drosophila melanogaster*. Using the Drosophila Genetics Reference Panel (DGRP) as a starting point, we queried the genetic basis of starvation sensitivity in one of the most sensitive DGRP lines. We localize a major effect locus modifying starvation resistance to the phospholipase iPLA2-VIA. This finding is consistent with the work of others which demonstrate the importance of lipid regulation in starvation stress. Furthermore, we demonstrate that iPLA2-VIA plays a role in the maintenance of sugar reserves post-starvation, which highlights a key dynamic between lipid remodeling, sugar metabolism and resistance to starvation stress.

Introduction

Coping with periods of low nutrient availability is essential to the survival of any animal. At the simplest level, several metabolic processes need to integrate nutritional signals to successfully cope with periods of starvation. An animal has to be able to convert the energy obtained from its diet into a usable form. Energy that is not needed immediately has to be converted into a form amenable for efficient storage, and that stored energy has to be accessible for utilization during periods of starvation. For these reasons, the genetics of starvation resistance is an attractive trait for understanding the integrative aspects of animal physiology (Brown et al. 2019). Furthermore, a better understanding of the genetic basis of nutrient catabolism and storage will also shed a light on genes relevant to metabolic diseases and fitness traits in nature (Hardy et al. 2018).

Starvation resistance is a classic complex (or quantitative) trait (Harbison et al. 2004; Mackay et al. 2009). Complex traits refer to continuously distributed phenotypes across individuals within a population. These traits are believed to result from complex integration of genetic and environmental factors. While many contend that these phenotypes are primarily due to a multitude of relatively common variants with small to modest effect sizes(Lango Allen et al. 2010), others suggest that complex traits result from a smaller number of low frequency, but highly penetrant variants (Frazer et al. 2009). What remains clear is that effectively linking genetic variants to phenotypes is a central challenge of modern biology. A forward-genetic approach to establishing these associations would be to interrogate the role(s) of genes on phenotypes by assessing the effect of introducing mutations in specific genotypes. At a population level, genome wide association studies (GWAS) have contributed significantly to our understanding of the genetic basis of

complex traits (Mackay et al. 2012). However, the rapidly declining cost of next generation

sequencing technologies has made multiple tools available to associate genetic markers

with factors underlying complex traits. A variety of methods have been used to study

starvation resistance in *Drosophila melanogaster.*

The Drosophila Genetics Reference Panel (DGRP), is a collection of 205 fully sequenced

inbred *Drosophila* lines which has been used as a resource to study many complex traits.

Starvation resistance shows large variation across individual lines of the DGRP consistent

with a typical quantitative trait. Genome wide association analyses have identified

hundreds of statistically significant associations between nucleotide variants and

starvation resistance (Mackay et al. 2012). Studies in the DGRP have revealed that for

starvation resistance, the effect size of variants has an inverse relationship with minor

allele frequency (MAF) (Mackay et al. 2012). Interestingly, rarer variants appear to have

a large degree of pleiotropy, with different variants having opposing effects on various

complex traits (Mackay and Huang 2018). This is noteworthy as rare variants (Minor Allele

Frequency (MAF) <0.05) are often ignored in individual variant-based association tests.

One can therefore imagine a case where a variant present in only a handful of lines, with

a large effect size, would not be identified since the rare variant would not be included in

the study. As GWAS sample sizes have exceeded 10^5 in some cases, the definition of

rare alleles and their effect sizes has changed (Smith, et al. 2022)

The use of outbred advanced intercross populations (AIPs) is a useful tool to explore the

possibility that nucleotide variants have different effects on phenotype in different genetic

backgrounds and at different frequencies in mapping populations. A small set of parental

lines can be used to generate many genetically distinct individuals through recombination

over multiple generations. Variants that are rare in the source population will then be

present at appreciable levels in the recombinant individuals, allowing one to look for

associations in this generated population.

Whole-genome sequencing of pools of individuals (Pool-seq) allows for efficient estimates

of allele frequency differences across populations. This method can be particularly useful

when combined with extreme QTL mapping (Pool 2016; Mackay and Huang 2018). Since

causal alleles are expected to be enriched in the individuals at the tails of a given

phenotypic distribution, sequencing pools of individuals from these tails can be used to

identify alleles whose frequency differ significantly from each tail as candidate causal

alleles.

In the current study, we sought to use a combination of quantitative genetic and classical

genetic mapping to examine the genetic basis of starvation. Combining pool-seq of AIPs

with more classical approaches offers the best of both worlds, rapid and cost- effective

identification of genetic differences across populations, and finer resolution mapping to

distinguish between effect loci and linked mutations. Here, we present a study where

existing variation in the starvation resistance of two DGRP lines was used to identify the

genetic basis of starvation sensitivity in a specific sensitive line (DGRP-765). We created

replicate advanced intercross populations derived from the two DGRP lines. Pool-seq and

extreme QTL analyses revealed a polygenic basis for the difference between starvation

sensitive and starvation resistant flies, with a major factor on chromosome 3L and other

factors on 3R, 2L, but not X and 2R. Using classical mapping approaches and gene-

specific deficiencies, we were able localize the major effect locus to a single gene, the

phospholipase iPLA2-VIA. The correspondence of these approaches contributes to our

understanding of the genetic architecture of starvation resistance and identifies a novel

mutation in iPLA2-VIA that uncovers an association between starvation sensitivity, lipid

metabolism and carbohydrate metabolism.

Materials and Methods

Fly stocks

The fly lines selected for this study were based on our previously published work

demonstrating epistatic interactions between mitochondrial haplotypes and nuclear

genotypes from the DGRP that modify development time (Mossman et al. 2016). Briefly,

a panel of 72 mitochondria-nuclear combinations (using the notation: mitochondrial DNA

haplotype; nuclear DNA haplotype) was created through introgression 6 mitochondrial

haplotypes (3 *Drosophila melanogaster* (*Dmel*) and 3 *Drosophila simulans* (*Dsim*)), into

12 different DGRP lines via balancer chromosome substitution. We screened 12 of

these 72 lines (2 mtDNAs from *Dmel* and 2 mtDNAs from *Dsim* on each of three DGRP

nuclear backgrounds: 765, 315 and 820) of these lines to identify mito-nuclear

interactions which modify starvation resistance. The 765 nuclear background was

selected because it was the nuclear background most sensitive to starvation resistance

included in our panel. The 315 nuclear background was selected because the *Dsim*

mtDNAs were distinct from the *Dmel* mtDNAs in development time, demonstrating

species-specific mtDNA effects. The 820 nuclear background was selected because it

was the line most resistant to starvation included in our panel and demonstrated mtDNA

haplotype-specific effects on development time (Mossman et al 2016). During our

screen, we found one unique combination, *sil;765* which appeared to significantly

increase resistance to starvation and proceeded to validate whether the increase in

starvation resistance was due to the introgression of the *sil* mitochondrial haplotype or

residual heterozygosity from the creation of the lines.

Stock Name	Mitochondrial haplotype	Nuclear haplotype	Phenotype
Zim53;765	Zim53	DGRP-765	Starvation sensitive
line_765-hybrid (*sil;765*)	sil	DGRP-765/Unknown	Starvation resistant
Zim53;315	Zim53	DGRP-315	Starvation resistant
DGRP-765	Not altered	DGRP-765	Starvation sensitive
iPLA2-VIA[Delta174]([Lin et al. 2018](#))	Not altered	Dmel\P{EPgy2}iPLA2-VIA[EY05103]	Starvation sensitive

Table of main stocks used in this study. A table of all stocks used in this study can

be found in supplemental file 1.

Starvation Stress

Survival during starvation stress was measured by placing approximately 125 female flies

per genotype in each of three replicate demography cages. Flies were collected and

sorted over CO_2 and allowed to recover in quart-sized plastic demography cages affixed

with 25x95 mm vials of high sugar food (15% sucrose, 5% Yeast, 2% agar) for 7 days

unless otherwise indicated. After 7 days, any dead flies were removed, and the food

supplied to each demography cage is replaced with 2% Difco Bacto-agar prepared with

deionized H_2O. The number of dead flies was recorded every 12 hours and removed from

each demography cage until all flies were dead. The survival data from each cage were

modeled using a Kaplan-Meir survival model implemented in R (Therneau and Grambsch

2000). Significant differences in survivals were determined using a log rank test.

Whole genome sequencing of sensitive and resistant pools from Advanced

Intercross Populations

Population Cages. Five replicate advanced intercross populations (AIPs) were

established between the sensitive (*Zim53;765*) and resistant (*Zim53;315*) lines, as

follows: 25 *Zim53;765* males were crossed to 25 *Zim53;315* virgin females, in addition to

the reciprocal cross, to yield a total number of 50 male and 50 female parents of each line

as the founders of each replicate cage. These parents were introduced into Plexiglas

cages (20 x 20 x 20cm) and allowed to lay eggs in four 125mL culture bottles with 30mL

of Drosophila medium in each bottle. The four bottles in each cage were left with their

tops open for 3 days, and then each bottle was capped to allow flies to develop. Prior to

eclosion, the caps were removed from each bottle, and all flies were allowed to emerge

into the cage and mate at random. Four new culture bottles were placed into each

replicate cage, and the adults were allowed to lay eggs for 48 hours. After egg laying,

adults were cleared from each bottle, plugs were inserted, and cultures were allowed to

develop for 7 days following the termination of egg laying, after which the plugs were

removed from all bottles to allow eclosion of the next generation into each cage.

Population sizes exceeded 1000 individuals after the first generation. All five cages were

maintained at large population sizes (>2000 adults) for 10 generations to allow

recombination to take place between the sensitive (765) and resistant (315) DGRP

nuclear genotypes.

Starvation Stress. The female flies from each generation-10 population cage were

collected over CO_2 and moved to demography cages with a density of approximately 150

flies per cage. The flies were allowed to recover for 7 days with access to high sugar food

and then subjected to starvation conditions by replacing the food vials with a 2% agar and

water mix. Every six hours from the start of starvation, all the dead flies were collected,

flash frozen and stored at -80C.

Pool-sequencing. Pools of approximately 50 dead flies were made by combining flies from

the two earliest time points where deaths were observed (24 hour and 30 hour) and the

last two time points (96 hour and 102 hour). DNA from each replicate pool was isolated

using the DNeasy DNA purification kit from Qiagen (Cat No 69506). Each sample was

RNase treated prior to DNA elution using a 10-minute RNase A (Cat No 19101) treatment.

Libraries were prepared and sequenced using the DNBseq platform by the Beijing

Genomics Institute (BGI).

Estimation of genetic differentiation using Popoolation2. Raw reads were mapped to the

dm6 build of the *Drosophila* genome using *bwa* version 0.7.15 (Li and Durbin 2009).

Variants were called using *samtools* version 1.9 (Li et al. 2009). Allele frequency

differences and estimates of population subdivision (F_{ST}) were determined using

Popoolation2 (Kofler et al. 2011), and the significance of differences in allele frequencies

between the early-dying and late-dying sample across the five replicate populations was

determined using the Cochran-Mantel-Haenszel test. P-values were corrected using the

Bonferroni method and differences with FDR-corrected p-values < 0.05 were considered

to be significant. Linkage disequilibrium was estimated from paired end reads, following

the procedure described in ([Feder et al. 2012](#)).

Forward genetic mapping of starvation sensitivity

To generate *Zim53;765* selectively heterozygous for different autosomes, 20 *Zim53;765*

females were first crossed to 20 males from an in-house double balanced stock (*wg[Sp-*

1]/CyO; Dr[1]/TM3, Sb[1]). Twenty male progeny with both the 2nd chromosome *CyO* and

3rd chromosome *Dr* markers present were collected and sorted over CO_2 and crossed to

20 *Zim53;765* virgin females in eight replicate 20-female x 20-male crosses. Progeny from

this second cross were segregated into four groups based on the presence of the *CyO*

and *D*r markers. The four groups were, none (reconstituted wild-type *Zim53;765*),

chromosome 2 heterozygotes (*CyO/765*), chromosome 3 heterozygotes (*Dr/765*) and

heterozygotes for both chromosome 2 and 3 (*CyO/765;Dr/765*). Flies from each group

were moved to 3 replicate demography cages with approximately 125 flies each. The

survival of each group under starvation stress was assayed as described earlier.

Chromosome 3 recombination

The chromosome segregation experiments indicated that the majority of the difference in

starvation resistance between the lines mapped to chromosome 3. We attempted to

generate *Zim53;765* chromosome 3 recombinants by crossing *Zim53;765* virgin females

to 20 males from a multiply marked recessive stock (BSC_576). The marked stock

contains the *ru, h Diap, st, cu, sr, e* and *ca* markers. The progeny from this cross however

failed to eclose. As an alternative, we crossed the *siI;765* virgin females to *BSC_576*

males in an effort to allow recombination of the 765 alleles that is present in the *sil;765*

stock. Twenty F_1 virgin females were collected and crossed to 20 males from a similarly

multiply marked recessive stock with identical recessive markers in addition to a dominant

marker present on one copy of chromosome 3 (BSC_1711) in three replicate 20-female

x 20-male crosses. F_1 males from this second cross were collected and segregated based

on the recessive and dominant markers present. Five classes were initially designated

based on the visible markers present, class i*: ru, h*; class ii: *ru, h, Diap, st*; class iii: *ru, h,*

Diap, st, cu; class iv: *ru, h, Diap, st, cu, sr*; and class v: *ru, h, Diap, st, cu, sr, e*. Twenty

males from each group were crossed to 20 *Zim53;765* virgin females and allowed to mate

for 48 hours in three replicate 20-female x 20-male crosses. The males were removed

and similarly mated to *sil;765* virgin females. Females from each cross were collected

over CO_2. Sufficient flies for a starvation stress survival assay were only available from

the class i, class ii and class v crosses, so only these were considered. The survival of

these females under starvation stress was measured as described above.

Chromosome 3L Deficiency Screen

The recombination mapping implicated chromosome 3L as harboring the major starvation

factor, so deficiency mapping was pursued. Twenty males from each deficiency line

(Supplemental Table 1) were crossed to 20 *Zim53;765* virgin females in eight replicate

20-female x 20-male crosses. F_1 females were collected and sorted over CO_2. Females

were separated based on the presence of the respective balancer marker (flies

heterozygous for the deficiency and the 765 chromosome do not have the balancer

present) and moved to 3 replicate demography cages with approximately 125 flies each,

for each genotype. The flies were allowed to recover for 7 days on high sugar food and

survival under starvation stress measured as described above.

iPLA2-VIA Complementation

The chromosome 3L deficiency mapping, in conjunction with the available DGRP

sequences, identified the iPLA2-VIA as a candidate gene, and deficiencies within that

gene were obtained for further complementation mapping. Twenty

Zim53;765/BSC_25204 flies were crossed to twenty *BSC_80133/ iPLA2-VIA[Delta174]*

and independently to *BSC_80134/iPLA2-VIA[Delta192]* in reciprocal. Each cross was

done in replicates of 4. As a control, males from each *iPLA2-VIA* deficiency stock were

also crossed to *Zim53;315* females. F_1 females were collected and sorted over CO_2 and

moved to 3 replicate demography cages with approximately 125 flies each. The flies were

allowed to recover for 7 days on high sugar food and survival under starvation stress

measured as described above.

Glucose Assay

The free glucose levels were measured for *Zim53;765*, *Zim53;315*, *iPLA2-VIA[Delta174]*,

Zim53;315 crossed to *iPLA2-VIA[Delta174]* and *Zim53;765* crossed to iPLA2-

VIA[Delta174] as described in ([Tennessen et al. 2014](#)). 120 female flies from each

genotype were collected and sorted over CO_2 and allowed to recover in demography

cages affixed with 25x95 mm vials of high sugar food (15% sucrose, 5% Yeast, 2% agar)

for 5 days. Three replicates were done for each genotype. After 5 days, any dead flies

are removed. Half of the remaining flies were removed from the demography cage,

immediately flash frozen in liquid N_2 and stored at -80°C. The food supplied to each

demography cage was replaced with 2% Difco Bacto-agar prepared with deionized H_2O.

After 12 hours, dead flies were discarded and the remaining flies were removed from the

demography cage, immediately flash frozen in liquid N_2 and stored at -80°C. Six frozen

flies from each replicate were transferred to a 1.5uL micro-centrifuge tube and

homogenized in 120uL of Phosphate Buffered Saline (PBS) using a steel bead and

Qiagen TissueLyser II set to 30Hz for 2 minutes. A 10uL aliquot was removed to measure

protein content using the Pierce™ BCA Protein Assay Kit (23225). The remaining

homogenate was heat treated for 10 minutes at 70°C. The homogenate was diluted 1:4

with PBS. The free glucose was measured with the Glucose Oxidase (GO) kit (MAK097-

1KT) by following the protocol for free glucose measurement laid out in (Tennessen et al.

2014).

Data Availability

Drosophila strains, code and raw data are available at the Rand Lab GitHub site for this

publication (https://github.com/DavidRandLab/Williams-et-al.-Starvation-Genetics-2021).

The authors affirm that all analyzed data necessary for confirming the conclusions of the

article are present within the article, figures, tables and GitHub site. Raw sequences for

the pool-seq mapping are available at BioProject accession number: PRJNA1130456.

Results
Selection of Genotypes for this study

Based on our interest in the genetics of starvation resistance, we designed a screen to

examine the effect of gene x gene and gene x environment interactions on starvation

resistance. We created a panel of unique mito-nuclear introgression lines based on

introgression of six different mitochondrial DNA haplotypes into several individual fly

lines from the DGRP as previously described (Mossman et al. 2016). Using these

introgression lines, we examined how different mito-nuclear introgression strains

responded to starvation stress. *mito;765* lines were of particular interest because the

765 nuclear background represents one of the least resistant lines in terms of starvation

resistance in the DGRP, and it appeared that introgression of the *siI* mitochondrial DNA

haplotype significantly improved resistance to starvation in the 765 nuclear background.

To test whether the rescue of starvation resistance mapped to a cytoplasmic or nuclear

factor, we performed reciprocal crosses to examine if maternal inheritance of mtDNA

was the source of the rescue of starvation sensitivity. We observed that when we

crossed *Zim53;765* to this *siI;765*, the direction of the cross (*Zim53;765 female* crossed

siI;765 males vs *siI;765* females to *Zim53;765 males*) made no difference to starvation

resistance of its progeny. This implied that the *siI* mtDNA haplotype was not the source

of the rescue, as would be expected if the trait was being inherited maternally

(Supplemental figure 1a). This prompted us to analyze the reads from an RNA-

sequencing data set from each fly line. We compared the genotype calls for each SNP

present in *Zim53;765* reads (restricted to the coding regions) and the genotype calls at

the same locations in *siI;765*. We observed that at the vast majority of these sites, the

siI;765 was heterozygous for the 765 allele and an alternative allele (in most cases the

reference allele) (Supplemental figure 1b and 1c). We did a similar comparison of

siI;765 to an alternate DGRP line (*Zim53;315*) and did not see a similar overlap

between the *siI;765* allele and the *Zim53;315* allele (Supplemental figure 1b and 1c).

Based on these results we concluded that the *siI;765* was largely heterozygous for 765.

This suggested to us that starvation sensitivity in *Zim53;765,* and the other 4 *mito;765*

combinations, were homozygous recessive and the improvement in resistance to

starvation seen in *siI;765* was due to its nuclear heterozygosity. Indeed, we observed

that the offspring from a *Zim53;765* to *Zim53;315* cross were significantly more resistant

to starvation than *Zim53;765* (data not shown). For the rest of the study we refer to the

siI;765 as *line_765-hybrid* to represent the true nature of its nuclear genome.

Sites on chromosome 2R and 3 have significant differences in allele frequency
between early and late dying flies.

To understand the genetic basis of 765 starvation sensitivity, we sought to identify genetic

differences characteristic of starvation sensitive flies. We created an advanced intercross

population in which *Zim53;765* was allowed to recombine with *siI;315*. We set up five

independent populations where *Zim53;765* flies were crossed to *siI;315* in reciprocal and

the populations maintained for 10 generations at population sizes >1000 adults. Following

this, females from each population were subjected to starvation stress and dead flies were

collected every six hours until all flies were dead. Flies that died during the first three and

last two six-hour intervals were pooled and sequenced (24 - 36 hours vs. 96 and 102

hours, hereafter 'early' and 'late' dying pools). Consistent differences in allele frequencies

across the early and late dying populations were determined using the Cochran-Mantel-

Haenszel test implemented in Popoolation2 (Kofler et al. 2011). There was an enrichment

in significantly different variants across chromosome 3 and a localized region of

chromosome 2R (Figure 1). These results suggest that *Zim53;765* and as a consequence

DGRP-765, starvation sensitivity is based on variants present on chromosome 3 and

chromosome 2.

Chromosome 3 heterozygosity partially suppresses *Zim53;765* starvation
sensitivity

Based on the suppression of starvation sensitivity seen in *Zim53;765* heterozygotes, we

reasoned that starvation sensitivity in *Zim53;765* was a homozygous recessive trait and

that we could localize the effect locus/loci to chromosomes by using chromosome

segregation tests and observing which chromosome segregants masked the trait. We

selectively made *Zim53;765* heterozygous for individual autosomes. Using chromosome

segregation tests, we generated the four 2-chromosome segregant populations of flies as

follows: heterozygous for chromosome 2, chromosome 3, chromosome 2 and 3 in

combination or neither. We observed that flies heterozygous for chromosome 3 had an

80% increase in mean survival relative to the non-heterozygotes (Figure 2a). Whereas

flies heterozygous for chromosomes 2 and 3 had a 140% increase in mean survival

(Figure 2a). Taken together, these results suggest that the main effect locus/loci

controlling *Zim53;765* starvation sensitivity are present on chromosome 3 with a

secondary locus/loci present on chromosome 2. Importantly, these findings are consistent

with the extreme QTL analysis described earlier for the Advanced Intercross Populations.

Heterozygosity of chromosome 3L region partially suppresses Zim53;765
starvation sensitivity
340

To localize the main effect loci on chromosome 3 we decided to conduct a deficiency

screen along chromosome 3. As there are over 200 deficiencies spanning chromosome

3, we first sought to narrow the region of interest further. We generated a series of

Zim53;765 recombinants which were heterozygous for overlapping regions of

chromosome 3. Each recombinant was generated by crossing *line_765-hybrid* virgin

females to a multiply marked recessive stock, then crossing the F_1 virgin females to males

from a similarly marked recessive stock with one dominant marker added to distinguish

progeny. Three sets of recombinants (classes i, ii, v) were selected based on the

presence of visible recessive markers and sufficient flies for starvation analyses (Figure

2b). Recombinant class i is likely heterozygous for the region of 3L from the telomere up

to at least the regions marked by *h,* with recombination breakpoints existing anywhere

between h and the *cu* gene. Recombinant class ii is heterozygous for most of 3L and up

to *cu* gene on 3R, with breakpoints existing anywhere between the *cu* gene and the *ca*

gene. Recombinant class v is likely heterozygous for most of chromosome 3 with

breakpoints existing from the *ca* gene up to the telomere of 3R. It is important to note that

because these recombinants were generated from *line_765-hybrid*, the pool of

recombinants will have both the *Zim53;765* allele and the alternate (unknown) allele

present. Males from each recombinant class were initially crossed to *Zim53;765* females,

and then the same males crossed to *line_765-hybrid* as a control. The starvation

sensitivity of the female offspring of the *Zim53;765* and recombinant males was compared

to the female offspring of the *line_765-hybrid* and recombinant male controls. While we

cannot discern genotype at the individual fly level, 765 chromosome 3 alleles will be

enriched in the progeny from the *Zim53;765* vs recombinant cross, while the alternative

will be true for the *line_765-hybrid* vs recombinant cross. We expected that flies

homozygous for the causal *Zim53;765* allele would die earlier during starvation stress,

thus revealing the region of *Zim53;765* chromosome 3 responsible for its starvation

sensitivity. Although the heterogeneity of respective offspring populations will mask this

effect, the enrichment of the 765 allele in the *Zim53;765* vs recombinant cross population

and the alternate allele in the *line_765-hybrid* vs recombinant cross population should

result in appreciable differences in deaths early on during starvation stress. Indeed, we

observed a significant difference between the starvation sensitivity of class i offspring, but

not in the case of the cross ii offspring; class v offspring showed a small but significant

difference (see Figure 2C). These results indicate that heterozygosity of a region between

the *h* marker of recombinant i and the *cu* marker of recombinant ii is likely the largest-

effect factor required to suppress *Zim53;765* starvation sensitivity.

Hemizygosity of 3L: 9812381 – 9899255 prevents partial suppression of *Zim53;765*
376
377 **starvation sensitivity**
378
We employed a deficiency screen across the region between recombinant class i and

recombinant class ii markers. The objective of this screen was to identify the region(s) of

heterozygosity required for the suppression of *Zim53;765* starvation sensitivity.

Zim53;765 females were crossed to a series of flies deficient for various parts of

chromosome 3L. The deficient region of each line is maintained over a balancer

chromosome so the resulting offspring can be positive for the balancer, indicating

complete heterozygosity of chromosome 3 or only partially heterozygous for chromosome

3 and hemizygous for the deficient region. Significant differences in the starvation

resistance of the two classes of offspring indicates that heterozygosity of the missing

deficiency region is required for the suppression of starvation sensitivity. We initially

screened 10 deficiency lines and identified three where the starvation sensitivity was not

suppressed (failed to complement) in the partial chromosome 3L hemizygote (Figure 3a).

All three lines had deficient regions between 3L:9812381 – 3L:9899255 (Figure 3b),

indicating that heterozygosity of parts of the region is required for the suppression of

Zim53;765 starvation sensitivity.

Mutation in iPLA2-VIA is a candidate cause of *Zim53;765* starvation sensitivity

Next, we sought to identify *Zim53;765* variants in the chromosome 3L region required for

suppression of *Zim53;765* starvation sensitivity which could underlie its sensitivity to

starvation. We utilized the available genomic sequencing data of each DGRP line to

identify 765 variants (Mackay et al. 2012). As the causal variant is deleterious, we focused

our initial analysis on low frequency variants (present in less than 10 % in the DGRP

population). We identified 83 variants which fit our criteria (Figure 4a), of these only 3

were predicted to lead to non-synonymous changes (Figure 4b and Table 1). A C-to-A

mutation, predicted to result in a G-to-V amino acid residue change in the conserved

GXGXXG motif of the phospholipase iPLA2-VIA protein, appeared as the most likely

candidate. Mutations in iPLA2-VIA have previously been shown to cause starvation

sensitivity and reduced lifespan in *Drosophila* (Kinghorn et al. 2015; Kinghorn and

Castillo-Quan 2016; Lin et al. 2018). In addition, PolyPhen-2 predicted the mutation as

probably damaging, with a score of 0.999 out of 1 (Adzhubei et al. 2010). Based on this

analysis, we decided to determine if *Zim53;765* has a functional copy of the iPLA2-VIA

gene.

iPLA2-VIA[Del174] fail to complement Zim53;765/ DGRP-765 starvation sensitivity
and glucose homeostasis
To determine if *Zim53;765* and as a consequence DGRP-765, results from a defective

copy of iPLA2-VIA, we sought to determine if the 765 copy of iPLA2-VIA gene could

complement an iPLA2-VIA null. We crossed *Zim53;765* virgin females to two previously

characterized iPLA2-VIA null lines (*iPLA2-VIA[Del174]*, *iPLA2-VIA[Del192]*, caused by 2

different deletions generated by imprecise excision of a mobile element) (Lin et al. 2018).

To rule out the potential effects of mtDNA haplotype on starvation resistance, we also

utilized the DGRP-765 strain obtained from the Bloomington Stock center. We found that

progeny from DGRP-765 crossed *to iPLA2-VIA[Del174]* or *iPLA2-VIA[Del192]* were also

starvation sensitive (Figure 5). In contrast, crossing *Zim53;765*, *iPLA2-VIA[Del174]* or

iPLA2-VIA[Del192] to *Zim53;315* resulted in the suppression of starvation sensitivity

(Figure 5). Ablation or pharmaceutical inhibition of calcium independent phospholipases

have been shown to reduce insulin secretion in response to a glucose challenge (Song

et al. 2005; Ali et al. 2013). This raised the question of whether *Zim53;765* showed signs

of impaired glucose metabolism. We observed that following 12 hours of starvation,

Zim53;765 significantly depletes its available free glucose in contrast to *Zim53;315*

suggesting improper regulation of glucose during acute starvation (Figure 6).

Interestingly, this phenotype was not observed in iPLA2-VIA[Del174] (Figure 6). However,

we found that progeny from *Zim53;765* crossed to iPLA2-VIA[Del174] also significantly

deplete their available free glucose following 12 hours of starvation (Figure 6). Taken

together, these results implicate iPLA2-VIA in *Zim53;765* starvation sensitivity, with a

parallel effect on glucose homeostasis. In addition, its effects extend beyond a simple

loss of function as evidenced by the *Zim53;765* specific phenotypes observed.

Discussion

Polygenic basis of *Zim53;765* starvation sensitivity

Our advanced intercross population (AIP) design used a line at the extreme end of the

starvation resistance spectrum (*Zim53;765*, most sensitive) and another line more central

in the distribution. Though a compelling case could be made for the use a line at the

opposite end of the distribution (most resistant). We focused our effort on attempting to

identify why *Zim53;765* was starvation sensitive and reasoned that intercrossing

Zim53;765 with *Zim53;315* would result in populations with the starvation sensitivity

allele(s) and the more common allele. We sequenced pools of flies from the extreme ends

of the AIP's starvation resistance distribution and found that there were several variants

with significant differences in allele frequencies. It is likely that many of these different

variants are in linkage disequilibrium since the populations recombined over a period of

only 10 generations. Nevertheless, the clustering of significant variants on chromosome

2 and 3 are suggestive of a polygenic basis for *Zim53;765* starvation sensitivity. This is

supported by the results of the chromosome segregation analysis. There we found

chromosome 3 heterozygosity was necessary to suppress *Zim53;765* starvation

sensitivity. In addition, heterozygosity of chromosome 2 and 3 together further improved

starvation resistance. Notably, chromosome 2 heterozygosity in isolation was not

sufficient to even partially suppress *Zim53;765* starvation sensitivity. This indicates that

the chromosome 2 loci/locus acts as a modifier that genetically interacts with the

chromosome 3 locus/loci. This may provide a potential explanation for why the vast

majority of significant variants identified were localized to chromosome 3L, while there

were clusters with fewer significant variants on chromosome 2L and 3R.

Narrowing the main effect locus

We attempted to narrow the target region, by generating recombinants between

Zim53;765 and an unaffected genotype. We designed these crosses to be done with

multiple marked recessive stocks, as this would allow us to have a general idea of the

regions of recombination. We however failed to do this directly as crossing *Zim53;765*

females to males from each of the multiple marked recessive stocks failed to yield an

appreciable number of progeny. While not ideal, we decided to use the *line_765-hybrid*

to generate recombinants for a few reasons. Reciprocally crossing *line_765-hybrid* to

Zim53;765 revealed that starvation resistances was not maternally inherited, suggesting

that mitochondrial DNA was not responsible for the phenotypic differences between the

two. *Zim53;765* and *line_765-hybrid* were generated using the same method described

in the materials and methods section, so a priori we knew that *line_765-hybrid* was a

hybridized version of 765 nuclear chromosomes, a fact confirmed by comparing the

alleles of *Zim53;765* variants to *line_765-hybrid*'s. The preponderance of the evidence

from the deficiency screen supported heterozygosity of the region of 3L 9812381 –

9899255 being required for suppression of *Zim53;765* starvation sensitivity. Four of the

six lines deficient for the region showed the expected result where suppression of

starvation sensitivity was masked in the hemizygous progeny. However, when *Zim53;765*

was crossed with Df(3L)BSC282, there was only a modest difference in the starvation

resistance of the heterozygous and homozygous progeny (Supplemental figure 2b). In

the case of *Zim53;765* crossed with Df(3L)BSC283, there was no statistical difference in

the starvation resistance of the heterozygous and homozygous progeny. It is possible

that the predicted deficient region of Df(3L)BSC282 and Df(3L)BSC283 are not

completely deleted.

Complementation points to Zim53;765 iPLA2-VIA

Of all the variants in the 3L 9812381 – 9899255 region, the nonsynonymous mutation

present in the calcium-independent phospholipase iPLA2-VIA immediately stood out, due

to the location of the mutation in the functional domain of the enzyme, and the phenotypes

previously associated with iPLA2-VIA loss of function. Mutations in the human homolog

PLA2G6 gene are characteristic of multiple autosomal recessive disease (Guo et al.

2018). Several studies have independently characterized starvation sensitivity in iPLA2-

VIA mutants (Kinghorn et al. 2015; Iliadi et al. 2018; Lin et al. 2018; Mori et al. 2019). In

each case, targeted mutations were introduced in iPLA2-VIA to generate null models,

with the goal of better understanding its role in neurodegeneration. These studies identified aberrant lipid metabolism, lipid peroxidation and mitochondrial dysfunction as potential causes of stress sensitivity and neurodegeneration. We should not necessarily expect point mutations in iPLA2-VIA to universally have the same effect as these loss of function models, as (Engel et al. 2010) mutations in PLA2G6 have varying effects on catalytic activity. However, we can expect that if mutation(s) in iPLA2-VIA do in fact underlie *Zim53;765* starvation resistance, an iPLA2-VIA null should fail to complement *Zim53;765* starvation sensitivity. We showed that two separate nulls generated by (Lin et al. 2018) do in fact fail to complement *Zim53;765* starvation sensitivity, indicating iPLA2-VIA underlies *Zim53;765* starvation sensitivity.

Some phospholipases are have been associated with modulation of insulin secretion in response to glucose by releasing arachidonic acid (AA), and exogenous AA supplementation is sufficient to promote insulin release from rat islet cells (Metz 1988a). This insulin secretion is believed to promote phospholipid hydrolysis and accumulation of phospholipid-derived intermediates (Metz 1988b). This turnover of phospholipids could promote insulin release, creating an interesting dynamic between glucose and lipid metabolism, with phospholipases at the center. The premature depletion of glucose and glycogen in *Zim53;765* early during starvations could be one consequence of its mutated iPLA2-VIA. Consistent with this, we found that iPLA2-VIA[Del174] fail to complement the premature glucose and glycogen depletion phenotypes associated with *Zim53;765* (Figure 6). What is especially interesting is that the iPLA2-VIA null themselves do not display this phenotype. This raises the possibility that affected pathways which ultimately

513　result in starvation sensitivity differ between the deletion mutation of iPLA2-VIA[Del174]

514　and the point mutation unique to *Zim53;765*. It will be an interesting target of a future

515　study to characterize the additional molecular bases underlying the starvation sensitivity

516　of *Zim53;765,* as this could potentially highlight novel roles for iPLA2-VIA and asses the

517　effects of the *Zim53;765* specific mutations have on the function of iPLA2-VIA.

Acknowledgments

520　The authors thank the HHMI Summer Fellows Julia Dewey, Tyler Devlin, Brian Franklin,

521　Cynthia Hale-Phillips, Matthew P. McAteer, Zemplen Pataki, Chén Yé, and Denise Yoon

522　who conducted the initial screen for mito-nuclear interactions affecting starvation

523　resistance, which provided the impetus for this work.

Funding

526　Work here was supported in part by Gilliam Fellowship for Advanced Study awarded to

527　SBW and DMR and NIGMS R01 2R01GM067862 awarded to DMR. DMR

528　acknowledges support of COBRE award P20GM109035 and MIRA R35GM139607

Conflicts of Interest

531　None declared.

535
Figure 1. Significant allelic frequency differences between the tails of replicate
advanced intercross populations subjected to starvation stress.
An advanced intercross population was created by crossing *Zim53;765* and *sil;315* in
reciprocal. The hybrids were allowed to recombine freely for 10 generations before
being subjected to starvation stress. Dead flies were collected every 6 hours and dead
flies from the first 3 timepoints were combined and subjected to sequencing analysis.
The same was done with the last two timepoints. In total 5 replicates AIP populations
were generated and sequenced. (A) Consistent differences in allele frequencies across
replicates were determined using the Cochran-Mantel-Haenszel (CMH) test and
Popoolation2. The y-axis is the negative log-transformed, Bonferroni corrected p-value
of each SNP (represented by each circle). The x-axis is the position of each SNP along
the indicated chromosome. Blue and red lines highlight corrected p-values above 1x10^{-5}
and 1x10^{-6} respectively. (B) F$_{ST}$ between pools for each site averaged across replicate
populations. The location of elevated FST values is generally concordant with significant
CMH test statistics.
551

552
Figure 2. Heterozygosity of chromosome 3 partially suppresses *Zim53;765*
starvation sensitivity
(A) Survival of female *Zim53;765* flies heterozygous for no chromosome, chromosome
2, chromosome 3 or chromosome 2 and 3 following starvation stress. Differences in
survival relative to flies with no marker (none) were determined using a log-rank test
(implemented in the R survival package) and included in the supplemental relevant
statistics file.(B) Cartoon representation of recombination breakpoints for each
recombinant class generated through meiotic recombination of the chromosome 3
mapping line. (C) Survival of female offspring produced by crossing each recombinant
class to *Zim53;765* homozygotes or heterozygotes. Differences in survival were
determined using a log-rank test (implemented in the R survival package) and included
in the supplemental relevant statistics file.
565

Figure 3. Hemizygosity of 3L: 9812381 – 9899255 prevents partial suppression of *Zim53;765* **starvation sensitivity**

(A) Survival of offspring from the cross between *Zim53;765* and the respective deficient male following starvation stress. Heterozygous refers to progeny which inherited the balancer chromosome, hemizygous refers to progeny which inherited the deficient chromosome. Line numbers correspond to Bloomington Stock Center names. Differences in survival was determined using a log-rank test (implemented in the R survival package) and included in the supplemental relevant statistics file. (B) Linear representation of hemizygous regions of respective deficiency lines. Highlighted region is common to the three deficiency regions which failed to suppress *Zim53;765* starvation sensitivity.

Figure 4. Frequency of 765 variants among the Drosophila Genetics Reference Panel

(A) Allele frequency of 765 variants in the 3L candidate region among the lines of the Drosophila Genetics Reference Panel. Variants found below the red line are present in less than ten percent of the populations. (B) Analysis of the types of low frequency (minor allele frequency less that 0.1 in the Drosophila Genetics Reference Panel) variants in 3L candidate region present in 765.

588
589 **Figure 5. *Zim53;765* iPLA2-VIA fails to complement iPLA2-VIA null lines**
590 Survival of female progeny from indicated crosses following starvation stress. Panels
591 are divided by male parent and colors segregate progeny from each type of female
592 parent. Differences in survival were determined using a log-rank test (implemented in
593 the R survival package) and included in the supplemental relevant statistics file.
594

595
596

598
Figure 6. iPLA2-VIA[Delta174] fails to complement glucose and glycogen
depletion phenotype of *Zim53;765*
The free glucose and total glycogen levels were measured for *Zim53;765*, *Zim53;315*,
iPLA2-VIA[Delta174], *Zim53;315* crossed to iPLA2-VIA[Delta174] and *Zim53;765*
crossed to iPLA2-VIA[Delta174]. Values in the un-starved (control) was compared to the
values in the starved samples (12 hours) for 3 replicates of 6 flies using a student's t-
test. Significant differences are indicated by * <0.05, ** <0.01.
606
607
608

Supplemental Figure 1. *sil;765* is a hybrid of DGRP-765
(A) Survival of female offspring of *Zim53;765,* and *sil;765* crossed in reciprocal. There is
no statistical difference in survival based on a log-rank test for significance. (B and C)
Genotype of *sil;765* at homozygous variant sites in RNA sequencing reads from (C)
Zim53;765 and (D) *Zim53;315* Green indicates that *sil;765* was heterozygous for both
the variant and the reference allele, blue indicates that *sil;765* was homozygous for the
reference allele.

A

B

618
Supplemental Figure 2
(A) Linear representation of deficient region of respective deficient lines in region
identified in meiotic recombination mapping. (B) Survival of offspring from the cross
between Zim53;765 and the respective deficient male following starvation stress.
Differences in survival were determined using a log-rank test (implemented in the R
survival package) and included in the supplemental relevant statistics file. Heterozygous
refers to progeny which inherited balancer chromosome, hemizygous refers to progeny

which inherited deficient chromosome. Line numbers correspond to Bloomington Stock
Center names.

References

628
629
630
631
Adzhubei, I. A., S. Schmidt, L. Peshkin, V. E. Ramensky, A. Gerasimova *et al.*, 2010 A
method and server for predicting damaging missense mutations. Nat Methods 7**:**
248-249.
Ali, T., G. Kokotos, V. Magrioti, R. N. Bone, J. A. Mobley *et al.*, 2013 Characterization of
FKGK18 as inhibitor of group VIA Ca2+-independent phospholipase A2
(iPLA2beta): candidate drug for preventing beta-cell apoptosis and diabetes. PLoS
One 8**:** e71748.
Brown, E. B., M. E. Slocumb, M. Szuperak, A. Kerbs, A. G. Gibbs *et al.*, 2019 Starvation
resistance is associated with developmentally specified changes in sleep, feeding
and metabolic rate. J Exp Biol 222.
Engel, L. A., Z. Jing, D. E. O'Brien, M. Sun and P. T. Kotzbauer, 2010 Catalytic function
of PLA2G6 is impaired by mutations associated with infantile neuroaxonal
dystrophy but not dystonia-parkinsonism. PLoS One 5**:** e12897.
Feder, A. F., D. A. Petrov and A. O. Bergland, 2012 LDx: estimation of linkage
disequilibrium from high-throughput pooled resequencing data. PLoS One 7**:**
e48588.
Frazer, K. A., S. S. Murray, N. J. Schork and E. J. Topol, 2009 Human genetic variation
and its contribution to complex traits. Nat Rev Genet 10**:** 241-251.
Guo, Y. P., B. S. Tang and J. F. Guo, 2018 PLA2G6-Associated Neurodegeneration
(PLAN): Review of Clinical Phenotypes and Genotypes. Front Neurol 9**:** 1100.
Harbison, S. T., A. H. Yamamoto, J. J. Fanara, K. K. Norga and T. F. C. Mackay, 2004
Quantitative Trait Loci Affecting Starvation Resistance in Drosophila
melanogaster. Genetics 166**:** 1807-1823.
Hardy, C. M., M. K. Burke, L. J. Everett, M. V. Han, K. M. Lantz *et al.*, 2018 Genome-
Wide Analysis of Starvation-Selected Drosophila melanogaster-A Genetic Model
of Obesity. Mol Biol Evol 35**:** 50-65.
Iliadi, K. G., O. B. Gluscencova, N. Iliadi and G. L. Boulianne, 2018 Mutations in the
Drosophila homolog of human PLA2G6 give rise to age-dependent loss of
psychomotor activity and neurodegeneration. Sci Rep 8**:** 2939.
Kinghorn, K. J., and J. I. Castillo-Quan, 2016 Mitochondrial dysfunction and defects in
lipid homeostasis as therapeutic targets in neurodegeneration with brain iron
accumulation. Rare Dis 4**:** e1128616.
Kinghorn, K. J., J. I. Castillo-Quan, F. Bartolome, P. R. Angelova, L. Li *et al.*, 2015 Loss
of PLA2G6 leads to elevated mitochondrial lipid peroxidation and mitochondrial
dysfunction. Brain 138**:** 1801-1816.
Kofler, R., R. V. Pandey and C. Schlotterer, 2011 PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq).
Bioinformatics 27**:** 3435-3436.
Lango Allen, H., K. Estrada, G. Lettre, S. I. Berndt, M. N. Weedon *et al.*, 2010 Hundreds
of variants clustered in genomic loci and biological pathways affect human height.
Nature 467**:** 832-838.

Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25**:** 1754-1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence Alignment/Map format and SAMtools. Bioinformatics 25**:** 2078-2079.

Lin, G., P. T. Lee, K. Chen, D. Mao, K. L. Tan *et al.*, 2018 Phospholipase PLA2G6, a Parkinsonism-Associated Gene, Affects Vps26 and Vps35, Retromer Function, and Ceramide Levels, Similar to alpha-Synuclein Gain. Cell Metab 28**:** 605-618 e606.

Mackay, T. F., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The Drosophila melanogaster Genetic Reference Panel. Nature 482**:** 173-178.

Mackay, T. F., E. A. Stone and J. F. Ayroles, 2009 The genetics of quantitative traits: challenges and prospects. Nat Rev Genet 10**:** 565-577.

Mackay, T. F. C., and W. Huang, 2018 Charting the genotype-phenotype map: lessons from the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip Rev Dev Biol 7.

Metz, S. A., 1988a Exogenous arachidonic acid promotes insulin release from intact or permeabilized rat islets by dual mechanisms. Putative activation of Ca2+ mobilization and protein kinase C. Diabetes 37**:** 1453-1469.

Metz, S. A., 1988b Membrane Phospholipid Turnover as an Intermediary Step in Insulin-Secretion - Putative Roles of Phospholipases in Cell Signaling. American Journal of Medicine 85**:** 9-21.

Mori, A., T. Hatano, T. Inoshita, K. Shiba-Fukushima, T. Koinuma *et al.*, 2019 Parkinson's disease-associated iPLA2-VIA/PLA2G6 regulates neuronal functions and alpha-synuclein stability through membrane remodeling. Proc Natl Acad Sci U S A 116**:** 20689-20699.

Mossman, J. A., L. M. Biancani, C. T. Zhu and D. M. Rand, 2016 Mitonuclear Epistasis for Development Time and Its Modification by Diet in Drosophila. Genetics 203**:** 463-484.

Pool, J. E., 2016 Genetic Mapping by Bulk Segregant Analysis in Drosophila: Experimental Design and Simulation-Based Inference. Genetics 204**:** 1295-1306.

Smith, S. P., S. Shahamatdar, W. Cheng, S. Zhang, et al. , 2022 Enrichment analyses identify shared associations for 25 quantitative traits in over 600,000 individuals from seven diverse ancestries. Am J Hum Genet. 2022 May 5;109(5):871-884.

Song, K., X. Zhang, C. Zhao, N. T. Ang and Z. A. Ma, 2005 Inhibition of Ca2+-independent phospholipase A2 results in insufficient insulin secretion and impaired glucose tolerance. Mol Endocrinol 19**:** 504-515.

Tennessen, J. M., W. E. Barry, J. Cox and C. S. Thummel, 2014 Methods for studying metabolism in Drosophila. Methods 68**:** 105-115.

Therneau, T. M., and P. M. Grambsch, 2000 *Modeling survival data : extending the Cox model*. Springer, New York.

Chromosome	start	end	REF	ALT	GENEID	CONSEQUENCE	REFCODON
chr3L	9824166	9824166	C	T	FBgn0264489	nonsynonymous	CCA
chr3L	9833935	9833935	G	A	FBgn0004390	synonymous	AGC
chr3L	9854881	9854881	C	A	FBgn0036053	nonsynonymous	GGC
chr3L	9880027	9880027	A	G	FBgn0011836	synonymous	TTT
chr3L	9880564	9880564	G	T	FBgn0011836	synonymous	ATC
chr3L	9881643	9881643	C	A	FBgn0011836	nonsynonymous	GAG
chr3L	9884329	9884329	A	T	FBgn0025866	synonymous	ATA
chr3L	9884605	9884605	A	G	FBgn0025866	synonymous	GGA

VARCODON	REFAA	VARAA	SYMBOL
TCA	P	S	CG43897
AGT	S	S	RasGAP1
GTC	G	V	iPLA2-VIA
TTC	F	F	Taf2
ATA	I	I	Taf2
GAT	E	D	Taf2
ATT	I	I	CalpB
GGG	G	G	CalpB