Natural variation in phospholipase IPLA2-VIA contributes to starvation sensitivity
in *Drosophila melanogaster*

Shawn B. Williams *[†], Brian Franklin[†‡], Faye A. Lemieux[†], David M Rand[†]

*[*]Department of Molecular Biology, Cell Biology and Biochemistry, Brown University,
Providence, RI 02906, USA

[†]Department of Ecology and Evolutionary Biology, Brown University, Providence, RI
02906, USA

[‡]Current address: University of Pennsylvania Perelman School of Medicine,

Philadelphia, PA 19104, USA

Correspondence:

David M Rand

Brown University

Box G-W

80 Waterman Street

Providence, RI 02912

Phone: (401) 863-2890

Email: david_rand@brown.edu

Abstract

Resistance to starvation is a classic complex trait where genetic and environmental variables can greatly modify an animal's ability to survive without nutrients. Genetic analyses in *Drosophila* have shown that starvation resistance is highly polygenic with different genetic architectures in different mapping populations. In this study we sought to dissect the genetic basis of starvation resistance among a set of mitonuclear genotypes carrying different mtDNAs paired with specific nuclear genomes from the Drosophila Genetic Reference Panel (DGRP). We focused on differences between one of the most sensitive strains (DGRP-765) and a strain with more moderate resistance (DGRP-315) whose starvation phenotypes appeared to be modified by alternative mtDNAs. Using complementary pooled-sequencing and forward genetic mapping approaches, we identified regions of chromosomes 2L, 3L and 3R contributing to starvation sensitivity and localize a major effect locus modifying starvation resistance to the coding region of phospholipase iPLA2-VIA. These analyses further confirm that the alternative mtDNAs had little influence on variation in starvation resistance between the genotypes studied. The sensitive line shows a starvation-dependent depletion of glucose and glycogen that is modified by hemi- and heterozygosity in the iPLA2-VIA region. These findings contribute to our understanding of the complex genetic relationship between resistance to starvation stress and nutrient metabolism.

Introduction

Coping with periods of low nutrient availability is a fundamental fitness component of all organisms. In animals, multiple metabolic processes are required to convert dietary nutrients into usable resources to successfully cope with periods of starvation. Energy that is not needed immediately has to be converted into a form amenable for efficient storage and that stored energy has to be accessible for utilization during periods of starvation. For these reasons, the genetics of starvation resistance provides a compelling model for understanding the integrative aspects of animal physiology (BROWN *et al.* 2019). Furthermore, a better understanding of the genetic basis of nutrient catabolism and storage will also shed a light on genes relevant to metabolic diseases and fitness traits in nature (HARDY *et al.* 2018).

Much of the information on the genetics of starvation resistance comes from studies in Drosophila employing either direct selection on experimental outbred populations or genome wide association studies (GWAS) across samples of inbred lines. Variation in starvation resistance is an appealing trait for studies of quantitative and evolutionary genetics due to its highly polygenic nature and its correlation with other fitness and metabolic traits (SERVICE AND ROSE 1985; CHIPPINDALE *et al.* 1996; HARSHMAN *et al.* 1999; HARDY *et al.* 2018; EVERMAN *et al.* 2019). GWAS approaches using either the Drosophila Genetic Reference Panel (DGRP;(MACKAY *et al.* 2012)) or the Drosophila Synthetic Population Resource (DSPR;(EVERMAN *et al.* 2019)) have identified a number of candidate loci influencing variation in starvation resistance confirming the complexity of its polygenic nature (HARBISON *et al.* 2004; SORENSEN *et al.* 2007; HUANG *et al.* 2012; MACKAY *et al.* 2012; EVERMAN AND MORGAN 2018; EVERMAN *et al.* 2019). Importantly, there are clear differences

among studies in the location of QTLs and the putative candidate loci identified (HUANG

et al. 2012; MACKAY *et al.* 2012; EVERMAN AND MORGAN 2018; EVERMAN *et al.* 2019). These

differences have been attributed to the distinct genetic composition of the mapping

populations and presumably unique epistatic interactions among loci influencing

starvation resistance (*ops. cit.*). Notably, none of the QTL identified in the two subsets of

the mapping lines used by (EVERMAN *et al.* 2019) overlapped, likely due to the different

founder lines used to build each subset (DSPR pA and pB). Despite these differences,

there are shared loci that emerge from a comparison of multiple studies (table S9 in

(EVERMAN *et al.* 2019)).

A growing literature has identified a role for mitochondrial functions in stress responses

at multiple levels (UMA NARESH *et al.* 2022; ERKOSAR *et al.* 2025). Given the central roles

that mitochondria play in both anabolic and catabolic functions, the complex interactions

between mitochondrial and nuclear genomes may play important roles in adaptive traits,

and particularly in metabolically demanding conditions such as starvation, oxidative stress

and animal performance (RAND *et al.* 2004; HILL 2015). In previous work, we demonstrated

the impact of mitonuclear epistasis in development time among a panel of 72 mitonuclear

genotypes constructed from 6 mtDNAs introgressed into 12 nuclear genomes from the

Drosophila Genetic Reference Panel (DGRP) (MOSSMAN *et al.* 2016). We selected a

subset of these lines to examine starvation resistance based on significant differences in

development time between mtDNAs in each of several DGRP backgrounds. These

analyses revealed a putative mitonuclear epistatic interaction influencing starvation

resistance and its relation to diet composition. In the current study, we sought to use a

combination of quantitative genetic and classical genetic mapping approaches to examine

the genetic basis of starvation resistance between two DGRP lines carrying different

mtDNAs (DGRP-765 and -315). Genomic sequencing of sensitive and resistant pools of

adults from advanced intercross populations derived from these lines identified regions

of allelic differentiation. Chromosome segregation and recombination mapping identified

major factors on chromosome 3L, 3R and 2L, but not chromosome X, 2R or the

mitochondrial genome. Deficiency mapping indicated that the starvation sensitivity in

DGRP-315 localizes in part to the phospholipase iPLA2-VIA gene with associated effects

of carbohydrate metabolism. The correspondence of these approaches contributes to our

understanding of the polygenic nature of starvation resistance and points to

polymorphisms in the iPLA2-VIA region that motivate further study of the associations

between starvation sensitivity, lipid metabolism and carbohydrate metabolism.

Materials and Methods

Fly stocks

The current study focuses on the differences in starvation resistance between specific

mitonuclear genotypes carrying alternative mtDNAs placed onto nuclear genomic

backgrounds of the Drosophila Genetic Reference Panel (DGRP;(MACKAY *et al.* 2012)).

In a previous study we constructed a panel of 72 mitonuclear genotypes using balancer

chromosome substitution to place six mtDNAs onto each of 12 DGRP nuclear

backgrounds to study mitonuclear epistasis for development time and its dietary

modification (MOSSMAN *et al.* 2016). From these 72 lines we quantified starvation

resistance of 12 mitonuclear genotypes consisting of four different mtDNAs placed on

three DGRP nuclear backgrounds (lines DGRP-765, -315 and -820). The four mtDNAs

included two from *D. simulans* (*siI and siII*) and two from *D. melanogaster* (*OregonR =*

OreR and *Zimbabwe53 = Zim53*). These mtDNAs were chosen to test for the impact of

mtDNA polymorphism and divergence on starvation resistance. The DGRP

backgrounds were chosen based on published starvation data showing that DGRP-765,

-315 and -820 had low, medium and high values of starvation resistance, respectively

(AYROLES *et al.* 2009; MACKAY *et al.* 2012). The genotype notation used is

mtDNA;nuclearDNA, e.g., *Zim53;315* carries the *D. melanogaster Zim53* mtDNA on a

homozygous *DGRP-3*15 chromosomal background.

The results of this starvation assay are shown in Figure 1. The resistance levels were

consistent with previously published rankings of the three nuclear genotypes (820 > 315

> 765). The results further showed that one unique combination, *siI;765,* (a *D. simulans*

siI mtDNA on the *DGRP-765* nuclear background) had a significant increase in

starvation resistance compared to the other mitonuclear genotypes in the *DGRP-765*

background. The goal of the current study was to determine whether the apparent

'rescue' of starvation resistance was due to the introgression of the *siI* mitochondrial

haplotype or due to a nuclear factor or factors. The focal mitonuclear genotypes are

shown in Table 1. Additional genotypes for segregation, recombination and deficiency

mapping are listed in Table S1.

Starvation Stress

Survival during starvation stress was measured by placing approximately 125 female flies

per genotype in each of three replicate demography cages. Flies were collected and

sorted over CO_2 and allowed to recover in quart-sized plastic demography cages affixed

with a 25x95 mm vial containing one of three different isocaloric food media: high sugar

food (15% sucrose, 5% Yeast, 2% agar), balanced food (10% sucrose, 10% yeast, 2%

agar) or low sugar food (5% sucrose, 15% yeast, 2% agar; referred to as food type V, II

and IV, respectively, in (ZHU *et al.* 2014)). The three replicate cages for each genotype

were conditioned on these diets for 7 days unless otherwise indicated, with new food vials

added every other day. After 7 days, any dead flies were removed, and the vial for each

demography cage was replaced with 2% Difco Bacto-agar prepared with deionized H_2O

to initiation starvation. The number of dead flies was recorded every 12 hours and

removed from each demography cage until all flies were dead. As starvation resistance

values for the different genotypes were most distinct on the high sugar diet "V", most of

the genetic mapping was carried out on that diet. The survival data from each cage were

generally consistent so counts from each replicate cage were pooled and the combined

data modeled using a Kaplan-Meir survival model implemented in R (THERNEAU AND

GRAMBSCH 2000). Significant differences in survivals were determined using a log rank

test.

Forward genetic mapping of starvation sensitivity

Chromosomal localization of resistance. To determine if different mtDNAs were

associated with the difference in starvation resistance reciprocal crosses were done

between siI;765 and Zim53;765 as follows: virgin female siI;765 x male Zim53;765 and

virgin female Zim53;765 x male sil;765). Maternal inheritance of mtDNA provides a test

for the role of the *sil* mtDNA in the rescue of starvation sensitivity. To localize nuclear

factors associated with starvation sensitivity we generated genotypes that were

selectively heterozygous for different combinations of *Zim53;765* autosomes. Twenty

Zim53;765 virgin females were first crossed to 20 males from the double-balancer stock

(wg^{Sp-1}/*CyO; Dr^1/TM3, Sb^1*). Twenty F1 male progeny with both the 2nd chromosome

CyO and 3rd chromosome *Dr* markers present were sorted over CO_2 and crossed to 20

Zim53;765 virgin females in eight replicate 20-female x 20-male crosses. Progeny from

this second cross were segregated into four groups based on the presence or absence

of the *CyO* and *D*r markers. The four groups were: reconstituted wild-type *Zim53;765*,

chromosome 2 heterozygotes (*CyO/765*), chromosome 3 heterozygotes (*Dr/765*) and

heterozygotes for both chromosome 2 and 3 (*CyO/765;Dr/765*). Flies from each group

were moved to 3 replicate demography cages with approximately 125 flies each. The

survival of each group under starvation stress was assayed as described earlier.

Whole genome sequencing of sensitive and resistant pools from Advanced

Intercross Populations

Population Cages. Five replicate advanced intercross populations (AIPs) were

established between the sensitive (*Zim53;765*) and resistant (*Zim53;315*) lines. Each

replicate population was initiated using 25 *Zim53;315* virgin females crossed to 25

Zim53;765 males, in addition to the reciprocal cross, to yield a total number of 50 male

and 50 female parents of each line as the founders of each of the five replicate cages.

These parents were introduced into Plexiglas cages (20 x 20 x 20cm) and allowed to lay

eggs in four 125mL culture bottles with 30mL of Drosophila medium in each bottle. The

four bottles in each cage were left with their tops open for 3 days, and then each bottle

was plugged to allow flies to develop. Prior to eclosion, the plugs were removed from

each bottle, and all flies were allowed to emerge into the cage and mate at random. Four

new culture bottles replaced the old bottles in each replicate cage, and the adults were

allowed to lay eggs for 48 hours. After egg laying, adults were cleared from each bottle,

plugs were inserted, and cultures were allowed to develop for 7 days following the

termination of egg laying, after which the plugs were removed from all bottles to allow

eclosion of the next generation into each cage. Population sizes exceeded 1000

individuals after the first generation. All five cages were maintained at large population

sizes (>2000 adults) for 10 generations to allow recombination to take place between the

sensitive (*Zim53;765*) and resistant (*Zim53;315*) DGRP nuclear genotypes. These two

DGRP strains are homozygous for the standard (ST) chromosome arrangement

(http://dgrp2.gnets.ncsu.edu/) so this lack of inversions should have allowed free

recombination across the genome in these populations.

Starvation Stress. The female flies from each generation-10 population cage were

collected over CO_2 and moved to demography cages with a density of approximately 150

flies per cage. The flies were allowed to recover for 7 days with access to high sugar food

(Type V: 15% sucrose, 5% Yeast, 2% agar) and then subjected to starvation conditions

by replacing the food vials with a 2% agar and water mix. Every six hours from the start

of starvation, all the dead flies were collected, flash frozen and stored at -80C. The high

sugar treatment enhanced the difference among genotypes in starvation resistance. The

genotypic differences in starvation were also evident with moderate or low sugar

treatments but mapping experiments were more effective with the high sugar treatment

so that was used throughout these analyses.

Pool-sequencing. Pools of approximately 50 dead flies were made by combining flies from

the two earliest time points when deaths were first observed (24 hour and 30 hour) and

the last two time points when the last live flies were present (96 hour and 102 hour). DNA

from each replicate pool was isolated using the DNeasy DNA purification kit from Qiagen

(Cat No 69506). Each sample was RNase treated prior to DNA elution using a 10-minute

RNase A (Cat No 19101) treatment. Libraries were prepared and sequenced using the

DNBseq platform by the Beijing Genomics Institute (BGI).

Estimation of genetic differentiation using Popoolation2. Raw reads were mapped to the

dm6 build of the *Drosophila* genome using *bwa* version 0.7.15 (LI AND DURBIN 2009).

Variants were called using *samtools* version 1.9 (LI *et al.* 2009). Allele frequency

differences and estimates of population subdivision (F_{ST}) were determined using

*Popoolation2 (*KOFLER *et al.* 2011*)*, and the significance of differences in allele frequencies

between the early-dying and late-dying sample across the five replicate populations was

determined using the Cochran-Mantel-Haenszel test. P-values were corrected using the

Bonferroni method and differences with FDR-corrected p-values < 0.05 were considered

to be significant. Linkage disequilibrium was estimated from paired end reads, following

the procedure described in (FEDER *et al.* 2012). The expectation of this AIP pool-seq

experiment is that recombination would segregate major factors responsible for the

difference in starvation resistance between these strains so that extreme-phenotype

mapping by pooled sequencing might identify chromosomal regions associated with this

difference. We acknowledge that 10 generations, moderate populations sizes (~2000)

and founding samples of 50 extreme-phenotype adults may not have the power to

generate high resolution recombinants for localizing individual loci, but our intention was

to use this as a means to complement other classical mapping approaches (chromosome

segregation, marker-recombination assays and deficiency mapping). The use of five

replicate populations provided some level of replication that was appropriate for the

budget available at the time.

Chromosome 3 recombination

The chromosome segregation experiments indicated that the majority of the difference in

starvation resistance between the lines mapped to chromosome 3 and the pooled

sequence analyses of the AIP cages identified two broad candidate regions on

chromosome 3, one each on 3L and 3R. We performed recombination mapping using

three successive crosses to marked mapping stocks to further localize candidate regions

(cross 1, 2, 3; see Fig. S1). For cross 1 we attempted to generate *Zim53;765* chromosome

3 recombinants by crossing 20 *Zim53;765* virgin females to 20 males from a multiply

marked recessive stock obtained from the Bloomington Stock Center (BSC_576 = *ru^1*

hry^1 Diap^{th-1} st^1 cu^1 sr^1 e^s ca^1 hereafter referred to as BSC_576 for convenience). The

progeny from this cross however failed to eclose. As an alternative cross 1, we crossed

20 *sil;765* virgin females to 20 *BSC_576* males in an effort to allow recombination across

the *DGRP-765* 3rd chromosomes that are present in the *sil;765* stock. The *sil;765*

genotype was suspected to be heterozygous for the *DGRP-765* 3rd chromosome and a

3rd chromosome from the *TM3* balancer stock that may have recombined with the *DGRP-*

765 3rd chromosome during the construction of the *siI;765* original stock since *siI;765*

lacked the *TM3* marker *Sb^1* (Sb is at 89B4, band 58). This suspicion of balanced

heterozygosity was based on the observation that several generations of backcrossing of

male DGRP-765 to virgin female *siI;765* did not reduce starvation resistance implying that

the *siI;765* carried some heterozygosity and was in some way resistant to the impact of

recurrent backcrossing.

For cross 2, 20 F$_1$ virgin *siI;765* / BSC_576 females were collected and crossed to 20

males from a multiply marked recessive stock with the same recessive markers as

BSC_576 in addition to a dominant marker present on one copy of chromosome 3

balanced over the TM6 balancer (BSC_1711: *ru^1 hry^1 Diap^{th-1} st^1 cu^1 sr^1 e^s Pri^1 ca^1* /

TM6B, Bri^1, Tb^1). This cross was done in three replicate 20-female x 20-male vials. F$_1$

males from this second cross were collected and segregated into distinct recombinant

classes based on the recessive and dominant markers present.

For cross 3, males from each of these recombinant categories were each mated to virgin

starvation resistant *siI;765* females and to virgin starvation sensitive *Zim53;765* females

to generate offspring with different segments of the 3rd chromosome that was

homozygous for the starvation-sensitive *DGRP-765* chromosome. Comparison of the

offspring from these two different mitonuclear lines that showed significant differences in

starvation (*siI;765* and *Zim53;765*) could further resolve regions of chromosome 3

associated with starvation sensitivity. Due to the presumed heterozygosity of the *siI;765*

3rd chromosome we expected that recombination between the multiple-marked 3rd

chromosome with the *sil;765* 3rd chromosomes would be confounded if: 1) the

chromosome suspected to be the homologue of the 3rd chromosome in *sil;765* was a

broken balancer chromosome, or 2) if that homologue was some other wild type

chromosome that had been recombining with the *DGRP-765* 3rd chromosome. This would

reduce the efficiency of the recombination mapping as only an unknown fraction of the

offspring would have been derived from the actual pairing of the *sil;765* 3rd chromosome

derived from DGRP-765 and the recombinant-marked chromosomes derived from the F1

female parent. We acknowledge that this was not ideal but collected the mapping data to

see if it provided useful information consistent with the independent chromosome

segregation mapping and AIP pool-seq data.

Five classes of recombinant offspring phenotypes were designated based on the visible

markers present reflecting different breakpoints along the chromosome: class i*: ru, h*;

class ii: *ru, h, Diap, st*; class iii: *ru, h, Diap, st, cu*; class iv: *ru, h, Diap, st, cu, sr*; and class

v: *ru, h, Diap, st, cu, sr, e*. Twenty males from each group were crossed to 20 *Zim53;765*

virgin females and allowed to mate for 48 hours in three replicate 20-female x 20-male

crosses (the *Zim53;765* genotype was confirmed to be homozygous for the DGRP-765

chromosome). The males from each of these crosses were removed and then re-mated

to *sil;765* virgin females. Female offspring from each of these two sets of crosses were

collected over CO_2. Sorted by recombinant class and conditioned for starvation assays

on high sugar food. Sufficient flies for a starvation stress survival assay were only

available from the class i, class ii and class v crosses, so only these were considered.

The survival of these females under starvation stress was measured as described above.

Chromosome 3L Deficiency Screen

The recombination mapping and the AIP pool-seq data implicated chromosome 3L as harboring a major starvation factor, so deficiency mapping was pursued. Twenty males from each deficiency line (Supplemental Table 1) were crossed to 20 *Zim53;765* virgin females in eight replicate 20-female x 20-male crosses. F_1 females were collected and sorted over CO_2. Females were separated based on the presence of the respective balancer marker (flies heterozygous for the deficiency and the *765* chromosome do not have the balancer present) and moved to 3 replicate demography cages with approximately 125 flies each, for each genotype. The flies were allowed to recover for 7 days on high sugar food and survival under starvation stress measured as described above.

iPLA2-VIA Complementation

The chromosome 3L deficiency mapping, in conjunction with the available DGRP sequences, identified *iPLA2-VIA* as a candidate gene based on nucleotide polymorphisms private to *DGRP-765* and starvation sensitivity phenotypes of *iPLA2-VIA* alleles (LIN *et al.* 2018). Deficiencies within that gene were obtained from the Bloomington Stock Center for further complementation mapping: *BSC_80133 = y¹ w*; iPLA2-VIA^Delta174* and *BSC_80134 = y¹ w¹; iPLA2-VIA^Delta192* hereafter *80133* and *80134* respectively. These deficiencies were generated by imprecise excision of a P-element from *iPLA-VIA* that removed segments of 5' noncoding sequence and exons 1-5 and -7 (LIN *et al.* 2018). Twenty *Zim53;765* flies were crossed in reciprocal to 20 flies from each of these two *iPLA-*

VIA coding region deficiencies. Each cross was done in replicates of 4. As a control,

males from each *iPLA2-VIA* deficiency stock were also crossed to *Zim53;315* females

with the expectation that the *DGRP-315* nuclear genome would complement the iPLA2-

VIA deficiencies and restore starvation to normal levels . F_1 females were collected and

sorted over CO_2 and moved to 3 replicate demography cages with approximately 125

flies each. The flies were allowed to recover for 7 days on high sugar food and survival

under starvation stress measured as described above.

Glucose Assay

The free glucose levels, as described in (TENNESSEN *et al.* 2014), were measured for five

genotypes: *Zim53;765*, *Zim53;315*, *iPLA2-VIA^{Delta174}*, *Zim53;315* crossed to *iPLA2-*

VIA^{Delta174} and *Zim53;765* crossed to *iPLA2-VIA^{Delta174}*. 120 female flies from each

genotype were collected and sorted over CO_2 and allowed to recover in demography

cages affixed with 25x95 mm vials of high sugar food (15% sucrose, 5% Yeast, 2% agar)

for five days. Three replicates were done for each genotype. After five days, any dead

flies were removed and half of the remaining live flies were removed from the demography

cage, immediately flash frozen in liquid N_2 and stored at -80°C. The food supplied to each

demography cage was replaced with 2% Difco Bacto-agar prepared with deionized H_2O.

After 12 hours, dead flies were discarded and the remaining live flies were removed from

the demography cage, immediately flash frozen in liquid N_2 and stored at -80°C. Six

frozen flies from each replicate were transferred to a 1.5uL micro-centrifuge tube and

homogenized in 120uL of Phosphate Buffered Saline (PBS) using a steel bead and

Qiagen TissueLyser II set to 30Hz for 2 minutes. A 10uL aliquot was removed to measure

protein content using the Pierce™ BCA Protein Assay Kit (23225). The remaining

homogenate was heat treated for 10 minutes at 70°C. The homogenate was diluted 1:4

with PBS. The free glucose was measured with the Glucose Oxidase (GO) kit (MAK097-

1KT) by following the protocol for free glucose measurement laid out in (TENNESSEN *et al.*

2014).

Data Availability

Drosophila strains, code and raw data are available at the Rand Lab GitHub site for this

publication (https://github.com/DavidRandLab/Williams-et-al.-Starvation-Genetics-2021).

The authors affirm that all analyzed data necessary for confirming the conclusions of the

article are present within the article, figures, tables and GitHub site. Raw sequences for

the pool-seq mapping are available at BioProject accession number: PRJNA1130456.

Results
Starvation resistance among mitonuclear genotypes

Twelve mitonuclear genotypes from a panel of 72 previously-reported genotypes

(MOSSMAN *et al.* 2016) were chosen for starvation assays as described in the Methods

section. Each of these 12 mitonuclear introgression lines was subjected to starvation

assays on three different isocaloric diets: high sugar food (15% sucrose, 5% Yeast, 2%

agar), balanced food (10% sucrose, 10% yeast, 2% agar) or low sugar food (5%

sucrose, 15% yeast, 2% agar; referred to as food type V, II and IV, respectively (ZHU *et*

al. 2014)). The mitonuclear genotypes on the *DGRP-765* nuclear background were of

particular interest because the *DGRP-765* line represents one of the least resistant lines

in terms of starvation resistance in the DGRP, and it appeared that introgression of the

siI mitochondrial DNA haplotype significantly improved resistance to starvation only in

the *DGRP-765* nuclear background (Figure 1). To test whether the rescue of starvation

resistance mapped to a cytoplasmic or nuclear factor, we performed reciprocal crosses

to examine if maternal inheritance of mtDNA was the source of the rescue of starvation

sensitivity. When we crossed *Zim53;765* to *siI;765*, the direction of the cross (*Zim53;765*

female crossed *siI;765* males vs. *siI;765* females to *Zim53;765 males*) made no

difference to starvation resistance of its progeny. This implied that the *siI* mtDNA

haplotype was not the source of the rescue, as would be expected if the trait was being

inherited maternally (Fig. S2).

We analyzed the reads from an RNA-sequencing data set from each fly line and

compared the genotype calls for each SNP present in *Zim53;765* reads (restricted to the

coding regions) and the genotype calls at the same locations in *siI;765*. For the vast

majority of these sites, the *siI;765* was heterozygous for the *DGRP-765* allele and an

alternative allele (in most cases the reference allele) (Fig. S2A,B). We did a similar

comparison of *siI;765* to an alternate mitonuclear genotype (*Zim53;315*) and did not see

a similar overlap between the *siI;765* allele and the *Zim53;315* allele (Fig. S2A,B).

Based on these results we concluded that the *siI;765* genotypes was heterozygous

across the *DGRP-765* chromosomes. This suggested to us that starvation sensitivity in

Zim53;765, and the two other *mito;765* combinations (*Ore;765* and *siII;765*), were

homozygous recessive and the improvement in resistance to starvation seen in *siI;765*

was due to its nuclear heterozygosity. Indeed, we observed that the offspring from a

Zim53;765 to *Zim53;315* cross were significantly more resistant to starvation than

Zim53;765 (see Figure 6).

Chromosome 3 heterozygosity partially suppresses *Zim53;765* starvation
sensitivity
391
Based on the suppression of starvation sensitivity seen in *Zim53;765* heterozygotes,

(Zim53;765 x Zim53:315 F1 hybrids) we reasoned that starvation sensitivity phenotypes

in *Zim53;765* and other genotypes were homozygous recessive traits and that we could

localize the effect locus/loci to chromosomes using chromosome segregation tests and

observing which chromosome segregants masked the trait. Virgin females from three

different genotypes (the original *DGRP-765, Zim53;765* and *Zim53;315*) were mated to

males from the marked double-balancer stock (wg^{Sp-1}/CyO; Dr^1/TM3, Sb^1). F1 males

heterozygous for the maternal autosomes paired with the paternal *CyO* and Dr^1 markers

were mated back to virgin females from the three maternal genotypes, generating the

four 2-autosome segregant populations of F1 flies as follows: heterozygous for

chromosome 2, heterozygous for chromosome 3, heterozygous for both chromosome 2

and 3, or neither (the latter being reconstituted homozygous wild type 2nd and 3rd

chromosomes of the original maternal genotypes: *DGRP-765, Zim53;765* and

Zim53;315). *DGRP-765* and *Zim53;765* flies heterozygous for chromosome 3 had a

~200% increase in mean survival relative to the original stock (Figure 2A; Table

S2:SummaryTable_figure2a) and flies heterozygous for both chromosomes 2 and 3

realized additional increases in mean survival. Heterozygosity for chromosome 2 alone

contributed ~75% increase (range 50-116%; see Table S2:SummaryTable_figure2a).

For Zim53:315, chromosome 3 heterozygosity contributed nothing alone but a small

effect in combination with chromosome 2.

A repeat segregation experiment with *Zim53;765* and the heterozygous *sil;765*

genotype was conducted and revealed comparable results for *Zim53;765* (80-100%

increase from chromosome 3 heterozygosity with additional effects of chromosome 2)

and very different chromosomal influences for *sil;765*, as expected from its

heterozygous condition (Figure 2B; see Table S2:SummaryTable_figure2b; raw data

are in other tabs of Table S2). Taken together, these results suggest that the main

effect locus/loci controlling *Zim53;765* and DGRP-765 starvation sensitivity are present

on chromosome 3 with a secondary locus/loci present on chromosome 2.

Sites on chromosome 2R and 3 have significant differences in allele frequency
between early and late dying flies.
422
To understand the genetic basis of 765 starvation sensitivity, we sought to identify genetic

differences characteristic of starvation sensitive flies. We created replicate advanced

intercross populations (AIPs) in which *Zim53;765* was allowed to recombine with

Zim53;315. We set up five independent populations where *Zim53;765* flies were crossed

to *Zim;315* in reciprocal and the populations maintained for 10 generations at population

sizes >1000 adults. Following this, females from each population were subjected to

starvation stress and dead flies were collected every six hours until all flies were dead.

Flies that died during the first three and last two six-hour intervals were separately

combined into early- and late-dying pools of adult females (18-24, 24-30 and 30-36 hours

vs. 90-96 and 96-102 hours). Total genomic DNA was sequenced from each of the five

replicates from these two pooled samples, hereafter 'early' and 'late' dying pools.

Consistent differences in allele frequencies across the early and late dying populations

were determined using the Cochran-Mantel-Haenszel test implemented in Popoolation2

(KOFLER *et al.* 2011). There was an enrichment in significantly different variants across

chromosome 3 and a localized region of chromosome 2R (Figure 3). These results

suggest that *Zim53;765* and as a consequence *DGRP-765*, starvation sensitivity is based

on variants present on chromosome 3 and chromosome 2. Importantly, these advanced

intercross pool-seq analyses are consistent with the chromosome segregation analyses

where factors on chromosomes 3 and 2 were responsible for starvation sensitivity of the

Zim;765 genotype.

Heterozygosity of chromosome 3L region partially suppresses Zim53;765
starvation sensitivity
446

To further localize the main effect loci on chromosome 3 we used recombination and

deficiency mapping across chromosome 3. As there are over 200 deficiencies spanning

chromosome 3, we first generated a series of recombinants with break points at different

locations along chromosome 3. Cross 1 used *sil;765* virgin females mated to a multiply

marked recessive stock (*ru^1 hry^1 Diap^{th-1} st^1 cu^1 sr^1 e^s ca^1* hereafter referred to as

BSC_576 for convenience). Cross 2 used heterozygous F$_1$ virgin females from cross 1

mated to males from a similarly marked recessive stock with one dominant marker added

to distinguish progeny (BSC_1711: *ru^1 hry^1 Diap^{th-1} st^1 cu^1 sr^1 e^s Pri^1 ca^1 / TM6B, Bri^1,*

Tb^1). The F1 offspring from cross 2 revealed recombinant males that were pooled into

three classes of recombinants (classes i, ii, v) that were selected based on the presence

of visible recessive markers and sufficient flies for starvation analyses (Figure 4A).

Recombinant class i is likely heterozygous for the region of 3L from the telomere up to at

least the regions marked by *h,* with recombination breakpoints existing anywhere

between h and the *cu* gene. Recombinant class ii is heterozygous for most of 3L and up

to *cu* gene on 3R, with breakpoints existing anywhere between the *cu* gene and the *ca*

gene. Recombinant class v is likely heterozygous for most of chromosome 3 with

breakpoints existing from the *ca* gene up to the telomere of 3R. Other classes of

recombinants did not emerge in sufficient numbers to conduct the final mapping crosses.

It is important to note that because these recombinants were generated from *sil;765*, the

pool of recombinants will carry both the *DGRP-765* alleles from that chromosome which

was substituted onto the *D. simulans sil* mtDNA background during original stock

construction and alternate, unknown alleles from a chromosome that was likely

introduced during the balancer substitution process during original line construction. A

TM3 Sb^1 3rd chromosome balancer was used in these crosses and may have recombined

with segments of the *DGRP-765* 3rd chromosome during the replacement crosses (see

Methods).

To reveal the localization, cross 3 used males from the three recombinant classes

emerging from cross 2, each of which were mated to virgin *Zim53;765* females, and then

the same males were mated to virgin *sil;765* as a control. The starvation sensitivity of the

female offspring of the *Zim53;765* and recombinant males was compared to the female

offspring of the *sil;765* and recombinant male controls. While we cannot discern genotype

at the individual fly level, *DGRP-765* chromosome 3 alleles will be enriched in the progeny

from the *Zim53;765* vs recombinant cross, while the alternative will be true for the *sil;765*

vs recombinant cross. We expected that flies homozygous for the causal *Zim53;765* allele

would die earlier during starvation stress, thus revealing the region of *Zim53;765*

chromosome 3 responsible for its starvation sensitivity. Although the heterogeneity of

respective offspring populations will mask this effect, the enrichment of the *DGRP-765*

allele in the *Zim53;765* vs recombinant cross population and the alternate allele in the

sil;765 vs recombinant cross population should result in appreciable differences in deaths

early on during starvation stress. Indeed, we observed a significant difference between

the starvation sensitivity of recombinant class i offspring, but not in the case of the class

ii offspring; class v offspring showed a small but significant difference (see Figure 4B;

Table S3). These results indicate that heterozygosity of a region between the *h* marker of

recombinant class i and the *cu* marker of recombinant class ii carries a factor required to

suppress *Zim53;765* starvation sensitivity.

Hemizygosity of 3L: 9812381 – 9899255 prevents partial suppression of *Zim53;765*
starvation sensitivity
494
We employed a deficiency screen across the region between recombinant class i and

recombinant class ii markers. The objective of this screen was to identify the region(s) of

chromosome 3L that failed to complement *Zim53;765* starvation sensitivity. *Zim53;765*

virgin females were crossed to a series of flies deficient for various parts of chromosome

3L (Table S1). The deficient region of each line is maintained over a balancer

chromosome so the resulting offspring can be positive for the balancer, indicating

complete heterozygosity of chromosome 3, or offspring lacking the balancer are

hemizygous across the deficiency region and heterozygous for the rest of chromosome

3. Significant differences in the starvation resistance of the two classes of offspring

indicates that heterozygosity of the deficiency region is required for the suppression of

starvation sensitivity. We initially screened 15 deficiency lines and identified nine where

the starvation sensitivity was not suppressed (failed to complement) in the partial

chromosome 3L hemizygote (Table S4A and Fig.S3). Several of the deficiency stock

crosses with the most significant difference between Df and wt chromosome pairings

shared a common region of chromosome 3 between 3L:9812381 – 3L:9899255 (Fig. S3

and Figure 5), indicating that heterozygosity in this region is required for the suppression

of *Zim53;765* starvation sensitivity. Two deficiencies mapped to similar locations but

showed conflicting evidence for suppression of starvation sensitivity (Df282 and Df 283;

Fig. S3 and Table S1). The latter contained additional genes (lncRNA:CR45878,

CG46387) in the proximal region of the deficiency; the functional consequences of these

mapping differences were unclear. Three deficiencies on either side of the putative

overlap region also showed significant suppression of sensitivity (BSC113, 4470 and

4488). Despite the variation among the deficiency data, they were consistent with the

three   broader-scale   approaches   of   chromosome   segregation,   pool-seq   and

recombination mapping in identifying candidate genes in this region of chromosome 3L.

Variation in iPLA2-VIA as candidates for *Zim53;765* starvation sensitivity

Next, we sought to identify *Zim53;765* variants in the chromosome 3L region that might

underlie its sensitivity to starvation. We utilized the available genomic sequencing data of

each DGRP line to identify nucleotide variants unique to the *DGRP-765* sequence

(MACKAY *et al.* 2012). As the causal variant is deleterious, we focused our initial analysis

on low frequency variants (present in less than 10 % of the 200 DGRP2 sequences). We

identified 83 variants which fit this frequency cutoff and only thee SNPs were predicted to

lead to non-synonymous changes (Fig. S4 and Table 2). A C-to-A mutation, predicted to

result in a G-to-V amino acid residue change in the conserved GXGXXG DNA binding

motif of the phospholipase *iPLA2-VIA*, appeared as a strong candidate. Mutations in

iPLA2-VIA have previously been shown to cause starvation sensitivity and reduced

lifespan in *Drosophila (*KINGHORN *et al.* 2015*;* KINGHORN AND CASTILLO-QUAN 2016*;* LIN *et al.*

2018*)*. In addition, PolyPhen-2 predicted the mutation as probably damaging, with a score

of 0.999 out of 1 (ADZHUBEI *et al.* 2010). Based on this analysis, we performed finer

resolution deficiency mapping in the *iPLA2-VIA* gene.

iPLA2-VIA^{Del174} fails to complement Zim53;765/ DGRP-765 starvation sensitivity
and glucose homeostasis
To determine if the starvation phenotype of *Zim53;765* and as a consequence *DGRP-*

765, results from a defective copy of *iPLA2-VIA*, we sought to determine if the *Zim53;765*

copy of *iPLA2-VIA* gene could complement *iPLA2-VIA* null mutations. We crossed

Zim53;765 virgin females to two previously characterized *iPLA2-VIA* null lines (*iPLA2-*

VIA^{Del174}, *iPLA2-VIA^{Del192}*)*, generated by imprecise excision of a mobile element; Table

S1 (LIN *et al.* 2018). To rule out the potential effects of mtDNA haplotype on starvation

resistance, we also utilized the original *DGRP-765* strain obtained from the Bloomington

Stock center. We found that progeny from *DGRP-765* crossed to *iPLA2-VIA^{Del174}* or

iPLA2-VIA^{Del192} were also starvation sensitive (Figure 6). In contrast, crossing *iPLA2-*

VIA^{Del174} or *iPLA2-VIA^{Del192}* to the starvation resistant strain *Zim53;315* resulted in the

suppression of starvation sensitivity (i.e., restored starvation resistance; Figure 6).

Depletion of glucose and glycogen are associated with starvation sensitivity in

Zim53;765 and DGRP-765. Ablation or pharmaceutical inhibition of calcium independent

phospholipases have been shown to reduce insulin secretion in response to a glucose

challenge (SONG *et al.* 2005; ALI *et al.* 2013). This raised the question of whether *Zim53;765*

and *iPLA2-VIA* share defects in glucose metabolism. We observed that following 12 hours

of starvation, *Zim53;765* and the original *DGRP-765* line significantly deplete their levels

of free glucose and glycogen in contrast to *Zim53;315,* suggesting improper regulation of

glucose during acute starvation (Figure 7). Interestingly, this phenotype was not observed

in *iPLA2-VIA^{Del174}* homozygotes (Figure 7). However, we found that progeny from

Zim53;765 crossed to *iPLA2-VIA^{Del174}* also significantly deplete their available free

glucose following 12 hours of starvation (Figure 7; Table S6a,b,c). Taken together, these

results indicate shared genetic bases of *iPLA2-VIA* and *Zim53;765* in starvation sensitivity

but distinct roles in glucose metabolic effects on starvation. Notably these effects follow

a week of conditioning on high sucrose foods that may influence the time course of sugar

depletion during starvation and its impact on survival during starvation.

Discussion

In the current study we sought to identify the genetic basis of a striking difference in

starvation resistance between one particular genotype among a set of related mitonuclear

genotypes (see Figure 1). Given the central role that mitochondria play in both anabolic

and catabolic processes related to energy stores, the possibility that combinations of

mitochondrial and nuclear genes might contributed to quantitative variation for starvation

resistance warranted additional genetic analysis. This is further motivated by the highly

polygenic nature of starvation resistance and its correlation with other fitness and

metabolic traits (SERVICE AND ROSE 1985; CHIPPINDALE *et al.* 1996; HARSHMAN *et al.* 1999;

HARDY *et al.* 2018; EVERMAN *et al.* 2019). Our early crosses ruled out an effect of the mtDNA

on the strong differences in starvation between *siI;765* and *Zim53;765* which refocused

the analyses to the genetic architecture of starvation resistance between distinct nuclear

genomes. Because the nuclear genomes used to build the mitonuclear lines were part of

the well-characterized DGRP we sought to use the classical tools of the Drosophila model

and the genomic information available for the DGRP to carry out a quantitative trait locus

search between two distinct homozygous lines from the DGRP. The motivation was

twofold: the difference between the sensitive mitonuclear lines (in *DGRP-765*

backgrounds) and resistance lines (in *DGRP-315* or *-820* backgrounds) was very distinct,

and the sensitive *mitonuclear-765* lines also appeared less sensitive to the high

sugar:protein ratio in the food used prior to the starvation assay (Figure 1). Applying

forward genetic approaches to this classic quantitative genetic system offered the

opportunity to test the hypothesis that starvation sensitivity was associated with sugar

metabolism.

Polygenic basis of *Zim53;765* starvation sensitivity

The chromosomal segregation analyses and the Advanced Intercross Population (AIP)

pool-seq analyses show parallel results at a chromosomal scale and further confirm the

polygenic basis of the differences between the *DGRP-315* and *-765* genomes. Figure 2

shows a clear main effect of chromosome 3 and some additive or epistatic effects from

chromosome 2. Chromosome 3 heterozygosity was necessary to suppress *Zim53;765*

starvation sensitivity but heterozygosity of chromosome 2 and 3 together further improved

starvation resistance. Notably, chromosome 2 heterozygosity in isolation had limited

effect on suppressing *Zim53;765* starvation sensitivity (Figure 2A, B). This indicates that

the chromosome 2 locus/loci acts as a modifier that genetically interacts with the

chromosome 3 locus/loci. Importantly, *Zim53;315* has a distinct genetic basis of starvation

resistance, with chromosome3 having little effect (Figure 2A)

The pool-seq results from the AIP design (Figure 3) show corresponding patterns for

chromosomes 2 and 3: strong evidence on chromosome 3 for allele frequency differences

between early-and late dying pools of flies, but limited evidence for this on chromosome

2. Figure 2 shows a weak signal of starvation suppression by chromosome 2 for the very

last-dying flies but it is not significant. This could reflect heterozygosity of the few loci that

just pass significance on chromosome 2L in the pool-seq data (Figure 3), but this remains

speculative.

We note that the AIP analyses involved recombination between the focal line at the

extreme end of the starvation resistance spectrum (*Zim53;765*, most sensitive) and a line

more central in the distribution of starvation scores (Zim;315, with the DGRP-315 nuclear

chromosomes). Though a compelling case could be made for recombining Zim;315 with

a line at the opposite end of the distribution (most resistant; e.g., Zim;820 with the *DGRP-*

820 nuclear chromosomes), our starvation data did not show a large difference between

the *-315* and *-820* mitonuclear genotypes, but the *-315* mitonuclear genotypes had

somewhat tighter variance of starvation scores on the different sugar:protein diets, so we

chose that background (Figure 1). The AIP pool-seq data show strong allelic

differentiation across one broad peak on chromosome 3L and additional peaks on 3R,

with fewer significant peaks on chromosome 2L (4 SNPs above threshold). The resolution

of these analyses is admittedly low and resolving specific candidate loci was not an

expected goal of the AIP pool-seq. It is likely that many of these significant variants are

in linkage disequilibrium since the populations recombined over a period of only 10

generations. Nevertheless, the clustering of significant variants on chromosomes 2 and

3 are consistent with a polygenic basis for *Zim53;765* starvation sensitivity. The AIP

analyses served the important goal of confirming the results of the chromosome

segregation analyses and identifying where to focus subsequent higher resolution

recombination and deficiency mapping.

Dissecting the 3L candidate region

We sought to narrow the target region, by generating recombinants across chromomere

3 between *Zim53;765* and the multiply marked recessive stocks, as this would provide a

more localized region for deficiency analyses. We however failed to do this directly as

crossing *Zim53;765* females to males from each of the multiple marked recessive stocks

failed to yield an appreciable number of offspring. We use the *sil;765* to generate

recombinants because *sil;765* was constructed using balancer substitutions and

backcrosses to *DGRP-765* and thus likely carried *DGRP-765* chromosomes, and it

produced sufficient offspring to proceed with the remaining two recombination crosses.

Despite learning that *sil;765* was a hybridized version of *765* nuclear chromosomes, a

fact confirmed by comparing the alleles of *Zim53;765* variants to *sil;765*'s, it remained

possible that the recombinant offspring would help localize genetic effects between

chromosome arms 3L or 3R, which proved true. The 15 different deficiencies used for

mapping were chosen to span the regions of chromosome 3L that were most likely to

carry a candidate gene or genes contributing to the starvation differences between the

DGRP chromosomes present in *Zim;765* and *Zim;315*. The preponderance of the

evidence from these data supported the hypothesis that heterozygosity of the region of

3L 9812381 – 9899255 was required for suppression of *Zim53;765* starvation sensitivity,

with deficiencies BSC392, -394 and -673 being clear examples (Figures 5 and S3). Again,

however, the deficiency data included some inconsistent results. When *Zim53;765* was

crossed with Df(3L)BSC282, there was only a modest difference in the starvation

resistance of the heterozygous and homozygous progeny, and when crossed with

Df(3L)BSC283, there was no statistical difference in the starvation resistance (Fig. S3).

The two additional genes missing in Df(3L)BSC283 relative to Df(3L)BSC282 may explain

these conflicting data, but neither gene has any known function reported in Flybase

(lncRNA:CR45878, CG46387).

Complementation points to Zim53;765 iPLA2-VIA

There are over 30 genes in the candidate regions defined by 3L 9812381 – 9899255 (see

Table S1). Of all the variants in this region, the nonsynonymous mutation present in the

calcium-independent phospholipase iPLA2-VIA immediately stood out, due to the location

of the mutation in the functional domain of the enzyme, and the phenotypes previously

associated with iPLA2-VIA loss of function. Mutations in the human homolog PLA2G6

gene are characteristic of multiple autosomal recessive disease (Guo *et al.* 2018). The

amino acid mutation in the Drosophila iPLA2-VIA protein that is private to *DGRP-765* lies

in the GXGXXG consensus nucleotide binding motif similar to those found in other protein

kinases. This motif contributes to the binding of ATP to iPLA2-VIA which is essential for

its catalytic function (Ramanadham *et al.* 2015). Several studies have independently

characterized starvation sensitivity in *iPLA2-VIA* mutants (Kinghorn *et al.* 2015; Iliadi *et*

al. 2018; Lin *et al.* 2018; Mori *et al.* 2019). In each case, targeted mutations were introduced

in *iPLA2-VIA* to generate null models, with the goal of better understanding its role in

neurodegeneration. These studies identified aberrant lipid metabolism, lipid peroxidation

and mitochondrial dysfunction as potential causes of stress sensitivity and

neurodegeneration. We should not necessarily expect point mutations in *iPLA2-VIA* to

universally have the same effect as these deletion loss of function models, as mutations

in PLA2G6 have varying effects on catalytic activity (Engel *et al.* 2010). However, we can

expect that if mutation(s) in *iPLA2-VIA* do in fact underlie *Zim53;765* starvation resistance,

an *iPLA2-VIA* null should fail to complement *Zim53;765* starvation sensitivity. We showed

that two separate nulls generated by (Lin *et al.* 2018) do in fact fail to complement

Zim53;765 starvation sensitivity, indicating that natural variants in iPLA2-VIA contribute

some of the starvation sensitivity of *Zim53;765* and *DGRP-765*.

The correlation between starvation resistance and lipid content has been demonstrated

across species and as associated responses to selection for starvation resistance

(Chippindale *et al.* 1996; Harshman *et al.* 1999; Hardy *et al.* 2018; Everman *et al.* 2019).

While our current study does not have details on lipid content, there may be a link to the

glucose and glycogen utilization we report (Figure 7), and to other possible candidate

genes in the 3L 9812381 – 9899255 region: there are several insulin-like peptides (*Ilps*)

~40-100kb distal to the 3L 9812381 – 9899255 that show polymorphism among the

DGRP lines. Some phospholipases have been associated with modulation of insulin

secretion in response to glucose by releasing arachidonic acid (AA), and exogenous AA

supplementation is sufficient to promote insulin release from rat islet cells (Metz 1988a).

This insulin secretion is believed to promote phospholipid hydrolysis and accumulation of

phospholipid-derived intermediates (METZ 1988b). This turnover of phospholipids could

promote insulin release, creating an interesting dynamic between glucose and lipid

metabolism, with phospholipases at the center. The premature depletion of glucose and

glycogen in *Zim53;765* in the early stages of starvation could be one consequence of the

SNP(s) in iPLA2-VIA or an interaction with *Ilps*. Consistent with this, we found that *iPLA2-*

VIA^{Del174} fails to complement the premature glucose and glycogen depletion phenotypes

associated with *Zim53;765* (Figure 7). What is especially interesting is that the iPLA2-VIA

null itself does not display this phenotype. This raises the possibility that affected

pathways which ultimately result in starvation sensitivity differ between the deletion

mutation of *iPLA2-VIA^{Del174}* and the point mutation unique to *Zim53;765*. It will be an

interesting target of a future study to characterize the additional molecular bases

underlying the starvation sensitivity of *Zim53;765,* as the amino acid mutation in the

conserved GXGXXG motif of *iPLA2-VIA* could potentially highlight novel functions of

iPLA2-VIA.

Acknowledgments

The authors thank the HHMI Summer Fellows Julia Dewey, Tyler Devlin, Brian Franklin,

Cynthia Hale-Phillips, Matthew P. McAteer, Zemplen Pataki, Chén Yé, and Denise Yoon

who conducted the initial screen for mito-nuclear interactions affecting starvation

resistance, which provided the impetus for this work. Dr. Jim Mossman assisted with the

analyses of DGRP polymorphisms. Yevgeniy Raynes and Leah Darwin provided helpful

edits for R code.

Funding

719 The work was supported in part by Gilliam Fellowship for Advanced Study awarded to

720 SBW and DMR and NIGMS R01 2R01GM067862 awarded to DMR. DMR

721 acknowledges support of COBRE award P20GM109035 and MIRA R35GM139607

Conflicts of Interest

724 None declared.

Stock Name (mtDNA;nuclearDNA)	Mitochondrial haplotype	Nuclear haplotype	Phenotype
Zim53;765	*D. melanogaster Zim53*	DGRP-765	Starvation sensitive
sil;765	*D. simulans sil*	DGRP-765 recombinant	Starvation resistant
DGRP-765	From original DGRP-765 line	DGRP-765	Starvation sensitive
Zim53;315	*D. melanogaster Zim53*	DGRP-315	Starvation resistant

Table 1. Mitonuclear stocks used for genetic analyses in this study. A table of additional stocks used for chromosome substitution, recombination mapping and deficiency mapping are listed in supplemental file 1.

Chr.	start	end	REF	ALT	CONSEQUENCE	REFDNA	VARDNA	REFAA	VARAA	SYMBOL
chr3L	9824166	9824166	C	T	nonsynonymous	CCA	TCA	P	S	CG43897
chr3L	9833935	9833935	G	A	synonymous	AGC	AGT	S	S	RasGAP1
chr3L	9854881	9854881	C	A	nonsynonymous	GGC	GTC	G	V	iPLA2-VIA
chr3L	9880027	9880027	A	G	synonymous	TTT	TTC	F	F	Taf2
chr3L	9880564	9880564	G	T	synonymous	ATC	ATA	I	I	Taf2
chr3L	9881643	9881643	C	A	nonsynonymous	GAG	GAT	E	D	Taf2
chr3L	9884329	9884329	A	T	synonymous	ATA	ATT	I	I	CalpB
chr3L	9884605	9884605	A	G	synonymous	GGA	GGG	G	G	CalpB

Table 2. Sequence details of candidate SNPs in the genomic region identified by classical mapping approaches. See text and Fig. S4 for details.

735
736

Figure 1. Starvation time among mitonuclear genotypes on different diets. The x-axis lists the genotypes with the notation: mtDNA;nucDNA-Diet, with Diets I, IV and V being balanced, low and high sugar:yeast ratios, respectively (10g:10g, 5g:15g, 15g:5g; see Methods).

Figure 2. Chromosome 3 suppresses *Zim53;765* starvation sensitivity
(A) *DGRP-765* and *Zim53;765* flies are starvation sensitive while *Zim;315* flies are more resistant. In both *DGRP-765* and *Zim53;765*, heterozygosity for chromosome 3 (chr3) rescues the majority of starvation sensitivity with chromosome 2 heterozygotes (chr2) contributing additional resistance. In *Zim53;315* heterozygosity on chromosome 2 is the primary source of increases in starvation resistance, with little additional contribution from chromosome 3. (B) A replicate experiment with *Zim53;765* shows a similar major effect of chromosome 3 with additional effect of chromosome 2, while the *sil;765* stock with nuclear heterozygosity has increased starvation resistance and distinct chromosomal contributions. Differences in survival between pairs of genotypes were determined using a log-rank test (implemented in the R survival package) and included in Table_S2_Chromosome Segregation Mapping.

A

B

Figure 3. Significant allelic frequency differences between the tails of replicate advanced intercross populations subjected to starvation stress.
An advanced intercross population was created by crossing *Zim53;765* and *sil;315* in reciprocal. The hybrids were allowed to recombine freely for 10 generations before being subjected to starvation stress. Dead flies were collected every 6 hours and dead flies from the first 3 timepoints were combined and subjected to sequencing analysis. The same was done with the last two timepoints. In total 5 replicates AIP populations were generated and sequenced. (A) Consistent differences in allele frequencies across replicates were determined using the Cochran-Mantel-Haenszel (CMH) test and Popoolation2. The y-axis is the negative log-transformed, Bonferroni corrected p-value of each SNP (represented by each circle). The x-axis is the position of each SNP along the indicated chromosome. Blue and red lines highlight corrected p-values above 1×10^{-5} and 1×10^{-6} respectively. (B) F_{ST} between pools for each site averaged across replicate populations. The location of elevated FST values is generally concordant with significant CMH test statistics.

Figure 4. Recombination mapping localizes the majority of *Zim53;765* starvation sensitivity to chromosome 3L
(A) Cartoon representation of recombination breakpoints for each recombinant class generated through meiotic recombination of the chromosome 3 mapping line. (B) Survival of female offspring produced by crossing each recombinant class to *Zim53;765* homozygotes or heterozygotes. Differences in survival were determined using a log-rank test (implemented in the R survival package; *** = p , 0.001, * = p<0.05) and included in Table_S3_Meiotic Mapping Survival.

Figure 5. Deficiency mapping identifies the 3L: 9812381 – 9899255 region as a component of *Zim53;765* starvation sensitivity

(A) Survival of offspring from the cross between female *Zim53;765* and male deficiency stocks following starvation stress. Heterozygous refers to progeny that inherited the marked balancer chromosome, hemizygous refers to progeny that inherited the deficient chromosome. Line numbers correspond to Bloomington Stock Center names (see table S1). Differences in survival were determined using a log-rank test (implemented in the R survival package) and included in Table S4_Deficiency mapping: summary and statistics tabs. (B) Linear representation of hemizygous regions of respective deficiency lines. Highlighted region is common to the three deficiency regions which failed to suppress *Zim53;765* starvation sensitivity. See Fig. S3 for additional crosses

813
Figure 6. *Zim53;765* fails to complement iPLA2-VIA null lines.
Survival of female offspring from indicated crosses following starvation stress. Each
panel represents the male parent, and colored lines identify the female parent. *Zim;765*
is starvation sensitive as a homozygote and as a heterozygote over two different *iPLA2-*
VIA deficiency null alleles. Differences in survival were determined using a log-rank test
(implemented in the R survival package) and included in Table S5_iPLA-VIA
complementation).

823
824

826
Figure 7. iPLA2-VIA[Delta174] fails to complement glucose and glycogen
depletion phenotype of *Zim53;765*
The free glucose and total glycogen levels were measured for *Zim53;765*, *Zim53;315*,
iPLA2-VIA[Delta174], *Zim53;315* crossed to iPLA2-VIA[Delta174] and *Zim53;765*
crossed to iPLA2-VIA[Delta174]. Values in the un-starved (control) was compared to the
values in the starved samples (12 hours) for 3 replicates of 6 flies using a student's t-
test. Significant differences are indicated by * <0.05, ** <0.01.
834
835

837
838
839 **Supplemental Figure S1**. Recombination scheme for meiotic mapping. The scheme
840 includes three sequential crosses using the focal lines described in the text and two
841 multiply marked stocks to resolve recombination breakpoints. Note that *Zim53;765* is
842 homozygous for DGRP-765 chromosomes and is starvation sensitive while *sil;765* is
843 starvation resistant but has nuclear heterozygosity is starvation resistant (see Figure 1).
844

845
Supplemental Figure S2. *sil;765* is a hybrid of DGRP-765
(A) Survival of female offspring of *Zim53;765,* and *sil;765* crossed in reciprocal. There is
no statistical difference in survival based on a log-rank test for significance. (B and C)
Genotype of *sil;765* at homozygous variant sites in RNA sequencing reads from (C)
Zim53;765 and (D) *Zim53;315* Green indicates that *sil;765* was heterozygous for both
the variant and the reference allele, blue indicates that *sil;765* was homozygous for the
reference allele.

A

B

854
Supplemental Figure S3
(A) Linear representation of deficient region of respective deficient lines in region
identified in meiotic recombination mapping. (B) Survival of offspring from the cross
between *Zim53;765* and the respective deficient male following starvation stress.
Differences in survival were determined using a log-rank test (implemented in the R
survival package) and included in TableS4_Deficiency mapping. Heterozygous refers to
progeny that inherited the balancer chromosome, hemizygous refers to progeny that
inherited the deficient chromosome. Line numbers correspond to Bloomington Stock
Center names.
864

865
866

867
Supplemental Figure S4. Frequency of DGRP-765 variants among the Drosophila
Genetic Reference Panel
(A) Allele frequency of *DGRP-765* variants in the 3L candidate region among the lines
of the Drosophila Genetic Reference Panel. Variants found below the red line are
present in less than ten percent of the lines. (B) Proportions of low frequency (<10%)
variants in different functional categories for in the 3L candidate region present in
DGRP-765.
875

References

Adzhubei, I. A., S. Schmidt, L. Peshkin, V. E. Ramensky, A. Gerasimova *et al.*, 2010 A
method and server for predicting damaging missense mutations. Nat Methods 7**:**
248-249.
Ali, T., G. Kokotos, V. Magrioti, R. N. Bone, J. A. Mobley *et al.*, 2013 Characterization of
FKGK18 as inhibitor of group VIA Ca2+-independent phospholipase A2
(iPLA2beta): candidate drug for preventing beta-cell apoptosis and diabetes. PLoS
One 8**:** e71748.
Ayroles, J. F., M. A. Carbone, E. A. Stone, K. W. Jordan, R. F. Lyman *et al.*, 2009 Systems
genetics of complex traits in Drosophila melanogaster. Nat Genet 41**:** 299-307.
Brown, E. B., M. E. Slocumb, M. Szuperak, A. Kerbs, A. G. Gibbs *et al.*, 2019 Starvation
resistance is associated with developmentally specified changes in sleep, feeding
and metabolic rate. J Exp Biol 222.
Chippindale, A. K., T. J. F. Chu and M. R. Rose, 1996 Complex trade-offs and the
evolution of starvation resistance in Drosophila melanogaster. Evolution 50**:** 753-
766.
Engel, L. A., Z. Jing, D. E. O'Brien, M. Sun and P. T. Kotzbauer, 2010 Catalytic function
of PLA2G6 is impaired by mutations associated with infantile neuroaxonal
dystrophy but not dystonia-parkinsonism. PLoS One 5**:** e12897.
Erkosar, B., C. Dupuis, L. Savary and T. J. Kawecki, 2025 Shared genetic architecture
links energy metabolism, behavior and starvation resistance along a power-
endurance axis. Evol Lett 9**:** 150-162.
Everman, E. R., C. L. McNeil, J. L. Hackett, C. L. Bain and S. J. Macdonald, 2019
Dissection of Complex, Fitness-Related Traits in Multiple Drosophila Mapping
Populations Offers Insight into the Genetic Control of Stress Resistance. Genetics
211**:** 1449-1467.
Everman, E. R., and T. J. Morgan, 2018 Antagonistic pleiotropy and mutation
accumulation contribute to age-related decline in stress response. Evolution 72**:**
303-317.
Feder, A. F., D. A. Petrov and A. O. Bergland, 2012 LDx: estimation of linkage
disequilibrium from high-throughput pooled resequencing data. PLoS One 7**:**
e48588.
Guo, Y. P., B. S. Tang and J. F. Guo, 2018 PLA2G6-Associated Neurodegeneration
(PLAN): Review of Clinical Phenotypes and Genotypes. Front Neurol 9**:** 1100.
Harbison, S. T., A. H. Yamamoto, J. J. Fanara, K. K. Norga and T. F. Mackay, 2004
Quantitative trait loci affecting starvation resistance in Drosophila melanogaster.
Genetics 166**:** 1807-1823.
Hardy, C. M., M. K. Burke, L. J. Everett, M. V. Han, K. M. Lantz *et al.*, 2018 Genome-
Wide Analysis of Starvation-Selected Drosophila melanogaster-A Genetic Model
of Obesity. Mol Biol Evol 35**:** 50-65.
Harshman, L. G., A. A. Hoffmann and A. G. Clark, 1999 Selection for starvation resistance
in *Drosophila melanogaster*: physiological correlates, enzyme activities and
multiple stress responses. Journal of Evolutionary Biology 12**:** 370-379.

Hill, G. E., 2015 Mitonuclear Ecology. Mol Biol Evol 32**:** 1917-1927.

Huang, W., S. Richards, M. A. Carbone, D. Zhu, R. R. Anholt *et al.*, 2012 Epistasis dominates the genetic architecture of Drosophila quantitative traits. Proc Natl Acad Sci U S A 109**:** 15553-15559.

Iliadi, K. G., O. B. Gluscencova, N. Iliadi and G. L. Boulianne, 2018 Mutations in the Drosophila homolog of human PLA2G6 give rise to age-dependent loss of psychomotor activity and neurodegeneration. Sci Rep 8**:** 2939.

Kinghorn, K. J., and J. I. Castillo-Quan, 2016 Mitochondrial dysfunction and defects in lipid homeostasis as therapeutic targets in neurodegeneration with brain iron accumulation. Rare Dis 4**:** e1128616.

Kinghorn, K. J., J. I. Castillo-Quan, F. Bartolome, P. R. Angelova, L. Li *et al.*, 2015 Loss of PLA2G6 leads to elevated mitochondrial lipid peroxidation and mitochondrial dysfunction. Brain 138**:** 1801-1816.

Kofler, R., R. V. Pandey and C. Schlotterer, 2011 PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics 27**:** 3435-3436.

Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25**:** 1754-1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence Alignment/Map format and SAMtools. Bioinformatics 25**:** 2078-2079.

Lin, G., P. T. Lee, K. Chen, D. Mao, K. L. Tan *et al.*, 2018 Phospholipase PLA2G6, a Parkinsonism-Associated Gene, Affects Vps26 and Vps35, Retromer Function, and Ceramide Levels, Similar to alpha-Synuclein Gain. Cell Metab 28**:** 605-618 e606.

Mackay, T. F., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The Drosophila melanogaster Genetic Reference Panel. Nature 482**:** 173-178.

Metz, S. A., 1988a Exogenous arachidonic acid promotes insulin release from intact or permeabilized rat islets by dual mechanisms. Putative activation of Ca2+ mobilization and protein kinase C. Diabetes 37**:** 1453-1469.

Metz, S. A., 1988b Membrane Phospholipid Turnover as an Intermediary Step in Insulin-Secretion - Putative Roles of Phospholipases in Cell Signaling. American Journal of Medicine 85**:** 9-21.

Mori, A., T. Hatano, T. Inoshita, K. Shiba-Fukushima, T. Koinuma *et al.*, 2019 Parkinson's disease-associated iPLA2-VIA/PLA2G6 regulates neuronal functions and alpha-synuclein stability through membrane remodeling. Proc Natl Acad Sci U S A 116**:** 20689-20699.

Mossman, J. A., L. M. Biancani, C. T. Zhu and D. M. Rand, 2016 Mitonuclear Epistasis for Development Time and Its Modification by Diet in Drosophila. Genetics 203**:** 463-484.

Ramanadham, S., T. Ali, J. W. Ashley, R. N. Bone, W. D. Hancock *et al.*, 2015 Calcium-independent phospholipases A2 and their roles in biological processes and diseases. J Lipid Res 56**:** 1643-1668.

Rand, D. M., R. A. Haney and A. J. Fry, 2004 Cytonuclear coevolution: the genomics of cooperation. Trends Ecol Evol 19**:** 645-653.

Service, P. M., and M. R. Rose, 1985 Genetic Covariation among Life-History Components - the Effect of Novel Environments. Evolution 39**:** 943-945.

Song, K., X. Zhang, C. Zhao, N. T. Ang and Z. A. Ma, 2005 Inhibition of Ca2+-independent
phospholipase A2 results in insufficient insulin secretion and impaired glucose
tolerance. Mol Endocrinol 19**:** 504-515.
Sorensen, J. G., M. M. Nielsen and V. Loeschcke, 2007 Gene expression profile analysis
of *Drosophila melanogaster* selected for resistance to environmental stressors.
Journal of Evolutionary Biology 20**:** 1624-1636.
Tennessen, J. M., W. E. Barry, J. Cox and C. S. Thummel, 2014 Methods for studying
metabolism in Drosophila. Methods 68**:** 105-115.
Therneau, T. M., and P. M. Grambsch, 2000 *Modeling survival data : extending the Cox*
model. Springer, New York.
Uma Naresh, N., S. Kim, T. Shpilka, Q. Yang, Y. Du *et al.*, 2022 Mitochondrial genome
recovery by ATFS-1 is essential for development after starvation. Cell Rep 41**:**
111875.
Zhu, C. T., P. Ingelmo and D. M. Rand, 2014 GxGxE for lifespan in Drosophila:
mitochondrial, nuclear, and dietary interactions that modify longevity. PLoS Genet
10**:** e1004354.
984