Insights into the RNA virome of the corn leafhopper *Dalbulus maidis*, a major emergent threat of Maize in Latin America

Humberto Debat[1,2*], Esteban Simon Farrher[3*], Nicolas Bejerman[1,2]

[1]*Instituto de Patología Vegetal – Centro de Investigaciones Agropecuarias – Instituto Nacional de Tecnología Agropecuaria (IPAVE-CIAP-INTA), Camino 60 Cuadras Km 5,5 (X5020ICA), Córdoba, Argentina.*

[2]*Unidad de Fitopatología y Modelización Agrícola – Consejo Nacional de Investigaciones Científicas y Técnicas (UFYMA-CONICET)., Camino 60 Cuadras Km 5,5 (X5020ICA), Córdoba, Argentina.*

[3]*Facultad de Ciencias Químicas, Universidad Nacional de Córdoba, Córdoba, 5000, Argentina.*

* Both authors contributed equally to this manuscript

Corresponding authors: Debat Humberto, debat.humberto@inta.gob.ar 0000-0003-3056-3739; Esteban Farrher, esteban.farrher@mi.unc.edu.ar; Nicolas Bejerman, bejerman.nicolas@inta.gob.ar 0000-0002-7851-3506.

Abstract

The maize leafhopper (*Dalbulus maidis*) is a significant threat to maize crops in tropical and subtropical regions, causing extensive economic losses. While its ecological interactions and control strategies are well-studied, its associated viral diversity remains largely unexplored. Here, we employ high-throughput sequencing data mining to comprehensively characterize the *D. maidis* RNA virome, revealing novel and diverse RNA viruses. We characterized six new viral members belonging to distinct families, with evolutionary cues of beny-like viruses (*Benyviridae*), bunya-like viruses (*Bunyaviridae*) iflaviruses (*Iflaviridae*), orthomyxo-like viruses (*Orthomyxoviridae*), and rhabdoviruses (*Rhabdoviridae*). Phylogenetic analysis of the iflaviruses places them within the genus *Iflavirus* in affinity with other leafhoppers-associated iflaviruses. The five-segmented and highly divergent orthomyxo-like virus showed a relationship with other insect associated orthomyxo-like viruses. The rhabdo virus is related with a leafhopper associated rhabdo-like virus. Furthermore, the beny-like virus belonged to a cluster of insect-associated beny-like viruses, while the bi-segmented bunya-like virus was related with other bi-segmented insect-associated bunya-like viruses. These results highlight the existence of a complex virome linked to *D. maidis* and paves the way for future studies investigating the ecological roles, evolutionary dynamics, and potential biocontrol applications of these viruses on the *D. maidis*—maize pathosystem.

Key words: RNA virome; *Dalbulus maidis*; maize leafhopper; data mining

1. Introduction

The maize leafhopper *Dalbulus maidis* (DeLong & Wolcott) (Hemiptera: Auchenorrhyncha) is a significant threat to maize crops causing extensive economic losses [1]. This insect pest is widely distributed in the Americas, from the southern United States to northern Argentina [2]. In addition to the direct damage caused by its feeding, this insect also serves as the vector for the mollicute *Spiroplasma kunkelii* Whitcomb, the causal agent of Corn Stunt disease, and the maize bushy stunt (MBS) phytoplasma [1]. *D. maidis* is also the vector of two viruses: maize rayado fino virus (MRFV) [1], and maize striate mosaic virus (MSV) [3]. While its ecological interactions [4, 5] and control strategies, such as germplasm resistance, insecticide sprays and some cultural practices are well-studied [5, 6], its associated viral diversity remains largely unexplored.

Viruses are the most abundant microbes on the planet [7-9] and RNA viruses dominate the eukaryotic virome [10, 11]. Hundreds of viruses have been identified in recent years in arthropods [10-15], illustrating that invertebrates usually have a greater and more diverse virome than vertebrates [16]. Nevertheless, over the last years there was a rapid increase of vertebrate-associated virus entries in the NCBI-GenBank, while low numbers of invertebrate-associated virus were identified and uploaded into public repositories [16]. The availability of a great amount of RNA-seq data (metatrancriptomics) in public databases has revolutionized the field of virus discovery, fostering the identification of novel viruses even without a previous knowledge of their genome sequence [7, 17, 18]. The comprehensive analysis of the RNA virosphere led to the identification of a great number of novel viruses thus shedding light not only on the huge diversity of RNA viruses in arthropods, but also on their evolutionary histories as well as on their genomic organization complexity [11, 12, 15, 16, 19, 20]. Furthermore, this advances on the knowledge of the virosphere are leading to opportunities and challenge on virus discovery [21] and to develop a new, genome-based virus taxonomy [22-26].

This paper reports the analysis of 23 publicly available *D. maidis* transcriptome datasets from USA, Brazil and Argentina, that resulted in the discovery of six novel RNA viruses associated to this insect. To our knowledge, this is the first report of insect viruses associated to this economically important insect, revealing that the *D. maidis* RNA virome is rich and diverse.

2. Material and Methods

2.1 Exploration of the D. maidis virome from public RNA-seq datasets, sequence assembly and virus identification

We analyzed all *D. maidis* Sequence Read Archives (SRA) deposited in the NCBI (https://www.ncbi.nlm.nih.gov/sra). The raw nucleotide sequence reads from each SRA experiment were downloaded from their associated NCBI BioProjects (Table S1). The datasets were pre-processed by trimming and filtering with the Trimmomatic v0.40 tool as implemented in http://www.usadellab.org/cms/?page=trimmomatic with standard parameters. The resulting reads were assembled *de novo* with rnaSPAdes using standard parameters on the Galaxy server (https://usegalaxy.org/). The transcripts obtained from *de novo* transcriptome assembly were subjected to bulk local BLASTX searches (E-value < 1e^{-5}) against a curated database of the publicly available reference viral proteins. The resulting viral sequence hits of each dataset were explored in detail. Tentative virus-like contigs were curated (extended and/or confirmed) by iterative mapping of each SRA library's filtered reads. This strategy was used to extract a subset of reads related to the query contig, used the retrieved reads from each mapping to extend the contig and then repeat the process iteratively using as query the extended sequence [27]. The extended and polished transcripts were reassembled using Geneious v8.1.9 (Biomatters Ltd.) alignment tool with high sensitivity parameters.

2.2 Bioinformatics tools and analyses

2.2.1 Sequence analyses

Open reading frames (ORFs) were predicted with ORFfinder (minimal ORF length 120 nt, genetic code 1, https://www.ncbi.nlm.nih.gov/orffinder/), functional domains and architecture of translated gene products were determined using InterPro (https://www.ebi.ac.uk/interpro/search/sequence-search) and the NCBI Conserved domain database - CDD v3.20 (https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) with e-value = 0.1. Further, HHPred and HHBlits as implemented in https://toolkit.tuebingen.mpg.de/#/tools/ were used to complement annotation of divergent predicted proteins by hidden Markov models. Transmembrane domains were predicted using the TMHMM version 2.0 tool (http://www.cbs.dtu.dk/services/TMHMM/) and signal peptides were predicted using the SignalP version 6.0 tool (https://services.healthtech.dtu.dk/services/SignalP-6.0/), while nuclear localization signals (NLS) were predicted using cNLS Mapper tool (https://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi). The presence of gene junction sequences flanking ORFs was also included as a criterion to determine the potential coding

sequences. The predicted proteins were then subjected to NCBI-BLASTP searches against the non-redundant protein sequences (nr) database.

2.2.2 Phylogenetic analysis

Phylogenetic analysis based on the RdRp proteins, was done using MAFFT 7.505 with multiple aa sequence alignments using FFT-NS-i as the best-fit model. The aligned aa sequences were used as the input in Mega11 software [28] to generate phylogenetic trees by the maximum-likelihood method using the best-fit substitution model. Local support values were computed using bootstraps with 1,000 replicates.

3. Results and Discussion

RNA viruses associated with *D. maidis*

Insects are the most abundant group of animals on earth [29]. These animals have a rich and diverse virome which is being unraveled in the later years using high-throughput sequencing [10, 11, 14, 15, 30]; however, only a small portion of the insect viromes diversity has been described so far [31]. Several viruses were identified in some leafhoppers, mainly iflaviruses, [14]. In addition, a rhabdovirus [32], an orthomyxo-like virus [12], a chuvirus [33], a reovirus [34], a nido-like virus [35], and nine RNA viruses, belonging to seven viral clades (Picorna-Calici, Permutotetra, Bunya-Arena, Reo, Partiti-Picobirna, Luteo-Sobemo and Toti-Chryso) [36] were also identified, which illustrates how rich and diverse could be the virome associated to leafhoppers. There are more than 20,000 described species of leafhoppers (*Cicadellidae*) which illustrate that the handful of viruses linked to this invertebrate is scarce, poorly miss-represents the diversity of these invertebrates, andeven important agronomic pests as the maize leafhopper virome are still unexplored. Thus, here we analyzed transcriptome datasets of maize leafhopper collected in Argentina, USA and Brazil. The *de novo* assembling of those 23 libraries followed by similarity analyses resulted in the robust identification of six novel highly divergent RNA viruses, including a beny-like virus, a bunya-like virus, two iflaviruses, a orthomyxo-like virus and a rhabdovirus, which were tentatively named Chicharrita del maiz beny-like virus (ChMBLV), Chicharrita del maiz bunya-like virus (ChMBV), Chicharrita del maiz iflavirus 1 (ChMIfV1), Chicharrita del maiz iflavirus 2 (ChMIfV2), Chicharrita del maiz orthomixo-like virus (ChMOMLV) and Chicharrita del maiz rhabdoirus (ChMRV), respectively. This first glimpse of the virus repertoire reveals how rich and diverse is the virome associated to the maize leafhopper. We assembled the full-length coding region sequence for all viruses, but the rhabdovirus, which its nearly complete coding region was able to be assembled. Revealing the virus diversity in this

organism is crucial to further exploit the identified viruses as specific and targeted tools for the biological control of insect populations.

Molecular and phylogenetic characterization of the newly identified maize leafhopper Beny-like viruses

The *Benyviridae* is a family with members that have positive-sense, single stranded RNA viruses with only one genus, the *Benyvirus*, recognized by the ICTV [37]. Benyviruses are plant-infecting viruses with a multipartite genome [37]. Nevertheless, metagenomic studies as well as the metatrancriptomic analysis of publicly data deposited in NCBI resulted in the discovery of many benyviruses related sequences, that resemble to newly unassigned *Benyviridae* members, that likely infect a wide range of non-plant hosts, such as insects and fungi [38-41]. Moreover, the insect-associated beny-like viruses genomes are monopartite which is contrasting to the multipartite genomes displayed by the plant-associated counterparts. Here, a novel beny-like virus was identified and named as Chicharrita del maiz beny-like virus (ChMBLV), which was only detected in one transcriptome datasets of maize leafhopper, that was collected in USA (Table S1). The positive-sense, single stranded RNA ChMBLV sequence is 6,887 nucleotides (nt) long and has two ORFs encoding a non-structural polyprotein of 2,003 amino acids (aa) and a putative structural protein of 224aa (Fig. 1A) (Table 1). This genomic organization is similar to that reported for the beetle associated viruses Diabrotica undecimpunctata virus 2 (DuV2) [42] and Guiyang benyvirus 1 (GuBV1) (Feng, unpublished). ChMBLV non-structural polyprotein showed the highest BlastP similarity with that one encoded by DuV2, with 46.61% identity (Table 1); therefore, based on genetic distance ChMBLV is a new virus distantly related to other NCBI database entries of insect viruses. Two conserved domains, the Viral_helicase1, and the ps-ssRNAv_RdRp-like, were identified in the polyprotein at aa positions 617-848 and 1,673-1,857, respectively. Similar conserved domains were found in those non-structural polyproteins encoded by related beny-like viruses [38, 39, 41, 42] ChMBLV structural protein showed the highest BlastP similarity with that one encoded by DuV2, with 29.75% identity (Table 1). This protein encoded by the ChMBLV ORF2 is likely a coat protein, since the TMV_coat super family conserved domain was identified at aa position 108-180. A similar conserved domain was also identified in the ORF2 encoded proteins of DuV2 [42], and GuBV1 (Feng, unpublished).

Phylogenetically analysis of the non-structural polyprotein amino acid sequence placed ChMBLV in a clade containing several insect-associated beny-like viruses, where ChMBlV was clustered together with the beetle Lepidiota negatoria beny-like virus within the insect clade of beny-like viruses, which is separated from the clades composed of fungi-

associated beny-like viruses and plant-associated ones (Fig. 1B). Thus, we propose to create a new genus, to be named as "*Insebenyvirus*" within the family *Benyviridae*, to accommodate the insect-associated beny-like viruses.

To our knowledge, ChMBLV is the first reported beny-like virus associated to leafhoppers.

Molecular and phylogenetic characterization of the newly identified maize leafhopper Bunya-like virus

Bunyavirales is a viral order including several families of negative-sense single-stranded RNA (-ssRNA) viruses with a multi segmented genome [43]. Bunyavirids are highly diverse and have a wide host range and a considerable genome complexity. Many proposed members of this family have been discovered in several insect species [12]. Here, a novel bunya-like virus was identified and named as Chicharrita del maiz bunyan-like virus (ChMBV), which was detected in the three transcriptome datasets of maize leafhopper from Brazil (Table S1). The full-length ChMBV coding sequence comprised two segments (L and S) of single-stranded, negative-sense RNA of 6,651nt and 1,281nt, respectively. Segment L encodes a putative RdRp protein of 2,166 aa, and the segment S encodes a putative nucleoprotein (NP) of 327 aa (Fig. 2A) (Table 1).

As expected, the structural NP encoding segment was the most abundant one in the three libraries from Brazil, whereas the segment encoding the RdRp was the one that accumulated less on those datasets where ChMBV was identified (Table S1). ChMBV present low homology to other bunyavirids (Table 1). The RdRp protein shows the highest BlastP similarity with the leafhopper associated Hubei insect virus 1 (HbIV1) RdRp, with 49.84% identity (Table 1), and a Bunya_RdRp super family conserved domain was identified in its sequence at aa positions 603-1,303. The NP protein shows the highest BlastP similarity with the HbIV1 NP, with 38.01% identity (Table 1), and a Tenui_N super family conserved domain has been identified in its sequence at aa positions 40-191.

The phylogenetic tree based on ChMBV RdRp encoded protein, showed that this virus is evolutionary related with leafhopper associated bunya-like viruses. ChMBV clustered with HbIV1, Hemipteran phenui-related virus OKIAV285 and the rice green leafhopper associated virus Hangzhou nephotettix cincticeps phenuivirus 1, and this cluster is linked with the planthopper-associated Sanya nilaparvata lugens phenuivirus 1 (Fig.2B). Bunyavirids have different number of genome segments; however, those viruses phylogenetically related with ChMBV, are all bisegmented RNA viruses [11]. Thus, these leafhopper-associated bunya-like viruses appear to be a new cluster of bisegmented RNA viruses that could be classified into a new genus within the *Bunyavirales* order.

Molecular and phylogenetic characterization of the newly identified maize leafhopper Iflaviruses

Iflaviridae is a family composed of only one genus, the *Iflavirus,* of positive-sense single-stranded RNA (-ssRNA) viruses with a monopartite genome, that contains an ORF encoding a single polyprotein of approximately 3,000 aa. All reported iflaviruses are associated to arthropod hosts, mainly insects [44, 45], and are among the most common viruses identified in different species of arthropods [46]. Here, two iflaviruses were identified and named as Chicharrita del maiz iflavirus virus 1 (ChMIfV1) and Chicharrita del maiz iflavirus 2 (ChMIfV2). ChMIfV1 was detected in five transcriptome datasets of the maize leafhopper (one from Argentina, another from USA and three from Brazil), while ChMIfV2 was detected in 18 transcriptome datasets, all from USA (Table S1). The iflaviruses were the most prevalent and abundant among the six viruses identified in this study (Table S1).

The positive-sense single-stranded RNA ChMIfV1 genome is 11,104 nt in length, while ChMIfV2 genome is 10,318 nt in length (Fig. 3A) (Table 1) and are A/U rich, similar to other iflaviruses [47]. Both viruses have a single ORF which encodes a polyprotein of 3,306aa and 3,102aa, respectively (Fig. 3A) (Table 1).

ChMIfV1 and ChMuIfV2 polyproteins share only a 26.3% aa identity between them, and ChMIfV1 polyprotein showed the highest BlastP similarity with that one encoded by the rice green leafhopper associated virus Nephotettix cincticeps positive-stranded RNA virus 1 (NcPSRV1), with 64.03% identity (Table 1); while ChMIfV2 polyprotein showed the highest BlastP similarity with that one encoded by the American grapevine leafhopper associated virus Scaphoideus titanus iflavirus 1 (StIfV1) with 53.04% identity (Table 1). The prediction of the conserved domains present in both ChMIfV1 and ChMuIfV2 polyproteins, showed that both viruses have the typical iflavirus gene organization. The structural domains are located at the N-terminal region of the polyprotein, while the non-structural domains are located downstream in the C-terminal region [44]. Two rhinovirus-like capsid domains (Rhv-like, aa 338-518 and aa 597-785 in ChMIfV1, aa 327-508 and aa 602-786 in ChMIfV2) and a cricket paralysis virus–like capsid domain (CRPV_capsid super family, aa 997-1,148 im ChMIfV1 and aa 1,057-1,209 in ChMIfV2) were identified in the N-terminal region of the ChMIfV1 and ChMIfV2 polyproteins, while an helicase (RNA_Helicase, aa 1,550-1,656 in ChMIfV1 and aa 1,552-1,658 in ChMIfV2), a Picornavirales 3C/3C-like protease (3C-Pro, aa 2,529-2,741 in ChMIfV1 and aa 2,301-2,530 in ChMIfV2) and RNA-directed RNA polymerase (RdRp, aa 2,952-3,256 in ChMIfV1 and aa 2,738-3,055 in ChMIfV2) conserved domains were identified in the C-terminal region of the ChMIfV1 and ChMuIfV2 polyproteins (Fig. 3A).

Phylogenetic analysis based on the RdRp conserved region shows that ChMlfV1 and ChMlfV2 clustered within the *Iflaviridae* members with other leafhopper-associated iflaviruses; where both viruses were placed in different clades as predicted by the low identity observed between them (Fig. 3B). ChMlfV1clustered with NcPSRV1 and the rice pest "zig-zag leafhopper" associated virus Hangzhou recilia dorsalis iflavirus 1 (Fig. 3B); while ChMlfV2 clustered with the American grapevine leafhopper StlfV1 and StlfV2, Hangzhou recilia dorsalis iflavirus 2 and the rice green leafhopper associated virus Congyang nephotettix cincticeps iflavirus 1 (Fig. 3B). All in all, the branching of both viruses withing the group of Iflavirus members indicates that both viruses should be included in the family *Iflaviridae* (Fig.3B).

Iflavirus infections are usually asymptomatic; however, some ifllaviruses have been reported to be harmful to the host insects leading to morphological deformities, behavioral change, and death [48-50]. Moreover, recently the the armyworm Spodoptera exigua iflavirus 1 (SeIV1) was shown to enhance the susceptibility of the armyworm to the disease caused by Spodoptera exigua multiple nucleopolyhedrovirus (SeMNPV) [51], as well as its susceptibility to Bacillus thuringiensis-based insecticide formulations and the parasitism by the wasp Hyposoter didymator [52]. Moreover, iflaviruses have been proposed for VIGS to regulate the expression of insect metabolic genes [53]. Thus, it would be critical to further explore if the iflaviruses identified in this study are pathogenic to its insect host for their potential deployment as tools for biocontrol agents.

Molecular and phylogenetic characterization of the newly identified maize leafhopper Orthomyxo-like virus

Orthomyxoviridae is a family of negative-sense single-stranded RNA (-ssRNA) viruses with a multi segmented genome [54] . Members belonging to the *Orthomyxoviridae* family are highly diverse, with a wide host range and complex genomes. Many members of this family have been discovered in various insect species [12]. Here, a novel orthomyxo-like virus was identified and named as Chicharrita del maiz orthomyxo -like virus (ChMOMLV), which was detected in three transcriptome datasets of maize leafhopper, one from USA and two from Brazil. The library from USA showed the highest abundance of virus reads, and among segments, as expected, the ones encoding for the structural nucleoprotein (NP) and hemagglutinin protein (HA) showed the highest RNA titers in all ChMOMLV positive libraries (Table S1). The ChMOMLV genome sequence comprised 5 segments of single-stranded, negative-sense RNAs of 2,557nt, 2,460nt, 2,348nt, 1,817nt and 1,681nt (Fig. 4A). RNA1 and RNA2 encode polymerase PB1 and PB2 proteins of 794 and 777 aa, respectively, and RNA3 encodes a putative polymerase PA of 739aa. RNA4 encodes a putative NP of 572aa, while

RNA5 encodes a putative HA of 503 aa (Fig. 4A) (Table 1). ChMOMLV presents low homology to other orthomixo-like virus and members belonging to the *Orthomyxoviridae* family (Table 1). PB1 protein shows the highest BlastP similarity with the brown marmorated stink bug associated virus Halyomorpha halys orthomyxo-like virus 1 (HhOlV1) PB1, with 49.41% identity, and a Flu_PB1 super family conserved domain has been identified in its sequence at aa positions 41-747. PB2 protein shows the highest BlastP similarity with HhOlV1-PB2, with 26.32% identity and a Flu_PB2 super family conserved domain has been identified in its sequence at aa positions 15-409. The PA protein shows the highest BlastP similarity with the Bat faecal associated orthomyxo-like virus 1 PA, with 34.59% identity and a Flu_PA super family conserved domain was detected in its sequence at aa positions 126-731. The putative NP protein shows the highest BlastP similarity with the Hemipteran orthomyxo-related virus OKIAV183 NP, with 29.87% identity with a Flu_NP super family conserved domain in its sequence at aa positions 340-553. The HA protein shows the highest BlastP similarity with the parasitoid wasp associated virus Cotesiavirus orthomyxo HA, with 24.56% identity and an envelope glycoprotein conserved domain at aa positions 295-495. The phylogenetic tree based on ChMOMLV PB1 encoded protein, showed that this virus is clustered with the insect-associated orthomyxo-like viruses HhOlV1 and Hemipteran orthomyxo-related virus OKIAV191 and this cluster its related with a clade formed by putative quaranjavirus (Fig.4B). Orthomixovirids have different number of genome segments; however, those viruses phylogenetically related with ChMOMLV also have 5 RNA segments [12, 55]. Viruses belonging to the genus *Quaranjavirus* are known as argasid tick-borne viruses and have genomes with 5 to 6 RNA segments [56]. Many quaranji-like viruses were recently described in several insects; therefore, the diversity of quaranjaviruses might be greatly underestimated in arthropods other than the previously described hematophagous insects [39]. Thus, future studies should asses whether this group of unclassified quaranja-like viruses including ChMOMLV could be classified within the genus *Quaranjavirus*, or if would be more appropriate to encompass these fifth segmented viruses in a novel genus within the family *Orthomyxoviridae*.

To our knowledge, ChMOMLV is the first reported orthomyxo-like virus associated to leafhoppers.

Rhabdovirus

Rhabdoviridae is a family of negative-sense single-stranded RNA (-ssRNA) viruses with a monopartite genome as a hallmark of almost all rhabdoviruses. However, plant-associated rhabdoviruses with bi-segmented and tri-segmented genomes have been described [27, 57] . Members belonging to the *Rhabdoviridae* family are highly diverse and have a wide host

range and a great genome complexity and many members of this family have been discovered in many insect species, some of which may serve as unique hosts or may act as biological vectors for transmission to other animals or plants [12, 57, 58]. Here, a novel rhabdo-like virus was identified and named as Chicharrita del maiz rhabdovirus (ChMRV), which was only detected in the transcriptome dataset of maize leafhopper collected in Argentina (Table S1). The partial negative-sense single-stranded RNA ChMRV genome is 12,271nt in length (Table 1, Fig. 5A). ChMRV genome has the five canonical rhabdoviral structural protein genes nucleocapsid (N), phosphoprotein (P), matrix protein (M), glycoprotein (G) and the polymerase (L), as is the typical architecture within the family *Rhabdoviridae* [57, 58]. Two additional genes, named as U1 and U2, were detected between the G and L genes. Thus, the genomic organization of ChMRV is 3'-N-P-M-G-U1-U2-L-5´ (Fig. 5A), which is similar to the one reported for a few insect-associated rhabdoviruses, such as the mosquito associated Xiangyun mono-chu-like virus 11 [57] and the leafhopper associated viruses Guiyang nephotettix cincticeps rhabdovirus 1 (GuNCRV1) (Feng, unpublished).

The ORF1, which is 1,365nt in length, encodes a putative N protein of 454aa (Table 1). One nuclear localization signal (NLS) was identified at aa positions 405–440. A NLS was also identified in the N protein encoded by the leafhopper associated Nephotettix cincticeps negative-stranded RNA virus-1 (NcNSRV1) [32]. The ORF2, which is 936nt in length, encodes a putative P protein of 311aa (Table 1). One NLS was identified at aa positions 103–132. A NLS was also identified in the P protein encoded by NcNSRV1 and Spodoptera frugiperda rhabdovirus [32, 59]. The ORF3, which is 729nt in length, encodes a putative M protein of 242aa (Table 1). One NLS was identified at aa positions 232–241. A NLS was also identified in the M protein encoded by Vesicular stomatitis virus [60]. The ORF4, which is 1,989nt in length, encodes a putative G protein of 662aa (Table 1). A signal peptide was predicted in the N-terminus of this protein at aa positions 21-22, while a transmembrane domain was predicted in the C-terminus at aa positions 611-633. Signal peptides and transmembrane domains are present in every rhabdovirus glycoproteins, due to its membrane-associated function in the rhabdoviral life cycle [61].

The partial ORF7, encodes the putative L protein (Table 1). Two NLS were identified at aa positions 159–170 and 723-751. A NLS was also identified in the L protein encoded by NcNSRV1 and GuNCRV1 [32]. As stated above, two accessory ORFs were predicted between the G and L genes, which was not surprising, since the G-L gene junction is the most commonly occupied site of accessory genes within rhabdoviruses [62]. Moreover, the presence of two additional genes in both ChMRV and GuNCRV1 could be an evidence of

gene duplication, which has been mentioned as a mechanism involved in the evolutionary process in other rhabdoviruses [63]. The ORF5, which is 273nt in length, encodes a putative U1 protein of 90aa (Table 1). A transmembrane protein was predicted at aa positions 54-76. This protein has the structural characteristics of a class I viroporin, that has been speculated to display membrane-associated functions [64, 65]. The ORF6, which is 159nt in length, encodes a putative U1 protein of 52aa (Table 1). A transmembrane protein was predicted at aa positions 20-42. Thus, like U1, this protein also should have membrane-associated functions. As expected, transmembrane domains were also identified in the U1 and U2 proteins encoded by GuNCRV1.

BlastP analysis of each protein encoded by ChMRV genome showed that the highest identities of the N, M, G and L proteins was with the GuNCRV1 homologous proteins, with low sequence identities (Table 1). On the other hand, both P and P5 proteins have no hits with any protein present in the NCBI database.

The consensus gene junction 3´-UUAUUUUUUGGUAA-5´ was identified in the non-coding regions of ChMRV genome. This sequence is highly similar to that reported for GuNCRV1 and was used to identify the accessory genes as bona fide predictions.

To explore the relationships of ChMRV with members belonging to the family *Rhabdoviridae*, phylogenetic analysis based on the L protein sequence were carried out, which showed that ChMRV is clustered with the leafhopper associated GuNCRV1, and the froghopper associated hemipteran rhabdo-related virus OKIAV30 (Fig. 5B). These viruses should be classified into a novel genus within the subfamily *Deltarhabdovirinae* in the family *Rhabdoviridae*.

Concluding remarks

Analysis of publicly available RNA-seq data led to the identification of several novel viruses which have expanded the knowledge about the virosphere, shedding light on the diversity and phylogenetic relationships of viruses infecting a wide number of host including insects [10-15]. The six viruses associated with the maize leafhopper were related with other leafhopper-associated viruses, thus supporting the host assignments of those viruses identified in this study. Knowledge of the virome (insect viral community) of insect pests is essential to develop bio-insecticides based on viruses; therefore, the identification of the maize leafhopper virome would be useful to develop novel and sustainable approaches based on virocontrol to manage this agronomically important pest.

Author Contributions: H.D and N.B., conceptualization; H.D, S.F. and N.B., data analysis; N.B., writing—original draft and H.D, S.F. and N.B., writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Conflict of interest: All authors declare that they have no conflict of interest

Ethical Approval: This article does not contain any studies with human participants or animals performed by any of the authors.

Funding: This research received no external funding

Data availability statement: The sequences deposited in GenBank with accession numbers will be updated when available and have been uploaded as Supplementary Materials 1 of this work.

Acknowledgments: We would like to express sincere gratitude to the generators of the underlying data used for this work. By following open-access practices and supporting accessible raw sequence data in public repositories available to the research community, they promoted the generation of new knowledge and ideas.

References

1. Nault, L.R. Maize bushy stunt and corn stunt: a comparison of disease symptoms, pathogen host ranges, and vectors. *Phytopathol* **1980,** 70, 659–662

2. Maluta, N., Castro, T., Spotti Lopes, J. DC-electrical penetration graph waveforms for Dalbulus maidis (Hemiptera: Cicadellidae) and the effects of entomopathogenic fungi on its probing behavior. *Sci reports* **2023,** 13, 22033

3. Vilanova, E.S., Ramos, A., Souza de Oliveira, M.C., Esteves, M.B., Gonçalves, M.C., Lopes, J.R.S.First report of a mastrevirus (Geminiviridae) transmitted by the corn leafhopper. *Plant Dis* **2022,** 106, 1330–1333.

4. Nault L.R. Evolution of an insect pest: maize and the corn leafhopper, a case study. *Maydica* **1990,** 35, 165-175.

5. Pozebon, H., Stürmer, G.R., Arnemann, J.A. Corn stunt pathosystem and its leafhopper vector in Brazil. *J. Econ. Entomol.* **2022**, 115, 1817–1833.

6. Carpane, P., Catalano, M.I. Probing behavior of the corn leafhopper *Dalbulus maidis* on susceptible and resistant maize hybrids. *PLoS One* **2022,** 17, e0259481.

7. Edgar, R.C., Taylor, J., Lin, V., Altman, T., Barbera, P., Meleshko, D., et al. Petabase-scale sequence alignment catalyses viral discovery. *Nature*. **2022**, 602, 142-147.

8. Reche, I.D'Orta, G.; Mladenov, N., Winget, D.M., Suttle, C.A. Deposition rates of viruses and bacteria above the atmosperic boundary layer. *ISME J.* **2018**, 12, 1154–1162.

9. Zhang, Y.-Z., Shi, M., Holmes, E. Using metagenomics to characterize an expanding virosphere. *Cell* **2018**, 172, 1168–1172.

10. Li, C., Shi, M., Tian, J., Lin, X., Kang, Y., et al. Unprecedented genomic diversity of RNA viruses in arthropods reveals the ancestry of negative-sense RNA viruses. *Elife* **2015**, 4.

11. Shi, M., Lin, X.-D., Tian, J.-H., Chen, L.-J., Chen, X., et al. Redefining the invertebrate RNA virosphere. *Nature* **2016**, 540, 539–543.

12. Käfer, S., Paraskevopoulou, S., Zirkel, F., Wieseke, N., Donath, A., et al. Re-assessing the diversity of negative strand RNA viruses in insects. *PLOS Pathog*. 2019, 15, e1008224.

13. Nouri, S., Matsumura, E.E., Kuo, Y.W., Falk, B.W. Insect-Specific Viruses: From Discovery to Potential Translational Applications. *Curr. Opin. Virol*. **2018**, 33, 33–41.

14. Qi, Y.-H., Ye, Z.-X., Zhang, C.-X., Chen, J.-P., Li, J.-M. Diversity of RNA viruses in agricultural insects. *Comput. Struct. Biotechnol. J.* **2023**, 21, 4312–4321.

15. Wu, H., Pang, R., Cheng, T., Xue, L., Zeng, H., et al. Abundant and Diverse RNA Viruses in Insects Revealed by RNA-Seq Analysis: Ecological and Evolutionary Implications. *mSystems* **2020**, 5, e00039-20.

16. Harvey, E., Holmes, E.C. Diversity and evolution of the animal virome. *Nat. Rev. Microbiol.* **2022**, 20, 321–334.

17. Lauber, C., Zhang, X, Vaas, J., Klingler, F., Mutz, P., et al. Deep mining of the Sequence Read Archive reveals major genetic innovations in coronaviruses and other nidoviruses of aquatic vertebrates. *Plos Pathogens* **2024**, 20, e1012163.

18. Neri, U., Wolf, Y.I., Roux, S., Camargo, A.P., Lee, B., et al. Expansion of the global RNA virome reveals diverse clades of bacteriophages. *Cell* **2022**, 185, 4023–4037.e18.

19. Paraskevopoulou, S., Ka, S., Zirkel, F., Donath, A., Petersen, M., Viromics of extant insect orders unveil the evolution of the flavi-like superfamily. *Virus Evol* **2021**, *7*, veab030.

20. Rutar, S.O.; Kordis, D. Analysis of the RNA virome of basal hexapods. *PeerJ* **2020**, 8, e8336.

21. Charon, J., Olendraite, I., Forgia, M., Chong, L.C., Hillary, L.S., et al. Consensus statement from the first RdRp Summit: advancing RNA virus discovery at scale across communities. *Frontiers in Virology* **2024**, 4, 1371958.

22. Koonin, E.V., Dolja, V.V., Krupovic, M., Varsani, A., Wolf, Y.I., et al. Global organization and proposed megataxonomy of the virus world. *Microbiol Mol Biol Rev*. **2020** 84, e00061-19.

23. Kuhn, J.H., Wolf, Y.I., Krupovic, M., Zhang, Y.-Z., Maes, P., Dolja, V.V., Koonin, E.V. Classify viruses—The gain is worth the pain. *Nature* **2019**, 566, 318–320.

24. Siddell, S.G., Smith, D.B., Adriaenssens, E., Alfenas-Zerbini, P., Dutilh, B.E., et al. Virus taxonomy and the role of the International Committee on Taxonomy of Viruses (ICTV). *J. Gen. Virol*. **2023**, 104, 001840.

25. Simmonds, P., Adams, M.J., Benkő, M., Breitbart, M., Brister, J. R., Carstens, E. B., et al. Virus taxonomy in the age of metagenomics. *Nature Rev Microbiol*. **2017** 15, 161-168.

26. Simmonds, P.; Adriaenssens, E.M.; Zerbini, F.M.; Abrescia, N.G.A.; Aiewsakun, P.; et al. Four principles to establish a universal virus taxonomy. *PLoS Biol*. **2023**, 21, e3001922.

27. Bejerman, N.; Dietzgen, R.; Debat, H. Novel Tri-Segmented Rhabdoviruses: A Data Mining Expedition Unveils the Cryptic Diversity of cytorhabdoviruses. *Viruses* **2023**, 15, 2402.

28. Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol. Biol. Evol*. **2021**, 38, 3022–3027.

29. Stork, N.E.; NE, S. How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth? *Annu. Rev. Entomol*. **2018**, 63, 31–45.

30. Longdon, B.; Murray, G.G.R.; Palmer, W.J.; Day, J.P.; Parker, D.J.; Welch, J.J.; Obbard, D.; Jiggins, F.M. The evolution, diversity, and host associations of rhabdoviruses. *Virus Evol*. **2015**, 1, vev014.

31. Bonning B.C. The Insect Virome: Opportunities and Challenges. *Curr Issues Mol Biol*. **2020**, 34, 1–12.

32. Jia, W., Wang, F., Xiao, S., Yang, Y., Chen, L., et al. Identification and characterization of a novel rhabdovirus in green rice leafhopper, Nephotettix cincticeps. *Virus Res*. **2021**; 296, 198281.

33. Wang, H., Liu, Y., Liu, W., Cao, M., Wang, X. Sequence analysis and genomic organization of a novel chuvirus, Tàiyuán leafhopper virus. *Arch Virol*. **2019**, 164, 617–620.

34. Fu, Y., Cao, M., Wang, H., Du, Z., Liu, Y., et al. Discovery and characterization of a novel insect-specific reovirus isolated from Psammotettix alienus. *J Gen Virol*. **2020**, 101, 884–892.

35. Li, L.L., Ye, Z.X., Chen, J.P., Zhang, C.X., Huang, H.J., et al. Characterization of Two Novel Insect-Specific Viruses Discovered in the Green Leafhopper, Cicadella viridis. *Insects*. **2022**, 13:4.

36. Ottati, S.; Chiapello, M.; Galetto, L.; Bosco, D.; Marzachì, C.; Abbà, S. New viral sequences identified in the Flavescence Dorée phytoplasma vector Scaphoideus titanus. *Viruses* **2020**, 12, 287.

37. Gilmer, D., Ratti C., Consortium I.R. ICTV virus taxonomy profile: Benyviridae. J. Gen. Virol. **2017**, *98*, 1571–1572.

38. Etebari, K., Lenancker, P., Powell, K.S., Furlong, M.J. Transcriptomics Reveal Several Novel Viruses from Canegrubs (Coleoptera: Scarabaeidae) in Central Queensland, Australia. *Viruses* **2022**, 14, 649.

39. Huang, H.J., Ye, Z.X., Wang, X., Yan, X.T., Zhang, Y., et al. Diversity and infectivity of the RNA virome among different cryptic species of an agriculturally important insect vector: Whitefly Bemisia tabaci. *NPJ Biofilms Microbiomes* **2021**, 7, 43.

40. Kondo, H., Hirano, S., Chiba, S., Andika, I.B., Hirai, M., Maeda, T., Tamada, T. Characterization of burdock mottle virus, a novel member of the genus Benyvirus, and the identification of benyvirus-related sequences in the plant and insect genomes. *Virus Res.* **2013**, 177, 75–86.

41. Solovyev, A.G., Morozov, S.Y. Uncovering Plant Virus Species Forming Novel Provisional Taxonomic Units Related to the family *Benyviridae. Viruses* **2022**, 14, 2680.

42. Liu, S., Valencia-Jiménez, A., Darlington, M., Vélez, A.M., Bonning, B.C. Diabrotica undecimpunctata virus 2, a Novel Small RNA Virus Discovered from Southern Corn Rootworm, Diabrotica undecimpunctata howardi Barber (Coleoptera: Chrysomelidae). *Microbiol. Resour. Announc.* **2020**, 9, e00380-20.

43. Leventhal, S.S., Wilson, D., Feldmann, H., Hawman, D.W. A Look into Bunyavirales Genomes: Functions of Non-Structural (NS) Proteins. *Viruses* **2021**, 13, 314.

44. Valles, S., Chen, Y., Firth, A., Guerin, D., Hashimoto, Y., et al. ICTV virus taxonomy profile: Iflaviridae. J. Gen. Virol. **2017**, 98, 527–528.

45. van Oers M. (2010). "Genomics and biology of Iflaviruses" in Insect virology. eds. Asgari S., Johnson K. (Norfolk: Caister Academic Press), 231–250.

46. Daveu, R., Hervet, C., Sigrist, L., Sassera, D., Jex, A., Labadie, K., Aury, J.-M., Plantard, O., Rispe, C. Sequence diversity and evolution of a group of iflaviruses associated with ticks. *Arch. Virol.* **2021**, 166, 1843–1852.

47. Jia, W., Wang, F., Li, J.; Chang, X., Yang, Y., Yao, H., et al. A Novel Iflavirus Was Discovered in Green Rice Leafhopper. *Front. Microbiol.* **2020**, 11, 621141.

48. Geng, P., Li, W., de Miranda, J.R., Qian, Z., An, L., Terenius, O. Studies on the transmission and tissue distribution of Antheraea pernyi iflavirus in the Chinese oak silkmoth *Antheraea pernyi*. *Virology* **2017**, 502, 171–175.

49. Lanzi, G., de Miranda, J.R., Boniotti, M.B., Cameron, C.E.,Lavazza, A., Capucci, L., Camazine, S.M., Rossi, C. Molecular and biological characterization of deformed wing virus of honeybees (*Apis mellifera L.*). *J. Virol.* **2006**, 80, 4998–5009.

50. Wells, T., Wolf, S., Nicholls, E., Groll, H., Lim, K.S., Clark, S.J., Swain, J., Osborne, J.L, Haughton, A.J. Flight performance of actively foraging honey bees is reduced by a common pathogen. *Environ. Microbiol. Rep.* **2016**, 8, 728–737.

51. Carballo, A., Murillo, R., Jakubowska, A., Herrero, S., Williams, T., Caballero, P. Co-infection with iflaviruses influences the insecticidal properties of Spodoptera exigua multiple nucleopolyhedrovirus occlusion bodies: Implications for the production and biosecurity of baculovirus insecticides. *PLoS ONE* **2017**, 12, e0177301.

52. Mengual-Martí, A., Martínez-Solís, M., Yousef-Yousef, M., Garrido-Jurado, I., Delgado-Sanfiel, P., et al. Impact of covert infections with an RNA virus on the susceptibility of Spodoptera exigua to natural enemies *BioControl* **2022** 67, 605-615.

53. Ottati, S., Persico, A., Rossi, M., Bosco, D., Vallino, M., et al. Biological characterization of Euscelidius variegatus iflavirus 1. *J. Invertebr. Pathol.* **2020**, 173, 107370.

54. Shaw, M.L., Palese P. "Orthomyxoviridae," in Fields Virology, 6th Edn, eds Knipe D. M., Howley P. M., Cohen J. I., Griffin D. E., Lamb R. A., Martin M. A., et al. (Philadelphia, PA: Lippincott Williams and Wilkins), **2013**, 1151–1185.

55. Papa, G., Abbà, S., Galetto, L., Parise, C., Marzachì, C., Negri, I. Distribution and prevalence of viral genomes in Italian populations of the invasive brown marmorated stink bug Halyomorpha halys. *J Invertebr Pathol* **2023**, 200,107977.

56. Kobayashi, D., Murota, K., Faizah, A.N., Amoa-Bosompem, M., Higa, Y., et al. RNA virome analysis of hematophagous Chironomoidea flies (Diptera: Ceratopogonidae and Simuliidae) collected in Tokyo, Japan. *Med. Entomol. Zool.* **2020**, 71, 225–243.

57. Walker, P.J., Freitas-Astúa, J., Bejerman, N., Blasdell, K.R., Breyta, R., et al. ICTV Virus Taxonomy Profile: Rhabdoviridae 2022. *J. Gen. Virol.* **2022**, 103, 001689.

58. Dietzgen, R.G., Kondo, H., Goodin, M.M., Kurath, G., Vasilakis, N. The family Rhabdoviridae: Mono- and bi-partite negative-sense RNA viruses with diverse genome organization and common evolutionary origins. *Virus Res*. **2017**, 227:158–170.

59. Schroeder, L., Mar, T.B., Haynes, J.R., Wang, R., Wempe, L., Goodin, M.M. Host range and population survey of *Spodoptera frugiperda rhabdovirus*. *J. Virol*. **2019**, 93, 1–30.

60. Glodowski, D.R., Petersen, J.M., Dahlberg, J.E. Complex nuclear localization signals in the matrix protein of vesicular stomatitis virus. *J. Biol. Chem*. **2002**, *277*, 46864–46870.

61. Barrett, C.T.; Dutch, R.E. Viral Membrane Fusion and the Transmembrane Domain. *Viruses* **2020**, 12, 693.

62. Walker, P.J., Dietzgen, R.G., Joubert, D.A., Blasdell, K. Rhabdovirus accessory genes. *Virus Res*. **2011**, 162, 110–125.

63. Walker, P.J., Firth, C., Widen, S.G., Blasdell, K.R., Guzman, H., et al. Evolution of Genome Size and Complexity in the Rhabdoviridae. *PLoS Pathog*. **2015**, 11, e1004664.

64. Joubert, D.A., Blasdell, K.R., Audsley, M.D., Trinidad, L., Monaghan, P., et al. Bovine Ephemeral Fever Rhabdovirus 1 Protein Has Viroporin-Like Properties and Binds Importin 1 and Importin 7. *J. Virol*. **2013**, 88, 1591–1603.

65. Xia, X., Cheng, A., Wang, M., Ou, X., Sun, D., et al. Functions of Viroporins in the Viral Life Cycle and Their Regulation of Host Cell Responses. *Front. Immunol*. **2022**, 13, 890549.

FIGURE LEGENDS

Figure 1. (A) Schematic representation of the genome organization of Chicharrita del maiz beny-like virus. (B) Maximum likelihood phylogenetic trees reconstructed using the RdRp protein sequence of Chicharrita del maiz beny-like virus and of representative beny-like viruses. Bootstrap values above 50% are shown (1,000 replicates). Chicharrita del maiz beny-like virus is indicated with a green square. The scale bar the substitution per site.

Figure 2. (A) Schematic representation of the genome organization of Chicharrita del maiz bunyan-like virus. (B) Maximum likelihood phylogenetic trees reconstructed using the RdRp protein sequence of Chicharrita del maiz bunyan-like virus and of representative bunyan-

like viruses. Bootstrap values above 50% are shown (1,000 replicates). Chicharrita del maiz bunyan-like virus is indicated with a green square. The scale bar the substitution per site.

Figure 3. (A) Schematic representation of the genome organization of Chicharrita del maiz iflavirus 1 and Chicharrita del maiz iflavirus 2. (B) Maximum likelihood phylogenetic trees reconstructed using the RdRp protein sequence of Chicharrita del maiz iflavirus 1 and Chicharrita del maiz iflavirus 2 and of representative iflaviruses. Bootstrap values above 50% are shown (1,000 replicates). Chicharrita del maiz iflavirus 1 and Chicharrita del maiz iflavirus 2 are indicated with green squares. The scale bar the substitution per site.

Figure 4. (A) Schematic representation of the genome organization of Chicharrita del maiz orthomyxo-like virus. (B) Maximum likelihood phylogenetic trees reconstructed using the PB1 proteins sequence of Chicharrita del maiz orthomyxo -like virus and of representative orthomyxo -like viruses. Bootstrap values above 50% are shown (1,000 replicates). Chicharrita del maiz orthomyxo -like virus is indicated with a green square. The scale bar the substitution per site.

Figure 5. (A) Schematic representation of the genome organization of Chicharrita del maiz rhabdovirus. (B) Maximum likelihood phylogenetic trees reconstructed using the L protein sequence of Chicharrita del maiz rhabdovirus and of representative rhabdoviruses. Bootstrap values above 50% are shown (1,000 replicates). Chicharrita del maiz rhabdovirus is indicated with a green square. The scale bar the substitution per site.

A

```
5′  ┌─────────────────────────────────┐┌──────────┐  3′
    │  Non-structural polyprotein     ││ Structural│
    │                                 ││ protein   │
    └─────────────────────────────────┘└──────────┘
```

B

```
                    ┌─ red clover RNA virus 1 MG596242
          53 ┌──────┤
             │      └─ carrot associated RNA virus 1 OP889248
      100 ┌──┤
         │   │ ┌─ Arceuthobium sichuanense virus 3 BK059270
         │   └─┤
    94 ──┤     └─ Dactylorhiza hatagirea beny-like virus BK013327
        │
        │  ┌─ goji berry chlorosis virus MH791331
        │  │
        │  │     ┌─ wheat stripe mosaic virus MH151798
   100 ─┤  │ 100 ┤
        │        │ ┌─ rice stripe necrosis virus MT027255
        │        └─┤
        │      100 │  ┌─ Burdock mottle virus AB818898
        │          └──┤
   61 ──┤         100 │ ┌─ beet necrotic yellow vein virus MT227164
        │             └─┤
        │          100 └─ beet soil-borne mosaic virus AF280539
```

plants

```
                 ┌─ Hubei beny-like virus 1 OL700057
           100 ┌─┤
        100 ┌──┤ └─ Sichuan mosquito Beny-like virus MZ556264
           │  │
           │  └─ Bemisia tabaci beny-like virus 6 MW256699
           │
           │ 74 ┌─ ■ Chicharrita del maiz beny-like virus
   100 ────┤   ┌┤
           │   ││ └─ Lepidiota negatoria beny-like virus OM421673
           │   │
        52 │   │ ┌─ Sanya benyvirus 1 MZ209861
   ───┤    81 ─┤
      │           99 ┌─ Diabrotica undecimpunctata virus 2 MN646771
      │             └┤
      │           97 └─ Guiyang benyvirus 1 MZ209815
```

insects

```
   93 ┌─ Erysiphe necator associated beny-like virus 1 MN617775
     │
     │       ┌─ Lentinula edodes beny-like virus 1 MH744728
     │  100 ─┤
     │       └─ Monilinia fructicola beny-like virus 1 ON038406
     │
     │  ┌─ Monilinia benyvirus C MK231094
  100 ─┤
     │  │ ┌─ Rhizoctonia beny-like virus 1 OR227245
     └──┤
      92 └─ Sclerotium rolfsii beny-like virus 1 MH766487
```

fungi

```
        ┌─ Nudaurelia capensis beta virus AF102884
   100 ─┤
        │ ┌─ Dendrolimus punctatus virus AY594352
        └─┤
       100 └─ Helicoverpa armigera stunt virus U18246
```

Alphatetraviridae

├──────────┤
0.20

A

3' **RdRp** 5'   Segment L

3' **NP** 5'   Segment S

B

```
      68 ┌─ ■ Chicharrita del maiz bunyan-like virus
    81 ┌─┤
       │  └─ Hubei insect virus 1 KX884735
  100 ─┤
       └─ Hemipteran phenui-related virus OKIAV285 MT153431
  52
       ┌─ Hangzhou nephotettix cincticeps phenuivirus 1 MZ209637
  89
       └─ Sanya nilaparvata lugens phenuivirus 1 MZ209823
                ┌─ Lampyris noctiluca bunyan-like virus 1 MH620814
                ├─ Erthesina fullo bunyavirus 1 MW865704
  99            └─ Hubei bunya-like virus 13 KX884784
          85 ┌─ Gonipterus platensis bunyan-like virus MT435497
             └─ Thrips tabaci associated bunyavirales 1 MN764152
       91 100 ┌─ Degsystermes virus MW052074
  57         └─ Ogsystermes virus MW052081
          ┌─ Plecopteran phenui-related virus OKIAV280 MT153514
       99 │    ┌─ Wufeng shrew phenuivirus 9 OQ715525
          └──100┤
  71              └─ Wufeng shrew phenuivirus 7 OQ715523
       ┌─ Dipteran phenui-related virus OKIAV281 MT153358
    99 ┌─ Wuhan insect virus 16 KX884733
  82 ─┤
       └─ Linepithema humile bunyan-like virus 1 MH213237
  82   ├─ Hubei bunya-like virus 3 KX884778
  53   ┌─ barley aphid RNA virus 5 LC516850
    100│  ┌─ Hymenopteran phenui-related virus OKIAV255 MW288195
       100└─ Hubei bunya-like virus 2 KX884780
          ┌─ Lepidopteran phenui-related virus OKIAV270 MT153502
       100└─ Lepidopteran phenui-related virus OKIAV271 MT153406
       ┌─ Dipteran phenui-related virus OKIAV274 MT153385
  79   ┌─ Culex Bunyavirus 2 MW434596
    100└─ Fangshan bunya-like virus OR160351
           100 ┌─ Aphalara polygoni bunya-like virus BK062758
               └─ Gouleako virus HQ541738
  100       ┌─ Ceraphron bunya-like virus BK062755
       98   └─ Cumuto virus KF543244
```

Phenui-like viruses

Goukovirus

0.10

A

3´ PB1 5´ RNA 1
3´ PB2 5´ RNA 2
3´ PA 5´ RNA 3
3´ NP 5´ RNA 4
3´ HA 5´ RNA 5

B

Hymenopteran orthomyxo-related virus OKIAV175 MW288219
99
Hymenopteran orthomyxo-related virus OKIAV174 MW288228
79
100
Hymenopteran orthomyxo-related virus OKIAV171 MW288220
86
Hymenopteran orthomyxo-related virus OKIAV173 MT153296
100
Phasmatodean orthomyxo-related virus OKIAV172 MT153463
Hubei earwig virus 1 KX883858
100
Raphidiopteran orthomyxo-related virus OKIAV180 MW288202
Photinus pyralis orthomyxo-like virus 1 MG972988
76
69
Photinus pyralis orthomyxo-like virus 2 MG972993
100
Hemipteran orthomyxo-related virus OKIAV182 MT153551
100
Hemipteran orthomyxo-related virus OKIAV183 MT153375
87
Dermapteran orthomyxo-related virus OKIAV170 MT153548
Coleopteran orthomyxo-related virus OKIAV158 MT153412
100
Siphonapteran orthomyxo-related virus OKIAV157 MT153496
Phasmatodean orthomyxo-related virus OKIAV167 MT153421
Bat faecal associated orthomyxo-like virus 1 ON872578
100
Hemipteran orthomyxo-related virus OKIAV188 MT153485
Halyomorpha halys orthomyxo-like virus 1 OM650310
100
94
51
Hemipteran orthomyxo-related virus OKIAV191 MT153419
■ Chicharrita del maiz orthomyxo-like virus
Shuangao Insect Virus 4 KM817619
Dipteran orthomyxo-related virus OKIAV193 MT153481
100
Dipteran orthomyxo-related virus OKIAV195 MT153448
100
100
Jingshan Fly Virus 1 KM817615
87
Wuhan louse fly virus 4 KM817621
100
100
Wuhan louse fly virus 3 KM817620

Quaranja-like viruses

0,20

A

3´ — N — P — M — G — P4 — P5 — L — 5´

B

Apis rhabdovirus 4 MZ822105 — 100
Apis rhabdovirus 5 OP972869 — *Gammahymrhavirus*
Diachasmimorpha longicaudata rhabdovirus KP735609

Blattodean rhabdo-related virus OKIAV14 MT153532
Sanya conocephalus maculatus rhabdovirus 1 MZ209844

Entomophthora rhabdovirus A MK231099
Wuhan house fly virus 2 KM817649
Drosophila sturtevanti rhabdovirus 1 KR822823 — *Alphadrosrhavirus*
Wuhan fly virus 3 KM817647
Shayang Fly Virus 3 KM817636

Wenling crustacean virus 10 KX884450 — *Alphacrustrhavirus*
Wenling crustacean virus 11 KX884456

North fork virus PP415847
Fairlight virus MK026565
Blacklegged tick rhabdovirus 1 MF360790 — *Betaricinrhavirus*
Chimay rhabdovirus 975531
Mudanjiang Rhabd tick virus 1 ON746525

Hemipteran rhabdo-related virus OKIAV30 MT153400
■ Chicharrita del maiz rhabdovirus
Guiyang nephotettix cincticeps rhabdovirus 1 HWYCFY159 MZ209705
Guiyang nephotettix cincticeps rhabdovirus 1 HWYCFY172542 MZ209638
Guiyang nephotettix cincticeps rhabdovirus 1 HWYCFY134 MZ209819

Xiangshan rhabdo-like virus 4 OK491502 — *Betahymrhavirus*
Coleopteran rhabdo-related virus OKIAV20 MT153398

Nephotettix cincticeps negative-stranded RNA virus 1 MW084959
Hangzhou nephotettix cincticeps rhabdovirus 1 MZ209636

Guadeloupe Culex rhabdovirus MN013390 — *Stangrhavirus*
Grenada mosquito rhabdovirus 1 MG385079

Deltarhabdovirinae

Hirame rhabdovirus FJ376982
Infectious hematopoietic necrosis virus OQ801357 — *Novirhabdovirus* — *Gammarhabdovirinae*
Viral hemorrhagic septicemia virus MN038336
snakehead rhabdovirus AF147498

0,10