Nonergodicity and Simpson's paradox in neurocognitive dynamics of cognitive control

Percy K. Mistry[1]*, Nicholas K. Branigan[1]*, Zhiyao Gao[1], Weidong Cai[1,2], Vinod Menon[1,2,3]

[1] Department of Psychiatry & Behavioral Sciences
[2] Wu Tsai Neurosciences Institute
[3] Department of Neurology & Neurological Sciences

Stanford University, Stanford, CA, United States.

* These authors contributed equally to this work.

Corresponding authors:
Percy K. Mistry, Ph.D. & Nicholas K. Branigan & Vinod Menon, Ph.D.
email: percym@stanford.edu; branigan@stanford.edu; menon@stanford.edu

Abstract

Nonergodicity and Simpson's paradox present significant, yet underappreciated challenges in
neuroscience. Leveraging brain imaging and behavioral data from over 4,000 children and a
Bayesian computational model of cognitive dynamics, we investigated brain-behavior
relationships underlying cognitive control at both between-subjects and within-subjects levels.
Strikingly, we observed a reversal of associations of inhibitory control brain activations with
dynamic behavioral measures when comparing between-subjects and within-subjects analyses,
revealing the nonergodic nature of these processes. This nonergodicity was pervasive throughout
the brain but most pronounced in the salience network. Additionally, within-subjects analysis
uncovered dissociated brain representations of reactive and proactive control processes, as well
as distinct brain-behavior associations for individuals who adaptively versus maladaptively
regulated cognitive control. Our findings offer insights into dynamic neural mechanisms of
cognitive control during a critical developmental period. This work highlights the importance of
embracing nonergodicity in human neuroscience, with implications for both theoretical
understanding and applications to AI and psychopathology.

Introduction

Since the nineteenth century, the concept of ergodicity has been central to physics[1-3]. However, only recently have other disciplines such as economics[4] and psychology[5,6] begun to examine how their own theories and findings rest upon assumptions of ergodicity. Nonergodicity refers to a system's behavior where time averages and ensemble averages do not coincide[1]. In psychology, a classic case of nonergodicity is Simpson's paradox observed in the context of the speed-accuracy tradeoff: speed and accuracy are often positively correlated between individuals (faster people are more accurate)[7], but are negatively correlated within individuals (when an individual tries to respond faster, their accuracy decreases)[8]. This striking divergence between group-level patterns and individual-level dynamics underscores the nonergodic nature of behavioral processes.

The implications of nonergodicity are profound. Analyzing data from a single individual over time can yield different conclusions than analyzing data across multiple individuals at a single point in time[5,6,9]. This challenges traditional assumptions about the generalizability of findings from between-subjects analyses to individual processes[5,9-13] changing how we interpret and generalize results in the life sciences. While progress has been made in understanding nonergodicity within behavioral contexts[5,6,14-17], its application to human brain function remains unexplored. This gap in research is particularly striking given that most human neuroscience studies of brain-behavior relations rely heavily on between-subjects analysis. If neurocognitive processes are indeed nonergodic, considering within-subjects analysis of neuroscientific data could profoundly enhance our understanding of brain function.

Nonergodicity in neurocognitive function may manifest as a divergence between group-level and individual-level brain-behavior relationships. Specifically, the association between neural activity and behavior may differ substantially when examined between subjects after averaging over time (**Figure 1a**) compared to within subjects over time (**Figure 1b**), potentially leading to a neurocognitive manifestation of Simpson's paradox (**Figure 1c**)[11,16]. This phenomenon has profound implications for our understanding of brain function and cognition. Group-level relationships may obscure the dynamic processes occurring within each individual, leading to incomplete or even misleading conclusions about the neural mechanisms underlying cognitive processes[5,6,14-17].

The integration of nonergodic principles into human neuroscience not only promises to advance theoretical understanding but also holds practical implications for tailoring therapeutic strategies to better address individual neurological and cognitive differences. This approach could lead to a more precise and accurate understanding of brain function, paving the way for personalized interventions in clinical neuroscience and psychiatry. There is a pressing need to integrate nonergodic principles into the study of human cognition and brain function, particularly in complex cognitive domains such as inhibitory control, where individual variability and dynamic processes play crucial roles.

To address critical gaps in our knowledge of nonergodicity in neurocognitive processes, we leveraged a large sample (N~4,000) from the Adolescent Brain Cognitive Development (ABCD) study, a multisite longitudinal study of brain development and behavior in children and adolescents[18]. We used behavioral and brain imaging data to examine the neural mechanisms of

85 cognitive control at both the between-subjects and within-subjects levels, aiming to elucidate the
86 nonergodic nature of these processes and its implications for our understanding of brain function
87 and cognition.
88
An important component of human cognition is inhibitory control, which is the ability to
withhold or cancel maladaptive actions, thoughts, and emotions, and is a fundamental component
of goal-directed behavior[19,20]. This critical cognitive function allows individuals to navigate
complex environments, adapt to changing circumstances, and maintain focus on long-term goals
in the face of immediate temptations or distractions[21]. Given its central role in cognitive control,
understanding the neural mechanisms underlying inhibitory control has been a primary focus of
cognitive neuroscience research[22]. Inhibitory control engages a distributed network of cortical
and subcortical regions. Previous studies have implicated the involvement of the salience
network, particularly the anterior insula and dorsal anterior cingulate cortex, in detecting and
processing relevant stimuli and coordinating neural resources for inhibitory control[23-27].
Additionally, the inferior frontal gyrus, presupplementary motor area, and basal ganglia have
been shown to play crucial roles in implementing response inhibition[28-34]. While the dynamic
interplay between these cortical and subcortical regions is known to underpin successful
inhibitory control, our understanding of how these neural processes relate to moment-to-moment
fluctuations in cognitive control at the individual level remains limited.
104
The stop signal task (SST) is a widely used paradigm to assess inhibitory control. In this task,
participants are instructed to respond quickly to a primary "go" stimulus but to withhold their
response when an infrequent "stop" signal appears[20]. The ability to inhibit responses is quantified
using the stop signal reaction time (SSRT), which estimates the latency of the stopping
process[20,21]. Conventionally, studies investigating the neural basis of inhibitory control have
relied on between-subjects analyses of SSRT, comparing brain activity across individuals with
varying abilities to inhibit responses. These analyses typically correlate individual differences in
SSRT with differences in task-related brain activation patterns[24,35-41]. Such approaches aggregate
data at the individual level and make inferences about the neural mechanisms of inhibitory
control based on group-level relationships. However, this conventional approach implicitly
assumes ergodicity—specifically that the average trend in the group reflects the trend within
each individual[6,16,42]. If this assumption does not hold, then inferences made from group-level
associations cannot be generalized to explain the cognitive processes occurring within each
individual[11,16].
119
However, testing the ergodicity assumption in inhibitory control studies has been challenging,
particularly when using paradigms like the SST. This is because traditional models do not
provide trial-wise SSRT estimates, instead requiring a substantial number of trials for a single
estimate[43]. This limitation has constrained researchers' ability to examine within-subjects
dynamics directly, making it difficult to verify whether group-level relationships truly reflect
individual-level processes.
126
127 To address these limitations and capture the dynamic nature of inhibitory control, we developed
128 the Proactive Reactive and Attentional Dynamics (PRAD) model of SST behavior. This
129 computational model extends beyond conventional approaches to the SST by incorporating
130 proactive and reactive control mechanisms[44]. The PRAD model allows us to infer dynamic, trial-

level measures corresponding to elemental neurocognitive subprocesses for each subject, including SSRTs and measures of proactive delaying based on stop signal anticipation. PRAD parameters characterize the latent cognitive constructs governing action execution and inhibition in the SST, providing a more comprehensive view of inhibitory control processes than traditional models.

The PRAD model dissociates observed behavior into elemental latent neurocognitive subprocesses, that can characterize multiple dynamic measures of within-subjects variation. This is critical to conducting within-subjects analysis and evaluating ergodicity, since behavioral variability within a cognitive task is often governed by the complex interplay of multiple neurocognitive subprocesses, each of which may potentially have distinct underlying brain dynamics. By considering the various sequential dependencies and compensatory, competing, and interacting neurocognitive mechanisms underlying cognitive control, the PRAD model dissociates elemental processes and improves the interpretability of inferred within-subjects brain-behavior relationships. Specifically, the PRAD model can infer non-random and sequentially dependent trial-level variability in the probability and duration of proactive delayed responding[45], as well as in the effect of attentional modulation on the reactive SSRT.

We combined task fMRI data with the PRAD model to investigate the relationship between task-evoked brain responses and dynamic inhibitory control processes at both between-subjects and within-subjects levels. This approach allows us to directly compare inferences made from traditional group-level analyses (group-level associations) with those derived from a fine-grained examination of within-individual variation in neurocognitive dynamics. We leveraged the dynamic representations of subjects' behaviors from the PRAD cognitive model alongside simultaneous task fMRI data to probe how moment-to-moment variations in brain activity relate to fluctuations in inhibitory control processes. This analysis strategy promotes a richer understanding of the neural basis of inhibitory control, potentially revealing insights that would be obscured in conventional group-level analyses.

We had five interconnected aims (**Figure 1d**). First, we examined how key parameters of reactive and proactive control[46] from the PRAD model related to brain activity at both between-subjects and within-subjects levels, identifying networks showing significant associations. By comparing these levels, we show the limitations of assuming ergodicity in the study of inhibitory control and underscore the importance of considering within-subjects variability. Second, motivated by the replicability crisis in neuroscience research, we assessed the stability and robustness of our findings. We examined the stability of our within-subjects findings as a function of the sample and sample size through bootstrap resampling. Then, we tested whether our finding of nonergodicity was robust to alternative between-subjects analyses using different measures of brain activation and robust to the use of a directly observed behavioral measure in place of latent cognitive model parameters. Third, we probed the brain's implementation of proactive and reactive control processes underlying inhibitory control. Proactive control involves the anticipation and preparation for stopping, while reactive control involves the actual implementation of response inhibition[46]. By comparing the brain representations of these processes, we improve understanding of how their neural underpinnings relate. Fourth, we identified subgroups of within-subjects results associated with individual differences in adaptive regulation of inhibitory control. Finally, to characterize the spatial distribution of nonergodicity

across the brain, we quantified and compared nonergodicity levels across different functional networks. We defined a network's degree of nonergodicity as the proportion of individuals whose within-subjects brain-behavior associations exhibited a direction opposite to that observed in the between-subjects analysis. This approach allowed us to identify brain networks with significant levels of nonergodicity.

Our findings demonstrate a stark contrast between traditional between-subjects analyses and novel within-subjects analyses that account for dynamic cognitive processes. Results not only highlight the limitations of assuming ergodicity in cognitive control research but also provide a more precise understanding of the neural mechanisms underlying inhibitory control at the individual subject level. Our within-subjects analysis of neurocognitive dynamics reveals insights that are either obscured by or entirely unavailable from conventional between-subjects approaches. These discoveries have profound implications for both basic neurocognitive research and clinical applications.

Results

Dynamic cognitive process model of behavior

We used data from the SST (**Figure 2a**) to investigate dynamic cognitive processes underlying cognitive control in a large sample ($N\sim4000$) from the baseline visit of the ABCD study. We developed PRAD, a dynamic cognitive process model[44] that provides a multidimensional perspective of the elemental cognitive processes governing the reactive and proactive dynamics involved in inhibitory control. This model incorporates latent dynamics that respond to internal cognitive states (endogenous variables) and external environmental contingencies (exogenous variables), with interaction between these dynamics governed by latent trait measures. The latent trait (individual-level) and dynamic (trial-level) measures are simultaneously inferred within a hierarchical Bayesian framework (**Figure 2b**). For each subject and trial, the model infers a measure of reactive inhibitory control, SSRT, and 2 measures of proactive control, probability of proactivity and proactive delaying (**Figure 2c-d**). SSRT measures how long it takes for a subject to inhibit a response, probability of proactivity measures whether they use a strategy of proactive delaying, and proactive delaying measures the length of proactive delays when a proactive strategy is utilized[45,47]. Importantly, all 3 of these parameters are inferred for each trial (**Figure 2e**). **Supplementary Figure S1** provides a detailed illustration of the model. Validation of the PRAD model can be found in a separate study[44].

Between-subjects and within-subjects analysis of brain-behavior associations

We examined how 3 key parameters of the PRAD model that were inferred at a trial level (dynamic measures)—SSRT, probability of proactivity, and proactive delaying—related to brain activity at both between-subjects and within-subjects levels (**Figure 3**). To investigate brain-behavior relationships in inhibitory control, we conducted both between-subjects and within-subjects analyses using fMRI data from the stop signal task (SSRT and probability of proactivity $N = 4469$; proactive delaying $N = 4176$). For the between-subjects analyses, we Pearson correlated subject-average brain activation during successful stopping (correct stop versus correct go activation) with subject-average SSRT, probability of proactivity, and proactive

delaying across participants[24,37-41]. For the within-subjects analyses, we regressed the fMRI
signal on the parameters using fMRI general linear models with SSRT, probability of proactivity,
and proactive delaying included as regressors. We thus modeled the fMRI signal as a
combination of static trial-type effects and dynamic effects proportional to trial-by-trial
variations in the cognitive model parameters. This allowed us to examine how fluctuations in
cognitive processes covaried with brain activity within each individual, while adjusting for
stimulus types. These complementary approaches enabled us to compare group-level and
individual-level trends in brain-behavior relationships.
231
SSRT. Between subjects, SSRT showed widespread negative correlations with brain activity,
including in frontal, parietal, and temporal areas, and in regions implicated in cognitive control
(**Figure 3a**). This suggests that individuals with faster inhibitory responses (lower SSRT) show
greater activation in these regions during successful stopping, replicating previous findings[41]. In
contrast, within subjects, SSRT showed positive associations with brain activity, particularly in
frontal and parietal regions (**Figure 3b**). This suggests that on trials where individuals have
slower inhibitory responses, they show increased activation in these areas.
239
240 *Probability of proactivity.* Between subjects, probability of proactivity showed minimal
241 correlations with brain activity (**Figure 3c**). In contrast, within subjects, probability of
242 proactivity showed negative associations in parietal, temporal, and lateral frontal regions, and
243 positive associations in default mode network regions (**Figure 3d**).
244
245 *Proactive delaying*. Between subjects, proactive delaying showed widespread positive
246 correlations with brain activity, particularly in frontal and parietal regions (**Figure 3e**). This
247 indicates that individuals who engage in more proactive delaying exhibit higher activation in
248 these areas during successful stopping. In contrast, within-subjects, proactive delaying showed
249 negative associations in frontal and parietal cortex (**Figure 3f**).
250
These findings reveal a striking divergence of between-subjects and within-subjects brain-
behavior relationships in inhibitory control. The reversal of association directions, particularly
for SSRT and proactive delaying, suggests that inferences about the neural mechanisms
underlying inhibitory control do not generalize between group and individual levels. This
nonergodic pattern highlights the importance of considering both levels of analysis to more fully
understand the neurocognitive dynamics of inhibitory control.
257
258 ***Network-level visualization of between- and within-subjects brain-behavior associations***
259
260 To elucidate the patterns of brain-behavior relationships across different functional brain
261 networks, we visualized the between-subjects and within-subjects associations for SSRT,
262 probability of proactivity, and proactive delaying. We used the Shirer network atlas[48] for our
263 primary analysis as it includes the basal ganglia, a subcortical system important for the
264 implementation of inhibitory control[24,49].
265
SSRT (**Figure 4a**). Between-subjects analysis showed consistent negative correlations in most
networks, with effects in the posterior salience, frontoparietal, and default mode networks (all
$P_{FDR} < 0.01$). In contrast, within-subjects analysis revealed mixed negative and positive

associations, with opposite (positive) associations in the posterior salience, precuneus,
visuospatial, auditory, and sensorimotor networks (all $P_{FDR} < 0.01$). The reversal of association
directions underscores the nonergodic nature of SSRT-related brain activity.

Probability of proactivity (**Figure 4b**). Between-subjects analysis demonstrated no significantly
nonzero correlations in the networks (all $P_{FDR} \geq 0.05$). However, within-subjects analysis
unveiled a more complex pattern. Negative associations were observed in the salience and
frontoparietal networks, while positive associations were seen in the default mode network (all
$P_{FDR} < 0.01$). This disparity highlights the importance of examining within-subjects dynamics
for proactivity.

Proactive delaying (**Figure 4c**). Between-subjects analysis revealed positive correlations in all
the networks (all $P_{FDR} < 0.01$). Yet, within-subjects analysis showed predominantly negative
associations, notably in the salience and frontoparietal networks (all $P_{FDR} < 0.01$). The
retrosplenial cortex and parahippocampal gyrus nodes of the ventral default mode network
showed a positive within-subjects associations ($P_{FDR} < 0.01$).

These network-level visualizations emphasize the divergent patterns between group-level and
individual-level associations across different functional brain networks. The consistent reversals
observed, for both reactive and proactive control measures, reinforce that brain-behavior
relationships in inhibitory control are nonergodic. These findings indicate that both between-
subjects and within-subjects perspectives are needed to comprehensively understand the neural
dynamics underlying inhibitory control.

Stability of within-subjects brain-behavior associations

To assess the stability of our within-subjects findings, we performed bootstrap resampling at
varying sample sizes (**Figure 5**). Within-subjects associations between brain activity and the
model parameters (SSRT, probability of proactivity, and proactive delaying) were stable, even in
modest sample sizes. In each of 5 independent sets of brain areas, resampled results showed high
similarity with results in the full sample. Key findings were consistently observed in samples as
small as 25 subjects, with some effects requiring larger samples to emerge reliably. This stability
suggests the validity and potential generalizability of our within-subjects approach to
understanding neurocognitive mechanisms of inhibitory control. A detailed description of these
results is in the Supplementary Materials.

Robustness of nonergodicity to analytical choices

To test the robustness of our findings of nonergodicity, we conducted several control analyses
comparing between- and within-subjects brain-behavior relationships. We tested alternative
between-subjects analyses using different measures of brain activation and explored whether the
observed nonergodicity was specific to the latent model parameters or could also be seen using
reaction time on go trials, a directly observed behavioral measure. Nonergodicity persisted for all
approaches to between-subjects analysis of both model-derived parameters (**Supplementary
Figure S2**) and the observed behavioral measure (**Supplementary Figure S3**). Across all
analytical choices, we observed divergent patterns of between-subjects and within-subjects

associations. A detailed description of these control analyses and their results is in the
Supplementary Materials. Our results strongly suggest that the neurocognitive dynamics of
inhibitory control in children are fundamentally nonergodic, with implications for how we
interpret findings from traditional group-level associations.
319
320 ***Representational similarity analysis reveals dissociated reactive and proactive representations***
321
Building on our findings of nonergodic brain-behavior relationships, we sought to deepen our
understanding of how the brain implements proactive and reactive control at the individual level.
While the preceding results demonstrate the importance of within-subjects analyses, they leave
unresolved the question of how representations of reactive and proactive processes relate to each
other within brain networks. To investigate this, we used representational similarity analysis to
examine the overlap between brain representations of reactivity (SSRT) and proactivity
(probability of proactivity and proactive delaying) within individuals. For each subject and each
brain network, we computed pairwise Pearson correlations between the subject's brain maps of
SSRT and probability of proactivity, SSRT and proactive delaying, and probability of proactivity
and proactive delaying, over the voxels in each network (**Figure 6a**).
332
SSRT showed low similarity with both proactive measures (probability of proactivity and
proactive delaying) in all networks, with median correlations ranging from -0.07 to 0.06. In
contrast, the 2 proactive measures (probability of proactivity and proactive delaying) exhibited
high similarity in all networks, with median correlations ranging from 0.61 to 0.67. In each
network, the 3 similarity measures (SSRT and probability of proactivity, SSRT and proactive
delaying, and probability of proactivity and proactive delaying) were significantly different from
each other in their median values (**Figure 6b**) (all $P_{\text{FDR}} < 0.01$). These findings suggest that
representations of reactivity and proactivity are largely dissociated. This dissociation persists
across multiple brain networks, indicating a fundamental separation in how the brain encodes
reactive and proactive control processes.
343
344 ***Adaptive regulation of inhibitory control associated with distinct within-subjects results over***
345 ***subgroups***
346
To understand how our within-subjects associations related to between-subjects variation in
cognitive and task strategies, we examined the within-subjects results between subgroups
showing adaptive and maladaptive regulation of reactive and proactive behaviors. These
subgroups were identified using PRAD model parameters γ_1 (governing regulation of stopping
expectancy, and hence SSRT over trials) and θ_1 (governing regulation of probability of
proactivity over trials, in response to performance monitoring).
353
First, the cognitive model infers for each subject γ_1, which determines whether subjects
adaptively ($\gamma_1 > 0$) or maladaptively ($\gamma_1 < 0$) regulate reactivity. Adaptive (maladaptive)
regulation of reactivity involves increasing (decreasing) expectancy of a stop trial as the number
of successive go trials increases. Since expectancy of a stop trial is one of several determinants
of SSRT, γ_1 influences SSRT variation through time. Thus, we examined within-subjects
associations between SSRT and brain activity separately among subjects with $\gamma_1 < 0$ ($N =$
2513) and $\gamma_1 > 0$ ($N = 1956$) (**Figure 7a**). We found differences in the distributions of within-

subjects SSRT associations between the γ_1 subgroups in all networks examined (all $P_{\text{FDR}} <$ 0.01). The maladaptive regulation group had a larger (more positive) effect size in every network. In fact, across most networks, SSRT exhibited opposite associations with brain activity in the two subgroups. For example, in the frontoparietal and default mode networks, SSRT displayed a positive association among subjects with $\gamma_1 < 0$ and a negative association among subjects with $\gamma_1 > 0$ (all $P_{\text{FDR}} < 0.01$). Moreover, this analysis revealed that some of the effects in the full sample were driven by subjects belonging to one of the subgroups. For example, in the anterior salience network, brain activity's positive association with SSRT in the full sample was driven by $\gamma_1 < 0$ subjects; among $\gamma_1 < 0$, the effect in the anterior salience had a Cohen's d of ~0.3, twice that of the effect in the full sample, while among $\gamma_1 > 0$, there was no significant effect at all.

Second, the cognitive model infers for each subject θ_1, which determines whether subjects adaptively ($\theta_1 < 0$) or maladaptively ($\theta_1 > 0$) regulate proactivity. Adaptive (maladaptive) regulation of proactivity involves increasing (decreasing) the probability of proactivity following a failed stop trial and decreasing (increasing) the probability of proactivity following a no-response go-trial. Therefore, θ_1 influences the probability of proactivity's variation through time. Thus, we examined the probability of proactivity's within-subjects associations with brain activity separately among subjects with $\theta_1 < 0$ ($N = 3054$) and $\theta_1 > 0$ ($N = 1415$) (**Figure 7b**). We found consistent differences in the probability of proactivity's within-subjects associations between these subgroups. Generally, subjects with adaptive regulation ($\theta_1 < 0$) had more negative associations between probability of proactivity and brain activity. This negative coupling between proactivity and brain activity was pronounced within the adaptive regulation group in the anterior salience network (Cohen's d of ~0.4). This analysis also clarified how adaptive and maladaptive regulation of proactivity contributed to the effects observed in the full sample. The within-subjects positive association between probability of proactivity and activation in the retrosplenial cortex and parahippocampal gyrus network was observed in both θ_1 subgroups (both $P_{\text{FDR}} < 0.01$), but it was twice as large among subjects who maladaptively regulated proactivity.

These results demonstrate that population subgroups related to adaptive and maladaptive regulation of inhibitory control demonstrate different, and even opposite, within-subjects brain-behavior associations.

Brain networks exhibit varying degrees of nonergodicity and a hierarchical organization by nonergodicity

To further understand the distribution of nonergodicity across the brain, we quantified and compared nonergodicity levels in different brain networks. We defined a nonergodicity measure as the fraction of subjects for whom within-subjects brain-behavior association showed an opposite sign to the between-subjects association. Values above 0.5 indicate higher nonergodicity, while values below 0.5 suggest more ergodic behavior.

Our analysis revealed substantial variation in nonergodicity across brain networks (**Figure 8a**). Notably, the anterior salience network consistently demonstrated the highest level of nonergodicity for all three cognitive model parameters (SSRT, probability of proactivity, and

proactive delaying). The measures of network nonergodicity for associations with the probability of proactivity showed wide confidence intervals, reflecting the weak between-subjects associations between this parameter and brain activation.

To understand how networks related to each other in terms of nonergodicity, we performed hierarchical clustering on the joint nonergodicity measures of the networks with respect to all three cognitive model parameters (**Figure 8b-d**). This analysis revealed a hierarchical organization of brain networks based on their nonergodicity profiles. The anterior salience network emerged as the most dissimilar, forming its own cluster separate from all other networks. Additionally, the default mode network and retrosplenial cortex and parahippocampal gyrus clustered together, suggesting similarities in their nonergodic behavior.

These findings demonstrate that brain networks exhibit varying levels of nonergodicity and reveal a hierarchical organization of brain networks based on their nonergodic properties. This organization suggests a new perspective on the functional architecture of the brain and how it relates to cognitive control processes.

Discussion

A fundamental question we examined in this study is whether between- and within-subjects brain-behavior associations yield convergent findings in understanding neurocognitive processes underlying cognitive control. Divergence between these levels of analysis would provide evidence for nonergodicity, a phenomenon where inferences drawn from population-level data do not accurately represent individual-level processes. To investigate this, we leveraged a large community sample of children from the ABCD study and employed a dynamic computational model to elucidate the potentially nonergodic nature of neurocognitive dynamics underlying cognitive control. We combined task fMRI data with a hierarchical Bayesian model of cognitive dynamics to examine brain-behavior relationships at both between-subjects and within-subjects levels. Our findings reveal pervasive nonergodic patterns in brain-behavior relationships underlying cognitive control, with striking divergences between population-level and individual-level analyses. These results underscore the need to distinguish between-subjects and within-subjects inferences about the brain—a new paradigm we term "nonergodicity neuroscience".

At the between-subjects level, inhibitory control activations in key cognitive control networks were negatively correlated with subject-average SSRTs, aligning with previous studies[24,35-37]. This suggests that individuals with better inhibitory control (lower SSRTs) exhibit greater activation in regions associated with cognitive control. However, within-subjects analysis revealed a markedly different pattern: brain activity in some of these same networks was positively associated with trial-level SSRTs and negatively associated with trial-level proactive control. Thus, within-subjects associations were largely dissonant with the between-subjects findings, revealing the nonergodic nature of inhibitory control processes. Specifically, the opposing patterns demonstrated Simpson's paradox, a type of nonergodicity frequently observed in behavioral studies, where relationships at the group level are changed at the individual level[5,12,13,50]. This divergence underscores the importance of considering both group-level and individual-level associations in cognitive neuroscience research and highlights the complex, dynamic nature of inhibitory control processes.

Furthermore, representational similarity analysis uncovered distinct neural representations for reactive and proactive cognitive processes. Across all examined brain networks, representations of reactive control showed low similarity with both measures of proactive control, while the two proactive measures showed high similarity with each other. This dissociation suggests that reactive and proactive aspects of inhibitory control rely on distinct neural resources, potentially allowing for independent modulation and development of these strategies.

Modeling trial-level responses at the single-subject level

We used PRAD, a novel hierarchical Bayesian model of proactive and reactive control which represents a significant advance in the assessment of inhibitory control, surpassing the capabilities of conventional race models[44]. Unlike traditional approaches, which provide subject-aggregate SSRT estimates as an index of inhibitory control, PRAD estimates SSRT at the level of each individual trial. Briefly, PRAD models trial-specific SSRT estimates that account for both proactive control strategies and dynamic attentional modulation effects, providing dynamic measures of latent inhibitory control processes[44]. This feature is critical as it allows for precise, trial-specific inferences rather than broad generalizations across the entire task. Importantly, the trial-level model infers additional trial-level measures to the SSRT, such as the probability of proactive cognitive states and the length of proactive delaying of responses.

Trial-level granularity facilitated a more comprehensive examination of how neural responses are modulated across individual trials. Specifically, this model allowed us to identify brain areas that tracked neural activity in response to trial-specific SSRT estimated at each stop trial. Moreover, it enabled tracking ongoing neural dynamics associated with temporal fluctuations in proactive control, providing insights into the brain systems that support a key component of cognitive control.

By employing trial-level analyses, we could dissect proactive and reactive control processes to examine how they fluctuate over time within each individual. This approach deepens our understanding of the mechanisms underlying inhibitory control and enabled us to determine whether assumptions of brain-behavior ergodicity are justified.

Nonergodic brain-behavior associations reveal novel mechanisms of cognitive control

Nonergodicity, a concept originating from physics, describes situations where ensemble averages and time averages do not converge[1]. In the behavioral and life sciences, nonergodicity manifests when measures such as the mean, variance, or covariance differ between population-level analyses (between subjects) and individual-level analyses over time (within subjects)[5,10,11,16]. Our study focused on comparing the covariance of brain activity and behavior between these two levels of analysis, considering divergent associations as evidence of nonergodicity and Simpson's paradox[12,42,50,51].

Our findings reveal pervasive nonergodic patterns in brain-behavior relationships across multiple measures of inhibitory control, encompassing both proactive and reactive control processes. Proactive control, involving anticipation and preparation for stopping, and reactive control,

499 involving the actual implementation of response inhibition, showed markedly different patterns
500 at the within-subjects level compared to the between-subjects level.
501
We found a positive association within subjects between trial-level SSRTs and brain activity in
the anterior and posterior salience networks, which suggests that longer SSRTs, indicating poorer
reactive control, are associated with greater neural effort or engagement. This may reflect
compensatory mechanisms or the increased demand for cognitive resources when individuals
struggle to inhibit their responses. In contrast, between subjects, we observed no significant
association and a negative association, respectively, in the anterior and posterior salience
networks between SSRT and brain activity.
509
Within-subjects analysis revealed a negative association between trial-level engagement of
proactive control and brain activity in the frontoparietal, salience, and subcortical systems. This
is consistent with greater proactive control being associated with a lower need to engage
cognitive control networks that facilitate reactive control, to implement successful response
inhibition. This finding also aligns with recent theoretical frameworks proposing that proactive
control modulates reactive control via preparatory processes[52]. In contrast, we observed a
positive association between trial-level proactive control and default mode network activity. This
included the posterior medial cortex and the ventromedial prefrontal cortex, the two core cortical
nodes that anchor the default mode network[53]. This may reflect internally oriented processing
that supports proactive regulation. Between-subjects analysis failed to capture these dynamic
relationships, showing no association between one measure of proactive control and brain
activations.
522
Our findings highlight the importance of considering within-subjects variability and dynamics
when studying the neural mechanisms of cognitive control. Conventional between-subjects
analyses, which assume ergodicity, may not capture the complex and dynamic nature of
proactive and reactive control processes as they unfold within individuals over time during
development and in psychopathology. Nonergodicity neuroscience approaches using the
methodologies advanced in this study may be crucial for understanding neurocognitive functions
at the individual subject level.
530
531 ***Robustness of nonergodicity findings and stability of within-subjects brain-behavior***
532 ***associations***
533
Leveraging the large-scale ABCD dataset, we addressed the critical challenge of replicability in
human neuroscience[54,55]. Our analyses revealed that within-subjects associations were stable and
reliably detectable, even in sample sizes typical of cognitive neuroscience studies. Bootstrap
resampling analyses showed the reliability of key findings across different sample sizes. For
instance, the association between proactivity measures and right anterior insula suppression was
consistently observed in over 95% of samples, even with sample sizes of $N = 25$. Moreover, our
findings of nonergodicity were robust to various analytical strategies. When comparing the
results of various between- and within-subjects approaches, there were variations in the details of
brain-behavior inferences, but nonergodic dissociations persisted across every approach to brain-
behavior association, including the use of observed dynamic measures (go reaction time) rather
than model-based latent dynamic measures.

545
By demonstrating that these patterns are robust and detectable even in modest sample sizes, our
study provides a foundation for future research into nonergodicity in brain function. It also
suggests that meaningful insights into neurocognitive mechanisms can be gained from studies
with more typical sample sizes, although larger samples provide greater precision and the ability
to detect subtler effects. The stability and robustness of our findings suggests their applicability
to diverse research and clinical contexts, including understanding cognitive processes related to
inhibitory control and studying psychiatric disorders.[56]

553
554 *Nonergodicity between brain networks: Implications for understanding cognitive control*

555
Our analysis of the distribution of nonergodicity between brain networks has implications for
understanding the neural mechanisms underlying inhibitory control as well as for cognitive
neuroscience research broadly. The consistent finding of high nonergodicity in the anterior
salience network across all cognitive model parameters is particularly intriguing. The salience
network is a key brain network, known for its role in detecting behaviorally relevant stimuli and
coordinating brain network dynamics[57,58]. It is noteworthy that this core network, which is of
great interest in the study of cognition[23-26,59] and psychopathology[60-62], appears to exhibit the
most pronounced disconnect between group-level and individual-level inferences. Moreover, the
hierarchical clustering of networks based on nonergodicity profiles provides a fresh perspective
on brain organization. The distinct clustering of the anterior salience network and the default
mode network suggest that nonergodicity may be an important factor in understanding functional
brain architecture.

568
These findings have several significant implications for cognitive neuroscience research and
practice. Methodologically, our results underscore the importance of complementing group-level
analyses with individual-level investigations. The high degree of nonergodicity observed,
particularly in key networks involved in cognitive control, suggests that solely relying on group-
level analyses may lead to incomplete or misleading conclusions about brain-behavior
relationships. From the perspective of individual differences, the varying levels of nonergodicity
across networks highlight the importance of considering individual variability in brain function.
This may be particularly relevant for understanding individual differences in inhibitory control
abilities and for developing personalized interventions for disorders characterized by impaired
inhibitory control. Theoretically, the observed nonergodicity challenges simplistic models of
brain function and calls for quantitatively rigorous theories that can account for the complex,
context-dependent nature of brain-behavior relationships. This may require a shift toward more
dynamic, process-oriented models of cognition and brain function.

582
583 *Distinct neural representations for reactive and proactive cognitive processes*

584
To further elucidate the neural architecture underlying inhibitory control, we employed
representational similarity analysis, a powerful method for investigating the informational
content of brain activity patterns[63]. This approach allows us to compare the similarity of neural
representations across different cognitive processes, providing insights into how the brain
organizes and processes information[64]. In our study, we used representational similarity analysis
to examine the overlap between brain representations of reactivity (SSRT) and proactivity

(probability of proactivity and proactive delaying) within individuals. By comparing
representational patterns across different cognitive processes, we sought to determine whether
reactive and proactive control rely on shared or distinct neural resources.
594
Our analysis revealed a striking dissociation between the neural representations of reactive and
proactive control processes. Across all examined brain networks, we found low similarity
between representations of SSRT and representations of each proactive measure (probability of
proactivity and proactive delaying). In contrast, the two proactive measures showed high
similarity with each other. This pattern suggests that reactive and proactive control processes are
represented orthogonally in the brain. These measures of proactivity affecting the go process and
reactivity affecting the stopping process are more explicitly defined based on naturally occurring
dynamics as inferred by the PRAD model in the standard SST task, rather than being based on
modified versions of the standard task which may attempt to experimentally induce proactive
and reactive aspects of inhibitory control on different trials[29,65-67]. Theoretically, our findings
challenge simplistic models of inhibitory control and suggest that reactive and proactive
processes, while both contributing to inhibitory control, are implemented through distinct neural
mechanisms[46,68].
608
609 Given that our study focused on children, the clear separation of reactive and proactive
610 representations may reflect a developmental stage in the organization of cognitive control
611 processes. The separation we found may allow for independent development of reactive and
612 proactive strategies, potentially explaining individual differences in inhibitory control abilities[46].
613 Future studies could investigate whether this orthogonality persists or changes with age[69].
614
615 ***Attentional modulation and performance monitoring associated with distinct brain-behavior***
616 ***associations***
617
A separate behavioral investigation of our hierarchical Bayesian model highlighted the
significance of adaptive regulation of reactive and proactive control in shaping within-subjects
variability in SSRT and stop failure rates[44]. Two model parameters are decisive in controlling
these dynamics: γ_1 represents individual differences in sustained attention and regulates the trial-
level expectancy of stopping, and θ_1 represents individual differences in performance monitoring
and regulates the trial-level proclivity for proactive control. These findings point to the
importance of considering individual differences in attentional modulation and performance
monitoring systems when studying inhibitory control.
626
Building on these results, we investigated brain-behavior associations between subjects who
differed on these traits. We used γ_1 and θ_1 (separately) as a basis for creating subgroups within
our sample. By dividing our participants into subgroups, we could examine how individual
differences in attentional regulation and performance monitoring correlate with the relationship
between neural activity and cognitive processes within subjects. Subjects stratified based on
whether they adaptively or maladaptively regulated stopping expectancy showed distinct within-
subjects associations between trial-level SSRTs and brain activity, with different distributions of
associations between the subgroups in every network and opposite associations in most
networks. Among all subjects, we observed that anterior salience network activation
accompanied poorer reactive control at the trial level, but examining these results by γ_1 subgroup

revealed that this association only held for subjects who maladaptively regulated reactivity.
Similarly, subjects divided by whether they adaptively or maladaptively regulated proactivity
showed different within-subjects associations for a measure of proactivity in most networks.
With greater trial-level proactivity, the maladaptive regulation group showed weaker suppression
of the anterior salience network and stronger activation of the retrosplenial cortex and
parahippocampal gyrus network, which anchors the ventral posterior aspects of the default mode
network.
644
Collectively, the findings reveal that groups characterized by adaptive and maladaptive
regulation of reactivity and proactivity display notably different patterns of within-subjects
associations between brain activity and model parameters. These distinctions hint that
individuals' distinct cognitive strategies or profiles relate to the implementation of proactive and
reactive control processes in the brain. This variation highlights the personalized nature of
cognitive function and stresses the importance of considering individual differences in the neural
mechanisms of inhibitory control. Identifying heterogeneity based on cognitive model
parameters provides an interpretable approach for studying individual differences in inhibitory
control and their neural correlates. This approach moves beyond simple between-subjects
comparisons and allows for a theory- and mechanism-driven investigation of the heterogeneity in
brain-behavior relationships.
656
657 *Conclusions*
658
Our study provides evidence for nonergodicity in the neurocognitive processes underlying
inhibitory control using a large, community-representative sample of children from the ABCD
study. By combining task fMRI data with a dynamic cognitive model, we found that within-
subjects associations between brain activity and model parameters differed from between-
subjects associations, challenging the assumption of ergodicity in cognitive neuroscience
research. The findings demonstrate divergent group-level and individual-level brain-behavior
associations, reveal dissociated proactive and reactive control systems, and identify meaningful
individual differences in these control processes. Crucially, the study establishes the stability and
robustness of within-subjects measures, laying a foundation for characterizing nonergodic
processes. The work highlights the value of large, heterogeneous samples, dynamic
computational models, and analysis of within-subjects variability for studying nonergodic
phenomena. More broadly, it suggests a paradigm shift away from untested ergodic assumptions
toward nonergodicity neuroscience. Appreciating the nonergodic nature of neurocognitive
processes may be essential for advancing our understanding of cognition in health and disease, as
well as of artificial neural networks.
674

675 **Acknowledgements.** This work was supported by National Institutes of Health MH121069
676 (V.M.), MH124816 (W.C.), National Science Foundation No. 2024856 (V.M.), and the Stanford
677 Maternal and Child Health Research Institute (W.C., P.M.).
678
Contributions. Study design: P.K.M., N.K.B., Z.G., W.C., V.M.; Analysis: P.K.M., N.K.B., Z.G.;
Interpreting results: P.K.M., N.K.B., Z.G., W.C., V.M.; Writing: P.K.M., N.K.B., V.M.; Editing:
P.K.M., N.K.B., Z.G., W.C., V.M.; Supervision: P.K.M., V.M.

Data availability. Data used in this study were from the ABCD study (https://abcdstudy.org/),
held in the National Institute of Mental Health Data Archive. These data are publicly shared with
eligible researchers.
686
687   **Code availability.** All code used in this study will be made available upon publication
688   (https://github.com/scsnl).
689

Methods

Inclusion criteria

Data were from the baseline visit of the ABCD study[18] (Collection #2573), $N = 11817$. Subjects were excluded if they did not meet each of the following criteria: meet the ABCD study's SST task-fMRI inclusion recommendations (in abcd_imgincl01.txt, imgincl_sst_include==1; $N = 3546$ excluded); have 2 SST fMRI runs of good quality (in mriqcrp20301.txt, iqc_sst_total_ser==iqc_sst_good_ser==2; $N = 677$ excluded); are successfully fit with the cognitive model of the SST ($N = 562$ excluded); have 2 SST fMRI runs in the release 4.0 minimally processed data ($N = 16$ excluded); have enough volumes acquired to cover the SST experiment (the last SST trial must have happened no more than 2 seconds after the final volume was acquired; $N = 16$ excluded); have mean framewise displacement of less than 0.5 mm for both runs (calculated using the method of ref. [70]; $N = 1986$ excluded); have release 4.0 minimally processed events.tsv files of shape (181,3) for both runs ($N = 7$ excluded); and have consistent release 4.0 behavioral data (in release 4.0, for some subjects, the "sst.csv" files from ABCD Task fMRI SST Trial Level Behavior, abcd_sst_tlb01, disagreed with the minimally processed "events.tsv" files; for example, one trial might be labeled a go trial by one file and a stop trial by the other; $N = 102$ excluded). Then, we excluded siblings by randomly keeping one member from each family (using the genetic_paired_subjectid variables from gen_y_pihat; $N = 436$ excluded) and excluded subjects without scanner serial number recorded (in mri_y_adm_info, missing mri_info_deviceserialnumber; $N = 10$ excluded). Applying these inclusion criteria left us with a sample of $N = 4469$. For analyses involving the proactive delaying, a further 293 subjects were excluded who had no trials with probability of proactivity greater than 0.5 during at least one run, and therefore, by definition, a proactive delaying of 0 for all trials of at least one run. For these subjects, we were unable to examine within-subjects relationships between proactive delaying and brain activity. To maintain comparability of the between- and within-subjects analyses, we also excluded these subjects from the between-subjects analyses involving proactive delaying. Thus, analyses involving the proactive delaying used a sample of $N = 4176$.

Brain imaging

Imaging acquisition for the ABCD SST is detailed in other work.[71] We used the minimally processed data from ABCD release 4.0 (Collection #2573), which included distortion correction and motion correction[72]. We then further processed the images using Nilearn and FSL FLIRT: (1) initial volumes were removed (Siemens: 8, Philips: 8, GE DV25: 5, GE DV26 and other GE versions: 16); (2) the mean image in the time dimension was computed using mean_img from the Nilearn image module; (3) the mean image was registered to an echo-planar imaging template in MNI152 space (SPM12's toolbox/OldNorm/EPI.nii) using FSL FLIRT, and an affine of this transformation was obtained; (4) the time-series of images was spatially normalized to MNI152 space with the affine from the previous step using FSL FLIRT; and (5) the images were smoothed with a Gaussian filter with a full-width at half maximum of 6 mm using smooth_img from the Nilearn image module.

Bayesian modeling of cognitive dynamics

The PRAD model[44] incorporates latent dynamics that respond to endogenous and exogenous
variables, with trait measures governing the interaction of such endogenous and exogenous
variables with latent processes, giving rise to non-stationary dynamics. This allows the PRAD
model to account for violations of context and stochastic independence. Overall, the PRAD
model incorporates separate evidence accumulation (drift-diffusion) processes for the go and
stop processes, similar to a canonical horse-race model[73]. However, in addition to typical drift-
diffusion process parameters, PRAD includes additional individual trait-like and dynamic trial-
level measures. The full PRAD model is specified below.

(a) The go process is modeled as a drift diffusion process, with a trial-invariant non-decision
time (τ_G) and initial directional bias (β_G), but a trial-varying decision threshold $(\alpha_{G,t})$ and drift
rate $(\delta_{G,t})$. The dynamic decision threshold

760 $$\alpha_{G,t} = \alpha_{G1}\left(1 - \epsilon_{C,t-1}\right) + \alpha_{G2}\left(\epsilon_{C,t-1}\right),$$

749 where $(\alpha_{G1}, \alpha_{G2})$ are distinct threshold levels and $\epsilon_{C,t-1}$ is an indicator of whether the left vs
750 right choice on the previous trial was erroneous. Thus, the dynamic threshold implements a form
751 of performance monitoring and varies between two levels based on the outcome of the previous
752 trials, with α_{G1} being the default threshold and α_{G2} reflecting the threshold after post-error
753 adjustments. The dynamic drift rate

761 $$\delta_{G,t} = \frac{\delta_0\ \mathbb{S}_{LR,t}}{(1 + e^{-12(stim_t-\kappa_0)})} \text{ if } stim_t \neq 0, \text{else } 0.$$

Here δ_0 is a measure of the maximum drift rate for an individual, with the actual drift rate
depending on the duration of the go stimulus $(stim_t)$ and an individual parameter κ_0, which can
be interpreted as the stimulus duration at which the drift rate is half the maximum. $\mathbb{S}_{LR,t}$ assumes
values 1 or -1 depending on the direction of the go stimulus (left or right). PRAD allows for trial
level changes to the drift rate, overcoming the issues with variable go stimulus durations
highlighted in previous work[74].

In addition, in the PRAD model, the onset of the go process may be deliberately delayed in
anticipation of a stop signal. This dynamic adaptation is modeled by adding a further delay ω_t to
the go process to reflect proactive delayed responding to the go stimulus, where

$$\omega_t = \lambda_t\, \rho_t.$$

Here, λ_t reflects a trial-level belief updating process, based on the history of stop signal delays
(SSD) encountered, and is an internal noisy estimate of the prospective anticipated SSD. The
parameter μ $(0 < \mu < 1)$ reflects persistence in belief updating, with high persistence implying a
lower decay rate of older SSDs encountered. Further, letting SSD_t be the SSD and $\mathbb{I}_{S,t}$ be a stop
trial indicator,

$$\lambda_t = \frac{\left(\sum_{i=1}^{t-1} \mu^{t-i-1}\, SSD_i\, \mathbb{I}_{S,i}\right)}{\left(\sum_{i=1}^{t-1} \mu^{t-i-1}\, \mathbb{I}_{S,i}\right)}.$$

ρ_t is a binary variable representing cognitive state. ρ_t indicates the presence $(\rho_t = 1)$ or absence
$(\rho_t = 0)$ of a proactive cognitive state on trial t. The proactive delayed responding is only
initiated on proactive cognitive states. Proactive cognitive states are governed by a baseline
proclivity for proactivity (θ_0), and a performance-monitoring based modulation (θ_1).

777
$$\rho_t \sim \text{Bernoulli} \left(\frac{1}{1 + e^{-\left(\theta_0 + \theta_1 \left(\epsilon_{G,t-1}\right) - \theta_1 \left(\epsilon_{S,t-1}\right)\right)}} \right).$$

Here, $\epsilon_{G,t-1}$ is an indicator of a go-omission (incorrectly stopping on a go trial) on the previous
trial, and $\epsilon_{S,t-1}$ is an indicator of a stopping error (not stopping on a stop trial). The PRAD model
assumes that the correction in terms of increasing or decreasing the probability of a proactive
cognitive state following these two types of trials will be in opposite directions. The sign of θ_1 is
an indicator of adaptivity or maladaptivity of the performance monitoring mechanism, and the
absolute value of θ_1 denotes the sensitivity of the state-switching mechanism to errors.
784
785 (b) The stop process is modeled as a drift diffusion process with a trial-invariant non-decision
786 time (τ_S), decision threshold (α_S), and drift rate (δ_S), but a trial-varying bias ($\beta_{S,t}$). The stop
787 process begins at the onset of the stop signal. The initial bias is

788
$$\beta_{S,t} = \left(\frac{1}{1 + e^{-\gamma_1 (nSSD_t - \gamma_0)}} \right).$$

Here, $nSSD_t$ reflects the number of trials since a stop signal was last encountered. This reflects
an attentional mechanism that modulates the stopping bias $\beta_{S,t}$ (which varies from 0 to 1).
Positive values of γ_1 result in an increase in stopping bias as $nSSD_t$ increases and vice versa.
Similarly, negative values of γ_1 result in a decrease in stopping bias as $nSSD_t$ increases and vice
versa. The absolute value of γ_1 measures the sensitivity to attentional modulation. The γ_0
parameter is a measure of the value of $nSSD_t$ when stopping bias is neutral (0.5).
795
Both the go and stop processes are implemented within a hierarchical Bayesian modeling
framework in JAGS[75], using the Wiener distribution[76], which produces a joint distribution of the
reaction times and the decision choice on each trial. The reaction times of the go process
correspond to the reaction times for pressing the left or right buttons in response to the go
stimulus. The reaction times of the stop process correspond to the SSRT. The stop process is only
initiated on stop trials after the appearance of the stop stimulus (which appears after a delay
corresponding to the SSD). The SSRT is not manifested as a behavioral action. Rather, if the
SSRT, which is the duration of the stop process, plus the SSD on a stop trial is smaller than the
go process reaction time, then the go action can be successfully inhibited (successful stop). The
interaction of the basic go and stop processes can be influenced by the dynamics of the proactive
delayed responding as well as the dynamics of the attentional modulation of reactive stopping.
The PRAD model enables obtaining the full posterior distributions of SSRT, proactive delay in
responding of the go process, and the probability of proactive cognitive states at a trial level.
809
Markov Chain Monte Carlo was run using 4 chains with 30,000 samples each. For each chain,
we had a burn-in of 10,000 samples, and a thinning factor of 10 was used to record 2000 of the
remaining 20,000 samples, based on which final inferences were made. The following priors
were used. $\delta_G \sim \text{Uniform}(0.0001,12)$. $\alpha_{G1} \sim \text{Uniform}(0.5,4)$. $\alpha_{G2} \sim \text{Uniform}(0.5,4)$.
$\kappa_0 \sim \text{Normal}(0.1, \text{precision} = 0.3)$ truncated to $(0,0.3)$. $\beta_G \sim \text{Beta}(1,1)$ truncated to
$(0.0001,0.9999)$. $\delta_S = p/\alpha_S$, where $p \sim \text{Uniform}(4,24)$ and $\alpha_S \sim \text{Uniform}(0.5,6)$.
$\theta_0 \sim \text{Normal}(0, \text{precision} = 0.3)$. $\theta_1 \sim \text{Normal}(0, \text{precision} = 0.3)$. $\mu \sim \text{Beta}(0.5,0.5)$ truncated
to $(0.8,0.9999)$. $\gamma_1 \sim \text{Normal}(0, \text{precision} = 0.3)$. $\gamma_0 \sim \text{Uniform}(0,12)$.
$\tau_G \sim \text{Uniform}(0.0001, \min(0.1, RT^*))$, where RT^* is the reaction time after censoring for
extremely small values. $\tau_S \sim \text{Uniform}(0.0001, \tau_G)$. For further details of the model, see ref. [44].

We defined the 3 latent dynamic measures examined in this study as follows. SSRT was the
posterior mean of the duration of the stop process on correct or incorrect stop trials; it was not
defined for go trials or trials on which a participant responded before the stop signal was shown.
Probability of proactivity was the posterior mean of ρ_t, i.e., the posterior probability of $\rho_t = 1$.
Proactive delaying was the posterior mean of λ_t on trials with probability of proactivity greater
than 0.5, otherwise 0. Probability of proactivity and proactive delaying were defined for all trials.

Between- and within-subjects, general linear model analysis of fMRI

We fit general linear models to the fMRI BOLD recordings using Nilearn's FirstLevelModel.
Condition and parametric regressors were modeled as impulses, with a duration of 0, and
convolved with the SPM software's double gamma hemodynamic response function and the
function's time derivative. Before fitting, the BOLD signal was scaled to percent signal-change
from the mean in the time dimension. An AR1 model was used to whiten the data and design
matrices to account for temporal autocorrelation in the BOLD signal.

To determine between-subjects associations between subject-average brain activity and
behavioral measures, for each subject and voxel, we fit the model

$$\text{BOLD}(t) = \beta_0 + (\text{HRF} * \text{Conditions})(t) + \text{Nuissance}(t) + \epsilon(t). \qquad \text{(Model 1)}$$

$\text{BOLD}(t)$ is the BOLD signal of the voxel at time t ($t \in \{1, \ldots, T\}$ for T the total number of
volumes acquired); $(\text{HRF} * \text{Conditions})(t)$ is the value at t of the convolution with the
hemodynamic response function HRF of condition regressor(s) Conditions; $\text{Nuissance}(t)$ is the
effect at t of nuisance regressors, which were 6 motion parameters (translational and rotational
displacement along each of three axes) and 6 cosine basis functions (corresponding to high-pass
filtering at 0.01 Hz); and $\epsilon(t)$ is the model's error at t. We fit models with three sets of condition
regressors:

$$\text{Conditions} = \beta_1 (I_{\text{Go}} + I_{\text{Stop}}). \qquad \text{(Conditions 1)}$$

$$\text{Conditions} = \beta_1 I_{\text{Go}} + \beta_2 I_{\text{Stop}}. \qquad \text{(Conditions 2)}$$

$$\text{Conditions} = \beta_1 I_{\text{Correct go}} + \beta_2 I_{\text{Incorrect go}} + \beta_3 I_{\text{Correct late go}} + \beta_4 I_{\text{Incorrect late go}}$$
$$+ \beta_5 I_{\text{No response go}} + \beta_6 I_{\text{Correct stop}} + \beta_7 I_{\text{Incorrect stop}} + \beta_8 I_{\text{SSD stop}}. \qquad \text{(Conditions 3)}$$

$I_{\text{Condition}}$ is an indicator function indicating when the subject experiences Condition (for
example, I_{Go} is 0 except at the moment when a subject is presented with a go trial). For between-
subjects analyses, we used Model 1 with Conditions 1 to obtain task activation (β_1); used
Model 1 with Conditions 2 to obtain go activation (β_1) and stop activation (β_2); and used
Model 1 with Conditions 3 to obtain correct stop versus correct go activation ($\beta_6 - \beta_1$), correct
stop versus incorrect go activation ($\beta_6 - \beta_2$), and incorrect stop versus correct stop activation
($\beta_7 - \beta_6$). For correlations between these activations and subject-average behavioral measures,
we computed each subject-average measure as the mean of the measure over the trials from both
runs during which it assumed values; that is, SSRT was averaged over correct and incorrect stop
trials, probability of proactivity and proactive delaying over all trials, and go reaction time over
go trials with a recorded response.

To determine within-subjects associations between trial-level brain activity and behavioral
measures, for each subject and voxel, we fit the model

$$BOLD(t) = \beta_0 + (HRF * Conditions)(t) + (HRF * Modulation)(t)$$
$$+Nuissance(t) + \epsilon(t). \qquad \text{(Model 2)}$$

$(HRF * Modulation)(t)$ is the value at t of the convolution with the hemodynamic response function of the parametric regressor Modulation. To investigate within-subjects associations between brain activity and SSRT on stop trials, probability of proactivity on all trials, proactive delaying on all trials, and observed reaction time on go trials, we set Modulation $= \beta_3$SSRT, Modulation $= \beta_3$P(Proactive), Modulation $= \beta_3$Proactive delaying, and Modulation $= \beta_3$Go RT, respectively, and used Model 2 with Conditions 2. Each of SSRT, P(Proactive), Proactive delaying, and Go RT was standardized over the conditions during which it assumed values by subtracting its mean and dividing by its standard deviation; that is, SSRT had mean 0 and standard deviation 1 over correct and incorrect stop trials (and was 0 on all other trials), P(Proactive) and Proactive delaying had mean 0 and standard deviation 1 over all trials, and Go RT had mean 0 and standard deviation 1 over go trials with a recorded response (and was 0 on all other trials).

We fit these regression models for each subject and each of their 2 SST runs. For each model, subject, and voxel, we combined the regression results from the 2 runs with a fixed effects model through FirstLevelModel's compute_contrast method. For each model and voxel, we estimated the effect of each scanner as the mean of the regression coefficients of the subjects who were scanned by it minus the grand mean of the regression coefficients of all subjects. Then, we adjusted the regression coefficients by subtracting the estimated scanner effects. All analyses of the regression coefficients used these adjusted values. We used a 1-sample Cohen's d (sample mean divided by sample standard deviation) to measure the effect sizes of regression coefficients. Whole-brain correlations and Cohen's d values were visualized on the cortical surface[77]. The surface was obtained using the fetch_surf_fsaverage function in Nilearn's struct module (mesh="fsaverage"); data were sampled to the surface using the vol_to_surf function in Nilearn's surface module; and plotted using the plot_surf_stat_map function in the surf_plotting module.

Networks and regions of interest

We extracted the whole-brain regression coefficients in 2 networks and 3 sets of regions. We used the Shirer networks for our primary analyses and used the Yeo-17 networks, Shirer regions, cognitive-control regions, and subcortical regions to test the stability of our within-subjects findings. For each subject and each regression coefficient of interest, we obtained the coefficient's value in each area (network or region) by calculating the mean of the subject's coefficients over the voxels belonging to the area. We used these area-average regression coefficients of each subject: to compute Cohen's d and Pearson r values for the network-level comparison of between- and within-subjects associations (Figure 4) and for the measurement of nonergodicity (Figure 8); and to compute Cohen's d values for the stability (Figure 5) and subgroup (Figure 7) analyses.

The Shirer networks and regions were obtained from ref. [48]. To obtain the voxel-coordinates of the Yeo-17 networks, we used a mapping between the Brainnetome[78] and Yeo atlases[79]. We assembled the cognitive-control regions to include 8 areas activated by the SST, 2 core default mode areas, and 1 core salience network and error-processing area. The areas activated by the

SST were taken from a meta-analysis of 70 inhibitory control studies[23] (right anterior insula, right caudate, right inferior frontal gyrus, right middle frontal gyrus, right presupplementary motor area, and right supramarginal gyrus) and a study that segmented high-resolution structural MRI[80] (left and right subthalamic nucleus). To obtain the default mode areas (posterior cingulate cortex and ventromedial prefrontal cortex), we retrieved a Neurosynth automatic meta-analysis of 777 studies for the term "default mode" on 2024-02-05; extracted clusters from this map using the connected_regions function in Nilearn's region_extractor module with keyword argument "extract_type" set to "connected_components"; identified by eye the clusters corresponding to the posterior cingulate and ventromedial prefrontal cortex; and for each cluster defined the region to be the 6 mm cube centered on the voxel with the highest meta-analysis Z-score in the cluster. To obtain the salience network and error-processing area (dorsal anterior cingulate cortex), we retrieved a Neurosynth automatic meta-analysis of 464 studies for the term "error" on 2023-09-21 and defined the region to be the 6 mm cube centered on the voxel with the highest meta-analysis Z-score. We obtained subcortical regions from a subcortical probabilistic atlas[81]. We resampled the atlas's probabilistic subcortical labels in 1 mm cubed MNI152 2009c nonlinear asymmetric space to the 2 mm cubed MNI152 space of our SPM echo-planar imaging template using resample_to_img from Nilearn's resampling module, and then thresholded these probabilistic maps at 0.5 to obtain region masks.

Stability analysis

To assess the stability of the within-subjects results, regression coefficients were resampled at varying sample sizes and the correlation was evaluated against the results in the full sample. Specifically, for each of SSRT, probability of proactivity, and proactive delaying, in each set of networks or regions: 10,000 samples of n subjects were drawn with replacement; the Cohen's d's of each sample's regression coefficients were calculated and Pearson correlated with the Cohen's d's of the full sample over the regions or networks; and the mean and 95% and 99% bootstrap confidence intervals were calculated of the correlation (SSRT and probability of proactivity $n =$ 25, 40, 70, 120, 200, 335, 560, 945, 1585, 2660, 4469; proactive delaying $n = 25, 40, 70, 115,$ 195, 325, 540, 900, 1500, 2505, 4176). The 95% and 99% bootstrap confidence intervals were calculated, respectively, as the intervals covering the 2.5th to 97.5th percentiles and 0.5th to 99.5th percentiles of the 10,000 correlations at each n.

We also directly examined the distributions of the Cohen's d's of the resamples as a function of n in regions of interest. Specifically, for each of SSRT, probability of proactivity, and proactive delaying, in each region of interest: 10,000 samples of n subjects were drawn with replacement; the Cohen's d of each sample's regression coefficients was calculated; and the mean and 95% and 99% bootstrap confidence intervals were calculated of the Cohen's d (SSRT and probability of proactivity $n = 25, 40, 70, 120, 200, 335, 560, 945, 1585, 2660, 4469$; proactive delaying $n =$ 25, 40, 70, 115, 195, 325, 540, 900, 1500, 2505, 4176). The 95% and 99% bootstrap confidence intervals were calculated as the intervals covering, respectively, the 2.5th to 97.5th percentiles and 0.5th to 99.5th percentiles of the 10,000 Cohen's d's at each n.

Representational similarity analysis

For each Shirer network, and for each subject, we computed the correlation between the subject's
within-subjects brain maps of SSRT and probability of proactivity, SSRT and proactive delaying,
and probability of proactivity and proactive delaying over the voxels in the network. Density
estimates used Seaborn's kdeplot function with each distribution of correlations over subjects
normalized to 1 (common_norm=False) and limited to values between -1 and 1 (clip=[-1,1]); all
other parameters, including those determining the kernel smoothing bandwidth, were kept at
their defaults.
961
962   *Measuring nonergodicity of brain networks*
963
For each of SSRT, probability of proactivity, and proactive delaying, and for each of the Shirer
networks, we computed bootstrap distributions of a measure of nonergodicity: we drew 10,000
samples of n subjects with replacement; for each resample, using the resample's between- and
within-subjects associations, we computed the fraction of subjects whose (within-subjects) brain
association with the parameter had the opposite sign of the between-subjects brain association
with the parameter, in the network; then, we computed the mean and 95% confidence interval
over the resamples of the fraction of opposite signs (SSRT and probability of proactivity $n = $
4469; proactive delaying $n = 4176$). The between-subjects association was the Pearson
correlation between correct stop versus correct go activation and the subject-average parameter,
which was recomputed for the subjects in each resample. The 95% confidence interval was
calculated as the interval covering the 2.5th to 97.5th percentiles of the 10,000 fractions of
opposite signs. The goal of using bootstrapping was to account for the strength of the between-
subjects results. Next, the nonergodicity of each network was defined as the three-dimensional
vector whose ith coordinate was the mean (over bootstrap resamples) fraction of subjects with
opposite signs in the network for the ith cognitive model parameter. Then, the Euclidean
distances were computed between the vectors and hierarchical clustering was performed on the
distances using the linkage function in scipy's cluster subpackage, hierarchy module with
"method" set to "average".
982
983   *Statistical testing*
984
985   Permutation tests were used for all statistical testing. The tests used two-sided alternatives and
986   10,000 resamples and were performed with Scipy's permutation_test function. FDR correction
987   was performed using the Benjamini-Hochberg procedure with Scipy's false_discovery_control
988   function. P_{FDR} denotes an FDR-corrected P value.
989
For the Shirer networks and an fMRI regression, we tested the null hypothesis for each network
that the regression coefficients in the network had a mean of 0 (Figure 4) by computing the
means of resamples in which the signs of the coefficients were randomly chosen
(permutation_test permutation_type='samples'). Then, FDR correction was applied to the P
values of all networks (e.g., FDR correction was applied to the P's of SSRT's regression
coefficients over the Shirer networks).
996
For the Shirer networks, an fMRI regression, and a behavioral measure, we tested the null
hypothesis for each network that the Pearson correlation between subject-average regression
coefficients in the network and subject-average behavioral measures was 0 (Figure 4) by

computing the Pearson correlations of resamples in which regression coefficients were randomly
paired with behavioral measures (permutation_test permutation_type ='pairings'). Then, FDR
correction was applied to the P values of all networks (e.g., FDR correction was applied to the
P's of correlations between correct stop versus correct go activation and SSRT over the Shirer
networks).
1005
For the Shirer networks and a param γ_1 or θ_1, we tested the null hypothesis for each network
that the mutually exclusive subgroups of subjects with param < 0 and with param > 0 had
different mean regression coefficients (Figure 7) by computing the differences between the
means for resamples in which coefficients were randomly assigned to param < 0 and param $>$
0 (permutation_test permutation_type='independent'). Then, FDR correction was applied to the
P values of all networks (e.g., FDR correction was applied to the P's of mean differences
between $\gamma_1 < 0$ and $\gamma_1 > 0$ over the Shirer networks).
1013
For the Shirer networks, we tested the null hypothesis for each network that there was no
difference in the network between the median correlation of SSRT and probability of proactivity
and the median correlation of SSRT and proactive delaying; the median correlation of SSRT and
probability of proactivity and the median correlation of probability of proactivity and proactive
delaying; and the median correlation of SSRT and proactive delaying and the median correlation
of probability of proactivity and proactive delaying (Figure 6). We computed the differences
between the medians of resamples in which correlations were randomly exchanged within
subjects (permutation_test permutation_type='samples'). Then, FDR correction was applied to
the P values of all comparisons in all networks. (Since there are 14 Shirer networks and 3 tests
per network, FDR correction was applied over 14×3 P's.)
1024
1025 ***Software***
1026
1027 Data were processed and analyzed using Python (version 3.9.16), Scipy (version 1.11.4),
1028 Seaborn (version 0.13.2), Nilearn (version 0.10.1), FSL FLIRT (version 6.0), MATLAB (version
1029 R2020b), and JAGS[75] (version 4.3.0).

References

Lebowitz, J. L. & Penrose, O. Modern ergodic theory. *Physics Today* **26**, 23-29 (1973).

Badino, M. The foundational role of ergodic theory. *Foundations of science* **11**, 323-347 (2006).

Moore, C. C. Ergodic theorem, ergodic theory, and statistical mechanics. *Proceedings of the National Academy of Sciences* **112**, 1907-1911 (2015).

Peters, O. The ergodicity problem in economics. *Nature Physics* **15**, 1216-1221 (2019).

Fisher, A. J., Medaglia, J. D. & Jeronimus, B. F. Lack of group-to-individual generalizability is a threat to human subjects research. *Proc Natl Acad Sci U S A* **115**, E6106-E6115, doi:10.1073/pnas.1711978115 (2018).

Molenaar, P. C. On the implications of the classical ergodic theorems: Analysis of developmental processes has to focus on intra-individual variation. *Developmental Psychobiology: The Journal of the International Society for Developmental Psychobiology* **50**, 60-69 (2008).

Dhakal, V., Feit, A. M., Kristensson, P. O. & Oulasvirta, A. in *Proceedings of the 2018 CHI conference on human factors in computing systems.* 1-12.

Heitz, R. P. The speed-accuracy tradeoff: history, physiology, methodology, and behavior. *Frontiers in neuroscience* **8**, 86875 (2014).

Molenaar, P. & Newell, K. M. *Individual pathways of change: Statistical models for analyzing learning and development.* (American Psychological Association, 2010).

Molenaar, P. C. A manifesto on psychology as idiographic science: Bringing the person back into scientific psychology, this time forever. *Measurement* **2**, 201-218 (2004).

Medaglia, J. D., Ramanathan, D. M., Venkatesan, U. M. & Hillary, F. G. The challenge of non-ergodicity in network neuroscience. *Network: Computation in Neural Systems* **22**, 148-153 (2011).

Kievit, R. A., Frankenhuis, W. E., Waldorp, L. J. & Borsboom, D. Simpson's paradox in psychological science: a practical guide. *Frontiers in psychology* **4**, 513 (2013).

Hamaker, E. L. Why researchers should think" within-person": A paradigmatic rationale. (2012).

Castro-Schilo, L. & Ferrer, E. Comparison of nomothetic versus idiographic-oriented methods for making predictions about distal outcomes from time series data. *Multivariate Behavioral Research* **48**, 175-207 (2013).

Hamaker, E., Ceulemans, E., Grasman, R. & Tuerlinckx, F. Modeling affect dynamics: State of the art and future challenges. *Emotion Review* **7**, 316-322 (2015).

Molenaar, P. C. & Campbell, C. G. The new person-specific paradigm in psychology. *Current directions in psychological science* **18**, 112-117 (2009).

Curran, P. J. & Bauer, D. J. The disaggregation of within-person and between-person effects in longitudinal models of change. *Annual review of psychology* **62**, 583-619 (2011).

Volkow, N. D. *et al.* The conception of the ABCD study: From substance use to a broad NIH collaboration. *Dev Cogn Neurosci* **32**, 4-7, doi:10.1016/j.dcn.2017.10.002 (2018).

Diamond, A. Executive functions. *Annual review of psychology* **64**, 135-168 (2013).

Logan, G. D. & Cowan, W. B. On the ability to inhibit thought and action: A theory of an act of control. *Psychological review* **91**, 295 (1984).

21   Verbruggen, F. & Logan, G. D. Response inhibition in the stop-signal paradigm. *Trends*
in cognitive sciences **12**, 418-424 (2008).
22   Aron, A. R., Robbins, T. W. & Poldrack, R. A. Inhibition and the right inferior frontal
cortex: one decade on. *Trends in cognitive sciences* **18**, 177-185 (2014).
23   Cai, W., Ryali, S., Chen, T., Li, C. S. & Menon, V. Dissociable roles of right inferior
frontal cortex and anterior insula in inhibitory control: evidence from intrinsic and task-
related functional parcellation, connectivity, and response profile analyses across multiple
datasets. *J Neurosci* **34**, 14652-14667, doi:10.1523/JNEUROSCI.3048-14.2014 (2014).
24   Cai, W. *et al.* Hyperdirect insula-basal-ganglia pathway and adult-like maturity of global
brain responses predict inhibitory control in children. *Nature communications* **10**, 4798
(2019).
25   Cai, W., Griffiths, K., Korgaonkar, M. S., Williams, L. M. & Menon, V. Inhibition-related
modulation of salience and frontoparietal networks predicts cognitive control ability and
inattention symptoms in children with ADHD. *Molecular psychiatry* **26**, 4016-4025
(2021).
26   Cai, W., Chen, T., Ide, J. S., Li, C.-S. R. & Menon, V. Dissociable fronto-operculum-
insula control signals for anticipation and detection of inhibitory sensory cue. *Cerebral*
cortex **27**, 4073-4082 (2017).
27   Friedman, N. P. & Robbins, T. W. The role of prefrontal cortex in cognitive control and
executive function. *Neuropsychopharmacology* **47**, 72-89 (2022).
28   Xue, G., Aron, A. R. & Poldrack, R. A. Common neural substrates for inhibition of
spoken and manual responses. *Cereb Cortex* **18**, 1923-1932, doi:10.1093/cercor/bhm220
(2008).
29   Swann, N. C. *et al.* Roles for the pre-supplementary motor area and the right inferior
frontal gyrus in stopping action: electrophysiological responses and functional and
structural connectivity. *Neuroimage* **59**, 2860-2870,
doi:10.1016/j.neuroimage.2011.09.049 (2012).
30   Li, C. S., Yan, P., Sinha, R. & Lee, T. W. Subcortical processes of motor response
inhibition during a stop signal task. *Neuroimage* **41**, 1352-1363,
doi:10.1016/j.neuroimage.2008.04.023 (2008).
31   Leung, H. C. & Cai, W. Common and differential ventrolateral prefrontal activity during
inhibition of hand and eye movements. *J Neurosci* **27**, 9893-9900,
doi:10.1523/jneurosci.2837-07.2007 (2007).
32   Cai, W., Cannistraci, C. J., Gore, J. C. & Leung, H. C. Sensorimotor-independent
prefrontal activity during response inhibition. *Hum Brain Mapp* **35**, 2119-2136,
doi:10.1002/hbm.22315 (2014).
33   Aron, A. R. & Poldrack, R. A. Cortical and subcortical contributions to Stop signal
response inhibition: role of the subthalamic nucleus. *J Neurosci* **26**, 2424-2433,
doi:10.1523/jneurosci.4682-05.2006 (2006).
34   Aron, A. R., Behrens, T. E., Smith, S., Frank, M. J. & Poldrack, R. A. Triangulating a
cognitive control network using diffusion-weighted magnetic resonance imaging (MRI)
and functional MRI. *J Neurosci* **27**, 3743-3752, doi:10.1523/jneurosci.0519-07.2007
(2007).
35   Zhang, S. *et al.* Independent component analysis of functional networks for response
inhibition: Inter-subject variation in stop signal reaction time. *Human brain mapping* **36**,
3289-3302 (2015).

36   Boehler, C. N., Appelbaum, L. G., Krebs, R. M., Hopf, J.-M. & Woldorff, M. G. The
influence of different Stop-signal response time estimation procedures on behavior–
behavior and brain–behavior correlations. *Behavioural brain research* **229**, 123-130
(2012).
37   Aron, A. R. & Poldrack, R. A. Cortical and subcortical contributions to stop signal
response inhibition: role of the subthalamic nucleus. *Journal of Neuroscience* **26**, 2424-
2433 (2006).
38   Li, C. S., Huang, C., Constable, R. T. & Sinha, R. Imaging response inhibition in a stop-
signal task: neural correlates independent of signal monitoring and post-response
processing. *J Neurosci* **26**, 186-192, doi:10.1523/JNEUROSCI.3741-05.2006 (2006).
39   Li, C. S. *et al.* Neural correlates of impulse control during stop signal inhibition in
cocaine-dependent men. *Neuropsychopharmacology* **33**, 1798-1806,
doi:10.1038/sj.npp.1301568 (2008).
40   Chevrier, A. & Schachar, R. J. BOLD differences normally attributed to inhibitory control
predict symptoms, not task-directed inhibitory control in ADHD. *J Neurodev Disord* **12**,
8, doi:10.1186/s11689-020-09311-8 (2020).
41   Chaarani, B. *et al.* Baseline brain function in the preadolescents of the ABCD Study. *Nat
Neurosci* **24**, 1176-1186, doi:10.1038/s41593-021-00867-9 (2021).
42   Mangalam, M. & Kelty-Stephen, D. G. Point estimates, Simpson's paradox, and
nonergodicity in biological sciences. *Neurosci Biobehav Rev* **125**, 98-107,
doi:10.1016/j.neubiorev.2021.02.017 (2021).
43   Verbruggen, F. *et al.* A consensus guide to capturing the ability to inhibit actions and
impulsive behaviors in the stop-signal task. *Elife* **8**, doi:10.7554/eLife.46323 (2019).
44   Mistry, P. K., Warren, S. L., Branigan, N. K., Cai, W. & Menon, V. Computational
Modeling of Proactive, Reactive, and Attentional Dynamics in Cognitive Control.
bioRxiv, doi:10.1101/2024.10.01.615613 (2024).
45   Verbruggen, F. & Logan, G. D. Proactive adjustments of response strategies in the stop-
signal paradigm. *Journal of Experimental Psychology: Human Perception and
Performance* **35**, 835 (2009).
46   Braver, T. S. The variable nature of cognitive control: a dual mechanisms framework.
Trends in cognitive sciences **16**, 106-113 (2012).
47   Doekemeijer, R. A., Dewulf, A., Verbruggen, F. & Boehler, C. N. Proactively adjusting
stopping: response inhibition is faster when stopping occurs frequently. *Journal of
cognition* **6** (2023).
48   Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V. & Greicius, M. D. Decoding
subject-driven cognitive states with whole-brain connectivity patterns. *Cereb Cortex* **22**,
158-165, doi:10.1093/cercor/bhr099 (2012).
49   Aron, A. R. *et al.* Converging evidence for a fronto-basal-ganglia network for inhibitory
control of action and cognition. *Journal of Neuroscience* **27**, 11860-11864 (2007).
50   Sprenger, J. & Weinberger, N. Simpson's paradox. *The Stanford Encyclopedia of
Philosophy* (2021).
51   Pearl, J. in *Probabilistic and causal inference: The works of judea Pearl*   399-412
(2022).
52   van den Wildenberg, W. P., Ridderinkhof, K. R. & Wylie, S. A. Towards Conceptual
Clarification of Proactive Inhibitory Control: A Review. *Brain Sciences* **12**, 1638 (2022).
53   Menon, V. 20 years of the default mode network: A review and synthesis. *Neuron* (2023).

54   Button, K. S. *et al.* Power failure: why small sample size undermines the reliability of
neuroscience. *Nat Rev Neurosci* **14**, 365-376, doi:10.1038/nrn3475 (2013).
55   Munafò, M. R. *et al.* A manifesto for reproducible science. *Nat Hum Behav* **1**, 0021,
doi:10.1038/s41562-016-0021 (2017).
56   Meyer-Lindenberg, A. The non-ergodic nature of mental health and psychiatric disorders:
implications for biomarker and diagnostic research. *World Psychiatry* **22** (2023).
57   Craig, A. D., Chen, K., Bandy, D. & Reiman, E. M. Thermosensory activation of insular
cortex. *Nature neuroscience* **3**, 184-190 (2000).
58   Menon, V. Insular cortex: A hub for saliency, cognitive control, and interoceptive
awareness. (2024).
59   Molnar-Szakacs, I. & Uddin, L. Q. Anterior insula as a gatekeeper of executive control.
Neuroscience & Biobehavioral Reviews **139**, 104736 (2022).
60   Uddin, L. Q. & Menon, V. The anterior insula in autism: under-connected and under-
examined. *Neuroscience & Biobehavioral Reviews* **33**, 1198-1203 (2009).
61   Segal, A. *et al.* Regional, circuit and network heterogeneity of brain abnormalities in
psychiatric disorders. *Nature Neuroscience* **26**, 1613-1629 (2023).
62   Xie, C. *et al.* A shared neural basis underlying psychiatric comorbidity. *Nature medicine*
29, 1232-1242 (2023).
63   Kriegeskorte, N., Mur, M. & Bandettini, P. A. Representational similarity analysis-
connecting the branches of systems neuroscience. *Frontiers in systems neuroscience* **2**,
249 (2008).
64   Haxby, J. V., Connolly, A. C. & Guntupalli, J. S. Decoding neural representational spaces
using multivariate pattern analysis. *Annual review of neuroscience* **37**, 435-456 (2014).
65   Cunillera, T., Fuentemilla, L., Brignani, D., Cucurell, D. & Miniussi, C. A simultaneous
modulation of reactive and proactive inhibition processes by anodal tDCS on the right
inferior frontal cortex. *PloS one* **9**, e113537 (2014).
66   Messel, M. S., Raud, L., Hoff, P. K., Skaftnes, C. S. & Huster, R. J. Strategy switches in
proactive inhibitory control and their association with task-general and stopping-specific
networks. *Neuropsychologia* **135**, 107220 (2019).
67   Cai, W., George, J. S., Verbruggen, F., Chambers, C. D. & Aron, A. R. The role of the
right presupplementary motor area in stopping action: two studies with event-related
transcranial magnetic stimulation. *J Neurophysiol* **108**, 380-389,
doi:10.1152/jn.00132.2012 (2012).
68   Aron, A. R. From reactive to proactive and selective control: developing a richer model
for stopping inappropriate responses. *Biological psychiatry* **69**, e55-e68 (2011).
69   Luna, B., Marek, S., Larsen, B., Tervo-Clemmens, B. & Chahal, R. An integrative model
of the maturation of cognitive control. *Annual review of neuroscience* **38**, 151-170
(2015).
70   Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious
but systematic correlations in functional connectivity MRI networks arise from subject
motion. *Neuroimage* **59**, 2142-2154, doi:10.1016/j.neuroimage.2011.10.018 (2012).
71   Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD) study: Imaging
acquisition across 21 sites. *Dev Cogn Neurosci* **32**, 43-54, doi:10.1016/j.dcn.2018.03.001
(2018).

72   Hagler, D. J., Jr. *et al.* Image processing and analysis methods for the Adolescent Brain
Cognitive Development Study. *Neuroimage* **202**, 116091,
doi:10.1016/j.neuroimage.2019.116091 (2019).
73   Band, G. P., Van Der Molen, M. W. & Logan, G. D. Horse-race model simulations of the
stop-signal procedure. *Acta psychologica* **112**, 105-142 (2003).
74   Bissett, P. G., Hagen, M. P., Jones, H. M. & Poldrack, R. A. Design issues and solutions
for stop-signal data from the Adolescent Brain Cognitive Development (ABCD) study.
Elife **10**, e60185 (2021).
75   Plummer, M. in *Proceedings of the 3rd international workshop on distributed statistical
computing.* 1-10 (Vienna, Austria).
76   Wabersich, D. & Vandekerckhove, J. Extending JAGS: A tutorial on adding custom
distributions to JAGS (with a diffusion model example). *Behavior research methods* **46**,
15-28 (2014).
77   Fischl, B., Sereno, M. I., Tootell, R. B. & Dale, A. M. High-resolution intersubject
averaging and a coordinate system for the cortical surface. *Human brain mapping* **8**, 272-
284 (1999).
78   Fan, L. *et al.* The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional
Architecture. *Cereb Cortex* **26**, 3508-3526, doi:10.1093/cercor/bhw157 (2016).
79   Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic
functional connectivity. *J Neurophysiol* **106**, 1125-1165, doi:10.1152/jn.00338.2011
(2011).
80   Forstmann, B. U. *et al.* Cortico-subthalamic white matter tract strength predicts
interindividual efficacy in stopping a motor response. *Neuroimage* **60**, 370-375,
doi:10.1016/j.neuroimage.2011.12.044 (2012).
81   Pauli, W. M., Nili, A. N. & Tyszka, J. M. A high-resolution probabilistic in vivo atlas of
human subcortical brain nuclei. *Sci Data* **5**, 180063, doi:10.1038/sdata.2018.63 (2018).
1237
1238

Figures

Figure 1. Conceptual overview of the study and key findings. The figure illustrates the
methodology for between-subjects and within-subjects analyses, the concept of nonergodicity,
and previews the main results. **a,** Between-subjects analysis. Subject-average brain activation in
each voxel is correlated with a subject-average cognitive measure across the population. **b,**
Within-subjects analysis. For each individual, the time-series of brain activity in each voxel is
associated with the time-series of a cognitive measure. **c,** Simpson's paradox. Simpson's paradox
occurs when associations disagree between subjects and within subjects; it exemplifies
nonergodicity in the behavioral sciences. **d,** Study aims. We examined brain-behavior
associations for nonergodicity; tested our within-subjects results for stability and robustness;
used these results to probe the brain implementations of proactive and reactive control and
adaptive regulation of inhibitory control; and investigated how nonergodicity varied by brain
network. Nonergodic patterns in brain-behavior associations were consistently observed,
revealing that group-level (between-subjects) and individual-level (within-subjects) associations
yield divergent results for inhibitory control processes. This challenges the common assumption
that findings from such group-level analyses can be directly applied to understand individual-
level cognitive processes.

Figure 2. Computational modeling of inhibitory control dynamics in the stop signal task. a, Stop signal task. On go trials subjects should respond by pressing a button to indicate the arrow direction, and on stop trials should inhibit their response when the stop signal appears. **b,** Computational model schematic. The computational model infers individual-level and trial-level parameters that measure inhibitory control from observed stop failure and response times. SSRT is the time it takes for the stop process to complete. Probability of proactivity is the probability that a subject uses a proactive control strategy. Proactive delaying is the delay in initiating the go process when a subject uses a proactive control strategy. **c,d,** Within-trial dynamics. **c,** Increased proactivity improves stop trial performance through its effect on the go process. **d,** Increased stopping expectancy improves stop trial performance through its effect on SSRT. **e,** Trial-by-trial dynamics. Illustration of how SSRT, probability of proactivity, and proactive delaying might vary over time within an individual. **f-h,** Key model quantities and their relations. The computational model allows for a detailed, dynamic analysis of inhibitory control processes with trial-level temporal resolution, enabling the investigation of within-subjects variability and nonergodic patterns in cognitive control.

Between-subjects associations between cognitive model parameters and brain activity

Within-subjects associations between cognitive model parameters and brain activity

1274
Figure 3. Divergence of between-subjects and within-subjects brain-behavior associations
in inhibitory control. a,c,e, Between-subjects analysis. Whole-brain correlation maps showing
associations between subject-average brain activation (correct stop versus correct go activation)
and subject-average cognitive model parameters: SSRT **(a)**, probability of proactivity **(c)**, and
proactive delaying **(e)**. Thresholded at Pearson $r \geq 0.05$. **b,d,f,** Within-subjects analysis. Whole-
brain Cohen's d maps showing associations between trial-by-trial brain activity and cognitive
model parameters: SSRT **(b)**, probability of proactivity **(d)**, and proactive delaying **(f)**. SSRT
associations were computed on stop trials; probability of proactivity and proactive delaying
associations were computed on all trials. Thresholded at Cohen's $d \geq 0.1$. For both between- and
within-subjects analyses: SSRT and probability of proactivity $N = 4469$; proactive delaying
$N = 4176$. Striking differences were observed comparing between-subjects and within-subjects
associations across multiple brain regions. This divergence provides evidence for nonergodicity
in inhibitory control processes, challenging the assumption that group-level findings can be
directly applied to understand individual-level cognitive dynamics.

1289
Figure 4. Network-level comparison of between-subjects and within-subjects brain-
behavior associations in inhibitory control. The between-subjects analysis correlated subject-
average brain activation (correct stop versus correct go activation) and subject-average cognitive
model parameters: SSRT **(a)**, probability of proactivity **(b)**, and proactive delaying **(c)**. The
within-subjects analysis regressed trial-by-trial brain activity on trial-by-trial cognitive model
parameters: SSRT **(a)**, probability of proactivity **(b)**, and proactive delaying **(c)**. Effect sizes are
shown for both analyses (between subjects: Pearson r; within subjects: Cohen's d). Statistical
significance is indicated by colored asterisks: red for between subjects ($P_{\text{FDR}} < 0.01$) and blue
for within subjects ($P_{\text{FDR}} < 0.01$). Networks are based on the Shirer parcellation[48]. Differences
in the existence and direction of associations between between-subjects and within-subjects
analyses were observed across multiple brain networks, including the anterior and posterior
salience, left and right frontoparietal, and default mode networks. AntSalience: anterior salience
network; PostSalience: posterior salience network; lFPN: left frontoparietal network; rFPN: right
frontoparietal network; DMN: default mode network; RSC/PHG: retrosplenial cortex /
parahippocampal gyrus network; Precuneus: precuneus network; Visuospatial: visuospatial
network; Language: language network; VisualAssoc: visual association network; Visual: visual
network; Auditory: auditory network; Sensorimotor: sensorimotor network; BasalGanglia: basal
ganglia network.

Figure 5. Stability analysis of within-subjects associations. a-c, Stability plots for SSRT **(a)**, probability of proactivity **(b)**, and proactive delaying **(c)**. In each panel, the top plots show the distributions of correlation between the effect in resamples and the effect in the full sample, while the bottom plots show the distributions of the effect in selected regions. For each plot, 10,000 resamples of n subjects were drawn with replacement for each sample size n, and Cohen's d was calculated for each resample. For top panel plots, correlations were then computed for each resample over the areas belonging to the set of brain areas. The dashed lines depict 95% and 99% bootstrap confidence intervals. Within-subjects associations demonstrated stability for all 3 cognitive model parameters across 5 different collections of brain areas and in 5

1318 regions of interest, even at modest sample sizes. This stability supports the reliability of brain-
1319 behavior associations in inhibitory control processes.
1320

1321
Figure 6. Dissociated brain representations of reactive and proactive control processes. For
each subject, correlations were computed between pairs of within-subjects brain maps for the
model parameters (SSRT, probability of proactivity, and proactive delaying) over each brain
network. **a,** Distributions of correlations over subjects. Kernel density estimation was performed.
b, Average correlations over subjects. The height of the bars is the median. Statistical
significance of difference in medians is indicated by asterisks (* $P_{\text{FDR}} < 0.01$). Reactive (SSRT)
and proactive (probability of proactivity, proactive delaying) control processes showed
dissociated representations across all brain networks. This dissociation suggests that the brain
employs distinct neural resources for reactive and proactive aspects of inhibitory control. In
contrast, the two proactive measures showed high similarity, validating their representation of
related cognitive processes.

1333

Figure 7. Distinct brain-behavior associations for adaptive and maladaptive regulators of
inhibitory control. a, Within-subjects associations of SSRT with brain activity between γ_1
subgroups. We identified two distinct subgroups of subjects with opposite profiles of attentional
modulation. Subjects showed either maladaptive regulation ($\gamma_1 < 0$, $N = 2513$) or adaptive
regulation ($\gamma_1 > 0$, $N = 1956$) of their expectancy of stopping over time. The two groups showed
differences in within-subjects associations between SSRT and brain activity in various networks.
This demonstrates that individual differences in attentional dynamics play a role in shaping the
relationship between neural activity and inhibitory control processes. A colored asterisk indicates
that a network or region's associations were nonzero among subjects with $\gamma_1 < 0$ (blue * $P_{FDR} <$
0.01) or among subjects with $\gamma_1 > 0$ (purple * $P_{FDR} < 0.01$). **b,** Within-subjects associations of
probability of proactivity with brain activity between θ_1 subgroups. We identified two distinct
subgroups of subjects with opposite profiles of performance monitoring. Subjects showed either
adaptive regulation ($\theta_1 < 0$, $N = 3054$) or maladaptive regulation ($\theta_1 > 0$, $N = 1415$) of their
proactivity over time. The two groups showed differences in within-subjects associations
between probability of proactivity and brain activity in various networks. This demonstrates that
the neural correlates of proactive control are influenced by an individual's strategy for adjusting
proactivity in response to task outcomes. A colored asterisk indicates that a network or region's
associations were nonzero among subjects with $\theta_1 < 0$ (blue * $P_{FDR} < 0.01$) or among subjects
with $\theta_1 > 0$ (purple * $P_{FDR} < 0.01$). For both panels, a black asterisk indicates that associations
had different distributions between the two subgroups (* $P_{FDR} < 0.01$, n.s. $P_{FDR} \geq 0.01$).

Figure 8. Nonergodicity across brain networks. a, Degree of nonergodicity of brain networks. Bar plots show the extent of nonergodicity for each brain network across three cognitive model parameters (SSRT, probability of proactivity, proactive delaying). The height of the bars is the mean fraction of subjects showing opposite-sign associations compared to between-subjects results. The bars are shaded by the magnitude of the between-subjects correlations. Error bars show 95% bootstrap confidence intervals. **b,** 3-dimensional embedding of each network. Networks are represented as points based on their nonergodicity measures for the three parameters. **c-d,** Hierarchical clustering of networks based on their nonergodicity profiles. **c,** Dendrogram of clustering. **d,** Euclidean distances between network embeddings. Brain networks exhibited varying degrees of nonergodicity. The anterior salience network showed the highest level of nonergodicity and a unique profile of nonergodicity distinct from that of all other networks.