Title: Modulation of microglia activation after human donor retinal ganglion cell

transplantation

Volha V. Malechka[1], Monichan Hayes Phay[1*], Josy Augustine[1,2], John Dayron Rivera[1],*

Clemens Alt[3], Meredith Gregory-Ksander[1], Petr Baranov[1]

[1]Department of Ophthalmology, Schepens Eye Research Institute of Massachusetts

Eye and Ear, Harvard Medical School, Boston, Massachusetts, USA 02114

[2]Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry

and Biomedical Sciences, Queen's University Belfast, Belfast, United Kingdom

[3]Center for Systems Biology and Wellman Center for Photomedicine, Massachusetts

General Hospital and Harvard Medical School, Boston, Massachusetts, USA

*Contributed equally

Running Head: Modulation of microglia after RGC transplantation

Corresponding Author: Petr Baranov

e-mail: Petr_Baranov@meei.harvard.edu

Conflict of Interests: No conflicting relationship exists for any author

23 **Keywords:** transplantation, microglia, retinal ganglion cells, stem cell therapy

Abstract

Stem cell-derived retinal ganglion cell (RGC) transplantation therapy offers a promising avenue for restoring vision in patients with significant RGC loss. The major challenge in this therapeutic approach is ensuring the survival of transplanted human donor RGCs within the host retina. Previously, we demonstrated the pivotal role of host retinal microglia and macrophages in the acceptance and survival of human donor cells. This current study aimed to improve the survival and integration of human donor RGCs through modulation of the host retinal microglia. Pretreatment of human donor RGCs with annexin V and soluble FasL (sFasL) before transplantation significantly enhanced their acceptance and survival rate. Detailed analyses of the host mouse retina post-transplantation revealed morphological changes and activation patterns in microglia and macrophages. These findings indicate that sFasL and annexin V activate pro-survival pathways and foster a supportive environment for the acceptance and survival of transplanted human donor RGCs.

Introduction

The degeneration of retinal ganglion cells (RGCs) is a key factor in various eye disorders, including different forms of glaucoma, hereditary optic neuropathies, ischemic optic neuropathies, and demyelinating diseases, all of which result in progressive axonal degeneration and death of RGCs, ultimately leading to irreversible blindness[1,2,3,4,5]. Despite significant advances, the clinical needs for these retinal neurodegenerative conditions remain largely unmet. Consequently, a variety of therapeutic strategies are being actively explored, including stem cell-derived therapy, pharmacologic neuroprotection, and gene therapy. Among these, stem cell-derived RGC transplantation therapy holds considerable promise for restoring vision in patients with RGC loss and axonal degeneration[6,7]. However, one of the primary challenges with this therapeutic approach is ensuring the survival of the transplanted human RGCs within the host retina, due to the active surveillance and subsequent elimination by the host microglia[8].

In the healthy and normally developed retina, microglia reside in the outer plexiform layer (OPL) and inner plexiform layer (IPL), where they perform essential functions in immune defense, homeostasis, and development[9]. When exposed to damaging stimuli, microglia become activated and migrate to the sites of retinal injury[10].

We hypothesize that during cell isolation and transplantation, human donor RGCs undergo significant stress, which leads to the externalization of pro-apoptotic signals on their cell surface. These signals attract tissue microglia and macrophages to the site of injury and initiate clearance of the transplanted cells. By blocking these signals and modulating the host retinal microenvironment, we aim to prevent microglia and

macrophages from recognizing and eliminating the transplanted cells, thereby

enhancing the survival of human donor RGCs post-transplantation.

Annexin V emerges as a potential candidate for modulating the retinal

microenvironment and preventing microglia and macrophages from recognizing and

removing human donor RGCs after transplantation. Annexin V is a calcium-dependent,

membrane-bound protein with high affinity for phosphatidylserine (PS), an endogenous

phospholipid and a component of the plasma membrane that becomes exposed on the

cell surface during cellular stress and early apoptosis[11]. Given the distressing nature of

isolation and transplantation, human donor RGCs likely externalize PS under stress.

Therefore, we hypothesized that pretreating donor RGCs with annexin V to mask PS

would prevent recognition and clearance by microglia and macrophages following

transplantation.

Fas ligand (FasL), or CD95 ligand, is another possible candidate for modulating the

retinal microenvironment and improving survival of transplanted human donor RGCs.

FasL is a 40kDa type II transmembrane protein of the tumor necrosis factor (TNF)

family that is capable of inducing apoptosis upon binding to the Fas receptor[12]. Within

the eye, FasL can be expressed as a membrane-bound protein (mFasL), which is pro-

inflammatory and pro-apoptotic molecule, or cleaved and released as a soluble protein

(sFasL), which is non-inflammatory and non-apoptotic[13,14]. Previous studies

demonstrated that FasL deficiency or intravitreal administration of sFasL protects RGCs

from cell death in a mouse model of glaucoma[15]. Furthermore, sFasL has demonstrated

therapeutic benefits in reducing inflammation in experimental models of keratitis[16] and

autoimmune disease[17]. Moreover, there is evidence that post injury treatment with

intravitreal AAV2-soluble Fas ligand restores retinal microglia homeostasis and rescues

RGC function in a mouse model of glaucoma[18]. We hypothesize that Fas-FasL

interaction may protect human donor RGCs from immune attack by host microglia and

macrophages, thereby improving graft survival in the host retina.

Taken together, this study demonstrates that pretreating human donor RGCs with

annexin V and sFasL before transplantation modulates the host microenvironment and

improves graft survival in the host mouse retina. Additionally, our results provide the first

quantitative analysis of microglia and macrophages morphology in the mouse retina

following human donor RGC transplantation.

Materials and methods

Animals

All procedures conformed to the National Research Council's Guide for the Care and

Use of Laboratory Animals, the Association for Research in Vision and Ophthalmology

(ARVO) guidelines and were approved by the Institutional Animal Care and Use

Committee (IACUC) of the Schepens Eye Research Institute (SERI). All animals were

purchased from Jackson Laboratory (Bar Harbor, ME). The breeding of animals was

established in the Rodent Barrier Facility at the SERI. At the beginning of the

experiment equal numbers of male and female *Brn3b$^{-/-}$* and *Cx3cr1$^{-/-}$* mice were group-

housed in a temperature-controlled environment with 12-hour light/dark cycle and

received a standard diet and water ad libitum.

Organoid dissociation and RGC isolation

RGCs were differentiated from H9:BRN3B:tdTomatoThy1.2-hESCs in retinal organoid

3D culture, as was previously described[19,20]. Mature human retinal organoids were

dissociated, and RGCs were isolated using magnetic microbead sorting for Thy1.2+

cells. Following cell isolation, RGCs were formulated in RGC media at $1 - 5 \times 10^5$

cells/mL. Then, human donor RGCs were pretreated with 5 mg/mL annexin V

(BioLegend, San Diego, CA), a calcium-dependent membrane-bound protein with high

affinity to PS, and 0.2 mg/mL sFasL (R&D Systems, Minneapolis, MN), a non-apoptotic,

non-inflammatory protein, for at least 20 minutes on ice. Following human donor RGC

pretreatment, the cells were formulated for transplantation at 2×10^4 cell/µL in RGC

media.

In vivo human donor RGC transplantation

The *Brn3b$^{-/-}$* mice were anesthetized intraperitoneally with 100 mg/kg ketamine

hydrochloride and 20 mg/kg xylazine hydrochloride solution diluted with saline. Pupils

were dilated with 1% tropicamide ophthalmic solution (Bausch + Lomb, Bridgewater,

NJ). Following tropicamide application, one drop of 0.5% proparacaine hydrochloride

ophthalmic solution (Bausch + Lomb, Bridgewater, NJ) was applied onto the cornea for

5 minutes before transplantation. Up to 2×10^4 donor cells per 1 µL were injected into

one eye of *Brn3b$^{-/-}$* mice using a beveled glass microneedle with 100 µm inner diameter

at a 1 µL/min flow rate. Each mouse received a single subretinal injection of 1.0 µL of

human donor RGCs pretreated with annexin V and sFasL during transplantation.

Control group received a single subretinal injection of 1.0 µL of human donor RGCs with

no annexin V or sFasL pretreatment. Following human donor RGC transplantation, one

drop of GenTeal® Tears Lubricant eye gel (Alcon Laboratories, Fort Worth, TX) was

applied onto the cornea to protect the eye from dryness. Mice were returned to the

housing room and maintained on a standard 12-hour light/dark cycle for 3 days. On day

3 mice were euthanized, and eyes were immediately collected and fixed in 4 %

paraformaldehyde (PFA) for 12 hours before immunohistochemistry analysis.

Immunohistochemistry of whole retina mounts

After fixation with 4% PFA was complete, retina samples were incubated in blocking

buffer [10 % goat serum, 1 % bovine serum albumin (BSA), 0.1 % sodium citrate, 0.1 %

Tween 20, and 0.1 % Triton-X in 1X phosphate-buffered saline (PBS)] containing goat

anti-mouse antibodies in dilution 1:400 at 4°C for at least 12 hours overnight to limit

nonspecific staining. Following blocking, retina flat mounts were incubated with Iba-1

primary antibodies (ab178846, Abcam, Boston, MA) in staining buffer (1 % BSA, 0.25 %

Tween 20 and 0.25 % Triton-X in 1X PBS) for 72 hours at 4°C. Then, each retina

sample was washed thrice for 15 minutes with washing buffer (0.1 % Tween-20, 0.1 %

Triton-X in 1X PBS) at room temperature to remove any unbound primary antibodies

before adding secondary antibodies. Secondary antibodies were prepared in dilution

1:500 in staining buffer, and the retina samples were incubated with secondary

antibodies for 24 hours at 4°C. Then, each sample was washed with washing buffer

thrice for 15 minutes at room temperature and then incubated with 1 µg/mL DAPI (4',6-

diamidino-2-phenylindole) in 1X PBS for 20 minutes at room temperature to stain the

cell nuclei. Following final rinse with 1X PBS, retina samples were mounted and stored

in the dark at room temperature before imaging. Images of retina mounts were acquired

using confocal microscope (Olympus, Tokyo, Japan), and the Iba1-positive microglia

and macrophages were imaged within the OPL and IPL.

Quantification and analysis of morphology of retinal microglia and macrophages

after human donor RGC transplantation

The image of each retina flat mount was divided into four quadrants with one mid-

peripheral region being a representative image of each quadrant; therefore, a total of

four images per retina was used for quantitative analysis of retinal microglia and

macrophages. The analysis was applied to quantify retinal microglia and macrophages

and assess morphology of retinal microglia and macrophages according to metrics such

as cell shape, ramification, and complexity. Morphometric analysis of total number of

Iba1-positive retinal microglia and macrophages within the OPL and IPL was performed

on skeletonized images of Iba1-positive cells from the retinal flat mounts using

previously published protocols[21]. Briefly, eight Z-stack images (two images per retinal

quadrant) were captured at 20X magnification from one retina per animal and

processed using ImageJ software (provided in the public domain). The images were

binarized using the "adjust threshold" function and skeletonized using the "skeletonize"

function. The plugin "Analyze Skeleton" was then employed to generate the number of

branches, number of junctions, and average branch length of the microglia and

macrophages in the retina flat mounts. All morphometric parameter values were divided

by the total number of cells per unit area of retinal tissue in mm^2.

Results

Donor cell survival is improved by blocking phosphatidylserine residues on human donor RGCs with annexin V

We observed that pretreatment of human donor RGCs with annexin V prior to transplantation improves human donor RGC survival rate and overall transplant success (**Figure 1**). We also observed in the successful grafts, that the axons of human donor RGCs enter the optic nerve head (**Figure 1A**). Specifically, in the pretreated group, eleven out of twelve recipients of human donor RGCs (92%) had approximately 200 donor cells within the retina at day 4, representing more than 1% of all transplanted human donor RGCs in one eye. In contrast, in the control group, only six out of eleven recipients (55%) showed survival of human donor RGCs after transplantation (**Figure 1B**). Furthermore, in the annexin V pretreated group, four out of twelve transplants exhibited neurite outgrowth into the optic nerve head, compared to only one out of eleven transplants in the control group (**Figure 1C**).

Host microglia are activated after transplantation of human donor RGCs in Cx3cr1-GFP mice

To study microglia activation *in vivo*, we imaged the retina repeatedly by scanning laser ophthalmoscopy (SLO)[22] of *Cx3cr1-GFP* heterozygous mice (**Figure 2**). On day 1 post human donor RGC transplantation, host retinal microglia and macrophages become activated and initiate displacement toward injection site to remove the transplanted RGCs (**Figure 2A, 2B**). By days 2 and 3, microglia and macrophages began eliminating human donor RGCs, occupying the area of damage, and by day 7, all human donor RGCs were removed and processed by these activated cells (**Figure 2A, 2B**).

Retinal microglia and macrophages rapidly respond to physiological changes and

demonstrate distinct morphology based on their state of reactivity. As shown in **Figure**

2C, activated microglia and macrophages retract their processes and round their cell

bodies, whereas in their homeostatic state, they exhibit small cell bodies with ramified

processes.

Human donor RGCs are eliminated by primary microglia in vitro

To determine if donor RGCs are eliminated by microglia *in vitro*, we co-cultured primary

microglia with human donor RGCs (**Figure 3A**). We observed that primary microglia

become activated, move towards human donor RGCs and began eliminating them by

day 1 (**Figure 3B**). This finding is consistent with our *in vivo* observation, demonstrating

that human donor RGCs are recognized and eliminated by microglia in the host

microenvironment. Therefore, modulating this microenvironment might help suppress

microglia activity and improve the survival of human donor RGCs.

Host microglia is modulated by exogenous pretreatment of human donor RGCs

with annexin V and sFasL before transplantation

To investigate whether the host retinal microenvironment, where microglia and

macrophages play an essential role in active surveillance can be modulated, we

pretreated human donor RGCs before transplantation. Following dissociation and

isolation of human donor RGCs, we pretreated the cells with 5 mg/mL annexin V and

0.2 mg/mL sFasL for at least 20 minutes on ice. Following transplantation, our results

indicate improved survival and axonal elongation in human donor RGCs pretreated with

annexin V and sFasL as compared to non-pretreated cells (**Figure 4A, 4B**).

Microglia and macrophages are resident cells in the retina and are localized to the OPL

and IPL in the retina. To explore the effect of pretreating human donor RGCs with

annexin V and sFasL on microglia and macrophages, we analyzed Iba1-positive cells

within OPL and IPL after human donor RGC transplantation. At 3 days post

transplantation of human donor RGCs pretreated with Annexin V and sFasL, microglia

and macrophages displayed homeostatic morphology with prominent and longer

processes as compared to the control group with no pretreatment, where microglia and

macrophages displayed a more reactive phenotype with rounded cell bodies and

retracted processes (**Figures 4C-F**). Interestingly, we also found that microglia and

macrophages become more activated in the OPL as compared to the IPL after

subretinal delivery of human donor RGCs (**Figures 4C-F**). Moreover, the total number

of microglia and macrophages was reduced in the pretreatment group as compared to

the control group with no pretreatment (**Figures 4G, 4J**), again with more significant

changes observed in IPL. Additionally, the number of branches per microglia and

macrophage was significantly (P<0.01) increased in the pretreatment group, indicating

less activation when compared to the control group (no pre-treatment) (**Figures 4H,**

4K).

Moreover, microglia and macrophages in the OPL exhibit a significant difference in

branch number per cell between pretreated and control groups (**Figure 4H**), while those

in the IPL showed lower branch numbers per cell (**Figure 4K**). Similar trend is observed

in the branch length per microglia and macrophage cell with longer and more ramified

branches in OPL that in IPL, suggesting that microglia and macrophages are less

activated in the OPL than IPL after human donor RGC transplantation (**Figures 4I, 4L**).

In summary, pretreating human donor RGCs with annexin V and sFasL leads to

morphologic changes in microglia and macrophages within the OPL and IPL of the

mouse retina.

Discussion

Optic neuropathies and retinal neurodegenerative diseases, such as glaucoma, diabetic

retinopathy, and age-related macular degeneration, result in progressive damage of

retinal ganglion cells (RGCs)[23]. If left untreated, these diseases lead to irreversible

blindness. For the patients with optic neuropathies and neurodegenerative diseases of

the retina, cell replacement therapy[24,25,26] and treatment with neuroprotective agents are

promising strategies to halt disease progression and preserve vision[27,28]. One of the key

steps for improving cell transplantation success in the retina is to define the factors that

positively affect survival of donor RGCs and to modulate the host microenvironment by

enhancing the beneficial effect of these factors on donor cells.

The retina is populated by specialized immune resident macrophages called microglia.

Normally, retinal microglia reside within the outer plexiform layer (OPL), the inner

plexiform layer (IPL), and the retinal ganglion cell (RGC) layer. Microglia represent one

of the main barriers the donor cells need to overcome for successful survival and

integration within the host microenvironment, as microglia are responsible for

elimination of antigens that threaten the retinal integrity[29].

Prior studies demonstrated that phosphatidylserine (PS), a phospholipid that is localized

to the inner part of the cell membrane, is crucial for identification of apoptotic cells by

macrophages[30,31]. Considering donor RGCs experience a substantial stress during

dissociation, isolation, and transplantation events, it is predicted they might generate

pro-apoptotic signals and externalize a PS signal on the cell surface, allowing them to

be recognized and cleared by retinal microglia and macrophages[32]. Annexin V, a

calcium-dependent membrane-binding protein, efficiently and specifically attaches to

PS, thus, representing a selective probe for recognition and quantification of apoptotic

cells[33,34,35]. By pretreating donor RGCs with annexin V just after isolation and right

before transplantation events, a PS signal on the cell surface of donor RGCs can be

blocked, and hence preventing donor RGCs from being recognized and attacked by

retinal microglia and macrophages.

In addition to blocking the PS signal with annexin V, we previously demonstrated that

treatment with sFasL is also neuroprotective and prevents death of RGCs in chronic

and inducible mouse models of glaucoma[15,36]. Herein, we determined the combination

of annexin V and sFasL significantly improves survival of human donor RGCs in the

host mouse retina. FasL is a type II transmembrane protein that exists as both a

membrane bound protein (mFasL) that is pro-inflammatory and pro-apoptotic, and a

soluble form (sFasL) that is cleaved from the cell surface and released as a soluble

fragment that is non-inflammatory and non-apoptotic and can antagonize the activity of

mFasL[14,15,37]. In the healthy eye, the anti-apoptotic and anti-inflammatory sFasL is the

dominant isoform expressed[36], but with disease there is a shift and the pro-inflammatory

and pro-apoptotic mFasL is the dominant isoform expressed in a variety of ocular

pathologies[36,38,39,40].

In inducible and chronic models of glaucoma it has been shown that mFasL mediates

destruction of RGCs either by directly killing RGCs and/or by inducing the production of

pro-inflammatory chemokines or other mediators in Fas+ glial cells and thereby causing

neurotoxic inflammation[15,36,41]. By contrast, intravitreal AAV2-mediated gene delivery of

sFasL provided complete and long-term protection of RGCs and their axons[15,36].

Importantly, the sFasL-mediated neuroprotection of RGCs also correlates with reduced

microglia activation and reduced production of pro-inflammatory mediators. Therefore,

sFasL could be improving survival of donor RGCs by blocking FasL-mediated apoptosis

of RGCs and/or by blocking FasL-mediated activation of glia and the induction of

neurotoxic inflammation.

Normally, in the healthy eye, retinal microglia display a highly ramified morphology with

a small cell body and multiple branching processes. When responding to inflammation

or injury, microglia retract their processes and present round cellular morphology.

Although microglia are the most abundant immune cells in the retina, the retina also

contains a population of perivascular macrophages that are located within the glia

limitans of retinal vasculature[42]. The glia limitans acts as a physical barrier against

unwanted cells that attempt to enter the retina. During the subretinal delivery of donor

RGCs, the microinjury to small retinal blood vessels may occur causing the leakage of

perivascular macrophages to the site of injection. Both microglia and macrophages

express the ionized calcium-binding adaptor (Iba1) molecule, a 17-kDa actin-binding

cross-linking protein that is upregulated during activation of these cells[43].

Here, we investigated the morphology of Iba1+ microglia and macrophages following

pretreatment of human donor RGCs with annexin V and sFasL before transplantation.

We found that transplantation of human donor RGCs leads to activation of host retinal

microglia and macrophages causing them to move towards the injection site to eliminate

human donor RGCs. Pretreatment with annexin V and sFasL resulted in reduced

activation of microglia and macrophages, with Iba1+ microglia and macrophages

displaying a more ramified morphology with longer processes (**Figures 4E, 4F**). In

contrast, the Iba1+ microglia and macrophages in the control group exhibited a round

ameboid morphology with retracted processes (**Figures 4C, 4D**).

Our results demonstrate that modulating the host retinal microenvironment through the

microglia manipulation can improve the survival of human donor RGCs after

transplantation.

Statistical analysis

All graphs were generated by GraphPad PRISM 9 (GraphPad software, San Diego,

CA). Statistical significance was determined using an unpaired Welch's *t*-test. All data

are presented as average ± standard deviation (SD). Where normally distributed, the

data were analyzed with a *P* value of <0.05 (*p < 0.05, **p < 0.01, ***p < 0.001, ****p <

0.0001) as a statistically significant.

Acknowledgments

This work was supported by NEI/NIH U24 (PB), GFF (PB), and Iraty Award (PB). The

authors would like to thank Dr. Don Zack for providing the Brn3b-tdTomato hESCs.

Author contributions

MHP, MGK and PB initiated the project and designed experiments. VVM, MHP, JA

performed confocal microscopy, immunohistochemistry experiments. JDR differentiated

human donor RGCs. JA analyzed the images of microglia cells. CA performed scanning

laser ophthalmoscopy. VVM and PB performed cell transplantation, wrote the

manuscript draft. PB supervised the project. All authors reviewed and contributed to

editing the manuscript.

Declaration of interests

353 The authors declare no competing interests.

References

1. Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. Br J Ophthalmol. 2006 Mar;90(3):262-7. doi: 10.1136/bjo.2005.081224.
2. Cook C, Foster P. Epidemiology of glaucoma: what's new? Can J Ophthalmol. 2012 Jun;47(3):223-6. doi: 10.1016/j.jcjo.2012.02.003.
3. Newman NJ, Biousse V. Hereditary optic neuropathies. Eye (Lond). 2004 Nov;18(11):1144-60. doi: 10.1038/sj.eye.6701591.
4. Patil AD, Biousse V, Newman NJ. Ischemic Optic Neuropathies: Current Concepts. Ann Indian Acad Neurol. 2022 Oct;25(Suppl 2):S54-S58. doi: 10.4103/aian.aian_533_22.
5. Klistorner A, Sriram P, Vootakuru N, Wang C, Barnett MH, Garrick R, Parratt J, Levin N, Raz N, Van der Walt A, Masters L, Graham SL, Yiannikas C. Axonal loss of retinal neurons in multiple sclerosis associated with optic radiation lesions. Neurology. 2014 Jun 17;82(24):2165-72. doi: 10.1212/WNL.0000000000000522.
6. Singh MS, Park SS, Albini TA, Canto-Soler MV, Klassen H, MacLaren RE, Takahashi M, Nagiel A, Schwartz SD, Bharti K. Retinal stem cell transplantation: Balancing safety and potential. Prog Retin Eye Res. 2020 Mar;75:100779. doi: 10.1016/j.preteyeres.2019.100779.
7. Zhang J, Wu S, Jin ZB, Wang N. Stem Cell-Based Regeneration and Restoration for Retinal Ganglion Cell: Recent Advancements and Current Challenges. Biomolecules. 2021 Jul 5;11(7):987. doi: 10.3390/biom11070987.
8. Rashid K, Akhtar-Schaefer I, Langmann T. Microglia in Retinal Degeneration. Front Immunol. 2019 Aug 20;10:1975. doi: 10.3389/fimmu.2019.01975.
9. Karlstetter M, Scholz R, Rutar M, Wong WT, Provis JM, Langmann T. Retinal microglia: just bystander or target for therapy? Prog Retin Eye Res. 2015 Mar;45:30-57. doi: 10.1016/j.preteyeres.2014.11.004.
10. Chen M, Xu H. Parainflammation, chronic inflammation, and age-related macular degeneration. J Leukoc Biol. 2015 Nov;98(5):713-25. doi: 10.1189/jlb.3RI0615-239R.
11. Vermes I, Haanen C, Steffens-Nakken H, Reutelingsperger C. A novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. J Immunol Methods. 1995 Jul 17;184(1):39-51. doi: 10.1016/0022-1759(95)00072-i.
12. Kägi D, Vignaux F, Ledermann B, Bürki K, Depraetere V, Nagata S, Hengartner H, Golstein P. Fas and perforin pathways as major mechanisms of T cell-mediated cytotoxicity. Science. 1994 Jul 22;265(5171):528-30. doi: 10.1126/science.7518614.
13. Suda T, Hashimoto H, Tanaka M, Ochi T, Nagata S. Membrane Fas ligand kills human peripheral blood T lymphocytes, and soluble Fas ligand blocks the killing. J Exp Med. 1997 Dec 15;186(12):2045-50. doi: 10.1084/jem.186.12.2045.
14. Hohlbaum, A. M., Moe, S. & Marshak-Rothstein, A. Opposing effects of transmembrane and soluble Fas ligand expression on inflammation and tumor cell survival. The Journal of experimental medicine 191, 1209-1220 (2000).
15. Gregory, M. S. *et al.* Opposing roles for membrane bound and soluble Fas ligand in glaucoma-associated retinal ganglion cell death. *PloS one* **6**, e17659, doi:10.1371/journal.pone.0017659 (2011).
16. Rogge M, Yin XT, Godfrey L, Lakireddy P, Potter CA, Del Rosso CR, Stuart PM. Therapeutic Use of Soluble Fas Ligand Ameliorates Acute and Recurrent Herpetic

Stromal Keratitis in Mice. Invest Ophthalmol Vis Sci. 2015 Oct;56(11):6377-86. doi: 10.1167/iovs.15-16588.

17. Li NL, Nie H, Yu QW, Zhang JY, Ma AL, Shen BH, Wang L, Bai J, Chen XH, Zhou T, Zhang DQ. Role of soluble Fas ligand in autoimmune diseases. World J Gastroenterol. 2004 Nov 1;10(21):3151-6. doi: 10.3748/wjg.v10.i21.3151.

18. Gregory-Ksander MS, Krishnan A, Shrestha M, Nguyen K, Marshak-Rotstein A. Post injury treatment with intravitreal AAV2-soluble Fas ligand restores retinal microglia homeostasis and rescues retinal ganglion cell function in a mouse model of glaucoma. ARVO Annual Meeting Abstract, June 2023

19. Fligor CM, Huang KC, Lavekar SS, VanderWall KB, Meyer JS. Differentiation of retinal organoids from human pluripotent stem cells. Methods Cell Biol. 2020;159:279-302. doi: 10.1016/bs.mcb.2020.02.005.

20. Oswald J, Kegeles E, Minelli T, Volchkov P, Baranov P. Transplantation of miPSC/mESC-derived retinal ganglion cells into healthy and glaucomatous retinas. Mol Ther Methods Clin Dev. 2021 Mar 10;21:180-198. doi: 10.1016/j.omtm.2021.03.004.

21. Young K, Morrison H. Quantifying Microglia Morphology from Photomicrographs of Immunohistochemistry Prepared Tissue Using ImageJ. J Vis Exp. 2018 Jun 5;(136):57648. doi: 10.3791/57648.

22. Clemens Alt, Charles P. Lin. In vivo quantification of microglia dynamics with a scanning laser ophthalmoscope in a mouse model of focal laser injury. Proc. of SPIE. 2012. Vol. 8209, 820907doi: 10.1117/12.909141

23. Carelli V, La Morgia C, Ross-Cisneros FN, Sadun AA. Optic neuropathies: the tip of the neurodegeneration iceberg. Hum Mol Genet. 2017 Oct 1;26(R2):R139-R150. doi: 10.1093/hmg/ddx273.

24. Johnson TV, Bull ND, Hunt DP, Marina N, Tomarev SI, Martin KR. Neuroprotective effects of intravitreal mesenchymal stem cell transplantation in experimental glaucoma. Invest Ophthalmol Vis Sci. 2010 Apr;51(4):2051-9. doi: 10.1167/iovs.09-4509.

25. Chamling X, Sluch VM, Zack DJ. The Potential of Human Stem Cells for the Study and Treatment of Glaucoma. Invest Ophthalmol Vis Sci. 2016 Apr 1;57(5):ORSFi1-6. doi: 10.1167/iovs.15-18590.

26. da Silva-Junior AJ, Mesentier-Louro LA, Nascimento-Dos-Santos G, Teixeira-Pinheiro LC, Vasques JF, Chimeli-Ormonde L, Bodart-Santos V, de Carvalho LRP, Santiago MF, Mendez-Otero R. Human mesenchymal stem cell therapy promotes retinal ganglion cell survival and target reconnection after optic nerve crush in adult rats. Stem Cell Res Ther. 2021 Jan 19;12(1):69. doi: 10.1186/s13287-020-02130-7.

27. Chang EE, Goldberg JL. Glaucoma 2.0: neuroprotection, neuroregeneration, neuroenhancement. Ophthalmology. 2012 May;119(5):979-86. doi: 10.1016/j.ophtha.2011.11.003.

28. Wareham LK, Risner ML, Calkins DJ. Protect, Repair, and Regenerate: Towards Restoring Vision in Glaucoma. Curr Ophthalmol Rep. 2020;8(4):301-310. doi: 10.1007/s40135-020-00259-5.

29. Murenu E, Gerhardt MJ, Biel M, Michalakis S. More than meets the eye: The role of microglia in healthy and diseased retina. Front Immunol. 2022 Nov 29;13:1006897. doi: 10.3389/fimmu.2022.1006897.

30. Fadok VA, Savill JS, Haslett C, Bratton DL, Doherty DE, Campbell PA, Henson PM. Different populations of macrophages use either the vitronectin receptor or the phosphatidylserine receptor to recognize and remove apoptotic cells. J Immunol. 1992 Dec 15;149(12):4029-35.

31. Martin SJ, Reutelingsperger CP, McGahon AJ, Rader JA, van Schie RC, LaFace DM, Green DR. Early redistribution of plasma membrane phosphatidylserine is a general

feature of apoptosis regardless of the initiating stimulus: inhibition by overexpression of Bcl-2 and Abl. J Exp Med. 1995 Nov 1;182(5):1545-56. doi: 10.1084/jem.182.5.1545.

32. Logue SE, Elgendy M, Martin SJ. Expression, purification and use of recombinant annexin V for the detection of apoptotic cells. Nat Protoc. 2009;4(9):1383-95. doi: 10.1038/nprot.2009.143.

33. Gerke V. Annexins And Membrane Organisation In The Endocytic Pathway. Cell Mol Biol Lett. 2001;6(2):204.

34. Gerke V, Moss SE. Annexins: from structure to function. Physiol Rev. 2002 Apr;82(2):331-71. doi: 10.1152/physrev.00030.2001.

35. Koopman G, Reutelingsperger CP, Kuijten GA, Keehnen RM, Pals ST, van Oers MH. Annexin V for flow cytometric detection of phosphatidylserine expression on B cells undergoing apoptosis. Blood. 1994 Sep 1;84(5):1415-20.

36. Krishnan, A. *et al.* Overexpression of Soluble Fas Ligand following Adeno-Associated Virus Gene Therapy Prevents Retinal Ganglion Cell Death in Chronic and Acute Murine Models of Glaucoma. J Immunol 197, 4626-4638, doi:10.4049/jimmunol.1601488 (2016)

37. Tanaka, M., Itai, T., Adachi, M. & Nagata, S. Downregulation of Fas ligand by shedding. Nature medicine 4, 31-36 (1998).

38. Gregory, M. S. *et al.* Membrane Fas ligand activates innate immunity and terminates ocular immune privilege. J Immunol 169, 2727-2735 (2002).

39. Matsumoto, H. *et al.* Membrane-bound and soluble Fas ligands have opposite functions in photoreceptor cell death following separation from the retinal pigment epithelium. Cell Death Dis 6, e1986, doi:10.1038/cddis.2015.334 (2015).

40. Gregory-Ksander, M., Perez, V. L., Marshak-Rothstein, A. & Ksander, B. R. Soluble Fas ligand blocks destructive corneal inflammation in mouse models of corneal epithelial debridement and LPS induced keratitis. Exp Eye Res 179, 47-54, doi:10.1016/j.exer.2018.10.013 (2019).

41. Krishnan, A., Kocab, A. J., Zacks, D. N., Marshak-Rothstein, A. & Gregory-Ksander, M. A small peptide antagonist of the Fas receptor inhibits neuroinflammation and prevents axon degeneration and retinal ganglion cell death in an inducible mouse model of glaucoma. Journal of neuroinflammation 16, 184, doi:10.1186/s12974-019-1576-3 (2019).

42. Mendes-Jorge L, Ramos D, Luppo M, Llombart C, Alexandre-Pires G, Nacher V, Melgarejo V, Correia M, Navarro M, Carretero A, Tafuro S, Rodriguez-Baeza A, Esperança-Pina JA, Bosch F, Ruberte J. Scavenger function of resident autofluorescent perivascular macrophages and their contribution to the maintenance of the blood-retinal barrier. Invest Ophthalmol Vis Sci. 2009 Dec;50(12):5997-6005. doi: 10.1167/iovs.09-3515.

43. Sasaki Y, Ohsawa K, Kanazawa H, Kohsaka S, Imai Y. Iba1 is an actin-cross-linking protein in macrophages/microglia. Biochem Biophys Res Commun. 2001 Aug 17;286(2):292-7. doi: 10.1006/bbrc.2001.5388.

Figure legends:

Figure 1. Donor cell survival is improved by blocking phosphatidylcholine (PS) residues on human donor retinal ganglion cells (hRGCs) with annexin V. (**A**) Donor hRGCs survive and send their projections to the optic nerve head (ONH). (**B**) Donor hRGCs survive better after pretreatment with annexin V. (**C**) Donor hRGCs enter ONH more successfully after pretreatment with annexin V.

Figure 2. Studying microglia activation *in vivo* in the retina of *Cx3cr1-GFP* mice after donor human retinal ganglion cells (hRGCs) transplantation. (**A**) Imaging *in vivo* by scanning laser ophthalmoscopy (SLO). Image on day 1 represents injected donor tdTomato-positive hRGCs (red) and host microglia cells (green) after transplantation. Note that by day 7 all injected donor tdTomato hRGCs are eliminated by activated host microglia cells. (**B**) Interaction between host microglia and donor hRGCs on a single cell level. (**C**) Morphological characteristics of a microglia cell.

Figure 3. Elimination of human retinal ganglion cells (hRGCs) by primary microglia *in vitro* on day 0 (**A**) and day 1 (**B**).

Figure 4. Pretreatment of donor hRGCs with annexin V and sFasL inhibits microglia activation and increases survival of hRGCs after transplantation. (**A, B**) Donor hRGCs survive better after pretreatment with annexin V and sFasL. Activation of Iba1-positive microglia and macrophages in the OPL (**C, E**) and IPL (**D, F**) of the retina in the control group and group pretreated with annexin V and sFasL before donor RGC

531   transplantation. Microglia density (**G, J**), number of branches per microglia and

532   macrophage cell (**H, K**), branch length per microglia and macrophage cell (**I, L**) in the

533   OPL (**G, H, I**) and IPL (**J, K, L**) of the retina in the control group and group pretreated

534   with annexin V and sFasL. hRGCs, human retinal ganglion cells; IPL, inner plexiform

535   layer; OPL, outer plexiform layer; sFasL, soluble Fas ligand.

A

B

C

Figure 1. Malechka VV et al. 2024

A

B

C

Resting ←————————————————————————→ Activated

Figure 2. Malechka VV et al. 2024

Figure 3. Malechka VV et al. 2024

Figure 4. Malechka VV et al. 2024