Cnidaria XIAP activates caspase-mediated cell death

Yuan Chen[1,2,3], Meng Wu[1,2,3], Zihao Yuan[1,2,3], Qingyue Wang[1,2,3], Hang Xu[1,2]*, Li Sun[1,2,3]*

[1]CAS and Shandong Province Key Laboratory of Experimental Marine Biology, Institute of

Oceanology, Center for Ocean Mega-Science, Chinese Academy of Sciences, Qingdao, China

[2]Laboratory for Marine Biology and Biotechnology, Qingdao Marine Science and Technology Center,

Qingdao, China.

[3]College of Marine Sciences, University of Chinese Academy of Sciences, Qingdao, China

*Correspondence: xuhang@qdio.ac.cn; lsun@qdio.ac.cn

Abstract

In vertebrate, X-linked inhibitor of apoptosis (XIAP) is a potent inhibitor of apoptosis. XIAP inhibits apoptosis by interacting with proapoptotic caspases via the baculovirus IAP repeat (BIR) domains and mediating caspase ubiquitination via the really interesting new gene (RING) domain and ubiquitin-associated (UBA) domain. In invertebrate, mostly arthropods, XIAP is also known as an apoptosis inhibitor. To date, no study on basal metazoan XIAP has been documented. In the present work, we examined the biological activity of XIAP from jellyfish *Aurelia coerulea* (AcXIAP) and other non-bilaterians. AcXIAP possesses three BIRs and one RING domain but lacks the UBA domain. AcXIAP augmented the apoptosis-inducing activity of all of the four *A. coerulea* caspases, of both the initiator and the effector clades, identified in this study. AcXIAP activated caspase via one of the BIRs, which bound and stabilized the caspase, and the RING domain, which mediated ubiquitination on the p20 subunit of the caspase in a lysine-independent manner. Similar caspase-activating properties were also observed in the XIAP of hydra, coral, and sponge. In hydra, XIAP knockdown markedly decreased cell death induced by an apoptosis inducer. Together these results revealed the unconventional function and working mechanism of XIAP in Cnidaria, and shed new light into the functional and structural evolution of XIAP.

Keywords: XIAP; Cnidaria; jellyfish; apoptosis; caspase

Highlights

- Jellyfish XIAP (AcXIAP) promotes caspase-mediated apoptosis.

- AcXIAP stabilizes and promotes the activation of caspase via the BIR3 domain.

- AcXIAP mediates lysine-independent ubiquitination on caspase via the RING domain.

- Pro-apoptotic XIAP is common in non-bilaterians and involved in apoptosis in hydra.

1. Introduction

Apoptosis, a type of programmed cell death (PCD), is a fundamental process required for the maintenance of homeostasis under both physiological and pathological conditions. The operation of apoptosis relies on a class of cysteine aspartyl proteases called caspases (Fan, Han, Cong, & Liang, 2005). In humans and mice, a dozen caspases have been identified. These caspases are divided into two groups, i.e., apoptotic caspases, which are further divided into apoptosis initiator caspases (-2, -8, -9, 10) and effector caspases (-3, -6, -7), and inflammatory caspases (-1, -4, -5, -11, 12), which are involved in inflammatory responses (Ross et al., 2022). Caspases are synthesized as inactive zymogens consisting of a pro-domain and a protease domain (CASc), the latter comprising a large subunit (p20) and a small subunit (p10). The pro-domains of effector caspases generally exhibit no structure, while the pro-domains of inflammatory and initiator caspases typically contain either the caspase recruitment domain (CARD) domain or the death effector domain (DED) (Fan et al., 2005). During the process of apoptosis, active effector caspases, induced by the mitochondrial or death receptor-mediated cascade signaling, cleave various substrates, including DNA fragmentation factor 45, gelsolin, and poly (ADP-ribose) polymerase, cumulatively leading to

apoptosis (Cohen, 1997; Reed, 2000). Unlike other common regulatory enzymes, such as

phosphorylase, methylase, and acetylase (Gujral, Mahajan, Lissaman, & Ponnampalam, 2020;

Hao, Xu, Jiang, & Sun, 2024; Zhang, Jing, Li, He, & Guo, 2019), caspases hydrolyze their

substrates directly and hence exert an irreversible effect on the substrates (Cohen, 1997). As a

result, the activities of caspases are tightly controlled, either directly through binding to

regulatory proteins, or indirectly through transcription or translation (Hanifeh & Ataei, 2022).

The inhibitor of apoptosis protein (IAP) represents a family of negative regulators of

caspase and apoptosis (Budhidarmo & Day, 2015). All IAPs contain at least one copy of the

baculovirus IAP repeat (BIR) domain in their N-terminus. Most IAPs contain three BIR

domains, which confer on IAPs the ability to bind caspases and other substrates (Hanifeh &

Ataei, 2022). X-linked inhibitor of apoptosis (XIAP), the most potent and multifaceted

member of IAPs, also contains a Ub-associated domain (UBA) for interacting with poly-Ub

chains, and a RING (Really Interesting New Gene) domain pivotal for E3 ligase activity

(Hanifeh & Ataei, 2022). In humans, XIAP inhibits both the effector caspase-3/7 and the

initiator caspase-9 (Deveraux et al., 1999). The inhibitory effect of XIAP on caspase-3/7

depends on the binding of the second BIR (BIR2) to the caspase, while the inhibitory effect of

XIAP on caspase-9 is mediated by the third BIR (BIR3). The RING domain of XIAP mediates

the transfer of the ubiquitin moiety from the enzyme E2 to the substrate caspase, whereby

targeting the caspase for proteasomal degradation (Budhidarmo & Day, 2015; Hanifeh &

Ataei, 2022).

Caspases are conserved from invertebrate to vertebrate, and caspase genes have been

identified in the genomes of sponge, cnidarians, and all bilaterian clades (Kumar, 2007; Lasi,

David, & Böttger, 2010; Wiens, Krasko, Perovic, & Müller, 2003). The functions of caspases

have been studied in mammals and the major invertebrate model organisms, the nematode

Caenorhabditis elegans and the arthropod *Drosophila melanogaster*(Kumar, 2007). In

addition, two recent reports showed that coral and abalone caspase-3 activated pyroptosis, a

type of PCD different from that of apoptosis (Jiang, Zhou, Sun, Zhang, & Sun, 2020; Qin,

Jiang, Xu, Yuan, & Sun, 2023). In contrast, functional studies of IAPs in invertebrates are

documented mainly with arthropods (Bergmann, Yang, & Srivastava, 2003; Clavier,

Rincheval-Arnold, Colin, Mignotte, & Guénal, 2016; Hay, 2000; Tenev, Ditzel, Zachariou, &

Meier, 2007). In insects, as in mammals, IAPs exert an inhibitory effect on caspase-mediated

apoptosis (Hamajima et al., 2016; Huang et al., 2001; Liao, Yang, & Wu, 2002). In addition to

insects, two reports showed that IAPs from other members of Arthropod, i.e., tiger shrimp

Penaeus monodonthe and Chinese mitten crab *Eriocheir sinensis*, displayed anti-apoptotic

activity (Leu, Kuo, Kou, & Lo, 2008; Qu et al., 2019). These evidences indicate a highly

conserved role of IAPs as a negative regulator of apoptosis in both vertebrate and

invertebrate.

Cnidaria (corals, sea anemones, jellyfish, and hydroids) is an ancient phylum and a sister

group to Bilateria, the two having separated from their common ancestor over 500 million

years ago (Masuda-Ozawa et al., 2022). Genome sequence analyses revealed complex caspase

family members in cnidarians (Dunn, Phillips, Spatafora, Green, & Weis, 2006; Lasi, Pauly, et

al., 2010; Moya et al., 2016). However, the functions of the caspases and their regulatory

IAPs in Cnidaria remain to be explored. Particularly, in jellyfish, no research on caspase or

IAP has been documented. In the present study, we identified four caspases and one XIAP

ortholog (AcXIAP) from the moon jellyfish *Aurelia coerulea*. We found that the four jellyfish

caspases represented different phylogenetic clades but all exhibited apoptotic activities, which

were significantly enhanced, rather than inhibited, by AcXIAP. Similar apoptosis-promoting

effects were subsequently discovered in the XIAP homologues of other Cnidaria species and

Porifera. We further investigated the working mechanism of AcXIAP, and examined the *in*

vivo effect of XIAP knockdown on cell death. Our results uncovered the existence and the

functioning mechanism of pro-apoptotic XIAP in basal metazoans.

2. Materials and Methods

Animals and cell lines

Jellyfish *Aurelia coerulea* medusas were maintained in the laboratory as described previously

(Wang et al., 2020). The entire body of the animal was used for RNA isolation with a TRIZOL

reagent (Invitrogen, USA). Hydra polyps were collected from Beijing, China, and cultured at

18°C in Hydra medium composed of 1 mM NaCl, 1 mM $CaCl_2$, 0.1 mM KCl, 0.1 mM

$MgSO_4$, and 1 mM tris-HCl (pH 7.6). The animals were fed with fresh *Artemia nauplii* larvae

as previously reported (Lasi, Pauly, et al., 2010). HEK293T cells (American Type Culture

Collection) were cultured in Dulbecco's modified Eagle's medium supplemented with 10%

fetal bovine serum (Gibco, Renfrewshire, UK) at 37°C in a 5% CO_2 incubator.

Sequence and phylogenetic analysis

For sequence alignment, XIAP sequences (Table S1) were selected from the National Center

133   for Biotechnology Information (NCBI) (www.ncbi.nlm.nih.gov) and aligned using DNAMAN.

134   Phylogenetic analysis was performed as reported previously using IQ-TREE 2 v.2.1.2 with

135   JTT+R3 substitution models and 1000 bootstraps (Chen, Yuan, & Sun, 2023). Conserved

136   domain analysis was performed using the conserved domain database of NCBI

137   (www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi).

139   Gene cloning and mutagenesis

A. coerulea caspases and XIAP genes were identified by PCR with primers (Table S2) based

on the caspase and XIAP sequences of *Aurelia aurita*. The PCR products were cloned into

pEASY-T1 Simple using the T1 Simple-cloning Kit (TransGen, Beijing, China) and verified

by sequencing. The codon-optimized protein coding sequences (CDSs) of sponge, hydra,

coral and human caspases/XIAPs (Table S1) were synthesized by Sangon Biotech (Shanghai,

China). The CDSs of AcCaspA and AcXIAP truncates were obtained based on their respective

full-length CDSs. Site-directed mutagenesis of lysine to arginine and cysteine to alanine in

AcCaspA was performed using the Fast Mutagenesis System (TransGen, Beijing, China)

according to the manufacturer's instruction. For transfection, the CDSs were subcloned into

pmCherry-N1, pEGFP-N1, pCAGGS-Flag, or pCS2-6 × Myc expression vector (Clontech,

Mountain View, CA, USA). All plasmids were verified by sequencing.

152   Flow cytometry

HEK293T cells were seeded in 12-well plates (Costar, Corning, NY, USA) and transfected

with indicated plasmids. At 24 h post transfection, the cells were collected and washed with

ice-cold PBS. The cells were then doubly stained with Annexin V–fluorescein isothiocyanate

(FITC) and propidium iodide (PI) and analyzed with a FACSAria II flow cytometer (Becton

Dickinson Biosciences, NJ, USA).

Preparation of recombinant protein

The CDS of AcCaspA was subcloned into pET-28a (+) expression vector (Novagen, Madison,

WI, USA) and expressed as His-tagged protein. The recombinant plasmid was introduced into

Escherichia coli Transetta (DE3) (TransGen, Beijing, China) via transformation. The

recombinant protein was prepared from the *E. coli* transformant as described previously

(Chen, Yu, Jiang, & Sun, 2022) with Ni-NTA agarose (QIAGEN, Dusseldorf, Germany). The

protein purity was determined by SDS-PAGE (Fig. S3).

AcCaspA activity

To determine the proteolytic activity of AcCaspA, AcCaspA was incubated with various

colorimetric substrates as described previously (Xu, Yuan, Qin, Jiang, & Sun, 2024). After 1.5

h incubation, the release of ρ-nitroanilide (ρNA) was monitored at OD_{405} nm. To determine

the AcCaspA activity in AcCaspA-expressing HEK293T cells, the cell lysates were buffered

in a 100 μl reaction system [50 mM Hepes (pH 7.5), 3 mM EDTA, 150 mM NaCl, 0.005%

(v/v) Tween 20, and 10 mM dithiothreitol (DTT)] and incubated with Ac-VDVAD-AFC

(MedChem Express, NJ, USA) at a final concentration of 200 μM at 25°C for 1.5 h.

Fluorescence was then measured with a BioTek Synergy HT plate reader (BioTek Instruments,

VT, USA).

XIAP–caspase interaction

HEK293T cells were seeded in 24-well plates (Costar, Corning, NY, USA) overnight. The

cells were transfected with the plasmid expressing caspases, XIAP, or caspase plus XIAP

using PolyJet transfection reagent (SignaGen Laboratories, Ijamsville, MD, USA). After

various hours of transfection, the cells were observed with an inverted microscope (TiS/L100,

Nikon, Tokyo, Japan). For co-localization analysis, HEK293T cells were co-transfected with

the vectors expressing EGFP-tagged AcCaspA and mCherry-tagged AcXIAP for 24 h, and the

cells were observed with a Carl Zeiss LSM 710 confocal microscope (Carl Zeiss, Jena,

Germany).

HvXIAP knockdown and its effect on cell death

HvXIAP-specific small interfering RNA (siRNA) (5'-GGACCAUAAUCCAAUUCAATT-3')

and scrambled RNA (5'-UUCUCCGAACGUGUCACGUTT-3') (negative control) were

designed and synthesized by Sangon Biotech (Shanghai, China). Hydra polys were randomly

divided into different groups (10 animals/group). The groups were administered with or

without HvXIAP-specific siRNA or scrambled RNA with an electroporator (BTX Gemini X2,

USA) based on the reported method of electroporation (Bosch, Augustin, Gellner, Khalturin,

& Lohmann, 2002). After electroporation, the polyps were transferred into Hydra medium

supplemented with 20% (v/v) hyperosmotic dissociation medium consisting of 3.6 mM KCl,

6 mM $CaCl_2$, 1.2 mM $MgSO_4$, 6 mM sodium pyruvate, 6 mM sodium citrate, 12.5 mM N-tris

[hydroxymethyl] methyl-2aminoethanesulfonic acid (TES), 6 mM glucose, and 50 mg/mL

rifampicin. After 24 h incubation, the medium was replaced by fresh Hydra medium. To

examine the efficiency of electroporation, hydra polyps were electroporated as above with or

without 6-carboxy-fluorescein-labeled scrambled RNA and observed with a fluorescence

microscope (Olympus, Tokyo, Japan) at 24 h after electroporation. For HvXIAP knockdown,

hydra polyps were electroporated as above with HvXIAP-specific siRNA or scrambled RNA.

The knockdown efficiency was determined by Quantitative Real-Time Reverse

Transcription-PCR (qRT-PCR) as reported previously (Xu, Jiang, Yu, Yuan, & Sun, 2022)

with primers listed in Table S2. To examine the effect of HvXIAP knockdown on cell death

induced by colchicine, hydra polyps were administered with HvXIAP-specific siRNA or

scrambled RNA as above and then treated with colchicine (2 mg/ml) for 2 h, followed by

washing in Hydra medium. The polyps were stained with PI (2 μg/ml) for 15 min and then

observed with a fluorescence microscope (Olympus, Tokyo, Japan).

Immunoblotting

HEK293T cells transfected with indicated plasmids were lysed with lysis buffer (Beyotime,

Shanghai, China), and the lysate was boiled in SDS loading buffer and electrophoresed in 12%

SDS-PAGE (GenScript, Piscataway, NJ). The proteins were transferred to a NC membrane

and treated with the indicated antibodies. β-actin served as a control. The protein bands were

visualized with an ECL kit (Sparkjade Biotechnology Co. Ltd, Shandong, China). The

antibodies used for immunoblotting are as follows: HRP-conjugated mouse anti DDDDK-Tag

mAb, HRP-conjugated mouse anti Myc-Tag mAb, HRP-conjugated mouse anti HA-Tag mAb,

and HRP-conjugated β-actin mouse mAb. All antibodies were from ABclonal Technology,

Wuhan, China.

Co-immunoprecipitation (Co-IP)

For Co-IP, HEK293T cells were transfected with the indicated plasmids for 24 h and lysed

with lysis buffer. The cells lysates were incubated with anti-Flag or anti-Myc affinity gels

(Beyotime, Shanghai, China) 4°C overnight. The affinity gels were then washed with PBS for

10 times and boiled with loading buffer to elute the bound proteins. The eluted proteins were

subjected to immunoblotting as described above.

Statistical analysis

The GraphPad Prism software 6 was used for statistical analysis, with statistical significance

defined as $p < 0.05$. The unpaired Student's t-test was used for the comparison between two

groups. Error bars represent the mean ± SD.

3. Results

3.1 Identification of four *A. coerulea* caspases with apoptotic activity

Four caspases were identified from jellyfish *A. coerulea* (Fig. 1A). Two of these caspases

exhibit no apparent structure in the pro-domain and were named AcCaspA and AcCaspB,

respectively. The other two caspases possess a CARD domain besides the CASc domain and

were designated AcCCaspA (the first capital C indicates "CARD') and AcCCaspB,

respectively. Phylogenetic analysis showed that AcCaspA and AcCCaspA were relatively

closely related to human/mouse effector caspases (CASP3/7/6), while AcCaspB and

AcCCaspB had closer relationships with human/mouse initiator caspases (Fig. 1B). AcCaspA,

AcCaspB, AcCCaspA, and AcCCaspB contain 315, 462, 437, and 624 amino acid residues,

respectively, and share 23.08% identities with each other. All four caspases possess the

conserved catalytic motifs of "SHG" and "QACRG" in p20 and the substrate binding

"GSWF/Y" motif in p10 (Fig. 1A). When expressed in HEK293T cells, all of the four

caspases induced marked apoptosis (Fig. 1C, D).

3.2 AcXIAP enhances AcCasp-mediated apoptosis.

In addition to the caspases, a XIAP structural homologue, named AcXIAP, was also identified

in jellyfish (Fig. 2A; Fig. S1). AcXIAP contains 779 amino acid residues and forms three

conserved BIR domains and one RING domain. Comparing with human XIAP (HsXIAP),

AcXIAP is 282 amino acids longer, due mostly to the elongated linker regions, and lacks an

apparent UBA domain. The predicted three-dimensional (3D) structure of AcXIAP differed

considerably from HsXIAP in the C-terminal RING-containing region and in the linkers

connecting the BIRs and the RING domains (Fig. 2B). To investigate whether AcXIAP, like

mammalian XIAP, inhibited caspase activity, ectopic co-expression of AcXIAP and each of

the four jellyfish caspases was conducted. AcXIAP proved to significantly increase, rather

than decrease, the apoptosis activities of all four jellyfish caspases (Fig. 2C, D).

3.3 AcXIAP promotes caspase activation via BIR3 as well as RING

AcCaspA, the simplest of the four jellyfish caspases in structure, was used to explore the

molecular mechanism underlying the caspase-activation effect of AcXIAP. AcXIAP was

found to interact and co-localize with AcCaspA (Fig. 3B, Fig. S2). Furthermore, in the

presence of AcXIAP, both the amount and the activation cleavage of AcCaspA were markedly

increased (Fig. 3B, panel 4). To determine the specific domain(s) involved in

AcXIAP-AcCaspA interaction, a series of AcXIAP truncates were created, i.e., BIR1, BIR2,

BIR3, and AcXIAP without the RING domain (AcXIAP-ΔRING) (Fig. 3A). The results of

co-expression analysis showed that like the wild type AcXIAP, the truncates AcXIAP-ΔRING

and BIR3 promoted the activation cleavage of AcCaspA (Fig. 3C, panel 3), which was

consistent with the Co-IP result showing that AcXIAP-ΔRING and BIR3 were able to interact

with and enhanced the activation of AcCaspA (Fig. 3C, panel 1). In contrast, BIR1 and BIR2

could not interact with AcCaspA. The activation effect of AcXIAP-ΔRING was apparently

dose-dependent (Fig. 3D). To further examine the effect of AcXIAP on AcCaspA activity,

recombinant AcCaspA was prepared, and its substrate cleavage specificity was determined

(Fig. S3). The specific cleavage sequence motif was then served as a substrate to determine

the cleavage activity of AcCaspA. The results showed that both AcXIAP and

AcXIAP-ΔRING significantly enhanced AcCaspA activity and, consistently,

AcCaspA-mediated apoptosis (Fig. 3E, F). However, the enhancing effect of AcXIAP-ΔRING

was significantly lower than that of AcXIAP, indicating the importance of the RING domain.

3.4 AcXIAP mediates caspase ubiquitination on the p20 subunit

Since in mammalian XIAP, the RING domain functions in ubiquitination, we examined the

effect of AcXIAP on AcCaspA ubiquitination. For this purpose, AcCaspA-M, an inactive

AcCaspA with mutation at the catalytic site, was created. Co-IP showed that both AcCaspA

and AcCaspA-M were ubiquitinated after co-expression with AcXIAP, but not after

co-expression with AcXIAP-ΔRING (Fig. 4A, B). To determine the AcCaspA region required

for ubiquitination, AcXIAP was co-expressed with the p20 or p10 region of AcCaspA. The

results showed that p20 was ubiquitinated in the presence of AcXIAP but not in the presence

of AcXIAP-ΔRING (Fig. 4C). In contrast, AcXIAP or AcXIAP-ΔRING did not affect the

ubiquitination of p10, but markedly increased the amount of p10 (Fig. 4C). To examine the

importance of the lysine residues in AcXIAP-mediated ubiquitination, all of the 28 lysine

residues in AcCaspA were mutated. The resulting mutant, AcCaspA-K28, could still be

ubiquitinated by AcXIAP (Fig. 4D).

3.5 Activation of caspase-mediated apoptosis is a common feature of non-bilaterian

XIAP

Since, as shown above, AcXIAP differed completely from mammalian XIAP in the role of a

caspase regulator, we wondered whether this was a general property of basal metazoan XIAP.

A search of the available database identified XIAP structural homologs in Porifera

(*Amphimedon queenslandica*) and Cnidaria (*Hydra vulgaris* and *Stylophora pistillata*) but not

in Ctenophora and Placozoa (Fig. 5A, Fig. S1). Like AcXIAP, the Porifera and Cnidaria XIAP

homologs possess 2 to 3 BIRs and one RING domain, but lack the UBA domain (Fig. S1). To

examine their regulatory effects, the XIAP of sponge, hydra, and coral (named AqXIAP,

HvXIAP, and SpXIAP, respectively) were each co-expressed with their respective caspases

(AqCasp, HvCasp, and SpCasp, respetively) in HEK293T cells. Subsequent analyses showed

that AqXIAP, HvXIAP, and SpXIAP all significantly enhanced caspase-mediated apoptosis

(Fig. 5B-E). To examine the effect of XIAP under the *in vivo* condition, hydra polyps with

XIAP knockdown were created (Fig. S4). Following treatment with colchicine, an inducer of

apoptosis (Cikala, Wilm, Hobmayer, Böttger, & David, 1999), the XIAP-knockdown polyp

exhibited much less cell death than the normal polyp (Fig. 5F).

314   4. Discussion

The moon jellyfish *Aurelia* is an emblematic medusa in Cnidaria, with a complex life cycle

like most medusozoans (Gold et al., 2019). Over twenty species of *Aurelia* have been

recognized (Kraus, Osadchenko, & Kosevich, 2022; Lawley et al., 2021). Among them, *A.*

coerulea and *Aurelia aurita* are invasive species widely distributed around the Northwest

Pacific and Atlantic coast of Europe (Kraus et al., 2022). In contrast to the invertebrate model

organisms *C. elegans* and *Drosophila*, cnidarians (hydra and coral) appear to have complex

caspase repertoires (Lasi, David, et al., 2010; Shrestha et al., 2020), but the functions of these

caspases remain elusive. In jellyfish, whose highly watery and gelatinous nature renders

cellular and *in vivo* studies difficult, no study on caspase or IAP has been documented. In the

present study, we identified and characterized four *A. coerulea* caspases, all of which were

found to induce robust apoptosis when expressed in human cells, suggesting that these

caspases probably worked via conserved mechanisms similar to that of mammalian caspases.

This is consistent with the presence of highly conserved catalytic motifs and substrate binding

sequences in the four jellyfish caspases. Nevertheless, it should be noted that although

AcCCaspA, one of the four identified jellyfish caspases, was grouped with human/mouse

effector caspases, it differs from mammalian effector caspases by possessing an N-terminal

CARD domain. On the other hand, AcCaspB, which possesses neither CARD nor DED

domain, proved to be more closely related to human/mouse initiator caspases than to the

effector caspases. These results indicate that the caspase classification criteria of the

vertebrate cannot be mechanically applied to invertebrate species.

XIAP is the most studied members of the IAP family and well-known as a potent

inhibitor of apoptosis in vertebrates and some invertebrates, mostly arthropods (Hanifeh &

Ataei, 2022; Hay, 2000; Miao, Qi, Li, Que, & Zhang, 2016). In this study, we found that

AcXIAP significantly enhanced, rather than inhibited, the apoptotic activities of all of the four

examined jellyfish caspases. Since the four jellyfish caspases differ considerably in sequence

characteristics and fall into different phylogenetic clades, the activating effect of AcXIAP is

probably not specific to a certain type of caspase. Similar to human XIAP, AcXIAP contains

the conserved RING domain and three BIR domains, however, AcXIAP differs noticeably

from human XIAP in the predicted 3D structures over the RING region and the linker regions,

especially in the linkers bridging BIR2-BIR3 and BIR3-RING. These linkers, unlike that in

human XIAP, are much longer and adopt, in parts, well-defined secondary structures, and

connect the BIRs and RING modules loosely into a higher-ordered structure more extended

than that of human XIAP. It is reasonable to speculate that these structural "peculiarities" may

account for, or at least contribute to, the unconventional quality of AcXIAP.

A previous study showed that the BIR3 domain of human XIAP directly interacted with

and inhibited caspase 9 (Deveraux et al., 1999). For AcXIAP, we found that it was also BIR3

that mediated the binding to the target caspase, though in this case the binding leading to a

stimulating, rather than inhibiting, effect on caspase activation. The observation that BIR3

alone could reproduce the caspase-activating effect of AcXIAP not only highlighted the vital

role of BIR3, but also indicated that BIR3 could play its role in an autonomous manner that

did not require the cooperation of the other domains of AcXIAP. Like AcXIAP, BIR3 enabled

more AcCaspA, in both uncleaved and cleaved forms, to be detected, suggesting that BIR3

probably facilitated caspase activation by holding the caspase in a more stable state. In

addition to BIR3, another AcXIAP truncate, AcXIAP-ΔRING, also exerted an augmenting

effect on AcCaspA activation, but this effect was significantly lower than that of AcXIAP.

Since the RING domain in mammalian XIAP mediates ubiquitination (Galbán & Duckett,

2010), it is possible that RING-associated ubiquitination is involved in AcXIAP regulation of

caspase activation. Indeed, we found that AcXIAP ubiquitinated AcCaspA at p20 in a

RING-dependent manner, thus indicating a conserved function of RING in invertebrate and

vertebrate XIAP. However, the ubiquitination mechanism of the RING of AcXIAP is probably

different from that of vertebrate XIAP, since mutation analyses demonstrated that none of the

28 lysine residues in AcCaspA was required for AcXIAP-mediated ubiquitination.

The IAPs, characterized by at least one BIR domain and a RING motif, occurred early in

metazoan evolution (Cao, Wang, Yang, Xie, & Yu, 2008; Deveraux & Reed, 1999). In this

study, we found that besides jellyfish, XIAP structural homologs existed in other

non-bilaterian organisms, including sponge, hydra, and coral. It is striking that, like AcXIAP,

all of these non-bilaterian XIAPs exhibited caspase-activating property, suggesting a

conserved pro-apoptosis function of XIAP in basal metazoans. This was supported by the *in*

vivo study in hydra, which showed that XIAP knockdown markedly weakened

colchicine-induced cell death. With these observations, one may speculate that XIAP in its

early forms probably act as a caspase activator via its BIR-mediated substrate binding and

RING-mediated ubiquitination, the latter in a lysine-independent manner. It is possible that

the BIR may exert, in a chaperon-like manner, on the substrate caspase a stabilizing effect that

is further strengthened by subsequent ubiquitination. It is also possible that both BIR

interaction and RING-mediated ubiquitination may alter the structure of the bound caspase in

a way that directly affects the activation cleavage of the caspase.

In conclusion, in this study we discovered the existence and the working mechanism of

pro-apoptosis XIAP in Cnidaria. Our results indicated that the basic modular organization of

XIAP had come into existence in non-bilaterians, and the co-operation of the different

task-oriented domains enables XIAP to play the role of a broad range caspase activator. In

Bilateria, the general domain-based functional division of the primitive XIAP is retained,

which preserves the nature of XIAP as a caspase regulator, but evolution-selected structural

alterations have led to a role-reversal of the XIAP from a positive regulator to a negative

regulator. These findings shed new light on the function, evolution, and regulation mechanism

of XIAP.

Acknowledgements

This work was supported by the Science & Technology Innovation Project of Laoshan

Laboratory (LSKJ202203000) and the Innovation Research Group Project of the National

Natural Science Foundation of China (42221005). The authors acknowledge the

Oceanographic Data Center, IOCAS, and System Simulation of Laoshan Laboratory for

providing the computing resources.

Author Contributions

Conceptualization and Funding acquisition: LS; Data curation: YC and HX; Methodology:

YC; Validation: YC and HX; Investigation: YC, MW, ZHY, and QYW; Formal analysis: YC;

Writing - original draft: YC; Writing - review & editing: LS and HX.

Declaration of interests

The authors declare no competing interests.

Data Availability

The data presented in this study can be accessed in the article/Supplemental Materials.

References

Bergmann, A., Yang, A. Y.-P., & Srivastava, M. (2003). Regulators of IAP function: coming to grips with the grim reaper. *Current opinion in cell biology, 15*(6), 717-724.

Bosch, T. C., Augustin, R., Gellner, K., Khalturin, K., & Lohmann, J. U. (2002). In vivo electroporation for genetic manipulations of whole Hydra polyps. *Differentiation, 70*(4-5), 140-147.

Budhidarmo, R., & Day, C. L. (2015). *IAPs: Modular regulators of cell signalling.* Paper presented at the Seminars in cell & developmental biology.

Cao, L., Wang, Z., Yang, X., Xie, L., & Yu, L. (2008). The evolution of BIR domain and its containing proteins. *FEBS letters, 582*(27), 3817-3822.

Chen, Y., Yu, C., Jiang, S., & Sun, L. (2022). Japanese Flounder HMGB1: A DAMP Molecule That Promotes Antimicrobial Immunity by Interacting with Immune Cells and Bacterial Pathogen. *Genes, 13*(9), 1509.

Chen, Y., Yuan, Z., & Sun, L. (2023). The evolutionary diversification and antimicrobial potential of MPEG1 in Metazoa. *Computational and Structural Biotechnology Journal, 21*, 5818-5828.

Cikala, M., Wilm, B., Hobmayer, E., Böttger, A., & David, C. N. (1999). Identification of caspases and apoptosis in the simple metazoan Hydra. *Current Biology, 9*(17), 959-962.

Clavier, A., Rincheval-Arnold, A., Colin, J., Mignotte, B., & Guénal, I. (2016). Apoptosis in

Drosophila: which role for mitochondria? *Apoptosis, 21*(3), 239-251.

Cohen, G. M. (1997). Caspases: the executioners of apoptosis. *Biochemical Journal, 326*(1), 1-16.

Deveraux, Q. L., Leo, E., Stennicke, H. R., Welsh, K., Salvesen, G. S., & Reed, J. C. (1999). Cleavage of human inhibitor of apoptosis protein XIAP results in fragments with distinct specificities for caspases. *The EMBO journal*.

Deveraux, Q. L., & Reed, J. C. (1999). IAP family proteins—suppressors of apoptosis. *Genes & development, 13*(3), 239-252.

Dunn, S. R., Phillips, W. S., Spatafora, J. W., Green, D. R., & Weis, V. M. (2006). Highly conserved caspase and Bcl-2 homologues from the sea anemone Aiptasia pallida: lower metazoans as models for the study of apoptosis evolution. *Journal of Molecular Evolution, 63*, 95-107.

Fan, T.-J., Han, L.-H., Cong, R.-S., & Liang, J. (2005). Caspase family proteases and apoptosis. *Acta biochimica et biophysica Sinica, 37*(11), 719-727.

Galbán, S., & Duckett, C. S. (2010). XIAP as a ubiquitin ligase in cellular signaling. *Cell Death & Differentiation, 17*(1), 54-60.

Gold, D. A., Katsuki, T., Li, Y., Yan, X., Regulski, M., Ibberson, D., . . . Greenspan, R. J. (2019). The genome of the jellyfish Aurelia and the evolution of animal complexity. *Nature ecology & evolution, 3*(1), 96-104.

Gujral, P., Mahajan, V., Lissaman, A. C., & Ponnampalam, A. P. (2020). Histone acetylation and the role of histone deacetylases in normal cyclic endometrium. *Reproductive Biology and Endocrinology, 18*, 1-11.

Hamajima, R., Iwamoto, A., Tomizaki, M., Suganuma, I., Kitaguchi, K., Kobayashi, M., . . . Ikeda, M. (2016). Functional analysis of inhibitor of apoptosis 1 of the silkworm Bombyx mori. *Insect biochemistry and molecular biology, 79*, 97-107.

Hanifeh, M., & Ataei, F. (2022). XIAP as a multifaceted molecule in Cellular Signaling. *Apoptosis, 27*(7-8), 441-453.

Hao, K., Xu, H., Jiang, S., & Sun, L. (2024). Paralichthys olivaceus MLKL-mediated necroptosis is activated by RIPK1/3 and involved in anti-microbial immunity. *Frontiers in immunology, 15*, 1348866.

Hay, B. (2000). Understanding IAP function and regulation: a view from Drosophila. *Cell Death & Differentiation, 7*(11), 1045-1056.

Huang, Q., Deveraux, Q. L., Maeda, S., Stennicke, H. R., Hammock, B. D., & Reed, J. C. (2001). Cloning and characterization of an inhibitor of apoptosis protein (IAP) from Bombyx mori. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research, 1499*(3), 191-198.

Jiang, S., Zhou, Z., Sun, Y., Zhang, T., & Sun, L. (2020). Coral gasdermin triggers pyroptosis. *Sci Immunol, 5*(54). doi:10.1126/sciimmunol.abd2591

Kraus, Y., Osadchenko, B., & Kosevich, I. (2022). Embryonic development of the moon jellyfish Aurelia aurita (Cnidaria, Scyphozoa): another variant on the theme of invagination. *PeerJ, 10*, e13361.

Kumar, S. (2007). Caspase function in programmed cell death. *Cell Death & Differentiation, 14*(1), 32-43.

Lasi, M., David, C. N., & Böttger, A. (2010). Apoptosis in pre-Bilaterians: Hydra as a model.

Apoptosis, 15, 269-278.

Lasi, M., Pauly, B., Schmidt, N., Cikala, M., Stiening, B., Käsbauer, T., . . . Knapp, R. T.
(2010). The molecular cell death machinery in the simple cnidarian Hydra includes an
expanded caspase family and pro-and anti-apoptotic Bcl-2 proteins. *Cell research,*
20(7), 812-825.

Lawley, J. W., Gamero-Mora, E., Maronna, M. M., Chiaverano, L. M., Stampar, S. N.,
Hopcroft, R. R., . . . Morandini, A. C. (2021). The importance of molecular characters
when morphological variability hinders diagnosability: systematics of the moon
jellyfish genus Aurelia (Cnidaria: Scyphozoa). *PeerJ, 9*, e11954.

Leu, J.-H., Kuo, Y.-C., Kou, G.-H., & Lo, C.-F. (2008). Molecular cloning and
characterization of an inhibitor of apoptosis protein (IAP) from the tiger shrimp,
Penaeus monodon. *Developmental & Comparative Immunology, 32*(2), 121-133.

Liao, W.-T., Yang, Y., & Wu, X.-F. (2002). Expression and functional analysis of an inhibitor
of apoptosis protein from Trichoplusia ni. *Biochemical and biophysical research*
communications, 293(2), 675-679.

Masuda-Ozawa, T., Fujita, S., Nakamura, R., Watanabe, H., Kuranaga, E., & Nakajima, Y.-i.
(2022). siRNA-mediated gene knockdown via electroporation in hydrozoan jellyfish
embryos. *Scientific reports, 12*(1), 16049.

Miao, G., Qi, H., Li, L., Que, H., & Zhang, G. (2016). Characterization and functional
analysis of two inhibitor of apoptosis genes in Zhikong scallop Chlamys farreri.
Developmental & Comparative Immunology, 60, 1-11.

Moya, A., Sakamaki, K., Mason, B. M., Huisman, L., Forêt, S., Weiss, Y., . . . Hayward, D. C.
(2016). Functional conservation of the apoptotic machinery from coral to man: the
diverse and complex Bcl-2 and caspase repertoires of Acropora millepora. *BMC*
genomics, 17(1), 1-20.

Qin, K., Jiang, S., Xu, H., Yuan, Z., & Sun, L. (2023). Pyroptotic gasdermin exists in
Mollusca and is vital to eliminating bacterial infection. *Cell Rep, 42*(5), 112414.
doi:10.1016/j.celrep.2023.112414

Qu, C., Sun, J., Xu, Q., Lv, X., Yang, W., Wang, F., . . . Wang, L. (2019). An inhibitor of
apoptosis protein (Es IAP1) from Chinese mitten crab Eriocheir sinensis regulates
apoptosis through inhibiting the activity of Es Caspase-3/7-1. *Scientific reports, 9*(1),
20421.

Reed, J. C. (2000). Mechanisms of Apoptosis. *The American Journal of Pathology, 157*(5),
1415-1430. doi:10.1016/s0002-9440(10)64779-7

Ross, C., Chan, A. H., von Pein, J. B., Maddugoda, M. P., Boucher, D., & Schroder, K. (2022).
Inflammatory caspases: toward a unified model for caspase activation by
inflammasomes. *Annual review of immunology, 40*, 249-269.

Shrestha, S., Tung, J., Grinshpon, R. D., Swartz, P., Hamilton, P. T., Dimos, B., . . . Clark, A.
C. (2020). Caspases from scleractinian coral show unique regulatory features. *Journal*
of Biological Chemistry, 295(43), 14578-14591.

Tenev, T., Ditzel, M., Zachariou, A., & Meier, P. (2007). The antiapoptotic activity of insect
IAPs requires activation by an evolutionarily conserved mechanism. *Cell Death &*
Differentiation, 14(6), 1191-1201.

Wang, P., Zhang, F., Sun, S., Wang, W., Wan, A., & Li, C. (2020). Experimental clearance

rates of Aurelia coerulea ephyrae and medusae, and the predation impact on
zooplankton in Jiaozhou Bay. *Journal of Oceanology and Limnology, 38*(4),
1256-1269.
Wiens, M., Krasko, A., Perovic, S., & Müller, W. E. (2003). Caspase-mediated apoptosis in
sponges: cloning and function of the phylogenetic oldest apoptotic proteases from
Metazoa. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research, 1593*(2-3),
179-189.
Xu, H., Jiang, S., Yu, C., Yuan, Z., & Sun, L. (2022). GSDMEa-mediated pyroptosis is
bi-directionally regulated by caspase and required for effective bacterial clearance in
teleost. *Cell Death & Disease, 13*(5), 491.
Xu, H., Yuan, Z., Qin, K., Jiang, S., & Sun, L. (2024). The molecular mechanism and
evolutionary divergence of caspase 3/7-regulated gasdermin E activation. *Elife, 12*,
RP89974.
Zhang, J., Jing, L., Li, M., He, L., & Guo, Z. (2019). Regulation of histone arginine
methylation/demethylation by methylase and demethylase. *Molecular medicine*
reports, 19(5), 3963-3971.

Figure Legends

Figure 1. Phylogenetic analysis and apoptotic activity of jellyfish caspases. (**A**) Schematic representation of the four jellyfish caspases. (**B**) Phylogenetic analysis of jellyfish and human/mouse caspases. Branch support values are expressed as percentages. (**C**) HEK293T cells were transfected with or without (control) the backbone vector (negative control), or the vector expressing AcCaspA, AcCaspB, AcCCaspA, or AcCCaspB. The cells were stained with Annexin V-FITC/propidium iodide (PI) and subjected to flow cytometry. (**D**) Bright filed images of HEK293T cells transfected as above in (C). Scale bars, 20 μm.

Figure 2. The effect of AcXIAP on caspase-mediated apoptosis. (**A**) Schematic representation of AcXIAP and HsXIAP. The positions of the BIR and RING domains are indicated on the protein. (**B**) The three-dimensional structures of AcXIAP and HsXIAP. The conserved domains are circled. (**C** and **D**) HEK293T cells were transfected with the backbone vector (control) or the vector expressing AcXIAP, AcCaspA, AcCaspB, AcCCaspA, or AcCCaspB, or co-transfected with the vector expressing AcXIAP and the vector expressing AcCaspA, AcCaspB, AcCCaspA, or AcCCaspB. The cells were subjected to microscopy (C) and analysis of apoptotic cell ratios (D). Scale bar, 20 μm. Values are the means of quadruplicate experiments and shown as means ± SD. **$p < 0.01$.

Figure 3. The interaction of AcXIAP with AcCaspA and its effect on AcCaspA activation. (**A**) Schematic representation of AcXIAP truncates. (**B**) HEK293T cells were transfected with or without the vector expressing Myc-tagged AcCaspA or Flag-tagged AcXIAP, or co-transfected with the vectors expressing Myc-tagged AcCaspA and Flag-tagged AcXIAP. The cell lysates were immunoprecipitated (IP). The IP and input samples were immunoblotted (IB) with the antibodies against the indicated tags. (**C**) HEK293T cells expressing Myc-tagged AcCaspA alone or AcCaspA plus various Flag-tagged AcXIAP truncates were immunoblotted as above. (**D**) AcCaspA was expressed alone or together with different doses of AcXIAP-ΔRING in HEK293T cells for 36 h. The expression levels of AcCaspA and AcXIAP-ΔRING were determined. (**E** and **F**) HEK293T cells were transfected with or

without the vector expressing AcCaspA, AcXIAP, or AcXIAP-ΔRING, or co-transfected with

the vectors expressing AcCaspA and AcXIAP or AcCaspA and AcXIAP-ΔRING. The

proteolytic activity of AcCaspA was assessed using fluorogenic substrate (E); the percentages

of apoptotic cells in the transfectants were determined (F). Values are the means of triplicate

(E) or quadruplicate (F) experiments and shown as means ± SD. **$p < 0.01$; *$p < 0.05$.

Figure 4. The effects of AcXIAP and AcXIAP-ΔRING on the ubiquitination of AcCaspA

variants. For all subfigures, HEK293T cells were transfected with the plasmids expressing

the indicated proteins that were tagged as follows: Ub (ubiquitin), HA-tagged; AcCaspA

variants, Myc-tagged; AcXIAP and AcXIAP-ΔRING, Flag-tagged. The cell lysates were

immunoprecipitated (IP). The IP and input (left and right panels, respectively, in all subfigures)

samples were immunoblotted (IB) with the antibodies against the indicated tags. (**A and B**)

The effects of AcXIAP and AcXIAP-ΔRING on the ubiquitination of AcCaspA (A) and

AcCaspA-M (B). (**C**) The effects of AcXIAP and AcXIAP-ΔRING on the ubiquitination of

AcCaspA p20 and p10. The relative densities of the p20 and p10 bands in the input samples

were calculated with ImageJ. (**D**) The effects of AcXIAP and AcXIAP-ΔRING on the

ubiquitination of AcCaspA-K28.

Figure 5. The effects of hydra, coral, and sponge XIAP on caspase-induced apoptosis. (**A**)

Metazoan phylogeny with the presence (√) or absence (×) of XIAP ortholog. (**B-E**) HEK293T

cells were transfected with or without (control) the vector or vectors expressing the indicated

caspase/XIAP. The cells were subjected to microscopy (B-D) and analysis of apoptotic cell

ratios (E). Scale bar, 20 μm. Values in (E) are the means of quadruplicate experiments and

shown as means ± SD. **$p < 0.01$. (**F**) Hydra polyps were treated with or without (control)

scrambled siRNA or HvXIAP-specific siRNA and then stained with propidium iodide (PI).

The polyps were observed with a fluorescence microscope. Scale bar, 200 μm.

Figure 1

602   Figure 5

603