Diel changes in the expression of a marker gene and candidate genes for intracellular amorphous CaCO₃ biomineralization in *Microcystis*

Apolline Bruley[1,#], Juliette Gaëtan[1,#], Muriel Gugger[2], Claire Pancrace[2,3], Maxime Millet[1], Geoffroy Gaschignard[1], Manuela Dezi[1], Jean-François Humbert[3], Julie Leloup[3], Fériel Skouri-Panet[1], Isabelle Callebaut[1], Karim Benzerara*[1], Elodie Duprat*[1]

[1] Sorbonne Université, Muséum National d'Histoire Naturelle, UMR CNRS 7590, Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, IMPMC – Paris, France
[2] Institut Pasteur, Université Paris Cité, Collection of Cyanobacteria – Paris, France
[3] Sorbonne Université, CNRS, INRAE, IRD, Université Paris Est Créteil, Université Paris Cité, UMR 7618. Institut d'Ecologie et des Sciences de l'Environnement de Paris (iEES-Paris) – Paris, France

#Equal contributions
*Corresponding authors
Correspondence: karim.benzerara@sorbonne-universite.fr & elodie.duprat@sorbonne-universite.fr

ABSTRACT

Many phylogenetically diverse cyanobacteria biomineralize intracellular amorphous calcium carbonate (iACC) inclusions. This includes several genotypes of the *Microcystis* genus, a potentially toxic, bloom-forming cyanobacterium found worldwide in freshwater ecosystems. While we ignore the biological function of iACC and the molecular mechanisms driving their formation, this process may impact local geochemical cycles and/or be used for bioremediation strategies. Recently, a marker gene of this biomineralization pathway, named *ccyA*, was discovered. However, the function of the calcyanin protein encoded by *ccyA* remains unknown. Here, based on an RNA-Seq approach, we assess the expression of the *ccyA* gene in *Microcystis aeruginosa* PCC 7806 during a 24 h day/night cycle. The *ccyA* gene shows a clear day/night expression pattern with a maximum transcript abundance at the end of the night. This is consistent with the assumption that iACC biomineralization is related with photosynthesis and may therefore follow a day/night cycle as well. Moreover, several genes directly co-localized upstream and downstream of *ccyA*, on the same DNA strand show a similar expression pattern, including a *cax* gene encoding a calcium/proton exchanger and a gene encoding a protein with a CoBaHMA domain, which is also present in the N-terminal region of calcyanins in many iACC-forming cyanobacteria. This suggests that they all could be part of an operon, and may play a concerted role in iACC formation. Last, several other genes involved in carbon concentrating mechanisms and calcium transport show an expression pattern similar to that of *ccyA*. Overall, this study provides a list of candidate genes that may be involved in the biomineralization of iACC by cyanobacteria and whose role could be, in the future, analyzed by biochemistry and genetics approaches.

Keywords: Biomineralization, amorphous calcium carbonate, cyanobacteria, RNA-seq, *cax* gene, CCM

Introduction

Biomineralization is a process through which some living organisms form mineral phases. It is widespread in all the domains of life (Boskey, 2003). In some cases, biomineralization is genetically controlled, *i.e.* it is tightly controlled by specific genes, as documented in diverse eukaryotic organisms (Brownlee et al., 2015; Gilbert et al., 2022; Knoll, 2003; Marin & Luquet, 2004; Monteiro et al., 2016). For example, key genes encoding, *e.g.*, carbonic anhydrases (Chen et al., 2018; Le Roy et al., 2014), Ca^{2+}, H^+ and HCO_3^- transporters (Mackinder et al., 2011), and/or molecules inhibiting mineralization (Marin et al., 2000, 1996), play pivotal roles in the control of calcium carbonate ($CaCO_3$) precipitation by eukaryotic organisms. However, there are much fewer documented cases of genetically controlled biomineralization by prokaryotes (Cosmidis & Benzerara, 2022; Görgen et al., 2020). For a long time, the only well documented case was the intracellular formation of magnetites (Fe_3O_4) and/or greigites (Fe_3S_4) by magnetotactic bacteria (Monteil et al., 2018). This process is regulated by a suite of more than 40 genes involved in the control of the size, shape and crystallochemistry of the mineral byproducts (Liu et al., 2023; Taoka et al., 2023; Uebe & Schüler, 2016). More recently, the formation of intracellular amorphous calcium carbonates (iACC) by several species of cyanobacteria has been evidenced as an additional case of genetically controlled biomineralization in bacteria (Benzerara et al., 2014; Couradeau et al., 2012). Before this discovery, cyanobacteria were usually thought to induce the precipitation of $CaCO_3$ as an indirect by-product of photosynthesis, with no apparent involvement of specific genes (Altermann et al., 2006). Cyanobacteria biomineralizing iACC are cosmopolitan (Ragon et al., 2014) and can massively sequester alkaline earth elements such as Ca, Sr, Ba and Ra, including their radioactive isotopes (Blondeau et al., 2018a; Cam et al., 2016; Mehta et al., 2022a). As a result, these bacteria may be interesting candidates for bioremediating some radioactive pollutions (Mehta et al., 2022a). Morever, the biomineralization of iACC is phylogenetically widespread, within the phylum of Cyanobacteria as well as in other bacterial phyla (Monteil et al., 2021). It is also found in several strains (but not all) of the well-studied *Microcystis* genus (Gaëtan et al., 2023). This cyanobacterium is particularly interesting since it can form blooms, with some genotypes being toxic for human and wildlife (Harke et al., 2016), causing numerous environmental issues in eutrophic freshwater ecosystems (Dick et al., 2021). Whether the formation of iACC by some *Microcystis* strains may impact their ecological success under certain conditions remains to be deciphered. Some putative biological functions have been suggested for iACC such as intracellular storage of inorganic carbon, intracellular pH buffering or their use as a ballast modifying cell buoyancy (Cosmidis & Benzerara, 2022). While all these functions could make sense considering the ecology of *Microcystis*, they presently remain speculative. Last, the formation of iACC, a concentrated intracellular sink of Ca, by the large biomasses of *Microcystis* during blooms may impact the local geochemical cycle of Ca in these environments (Gaëtan et al., 2023).

Despite the environmental importance of this process, the molecular mechanisms driving the biomineralization of iACC in cyanobacteria remain poorly known. It has been shown that it costs some energy to the cells (Cam et al., 2018; De Wever et al., 2019). Moreover, a genetic control has been proposed (Benzerara et al., 2022). Indeed, based on a comparative genomics approach, one family of genes, named *ccyA*, was found in phylogenetically diverse iACC-forming cyanobacteria and was absent from cyanobacteria not biomineralizing iACC. The *ccyA* gene was found in about one third of all publicly released genomes of *Microcystis* (Gaëtan et al., 2023). The protein encoded by *ccyA*, called calcyanin, lacks detectable full-length homologs with a known function in public databases. Therefore, its function remains unknown. Several clues suggest that calcyanin may play a role in calcium homeostasis (Benzerara et al., 2023, 2022). However, since the expression of the *ccyA* gene in mutants of cyanobacteria not forming iACC did not result in the formation of iACC, this biomineralization process likely involves additional genes (Benzerara et al., 2022). For example, some of these genes may play a role in the fact that iACC inclusions form within an intracellular compartment, the envelope of which remains of unknown chemical composition (Blondeau et al., 2018b). Such a compartmentalization may serve to achieve chemical conditions allowing the precipitation and the stabilization of ACC. Additional examples of molecular processes that may favor iACC precipitation are carbon-concentrating mechanisms (CCMs) (Li et al., 2016; Görgen et al., 2021), which have been abundantly

described in cyanobacteria and involve diverse genes (Badger & Price, 2003). Last, calcium transport genes
may also be involved in the accumulation of high amounts of calcium within iACC (De Wever et al., 2019).
Consistently, the presence/absence of the *ccyA* gene in genomes was correlated with the presence/absence
of genes involved in the transport and homeostasis of calcium and inorganic carbon (Benzerara et al., 2022).
Moreover, in some genomes, *ccyA* was even colocalized with such additional genes.
Overall, one way of gaining a better understanding about the molecular mechanisms involved in iACC
biomineralization is to look at the expression of the *ccyA* gene together with other genes, including those
neighboring *ccyA* in the genome and those mentioned above. The diurnal (day/night) cycle is an important
parameter driving large variations of gene expression in cyanobacteria (Stöckel et al., 2008; Welkie et al.,
2019; Zinser et al., 2009). For example, the diurnal cycle influences the expression of numerous genes
involved in key cellular processes such as photosynthesis, carbon fixation, nitrogen metabolism, and
circadian rhythms. Therefore, we have investigated the expression of the whole genome of *Microcystis*
aeruginosa PCC 7806, which contains the *ccyA* gene and forms iACC under laboratory culture conditions,
over a 24 h day/night cycle. In particular, we assessed whether the expression of genes either neighboring
ccyA in the genome and/or involved in CCM and calcium transport might be correlated with that of *ccyA*.
109

110   **Methods**

111 **Cultures and sampling**
112
The culture, sampling, nucleic acid extraction, and RNA sequencing were carried out at the Pasteur
culture of cyanobacteria (PCC) collection at the institut Pasteur in 2017. The photosynthetic photon flux
density provided during the day (50 mmol photons.$m^{-2}.s^{-1}$; cool white OSRAM L 18/640) was measured using
a LICOR LI-185B quantum/radiometer/photometer equipped with a LICOR LI-193SB spherical sensor. A
preculture of *M. aeruginosa* strain PCC 7806 was grown at 22 °C in 40 mL of the BG11 growth medium
(Rippka et al., 1979) amended with 10 mM $NaHCO_3$ under a 13/11 h day/night cycle. Then, the culture was
transferred into three separate flasks (named A, B and C). The triplicate cultures of 40 mL were progressively
transferred to a final set of three cultures with a volume of 1250 mL each, in fresh BG11 medium
supplemented with 20 mM $NaHCO_3$ and synchronized in an INFORS incubator with 1% CO_2 and agitation at
50-100 rpm under the same day/night cycle and temperature conditions. Synchronization involved three
consecutive transfers, aiming for an initial optical density (OD) at 750 nm of 0.05 and growing for a month.
Sampling was performed during the mid-exponential phase (OD of 0.6-0.7) for each biological replicate, over
a period of 24 h at eight time points: (i) 7 am, 60 min before the transition from night to day (t1_N); (ii) 9 am,
60 min after the transition from night to day (t2_D); (iii) 12 am (t3-D); (iv) 4 pm (t4_D); (v) 8 pm, 1 h before
the transition from day to night (t5_D); (vi) 10 pm, 60 min after the transition from day to night (t6_N); (vii) 2
am, *i.e.* the middle of the night (t7_N), and (viii) in the end of the night period (t8_N), 24 h after the first
sampling point. The sampling procedure involved the removal of 150 mL of culture, which were centrifuged,
rinsed with sterile water, and centrifuged again. The resulting pellets were flash-frozen in liquid nitrogen and
stored at -80 °C until further processing. To minimize batch effects, the samples were randomized before
subsequent processing.
133
Nucleic acid extraction
RNA extraction was carried out using a Trizol reagent (Life Technologies) and the NucleoSpin miRNA kit
(Macherey-Nagel). Firstly, 1 mL of Trizol preheated at 65 °C was added to the frozen pellets and incubated at
65 °C for 12 min with vortexing three times. After centrifugation (10 min at 4 °C and 12,000 rcf), the upper
phase was transferred to a new tube and mixed with 500 µL of Trizol preheated at 65 °C. RNA extraction was
achieved by adding 400 µL of chloroform, and incubating samples for 3 min at room temperature before
centrifugation (10 min at 4°C and 12,000 rcf). The upper phase was again transferred to a new tube for
extraction with 400 µL of Trizol and 400 µL of chloroform with similar incubation and centrifugation
conditions. Then, the upper phase obtained from this step was further processed using the NucleoSpin
miRNA kit according to the manufacturer's instructions. The final purified RNA was eluted in 20 µL of
RNase/DNase-free sterile water. The absence of genomic DNA contamination was confirmed by PCR

targeting the 16S rRNA gene using specific primers. The quality and quantity of RNA were assessed using a
Nanodrop spectrophotometer (ThermoFisher Scientific) and a Bioanalyzer (Agilent) with RNA Nano chips.
147
Library Preparation and RNA Sequencing
Library preparation was conducted using the TruSeq Stranded mRNA sample preparation kit (Illumina,
San Diego, California) following the manufacturer's instructions. Ribosomal RNA (rRNA) depletion was
performed using the Ribo-Zero rRNA removal kit (bacteria, #MRZB12424, Illumina) with 5 µg of total RNA.
The rRNA-depleted RNA was then fragmented using divalent ions at 94°C for 8 min. The resulting fragmented
RNA samples were reverse-transcribed using random primers, followed by complementary-strand synthesis
to generate double-stranded cDNA fragments. There was no need for an end repair step in this process. A 3'-
end adenine was added, and specific Illumina adapters were ligated to the cDNA fragments. The ligated
products were amplified by PCR. The quality of the resulting oriented libraries was assessed using
Bioanalyzer DNA1000 Chips (Agilent, # 5067-1504) and quantified using spectrofluorimetry (Quant-iT™ High-
Sensitivity DNA Assay Kit, #Q33120, Invitrogen). Sequencing was performed on the Illumina HiSeq2500
platform of the Institut Pasteur, generating single-end 65 bp reads with strand specificity. Each lane of
sequencing accommodated a mixture of 18 multiplexed samples.
161
Read processing
Raw RNA-seq reads (available online, see reference Raw transcriptomics data) were cleaned from
adapter sequences and low-quality sequences using an in-house program
(https://github.com/baj12/clean_ngs). Only sequences of at least 25 nt in length were considered for further
analysis. Transcript abundance was quantified from the cleaned read dataset using the Salmon software
(Patro et al., 2017). Pseudo-mapping of reads was done considering the coding sequences (CDS, including
pseudogenes) of the complete RefSeq genome assembly of *M. aeruginosa* PCC 7806 as a reference
(GCF_002095975.1). For each replicate, the estimated transcript abundances were normalized by Salmon
according to the transcript size, genome size (number of CDS) and sample size (number of reads).
171
Differential expression analysis
Total abundance data were further processed using the DiCoExpress R-script-based tool (Lambert et al.,
2020) in order to detect genes expressed differentially between different time points. First, normalization
was achieved using Trimmed Mean of the M-value (Robinson and Oshlack, 2010). During this procedure, 220
low-count transcripts were discarded, resulting in a total of 4,440 genes with normalized counts. Principal
component analysis (PCA) was performed on this normalized dataset using the R packages FactoMineR and
Factoshiny (Lê et al., 2008). Then, the differential expression between time points was tested using a
generalized linear model. For comparison between each time point, statistics of differential gene expression
are provided as Fold Change (FC) and p-values.
181
Gene functional annotation
In addition to the annotations provided by the NCBI prokaryotic genome annotation pipeline for the
genome assembly of *M. aeruginosa* PCC 7806
(https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_002095975.1), protein sequences translated from
CDS were processed using DeepNOG protein orthologous groups assignment (Feldbauer et al., 2021) as
implemented in the Genovi tool (Cumsille et al., 2023). An orthologous group with a COG accession number
is assigned to each gene, as well as a confidence score ranging within [0, 1]. The stringent threshold was set
at 0.8. Orthologs of genes involved in carbon-concentrating mechanisms (CCM) and calcium transport were
specifically detected as bidirectional best BLASTp hits using protein sequence queries described in (Tang et
al., 2022) (15 genes from *Synechocystis* sp. PCC 6803, and the *ecaA* gene from *Anabaena* PCC 7120) and (De
Wever et al., 2019) (13 genes from *Chroococcidiopsis thermalis* PCC 7203), respectively. The following
thresholds were used: E-value cut-off of 1E-6, ≥30% identity and 70% coverage.
194
Sequence and structure analysis of unknown proteins
General information on the three domains of unknown function (DUF) encoded by genes located in the
genomic neighborhood of the *ccyA* gene was extracted from their respective InterPro entries (Paysan-Lafosse
et al., 2023; DUF5132: IPR033456; DUF454: IPR00740; DUF1269: IPR009200). AlphaFold2 (AF2) 3D structure

models of the three DUF-containing proteins and two hypothetical proteins neighboring *ccyA* were build
using ColabFold (Mirdita et al., 2022). Foldseek (van Kempen et al., 2023) was used to search for structural
similarities against the Protein Data Bank (PDB) and AlphaFold Database (AFDB). 3D structures were
manipulated using Chimera (Pettersen et al., 2004). Hydrophobic cluster analysis (HCA) (Callebaut et al.,
1997) was used to analyze protein secondary structure features.
204

Results

Overall expression pattern
A principal component analysis (PCA) was performed on the transcript abundances of the 4440 genes, for
each replicate at each sampling time (Figure 1). All the data points were projected onto a two-dimensional
plane defined by the first two principal components, which accounted for 64.69% of the total variance
(Figure 1). Each replicate (*i.e.* three independent cultures) at a single time showed only minor variations
between them, at least along axis 1 (accounting for almost 50% of the variance), and were distinctly
separated from the replicates at other time points. Furthermore, the resulting plot exhibited a clear
separation along axis 1 between night (N) and day (D) points.
214

Figure 1 - **Principal component analysis of the transcript abundances of 4440 *M. aeruginosa* genes**
during a day/night cycle. Abundances are expressed as log base 2 of the normalized counts, for each
replicate (a-c) at each sampling time (t1-t8). The 24 samples are projected onto the first two principal
components (Dim. 1 and Dim. 2), explaining 48.9% and 15.8% of the variance of the dataset,
respectively. Samples/replicates collected at different times over the day/night cycle appear with a
different color code. The "_N" and "_D" codes in the sample names refer to night and day, respectively.

Expression of the *ccyA* gene
In order to analyze the expression profile of the *ccyA* gene during the 24 h day/night cycle, we plotted
the normalized expression of *ccyA* in all replicates at the different time points (Figure 2). A clear day/night
pattern can be observed in the expression of *ccyA*, with a maximum transcript abundance at the end of the
night (t8_N, Table 1) and a minimum transcript abundance at 12 am during daytime (t3_D, Table 1). Pairwise
comparisons of *ccyA* transcript abundance at different time points show that the expression of *ccyA* at
daytime points (t2_D to t5_D) did not significantly differ between each other, but were significantly different
(p-value < 0.01) from some of the nightime points, t7_N and t8_N, respectively (Table S1). The 10 pm
nighttime point (t6_N) did not significantly differ from any others except from the 12 am daytime point

(t3_D; p-value < 0.05) and can therefore be considered as a transition point between day and night. When
compared with the mean expression of the 4440 genes of the whole transcriptome, the mean expression of
the *ccyA* gene was significantly higher (p-value < 0.05) at two-time points: 2 am (t7_N) and 7 am (t8_N)
(Table 1).
236
237
238

240 **Figure 2** - **Abundance profile of *ccyA* transcripts during a day/night cycle.** Abundances are expressed as
241 log base 2 of the normalized counts. The abundance mean for the three replicates is shown by a black
242 filled circle. Replicates a, b and c are represented by different symbols. For each sampling time (t1-t8),
243 error bar represents the standard error on the mean of the three replicates. Grey shaded areas outline
244 the night periods.

	t1_N(*) 7 am	t2_D 9 am	t3_D 12 am	t4_D 4 pm	t5_D(*) 8 pm	t6_N(*) 10 pm	t7_N(**) 2 am	t8_N(**) 7 am
Mean (*ccyA* gene, n=3 replicates)	10.31	9.07	8.88	9.12	9.31	9.73	10.39	10.43
Standard deviation (*ccyA* gene, n=3 replicates)	0.15	0.22	0.23	0.19	0.14	0.34	0.33	0.12
Mean (4440 genes, n=3 replicates)	6.87	7.35	7.39	7.44	7.44	7.36	7.23	7.18
Standard deviation (4440 genes, n=3 replicates)	3.79	2.45	2.57	2.67	2.61	2.87	3.21	3.33

245 **Table 1** - **Comparison between the transcript abundances of *ccyA* and the whole transcriptome, for
246 each time point.** Abundances are expressed as log base 2 of the normalized counts. When means
247 significantly differ, p-values of Wilcoxon signed rank test with continuity correction are depicted as
248 follows: (**): 0.01 − 0.05, (*): 0.05 − 0.1. Night points are shaded in grey.

249
Search of differential gene expression during the day/night cycle

Subsequently, a comprehensive analysis was conducted throughout the genome to identify all genes
exhibiting diurnal variations in their expression, using the minima and maxima of *ccyA* gene expression (t3_D
and t8_N) as reference time points (Figure 3). Overall, 1799 of the 4440 total genes showed a significant
difference in expression (at least 2-fold) between 12 am (during the day, t3_D) and 7 am (during the night,
t8_N) (Figure 3 and Table S2). Among these 1799 genes, 906 were significantly more expressed during the
day (t3_D), and conversely, 893 were over-expressed during the night (t8_N).
Figure 3 provides the annotations for the genes with the most substantial expression differences between
these two time points (-log10(p-value)>60): genes overexpressed during the day included a NAD(P)H-
quinone oxidoreductase and a protein of the UDP-glucose/GDP-mannose dehydrogenase family. The genes
overexpressed at night included: two genes encoding hypothetical proteins, a pseudo-gene encoding a
protein containing a helix-turn-helix domain, a gene encoding a protein from the SDR oxidoreductase family,
and a gene encoding a dihydroorotate dehydrogenase-like protein.

Figure 3 - **Differential expression of the whole transcriptome between sampling time t3_D (12 am,**
during the day period) and t8_N (7 am, end of the night period). Binary logarithms of the fold change
(FC) values (x-axis) are defined for 3116 CDS. CDS with a negative log2(FC) value correspond to
transcripts significantly more abundant at 12 am (t3_D) compared with 7 am (t8_N) (gold area; 906
CDS). CDS with a positive log2(FC) value correspond to transcripts significantly more abundant at 7 am
(t8_N) compared with 12 am (t3_D) (grey area; 893 CDS). The *ccyA* gene is indicated by a red dot. The
horizontal red dashed line indicates the significance p-value threshold after Bonferroni correction (-
log10(P-value)=5). The vertical red dashed lines indicate FC=2 and FC=0.5, *e.g.*, 2-fold abundance ratio
between the two sampling times. RefSeq accessions and annotations are indicated for CDS with P<2.0e-
66). See Suppl. Table S2 for details.

Genomic neighborhood of *ccyA*
Figure 4 displays the map of the 20 genes (18 genes and 2 pseudo-genes) located upstream (10 genes)
and downstream (10 genes) to *ccyA*. This number of 20 genes was set arbitrarily and considered as a close
neighborhood. When available, the functional annotations were extracted from the NCBI RefSeq database
(Table 2). In this database, the *ccyA* gene is incorrectly annotated as a gene coding for the cell envelope
biogenesis protein OmpA (Benzerara et al., 2022). Interestingly, three transposition elements were found
near the *ccyA* gene: an ISL3 family and an IS1 family transposases, both pseudo-genes located at -6 and -5
genes compared to *ccyA*, and an IS1634 family transposase located at +2 genes compared with *ccyA*. The
gene located directly upstream of *ccyA* was annotated as encoding a domain-of-unknown-function
(DUF5132)-containing protein, and the one directly downstream a calcium/proton exchanger (*cax* gene),

previously characterized in eukaryotes (Hirschi et al., 1996; Waight et al., 2013). At position -3 (upstream of *ccyA*), another DUF-containing protein (DUF454) was detected.

Figure 4 - **Schematic representation of the *ccyA* gene neighborhood in *M. aeruginosa* PCC 7806 complete genome assembly (NCBI genome browser display).** The *ccyA* gene is outlined by a green box. RefSeq protein accession, gene name (if defined), gene orientation (red for minus; green for plus) and locus tag are indicated for the 10 CDS located upstream (-10 to -1) and the 10 CDS located downstream (+1 to +10) with respect to *ccyA*. Protein functional annotation is provided in Table 2. This genomic region spans within the nucleotide position interval [4515650, 4535259] in the NCBI genome assembly NZ_CP020771.1. CDS boundaries and features correspond to RefSeq annotations (GCF_002095975.1). Gene symbols stand for *shc*: squalene--hopene cyclase; *tsf*: translation elongation factor Ts; *rspsB*: 30S ribosomal protein S2 ; *cax*: calcium/proton exchanger; *acs*: acetate--CoA ligase; *psaC*: photosystem I iron-sulfur center protein PsaC.

Table 2 - **Neighborhood of the *ccyA* gene in *M. aeruginosa* PCC 7806 complete genome assembly**. General features and annotations are listed for the ten CDS located upstream (-10 to -1) and the ten CDS located downstream (+1 to +10) of the position of *ccyA* (in bold, protein improperly annotated in RefSeq as OmpA). RefSeq annotations correspond to the assembly accession GCF_002095975.1 (NZ_CP020771.1).

Protein accession (NCBI RefSeq)	CDS position		Gene orientation	Start (nt)	End (nt)	Gene name	Protein functional annotation (NCBI RefSeq)
	NCBI RefSeq	relative to *ccyA*					
WP_002741645.1	4262	-10	+	4515650	4516447	*rpsB*	30S ribosomal protein S2
WP_002741643.1	4263	-9	+	4516564	4517319	*tsf*	translation elongation factor Ts
WP_002741642.1	4264	-8	+	4517577	4519484	*shc*	squalene--hopene cyclase
WP_051048450.1	4265	-7	-	4519545	4519886		transcriptional regulator
pseudo (partial=3')	4266	-6	+	4520389	>4520916		ISL3 family transposase
pseudo (partial=3')	4267	-5	-	<4520947	4521127		IS1 family transposase
WP_002753396.1	4268	-4	-	4521343	4521633		DUF5132 domain-containing protein
WP_036399769.1	4269	-3	+	4521771	4522358		DUF454 family protein
WP_228036328.1	4270	-2	+	4522489	4522689		hypothetical protein
WP_002741624.1	4271	-1	+	4522868	4523182		DUF5132 domain-containing protein
WP_084990073.1	**4272**	**0**	**+**	**4523229**	**4524278**		**cell envelope biogenesis protein OmpA**
WP_084990074.1	4273	+1	+	4524268	4525377	*cax*	calcium/proton exchanger
WP_084990075.1	4274	+2	-	4525733	4527469		IS1634 family transposase
WP_002734319.1	4275	+3	-	4527740	4527985	*psaC*	photosystem I iron-sulfur center protein PsaC
WP_002741619.1	4276	+4	+	4528237	4530207	*acs*	acetate--CoA ligase
WP_084990076.1	4277	+5	+	4530562	4531158		hypothetical protein
WP_002741615.1	4278	+6	+	4531198	4531722		GNAT family N-acetyltransferase
WP_002731545.1	4279	+7	-	4531850	4532356		DUF1269 domain-containing protein
WP_084990077.1	4280	+8	-	4532613	4533059		DUF1269 domain-containing protein
WP_002756663.1	4281	+9	-	4533361	4533978		CBS domain-containing protein
WP_002741611.1	4282	+10	-	4534069	4535259		glycosyltransferase family 4 protein

AlphaFold 2 (AF2) 3D structure models of two hypothetical proteins, as well as five proteins containing
DUFs present in the *ccyA* genomic neighborhood were analyzed (Figure 5) and compared with experimental
3D structures using Foldseek, in order to highlight possible structural and functional relationships. (i) Analysis
of the DUF454 family protein (WP_036399769.1; position -3 relatively to *ccyA*) indicates that its C-terminal
end matches the C-terminal end of the DUF454 IPR profile (inner membrane protein YbaN,from *E.coli*).
Meanwhile, its N-terminal part matches the Pfam HMA_2 profile (PF19991), corresponding to a CoBaHMA
domain (Gaschignard et al., 2024). The AF2 3D structure model also consistently highlights this two-domain
architecture, with the C-terminal, DUF454-like domain folding as a helical hairpin, while the CoBaHMA
domain possesses the basic conserved amino acids characteristic of this family (Figure 5-A). (ii) The DUF5132-
containing protein (WP_002741624.1; position -1 relatively to *ccyA*) is covered by the DUF5132 profile (aa
48-89) over a little less than half of its length and contains a predicted transmembrane segment in its N-
terminal part (aa 12-32). The AF2 3D structure model, characterized overall by good pLDDT values, indicates
a putative assembly, in a hairpin-like conformation, of the helical transmembrane segment and another
alpha-helix preceding a long, soluble helix matching the DUF5132 profile (Figure 5-B). (iii) A second member
of this family, with similar structural features is also present in the *ccyA* neighborhood (WP_002753396.1;
position -4 relatively to *ccyA*). No significant relationship with any known 3D structures could be detected by
FoldSeek. (iv) A hypothetical protein (WP_228036328.1; position -2 relatively to *ccyA*) corresponds to a
truncated part of a hypothetical protein referenced under RefSeq WP_271988195.1, which is also composed
of a CoBaHMA domain sharing the basic signature of this family (Figure 5-C). (v) Another hypothetical protein
(WP_084990076.1; position +5 relatively to *ccyA*) is predicted as a long coiled-coil (data not shown). (vi) and
(vii) Finally, two DUF1269-containing proteins (WP_002731545.1 and WP_084990077.1; genes at +7 and +8
relatively to *ccyA*), corresponding to two isoforms differing by their C-terminal end, match the DUF1269 IPR
profile over their whole lengths and include a long glycine-zipper motif embedded in a soluble domain, which
based on Foldseek searches, corresponds to a ferredoxin fold, as found for instance in the copper tolerance
CutA1 protein (pdb 1V6H, Bagautdinov, 2014) and in the nickel-responsive transcription factor NikR (pdb
2BJ3, Chivers and Tahirov 2003) (Figure 5-D). This glycine-zipper motif is similar, albeit more hydrophobic, to
the GlyZip motif in the C-ter domain of calcyanins, composed of two parts separated by a proline preceded
by a small amino acid (here a serine, while it is a glycine in calcyanins) (Figure 5-D). Overall, this analysis
highlighted some features in the neighborhood of the *ccya* gene, which were not captured by automatic
annotations, including motifs that are found in calcyanins of all cyanobacteria (gly-zip and hairpin-like
helices) as well as calcyanins of some species (CoBaHMA domains).

A) WP_036399769.1 DUF454 family protein

CoBaHMA
HMA_2 N-ter

DUF454-Cter

DUF454-Nter

DUF454-Cter

B) WP_002741624.1 DUF5132-domain containing protein

TM

DUF5132

C) WP_2719881195.1 hypothetical protein
(full length of WP_271988195.1)

CoBaHMA

D) WP_002731545.1
DUF1269 domain-containing protein

GlyZip

ferredoxin fold

pIDDT: ■ Very low (<50) ■ Low (60) ■ OK (70) ■ Confident (80) ■ Very high (>90)

Figure 5 - AlphaFold2 3D structure models of some proteins encoded by the genomic neighbors of *ccya*. The
3D structure models are represented as ribbons colored according to the pIDDT values (color code given at
bottom). In panels A and C, the basic amino acids constituting the signature of the CoBaHMA family are shown as
balls and sticks. (A) The CoBaHMA_DUF454-Cter architecture (encoding gene in position -3 relatively to *ccyA*) is
compared with a protein entirely covered by the DUF454 IPR profile (inner membrane protein YbaN (YABN_ECOLI,
UniProt P0AAR5); insert). (B) DUF5132 protein (encoding gene in position -1 relatively to *ccyA*). TM stands for
TransMembrane. (C) Hypothetical protein (encoding gene in position -2 relatively to *ccyA*) with a typical CoBaHMA
domain. (D) DUF1269 domain-containing protein (encoding gene in position +7 relatively to *ccyA*) containing a
GlyZip motif, predicted to fold as a helical hairpin (see HCA plot and sequence alignment in Figure S1).

Correlated expression of other genes with the *ccyA* gene
The expression patterns of genes exhibiting a significant difference in their circadian expression between
daytime (t3_D, 12 am) and nighttime (t8_N, 7 am) were compared with that of *ccyA* to figure out which ones
may have a maximum expression at the same time (Figure 6; Table S3). We note that such gene expression
variations over a day/night cycle are expected in an autotrophic organisms and at least some correlations are
likely not causal. However, their identification offers one possibility to better understand which other
pathways may be activated at the same time as *ccyA* is expressed. As a result, two subsets were identified:
on the one hand, 773 genes showed a highly positive correlation with the expression pattern of *ccyA* (with a
Pearson correlation coefficient rho > 0.75, this threshold being arbitrarily fixed; grey subset in Figure 6; see
also Figure S2 and Table S3); on the other hand, 706 displayed a negative correlation with the expression
pattern of *ccyA* (rho < -0.75, yellow subset in Figure 6; see also Figure S2 and Table S3). Figure 6 displays the
significance level (p-value) of the correlations between the expression profiles of these 1479 genes and that
of *ccyA*, and provides annotation for the genes with the most significant correlation.
The genes for which the expression patterns were the most significantly positively correlated with that of
the *ccyA* gene were: (i) an intermembrane mitochondrial space (IMS) domain-containing protein, (ii) a GTP-
binding protein, and (iii) a DUF-containing protein (DUF4278). The genes the most significantly anticorrelated

with *ccyA* expression profile were annotated as (i) an ABC transporter ATP-binding protein/permease, (ii) an
adenylyl-sulfate kinase, and (iii) a hypothetical protein.

Among the 20 neighboring genes of *ccyA*, eight showed an expression profile either positively or
negatively correlated with that of *ccyA* (see Figure S3 for a time plot of expression of all 20 genes).
Interestingly, the expression profile of the two genes adjacent to *ccyA*, located on the same DNA strand, *i.e.*
encoding for a calcium/proton exchanger (CAX) and DUF5132 (Table 2), displayed a strong positive
correlation with the expression profile of *ccyA* (Figure 6 and 7). The expression profile of the gene coding for
a DUF454 family protein (position -3 relatively to *ccyA*; Table 2), exhibited one of the strongest positive
correlations with the expression profile of *ccyA* (Figure 6), with a nearly identical expression profile (Figure
7). Finally, the expression profiles of two other genes, located at +10 and +5 genes with respect to *ccyA*
showed a weaker but significant positive correlation with the expression profile of *ccyA*. The gene at +10 was
annotated as a glycosyltransferase family 4 protein. The gene at +5 codes for a hypothetical protein
predicted as a long coiled-coil (Table 2).

Moreover, proteins involved in calcium transport and in CCM were searched in the genome of PCC 7806
(Table S4). In total, 24 genes involved in CCM and/or their homologs and 24 genes involved in calcium
passive or active transport and their homologs were identified. The expression patterns of seven out of the
24 genes involved in CCM were correlated negatively with the expression pattern of *ccyA*, whereas the
expression patterns of five of CCM genes were positively correlated with the expression pattern of *ccyA*
(Figure 6, in magenta; see Figure S4 for a time plot of the expression of all CCM genes). Among the 24 genes
involved in calcium transport, the expression patterns of five of them were negatively correlated with that of
ccyA, while the expression patterns of ten of them were positively correlated with that of *ccyA* (Figure 6, in
cyan except the *cax* gene in green; see Figure S5 for a plot of time expression of all "Ca genes").

385
386
Figure 6 - CDS with a day/night expression pattern strongly (anti-)correlated with that of *ccyA*.
The Pearson correlation coefficient (rho) is highly negative for 706 CDS (left). The Pearson correlation
coefficient is highly positive for 773 CDS (right). The distribution of correlation coefficients over the
4440 CDS is shown in Figure S2. RefSeq accessions and annotations are indicated for the 3 CDS with the
strongest negative and positive correlations (see Table S3 for details). CDS belonging to the carbon
concentrating mechanism or the calcium transport system are represented as magenta and cyan dots,
respectively. CDS located in the [-10, +10] neighborhood of *ccyA* (Table 2 and Figure 4) are represented
by green and red dots (for positive and negative DNA strands, respectively). The *cax* gene located at a
genomic position of +1 from the *ccyA* gene also belongs to the calcium transport system and appears in
green (see Table S4 for details).

Figure 7 - **Abundance profiles of three genomic neighbors of the *ccyA* gene during a day/night cycle.** The CDS located at positions -3 (blue), -1 (purple) and +1 (orange) correspond to the *ccyA* neighbors with transcript abundance profiles the most correlated to *ccyA* transcript abundance profile: Pearson correlation coefficient of 0.97 (p-value=1.7e-15), 0.95 (p-value=8.0e-13), 0.94 (p-value=1.2e-11), respectively.

Discussion

The identification of a marker gene of iACC biomineralisation in cyanobacteria was a first step towards understanding the molecular mechanisms controlling the formation of these biomineral phases (Benzerara et al., 2022). However, up to now, only the presence/absence of the gene and the structural features of the encoded protein have been investigated (Benzerara et al., 2022; Gaëtan et al., 2023; Gaschignard et al., 2024). Here, we explored whether this marker gene is transcribed in an organism capable of forming iACC, how this may be influenced by day/night cycles and how the expression of neighboring genes compares.

Several tools can be used for assessing the transcription of a large number of genes within a genome, including DNA microarray and high-throughput RNA sequencing (RNAseq) (Wang et al., 2009). Microarrays are limited by the use of probes: the number of studied genes remains limited by the available knowledge of the targeted genome. By contrast, RNAseq allows targeting all the mRNAs in a cell and has the advantage of measuring the level of transcriptional activity in the organisms (Mantione et al., 2014). Moreover, RNAseq can achieve higher resolution (*i.e.* higher accuracy in the estimation of gene expression) than microarrays. Because this technique does not suffer from microarray-based limitations such as background noise and saturation, it can also achieve a lower detection limit, and thus detect very low gene expression (Zhao et al., 2014). Straub et al. (2011) already studied the expression of the whole genome of *M. aeruginosa* PCC 7806 over a 24-h day/night cycle, using DNA microarray. Using their publicly available dataset, we determined that the expression level of the *ccyA* gene (MIC_2500) was not significantly different from the background noise. Here, the use of RNA-seq on the same organism enabled the detection and analysis of the *ccyA* gene expression dynamics. Indeed, the expression level of *ccyA* in our dataset was above the mean expression of the 4440 genes of *Microcystis* at all time and this was statistically significant for several time points during the night. We can thus conclude that the *ccyA* gene is clearly expressed in an iACC-forming strain.

In line with the study by Straub et al. (2011) on the dynamics of gene expression in *M. aeruginosa* PCC 7806, we found, as expected, that the day/night cycle is an important factor driving variations in global gene expression in PCC 7806. Cyanobacteria have a highly defined circadian clock, and this day/night gene expression pattern has been demonstrated in several strains (Pattanayak & Rust, 2014; Stöckel et al., 2008; Zinser et al., 2009). Straub et al. (2011) showed that the metabolism of *M. aeruginosa* undergoes compartmentalization between the light and dark periods. Specifically, during the light period, processes including carbon uptake, photosynthesis, and the reductive pentose phosphate pathway result in glycogen synthesis. Conversely, during the dark period, glycogen degradation, the oxidative pentose phosphate pathway, the TCA branched pathway, and ammonium uptake promote amino acid biosynthesis. Furthermore, the biosynthesis of secondary metabolites, such as microcystins, aeruginosin, and cyanopeptolin, predominantly occurs during the light period.

A clear day-night expression pattern was also observed for the *ccyA* gene: the expression was significantly higher than the average genome expression from the middle (2 am, t7_N) to the end of the night (7 am, t8_N). In all other time points of the 24-h cycle, the expression level of this gene did not statistically differ from the average expression of all genes. The gene is therefore overtranscribed during the second half of the night. While mRNA abundance provides valuable information about gene expression, it does not always directly correlate with the protein levels in the cell (Ingolia et al., 2009; Riba et al., 2019; Schwanhäusser et al., 2011). Several factors, including translation, initiation and elongation rate (Riba et al., 2019), post-translational modifications, protein stability, and degradation rates (Christiano et al., 2014; Gancedo et al., 1982) may impact the relationship between mRNA abundance and protein content. As such, it is difficult to infer the abundance of calcyanin at a given time from the transcript expression level only. However, one can speculate that the protein may be more produced during the second part of the night, and remain at an optimal abundance during the day when it may most function. In the future, direct measurements of the protein abundance should be performed using proteomics in order to quantify the calcyanin content in cells.

Interestingly, we showed that a Ca^{2+}/H^+ antiporter (*cax* gene) and two DUF-containing proteins were not only directly co-localized upstream and downstream of *ccyA*, on the same DNA brand but they were also co-expressed with *ccyA*. This suggests that those three genes could be part of a functioning unit of

DNA that could be transcribed together into a single mRNA strand, *i.e.* an operon, and may play a
concerted role in iACC formation.
The process of iACC formation necessitates the accumulation of calcium ions at concentrations higher
than the cytosolic concentrations typically measured in cells (Cam et al., 2015). Therefore, it may be
suggested that the protein encoded by the *cax* gene is involved in the build-up of these accumulations.
However, the functioning of CAX proteins raises questions. Indeed, the single Ca^{2+}/H^+ exchanger found in
Synechocystis PCC 6803 has been shown to be located in the plasma membrane and to catalyze Ca^{2+}
efflux out of the cells (Waditee et al., 2004). By contrast, it has been suggested that CAX proteins in
plants may have different locations and efflux Ca^{2+} from the cytoplasm either to the extracellular space or
to intracellular vacuoles (Shigaki et al., 2006). Investigating further the similarities/dissimilarities of the
CAX protein of *M. aeruginosa* PCC 7806 with already known CAX proteins and locating these transporters
within *M. aeruginosa* PCC 7806 cells would therefore be crucial to better understand their potential
involvement in iACC formation.
The gene encoding a DUF454 protein that co-localizes (position -3 relatively to *ccyA*) and co-expresses
with *ccya* is particularly interesting. Indeed, we show here that this DUF454 protein bears a CoBaHMA
(after Conserved Basic residues HMA) domain (Gaschignard et al., 2024). Strikingly, the calcyanins of
many cyanobacteria are similarly composed of a CoBaHMA domain in their N-terminal region and a C-
terminal three-fold repeat of a long glycine zipper motif conserved in all iACC-forming cyanobacteria
(Benzerara et al 2022). This is not the case for the calcyanins of *Microcystis,* which does not contain a
CoBaHMA domain but instead displays an unknown domain in their N-terminal region. Since the
discovery of the calcyanin family, it has been surprising to observe such a broad diversity in their N-
terminal domains despite an expected common function. Here, although we still ignore the function of
calcyanin, the discovery of a CoBaHMA-encoding gene neighboring the *ccyA* gene suggests that
Microcystis calcyanin may work in concert with other proteins, including a DUF454 protein and resulting
in a protein complex with a possibly similar architecture in *Microcystis* and other iACC-forming
cyanobacteria. Moreover, the presence in the close genomic neighborhood of *ccyA* of genes encoding for
another protein of the CoBaHMA family (-2 position relative to *ccyA*), as well as proteins possessing a
GlyZip-like motif (+7 and +8 positions relative to *ccyA*) relatively similar to the GlyZip motif in the C-ter
domain of calcyanins additionally suggests the possibility of complementary roles of these proteins for a
function to identify.
Moreover, building upon the approach followed by De Wever et al. (2019), we conducted a
comprehensive search for annotated genes involved in CCM and calcium transport, as well as their
homologous counterparts in the genome of *M. aeruginosa* PCC 7806. Firstly, a greater number of genes
associated with CCM or their homologs appeared to be expressed preferentially during the day instead of
night (nine *vs* four, respectively), compared with genes associated with calcium transport (three *vs* nine,
respectively) (Table S4). Similarly, Straub et al. (2011) showed that the transcript abundance of CCM
genes showed a peak in expression just after the night/day transition and then a rapid decrease after the
day/night transition. The elevated concentration of CO_2 in carboxysomes resulting from the CCM is a way
to improve carbon fixation (Price and Howitt, 2011). The expression of CCM genes during the day is
therefore expected. Yet, among the 24 genes identified as involved or potentially involved in CCM, 5 had
an expression pattern positively correlated with that of *ccyA*, *i.e.* maximum at night (Table S4). Notably,
two of these genes (*CmpC* and *CmpD*) encode two subunits of the inorganic carbon transporter BCT1
involved in the cyanobacterial CCM (Price & Howitt, 2011). BCT1 is a high-affinity but low-flux HCO_3^-
transporter. It is composed of four subunits, and among them, CmpC and CmpD are extrinsic cytoplasmic
proteins that possess binding sites for ATP (Price & Howitt, 2011). The co-expression of these five genes
with *ccyA* suggest that they may be bicarbonate transporters feeding iACC biomineralization.
Alternatively, it has been shown that the gene expression of these genes in *M. aeruginosa* PCC 7806
depends on the ambient pCO_2 (Sandrini et al., 2015). Notably, under high pCO_2 conditions (1450 ppm in
their experiments), the expression of high-affinity bicarbonate uptake systems (notably *cmpA*, *cmpB*, and
cmpC genes) is down-regulated, and cells shift towards CO_2 and low-affinity bicarbonate uptake
compared with cells grown under low pCO_2 conditions (200 ppm) that the authors considered as C_i-
limited. For the production of the transcriptomes that we studied here, pCO_2 was not regulated nor
monitored, and the change of expression in CCM-involved genes such as *cmpD* and *cmpC* could be due to
changes in pCO_2 conditions during day/night transitions, even if there are expected to be lower than in

the Sandrini et al. (2015)'s study. In future studies, exploring the molecular controls of biomineralization,
a special focus on these parameters would be appropriate.
In contrast to the bicarbonate transporters and their homologs, the number of genes possibly linked
with calcium transport (24 genes) and having an expression pattern positively correlated with that of
ccyA (10 genes) was higher than genes with an expression pattern negatively correlated with *ccyA* (5
genes) (Table S4). The transport of Ca^{2+} ions into the cytosol is commonly considered as a passive process
facilitated by channels that exhibit low ionic specificity (Domínguez et al., 2015). Here, only one out of
the ten genes whose expression was positively correlated with that of *ccyA* was identified as a passive
transporter (hB1-1). The other nine genes include the *cax* gene, neighboring *ccyA*, two *apnhaP* genes
(Na^+/H^+ antiporter) and three of their homologs, a UPF0016 gene (putative Ca^{2+}/H^+ transporter) and a
Ca^{2+}-ATPase and one of their homologs. They could thus also be candidate genes contributing to the
intracellular accumulation of calcium with the same question arising about the polarity of the transport
as for *cax* (towards the exterior of the cells or inside compartments containing iACC).
In a comparative genomics approach performed at the scale of the whole Cyanobacteria phylum,
ccyA was the only gene identified as shared by iACC+ strains and absent in iACC- strains (Benzerara et al.,
2022). However, some genes from both phenotypes (iACC- and iACC+), and therefore not highlighted by
the above-mentioned comparative genomics approach, could also be involved in iACC formation. Indeed,
these genes may be recruited in iACC+ strains for iACC formation, whereas they would have a different
function in iACC- strains. Moreover, there could also be genes specific to *Microcystis* involved in iACC
formation, with other non-homologous genes but of similar functions involved in other cyanobacterial
genera. The present study is a step forward towards identifying possible partners in iACC formation.
However, only future studies specifically targeting these genes will be able to provide definitive answers
about their implications in the iACC formation process. Furthermore, investigations into the
comprehensive response of *Microcystis* to fluctuations in calcium and/or pCO_2 levels could provide
valuable insights into the molecular mechanisms underlying the biomineralization of iACC. These studies
should encompass monitoring of the physicochemical parameters of the environment and the
quantification of iACC formation. Notably, certain semi-quantitative techniques, such as Fourier
transform infrared spectroscopy (FTIR) (Mehta et al., 2022b) and X-ray absorption near edge structure
(XANES) spectroscopy (Mehta et al., 2023), now offer the capability to detect cyanobacterial iACC and
assess their relative abundance.
Finally, it can be noted that the correlation between the *ccyA* gene and genes involved in calcium
transport and the CCM is not restricted to *Microcystis*. For example, the combined presence of homologs
of a Ca^{2+}/H^+ antiporter gene and the Na^+-dependent bicarbonate *bicA* gene was shown to be significantly
correlated with that of *ccyA* in the Cyanobacteria phylum (Benzerara et al., 2022). Moreover, the *ccyA*
gene was co-localized with a Ca^{2+}/H^+ antiporter gene in several other cyanobacteria (*C. fritschii* PCC 9212
and PCC 6912, and *Fischerella* sp. NIES-4106), mirroring the situation observed in *M. aeruginosa* PCC
7806. Overall, this offers some target genes for future genetic studies to better depict the molecular
mechanisms of iACC biomineralization.
546

Acknowledgements

We thank Varet H, Sismeiro O, Dillies MA, Legendre R and Coppe JY who run the RNA platform of the
Institut Pasteur and provided us with the processed trancriptomics data.

Data, scripts, code, and supplementary information availability

Raw transcriptomics data are available online on NCBI, Gene expression Omnibus, GEO accession
number: GSE255450. Accession "GSE255450" is currently private but a secure token will be provided to
reviewers. Supplementary information, supplementary data (output data from Salmon, DICOEXPRESS and
DeepNOG analyses as well as gene expression correlation) and supplementary figures and tables are
available online on bioRxiv, BIORXIV/2024/602159 - Version 1, https://doi.org/10.5802/fake2.doi. Links
to the tools used in this study are noted in the Material and methods section.

Conflict of interest disclosure

558   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
559   relation to the content of the article.

Funding

561   This work was supported by the Agence Nationale de la Recherche (ANR Harley, ANR-19-CE44-0017-
562   01; ANR PHOSTORE, ANR-19-CE01-0005). The PCC collection is funded by the Institut Pasteur. The PhD of
563   C.P. was supported by the Ile-de-France ARDoC Grant. Juliette Gaëtan PhD grant was funded by the
564   Learning Planet Institute and University Paris Cité grant Frontiers of Innovation in Research and
565   Education.

References

Raw transcriptomics data are available on NCBI, Gene expression Omnibus, GEO accession number:
GSE255450. https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE255450

Altermann, W., Kazmierczak, J., Oren, A., Wright, D.T., 2006. Cyanobacterial calcification and its
rock-building potential during 3.5 billion years of Earth history. *Geobiology* **4**, 147–166.
https://doi.org/10.1111/j.1472-4669.2006.00076.x

Badger, M.R., Price, G.D., 2003. CO_2 concentrating mechanisms in cyanobacteria: molecular
components, their diversity and evolution. *Journal of Experimental Botany* **54**, 609–622.
https://doi.org/10.1093/jxb/erg076

Benzerara, K., Duprat, E., Bitard-Feildel, T., Caumes, G., Cassier-Chauvat, C., Chauvat, F., Dezi, M.,
Diop, S.I., Gaschignard, G., Görgen, S., Gugger, M., López-García, P., Millet, M., Skouri-
Panet, F., Moreira, D., Callebaut, I., 2022. A new gene family diagnostic for intracellular
biomineralization of amorphous Ca carbonates by cyanobacteria. *Genome Biology and
Evolution* **14**, evac026. https://doi.org/10.1093/gbe/evac026

Benzerara, K., Görgen, S., Athar, K.M., Chauvat, F., March, K., Menguy, N., Mehta, N., Skouri-
Panet, F., Swaraj, S., Travert, C., Cassier-Chauvat, C., Duprat, E., 2023. Quantitative
mapping of calcium cell reservoirs in cyanobacteria at the submicrometer scale. *Journal of
Electron   Spectroscopy   and   Related   Phenomena   **267**,   147369.
https://doi.org/10.1016/j.elspec.2023.147369

Benzerara, K., Skouri-Panet, F., Li, J., Férard, C., Gugger, M., Laurent, T., Couradeau, E., Ragon, M.,
Cosmidis, J., Menguy, N., Margaret-Oliver, I., Tavera, R., López-García, P., Moreira, D.,
2014. Intracellular Ca-carbonate biomineralization is widespread in cyanobacteria.
Proceedings   of   the   National   Academy   of   Sciences   **111**,   10933–10938.
https://doi.org/10.1073/pnas.1403510111

Blondeau, M., Benzerara, K., Ferard, C., Guigner, J.-M., Poinsot, M., Coutaud, M., Tharaud, M.,
Cordier, L., Skouri-Panet, F., 2018a. Impact of the cyanobacterium *Gloeomargarita
lithophora* on the geochemical cycles of Sr and Ba. *Chemical Geology* **483**, 88–97.
https://doi.org/10.1016/j.chemgeo.2018.02.029

Blondeau, M., Sachse, M., Boulogne, C., Gillet, C., Guigner, J.-M., Skouri-Panet, F., Poinsot, M.,
Ferard, C., Miot, J., Benzerara, K., 2018b. Amorphous calcium carbonate granules form
within an intracellular compartment in calcifying cyanobacteria. *Front. Microbiol.* **9**, 1768.
https://doi.org/10.3389/fmicb.2018.01768

Boskey, A.L., 2003. Biomineralization: an overview. *Connective Tissue Research* **44**, 5–9.
https://doi.org/10.1080/03008200390152007

Brownlee, C., Wheeler, G.L., Taylor, A.R., 2015. Coccolithophore biomineralization: New questions,
new   answers.   *Seminars   in   Cell   &   Developmental   Biology*,   **46**,   11–16.
https://doi.org/10.1016/j.semcdb.2015.10.027

Callebaut, I., Labesse, G., Durand, P., Poupon, A., Canard, L., Chomilier, J., Henrissat, B., Mornon,
J.P., 1997. Deciphering protein sequence information through hydrophobic cluster analysis

(HCA): current status and perspectives. *Cell Mol Life Sci*, **53**, 621–645. https://doi.org/10.1007/s000180050082

Cam, N., Benzerara, K., Georgelin, T., Jaber, M., Lambert, J.-F., Poinsot, M., Skouri-Panet, F., Cordier, L., 2016. Selective uptake of alkaline earth metals by cyanobacteria forming intracellular carbonates. *Environ. Sci. Technol.* **50**, 11654–11662. https://doi.org/10.1021/acs.est.6b02872

Cam, N., Benzerara, K., Georgelin, T., Jaber, M., Lambert, J.-F., Poinsot, M., Skouri-Panet, F., Moreira, D., Lopez-Garcia, P., Raimbault, E., Cordier, L., Jezequel, D., 2018. Cyanobacterial formation of intracellular Ca-carbonates in undersaturated solutions. *Geobiology* **16**, 49–61. https://doi.org/10.1111/gbi.12261

Cam, N., Georgelin, T., Jaber, M., Lambert, J.-F., Benzerara, K., 2015. In vitro synthesis of amorphous Mg-, Ca-, Sr- and Ba-carbonates: What do we learn about intracellular calcification by cyanobacteria? *Geochimica et Cosmochimica Acta* **161**, 36–49. https://doi.org/10.1016/j.gca.2015.04.003

Chen, S., Gagnon, A.C., Adkins, J.F., 2018. Carbonic anhydrase, coral calcification and a new model of stable isotope vital effects. *Geochimica et Cosmochimica Acta* **236**, 179–197. https://doi.org/10.1016/j.gca.2018.02.032

Christiano, R., Nagaraj, N., Fröhlich, F., Walther, T.C., 2014. Global proteome turnover analyses of the yeasts *S. cerevisiae* and *S. pombe*. *Cell Reports* **9**, 1959–1965. https://doi.org/10.1016/j.celrep.2014.10.065

Cosmidis, J., Benzerara, K., 2022. Why do microbes make minerals? *Comptes Rendus. Géoscience* **354**, 1–39. https://doi.org/10.5802/crgeos.107

Couradeau, E., Benzerara, K., Gerard, E., Moreira, D., Bernard, S., Brown, G.E., Lopez-Garcia, P., 2012. An early-branching microbialite cyanobacterium forms intracellular carbonates. *Science* **336**, 459–462. https://doi.org/10.1126/science.1216171

Cumsille, A., Durán, R.E., Rodríguez-Delherbe, A., Saona-Urmeneta, V., Cámara, B., Seeger, M., Araya, M., Jara, N., Buil-Aranda, C., 2023. GenoVi, an open-source automated circular genome visualizer for bacteria and archaea. *PLOS Computational Biology* **19**, e1010998. https://doi.org/10.1371/journal.pcbi.1010998

De Wever, A., Benzerara, K., Coutaud, M., Caumes, G., Poinsot, M., Skouri☐Panet, F., Laurent, T., Duprat, E., Gugger, M., 2019. Evidence of high Ca uptake by cyanobacteria forming intracellular $CaCO_3$ and impact on their growth. *Geobiology* **17**, 676–690. https://doi.org/10.1111/gbi.12358

Dick, G.J., Duhaime, M.B., Evans, J.T., Errera, R.M., Godwin, C.M., Kharbush, J.J., Nitschky, H.S., Powers, M.A., Vanderploeg, H.A., Schmidt, K.C., Smith, D.J., Yancey, C.E., Zwiers, C.C., Denef, V.J., 2021. The genetic and ecophysiological diversity of *Microcystis*. *Environmental Microbiology* **23**, 7278–7313. https://doi.org/10.1111/1462-2920.15615

Domínguez, D.C., Guragain, M., Patrauchan, M., 2015. Calcium binding proteins and calcium signaling in prokaryotes. *Cell Calcium, Evolution of Calcium Signaling* **57**, 151–165. https://doi.org/10.1016/j.ceca.2014.12.006

Feldbauer, R., Gosch, L., Lüftinger, L., Hyden, P., Flexer, A., Rattei, T., 2021. DeepNOG: fast and accurate protein orthologous group assignment. *Bioinformatics* **36**, 5304–5312. https://doi.org/10.1093/bioinformatics/btaa1051

Gaëtan, J., Halary, S., Millet, M., Bernard, C., Duval, C., Hamlaoui, S., Hecquet, A., Gugger, M., Marie, B., Mehta, N., Moreira, D., Skouri-Panet, F., Travert, C., Duprat, E., Leloup, J., Benzerara, K., 2023. Widespread formation of intracellular calcium carbonates by the bloom-forming cyanobacterium *Microcystis*. *Environmental Microbiology* **25**, 751–765. https://doi.org/10.1111/1462-2920.16322

Gancedo, J.M., López, S., Ballesteros, F., 1982. Calculation of half-lives of proteins in vivo. Heterogeneity in the rate of degradation of yeast proteins. *Mol Cell Biochem* **43**, 89–95. https://doi.org/10.1007/BF00423096

Gaschignard, G., Millet, M., Bruley, A., Benzerara, K., Dezi, M., Skouri-Panet, F., Duprat, E., Callebaut, I., 2024. AlphaFold2-guided description of CoBaHMA, a novel family of bacterial domains within the heavy-metal-associated superfamily. *Proteins*, 1-19. https://doi.org/10.1002/prot.26668

Gilbert, P.U.P.A., Bergmann, K.D., Boekelheide, N., Tambutté, S., Mass, T., Marin, F., Adkins, J.F., Erez, J., Gilbert, B., Knutson, V., Cantine, M., Hernández, J.O., Knoll, A.H., 2022. Biomineralization: Integrating mechanism and evolutionary history. *Science Advances* **8**, eabl9653. https://doi.org/10.1126/sciadv.abl9653

Görgen, S., Benzerara, K., Skouri-Panet, F., Gugger, M., Chauvat, F., Cassier-Chauvat, C., 2020. The diversity of molecular mechanisms of carbonate biomineralization by bacteria. *Discov Mater* **1**, 2. https://doi.org/10.1007/s43939-020-00001-9

Harke, M.J., Steffen, M.M., Gobler, C.J., Otten, T.G., Wilhelm, S.W., Wood, S.A., Paerl, H.W., 2016. A review of the global ecology, genomics, and biogeography of the toxic cyanobacterium, *Microcystis* spp. *Harmful Algae* **54**, 4–20. https://doi.org/10.1016/j.hal.2015.12.007

Hirschi, K.D., Zhen, R.G., Cunningham, K.W., Rea, P.A., Fink, G.R., 1996. CAX1, an H^+/Ca^{2+} antiporter from *Arabidopsis*. *Proceedings of the National Academy of Sciences* **93**, 8782–8786. https://doi.org/10.1073/pnas.93.16.8782

Ingolia, N.T., Ghaemmaghami, S., Newman, J.R.S., Weissman, J.S., 2009. Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science* **324**, 218–223. https://doi.org/10.1126/science.1168978

Knoll, A.H., 2003. Biomineralization and evolutionary history. *Reviews in Mineralogy and Geochemistry* **54**, 329–356. https://doi.org/10.2113/0540329

Lambert, I., Paysant-Le Roux, C., Colella, S., Martin-Magniette, M.-L., 2020. DiCoExpress: a tool to process multifactorial RNAseq experiments from quality controls to co-expression analysis through differential analysis based on contrasts inside GLM models. *Plant Methods* **16**, 68. https://doi.org/10.1186/s13007-020-00611-7

Le Roy, N., Jackson, D.J., Marie, B., Ramos-Silva, P., Marin, F., 2014. The evolution of metazoan α-carbonic anhydrases and their roles in calcium carbonate biomineralization. *Frontiers in Zoology* **11**, 75. https://doi.org/10.1186/s12983-014-0075-8

Lê, S., Josse, J., Husson, F., 2008. FactoMineR: an R package for multivariate analysis. *Journal of Statistical Software* **25**, 1–18. https://doi.org/10.18637/jss.v025.i01

Li, J., Margaret Oliver, I., Cam, N., Boudier, T., Blondeau, M., Leroy, E., Cosmidis, J., Skouri-Panet, F., Guigner, J.-M., Férard, C., Poinsot, M., Moreira, D., Lopez-Garcia, P., Cassier-Chauvat, C., Chauvat, F., Benzerara, K., 2016. Biomineralization patterns of intracellular carbonatogenesis in cyanobacteria: molecular hypotheses. *Minerals* **6**, 10. https://doi.org/10.3390/min6010010

Liu, P., Zheng, Y., Zhang, R., Bai, J., Zhu, K., Benzerara, K., Menguy, N., Zhao, X., Roberts, A.P., Pan, Y., Li, J., 2023. Key gene networks that control magnetosome biomineralization in magnetotactic bacteria. *National Science Review* **10**, nwac238. https://doi.org/10.1093/nsr/nwac238

Mackinder, L., Wheeler, G., Schroeder, D., von Dassow, P., Riebesell, U., Brownlee, C., 2011. Expression of biomineralization-related ion transport genes in *Emiliania huxleyi*. *Environmental Microbiology* **13**, 3250–3265. https://doi.org/10.1111/j.1462-2920.2011.02561.x

Mantione, K.J., Kream, R.M., Kuzelova, H., Ptacek, R., Raboch, J., Samuel, J.M., Stefano, G.B., 2014. Comparing bioinformatic gene expression profiling methods: microarray and RNA-Seq. *Med Sci Monit Basic Res* **20**, 138–141. https://doi.org/10.12659/MSMBR.892101

Marin, F., Corstjens, P., de Gaulejac, B., de Vrind-De Jong, E., Westbroek, P., 2000. Mucins and molluscan calcification: molecular characterization of mucoperlin, a novel mucin-like protein from the nacreous shell layer of the fan mussel *Pinna nobilis* (Bivalvia, pteriomorphia). *Journal of Biological Chemistry* **275**, 20667–20675. https://doi.org/10.1074/jbc.M003006200

Marin, F., Luquet, G., 2004. Molluscan shell proteins. *Comptes Rendus Palevol* **3**, 469–492. https://doi.org/10.1016/j.crpv.2004.07.009

Marin, F., Smith, M., Isa, Y., Muyzer, G., Westbroek, P., 1996. Skeletal matrices, muci, and the origin of invertebrate calcification. *Proceedings of the National Academy of Sciences* **93**, 1554–1559. https://doi.org/10.1073/pnas.93.4.1554

Mehta, N., Bougoure, J., Kocar, B.D., Duprat, E., Benzerara, K., 2022a. Cyanobacteria accumulate radium (^{226}Ra) within intracellular amorphous calcium carbonate inclusions. *ACS EST Water* **2**, 616–623. https://doi.org/10.1021/acsestwater.1c00473

Mehta, N., Gaëtan, J., Giura, P., Azaïs, T., Benzerara, K., 2022b. Detection of biogenic amorphous calcium carbonate (ACC) formed by bacteria using FTIR spectroscopy. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* **278**, 121262. https://doi.org/10.1016/j.saa.2022.121262

Mehta, N., Vantelon, D., Gaëtan, J., Fernandez-Martinez, A., Delbes, L., Travert, C., Benzerara, K., 2023. Calcium speciation and coordination environment in intracellular amorphous calcium carbonate (ACC) formed by cyanobacteria. *Chemical Geology* **641**, 121765. https://doi.org/10.1016/j.chemgeo.2023.121765

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., Steinegger, M., 2022. ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679–682. https://doi.org/10.1038/s41592-022-01488-1

Monteil, C.L., Benzerara, K., Menguy, N., Bidaud, C.C., Michot-Achdjian, E., Bolzoni, R., Mathon, F.P., Coutaud, M., Alonso, B., Garau, C., Jézéquel, D., Viollier, E., Ginet, N., Floriani, M., Swaraj, S., Sachse, M., Busigny, V., Duprat, E., Guyot, F., Lefevre, C.T., 2021. Intracellular amorphous Ca-carbonate and magnetite biomineralization by a magnetotactic bacterium affiliated to the Alphaproteobacteria. *ISME J* **15**, 1–18. https://doi.org/10.1038/s41396-020-00747-3

Monteil, C.L., Perrière, G., Menguy, N., Ginet, N., Alonso, B., Waisbord, N., Cruveiller, S., Pignol, D., Lefèvre, C.T., 2018. Genomic study of a novel magnetotactic Alphaproteobacteria uncovers the multiple ancestry of magnetotaxis. *Environmental Microbiology* **20**, 4415–4430. https://doi.org/10.1111/1462-2920.14364

Monteiro, F.M., Bach, L.T., Brownlee, C., Bown, P., Rickaby, R.E.M., Poulton, A.J., Tyrrell, T., Beaufort, L., Dutkiewicz, S., Gibbs, S., Gutowska, M.A., Lee, R., Riebesell, U., Young, J., Ridgwell, A., 2016. Why marine phytoplankton calcify. *Science Advances* **2**, e1501822. https://doi.org/10.1126/sciadv.1501822

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., Kingsford, C., 2017. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417–419. https://doi.org/10.1038/nmeth.4197

Pattanayak, G., Rust, M.J., 2014. The cyanobacterial clock and metabolism. *Current Opinion in Microbiology* **18**, 90–95. https://doi.org/10.1016/j.mib.2014.02.010

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B.L., Salazar, G.A., Bileschi, M.L., Bork, P., Bridge, A., Colwell, L., Gough, J., Haft, D.H., Letunić, I., Marchler-Bauer, A., Mi, H., Natale, D.A., Orengo, C.A., Pandurangan, A.P., Rivoire, C., Sigrist, C.J.A., Sillitoe, I., Thanki, N., Thomas, P.D., Tosatto, S.C.E., Wu, C.H., Bateman, A., 2023. InterPro in 2022. *Nucleic Acids Research* **51**, D418–D427. https://doi.org/10.1093/nar/gkac993

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., Ferrin, T.E., 2004. UCSF Chimera-a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605–1612. https://doi.org/10.1002/jcc.20084

Price, G.D., Howitt, S.M., 2011. The cyanobacterial bicarbonate transporter *BicA*: its physiological role and the implications of structural similarities with human SLC26 transporters. *Biochem. Cell Biol.* **89**, 178–188. https://doi.org/10.1139/O10-136

Ragon, M., Benzerara, K., Moreira, D., Tavera, R., Lopez-Garcia, P., 2014. 16S rDNA-based analysis reveals cosmopolitan occurrence but limited diversity of two cyanobacterial lineages with contrasted patterns of intracellular carbonate mineralization. *Front. Microbiol.* **5,** 331. https://doi.org/10.3389/fmicb.2014.00331

Riba, A., Di Nanni, N., Mittal, N., Arhné, E., Schmidt, A., Zavolan, M., 2019. Protein synthesis rates and ribosome occupancies reveal determinants of translation elongation rates. *Proceedings of the National Academy of Sciences* **116**, 15023–15032. https://doi.org/10.1073/pnas.1817299116

Rippka, R., Deruelles, J., Waterbury, J.B., Herdman, M., Stanier, R.Y., 1979. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *Microbiology* **111**, 1–61. https://doi.org/10.1099/00221287-111-1-1

Robinson, M.D., Oshlack, A., 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology* **11**, R25. https://doi.org/10.1186/gb-2010-11-3-r25

Sandrini, G., Cunsolo, S., Schuurmans, J.M., Matthijs, H.C.P., Huisman, J., 2015. Changes in gene expression, cell physiology and toxicity of the harmful cyanobacterium *Microcystis aeruginosa* at elevated CO_2. *Frontiers in Microbiology* **6**, 401. https://doi.org/10.3389/fmicb.2015.00401

Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., Selbach, M., 2011. Global quantification of mammalian gene expression control. *Nature* **473**, 337–342. https://doi.org/10.1038/nature10098

Shigaki, T., Rees, I., Nakhleh, L., Hirschi, K.D., 2006. Identification of three distinct phylogenetic groups of CAX cation/proton antiporters. *J Mol Evol* **63**, 815–825. https://doi.org/10.1007/s00239-006-0048-4

Stöckel, J., Welsh, E.A., Liberton, M., Kunnvakkam, R., Aurora, R., Pakrasi, H.B., 2008. Global transcriptomic analysis of *Cyanothece* 51142 reveals robust diurnal oscillation of central metabolic processes. *Proceedings of the National Academy of Sciences* **105**, 6156–6161. https://doi.org/10.1073/pnas.0711068105

Straub, C., Quillardet, P., Vergalli, J., Marsac, N.T. de, Humbert, J.-F., 2011. A day in the life of *Microcystis aeruginosa* strain PCC 7806 as revealed by a transcriptomic analysis. *PLOS ONE* **6**, e16208. https://doi.org/10.1371/journal.pone.0016208

Tang, J., Zhou, H., Yao, D., Riaz, S., You, D., Klepacz-Smółka, A., Daroch, M., 2022. Comparative genomic analysis revealed distinct molecular components and organization of CO_2-concentrating mechanism in thermophilic cyanobacteria. *Frontiers in Microbiology* **13**, 876272. https://doi.org/10.3389/fmicb.2022.876272

Taoka, A., Eguchi, Y., Shimoshige, R., Fukumori, Y., 2023. Recent advances in studies on magnetosome-associated proteins composing the bacterial geomagnetic sensor organelle. *Microbiology and Immunology* **67**, 228–238. https://doi.org/10.1111/1348-0421.13062

Uebe, R., Schüler, D., 2016. Magnetosome biogenesis in magnetotactic bacteria. *Nat Rev Microbiol* **14**, 621–637. https://doi.org/10.1038/nrmicro.2016.99

van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C.L.M., Söding, J., Steinegger, M., 2023. Fast and accurate protein structure search with Foldseek. *Nat Biotechnol* 1–4. https://doi.org/10.1038/s41587-023-01773-0

Waditee, R., Hossain, G.S., Tanaka, Y., Nakamura, T., Shikata, M., Takano, J., Takabe, Tetsuko, Takabe, Teruhiro, 2004. Isolation and functional characterization of Ca^{2+}/H^+ antiporters from Cyanobacteria. *J. Biol. Chem.* **279**, 4330–4338. https://doi.org/10.1074/jbc.M310282200

Waight, A.B., Pedersen, B.P., Schlessinger, A., Bonomi, M., Chau, B.H., Roe-Zurz, Z., Risenmay, A.J., Sali, A., Stroud, R.M., 2013. Structural basis for alternating access of a eukaryotic calcium/proton exchanger. *Nature* **499**, 107–110. https://doi.org/10.1038/nature12233

Wang, Z., Gerstein, M., Snyder, M., 2009. RNA-Seq: a revolutionary tool for transcriptomics. *Nat Rev Genet* **10**, 57–63. https://doi.org/10.1038/nrg2484

Welkie, D.G., Rubin, B.E., Diamond, S., Hood, R.D., Savage, D.F., Golden, S.S., 2019. A Hard Day's Night: cyanobacteria in diel cycles. *Trends in Microbiology* **27**, 231–242. https://doi.org/10.1016/j.tim.2018.11.002

Zhao, S., Fung-Leung, W.-P., Bittner, A., Ngo, K., Liu, X., 2014. Comparison of RNA-Seq and microarray in transcriptome profiling of activated T cells. *PLOS ONE* **9**, e78644. https://doi.org/10.1371/journal.pone.0078644

Zinser, E.R., Lindell, D., Johnson, Z.I., Futschik, M.E., Steglich, C., Coleman, M.L., Wright, M.A., Rector, T., Steen, R., McNulty, N., Thompson, L.R., Chisholm, S.W., 2009. Choreography of the transcriptome, photophysiology, and cell cycle of a minimal photoautotroph, *Prochlorococcus*. *PLOS ONE* **4**, e5135. https://doi.org/10.1371/journal.pone.0005135