Genomics of Neotropical biodiversity indicators: two butterfly radiations with rampant chromosomal rearrangements and hybridisation

Eva SM van der Heijden[1,2], Karin Näsvall[1], Fernando A. Seixas[3], Carlos Eduardo Beserra Nobre[4], Artur Campos D Maia[4,5], Patricio Salazar-Carrión[1,2], Jonah M Walker[1,2], Daiane Szczerbowski[6], Stefan Schulz[6], Ian A Warren[2], Kimberly Gabriela Gavilanes Córdova[7], María José Sánchez-Carvajal[7], Franz Chandi[7], Alex P Arias-Cruz[7], Nicol Rueda-M[1,8], Camilo Salazar[8], Kanchon K Dasmahapatra[9], Stephen H Montgomery[10], Melanie McClure[11], Dominic E Absolon[1], Thomas C Mathers[1], Camilla A Santos[1], Shane McCarthy[1], Jonathan MD Wood[1], Gerardo Lamas[12], Caroline Bacquet[7,13], André Victor Lucci Freitas[14], Keith R. Willmott[15], Chris D Jiggins[2], Marianne Elias[13,16], Joana I Meier[1,2]

1. Tree of Life Programme, Wellcome Sanger Institute, UK
2. Department of Zoology, University of Cambridge, UK
3. Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA
4. Graduate Program in Animal Biology, Federal University of Pernambuco, Brazil
5. Graduate Program in Plant Biology, Federal University of Pernambuco, Brazil
6. Institute of Organic Chemistry, Technische Universität Braunschweig, Germany
7. Universidad Regional Amazónica Ikiam, Ecuador
8. Biology Program, Faculty of Natural Sciences, Universidad del Rosario, Bogotá, Colombia
9. Department of Biology, University of York, UK
10. University of Bristol, UK
11. Laboratoire Écologie, Évolution, Interactions des Systèmes Amazoniens (LEEISA), Université de Guyane, CNRS, IFREMER, Cayenne, France
12. Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima, Peru
13. Smithsonian Tropical Research Institute, Gamboa, Panama
14. Universidade Estadual de Campinas, Brazil
15. Florida Museum of Natural History, USA
16. Institut de Systématique Évolution Biodiversité (ISYEB), CNRS, MNHN, EPHE, Sorbonne Université, Université des Antilles, Paris, France

Author contributions:

PSC, JIM, ESMH, KGGC, MJSC, DS, FC, APAC, CB, NRM, CS, KKD, NN, SHM, MM, KRW, AVLF, CDJ and ME collected samples. GL, KRW and ME oversaw the butterfly identification and taxonomic revision. ESMH, JIM, JMW and IAW performed lab work. ESMH mapped and filtered the sequencing data, and ran the phylogenetic and hybridisation analyses. KN did the chromosomal rearrangement analysis. CEBN, ACDM, DS, SS and AVLF analysed the androconial semiochemicals. FAS ran the MSCi analysis. DEA, TCM, CAS, SM, KN and JW assembled and curated the new genomes. JIM designed the study with contributions from ESMH, ME and CDJ. ESMH, JIM and KN wrote the manuscript with contributions from all authors.

Corresponding author: Joana Isabel Meier, joana.meier@sanger.ac.uk

Key words: Speciation, biodiversity, butterflies, neotropics, genomics

Classification: Biological Sciences - Evolution

Abstract

A major question in evolutionary biology is what drives the diversification of lineages. Rapid, recent
radiations are ideal systems for addressing how new species arise because they still show key
morphological and ecological adaptations associated with speciation. While most studied recent
radiations have evolved in an insular environment, less research has been carried out on continental
radiations with complex species interactions. *Melinaea* and *Mechanitis* butterflies (Nymphalidae:
Ithomiini) have rapidly radiated in the Neotropics. They are classical models for Amazonian biogeography
and colour pattern mimicry and have been proposed as biodiversity indicators. We generated reference
genomes for five species of each genus, and whole-genome resequencing data of most species and
subspecies covering a wide geographic range to assess phylogeographic relationships, patterns of
hybridisation and chromosomal rearrangements. Our data help resolve the classification of these
taxonomically challenging butterflies and reveal very high diversification rates. We find rampant evidence
of historical hybridisation and putative hybrid species in both radiations, which may have facilitated their
rapid diversification. Moreover, dozens of chromosomal fusions and fissions were identified between
congeneric species, and even some within species. We conclude that interactions between geography,
hybridisation and chromosomal rearrangements have contributed to these two rapid radiations in the
highly diverse Neotropical region. We suggest that rapid radiations may be spurred by repeated periods
of geographic isolation during Pleistocene climate oscillations, combined with lineage-specific rapid
accumulation of incompatibilities during allopatric phases, followed by secondary contact with some gene
exchange.

Significance Statement

Understanding factors contributing to rapid speciation is a key aim of evolutionary biology. Here we focus
on two rapid radiations of Neotropical butterflies. Our genomic data with broad taxonomic and
geographic coverage reveal rampant hybridisation and chromosomal rearrangements, each likely
contributing to the high diversification rates. Our study highlights the use of genomic data to resolve
taxonomically challenging species groups and elucidate drivers of diversification in rapid radiations. We
show that for biodiversity hotspots with recent radiations, barcoding is insufficient to characterise species
richness due to gene flow and recent speciation. The taxonomic implications of both introgression and
karyotype diversity for species delimitation are important to consider during monitoring and management
of biodiversity in these vulnerable habitats.

Introduction

Rapid radiations, where a lineage diversifies into many different species over a short time period, are ideal systems for studying how new species evolve (1, 2). They can be driven both by non-adaptive processes, such as the accumulation of differences during periods of allopatry leading to incompatibilities upon secondary contact, and by adaptive processes such as adaptation to different ecological niches or sexual selection for different traits and preferences (3, 4). Sympatric radiations require some degree of niche differentiation among the species for stable coexistence and sufficient reproductive isolation such that incipient lineages do not merge (2).

While most lineages do not readily radiate even in the face of ecological opportunity, some lineages are particularly prone to rapid radiations, and do so repeatedly. We are only starting to understand the factors explaining these lineage-specific differences (5). Most knowledge stems from well-studied radiations that evolved in insular environments with little competition with other species and a relatively simple and geographically limited environment (e.g. Darwin's finches on the Galapagos Islands, cichlid fishes in lakes, *Anolis* lizards on the Caribbean islands or Hawaiian silverswords) (2, 6). However, many rapid radiations evolved on the much more complex continents and much less is known about drivers leading to their diversification (7–9). Reduced competition in insular environments allows niche specialisation without being immediately outcompeted when a lineage is not yet well-adapted to its environment. However, in large continental areas such as the hyperdiverse Neotropics, competition is much stronger limiting ecological speciation. On the other hand, large and complex environments on continents may provide more opportunity for allopatric divergence than a small island. The speed of accumulating incompatibilities in allopatry may thus be more important in rapid radiations on continents than insular environments.

Here, we study the drivers of diversification in two rapid continental radiations of the Neotropical butterfly tribe Ithomiini. Ithomiine butterflies (Nymphalidae: Danainae, ca. 400 species in 42 genera) are found across Central and South America (10, 11). They constitute a substantial part of the butterfly species assemblage and are regarded as good indicators of spatial patterns of biodiversity in the Neotropics, the most biodiverse areas in the world (10, 12, 13). Sequestration of pyrrolizidine alkaloids from Asteraceae and Boraginaceae plants render most Ithomiini unpalatable (14–18), and their colour patterns advertise this unpalatability to predators. They form Müllerian mimicry rings, where locally co-occurring species have converged in colour pattern, and thus share the cost of educating predators (10, 19, 20). We focus on two ithomiine genera, *Melinaea* and *Mechanitis*, which have diversified exceptionally fast with most species younger than a million years (11, 21). Hitherto, the study of these radiations has been hampered

by taxonomic challenges. *Melinaea* and *Mechanitis* are among the most taxonomically difficult of
Ithomiini, as Fox noted (1967) (22): 'these insects [are] so thoroughly confusing and so thoroughly
confused by my predecessors'. The species do not differ in genital or other morphological characteristics
and show substantial intraspecific wing pattern variation and mimicry between taxa. Barcoding does not
reliably distinguish species either (23, 24). As prior studies have only used few or no genetic markers or
did not have broad geographic coverage, the taxonomy is still partially unresolved, despite many
taxonomic revisions (22, 24–26, e.g. 27–29).

While the exact causes of their rapid radiations are unknown, different contributing factors have
been proposed. Ecological adaptation may be relevant as species show differences in microhabitats, host
plants, mimicry rings, and altitude, but on the other hand many species share habitats and host plants,
and the majority of species occurs in the lowlands (25, 30–33). Moreover, where species co-exist, they
converge in colour patterns and thus assortative mating may rely more strongly on chemical cues.
Ithomiine species differ in male-specific androconial compounds (chemical compounds secreted from
specialised wing scales where the fore- and hindwing overlap), which likely act as pheromones (Trigo et
al. 1996; Schulz et al. 2004; Blow et al. 2023).

Allopatric accumulation of differences could also have played a role in the rapid diversification of
ithomiini, as this could have occurred in different rainforest refugia during climatic oscillations e.g. in the
Pleistocene (26, 34, but see 35, 36) or on opposite sides of geographic barriers such as the Andes. Both
climatic refugia and the Andes have been proposed as "speciation pumps" in the Neotropics (37), also for
Ithomiini (38), where periods of allopatry followed by secondary contact create favourable circumstances
for speciation.

Another factor that might contribute to the diversification of ithomiine butterflies is hybridisation.
Phylogenetic studies using a limited number of markers have revealed mito-nuclear discordances and
paraphyletic taxa in Ithomiini (23, 36). This could be due to limited geographic or genetic resolution,
incomplete lineage sorting in the rapidly speciating lineages, or introgressive hybridisation. While gene
flow between sister lineages can homogenise their gene pools, opposing speciation, recent studies have
shown that sometimes introgressive hybridisation from more distant relatives can facilitate rapid
diversification by enriching the genetic diversity with novel, potentially adaptive variants or contributing
to the origin of new hybrid species (39–43). Admixture has been shown to kickstart adaptive radiation
(e.g. 44, 45), facilitate parallel adaptation (e.g. 42, 46), and novel adaptations (e.g. 47), but the role in
ithomiini diversification is hitherto unknown.

Ithomiini butterflies have an unusually high diversity in chromosome number (48), which could

also contribute to their rapid diversification. Offspring from parents with different karyotypes may suffer

reduced fitness, due to mismatch in pairing of homologous chromosomes that results in aneuploidy,

meiotic failure or hybrid sterility (49, 50). Furthermore, chromosomal rearrangements might facilitate

divergence in the face of gene flow by accumulation of incompatibilities in low recombining regions (51)

or if they link together co-adapted variants (52, 53). In *Melinaea* and *Mechanitis* butterflies, chromosome

counts range from 13 to 30 (48), and chromosomal rearrangements likely contribute to reproductive

isolation, as a cross between two closely-related *Melinaea* species with different karyotypes resulted in

nearly sterile hybrids (54). However, pervasive intraspecific variation in chromosome counts (48, 54)

indicates that not all rearrangements reduce fitness and their role in speciation thus remains an open

question.

Here, we use five reference genomes of each genus and whole-genome resequencing data of

almost all species and many subspecies, to resolve taxonomic uncertainties and explore whether

geography, introgressive hybridisation or chromosomal rearrangements may have played a role in the

rapid diversification of *Mechanitis* and *Melinaea* butterflies.

Results

Taxonomic revision

In this manuscript, we adhere to the 'genotypic cluster' species concept, which defines a species as 'a

morphologically or genetically distinguishable group of individuals that has few or no intermediates when

in contact with other such clusters' (55). Through whole-genome resequencing of 135 *Mechanitis* and 109

Melinaea individuals from across South and Central America, with additional individuals from the

outgroup genera *Forbestra*, *Eutresis* and *Olyras*, we shed light on the phylogenomic relationships and

taxonomy (Fig. 1). Our results, laid out in the following sections (see also Text S1), confirm species versus

subspecies status for most known taxa in agreement with (29), support two recent species

reclassifications (*Mel. tarapotensis* (54), *Mel. mothone* (56)) and reveal three additional taxa that need to

be elevated to species level (*Mel. maeonis* **stat rest** (Hewitson 1869), *Mec. nesaea* **stat rest** (Hübner 1820),

Mec. macrinus **stat rest** (Hewitson 1860), Fig. 1; Text S1). The placement of *Mel. menophilus mediatrix*

(French Guiana) has been uncertain (56); our dataset places it as the most divergent subspecies of *Mel.*

menophilus. A revised, annotated taxonomic list for these two genera can be found in Text S2.

Mitonuclear discordance

In both *Mechanitis* and *Melinaea*, we find rampant mitonuclear discordance, *i.e.* mismatches between

mitochondrial and nuclear phylogenies (IQtree2; maximum likelihood) (Fig. 1). For instance, *Mec. nesaea*

is sister to *Mec. polymnia* in the nuclear phylogeny, but to *Mec. lysimnia* in the mitochondrial phylogeny

(Fig. 1A - box III; Fig. S1-2). This result could either be indicative of incomplete lineage sorting (ILS) or

admixture (more details below). *Mec. messenoides* harbours two divergent mitochondrial haplotypes,

consistent with barcoding results (23, 25) (Fig. 1A - box II), one clustering with the *polymnia-lysimnia* clade,

and the other one with its nuclear sister species, *Mec. menapis*. The mitochondrially divergent *Mec.*

messenoides individuals do not form separate clades in the nuclear phylogeny, nor do they differ in

collection location or subspecies (Table S1; Fig. S1-2).

While the *Melinaea* species are clearly differentiated in the nuclear phylogeny, the mitochondrial

phylogeny shows almost no variation among species, confirming previous barcoding results (56) (Fig. 1B;

Fig. S3-4). *Mel. ludovica*, *Mel. tarapotensis* and *Mel. lilis* are the only species that form monophyletic

mitochondrial clades that are no part of the shallow clade without species differentiation. Notably, *Mel.*

ludovica is the only species that is also an outgroup to the other *Melinaea* species in the nuclear genome,

whereas *Mel. lilis* and *Mel. tarapotensis* are sister to *Mel. isocomma* and *Mel. satevis*, respectively.

Calibrated phylogenetic tree

We approximated the divergence times in the two genera using a Bayesian multispecies coalescent

method for inferring species trees (StarBEAST2), with one individual representing each lineage to produce

a phylogeny calibrated with divergence times from (11) (Fig. 2). The seven *Mechanitis* species are

estimated to have diversified within the past 1.36 million years, resulting in a speciation rate of 1.431

speciation events per lineage/My (assuming a pure birth model with a constant speciation rate). The ten

species of the core *Melinaea* clade (excluding the most divergent species, *Mel. ludovica*) have diversified

in the past 1.41 million years (Fig. 2B), giving a speciation rate of at least 1.633. Of the four potential

Melinaea species missing in our analysis (*Mel. ethra*, *mnasias, mnemopsis* and *scylax*), two likely form part

of the core clade (11, 57), which would increase the speciation rate to 1.672.

However, note that gene flow could affect our estimates, as suggested by discordance between

the StarBEAST2 and IQtree2 topologies (Fig. 2 vs Fig. 1 (note low concordance factors in Fig. 1)): *Mec.*

messenoides is not sister to *Mec. menapis* anymore, and *Mec. nesaea* has shifted to be sister to *Mec.*

lysimnia instead of *Mec. polymnia*. The relationships among lineages within *Mec. polymnia* also changed.

While ILS could partially explain this, introgression could also cause it (see details below). Gene flow

between sister lineages will make their apparent divergence times shorter than the initial split time, and

gene flow with non-sister lineages will make them appear longer.

Phylogeographic patterns

Through combining our sampling locations with those compiled by (10) adjusted according to our

taxonomic revision (Fig. 2), we assessed the biogeographic distribution of the species. We find four main

biogeographic regions (Fig. 2; Fig. S6-7), with most species restricted to one of them. Some sister-species

are separated by the Andes: *Mec. messenoides* and *mazaeus* (Western Amazonia) versus their respective

sisters *Mec. menapis* and *macrinus* (West of the Andes). However, note that the placement of *Mec.*

messenoides differs in the StarBEAST2-topology and is affected by hybridisation (see next section). Within

Mec. polymnia, Ecuadorian and Colombian individuals from opposite sides of the Andes are highly

divergent, indicating little gene flow, though our study includes one putative hybrid (Fig. S8). This pattern

of species separation by the Andes was not previously as apparent due to species misclassification.

Most individuals in our *Melinaea* dataset are from Western Amazonia (east of the Andes) and we

find that many sister species are sympatric. Only two species in our dataset occur west of the Andes (*Mel.*

lilis and *idae*) and they form a clade with a third species from east of the Andes (*Mel. isocomma*). Our

dataset lacks a potential third species occurring west of the Andes, *Mel. scylax* (10), which may represent

a subspecies of *Mel. lilis* (26, 57). Other potential species missing in our dataset are *Mel. mnemopsis* from

Western Amazonia, *Mel. ethra* from the Atlantic forest and *Mel. mnasias* from Western Amazonia and the

Atlantic Forest (10, 57).

Signatures of rampant introgression throughout both genera

Phylogenetic discordance between phylogenies constructed for the nuclear and mitochondrial genome

(Fig. 1), for different genomic regions inferred with concordance factors (Fig. 1), and with IQtree2 and

StarBEAST2 (Fig. 1 vs Fig. 2) is indicative of a history of introgression and/or ILS in both genera. We

assessed hybridisation throughout both radiations with windowed species tree inference with BPP (Fig.

S5), excess allele sharing between non-sister taxa estimated from Fbranch (Fig. S9) and joint inference of

species tree with gene flow using the approximate isolation with migration (AIM) model (Fig. S10). The

resulting phylogenies and introgression histories are summarised in Fig. 2 (Text S3), but we stress that this

is only one of multiple possible scenarios consistent with our observed patterns of excess allele sharing

and gene tree discordance.

In *Mechanitis*, two lineages show uncertain placements. As mentioned, the placement of *Mec.*
messenoides and *nesaea* varies depending on the methodology (IQtree2 and StarBEAST2) and genomic
regions (Fig. 1A – box II and III). Contrary to the nuclear phylogeny inferred with StarBEAST2, but in line
with the mitochondrial phylogeny, the species phylogeny inferred allowing for gene flow (AIM) places
messenoides closer to the *polymnia-nesaea-lysimnia* clade rather than with *menapis-mazaeus-macrinus*.
AIM indicates admixture between these clades (Fig S10A – arrows B-D), and Fbranch also confirms excess
allele sharing between *messenoides* and *polymnia-nesaea-lysimnia* (Fig. S9A - #1). According to the BPP-
analysis, *Mec. messenoides* groups with menapis-mazaeus-macrinus in ~51% of the genome, and with
polymnia-lysimnia-nesaea in ~28% of the genome (sometimes still with *Mec. menapis*) (Fig. 5B). The
branching order within the *polymnia-lysimnia-nesaea* clade is also highly variable across the genome:
polymnia and nesaea are sister in 46.3% of the genome grouping, while 31.7% of trees group *polymnia*
with *lysimnia*, and 14.7% group *nesaea* with *lysimnia* (Fig. 5C) (see also Fbranch - Fig. S9A #3). In short,
these results are consistent with *Mec. messenoides* being introgressed (Fig 2A - arrows a-d) and gene flow
between *polymnia-lysimnia-nesaea*.

In *Melinaea*, IQtree2 and StarBEAST2 produce the same topology (Fig. 1 vs Fig. 2), with *Mel. idae*
sister to *Mel. lilis* and *Mel. isocomma*. However, the AIM analysis groups *Mel. idae* with the ingroup clade,
although showing extensive gene flow from the lineage of *Mel. lilis* and *Mel. isocomma* (Fig. 2B - arrow
a; Fig S10B – arrow A,D; see also Fbranch (Fig. S9B - #1)). To resolve the position of *Mel. idae*, as well as
putative introgression between deeper branches of the phylogeny, we ran BPP focusing on the
relationships between the *lilis-isocomma-idae* clade and representatives of the ingroup clade (*mneme*,
marsaeus and *mothone*) (Fig. 5E-F). In the majority of the genome (43.4%) *lilis, idae* and *isocomma* group
together (like IQTree2 and StarBEAST2), while in 34.5% of the genome *idae* groups with *mothone-*
marsaeus-mneme (like AIM). Notably, the Z chromosome supports almost exclusively the latter
relationship. In 4.3% of the genome, *isocomma* clusters with *mothone-marsaeus-mneme*. *Mel. lilis* is
almost as often sister to *idae* (30.3%) as to *isocomma* (38.3%), and less commonly an outgroup to both
(4.1%), indicating more recent shared ancestry between *lilis* and *idae*, which is confirmed by Fbranch
excess allele sharing (Fig. S10B; Fig S9B - #2). *Mel. mothone, marsaeus* and *mneme* also vary in their
respective relationships (Fig. 5E).

We further investigated the timing of divergence and introgression for the three species showing
the strongest signals of introgression (*Mec. messenoides*, *Mec. nesaea* and *Mel. idae*), using a multispecies
coalescent-with-introgression (MSCi) approach. In all three cases, introgression is estimated to be old
(200-425 kya) (Fig. 3A-B scenario 1; Fig. 3C). For *Mec. messenoides* and *Mel. idae,* different replicate runs

produce different outcomes reflecting the uncertainty in the placement of these two species in the
phylogeny. Notably, some replicates indicate the origin of the introgressed lineages closely coincides with
the split time from both parents, suggestive of a hybrid origin (Fig. 3A-B scenario 2; Fig. S11; Table S2).
265

A focus on *Mec. nesaea*
As we propose to re-elevate *Mec. nesaea* to a species from its previous classification as a subspecies of
Mec. lysimnia, we investigated the relationships and reproductive isolation between *Mec. nesaea*, *Mec.*
lysimnia and *Mec. polymnia* in more detail. We resequenced 15 *Mec. nesaea*, 19 *Mec. l. lysimnia* and 9
Mec. p. casabranca, of which 24 were sampled from sympatry, and assessed the androconial compounds
that might act as pheromones, potentially contributing to assortative mating. Even though *Mec. lysimnia*
and *Mec. polymnia* have been observed to interbreed in nature (58) and putative hybrids have been
observed (Fig. S12), we find relatively high F_{ST} throughout the genome between all three species and no
evidence of ongoing gene flow based on ADMIXTURE analyses (Fig. S8; S13). Furthermore, they have been
shown to differ in number of chromosomes (Fig. 3D) (48), and we find their androconial bouquets to be
clearly distinct (Fig. 3E-F, Text S4; Table S3-4). All these lines of evidence suggest strong reproductive
isolation between *Mec. nesaea* and both *Mec. polymnia* and *Mec. lysimnia.* Genome scans for
introgression (f_{dM}) between *Mec. lysimnia* and *Mec. nesaea* compared to allopatric *Mec. polymnia* show
that this introgression is restricted to few genomic regions (Fig. 3G; Fig. S14-S15). The MSCi model for
Mec. nesaea suggests *Mec. nesaea* diverged from *Mec. polymnia* prior to *Mec. lysimnia* introgression.
However, *lysimnia*-introgression could have contributed key genetic variation to *Mec. nesaea* and
strengthened reproductive isolation to *Mec. polymnia*. Consistent with this hypothesis, we find that
introgression peaks coincide with peaks of elevated D_{xy} and F_{ST} between *Mec. nesaea* and *Mec. polymnia*
and mostly show no evidence of excess allele sharing between *Mec. nesaea* and sympatric *Mec. polymnia*
compared to allopatric *Mec. polymnia* (Fig. 3G; Fig. S13-S15), but the low levels of ongoing gene flow make
these measures poor predictors of reproductive isolation barriers.
287

Chromosomal rearrangements
To study chromosomal rearrangements, we generated haplotype-resolved genomes of five *Melinaea* and
five *Mechanitis* species using PacBio and HiC-data (Fig. S16), resulting in high quality genomes with >96%
BUSCO completeness and >94% of the genome assembled to chromosomes (Table S5-S6). The inferred
chromosome numbers match the chromosome counts from karyotyping (48). The *Mechanitis* genomes
are substantially shorter (291-320 Mb) than the *Melinaea* genomes (496-661 Mb, Table S5).

Whereas most Lepidoptera have conserved karyotypes with 31 chromosomes (59), our synteny
analysis using BUSCO genes show that *Mechanitis* and *Melinaea* genomes are highly rearranged compared
to the ancestral karyotype (Fig. S17-S19), corroborating earlier findings from two *Melinaea* genomes (60).
The median length of conserved syntenic blocks is 32-36 genes, compared to 168 genes in the outgroup
Danaus plexippus (Table S7; Fig. S19). The canonical Z has not undergone any fissions and has retained
the longest conserved syntenic blocks (225-227 genes; Fig. 4A; Table S7; Fig. S19) in our genomes.
However, both genera share a fusion of Z with parts of ancestral autosome 10 and *Mechanitis* has a further
fusion with parts of ancestral autosome 6 (Table S8; Fig. S18). In addition, a fusion-fission polymorphism
involving the Z and another autosome was observed between the Zs of the male *Mel. isocomma* (Table
S8; Fig. S16). A W-chromosome was identified in all females as a segment of varying size depleted of
BUSCO genes with moderate sequence similarity within genera, but none between (Fig. 4A). We found
W-autosome fusions in four species (Table S8). *Mec. macrinus* and *mazaeus* show complex fusions
between the W and multiple autosomes that are partially shared. Four species have two Z chromosomes,
as by definition, the homologue of the chromosome fused to the W becomes a Z chromosome. Similarly,
three species have two or three W chromosomes. Surprisingly, seven individuals were heterozygous for
one or more simple autosomal rearrangement and in *Mel. ludovica* we detected a complex chain
rearrangement involving four autosomes in one haplotype and three autosomes in the other haplotype
(Table S8; Fig. S16). None of the simple autosomal polymorphisms are shared among taxa.

While chromosome spreads had revealed species differences in chromosome counts (48), we
here show that these are not due to a few fissions or fusions, but complex rearrangements. Closely related
sister species show 3-47 chromosomal rearrangements (Fig. 4), which could contribute to reproductive
isolation. To assess if these chromosomal rearrangements confer reproductive isolation barriers, we
mapped the location of the chromosomal breakpoints between all species pairs to test for an association
between breakpoints and reduced gene flow. We found increased F_{ST} in all *Mechanitis* comparisons, and
reduced diversity (π) in breakpoint regions in seven of the ten comparisons (Fig. 4B, Fig. S20, Table S9). In
principle, elevated F_{ST} could be caused by increased background selection as recombination tends to be
reduced at chromosome ends and levels of ongoing gene flow are low (61, 62). However, we also find
elevated absolute divergence (D_{XY}) in windows coinciding with breakpoints in four of the ten comparisons,
which indicates that background selection alone cannot explain the pattern (63). In *Melinaea*, we did not
observe significantly elevated F_{ST} in the breakpoint regions, but we detected an increase in D_{XY} especially
in the comparisons involving the more distantly related *Mel. ludovica* (Table S9).

Discussion

Our results confirm that the two ithomiine genera *Melinaea* and *Mechanitis* represent fast and recent

radiations. They diverged in the past 1-2 million years, and speciated much faster than the other well-

studied Neotropical butterfly radiation of *Heliconius* (respectively 1.633 and 1.431 versus 0.324 speciation

events per lineage/My (46 species in 11.8 My)) (64). Kawahara et al. (21) previously found a significant

rate shift towards high diversification rates in ithomiine butterflies (clade L; 0.23 speciation events per

lineage/My) and we show that among Ithomiini, *Mechanitis* and *Melinaea* have an even higher speciation

rate, consistent with previous results (11). Our results shed light on potential drivers of their rapid

diversification, detailed below.

Given the high number of ancient introgression events across both genera, hybridisation may

have sped up speciation by boosting genetic variation as seen in other systems (e.g. 42). We identified

three species showing ancient introgression that might have a hybrid origin. Further research on which

genomic regions are more or less likely to introgress could inform us as to what extent introgression

contributed to reproductive isolation between those taxa and their parental lineages. For instance, similar

to *Heliconius* butterflies, introgression might have contributed to both mimicry ring switches and colour-

based assortative mating, which generate reproductive isolation, thereby driving speciation (65).

Despite ancient hybridisation, we find little evidence for ongoing gene flow between the species,

suggesting strong reproductive isolation. Some of this reproductive isolation is likely attributed to the

exceptionally high rates of chromosomal rearrangements across both genera, as complex chromosomal

rearrangements are expected to constitute barriers to gene flow (66, 67). While Lepidoptera

chromosomes are holocentric and may be able to tolerate simple fusions and fissions during meiosis, most

Lepidoptera have retained the same highly conserved karyotypes of 31 chromosomes (59, 68). The

massive chromosomal rearrangements we found in ithomiini are thus unusual, but there are other

lineages that also show high rates of fissions and fusions (e.g. *Leptidea* (69); and *Erebia* (70)) and there is

a slight association with variation in chromosome number and speciation rates (71).

Even though chromosomal rearrangements are thought to limit gene flow, many of our genomes

showed heterozygosity for chromosomal rearrangements. The heterozygote fission-fusions are

unambiguous in the HiC-data, matching the intraspecific variation in chromosome counts documented

previously (48) and cytogenetic evidence for pervasive polymorphism in *Mel. satevis cydon* (54). This

may be akin to *Leptidea sinapis*, where, despite multiple fusion/fission polymorphisms segregating within

populations (69, 72), crosses between chromosomal extremes show low survival of F2 hybrids (73). That

complex chromosomal rearrangements involving multiple chromosomes likely constitute stronger

barriers to gene flow compared to simple fusions and fissions has also been found in other taxa such as

shrews (74) and wallabies (75) and is in line with findings of nearly sterile hybrids in a cross between two

Melinaea sister species that differ in chromosome count (54). It is thus possible that simple fusions and

fissions are not strongly selected against, but if too many sequential fusions accumulate in isolated

populations, they cause reproductive isolation upon secondary contact.

Furthermore, we find many fusions between sex chromosomes and autosomes. Given the

disproportionate role sex chromosomes are proposed to have in speciation (Haldane's rule and/or large-

Z effect) (76–78), they may play a strong role in reproductive isolation of ithomiini butterflies. We found

two Z-autosome fusions shared among all congenerics, but no fission in the canonical Z, consistent with

conserved Z chromosomes in other Lepidoptera lineages (68). The heterozygote Z-autosome fusion in

Mel. isocomma potentially represents an ongoing fixation of a novel Z-autosome fusion, which might

rapidly rise in frequency if there are advantageous loci involved (79). The simple W-autosome fusion

observed in *Mel. menophilus* resemble those described in some *Heliconius* butterflies (80). The complex

fusion-fissions with multiple Ws and Zs in *Mec. macrinus* and *Mec. mazaeus* are similar to *Leptidea*

butterflies (81) and may constitute particularly strong barriers to gene flow (82).

Since the divergence times of *Mechanitis* and *Melinaea* species match Pleistocene climatic

fluctuations (2.58 Ma-11.7 ka), and many sister species are separated by the Andes or restricted to

different geographic regions corresponding to postulated Pleistocene refugia (26, 34, 37), we believe that

our findings are consistent with the speciation pump hypothesis. This hypothesis postulates that periods

of allopatry, followed by secondary sympatry, can facilitate diversification (37), as proposed in e.g. fishes

(45, 83) and Andean plants (84). The high rates of chromosomal rearrangements in *Mechanitis* and

Melinaea might accelerate the accumulation of chromosomal differences in isolated populations. On

coming together in secondary sympatry, these rearrangements likely cause low hybrid fitness, which in

turn might facilitate adaptation to different niches due to low levels of gene flow. Selection against

interbreeding might also lead to reinforcement of reproductive isolation via assortative mating, e.g. based

on pheromones (18, 85).

Our data suggests that hybridisation-derived genetic variation and high rates of chromosomal

rearrangements both may have played key roles in the fast diversification of *Melinaea* and *Mechanitis*,

and potentially contributed to adaptation and assortative mating facilitating species coexistence. Notably,

while an important role of hybridisation in diversification has been found in both insular environments

(e.g. silverswords (44), or cichlids (45)) and continental radiations (e.g. this study, *Heliconius* (86), or

Rhagoletis flies (87)), rapid radiations with high rates of chromosomal rearrangements seem to be

restricted to continents (e.g. this study, shrews (74), or wallabies (75)). Due to more opportunities for
geographical isolation on large land masses than in insular environments, factors such as high rates of
chromosomal rearrangements that speed up the allopatric accumulation of incompatibilities may thus be
particularly important in continental radiations.

Our study not only sheds light on drivers of continental radiations, but also largely resolves the
taxonomy of important biodiversity indicators. Hitherto, the study of these radiations has been hampered
by taxonomic challenges, whereas our combination of whole-genome resequencing with vast taxonomic
and geographic coverage, genome assemblies and androconial chemical analysis allowed us to resolve
taxonomic issues. Our study confirms that DNA barcoding can be misleading, massively underestimating
species richness, and should only be used with care to assess biodiversity. The taxonomic implications of
both introgression and karyotype diversity for species delimitation and designation of conservation units
are important to consider during monitoring and management of biodiversity in these vulnerable habitats.
402

Materials and methods

Collecting butterfly specimens

157 specimens of *Mechanitis*, 9 specimens of *Forbestra*, 109 specimens of *Melinaea,* 1 *Eutresis* and 1
Olyras were collected over the years 2000 to 2023 across Central and South America (Table S1). Adult
butterflies were caught with a net, and their bodies were subsequently preserved in ethanol, DMSO or
flash-frozen and stored at -70°C. Moreover, a few legs from dried museum specimens were used (Florida
Natural History Museum; Natural History Museum London). Wings were photographed and stored in
envelopes. The resulting dataset covers almost all species of *Melinaea* and *Mechanitis* from a wide
geographical range. In addition, 65 individuals of *Mec. lysimnia nesaea* (status restored to *Mec. nesaea* in
this paper; hereafter called *Mec. nesaea*), 19 *Mec. lysimnia lysimnia* and 8 *Mec. polymnia casabranca*
were collected for the androconial chemical analysis (Table S3).
414

DNA extractions & whole genome resequencing

DNA extractions were done with either the Qiagen MagAttract High Molecular Weight kit (Qiagen ID
67563), or the Qiagen QiaAmp DNA mini kit (51304), or a PureLink digestion and lysis step followed by a
magnetic bead DNA extraction (88). The dried museum specimens were extracted using a Lysis-C buffer
and a MinElute DNA extraction kit (protocol adapted from (89); Qiagen ID 28006). Library preparations
were performed using homemade TN5-transposase-mediated tagmentation (protocol adapted from
(90)), or following the manufacturer's guidelines with the Illumina DNA PCR-free library prep kit and

sequenced (150 bp paired-end) on Illumina NovaSeq 6000 or NovaSeq X machines at Novogene or the

Wellcome Sanger Institute.

Reference genomes

Haplotype-level chromosomally resolved reference genomes were assembled for five species of

Mechanitis (*Mec. messenoides, menapis, mazaeus, macrinus* and *polymnia*) and *Melinaea* (*Mel. ludovica,*

marsaeus, mothone, isocomma and *menophilus*). Note that earlier versions of two *Melinaea* genomes

were published previously (60). In short, we combined 12-57x PacBio HiFi sequencing and 33-197x

Illumina sequencing of HiC libraries (haploid coverages, Table S5) and assembled the genomes according

to the Tree of Life pipelines (https://github.com/sanger-tol/genomeassembly) (Text S5).

Whole genome mapping

To prepare the whole genome data for analysis, read quality was checked with FastQC (v0.11.9) (91).

Sequences below 50 bp were discarded and adapters and PolyG-tails were trimmed with FastP (v0.23.2)

(92), before they were aligned to *Melinaea marsaeus* (60) or *Mechanitis messenoides* using BWA-mem

(v.0.7.17) (93). Picard was used to remove PCR duplicates (v3.0.0) (94). Samtools (v1.17) (95) and GATK3

HaplotypeCaller (v3.8.1.0) (96, 97) were used for variant calling, with a minimum base quality score of 20.

VCFtools (v0.1.16) (98) was used for filtering. Based on the distribution of sequencing depth

(mean *Melinaea*: 7; *Mechanitis*: 15), all sites with a mean depth below 3 (*Melinaea*) or 5 (*Mechanitis*), and

above 15 (*Melinaea*) or 30 (*Mechanitis*) were removed. Insertions, deletions, sites with >50% missing data,

as well as genotypes with a depth below 2 (*Melinaea*) or 3 (*Mechanitis*) were removed. The mitochondrial

DNA was filtered separately, with a maximum depth of 1700 (*Melinaea*) or 1200 (*Mechanitis*).

Phylogenetic analyses

For each genus, we ran a Principal Component Analysis (PCA) using Plink (V1.9) to explore population

structure (99) (Fig. S21). Next, we inferred a phylogenetic tree based on a filtered subset of the whole

genome sequence data (also including monomorphic sites, thinned to 1 in 500 sites, with a minimum

genotype quality of 10). Our filtered VCF-files were converted to phylip with a custom script which calls

heterozygous sites as 'ambiguous' (equal likelihood for both alleles) to generate one sequence per

individual (vcf2phylip.py, http://www.github.com/joanam/scripts), and subsequently, IQtree2 (v2.1.2)

(100) produced phylogenetic trees with ultrafast bootstrap approximation (-B 1000; UFBoost) (101) and

the GTR-model.

We inferred separate phylogenies for mitochondrial and nuclear DNA. The nuclear trees are based

on 537,500 SNPs for *Mechanitis* and 784,526 SNPs for *Melinaea*. The mitochondrial phylogenies are based

on the full mitochondrial genome (not thinned) including 11,818 bp for *Mechanitis* and 11,815 bp for

Melinaea. For *Mechanitis*, we included a maximum of six individuals of the same subspecies and country

in the phylogenetic analyses, thus excluding several Brazilian *Mec. polymnia*, *Mec. nesaea* and *Mec.*

lysimnia. These individuals were included in hybridisation-analyses. For *Melinaea*, all individuals were

used.

In addition, a phylogenetic tree calibrated with divergence times from (11) was produced using

StarBEAST2 (102) (following https://beast2-dev.github.io/beast-

docs/beast2/DivergenceDating/DivergenceDatingTutorial.html) for the same dataset as the nuclear

phylogenetic tree, but thinned further to 1 in 5000 sites and with only one individual per lineage.

Divergence times between *Mec. mazaeus* and *Mec. macrinus* (0.39 MYA) and *Mel. isocomma* and *Mel.*

lilis (0.65 MY) were used for calibration. We used the HKY gamma-4 site model with a strict molecular

clock and the Calibrated Yule model. The model was run for 15.000.000 chains, stored every 5000 trees.

Phylogenies were visualised using the packages 'ape' (v5.7-1) and 'phytools' (v2.1-1) in R (Paradis

and Schliep 2019; Revell 2024) and FigTree (v1.1.4) (http://tree.bio.ed.ac.uk/software/figtree/). To

calculate a constant speciation rate (λ) we assumed a pure birth model, with n as the number of species

and T the time from root to tip: $n = e^{\lambda*T}$ which gives $\lambda = ln(n)/T$.

Distribution maps

The coordinates of our individuals and the individuals from (10) were plotted using the libraries 'sf',

'tmap', 'tmaptools' and 'mapview' in R (103–105). We classified the subspecies of (10) into species

following our taxonomic revision. Our individuals were plotted as dots on top of the overall distribution

combining data with (10). The basemap 'Esri.WorldTerrain' was used, provided through 'Leaflet' (ESRI,

ARCGIS; https://github.com/leaflet-extras/leaflet-providers); Fig. S6-7 has maps including the attribution.

Window trees

IQtree2 was used to produce phylogenetic trees in windows across the genome (one 20kb window every

200kb) and to calculate gene concordance factors between these window-trees and the whole-genome

phylogenetic tree (106).

DSuite analysis

We explored excess allele sharing between species or divergent subspecies using DSuite (107). We filtered

our genomic dataset to be 1 in 100 sites, only including variable sites. Dtrios calculated D and f4-ratio

statistics for all trios and Fbranch then summarised them as f-branch statistics using a species tree based

on the nuclear phylogeny of Fig. 2 (Table S1). The results of this analysis were visualised with a python

script provided by DSuite.

AIM with StarBEAST2

We followed the Approximate Isolatoin with Migration (AIM) in StarBEAST2 tutorial to obtain a

phylogenetic tree with admixture arrows (108, 109). For each genus, we picked one individual per species

(highest depth), and randomly extracted 150 (*Mechanitis*) or 200 (*Melinaea*) 800-1000bp windows across

the genome. We used the HKY gamma-4 site model with a strict molecular clock and the Yule model. The

migration rate was set to 25 with an initial value of 0.1, and we ran the model for 100.000.000 chains,

storing every 5000 trees. We updated the parameters according to the suggestions in the output of the

first run and re-ran the model to improve the results.

Species-tree inference

Phylogenetic relationships across the genome between species of *Melinaea* and *Mechanitis* were inferred

using the multispecies coalescent (MSC) approach implemented in BPP v.4.6.2 (110), while accounting for

incomplete lineage sorting. For each species, only the individual with the highest coverage was included.

To minimise the effect of linked selection, loci were required to be at least 2 kb from annotated exons.

Because the analysis assumes no intra-locus recombination and independence between loci, we selected

loci of 300 bp and at least 2 kb from neighbouring loci. Sequence alignments were produced for all loci,

masking repetitive elements annotated in the reference genome using RepeatMasker v4.1.5

(http://repeatmasker.org/). For each locus, individuals with more than 80% missing genotype calls were

excluded from the alignment and only loci with all individuals passing filters were considered.

Furthermore, sites with missing genotype calls were removed and loci with less than 30 bp passing filters

were excluded. Heterozygous sites were assigned IUPAC codes. Loci were grouped into blocks of 100 loci.

Species-tree estimation was then performed in BPP v.4.6.2 using the A01 analysis (species-tree inference

assuming no gene flow). Inverse gamma priors (invGs) were applied both to the root age ($\tau 0$) and to

effective population sizes (θ) – invG(3, 0.06) and invG(3, 0.04), respectively. Parameters were scaled

assuming a mutation rate of 2.9×10^{-9} substitutions per site per generation and 4 generations per year.

The MCMC was run for 1,000,000 iterations after 50,000 iterations of burn-in, sampling every 10

iterations.

Demographic modelling

For the three highly introgressed species *Mel. idae, Mec. messenoides* and *Mec. nesaea, we ran* a

multispecies-coalescent-with-introgression (MSCi) model implemented in BPP v.4.6. 2 (110) to better

estimate their position in the phylogeny and divergence timing while accounting for admixture. For *Mel.*

idae we considered *Mel. lilis* and *Mel. marsaeus* as sister/donor species, while for *Mec. messenoides*, *Mec.*

menapis and *Mec. polymnia* (West) were chosen. For *Mec. nesaea*, we used one individual of each of the

Brazilian populations (*Mec. nesaea, Mec. lysimnia lysimnia*, *Mec. polymnia casabranca*). Loci were

selected randomly from autosomes, and required to be at least 2 kb from annotated exons and 10 kb from

the nearest locus, and a maximum size of 2 kb. For each locus, individuals with more than 50% missing

data and sites containing missing genotypes or overlapping annotated repetitive elements were removed.

Only loci at least 800 bp long after filters and without missing individuals were considered. Heterozygous

sites were assigned IUPAC codes. Demographic parameters were estimated using a fixed species tree with

introgression events (Fig. S11). An inverse gamma prior (invG) was applied for all population size

parameters θ (α=3; β=0.04) and root age parameter τ (α=3; β=0.06). A beta prior was applied to the

introgression probability (ϕ) (α=1; β=1). Three replicate MCMC runs of 1,000,000 iterations, after a burn-

in period of 50,000 iterations, sampling every 50 iterations were performed for each dataset. Divergence

time was calculated based on 4 generations per year, and a mutation rate of 2.90E-09 as in *Heliconius* (43)

(T=τ/mutation rate/generations/million years).

Genome scans and ADMIXTURE

F_{ST}, D_{XY} and pi (π) were calculated in all species pairs in 20kb windows (including monomorphic sites, with

a minimum of 10,000 sites per window) across the genome with the scripts from Simon Martin

(https://github.com/simonhmartin/genomics_general). We also ran ADMIXTURE in subsets of *Mechanitis*

and *Melinaea* to see clustering between individuals with k ranging from 2 to 9 (111). For this dataset, we

used the same filter settings as for the Fbranch analysis (see above).

Introgression (f_{dM}*) scans*

To examine gene flow across the genome of *Mec. nesaea*, we computed f_{dM} with the script by Simon

Martin (ABBABABAwindows.py, https://github.com/simonhmartin/genomics_general) (112) with various

populations for P1,2,3,4. If for example P1=*Mec. polymnia casabranca* (Brazil), P2=*Mec. nesaea*, P3=*Mec.*

lysimnia Brazil, and P4=*Forbestra*, high f_{dM} values indicate introgression between *Mec. lysimnia* and *Mec.*

nesaea. We also tested other outgroups (*Mec. messenoides*) and with five single *Mec. nesaea* individuals,

to see if the signal was consistent in different individuals (Fig. S14).

Androconial Chemistry

We obtained samples of the androconial secretions from 92 males (Text S5). To exclude non-androconial

compounds from our analyses, for each population the same extraction procedure was adopted with

wings of conspecific females and two non-androconial wing areas of males. Eight solvent negative controls

were also taken for each sampling event.

The peak areas of each chromatogram were integrated with MSD ChemStation E.02.01.1177

(Agilent Technologies, USA) to obtain the total ion current signals. A series of linear alkanes (C7–C40) was

used to determine the linear retention indices (RI) of each compound (113). Compounds were identified

by comparing their mass spectra and retention indices with those of reference samples available from

personal and commercial mass spectral libraries (FFNSC 2, MassFinder 4, NIST17, MACE v.5.1 (114), and

Wiley Registry™ 9th ed.). The peaks exclusive to the androconial samples were used to determine the

relative percentages of each compound per sample.

Dihydropyrrolizines, such as hydroxydanaidal or methyl hydroxydanaidoate, are typically

accompanied by smaller peaks formed by degradation during GC/MS analyses and/or storage (115). These

degradation peaks were excluded from the generated ion chromatograms and statistical analysis. To avoid

non-evident contaminants, we only considered compounds present in more than three individuals for any

given taxon.

The data were plotted in R (116) using the vegan package

(https://doi.org/10.32614/CRAN.package.vegan) for nonmetric multidimensional scaling with

'monoMDS', specifying a global model, square root transformation and Wisconsin double standardisation

(autotransform=TRUE).

Chromosome rearrangements

Synteny and breakpoint analysis was performed using single copy orthologous genes identified with

BUSCO version 5.7.1 with the lineage database lepidoptera_odb10 and otherwise default options (117).

To compare large scale rearrangements in the *Mechanitis* and *Melinaea* versus the butterfly ancestral

linkage groups (Merian elements), we used two outgroup genomes, *Melitaea cinxia* (118) and *Danaus*

plexippus (119). The output from BUSCO was filtered with a custom script to contain only complete genes
located in chromosome-sized scaffolds and excluding W. Only single copy genes were included with the
exception of genes on the neo-Z2*,* where we also included genes classified as duplicated. The sex linking
of Z2 appears to be recent and most of the genes had high similarity to the genes on the corresponding
W's and were therefore classified as duplicated by BUSCO. Genes that were actually duplicated (occurring
more than once or present on other chromosomes than W) on the Z2 were removed. We determined
syntenic blocks, excluding single gene translocations, by comparing the position of the BUSCO-genes in
each genome against the Merian elements (*Melitaea cinxia*), and visualised the syntenic relationship with
the R-package gggenomes (120). Minimum number of fusions were determined by the number of
different Merian elements located in every query chromosome and fissions were determined as the
number of query chromosomes containing parts of each Merian element for each species. The
breakpoints between all species within *Mechanitis* and *Melinaea* was determined by the same principle
as above using an all against all approach for each genome to compare the BUSCO-gene positions. Synteny
analysis of the sex chromosomes and between haplotypes were performed with whole genome alignment
using minimap2/2.27 with default settings and -x asm10 (1% sequence divergence) (121) and visualised
after removing short alignments (<100kb for multispecies alignment, <500kb for haplotype alignment)
using a modified version of the R-package Farre-lab/syntenyPlotteR (122).

To investigate the association between chromosomal rearrangements and species divergence we
mapped the location of the breakpoints between each comparison to the reference genomes *Mec.*
messenoides and *Mel. marsaeus*. Divergence and diversity was estimated in 20 kb windows along the
genome (detailed above). To determine whether the observed statistics in the breakpoint regions were
different from the random expectation of regions of the same size and number across the genome, we
used permTest with the randomise function option 'randomizeRegions' and evaluate function option
'meanInRegions' for 50,000 permutations, as implemented in the R-package regioneR (123). The analyses
above were performed in R v4.4.0 (116). For conservative interpretation multiple correction was
performed with Bonferroni adjustment (α/n), where α = 0.05 and n = 10 for the comparisons within each
genus, resulting in values considered significantly different if their p-value from the randomisation test
was less than 0.005.

Funding

John's College, Cambridge awarded to JIM. ESMH was supported by NERC DTP C-CLEAR, the Zoology Department of the University of Cambridge, St John's College, Cambridge, and the Wellcome Sanger Institute PhD programme. CEBN had a PhD Scholarship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior. Sample collection was further supported by Leverhulme trust (KW and ME), a Phyllis and Eileen Gibbs Fellowship, ATIP grant, ANR SPECREP and CLEARWING, and HFSP RGP0014/2016 (ME and MM), a Research Fellowship from the Royal Commission for the Great Exhibition of 1851 and a Royal Society Research Grant (SHM) and Deutsche Forschungsgemeinschaft, Schu984/12-2 (SS).

Acknowledgements

We thank Ismael Aldas and Raúl Aldaz for assistance with catching butterflies in Ecuador, Mario Tuanama and Ronald Mori-Pezo for help with rearing and collecting butterflies in Peru, Augusto Rosa for sampling support in São Paulo, Erika de Castro for sampling in Brazil, Mathieu Joron for sampling in French Guiana and John R. MacDonald for sampling in Panama. We thank the Peruvian, Ecuadorian and Brazilian authorities as well as the Museo de Historia Natural in Peru and the Museo Ecuatoriano de Ciencias Naturales in Ecuador for their support. Many thanks to Dr Blanca Huertas and Robyn Crowther from the Natural History Museum in London (NHMUK) for providing butterfly legs of *Mel. isocomma* and *Mel. mothone* and to Dr Petra Korlevic for assistance with extracting DNA from museum specimens. We are grateful to the editor and reviewers for their helpful comments that have greatly improved our manuscript.

Data, materials and software availability

All code and tables underlying the figures are found in our GitHub repository: https://github.com/rapidspeciation/mechanitis_melinaea.

References

1. J. B. Losos, Adaptive radiation, ecological opportunity, and evolutionary determinism : American society of naturalists E. O. Wilson award address. *American Naturalist* **175**, 623–639 (2010).

2. R. G. Gillespie, *et al.*, Comparing Adaptive Radiations Across Space, Time, and Taxa in *Journal of Heredity*, (Oxford University Press, 2020), pp. 1–20.

3. E. Gittenberger, Radiation and adaptation, evolutionary biology and semantics. *organisms, diversity & evolution* **4**, 135–136 (2004).

4. J. E. Czekanski-Moir, R. J. Rundell, The Ecology of Nonecological Speciation and
Nonadaptive Radiations. *Trends in Ecology & Evolution* **34**, 400–415 (2019).

5. R. De-Kayne, *et al.*, Why Do Some Lineages Radiate While Others Do Not? Perspectives
for Future Research on Adaptive Radiations. *Cold Spring Harb Perspect Biol* a041448
(2024). https://doi.org/10.1101/cshperspect.a041448.

6. J. Cerca, *et al.*, Evolutionary genomics of oceanic island radiations. *Trends in Ecology &*
Evolution **38**, 631–642 (2023).

7. S. Claramunt, Discovering exceptional diversifications at continental scales: the case of
the endemic families of neotropical suboscine passerines. *Evolution* (2010).
https://doi.org/10.1111/j.1558-5646.2010.00971.x.

8. R. Maestri, *et al.*, The ecology of a continental evolutionary radiation: Is the radiation of
sigmodontine rodents adaptive? *Evolution* **71**, 610–632 (2017).

9. S. Tiatragul, A. Skeels, J. S. Keogh, Morphological evolution and niche conservatism
across a continental radiation of Australian blindsnakes. *Evolution* **78**, 1854–1868 (2024).

10. M. Doré, *et al.*, Anthropogenic pressures coincide with Neotropical biodiversity hotspots in
a flagship butterfly group. *Diversity and Distributions* **28**, 2912–2930 (2022).

11. N. Chazot, *et al.*, Renewed diversification following Miocene landscape turnover in a
Neotropical butterfly radiation. *Global Ecology and Biogeography* **28**, 1118–1132 (2019).

12. G. W. Beccaloni, K. J. Gaston, Predicting the species richness of neotropical forest
butterflies: Ithomiinae (Lepidoptera: Nymphalidae) as indicators. *Biological Conservation*
71, 77–86 (1995).

13. K. S. Brown, "Conservation of Neotropical Environments: Insects as Indicators" in *The*
Conservation of Insects and Their Habitats, (Elsevier, 1991), pp. 349–404.

14. K. S. Brown, Adult-obtained pyrrolizidine alkaloids defend ithomiine butterflies against a
spider predator. *Nature* **309**, 707–709 (1984).

15. P. J. Devries, F. G. Stiles, Attraction of Pyrrolizidine Alkaloid Seeking Lepidoptera to
Epidendrum paniculatum Orchids. **22**, 290–297 (1990).

16. J. R. Trigo, K. S. Brown, Variation of pyrrolizidine alkaloids in Ithomiinae: a comparative
study between species feeding on Apocynaceae and Solanaceae. *Chemoecology* **22**,
22–29 (1990).

17. A. V. L. Freitas, *et al.*, Tropane and pyrrolizidine alkaloids in the ithomiinesPlacidula
euryanassa andMiraleria cymothoe (Lepidoptera: Nymphalidae). *Chemoecology* **7**, 61–67
(1996).

18. M. McClure, *et al.*, Why has transparency evolved in aposematic butterflies? Insights from
the largest radiation of aposematic butterflies, the Ithomiini. *Proceedings of the Royal*
Society B: Biological Sciences **286** (2019).

19.   F. Müller, Ituna and Thyridia: a remarkable case of Mimicry in Butterflies. *Kosmos* **May**,
682   100 (1879).

683   20.   G. W. Beccaloni, Ecology, natural history and behaviour of ithomiine butterflies and their
684   mimics in Ecuador (Lepidoptera: Nymphalidae: Ithomiinae). *Tropical Lepidoptera* **8**, 103–
685   124 (1997).

686   21.   A. Y. Kawahara, *et al.*, A global phylogeny of butterflies reveals their evolutionary history,
687   ancestral hosts and biogeographic origins. *Nature Ecology and Evolution* **7**, 903–913
688   (2023).

689   22.   R. M. Fox, A monograph of the Ithomiidae (Lepidoptera). Part III. The Tribe Mechanitini.
690   *Memoirs of the American Entomological Society* **22** (1967).

691   23.   K. K. Dasmahapatra, M. Elias, R. I. Hill, J. I. Hoffman, J. Mallet, Mitochondrial DNA
692   barcoding detects some species that are real, and some that are not. *Molecular Ecology*
693   *Resources* **10**, 264–273 (2010).

694   24.   M. Elias, *et al.*, Limited performance of DNA barcoding in a diverse community of tropical
695   butterflies. *Proc. R. Soc. B.* **274**, 2881–2889 (2007).

696   25.   R. I. Hill, *et al.*, Ecologically relevant cryptic species in the highly polymorphic Amazonian
697   butterfly Mechanitis mazaeus s.l. (Lepidoptera: Nymphalidae; Ithomiini). *Biological*
698   *Journal of the Linnean Society* **106**, 540–560 (2012).

699   26.   K. S. Brown, Geographical patterns of evolution in Neotropical Lepidoptera: differentiation
700   of the species of Melinaea and Mechanitis (Nymphalidae, Ithomiinae). *Systematic*
701   *Entomology* **2**, 161–197 (1977).

702   27.   W. T. M. Forbes, A second review of Melinaea and Mechanitis (Lepidoptera, Ithomiinae).
703   *Journal of the New York Entomological Society* **56**, 1–24 (1948).

704   28.   R. M. Fox, A monograph of the Ithomiidae (Lepidoptera). Part II. The Tribe Melinaeini
705   Clark. *Transactions of the American Entomological Society* **86**, 109–171 (1960).

706   29.   G. Lamas, Checklist: Part 4A. Hesperioidea - Papilionoidea. *Atlas of Neotropical*
707   *Lepidoptera* **5** (2004).

708   30.   K. R. Willmott, J. Mallet, Correlations between adult mimicry and larval host plants in
709   ithomiine butterflies. *Proceedings of the Royal Society B: Biological Sciences* **271** (2004).

710   31.   N. Chazot, *et al.*, Mutualistic Mimicry and Filtering by Altitude Shape the Structure of
711   Andean Butterfly Communities. *The American Naturalist* **183**, 26–39 (2014).

712   32.   M. McClure, M. Elias, Unravelling the role of host plant expansion in the diversification of
713   a Neotropical butterfly genus. *BMC Evolutionary Biology* **16** (2016).

714   33.   I. Birskis-Barros, A. V. L. Freitas, P. R. Guimarães, Habitat generalist species constrain
715   the diversity of mimicry rings in heterogeneous habitats. *Sci Rep* **11**, 5072 (2021).

34.   K. S. Brown, Quaternary refugia in tropical America: evidence from race formation in
Heliconius butterflies. *Proc. R. Soc. Lond. B.* **187**, 369–378 (1974).

35.   A. Whinnett, *et al.*, Strikingly variable divergence times inferred across an Amazonian
butterfly 'suture zone.' *Proc. R. Soc. B.* **272**, 2525–2533 (2005).

36.   K. K. Dasmahapatra, G. Lamas, F. Simpson, J. Mallet, The anatomy of a "suture zone" in
Amazonian butterflies: A coalescent-based test for vicariant geographic divergence and
speciation. *Molecular Ecology* **19**, 4283–4301 (2010).

37.   J. Haffer, Speciation in Amazonian Forest Birds: Most species probably originated in
forest refuges during dry climatic periods. *Science* **165**, 131–137 (1969).

38.   N. Chazot, *et al.*, Contrasting patterns of Andean diversification among three diverse
clades of Neotropical clearwing butterflies. *Ecology and Evolution* **8**, 3965–3982 (2018).

39.   J. Mallet, Hybrid speciation. *Nature* **446**, 279–283 (2007).

40.   The Heliconius Genome Consortium, Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. *Nature* **487**, 94–98 (2012).

41.   R. Abbott, *et al.*, Hybridization and speciation. *J of Evolutionary Biology* **26**, 229–246
(2013).

42.   D. A. Marques, K. Lucek, V. C. Sousa, L. Excoffier, O. Seehausen, Admixture between
old lineages facilitated contemporary ecological speciation in Lake Constance
stickleback. *Nature Communications* **10** (2019).

43.   N. Rosser, *et al.*, Hybrid speciation driven by multilocus introgression of ecological traits.
Nature **628**, 811–817 (2024).

44.   M. Barrier, B. G. Baldwin, R. H. Robichaux, M. D. Purugganan, Interspecific hybrid
ancestry of a plant adaptive radiation: allopolyploidy of the Hawaiian silversword alliance
(Asteraceae) inferred from floral homeotic gene duplications. *Molecular Biology and*
Evolution **16**, 1105–1113 (1999).

45.   J. I. Meier, *et al.*, Cycles of fusion and fission enabled rapid parallel adaptive radiations in
African cichlids. *Science* **381**, eade2833 (2023).

46.   R. W. R. Wallbank, *et al.*, Evolutionary Novelty in a Butterfly Wing Pattern through
Enhancer Shuffling. *PLoS Biology* **14** (2016).

47.   L. H. Rieseberg, *et al.*, Major Ecological Transitions in Wild Sunflowers Facilitated by
Hybridization. *Science* **301**, 1211–1216 (2003).

48.   K. S. Brown, B. V. Schoultz, E. Suomalainen, Chromosome evolution in Neotropical
Danainae and Ithomiinae (Lepidoptera). *Hereditas* **141**, 216–236 (2004).

49.   C. A. Everett, J. B. Searle, B. M. N. Wallace, A study of meiotic pairing, nondisjunction
and germ cell death in laboratory mice carrying Robertsonian translocations. *Genet. Res.*
67, 239–247 (1996).

50.   F. J. Ayala, M. Coluzzi, Chromosome speciation: Humans, Drosophila, and mosquitoes.
(2005).

51.   K. Yoshida, *et al.*, Chromosome fusions repatterned recombination rate and facilitated
reproductive isolation during Pristionchus nematode speciation. *Nat Ecol Evol* (2023).
https://doi.org/10.1038/s41559-022-01980-z.

52.   M. A. F. Noor, K. L. Grams, L. A. Bertucci, J. Reiland, Chromosomal inversions and the
reproductive isolation of species. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 12084–12088 (2001).

53.   L. H. Rieseberg, Chromosomal rearrangements and speciation. (2001).

54.   M. McClure, B. Dutrillaux, A.-M. Dutrillaux, V. Lukhtanov, M. Elias, Heterozygosity and
Chain Multivalents during Meiosis Illustrate Ongoing Evolution as a Result of Multiple
Holokinetic Chromosome Fusions in the Genus *Melinaea* (Lepidoptera, Nymphalidae).
Cytogenet Genome Res **153**, 213–222 (2017).

55.   J. Mallet, A species definition for the Modern Synthesis. *TREE* **10**, 294–299 (1995).

56.   M. McClure, M. Elias, Ecology, life history, and genetic differentiation in Neotropical
Melinaea (Nymphalidae: Ithomiini) butterflies from north-eastern Peru. *Zoological Journal*
of the Linnean Society **179**, 110–124 (2016).

57.   A. H. B. Rosa, E. P. Barbosa, N. Wahlberg, A. V. L. Freitas, Systematic position and
conservation aspects of Melinaea mnasias thera (Lepidoptera: Nymphalidae: Danainae).
Nat. Conserv. Res. **9** (2024).

58.   J. Vasconcellos-Neto, K. S. Brown, Interspecific Hybridization in Mechanitis Butterflies
(Ithomiinae): a Novel Pathway for the Breakdown of Isolating Mechanisms. *Biotropica* **14**,
288 (1982).

59.   R. Robinson, *Lepidoptera Genetics* (Pergamon, 1971).

60.   J. Gauthier, *et al.*, First chromosome scale genomes of ithomiine butterflies
(Nymphalidae: Ithomiini): Comparative models for mimicry genetic studies. *Molecular*
Ecology Resources **23**, 872–885 (2023).

61.   S. H. Martin, J. W. Davey, C. Salazar, C. D. Jiggins, Recombination rate variation shapes
barriers to introgression across butterfly genomes. *PLoS Biol* **17**, e2006288 (2019).

62.   K. Näsvall, *et al.*, Nascent evolution of recombination rate differences as a consequence
of chromosomal rearrangements. *PLoS Genet* **19**, e1010717 (2023).

63.   T. E. Cruickshank, M. W. Hahn, Reanalysis suggests that genomic islands of speciation
are due to reduced diversity, not reduced gene flow. *Molecular Ecology* **23**, 3133–3157
(2014).

64.   K. M. Kozak, *et al.*, Multilocus species trees show the recent adaptive radiation of the
mimetic heliconius butterflies. *Systematic Biology* **64**, 505–524 (2015).

65.   M. Rossi, *et al.*, Adaptive introgression of a visual preference gene. (2024).

66.   M. Kirkpatrick, N. Barton, Chromosome Inversions, Local Adaptation and Speciation.
789   *Genetics* **173**, 419–434 (2006).

790   67.   A. Mackintosh, Chromosome Fissions and Fusions Act as Barriers to Gene Flow between
791   Brenthis Fritillary Butterflies. *Molecular Biology and Evolution* **40** (2023).

792   68.   C. J. Wright, L. Stevens, A. Mackintosh, M. Lawniczak, M. Blaxter, Comparative
793   genomics reveals the dynamics of chromosome evolution in Lepidoptera. *Nat Ecol Evol* **8**,
794   777–790 (2024).

795   69.   V. A. Lukhtanov, V. Dincă, G. Talavera, R. Vila, Unprecedented within-species
796   chromosome number cline in the Wood White butterfly Leptidea sinapis and its
797   significance for karyotype evolution and speciation. *BMC Evol Biol* **11**, 109 (2011).

798   70.   H. Augustijnen, *et al.*, A macroevolutionary role for chromosomal fusion and fission in
799   Erebia butterflies. *Science Advances* (2024).

800   71.   J. M. De Vos, H. Augustijnen, L. Bätscher, K. Lucek, Speciation through chromosomal
801   fusion and fission in Lepidoptera. *Phil. Trans. R. Soc. B* **375**, 20190539 (2020).

802   72.   L. Höök, K. Näsvall, R. Vila, C. Wiklund, N. Backström, High-density linkage maps and
803   chromosome level genome assemblies unveil direction and frequency of extensive
804   structural rearrangements in wood white butterflies (Leptidea spp.). *Chromosome Res* **31**,
805   2 (2023).

806   73.   V. A. Lukhtanov, *et al.*, Versatility of multivalent orientation, inverted meiosis, and rescued
807   fitness in holocentric chromosomal hybrids. *Proc. Natl. Acad. Sci. U.S.A.* **115** (2018).

808   74.   S. Garagna, M. Zuccotti, J. B. Searle, C. A. Redi, P. J. Wilkinson, Spermatogenesis in
809   heterozygotes for Robertsonian chromosomal rearrangements from natural populations of
810   the common shrew, Sorex araneus. *Reproduction* **87**, 431–438 (1989).

811   75.   S. Potter, *et al.*, Limited Introgression between Rock-Wallabies with Extensive
812   Chromosomal Rearrangements. *Molecular Biology and Evolution* **39**, msab333 (2022).

813   76.   J. B. S. Haldane, Sex ratio and unisexual sterility in hybrid animals. *Journ. of Gen.* **12**,
814   101–109 (1922).

815   77.   L. Z. Carabajal Paladino, *et al.*, Sex Chromosome Turnover in Moths of the Diverse
816   Superfamily Gelechioidea. *Genome Biology and Evolution* **11**, 1307–1319 (2019).

817   78.   A. J. Mongue, M. E. Hansen, J. R. Walters, Support for faster and more adaptive Z
818   chromosome evolution in two divergent lepidopteran lineages[*]. *Evolution* **76**, 332–345
819   (2022).

820   79.   D. Charlesworth, B. Charlesworth, Sex differences in fitness and selection for centric
821   fusions between sex-chromosomes and autosomes. *Genet. Res.* **35**, 205–214 (1980).

822   80.   N. Rueda-M, *et al.*, Genomic evidence reveals three W-autosome fusions in Heliconius
823   butterflies. *PLoS Genet* **20**, e1011318 (2024).

81. J. Šíchová, *et al.*, Fissions, fusions, and translocations shaped the karyotype and multiple
sex chromosome constitution of the northeast-Asian wood white butterfly, *Leptidea*
amurensis. Biol. J. Linn. Soc. **118**, 457–471 (2016).

82. T. Xiong, *et al.*, A polygenic explanation for Haldane's rule in butterflies. *Proc. Natl. Acad.*
Sci. U.S.A. **120**, e2300959120 (2023).

83. J. April, R. H. Hanner, A.-M. Dion-Côté, L. Bernatchez, Glacial cycles as an allopatric
speciation pump in north-eastern American freshwater fishes. *Molecular Ecology* **22**,
409–422 (2013).

84. B. Nevado, N. Contreras-Ortiz, C. Hughes, D. A. Filatov, Pleistocene glacial cycles drive
isolation, gene flow and speciation in the high-elevation Andes. *New Phytologist* **219**,
779–793 (2018).

85. P. M. B. Bacquet, *et al.*, Selection on male sex pheromone composition contributes to
butterfly reproductive isolation. *Proc. R. Soc. B.* **282**, 20142734 (2015).

86. R. M. Merrill, *et al.*, The diversification of Heliconius butterflies: What have we learned in
150 years? *Journal of Evolutionary Biology* **28**, 1417–1438 (2015).

87. T. H. Q. Powell, *et al.*, The Build-Up of Population Genetic Divergence along the
Speciation Continuum during a Recent Adaptive Radiation of Rhagoletis Flies. *Genes* **13**,
275 (2022).

88. M. Kucka, Chan, Yingguang Frank, HMW DNA extraction using magnetic beads.
protocols.io (2022). https://dx.doi.org/10.17504/protocols.io.b46bqzan.

89. P. Korlević, *et al.*, A Minimally Morphologically Destructive Approach for DNA Retrieval
and Whole-Genome Shotgun Sequencing of Pinned Historic Dipteran Vector Species.
Genome Biology and Evolution **13**, evab226 (2021).

90. S. Picelli, *et al.*, Tn5 transposase and tagmentation procedures for massively scaled
sequencing projects. *Genome Research* **24**, 2033–2040 (2014).

91. Andrews, S, FastQC: A quality control tool for high throughput sequence data. (2010).

92. S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ preprocessor in
Bioinformatics, (Oxford University Press, 2018), pp. i884–i890.

93. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754–1760 (2009).

94. Picard, Picard Toolkit. *Broad Institute* (2019). https://broadinstitute.github.io/picard/.

95. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10** (2021).

96. A. McKenna, *et al.*, The genome analysis toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Research* **20**, 1297–1303 (2010).

97. V. Ruano-Rubio, *et al.*, Scaling accurate genetic variant discovery to tens of thousands of
samples. (2017). https://doi.org/10.1101/201178.

98.   P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

99.   S. Purcell, *et al.*, PLINK: A tool set for whole-genome association and population-based
linkage analyses. *American Journal of Human Genetics* **81**, 559–575 (2007).

100. B. Q. Minh, *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534 (2020).

101. D. T. Hoang, *et al.*, UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol. Biol.*
Evol **35**, 518–522 (2017).

102. H. A. Ogilvie, R. R. Bouckaert, A. J. Drummond, StarBEAST2 Brings Faster Species Tree
Inference and Accurate Estimates of Substitution Rates. *Molecular Biology and Evolution*
34, 2101–2114 (2017).

103. E. Pebesma, Simple Features for R: Standardized Support for Spatial Vector Data. *The R*
Journal **10**, 439 (2018).

104. M. Tennekes, tmap: Thematic Maps in R. *J. Stat. Soft.* **84** (2018).

105. T. Appelhans, F. Detsch, C. Reudenbach, S. Woellauer, Mapview: Interactive viewing of
spatial data in R. *R package* **version 2.11.2** (2023).

106. B. Q. Minh, M. W. Hahn, R. Lanfear, New methods to calculate concordance factors for
phylogenomic datasets. *Molecular Biology and Evolution* **37**, 2727–2733 (2020).

107. M. Malinsky, M. Matschiner, H. Svardal, Dsuite - Fast D-statistics and related admixture
evidence from VCF files. *Molecular Ecology Resources* **21**, 584–595 (2021).

108. R. Bouckaert, *et al.*, BEAST 2: A Software Platform for Bayesian Evolutionary Analysis.
PLoS Comput Biol **10**, e1003537 (2014).

109. N. F. Müller, *et al.*, Joint inference of species histories and gene flow. [Preprint] (2018).
Available at: http://biorxiv.org/lookup/doi/10.1101/348391 [Accessed 7 November 2024].

110. T. Flouri, X. Jiao, B. Rannala, Z. Yang, A Bayesian Implementation of the Multispecies
Coalescent Model with Introgression for Phylogenomic Analysis. *Molecular Biology and*
Evolution **37**, 1211–1223 (2020).

111. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

112. S. H. Martin, J. W. Davey, C. D. Jiggins, Evaluating the Use of ABBA–BABA Statistics to
Locate Introgressed Loci. *Molecular Biology and Evolution* **32**, 244–257 (2015).

113. H. van D. Dool, P. D. Kratz, A generalization of the retention index system including linear
temperature programmed gas-liquid partition chromatography. *Journal of*
Chromatography A **11**, 463–471 (1963).

114. S. Schulz, A. Möllerke, MACE – An Open Access Data Repository of Mass Spectra for
Chemical Ecology. *J Chem Ecol* **48**, 589–597 (2022).

115. S. Schulz, *et al.*, Semiochemicals derived from pyrrolizidine alkaloids in male ithomiine
butterflies (Lepidoptera: Nymphalidae: Ithomiinae). *Biochemical Systematics and Ecology*
32, 699–713 (2004).

116. R Core Team, A language and environment for statistical computing. *R foundation for
statistical computing* (2024). https://doi.org/10.1080/01621459.1972.10481279.

117. M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology
and Evolution* **38**, 4647–4654 (2021).

118. R. Vila, *et al.*, The genome sequence of the Glanville fritillary, Melitaea cinxia (Linnaeus,
1758). *Wellcome Open Res* **6**, 266 (2021).

119. L. Gu, *et al.*, Dichotomy of Dosage Compensation along the Neo Z Chromosome of the
Monarch Butterfly. *Current Biology* **29**, 4071-4077.e3 (2019).

120. T. Hackl, M. J. Ankenbrand, B. van Adrichem, gggenomes: a grammar of graphics for
comparative genomics. *R package* (2024). https://github.com/thackl/gggenomes.

121. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

122. S. Quigley, J. Damas, D. M. Larkin, M. Farré, syntenyPlotteR: a user-friendly R package
to visualize genome synteny, ideal for both experienced and novice bioinformaticians.
Bioinformatics Advances **3**, vbad161 (2023).

123. B. Gel, *et al.*, regioneR: an R/Bioconductor package for the association analysis of
genomic regions based on permutation tests. *Bioinformatics* **32**, 289–291 (2016).

124. M. Carvalho, A. Victor, L. Freitas, E. Barbosa, Immature stages of Mechanitis polymnia
casabranca (Nymphalidae, Danainae). (2019). https://doi.org/10.5281/zenodo.2650300.

125. D. H. A. Melo, A. V. L. Freitas, Immature stages of Mechanitis lysimnia nesaea
(Nymphalidae: Danainae: Ithomiini). *Tropical Lepidoptera* **33**, 79–85 (2023).

Figures

Figure 1 - Rampant cytonuclear discordance in *Mechanitis and Melinaea* and a need for taxonomic revision

Co-phyloplot showing the nuclear and mitochondrial phylogenies of 135 Mechanitis *and 109* Melinaea *individuals.*

Nuclear phylogeny (left) based on 537,500 SNPs for Mechanitis *(A) and 784,526 SNPs for* Melinaea *(B) and full*

mitochondrial genome phylogeny (right). The coloured circles and connecting lines indicate the currently classified

species (25, 29)*. Br=Brazil; FG=French Guiana; E-Co=eastern Colombia; W-Co=western Colombia; E-Ec=eastern*

Ecuador; W-Ec=western Ecuador; E-Pe=eastern Peru; Pa=Panama, Su=Suriname. The coloured boxes highlight key

findings. The node labels show concordance factors, indicating the percentage of trees produced for windows across

the genome that contain that clade.

Figure 2 - Calibrated phylogeny of *Mechanitis and Melinaea* butterflies with evidence of introgression and biogeographic patterns.

Time-calibrated StarBEAST2 phylogenies of Mechanitis *(A) and* Melinaea *(B) with the newly proposed species*

classification and secondary calibrations from (11)*. One individual was included at the species level, or subspecies-*

level if they were very divergent. The regions in the overview map are based on the combined distribution of our

subspecies and samples from (10) *with region names adapted from* (11)*. Arrows between clades indicate potential*

hybridisation events (based on AIM, Fbranch and BPP). The node labels indicate the age as obtained by the

calibration. 'Core' in the Melinaea *phylogeny indicates the core clade of fast diverging* Melinaea, *and 'ingroup' is a*

clade referred to in the text. For each clade, cartoon wings based on representative colour patterns are shown, and

as an example, pink stars designate taxa part of the same mimicry ring. The distribution of our individuals is indicated

by coloured dots and coloured rings on a distribution map based on the subspecies distribution from (10)*); map from*

USGS, Esri, TANA, DeLorme and NPS; see Fig. S6-7 for larger maps). The chromosome numbers in the right column

are based on (48) *or our reference genomes (asterisk).*

Figure 3 - *Three ancestrally introgressed species, with a focus on* **Mechanitis nesaea**

A multispecies coalescent-with-introgression model explored the relation and timing of introgression relative to the

divergence times, in **A)** Mel. lilis*,* Mel. idae *and* Mel. marsaeus*;* **B)** Mec. menapis*,* Mec. messenoides*, and* Mec.

polymnia*; and* **C)** *the Brazilian* Mec. lysimnia lysimnia, Mec. nesaea *and* Mec. polymnia casabranca. **D)** *A closer look*

into the restored species Mec. nesaea*: phylogenetic relationships, chromosome numbers* (48)*, a photo of a*

representative adult and an early fifth instar larva (124, 125)*. The photo of* M. lysimnia lysimia *is a courtesy of*

Augusto Rosa. **E)** *Overlaid chromatograms of androconial extracts of representative individuals of* Mec. nesaea

(yellow line), Mec. l. lysimnia *(blue) and* Mec. polymnia casabranca *(orange). Peaks: (1) 4-Hydroxy-3,5,5-*

trimethylcyclohex-2-enone, (2) Hydroxydanaidal, (3) Methyl hydroxydanaidoate, (4) Methyl farnesoate isomer, (5)

Methyl (E,E)-farnesoate, (6) m/z *57, 43, 55, 56, 85, (7) Octadecatrienoic acid (cf.), (8) Octadecanoic acid, (9) Ethyl*

linolenate, (10) (E)-*Phytyl acetate (11) Hexacosene, (12) Heptacosene, (13) Nonacosene (not all compounds of Table*

S3 are found in these three individuals). **F)** *NMDS shows the androconial chemical bouquet of* Mec. nesaea *is clearly*

distinct from both putative parental lineages, most similar to Mec. polymnia. **G)** *A genome scan of f_{dM} across the*

genome (in 20kb windows) reveals that strong signatures of introgression (f_{dM}) between Mec. nesaea *and* Mec.

lysimnia *(P1=allopatric polymnia,P2=nesaea,P3=lysimnia,P4=Forbestra) overlaps with regions of high differentiation*

(F_{ST}) between Mec. nesaea *and its sister species* Mec. polymnia *(orange vertical lines - high F_{ST}, red dots - high f_{dM}*

and high F_{ST}). Chromosomal breakpoints between Mec. polymnia *and the four other reference genomes are shown*

with blue bars on top.

Figure 4 - Chromosomal rearrangements

A) Synteny between Melinaea *and* Mechanitis *genomes based on whole genome alignments. Horizontal bars represent individual chromosomes, with sex-chromsomes (black bar) and chromosomes involved in within-species polymorphic fusion-fissions (purple bar) highlighted. The cladogram is based on Fig. 2 and shows haploid chromosome numbers in parentheses. B) Example of differentiation (F_{ST}) and breakpoints (blue vertical lines) between* Mec. mazaeus *and* Mec. menapis *along the genome. The red dots indicate windows coinciding with breakpoints. C) Examples of HiC-contact maps. Top panel: Sex-chromosomes in* Mec. mazaeus. *Lower panel: Autosomal fission-fusion heterozygote in* Mel. mothone. *D) Matrix displaying number of fusion-fission rearrangements between species in each genus.*