Microbial single-cell omics in situ

Xihong Lan[1,2,3#], Qiaoxing Liang[1,2,3#], Jinhua He[4#], Jiayi Wu[1], Xiaoying Zhang[4], Fei

Li[1,2,3], Guoping Zhao[1,2], Ruidong Guo[1], Lili Li[1], Huijue Jia[1,2,3*]

[1]Greater Bay Area Institute of Precision Medicine (Guangzhou), Fudan University, Guangzhou, China.

[2]School of Life Sciences, Fudan University, Shanghai, China.

[3]State Key Laboratory of Genetic Engineering, Fudan University, Shanghai, China.

[4]Central Laboratory of Panyu Central Hospital, The Affiliated Panyu Central Hospital of Guangzhou Medical University, Guangzhou, China.

[#]These authors contributed equally to this work.

[*]Corresponding author. Email: huijue_jia@fudan.edu.cn

Abstract

Metagenomics has enabled understanding of the microbial composition and functional potential in various environments. Using Laser-Induced Forward Transfer (LIFT) technology, we report high-quality microbial single-cell genome or transcriptome in complex samples such as saliva, mouse gut and human tumor sections. Neighboring cells could be separately retrieved and bacteria could be fluorescently labelled. We illustrate cell-fate commitment with sporulation. The method is scalable and precise, and would empower fundamental insight for microbial populations and single-cell interactions with the host.

Introduction

Single-cell transcriptome analyses for mammalian cells are reaching their prime, further boosted by spatial omics technology[1]. Single-cell omics for bacterial cells, however, have remained a daunting task, due to the orders of magnitude lower concentration of nucleic acids, the lack of a poly(A) tail, and the Poisson distribution of droplet microfluidics [2]. Marine cyanobacteria were among the first bacteria to be analyzed using single-cell genomes [3], due to their dominance in the global ocean. Single-cell genomes of extreme environments have revealed genomic features of uncultured bacteria and archaea [4]. Other than low-diversity communities, microbial single-cell studies have mostly been illustrated for model organisms such as *E. coli*, and the genome or transcriptome analyzed, often using targeted methods, has remained far from complete [5-9].

Laser-Induced Forward Transfer (LIFT) is a straight-forward way of manipulating microparticles including cells [10-13]. With a laser beam that

evaporates or deforms a thin metal coating[14, 15], LIFT is ideally suited for picking

particles in the size range of bacteria. It can potentially tolerate food particles,

cell debris, and other objects commonly present in the human microbiome that

could undermine analyses using droplet microfluidics.

Metagenomic analyses of low-biomass environment such as tumors and

placenta has been costly and controversial [16]. Thus, a direct approach to

characterize the microbes in tissues with over 99.9% host sequences would

also be very much welcomed.

Here we present microbial single-cell genomes and transcriptomes using

a commercially available optics equipment, and demonstrate their superb utility

in a number of scenarios.

Results

Draft genome quality assembly for microbial single cells

Consisted of several key modules[17], our LIFT equipment was capable of image

capture, Raman spectral acquisition and single cell isolation from complex

samples after previous processing steps (Sup. Fig. 1a-d; Sup Movies S1). Next,

we explored the energy and mode required for microbial cell ejection using

bacteria of different sizes and shapes. Low energy resulted in cell ejection

failure including cell shifting and cell rupture (Sup. Fig. 2a and b), while

appropriate energy for cell ejection successfully isolated the selected cell

without affecting the closely adjacent cells (Sup. Fig. 2c). Further separation of

cells with different morphological characters suggested that appropriate energy

for cell LIFT was related to the size of the cell rather than its shape (Sup. Fig.

2d). For the larger bacteria, multiple LIFT attempts could also successfully

separate the cells without increasing energy (Sup. Fig. 2e).

To develop microbial single-cell technology for complex samples, we

started with saliva samples which are even more diverse than feces. A fresh

saliva sample was centrifuged and spotted onto the slide (Sup. Fig. 1c), and

bacteria cells were visualized before LIFT (Fig1.a). Single cells were ejected by

LIFT without affecting neighboring cells (Fig1.a and b, no burnt circle with the

low energy 60nJ) and Multiple Displacement Amplification (MDA) was

performed in a PCR tube. Without the need for sophisticated algorithms to pool

assembly from fragmental genomes, as was reported recently [5], here we used

state-of-the-art algorithms developed for metagenomes [18], and obtained a

bacterial genome of 99.99% completeness and low contamination 0.33%

(Fig1.c, Sup. Fig. 3a-d). This genome was identified as *Megamonas funiformis*.

Gene annotation through Bakta revealed the detection of coding sequences

(CDSs), tRNAs, rRNAs, ncRNAs, and CRISPRs (Fig1.d, Table S1). Calculating

the coding density showed that the genome sequence is >90% coding (Sup.

Fig. 3e and f), which is within the range for commensal bacteria (on average

87% for human gut [19]). Thus, LIFT from complex samples could obtain high-

quality single-cell bacteria genomes (single-cell assembled genomes, SAGs)

that far exceeds typical single-cell genomes [5] or metagenomically assembled

genomes [20], and are on-par with genomes from cultured isolates.

To validate the automatic process, *Sphingomonas sp000797515* labeled

with TADA were precisely ejected from sparse regions and densely packed

regions in mixture of several species, sequenced using two different reaction

volume, both of which yielded high-quality genomes (Sup. Fig. 4a-d; Sup

Movies S2). There were no significant differences in genome completeness,

contamination and coverage between the genomes obtained by the two

methods suggesting that downsizing the reaction volume does not lead to a

decline in SAGs quality (Sup. Fig. 4e). The single-cell genomes from LIFT were

also highly consistent (ANI>95%) with the known input genomes (Sup. Fig. 4f),

validating this approach.

To increase the throughput and eliminate individual differences in operation,

we made use of an automated pipetting platform to take over the molecular

biology work after automated single-cell ejection. With the aid of automatic

identification by adjusting parameters such as diameter, area and fluorescence

intensity, 34-62 bacteria from single strain suspensions and complex samples

could be precisely ejected within one minute for the next step of library

construction (Sup. Fig. 5a-c; Sup Movies S3-S4). Salivary microbes of three

fields from a single donor were marked for collection (0.1s per cell), which

consisted of diverse morphological characters (Fig2.a). After de novo assembly,

4 of the 39 SAGs have completeness >90 and contamination <5, which are

regarded as high quality [21]. 27 of the 39 SAGs have completeness >50 and

contamination <10, which are medium quality (Table S2). The species were

identified according to GTDB-Tk with ANI >95%. With this relatively small-scale

attempt, we already obtained oral bacteria from all major phyla, including

Actinobacteriota, *Bacteroidota*, *Campylobacterota*, *Firmicutes*, *Fusobacteriota*,

Patescibacteria and *Proteobacteria* (Fig.2b). Species from the oral cavity and

related body sites were seen, such as *TM7x*, *Haemophilus sp.*, *Neisserria sp.*,

Porphyromonas gingivalis, *Filifactor alocis*, *Fusobacterium peridonticum*,

Bifidobacterium vaginale (renamed from *Gardnerella vaginalis*) and

Parabacteroides merdae (Fig.2b). These results indicate that the single-cell

genomes could be reliably obtained with medium to high quality in an

automated and selective manner to reveal the microbial landscape in a complex

sample.

127 Cell-fate commitment revealed by single-cell transcriptome

Sporulation is a strategy for bacteria to survive harsh environments [22, 23], and

to potentially distribute to other hosts. We cultured a strain of *Bacillus*

licheniformis isolated from human saliva, and LIFT-picked single cells to

analyze their transcriptome following citric acid treatment (Sup. Fig. 6a).

Thousands of transcripts were detected from cells stimulated by citric acid (Sup.

Fig. 6b), which is consistent with widespread yet incomplete expression of the

bacterial genome per cell cycle, known for model organisms such as *E. coli* [24].

Besides protein coding genes, tRNAs and regulatory sRNAs were also detected

(Sup. Fig. 6c), favoring the robustness of the experimental protocol.

In addition to scRNA analysis according to the bacteria reference genome

(Sup. Fig. 6d-g), the high-quality reads obtained could also be de novo

assembled. Gene counts obtained by these two analytical methods were similar

(significance *p < 0.05*) and the expression profile was highly correlated (Sup.

Fig. 6h). Thus, both known and unknown bacteria could be investigated using

this scRNA approach.

Oligo(A) tails target bacterial mRNAs for degradation [25]; this coupling to

exonucleases in bacteria is evolutionarily similar to exosome degradation of

transcripts in the eukaryotic nucleus. We detected stretches of three or more

adenylates in a number of transcripts (Sup. Fig. 7a-f, Table S3). TctA (Tripartite

tricarboxylate transporter TctA family), a large transmembrane protein involved

in citrate uptake, for example, showed long stretches of poly(A) spanning the

entire gene, which led to three contigs in de novo assembly (Sup. Fig.7g and i).

The shorter form was more abundant in five of the five cells analyzed (Sup.

Fig.7h), consistent with degradation from 3' to 5'. These results highlight the

fundamental questions that can be addressed using high-quality microbial

single-cell transcriptome.

We next explored the dynamic transcription changes during the

commitment to sporulation in *B. licheniformis* following citric acid treatment. In

addition to the upright laser for LIFT, the commercially available equipment we

used also harbor an inverted laser for single-cell Raman spectrometry (Fig.3a).

Both the cell morphology and Raman spectra could be rapidly obtained from a

sample. During the citric acid-induced sporulation, the day 0, day 3 and day 9,

B. licheniformis cells can be visually and spectrally divided into three groups,

from day 0 rod (I), day 3 intermediate state (II) to day 9 oval spore (III) (Fig.3a

and b). The mean spectrum of cells I, II, III were different, and spectra from the

individual microbial cells formed three clusters (Fig.3c). Thus, the

morphological features of bacteria as well as Raman spectrum indicated

alterations in cellular gene expressions. Consistent with the decrease in growth,

while ribosomal RNAs, ribosomal proteins, and nucleoid associated proteins

dominated the day 0 transcriptome, the reads proportion of other genes notably

increased as the cells undergo morphological changes towards sporulation

(Sup. Fig. 8a). Comparison of the gene expression of cells I, II, III allowed us to

identify 194 differentially expressed genes, with these transcripts mainly

clustered into three groups according to variations in their reads counts (Fig.3d).

Gene ontology (GO) analysis of Biological Process (BP), Molecular Function

(MF), Cell Component (CC) showed that the genes preferentially expressed in

the intermediate state of group II are linked to the pathways such as cellular

response to stress, sporulation, hydrolase activity and intrinsic component of

membrane (Sup. Fig. 8b). Consistently, electron microscopy confirmed that

cellular component assembly occurred in cells II, which was also observed

before Raman microscopy and LIFT (Sup. Fig. 8c). Comparing cells I and cells

II, we found that germination gene *gerD* co-present with sporulation related

genes such as spore coat protein and *YjcZ* family sporulation protein were

upregulated after citric acid treatment at day 3 (Fig.3e). When cells I

transformed into spore, *spoIIIAC* (stage III sporulation protein AC), *YtvI*

(sporulation integral membrane protein) and spore-specific gene *splB* (spore

photoproduct lyase) were highly expressed (Fig.3f).

Further looking into the day 3 samples led to five types of cells with distinct

cell morphology, and the scRNA also formed five groups (Fig.4a). Principal

Component Analysis of gene expression profiles revealed that cells of type I,

type IV and type V were relatively clustered independently, while cells of type II

and type III were relatively dispersed (Fig.4b). Comparing gene counts in each

type, we found that cells of type II in cell division state were active in gene

transcription (Fig.4c). Genes whose expression were enriched in the different

cell types included enzymes, transcription regulator and biosynthesis proteins

(Fig.4d). Genes highly expressed in type I are FdhF/YdeP family

oxidoreductase, manA (mannose-6-phosphate isomerase, class I), PTS

mannose/fructose/sorbose transporter subunit IIC,

phosphoribosylaminoimidazolesuccinocarboxamide synthase, *ComGE* and 2-

dehydropantoate 2-reductase, which enable type I cells resistance to low PH

caused by citric acid, transporting sugar molecules, pantothenate biosynthesis,

DNA uptake and purine biosynthesis [26-30]. Genes highly expressed in type II

cells included DEAD/DEAH box helicase, primosomal protein DnaI, DNA

polymerase/3'-5' exonuclease PolX, DNA helicase RecQ, RNA-binding protein

and sigma-54-dependent Fis family transcriptional regulator, which indicated

DNA replication and gene transcription process in cell division [31, 32]. Genes

observed in type II were also expressed in type III, but relatively lower. When

cells changed into type IV, GTP associated genes MnmE, elongation factor Tu

and GTP 3',8-cyclase MoaA, 50S proteins and GntR family transcriptional

regulator were highly expressed, which regulated biological process such as

gene translation and cell motility [33, 34]. Genes in type V were mainly transporters,

Na+/H+ antiporter NhaC, permease and bacteriocin, which functioned in stress

resistance, substance exchange and inhibition of spore germination [35-37].

Comparing mean spectrum of five type cells, we observed Raman signal at

band 750nm-785nm contributed by Cytochrome c and metabolites of purine

degradation decreased (Fig.4e) [38, 39], which is consistent with variation of

metabolic and gene transcriptional activity in five type cells. In addition,

transcription of sporulation and spore related genes also varied with cell state

(Fig.4f). These results illustrate the discovery power of high-quality scRNA, and

indicate that visual and/or label-free Raman spectra-guided selection of

microbial single cells could greatly facilitate the investigation of heterogeneous

microbial populations. All cells could be rapidly scanned in seconds, before

more costly amplification and analyses.

Single-cell microbial genome and transcriptome from tissue-resident

microbiota *in situ*

Having demonstrated single-cell draft genomes from complex samples and

high-quality precision single-cell transcriptome from heterogenous populations,

we went on to directly study microbial single-cells in tissue samples, which has

been impossible with other methods. After gavage of mice with Cy3-amino-D-

alanine (Cy3ADA) and Fam-amino-D-alanine (FADA), various morphologies of

fluorescent bacteria were observed in intestinal lavage fluid, indicating that

intestinal bacteria can be specifically labeled with amino-D-alanine (Fig.5a).

Therefore, we further observed fluorescent bacteria near the mucus of a mouse

colon and inside host cells (Fig.5b). Single bacteria could be extracted from

these regions either in a fluorescence-dependent or -independent manner

(Fig.5c and d). We selected 25 bacteria from three fields for genome

sequencing, obtaining 3 high-quality and 5 medium-quality SAGs. Among these

bacteria, there are typical mouse gut bacteria like *Prevotella copri A*,

Escherichia flexneri, *Turicimonas muris*, *Bacteroides intestinalis*, *Bacteroides*

dorei as well as more aerobic and peritoneal species such as *Cutibacterium*

acnes, *Staphylococcus capitis* (Fig.5e)[40, 41]. These results suggested that LIFT

could be applied to tissue samples.

We next applied the method to tumor samples from human patients.

Resected tumor tissues from a patient diagnosed as colorectal cancer (Sup.

Fig. 9a) were cut into tissue sections and loaded onto the slide for microbial

single-cell omics (Fig.6a). We confirmed cancer infiltrating and staging by

immunohistochemical staining of marker genes such as P53, BrafV600E, Ki67

(Sup.Fig.9b). Using antibodies against bacterial lipopolysaccharide (LPS) and

vancomycin to detect Gram-negative and Gram-positive bacteria respectively,

we observed microbes with diverse morphological features including rod shape

and spherical shape, distributed in colorectal cancer (Sup. Fig. 9c). Without

special treatment, we searched for microbes in the bright field. The microbes

have distinct Raman spectral fingerprints compared to histiocytic Raman

spectrum (Sup. Fig. 9d), which means this technology can very effectively avoid

sequencing of host cells or collection of non-microbial particles. For this

colorectal cancer sample, a 1.33μm long and rod-shaped single cell was

spotted and separated for genomic sequencing combined with Raman

spectrum acquisition (Fig.6b and c). We obtained a medium-quality SAG

identified as *Bacteroides intestinalis* based on Mash analysis (Fig.6d).

Phylogenetic analysis of all *B. intestinalis* genomes from metadata-gut of Mash

database and our SAG revealed that SAG was similar to MGYG000028343

which is also from China (Fig.6e).

Moreover, a rod-shaped single cell adhered to intestinal mucosal was

separated for RNA sequencing (Fig.7a). After removal of host gene reads, we

obtained de novo transcriptome assembly identified as *Bacteroides*

xylanisolvens by GTDB-Tk, in which 70% contigs and 75% transcripts (2756

transcripts) belonged to *Bacteroides* (Fig.7b). Of note, we observed dust

particles in the lungs of this colorectal cancer patient with 50 years of smoking

when cancer metastasized to the lungs (Sup. Fig. 10a and Sup. Fig. 10b), which

is consistent with the finding that *Bacteroides xylanisolvens* is involved in

nicotine degradation in smokers [42]. Among the ~3616 genes expressed were

nicotine-degrading gene nicX, membrane proteins SusC/RagA family TonB-

linked outer membrane protein, ABC transporter, sialic acid-specific

acetylesterase and porin (Fig.7c), which are crucial for nicotine degradation,

nutrients uptake and thrive of *Bacteroidetes* in the gut [43-45]. Gene ontology

analysis further verified that these genes were enriched in pathways of passive

transmembrane transporter activity, sialic acid transmembrane transporter

activity and porin activity (Fig.7d). These results suggest the application

potential of this method for tissue microbe detection *in situ* and single cell spatial

microbiome.

Discussion

Our study provides a method based on LIFT technique for cell isolation and

further high-quality single cell sequencing. This method can precisely and

automatically isolate individual bacteria from complex samples, either in a label-

dependent or label-free manner. After rapid automated sorting, the cells were

transferred to an automated platform for nucleic acid extraction and

amplification, in preparation for subsequent single-cell genome or

transcriptome sequencing to enhance throughput. We are positive that further

reducing the reaction volumes, optimizing the reagents, the plasticware and the

precision selection process will further demonstrate the advantage of this

single-cell omics approach that does not rely on microfluidics and exhaustive

sequencing of abundant cells.

Knowing the complete single-cell genome and transcriptome would be very

useful to understanding fundamental questions of the microbiome beyond

model organisms. Both the single-cell genome and the transcriptome could be

either reference-based or de novo assembled, the latter means applicability to

any new sample instead of a model bacterium. As is commonly seen for single-

cell studies of mammalian cells, merely increasing the number of cells

sequenced by a few fold would not necessary ensure identification of rare populations. With our approach, rare populations could be enriched before sequencing. Moreover, our method provided multidimensional information for the study of single cell microbe, including bacterial shape, size, and Raman spectrum. Visually, the existing software could already be set to select for cell size, shape, area, etc (Sup. Fig.5b). Further standardization and algorithm development for the single-cell Raman spectra would allow more precision in sub-species analyses.

Another major advantage of this LIFT-based single-cell approach is that complex samples require minimum pretreatment, and can contain other particles or other cells. The separation of neighboring cells would also allow elucidation of microscopic cooperation in a native environment in future studies. We have previously reported multiple *Bacteroides* species in an Austrian cohort of colorectal adenoma and carcinoma [46, 47] and *B. fragilis* were typically studied in mice models of colitis [48]. Here the scRNA analyses have tentatively identified metabolism of nicotine, mucin, etc. We have not yet tried FFPE samples, which should be theoretically possible, so long as the nucleotides have not been too degraded. Further application of our microbial single-cell omics method to clinical samples would facilitate patient stratification and treatment.

In summary, here we present an optics-based strategy for precision single-cell microbial genomics, and transcriptomes that could fully capture the genomic landscape of host-associated microbiome, and allow nucleotide-resolution understanding of microbial gene transcription and turnover in individual cells in their native environment.

Methods

Animals

Specific pathogen-free C57BL/6 mice (female, 8 weeks old) were purchased from BESTEST Biotechnology Co., Ltd (Zhuhai, China) and maintained in the laboratory Animal Center of Greater Bay Area Institute of Precision Medicine (Guangzhou, China). These mice were bred in a temperature-controlled environment (25℃, 12-hour light/dark cycle) with free access to food and clean drinking water. All animal experiments were approved by the Animal Ethics Committee at Greater Bay Area Institute of Precision Medicine and were performed in accordance with guidelines approved by the Institutional Review Board of Fudan University.

Study participants and sample collection

All the adult volunteers were not known to have chronic diseases, and did not take antibiotics within the past three months. All healthy volunteers and the patient had written informed consent. All the samples were self-collected into a sterile tube, and handed in for further processing on the same day. The study was approved by the Institutional Review Board of Fudan University. Tissue collection was approved by the Ethics Committee at The Affiliated Panyu Central Hospital of Guangzhou Medical University and was conducted following guidelines established by this committee.

Laser Induced Forward Transfer (LIFT) of bacterial cells

The equipment, PRECI SCS-R300 (Hooke Instruments, Ltd., China), was turned on and adjusted according to standards following the manufacturer's

handbook [49] (Sup. Movie 1). The optical setup is shown in Sup. Fig. 1. Laser

used for LIFT was nanosecond single pulse laser with 532nm wavelength, 5ns

pulse width and 3.4μJ energy, which could produce a single adjustable spot

capable of ejecting 0.5-20μm microorganisms. For complex shape and larger

microbes (greater than 20μm), LIFT combined with beam shaping could

successfully sort them. Laser for Raman spectral acquisition was continuous

laser with a maximum laser power of 50mW, 532nm wavelength. The

parameters of the laser for the acquisition of Raman spectra are set to 3mW,

3s.

The filtered and washed samples were spotted onto a quartz slide coated with

25nm thickness of aluminum film (HSC24, Hooke Instruments, Ltd.), and air-

dried in the hood. The Hooke chip was designed to incorporate a thermal

insulation structure to effectively protect the cell from heat-induced damage.

Visually selected bacterial cells (the manufacturer's software) were LIFT-

transferred into collectors (HSR04, Hooke Instruments, Ltd.) using a laser

energy below 75nJ (exceeding the minimum energy required by most microbes),

and then flipped into individual PCR tubes with brief spinning at 2000g (Yooning

Mini-6K).

Whole genome amplification and sequencing from single bacterial cells

Following cell lysis at 65℃ for 10 min, Multiple displacement amplification (MDA)

was performed using the REPLI-g scDNA Polymerase, according to kit

instructions (REPLI-g single cell kit, #150345, QIAGEN). Sequencing libraries

(PE150) were constructed without PCR, and sequenced on a DNBSEQ-T7

high-throughput sequencer (MGI).

Genome assembly and analyses

Quality control on paired-end sequencing reads of raw data was performed by fastp (0.23.2) tool. The filtered reads were assembled by metaSPAdes (3.13.0) in single cell mode. Based on binning information provided by Kraken2 (2.0.7), the obtained assembled bins were processed with bin refinement using BinSpreader. The quality of genome bins was analysed by Checkm2 and QUAST. GTDB-Tk (1.0.2) was used to classify and taxonomically assign genome bins. Phylogenetic trees were constructed by PhyloPhlAn (3.0.67) tool.

Whole transcriptome amplification and sequencing from single bacterial cells

In addition to phosphate buffered saline (PBS) and lysis buffer according to the QIAGEN kit (#150065), cell lysis was performed at 24℃ for 5 min, in the presence of murine RNase Inhibitor (Vazyme), and MetaPolyzyme (MAC4L, Invitrogen). Total RNA Reverse-transcription, ligation and MDA were performed using the REPLI-g WTA single cell kit (#150065, QIAGEN), with extended reaction time for bacteria instead of mammalian cells: 2 hours for ligation, 4 hours for MDA. Sequencing libraries (PE150) were constructed without PCR, and sequenced on a DNBSEQ-T7 high-throughput sequencer (MGI).

Single-cell transcriptome analyses

For de novo transcriptome analysis, filtered reads of scRNA sequencing sample was assembled using rnaSPAdes (3.15.5)[50]. Bakta (1.8.1) conducts a comprehensive annotation of assembled transcriptome. Protein sequences

generated by Bakta (version 1.8.1; AMRFinderPlus for AMR gene annotations; a generalized protein sequence expert system for identity, distinct coverage and identity and priority values for each sequence) were further annotated by eggnog-mapper (5.0.2) for gene function. KofamScan is used to annotate protein sequences with KEGG Orthology (KO). DeepFRI is used to predict Enzyme Commission (EC) numbers and Gene Ontology (GO) terms for Bakta annotated genes. For analyzing the distribution and characteristics of oligoA, bioawk tool is used to analyze oligoA positions in gene/contig level, and then process the results to obtain matrix containing gene/contig lengths, oligoA positions, gene start and gene end positions. All transcripts in de novo transcriptome were evenly divided into ten percentile according to gene length and the number of occurrences of oligoA positions within each percentile range of the gene lengths was recorded. Standard deviation of oligoA distribution was calculated according to the fraction of counts in each percentile range of the gene length. The proportion of oligoA occurrence was normalized by R package "Mfuzz". The analysis of poly(A) with nine adenylate residues (9A) or more followed these steps. Bowtie2 is used to align reads to transcripts and calculate gene FPKM values. For genome-based analysis, reads are referred against *Bacillus licheniformis* genome and genome annotation GTF file (GCF_002074095.1) downloaded from NCBI database. After alignment with Bowtie2, gene quantification was performed using featureCount (subread2.0.3). Genes expression with fold change ≥1.5 and pvalue < 0.05 were determined as significant change using DESeq2 (version 1.20.0). Gene enrichment of Biological Process (BP), Molecular Function (MF), Cell Component (CC) was analysed by R package Clusterprofiler with pvalueCutoff = 0.05. Heatmaps of

gene expression were using R package "pheatmap". The bar plots, bubble plots

of GO analysis, scatterplots, and boxplots were visualized using on-line tool

imageGP (http://www.ehbio.com/ImageGP/index.php/Home/Index/index.html).

Bacillus licheniformis culture and transmission electron microscopy

The stored *Bacillus licheniformis* strain was inoculated into the Columbia Broth

medium overnight at 37°C. 100μl *Bacillus licheniformis* suspension was

transferred to 2ml medium containing 10% Columbia Broth, in which PH was

adjusted to 4.0 by adding citric acid. *Bacillus licheniformis* were cultured in citric

acid-amended 10% Columbia Broth for day 3 (exhibited a variety of

morphologies) and day 9 (almost all transformed into spore at day 9). Then

Bacillus licheniformis suspension was passed through 40μm cell strainers to

eliminate cell clumps. The resulting filtrate was centrifuged at 12,000g for 3

minutes to collect bacterial pellets. These pellets were then washed three times

with 1.5 ml of PBS each and finally resuspended in 0.7ml ddH2O. Prior to using

PRECI SCS R300 devices, 1-2 μl of the cell suspension was dropped onto chip

and allowed to dry either by evaporation at room temperature or by exposure

to sterile airflow in a biosafety cabinet.

For transmission electron microscopy, citric acid treated *Bacillus*

licheniformis were fixed with 4% PFA and stored at 4℃. The copper mesh was

incubated in 10μl fixed *Bacillus licheniformis* solution for 2 minutes. After

removal of excessive solution using filter paper, copper mesh was incubated

with 1% phosphotungstic acid (PTA) for 2 minutes and observed by electron

microscope (FEI Tecnai G2 spirit).

Metabolic labeling with fluorescent D-alanine in culture

TADA (tetramethylrhodamine-amino-D-alanine), Cy3ADA (Cyanine3-amino-D-alanine) and FamADA (5-Carboxyfluorescein-amino-D-alanine) were purchased from GuoPing Pharmaceutical (Anhui province, China). *Staphylococcus capitis*, *Streptococcus oralis*, *Lactobacillus plantarum*, *Akkermansia muciniphila* and *Clostridium butyricum* were isolated from faeces, urine and saliva using our LIFT technology and cultured in Gifu Anaerobic Medium at anaerobic incubator. The *Sphingomonas sp000797515* were generously provided by HOOKE Instruments Ltd. The *Sphingomonas sp000797515* were cultured in Luria-Bertani Medium and incubated with TADA probes (40mM) at 37℃ for 3 hours. After incubation, the *Sphingomonas sp000797515* were washed with PBS thrice, and mixed with other single-strain bacteria in ddH2O for the following analyses. The *Sphingomonas sp000797515* genome provided was assembled from our single strain suspension sequencing.

Metabolic labeling with fluorescent D-alanine in mice

Cy3ADA and FamADA were administered to mice via gavage to label mouse gut microbiota. In brief, Mice were gavaged with Cy3ADA and FamADA probes dissolved in 200µl of PBS at 1 mM concentration. 6 hours later, Mice were anesthetized using sodium pentobarbital (50 mg/kg). A part of mouse large intestine was excised and minced using scissors in 2ml of PBS buffer. Subsequently, the minced tissue was passed through 40µm cell strainers to eliminate nonbacterial tissues and food debris. The resulting filtrate was centrifuged at 12,000g for 3 minutes to collect bacterial pellets. These pellets

were then washed three times with 1.5ml of PBS each and finally resuspended

in ddH2O. The bacterial suspension was placed under a Leica fluorescence

microscope (ECLIPSE Ni-U) to observe the labeling of the bacteria.

Tissue pretreatment for LIFT cell sorting and Hematoxylin-Eosin staining,

Immunofluorescence staining, Gomoris staining

After euthanizing the mice, a small segment of colon tissue (0.5-0.8cm length)

free of food debris was excised, embedded in OCT (optimal cutting temperature

compound), and frozen. The frozen colon tissue was sectioned into slices with

6μm thickness and mounted flat onto glass slides and LIFT chip.

Several pieces of 2mm thickness tissues (20×20×2mm) separated by

laparoscopic colorectal cancer surgery were placed on filter paper and dry

gauze pad. After removal of excessive water, tissues were transferred to

supporter of freezing microtome (Leica CM1860 UV) and embedded with OCT.

In order to ensure the operation of the tissue section, OCT embedded tissue

was placed on cryo-stage until it's a solid ice cube. Processed tissue was

sectioned into 7μm thickness for Raman spectrometry and LIFT on PRECI

SCS-R300 (Hooke Instruments, Ltd., China) with 75nJ laser energy.

Dehydrated and paraffin embedded tissues were cut into 5μm thickness

and were stained with hematoxylin-eosin after de-paraffinization. For

immunofluorescence staining, tissue sections were treated with antigen

retrieval buffers, and then permeabilized and blocked by mixed solution of 0.3%

Triton X-100 and 5% BSA for 1 hour. Primary antibodies were used to incubate

tissue sections overnight at 4°C. Phosphate Buffer Saline containing 0.1%

Tween-20 (PBST) was used to wash tissue sections five times. Well-washed

sections were carefully incubated with secondary antibodies for 1 hour at room temperature. DAPI (4',6-diamidino-2-phenylindole) diluted at PBS with concentration 1:1000 was used to cell nuclei staining. Immunofluorescence images were captured by Leica sp8 system. Antibodies and their working concentration are listed as followed: anti-CK20 (ZSGB-BIO, #ZA-0574, 1:200), CDX2 (ZSGB-BIO, #ZA-0520, 1:200), Villin (ZSGB-BIO, #ZA-0575, 1:300), MSH2 (ZSGB-BIO, #ZA-0622, 1:250), MSH6 (ZSGB-BIO, #ZM-0367, 1:250), MLH1 (ZSGB-BIO, #ZM-0152, 1:300), PMS2 (ZSGB-BIO, #ZA-0542, 1:200), P53 (ZSGB-BIO, #ZM-0408, 1:200), BrafV600E (Roche, #760-5095, 1:300), and Ki67 (ZSGB-BIO, #ZM-0166, 1:300), anti-LPS (Cloud-Clone, #MAB526Ge22, 1:150), anti-Vancomycin (Invitrogen, #V34850, 1:150).

Gomoris staining method followed the manufacturer's protocol (Baso diagnostics. Inc. Zhuhai. China, #BA4375A). Briefly, 5μm thickness paraffin tissue sections were brought to water via xylene and descending grades of ethanol for deparaffinization. Following nuclear staining by Weigert's Iron Hematoxylin, sections were placed in Solution A for 15 minutes. After rinsing off the stain with distilled water, sections were stained with Solution B. After dehydration with ethanol and clearing with xylene, sections were mounted with a resinous medium.

References

1. Stahl, P.L. et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78–82 (2016).
2. Kuchina, A. et al. Microbial single-cell RNA sequencing by split-pool barcoding. *Science* **371** (2021).
3. Kashtan, N. et al. Single-cell genomics reveals hundreds of coexisting subpopulations in wild Prochlorococcus. *Science* **344**, 416–420 (2014).
4. Rinke, C. et al. Insights into the phylogeny and coding potential of microbial dark matter. *Nature* **499**, 431–437 (2013).
5. Zheng, W. et al. High-throughput, single-microbe genomics with strain

527   resolution, applied to a human gut microbiome. *Science* **376**, eabm1483 (2022).

528 6. Dar, D., Dar, N., Cai, L. & Newman, D.K. Spatial transcriptomics of planktonic
529   and sessile bacterial populations at single-cell resolution. *Science* **373**
530   (2021).

531 7. Pountain, A.W. et al. Transcription-replication interactions reveal bacterial
532   genome regulation. *Nature* **626**, 661-669 (2024).

533 8. Lan, F. et al. Massively parallel single-cell sequencing of diverse microbial
534   populations. *Nat Methods* **21**, 228-235 (2024).

535 9. Ma, P. et al. Bacterial droplet-based single-cell RNA-seq reveals antibiotic-
536   associated heterogeneous cellular states. *Cell* **186**, 877-891 e814 (2023).

537 10. Florian, C. & Serra, P. Printing via Laser-Induced Forward Transfer and the
538   Future of Digital Manufacturing. *Materials (Basel)* **16** (2023).

539 11. Barron, J.A., Krizman, D.B. & Ringeisen, B.R. Laser printing of single cells:
540   statistical analysis, cell viability, and stress. *Ann Biomed Eng* **33**, 121-130
541   (2005).

542 12. Karakaidos, P., Kryou, C., Simigdala, N., Klinakis, A. & Zergioti, I. Laser
543   Bioprinting of Cells Using UV and Visible Wavelengths: A Comparative DNA
544   Damage Study. *Bioengineering (Basel)* **9** (2022).

545 13. Kryou, C. & Zergioti, I. Laser-Induced Forward Transfer on Regenerative
546   Medicine Applications. *Biomedical Materials & Devices* **1**, 5-20 (2022).

547 14. Deng, Y., Renaud, P., Guo, Z., Huang, Z. & Chen, Y. Single cell isolation
548   process with laser induced forward transfer. *J Biol Eng* **11**, 2 (2017).

549 15. Ye, K. et al. Spatial transcriptomic profiling of isolated microregions in
550   tissue sections utilizing laser-induced forward transfer. *PLoS One* **19**,
551   e0305977 (2024).

552 16. Jia H. Investigating Human Diseases with the Microbiome. (Elsevier, ed. 1st,
553   2022, https://www.elsevier.com/books/investigating-human-diseases-with-the-
554   microbiome/jia/978-0-323-91369-0) (2022).

555 17. 梁鹏. 基于激光诱导向前转移的微生物单细胞弹射分选技术研究. 中国科学院大学(长
556   春光学精密机械与物理研究所). (2022).

557 18. Meyer, F. et al. Critical Assessment of Metagenome Interpretation: the second
558   round of challenges. *Nat Methods* **19**, 429-440 (2022).

559 19. Li, J. et al. An integrated catalog of reference genes in the human gut
560   microbiome. *Nat Biotechnol* **32**, 834-841 (2014).

561 20. Zhu, J. et al. Over 50,000 Metagenomically Assembled Draft Genomes for the
562   Human Oral Microbiome Reveal New Taxa. *Genomics Proteomics Bioinformatics* **20**,
563   246-259 (2022).

564 21. Bowers, R.M. et al. Minimum information about a single amplified genome (MISAG)
565   and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat*
566   *Biotechnol* **35**, 725-731 (2017).

567 22. Greene, R.A. & Slepecky, R.A. Minimal requirements for commitment to
568   sporulation in Bacillus megaterium. *J Bacteriol* **111**, 557-565 (1972).

569 23. Schultz, D., Wolynes, P.G., Ben Jacob, E. & Onuchic, J.N. Deciding fate in
570   adverse times: sporulation and competence in Bacillus subtilis. *Proc Natl*
571   *Acad Sci U S A* **106**, 21027-21034 (2009).

572 24. Macklin, D.N. et al. Simultaneous cross-evaluation of heterogeneous E. coli
573   datasets via mechanistic simulation. *Science* **369** (2020).

574 25. Bandyra, K.J. & Luisi, B.F. Licensing and due process in the turnover of
575   bacterial RNA. *RNA Biol* **10**, 627-635 (2013).

576 26. Iobbi-Nivol, C. & Leimkuhler, S. Molybdenum enzymes, their maturation and
577   molybdenum cofactor biosynthesis in Escherichia coli. *Biochim Biophys Acta*
578   **1827**, 1086-1101 (2013).

579 27. McCoy, J.G., Levin, E.J. & Zhou, M. Structural insight into the PTS sugar

transporter EIIC. *Biochim Biophys Acta* **1850**, 577-585 (2015).

28.   Price, D.R. & Wilson, A.C. A substrate ambiguous enzyme facilitates genome
reduction in an intracellular symbiont. *BMC Biol* **12**, 110 (2014).

29.   Mom, J., Chouikha, I., Valette, O., Pieulle, L. & Pelicic, V. Systematic
functional analysis of the Com pilus in Streptococcus sanguinis: a
minimalistic type 4 filament dedicated to DNA uptake in monoderm bacteria.
mBio **15**, e0266723 (2024).

30.   Levdikov, V.M. et al. The structure of SAICAR synthase: an enzyme in the de
novo pathway of purine nucleotide biosynthesis. *Structure* **6**, 363-376 (1998).

31.   Bruand, C., Farache, M., McGovern, S., Ehrlich, S.D. & Polard, P. DnaB, DnaD
and DnaI proteins are components of the Bacillus subtilis replication restart
primosome. *Mol Microbiol* **42**, 245-255 (2001).

32.   Shingler, V. Signal sensing by sigma 54-dependent regulators: derepression
as a control mechanism. *Mol Microbiol* **19**, 409-416 (1996).

33.   Ramakrishnan, V. Ribosome structure and the mechanism of translation. *Cell*
108, 557-572 (2002).

34.   Liu, G.F., Wang, X.X., Su, H.Z. & Lu, G.T. Progress on the GntR family
transcription regulators in bacteria. *Yi Chuan* **43**, 66-73 (2021).

35.   Jack, D.L., Yang, N.M. & Saier, M.H., Jr. The drug/metabolite transporter
superfamily. *Eur J Biochem* **268**, 3620-3639 (2001).

36.   Zhang, R. et al. A Novel MFS-MDR Transporter, MdrP, Employs D223 as a Key
Determinant in the Na(+) Translocation Coupled to Norfloxacin Efflux. *Front*
Microbiol **11**, 955 (2020).

37.   Egan, K. et al. Bacteriocins: Novel Solutions to Age Old Spore-Related
Problems? *Front Microbiol* **7**, 461 (2016).

38.   Premasiri, W.R. et al. The biochemical origins of the surface-enhanced Raman
spectra of bacteria: a metabolomics profiling by SERS. *Anal Bioanal Chem* **408**,
4631-4647 (2016).

39.   Kaila, V.R.I. & Wikstrom, M. Architecture of bacterial respiratory chains.
Nat Rev Microbiol **19**, 319-330 (2021).

40.   Xiao, L. et al. A catalog of the mouse gut metagenome. *Nat Biotechnol* **33**,
1103-1108 (2015).

41.   Liu, L. et al. Rotenone Induces Parkinsonism with Constipation Symptoms in
Mice by Disrupting the Gut Microecosystem, Inhibiting the PI3K-AKT Signaling
Pathway and Gastrointestinal Motility. *Int J Mol Sci* **26** (2025).

42.   Chen, B. et al. Gut bacteria alleviate smoking-related NASH by degrading gut
nicotine. *Nature* **610**, 562-568 (2022).

43.   Robinson, L.S., Lewis, W.G. & Lewis, A.L. The sialate O-acetylesterase EstA
from gut Bacteroidetes species enables sialidase-mediated cross-species
foraging of 9-O-acetylated sialoglycans. *J Biol Chem* **292**, 11861-11872 (2017).

44.   Papamichail, D. & Delihas, N. Outer membrane protein genes and their small
non-coding RNA regulator genes in Photorhabdus luminescens. *Biol Direct* **1**,
12 (2006).

45.   Hall, L.M. et al. Sequence diversity and antigenic variation at the rag locus
of Porphyromonas gingivalis. *Infect Immun* **73**, 4253-4262 (2005).

46.   Feng, Q. et al. Gut microbiome development along the colorectal adenoma-
carcinoma sequence. *Nat Commun* **6**, 6528 (2015).

47.   Wang, J. & Jia, H. Metagenome-wide association studies: fine-mining the
microbiome. *Nat Rev Microbiol* **14**, 508-522 (2016).

48.   Wong, C.C. & Yu, J. Gut microbiota in colorectal cancer development and
therapy. *Nat Rev Clin Oncol* **20**, 429-452 (2023).

49.   Instruments, H. PRECI SCS-R300 Raman Single Cell Sorter. (https://www.hooke-
instruments.com/ProductCenter/). (2022).

50. Bushmanova, E., Antipov, D., Lapidus, A. & Prjibelski, A.D. rnaSPAdes: a de novo transcriptome assembler and its application to RNA-Seq data. *Gigascience* **8** (2019).

Acknowledgements

The authors were grateful to Zhong Yang, Lili Li, Jiong Ma, Xu Zhang, Hang Li, Meiya Lin for equipment support. Ping Sun, Miaoyi Kuang for sequencing support. We thanked Prof. Hao Chen (Shanghai Institute of Materia Medica, Chinese Academy of Sciences) and Prof. Xin Feng (Jilin University) for their suggestions on mouse gut microbiota labeling.

Author contributions

Xihong Lan and Huijue Jia conceived and designed the methodology; Xihong Lan developed and performed the experiments with assistance from Jiayi Wu; Xihong Lan and Jiayi Wu optimized sequencing library preparation; Jinhua He and Xiaoying Zhang prepared tissue sections; Xihong Lan, Qiaoxing Liang and Ruidong Guo performed the sequencing data analysis; Xihong Lan wrote the original manuscript; Huijue Jia and Guoping Zhao revised the manuscript; Huijue Jia supervised the study.

Competing interests

H.J. was a shareholder of MGITech Inc. The remaining authors declare no competing interests.

Correspondence and requests for materials

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact: Huijue Jia (huijue_jia@fudan.edu.cn)

Figure captions

Figure.1 High-quality microbial single-cell draft genome obtained from saliva. (a) A view of the sample before and after ejection of the selected microbial cell. A single cell as indicated was selected for genomic sequencing. Scale bar 10μm. **(b)** Diagram illustrating the imaging by the inverted microscope and LIFT by the upright microscope, on the commercially available PRECI-SCS equipment. The collectors are placed on a wheel, whose rotation is controlled by the manufacturer's software (Multiple cells could also be transferred into the same collector, depending on the experimental design). **(c)** Visualization of single-cell assembled genome by CGView tool displaying GC skew, feature labels, GC content, divider rings, and tick density. **(d)** The pie chart of single-cell assembled genome annotation display different categories including coding sequence (CDS), CRISPR, ncRNA region, ncRNA, replication origin (oriC), rRNA, tetracycline resistance, tRNA with their respective proportions.

Figure.2 Automated microbial single-cell genomics. (a) Individual microbial cells of saliva distributed on the chip were observed before LIFT. The selected microbes numerically marked with variable morphology are ejected for single cell sequencing. Scale bar, 10μm. **(b)** A phylogeny constructed from 39 obtained SAGs genome sequence is represented by the inner dendrogram. The phylum of each SAGs is indicated by the branch color. Numbers for the tip nodes of the dendrogram are corresponding to numbers labeled in panel. Completeness and contamination of each SAGs are represented by two heatmaps overlay on the phylogenetic tree. The species name of each SAGs is listed in the outermost ring.

Figure.3 Coherent grouping of *B. licheniformis* cells visually, spectrally and with scRNA transcriptome. (a) Individual *B. licheniformis* cells I (treated without citric acid), II (treated with citric acid for 3 days), III (treated with citric acid for 9 days) distributed on the chip were observed in bright field before Raman spectrometry and LIFT. Scale bar, 10μm. **(b)** More than 50 cells from each group (Cells I, Cells II and Cells III) are selected for Raman spectra acquisition. Dimensionality reduction is performed on the obtained Raman spectra by UMAP. **(c)** The means and standard deviations of Raman spectra were calculated for each group. **(d)** *B. licheniformis* of morphological forms I (treated without citric acid), II (treated with citric acid for 3 days) and III (treated with citric acid for 3 days) were ejected (two cells for each group) for scRNA sequencing. Heatmap of differentially expressed transcripts in Cells I, Cells II and Cells III. The candidate transcripts are mainly clustered into three groups (left panel). Violin plot of normalized transcripts expression in Group I, Group II and Group III (right panel). Data are represented as mean ± SE (ns: $P >0.05$, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$) and calculated using ANOVA with post-hoc Tukey HSD Test. **(e)** Volcano plot of differentially expressed genes of Cells II

versus Cells I. Gene counts normalized and compared by DeSeq2. **(f)** Volcano
plot of differentially expressed genes of Cells III versus Cells I. Gene counts
normalized and compared by DeSeq2.

Figure.4 Single-cell transcriptome reveal heterogeneity in sporulation. (**a**)
Image of five type (four cells for each type) *B. licheniformis* cells in sample
treated with citric acid for 3 days and cells without citric acid treatment as control.
Scale bar, 1µm. (**b**) Principal Component Analysis of gene expression profile
from the five cell types. **(c)** Calculation of gene counts in each cell type. Data
are represented as mean ± SE (*P < 0.05, **P < 0.01, ***P < 0.001) and
calculated using two-tailed unpaired Student's t-test. **(d)** Heatmap of
representative gene expression (z-score of log-transformed values) from 20
individual *B. licheniformis* cells grown in day 3 citric acid-treated media. (**e**)
Raman spectra of five type *B. licheniformis* cells. The spectra are averages of
ten individual spectra and normalized to the intensity of the strongest feature in
each spectrum. **(f)** Dot plot of genes involved in *B. licheniformis* sporulation.
Gene expression and pvalue were calculated between the selected cell type
and the remaining cell type.

Figure.5 LIFT ejection of bacteria in mouse gut in bright and fluorescent
field. (a) Representative images of bacterial morphology in mouse large
intestinal lavage suspension after gavage with Cy3-ADA in mice. Scale bar,
10µm. **(b)** Cy3-ADA and Fam-ADA labeled bacteria on mouse large intestine
sections. Scale bar, 100µm. **(c)** Localized composite image of mouse large
intestinal tissue sections (upper panel). Scale bar, 200µm; LIFT ejection of
bacteria on mouse large intestine sections in bright field (lower panel). Scale
bar, 10µm. **(d)** LIFT ejection of bacteria on mouse large intestine sections in
fluorescent field. Scale bar, 10µm. **(e)** 25 single bacteria were isolated from
three fields of view in the mouse large intestinal tissue sections.   A phylogeny
constructed from 25 obtained SAGs genome is represented by the inner
dendrogram. The phylum of each SAGs is indicated by the branch color.
Completeness and contamination of each SAGs are represented by two
heatmaps overlay on the phylogenetic tree. The species name of each SAGs
is listed in the outermost ring.

Figure.6 Genomic sequencing for single microbe in tissue. (a) Diagram for
the colorectal cancer sample. **(b)** Tissue section containing the microbial single-
cell ejected and genome sequenced. Scale bar, 10µm. **(c)** Raman spectrum of
the indicated single cell. **(d)** Visualization of obtained SAG (bin Bacteroides) by
CGView tool displaying GC skew, feature labels, GC content, divider rings, and
tick density. **(e)** Phylogenetic tree was constructed by obtained SAG (bin
Bacteroides) and all *Bacteroides intestinalis* genomes downloaded from
metadata-gut of Mash database. Among Bacteroides intestinalis genomes from
Mash database, genome *MGYG000028343* was isolated from China.

Figure.7 Transcriptome sequencing for single microbes from tissue. (a)
Tissue section as in Figure 6, with inlets indicating the position of the bacteria
for scRNA analyses. Scale bar, 10μm. **(b)** Count ratio of contigs, transcripts in
single cell assembled transcriptome and genes by genome-based analysis.
Contig classification was annotated by Kraken2. Transcript annotation was
referred against de novo transcriptome annotated by eggnog-mapper software.
Gene annotation was referred against *Bacteroides xylanisolvens* genome (total
5306 genes). **(c)** The detection of nicotine degradation gene nicX and proteins
located on cell membrane. Gene FPKM was calculated by Bakta. **(d)** Gene
ontology analysis of transcripts classified to *Bacteroidetes*.

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6

a
Colorectal cancer

tumor site

tumor section

mucosal surfaces and bacteria

b
Tiusse bright field image

Enlarge

c
Raman spectrum of selected bacteria

d
bin_Bacteroides

- CDS
- tRNA
- rRNA
- GC content
- Other
- GC skew+
- GC skew-

Bacteroides intestinalis

Genome shotgun sequence

e
clade N=68

clade N=36

clade N=21

clade N=3

clade N=21

clade N=8

Fig.7

a

Tiusse bright field image   Enlarge

b

c

d