Spectral algal fingerprinting and long sequencing in synthetic algal-microbial communities

Ayagoz Meirkhanova[1], Sabina Marks[2], Nicole Feja[2], Ivan A. Vorobjev[1,3], Natasha S. Barteneva[*,1,4]

[1]School of Science and Humanities, Nazarbayev University, Astana 010000, Kazakhstan

[2]Faculty of Biology, University of Duisburg-Essen, Campus Essen, 45141 Essen, Germany

[3]National Laboratory Astana, Nazarbayev University, Astana 010000, Kazakhstan

[4]The Environmental Research and Efficiency Cluster, Nazarbayev University, Astana 010000, Kazakhstan

Abstract

1. Synthetic biology has made progress in creating artificial microbial and algal communities, but technical and evolutionary complexities still pose significant challenges.

2. Traditional methods for studying microbial and algal communities, such as microscopy and pigment analysis, are limited in throughput and resolution. In contrast, advancements in full-spectrum cytometry enabled high-throughput, multidimensional analysis of single cells based on their size, complexity, and spectral fingerprints, offering more precise and comprehensive analysis than conventional flow cytometry.

3. This study demonstrates the use of full-spectrum cytometry for analyzing synthetic algal-microbial communities, facilitating rapid species identification and enumeration. The workflow involves recording individual spectral signatures from monocultures, utilizing autofluorescence to distinguish them from noise, and subsequent creation of a spectral library for further analysis. The obtained library is used then to analyze mixtures of unicellular cyanobacteria and synthetic phytoplankton communities, revealing differences in spectral signatures. The synthetic consortium experiment monitored algal growth, comparing results from different instruments and highlighting the advantages of the spectral virtual filter system for precise population separation and abundance tracking. This approach demonstrated higher flexibility and accuracy in analyzing multi-component algal-microbial assemblages and tracking temporal changes in community composition.

4. By capturing the complete emission spectrum of each cell, this method enhances the understanding of algal-microbial community dynamics and responses to environmental stressors. With development of standardized spectral libraries, our work demonstrates an improved characterization of algal communities, advancing research in synthetic biology and phytoplankton ecology.

Keywords: synthetic algal-microbial communities; spectral flow cytometry; imaging flow cytometry; long sequencing; nanopore-based sequencing

Introduction

In recent decades, synthetic biology has made significant progress in building artificial microbial and algal communities (Deng et al., 2022; Fu et al., 2020); however, the difficulty of engineering complex biological systems remains significant due to technical hurdles and the intricate nature of biosystems, which evolve and refine themselves over time (Hartwell et al., 1999). Community selection experiments address fundamental questions in ecology and evolution, as well as applied biotechnological issues, yet constructing more complex stable communities to study the effects of different bacteria on community dynamics remains challenging. Some of the challenges include the fact that uni-algal systems are not stable, and bacterial contamination is likely if the algal community is maintained heterotrophically on a medium with a fixed source of carbon (Andersen, 2005) or in the presence of dead cells. Several studies have demonstrated that algal-microbial synthetic consortia, which exhibit mutualism, can be engineered (De-Bashan et al., 2002; Cho et al., 2015; Xu et al., 2016). However, algal-microbial consortia development is usually limited by studying a few algae (often, systems are over-dominated by certain algae) or only the bacterial part of the consortia, with some exceptions (Segev et al., 2016; Kolter, 2024). Moreover, long-term homeostasis of algal-microbial consortia could be difficult to maintain since evolutionary constraints are known to limit the long-term stability of synthetic systems (Castle et al., 2021), and behavior and composition of engineered communities is "unpredictable" (Brenner et al., 2008).

Finally, the absence of methods allowing to follow the dynamics of mixed algal-microbial systems further complicates the development of complex synthetic systems. Identification and characterization of individual microalga and bacteria in the consortium are essential for engineering such consortia and development of subsequent applications (Perera et al., 2019). While microbial community composition analysis has advanced significantly, including classification of reads up to the species level (Curry et al., 2022), effective methods for plankton identification and characterization are still limited. Traditionally, microbiome analysis in such systems was conducted at the genus and class levels due to limitations in next-generation sequencing. However, with the development of long sequencing techniques, more detailed analyses have become possible. For instance, in a long-term mesocosm experiment, species-level visualization using imaging cytometry combined with long sequencing was employed (Meirkhanova et al., 2024). The integration of a full-spectrum approach would, therefore, provide a distinct advantage by enabling more precise and comprehensive analysis.

An algal community can be categorized into subgroups based on morphology, size, cellular functions, and interactive relationships. However, morphology-based taxonomy has limitations, particularly in distinguishing cell types with similar morphologies. Traditional methods for studying microbial and algal communities, such as microscopy and pigment analysis, are constrained by throughput, taxonomic resolution, and functional characterization, primarily differentiating specific taxonomic groups. Spectrofluorimetry, spectroscopy, and HPLC-based approaches provide averaged integrated data at a population level (Poryvkina et al., 1994; Falkowsky, Oliver, 2007; Roy et al., 2011; Swan et al., 2016; Liu et al., 2021). The maximum of taxonomic groups identified by HPLC- pigments analysis typically limited by 4-7 distinct groups (Catlett, Siegel, 2018; Kraemer, Siegel, 2019). Though full-resolution reflectance spectra are used in discriminant and classification analyses of algae by airborne hyperspectral sensors, applying spectral unmixing algorithms based on reference data (Hochberg, Atkinson, 2003; Rodriguez et al., 2017; Kramer, Siegel, 2019; Cause-Nicholson et al., 2021), there is still a need for adequate *in situ* method. Conventional flow cytometry struggles to distinguish cells from other particulate matter in the water matrix, and imaging flow cytometry can differentiate only a few fluorescent channels. In contrast, full-spectrum cytometry has recently transformed the field by offering high-throughput, multidimensional analysis of single cells based on their size, complexity, and spectral fingerprint (Futamura et al., 2015; Robinson, 2022; Dott et al., 2024).

This study introduces full-spectrum cytometry as a powerful tool for analyzing synthetic algal-microbial communities, facilitating rapid, high-throughput analysis from species identification to enumeration without extensive sample preparation. By capturing the complete emission spectrum of each cell, full-spectrum cytometry provides valuable insights into phytoplankton community composition, structure, and dynamics, as well as their responses to environmental stressors that influence community structure (Barteneva et al., 2019, 2023; Jeppesen et al., 2005). Algal cells, distinguished by autofluorescent spectral signatures, are separated taxonomically, while light scattering parameters assess cell size and complexity, distinguishing target events from debris. Spectral cytometry identifies highly autofluorescent subpopulations using virtual filtering (Barteneva et al., 2019) or only autofluorescence finder algorithm (Wanner et al., 2022). Characterizing microbiota associated with algae from in-house cultures or natural consortia is crucial as microalgae growth phases correlate with shifts in microbial phylotypes (Geng et al., 2016), although microbial dynamics in synthetic systems remain understudied. This research demonstrates that modern spectral cytometry, coupled with species-level 16S rRNA sequencing, effectively identifies and enumerates algae based on their spectral signatures, advancing our understanding of phytoplankton ecology, synthetic biology, and evolution.

Methods

Sample preparation and instrument setup

Algal cultures (**Suppl. Table 1**) were obtained from Culture Collection of Algae at the University of Göttingen (Germany) and Central Collection of Algal Cultures at the University of Duisburg-Essen (Germany). Cultures were maintained in Basal Medium (BM) at 21°C, under 12/12 L/D cycle in algae growth chamber AL-30L2 (Percival, USA). Samples were recorded in parallel on three instruments: 7-laser ID7000 Spectral Cell Analyzer (Sony Biotechnology, USA), 5 laser BD FACSAria Special Order cell sorter (BD Biosciences, USA) and 4-lasers ImageStreamX MarkII (Amnis - Cytek, USA) imaging flow cytometer equipped with 40X and 60X objectives. Samples prepared for cytometric analysis included mono- and mixed cultures and were recorded using a standard 24-tube rack. Each sample was placed in round-bottom 5mL tubes, with a minimum volume of 250μL, and arranged in specified positions on the rack. Prior to data acquisition, instrument settings were adjusted and set in accordance with each sample (FSC and SSC gains depending on algal culture, threshold value – 11%, PMT voltage ranging from 40% to 70%). Samples were recorded at minimum flow rate (=1) and 50,000 events were set for data acquisition. FSC_A vs SSC_A plots were used for the initial assessment of instrument settings in the "Preview" mode prior to recording.

Next, synthetic consortium was set up and included representatives of three major phytoplankton groups – Cyanobacteria (*Microcystis* sp., *Synechococcus* sp., *Synechocystis* sp.), Chlorophyta (*Chlamydomonas* sp., *Scherffelia* sp., *Haematococcus pluvialis*) and Cryptophyta (*Rhodomonas* sp., *Cryptomonas* sp., *Chroomonas* sp.) – with size ranging from 2 to 25 µm (**Table 1, Suppl. Fig. 1**), growth of which was monitored daily for the period of 9 days. Triplicates were also analyzed in parallel on BD FACSAria cell sorter (BD Biosciences, USA) and ImageStreamX MarkII (Amnis - Cytek, USA) imaging flow cytometer. Microbial community composition during the last day of the experiment was assessed using nanopore-based full-length 16S rRNA gene sequencing.

Table 1. List of algal cultures used for recordings during artificial mix experiment and library construction using ID7000 spectral cell analyzer.

Phylum	Order	Species	Habitat	Strain number
Cyanobacteria	Nostocales	*Dolichospermum urugayensis*	Freshwater	2498
		Nodularia sphaerocarpa	Freshwater	50.79
	Gloeobacterales	*Gloeobacter violaceus*	Freshwater	35.87
	Chroococcales	*Microcystis* sp.	Freshwater	CCAC 7146B
	Synechococcales	*Synechococcus* sp.	Freshwater	CCAC 2944B
		Synechocystis sp.	Freshwater	CCAC 4059B
Chlorophyta	Chlorodendrales	*Scherffelia dubia*	Freshwater	CCAC 1398 B
	Chlamydomonadales	*Chlamydomonas* sp.	Freshwater	CCAC 4527 B
		Haematococcus pluvialis	Freshwater	CCAC 2072B
	Chlorellales	*Chlorella vulgaris*	Freshwater	211-12
	Sphaeropleales	*Acutodesmus obliquus*	Freshwater	22.81
Cryptophyta	Pyrenomonadales	*Chroomonas* sp.	Freshwater	CCAC 2614B
		Rhodomonas sp.	Freshwater	CCAC 1479B
	Cryptomonadales	*Cryptomonas* sp.	Freshwater	CCAC 2345B
Rhodophyta	Stylonematales	*Chroothece richteriana*	Marine	104.79
	Porphyridiales	*Porphyridium cruentum cf.*	Marine	CCAC 3771 B
Dinophyta	Peridiniales	*Peridinium* sp.	Freshwater	CCAC 1612 B
Euglenophyta	Euglenida	*Euglena sanguinea*	Freshwater	CCAC 3518 B
Bacillariophyta	Fragilariales	*Fragilaria* sp.	Freshwater	CCAC 5509 B

Autofluorescence Finder Tool and Virtual Filters setup

The ID7000 software's Autofluorescence Finder Tool was used to analyze the algae samples, treating autofluorescence as an independent fluorescent parameter. This feature is crucial for the taxonomic identification of different algae. By incorporating excitation lasers ranging from 320nm to 808nm, we included unique emission regions in the algal spectral signature (fluorescence intensity and shape), distinct from the chlorophyll peak and highly variable among different algae. This setup allowed for the detection of individual autofluorescent spectral signatures for each of the recorded algal monocultures. Following the identification of the best separation between autofluorescent populations, subpopulations of interest were gated and examined for their spectral signatures. Each population demonstrated a unique autofluorescence spectrum. Furthermore, Autofluorescence Finder tool was used for virtual filters setup. In a similar manner, highly variable regions between different algal taxonomic groups are defined and used to set a pair of virtual filters (VFs). Defined VFs are next used to separate subpopulations of interest within multi-component mixes.

Data analysis

Initial cytometric data analysis and population identification were performed using the ID7000 software. Identification of subpopulations within an artificial mix involved consecutive gating steps using a combination of 2D plots with specified virtual filters for best separation. In addition, t-Distributed Stochastic Neighbor Embedding, otherwise known as t-SNE, was done using FCS Express

(De Novo Software, USA). A modified version of t-SNE - opt-SNE, a tool allowing the mapping of
high-dimensional cytometry data onto a two-dimensional plot while conserving the original high-
dimensional structure, was used to independently identify subpopulations within the artificial mix,
minimizing gating bias. Obtained clusters were then used to compare data acquired on different
cytometers. To display fluorescence intensity dispersion, coefficient of variation was calculated and
plotted across all detection channels.

Sequencing library preparation and bioinformatic analysis
Library preparation, the sequencing run on the MinION Mk1C device and bioinformatic analysis were
performed in laboratory following the end of 9-day synthetic consortium experiment. Firstly, DNA
was extracted using the DNEasy Power Water Kit (Qiagen, USA) according to the manufacturer's
instructions, with an additional heating lysis step. The full-length 16S gene was then amplified using
the universal 16S primer pair 27F (5′-AGAGTTTGATCCTGGCTCAG-3′) and 1492R (5′-
TACGGYTACCTTGTTACGACTT-3′). Library preparation followed the manufacturer's protocol
(ONT, UK). The PCR reaction mixture included nuclease-free water, input DNA, LongAmp Hot Start
Taq 2X Master Mix (NEB, USA), and the respective 16S barcode primer. PCR products were next
purified using AMPure XP beads (Beckman Coulter, USA). In the final library preparation step,
sequencing adapters were added to the barcoded samples. Raw data underwent basecalling using the
guppy basecaller (ONT, UK). The resulting FASTQ files were processed using Python algorithms
beginning with read assessment, followed by quality (Q-score>10) and read length filtering. The next
steps involved adapter sequence removal and demultiplexing of reads into their respective barcodes.
The demultiplexed reads were then classified using the Emu taxonomic abundance estimator (Curry
et al., 2022) along with a custom reference database (Len et al., 2023) up to the species level
(https://gitlab.com/treangenlab/emu). Lastly, the data were rarefied prior to further analysis.

224 Results

The first step in the analysis of complex communities requires construction of a library with unique
spectral signatures for different phytoplankton groups. For this purpose, single phytoplankton cultures
were recorded first on ID7000 Spectral cell analyzer (Sony Biotechnology), and individual spectral
signatures were then extracted. **Fig. 1** demonstrates the general workflow for the analysis of
Microcystis sp. monoculture as an example. During the recording, emission spectra across 7 lasers
were obtained, from algal cultures with presence of background noise/cellular debris (**Fig. 1a**).

Autofluorescence was then utilized in order to discriminate the spectra of interest from the debri/noise.

In some cases, direct gating based on light scattering is possible as well, however, for picoplankton

(e.g., *Synechococcus*), with cell size ranging from 0.2 to 2 µm, the gating is not precise. Instead,

intrinsic cell fluorescence was used to distinguish events of interest from low-intensity/no-intensity

events. With "Autofluorescence Finder" tool clear separation between the two autofluorescent

populations was observed across all lasers, except for 808 nm, given the optimal set of virtual filters

(VFs). **Fig. 1b,c** demonstrates pairwise density plots of [VF-320]_A against VFs of 6 remaining

excitation lasers as an example, with detection of two populations: AF-A – *Microcystis* sp. – 98.9%

and AF-B – debris – 1.06% ([VF-320]_A vs. [VF-561]_A). In this case, filters were set around channel

20, where emission intensity was the highest. Unique spectral signature for *Microcystis* sp. was then

extracted by plotting the emission of autofluorescent population AF-A on a ribbon plot (**Fig. 1d**). The

same workflow was applied for the acquisition of spectral signatures of the remaining algal

monocultures (**Figure 1e**), resulting in a spectral library used for further analysis.

244
Figure 1. Schematic overview of the proposed experimental workflow for the analysis combining full-spectrum cytometry and sequencing. The workflow involves parallel analysis of obtained samples using MinION Mk1C-based sequencing platform (ONT) and ID7000 Spectral cell analyzer (Sony Biotechnology). The first step in the analysis of spectral data requires **(a)** recording and isolation of single spectral signatures, for which "Autofluorescence Finder" tool is utilized. During this step **(b)** a set of optimal virtual filters for discrimination of autofluorescent populations is determined. As determined using the tool, **AF-A** corresponds to *Microcystis* sp. and **AF-B** to debris

252 within the sample; **(c)** respective emission spectra of two defined autofluorescent populations
253 displayed below (red line – **AF-A**, blue line – **AF-B**). **(d)** Single spectral signature is then extracted
254 and **(e)** a library, containing unique spectral signatures corresponding to respective phytoplankton
255 species, is then constructed and utilized during subsequent analysis, including unsupervised
256 clustering techniques (e.g., t-SNE).
257
To discriminate multiple components in a suspension, a mixture of unicellular cyanobacteria was used

and recorded for further analysis (**Fig. 2**). In addition, components of the mix represented variable

taxonomic groups at the Order level (*Microcystis* sp., belonging to Chroococcales), as well as

members of the same Order (*Synechococcus* sp., *Synechocystis* sp., belonging to Synechococcales).

Individual spectral signatures were recorded prior to the mix; **Fig. 2b** demonstrates overlay plot of the

acquired spectra across 7 excitation lasers. As seen from the plot, *Microcystis* sp. (red) and

Synechococcus sp. (blue) possess nearly identical spectral signatures, with slight differences in

fluorescence intensity, while distinct region around channels 10-16 (566-646 nm) in the spectra for

Synechocystis sp. (violet) was recorded. Based on the obtained information, set of optimal VFs was

chosen to firstly discriminate autofluorescent populations from unwanted events, and to further

separate these populations into respective picoplanktonic groups. Gating strategy is presented in **Fig**

2c, where a combination of [VF-647]_A vs [VF-488]_A reveals two subpopulations, corresponding

to: *Synechocystis* sp. and population "A". Due to the similarity in the spectra of *Microcystis* sp. and

Synechococcus sp. subpopulation "A" was then differentiated based on SSC values. Identified

populations could not be discriminated based on FSC and SSC only (**Fig. 2d**). At this point, the

maximum number of individual phytoplankton species that could be discriminated reached 14 (**Fig.**

3).

Figure 2. Discrimination of cyanobacteria in a synthetic mix based on autofluorescence. **(a)** Ribbon plot demonstrating combined spectral signatures of three cyanobacterial representatives across 7 excitation lasers: *Synechococcus* sp., *Synechocystis* sp., and *Microcystis* sp. **(b)** Overlay plot of individual spectral signatures for each cyanobacterial representative (*Synechococcus* sp. - blue, *Synechocystis* sp. - violet, and *Microcystis* sp. - red); red regions indicate regions of interest, used for setting an optimal pair of VFs. **(c)** Dot plot of **[VF-637]**_A against **[VF-488]**_A with two autofluorescent populations identified: *Synechocystis* sp., population **A** was then plotted with **[VF-561]**_A against **SSC**_A to resolve two subpopulations *of Microcystis* sp. and *Synechococcus* sp. **(d)** Dot plot demonstrating relative position of each of the identified cyanobacterial species on FSC_A against SSC_A plot.

287
288 **Figure 3.** Gating strategy for discrimination of 14 groups of phytoplankton species based on
289 autofluorescence in a complex synthetic mix. **(a)** Ribbon plot demonstrating combined spectral

signatures of 16 phytoplankton species across 7 excitation lasers. **(b)** Overlay plot of individual spectral signatures for each phytoplankton representative. (c) Gating strategy for discrimination of each subpopulation within the synthetic mix: 1 – Chlorophyta (*Chlorella vulgaris* and *Acutodesmus obliquus*), 2 – *Euglena sanguinea*, 3 – *Peridinium* sp., 4 – *Cryptomonas* sp., 5 – *Fragilaria* sp., 6 – *Haematococcus pluvialis*, 7 – *Chroothece richteriana*, 8 – *Porphyridium cruentum* cf., 9 – filamentous Cyanobacteria (*Dolichospermum urugayensis* and *Nodularia sphaerocarpa*), 10 – *Chroomonas* sp., 11 – *Synechocystis* sp., 12 – *Gloeobacter violaceus*, 13 – *Microcystis* sp., 14 – *Synechococcus* sp.

Next, applicability of full-spectrum cytometry in monitoring of dynamics of synthetic phytoplankton consortium was assessed. For this purpose, an artificial mix experiment was set up; synthetic mix included 3 representatives of major phytoplankton groups – Cyanobacteria, Chlorophyta and Cryptophyta. Firstly, a library, containing individual spectral signatures covering each phytoplankton species within the mix was constructed. **Fig. 4a** represents 9 spectral signatures across three phytoplankton groups: Cyanobacteria (*Microcystis* sp., *Synechococcus* sp., *Synechocystis* sp.), Chlorophyta (*Chlamydomonas* sp., *Scherffelia* sp., *Haematococcus pluvialis*) and Cryptophyta (*Rhodomonas* sp., *Cryptomonas* sp., *Chroomonas* sp.). As seen from the graph, regions unique for each of the group are present, specifically between channels 10-15 (566-646nm) of the spectra. This information was further utilized to define a set of optimal VF pairs to discriminate these populations in a complex mix. VFs were defined in a way to cluster subpopulations within the same group, while maintaining the difference between the groups. Set of filters [VF-320]_A vs [VF-561]_A (**Fig. 4b**) was chosen as an optimal pair for the first step within the gating strategy (**Fig. 4c**), where three groups are identified and discriminated from the rest of the unwanted events (background noise/cellular debris). In addition to cytometric analysis, samples collected at the end of the experiment were also subjected to sequencing-based analysis**.**

Fig. 4d summarizes averaged changes in the relative abundance of the components of the artificial mix throughout the experiment. Cryptophytes, specifically *Chroomonas* sp., were recorded to be the dominant group within the mix starting from the fourth day, outnumbering cyanobacterial representatives. In addition to cytometric analysis, samples collected at the end of the experiment were also subjected to sequencing-based analysis. **Fig. 4e** demonstrates microbial community composition across three replicates at species levels. Three most abundant phyla besides Cyanobacteria were Proteobacteria and Bacteroidetes, with *Polaromonas sp. JS666* being the most abundant bacterial species. The results showed stability of bacterial composition in artificial algal-microbial mix during 9-days experiment resulting in unsignificant differences between sequenced triplicates.

326
327

328
329

Figure 4. Tracking temporal changes in synthetic community composition. **(a)** Overlay plot of individual spectral signatures for each phytoplankton group representative (Cyanobacteria: violet – *Microcystis* sp., red – *Synechococcus* sp., blue – *Synechocystis* sp.; Chlorophyta: yellow – *Scherffelia dubia*, orange – *Chlamydomonas* sp., green – *Haematococcus pluvialis*; Cryptophyta: pink – *Chroomonas* sp., blue – *Rhodomonas* sp., green – *Cryptomonas* sp.); red regions indicate regions of interest, used for setting an optimal pair of VFs. **(b)** "Autofluorescence Finder" tool with VFs set to regions of interest, demonstrating the best separation of subpopulations on [VF-320_A] vs [VF-561_A] plot. **(c)** Gating strategy, composed of sequential sub-gating steps, used to identify phytoplankton subpopulations within the synthetic community. **(d)** Changes in relative abundance of the artificial mix subpopulations during the nine-day experiment evaluated with ID7000 spectral cytometer. **(e)** Microbial community composition during the last day of the experiment in triplicates at species level revealed through full-length 16S rRNA sequencing.

The next steps involved pairwise combinations of defined filters to discriminate the components of each of the subpopulations. As mentioned previously, due to the high similarity of spectra between *Microcystis* sp. and *Synechococcus* sp., SSC was used to differentiate these populations. A comparison of the data obtained on three instruments revealed limitations of some of the instruments (**Fig. 5**). Opt-SNE dimensionality reduction algorithm was utilized to minimize "gating bias" and group data into clusters with similar characteristics in low-dimensional space. Three datasets were compared: synthetic mix data obtained from ID7000 instrument using a set of pre-defined VFs and a set of VFs matching the filters used on FACSAria, and lastly, data obtained from FACSAria. Both sets of filters utilized on the ID7000 resulted in clear separation of all 9 subpopulations within the mix, while a single population of *Scherffelia* sp. was not possible to discriminate from the rest on the FACSAria instrument. These results demonstrate the higher flexibility a virtual filter system offers in the analysis of multi-component mixes. In addition to more precise population separation obtained with the ID7000 instrument, it was possible to track the abundance of nearly all mix components, as demonstrated in **Suppl. Fig. 2**. Overlay of temporal, spectral signatures recorded for the period of 9 days for each mix member was used to assess the extent of signal intensity dispersion. As revealed through the coefficient of variation, regions most susceptible to differences in intensity generally included channels 1-13 (413.6-612.6 nm) and 30-34 (785.4-835.1 nm). High variation within these regions was also observed when comparing single spectral signatures between excitation lasers (**Fig. 6**).

1	*Microcystis* sp.	4	*Haematococcus* sp.	7	*Cryptomonas* sp.
2	*Synechocystis* sp.	5	*Chlamydomonas* sp.	8	*Rhodomonas* sp.
3	*Synechococcus* sp.	6	*Scherffelia* sp.	9	*Chroomonas* sp.

364
365
Figure 5. Identification of artificial mix subpopulations compared between SONY ID700 Spectral
cell analyzer and BD FACSAria II cytometer. First column represents data recorded on SONY ID7000
using a set of VFs best optimized for artificial mix discrimination. Opt-SNE demonstrates successful
discrimination of the 9 subpopulations within the mix based on VF data. Second column represents
data recorded on SONY ID7000 using a set of VFs matching optical filters used on BD FACSAria.
Similarly, all 9 subpopulations were discriminated. Last column represents data recorded on BD
FACSAria. Based on optical configuration of the instrument, only 8 subpopulations were identified
using opt-SNE.

374
375

376
Figure 6. Overlay plots for each phytoplankton species within synthetic mix across 7 excitation lasers to demonstrate variability of emission signal for each excitation source. The first row demonstrates overlay plots for Cyanobacterial representatives, second – Chlorophyta and third – Cryptophyta.

Discussion

Phytoplankton are extraordinarily diverse, comprising various phylogenetic groups such as diatoms, dinoflagellates, haptophytes, and cyanobacteria. Developing spectral-based technologies for rapidly evaluating the dynamics of microalgal communities is crucial for ecological monitoring and engineering artificial microbial and algal communities. Flow cytometry, a well-established tool in

immunological and oncological research, has also been applied to phytoplankton analysis (Trask et al., 1982; Yentsch et al., 1983; Sosik et al., 2010). However, conventional flow cytometers, which rely on narrow band-pass filters to detect specific fluorophores, are limited by spectral overlap and the number of distinguishable fluorophores. Advancements in cytometry, particularly full-spectrum cytometry, present significant improvements in studying synthetic algal-microbial communities. This approach addresses limitations in traditional methods such as microscopy, pigment analysis, and conventional flow cytometry by enabling rapid, high-throughput, and label-free analysis of single algal cells from various taxonomic groups. By capturing the full emission spectra, spectral cytometry allows precise identification and enumeration of phytoplankton, facilitating a comprehensive understanding of community composition, structure, and dynamics. Additionally, it offers insights into responses to environmental stressors and enhances the characterization of microbiota associated with growing algae, advancing our understanding of phytoplankton ecology, synthetic biology, and evolution.

One of the major challenges in conventional flow cytometry, specifically relevant to phytoplankton research, is the issue of autofluorescence. While in conventional flow cytometry, autofluorescence often interferes with fluorophore emission, and is, therefore, an unwanted signal; spectral cytometry utilizes this phenomenon as a separate parameter(s). In addition to this, the ability to measure the full spectrum of light emissions from each cell allows us to differentiate between various combinations of intrinsic fluorophores that conventional instruments cannot distinguish. This is particularly advantageous in phytoplankton research, due to the high diversity of algal pigments and their overlapping spectral signatures. Alternately, approaches that provide high taxonomic resolution favor methods that allow for genus- to species-level characterization, such as amplicon-sequencing (Sommeria-Klein et al., 2021).

In this work, we demonstrate the effectiveness of full-spectrum cytometry in differentiating between different algal taxa based on their autofluorescent signatures and light scattering parameters. Clear separation between picocyanobacteria (*Synechococcus* sp. and *Synechocystis* sp.), as well as between closely related phytoplankton taxa (Synechococcales order) was demonstrated. Recorded spectral signatures were then utilized in the construction of a spectral library, which is a necessary step in monitoring synthetic communities over time. It was reported before that the spectral shape of the functional absorption spectra is remarkably constrained within major phylogenetic groups of phytoplankton, including diatoms, haptophytes, dinoflagellates, and chlorophytes (Gorbunov et al., 2020) and variability is significantly smaller than the difference between groups. In our study, the

emission spectra, obtained from microalgae with a help of ID7000 spectral cell analyzer highlight the ability to discriminate between multiple phytoplankton species in a synthetic mix, even in the presence of closely related taxa. This precision is achieved using optimazed sets of virtual filters and gating strategies, which enable the clear separation of subpopulations. The temporal monitoring of synthetic phytoplankton communities further demonstrates the applicability of full-spectrum cytometry in tracking community dynamics and responses to environmental changes. Comparative analysis with other instruments, such as BD FACSAria cell sorter and ImageStreamX MarkII imaging flow cytometer, demonstrates the higher flexibility of the ID7000 spectral cell analyzer.

Development of biological applications of full spectrum cytometry in phytoplankton research is still in its early stages, with most published work focusing on clinical applications (Wanner et al., 2022; Peixoto et al., 2022; Dott et al., 2024). One of the limitations of this approach relates to the particle size range capabilities of spectral cytometry instrumentation similar to commercially available cytometers. As a result, bulky and lengthy plankton such as many colonial and filamentous cyanobacteria, diatoms, etc. cannot be captured during the analysis. In addition to this, *in situ* measurement of planktonic cells is not available. In contrast, a range of other cytometric instruments, such as FlowCAM and CytoBuoy designed for phytoplankton, have significantly extended particle size range allowing analysis of particles of 500 µm width and over 1000 µm length (Dubelaar et al., 1989; Sieracki et al. 1998; Dubelaar, Gerritzen, 2000). Implementing full-spectrum cytometry in phytoplankton research requires reproducible protocols and standardization throughout the workflow, from sample collection and handling to data acquisition and analysis. Additionally, natural variability in phytoplankton populations, including cell size, shape, and pigment composition differences, requires optimized controls to minimize data variability. With the latest ID7000 software update, system standardization for data reproducibility became available. In the context of phytoplankton research, this means that recorded algal libraries containing unique spectral signatures can be utilized across different experiments, time points and multiple instruments, minimizing data variability. Current work has resulted in a library, which includes spectral signatures of over 30 unique phytoplankton species across different taxa.

The stability of microbial part of artificial algal-microbial mix in accordance with data from other group (Li et al., 2022) reporting population stability with mutualistic co-culturing for up to 10 days. Microorganisms within the consortia could be unculturable, hence culture-independent methods such as high-throughput sequencing and molecular fingerprinting are used for characterization of microbial

part of consortia (De Roy et al., 2014). We suggest a nanopore-based sequencing at species-level to

be performed not at the end of experiment but in parallel with spectral cytometry evaluation or weekly.

In conclusion, by single-cell analysis and differentiating between different algal taxa based on their

autofluorescent spectral signatures and light scattering parameters, full-spectrum cytometry

overcomes the limitations of traditional methods such as microscopy, pigment analysis, and

conventional cytometry. Its application extends to constructing spectral libraries for various

phytoplankton groups and monitoring synthetic communities over time, providing detailed insights

into community structure, dynamics, and responses to environmental stressors. The potential for

parallel sequencing and associated microbial analysis at the multi-omics level is needed to fully

elucidate the microbial functions occurring in artificial algal communities. This technology provides

possibilities for a wide range of research applications, from addressing fundamental ecological and

evolutionary questions to applied biology, thereby advancing the field of environmental research.

464 Authors contributions

465   A.M. designed the study, performed the experiments, analyzed the data. S.M. and N.F. contributed

466   algal cells, in algal cells characterization. I.A.V. and N.S.B. initiated and supervised the project. A.M.

467   and N.S.B. wrote the manuscript with contributions from all authors.

469 Funding

470   This research was funded by MHES Kazakhstan, grant number AP14872028 to N.S.B.

472 Acknowledgements

473   We would like to acknowledge contribution of Aigul Kussanova during the initial stages of this work

474   and Polina Len during sequence classification. We would also like to thank Dmitry Malashenkov and

475   Aiym Duisen for helpful discussion and input, and Rudolf Bichele and Steffen Boechner from Sony

476   Biotechnology for their instrumental help and support.

478 Competing interests

479   The authors declare no competing financial interests.

Data availability

Raw sequences obtained during the analysis were deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under the BioProject ID PRJNA1131605. Spectral and cytometric data obtained during this study are available from the corresponding author on reasonable request.

References

Andersen, R. A. (Ed.) (2005) Algal culturing techniques. Elsevier.

Barteneva, N. S., Dashkova, V., & Vorobjev, I. (2019) Probing complexity of microalgae mixtures with novel spectral flow cytometry approach and "virtual filtering". *BioRxiv*, 516146.

Barteneva, N. S., Kussanova, A., Dashkova, V., Meirkhanova, A., & Vorobjev, I. A. (2023) Using Virtual Filtering approach to discriminate microalgae by spectral flow cytometer. In: *Spectral and Imaging Cytometry: Methods and Protocols*, 23-40. New York, Springer.

Brenner, K., You, L., Arnold, F.H., 2008. Engineering microbial consortia: a new frontier in synthetic biology. *Trends in Biotechnology*, *26*(9), 483-489.

Castle, S.D., Grierson, C.S., & Gorochowski, T.E., 2021. Towards an engineering theory of evolution. *Nature Communications*, 12(1), 3326.

Catlett, D., & Siegel, D.A., 2018. Phytoplankton pigment communities can be modeled using unique relationships with spectral absorption signatures in a dynamic coastal environment. *Journal of Geophysical Research: Oceans*, 123(1), 246-264.

Cawse-Nicholson, K., Townsend, P. A., Schimel, D., Assiri, A. M., Blake, P. L., Buongiorno, M. F.,

... & SBG Algorithms Working Group. (2021) NASA's surface biology and geology designated

observable: a perspective on surface imaging algorithms. *Remote Sensing of Environment*, 257,

112349.F

Cho, D.H., Ramanan, R., Heo, J., Lee, J., Kim, B.H., Oh, H.M., Kim, H.S. (2015). Enhancing

microalgal biomass productivity by engineering a microalgal–bacterial community. *Bioresource*

Technology, 175, 578-585.

Curry, K. D., Wang, Q., Nute, M. G., Tyshaieva, A., Reeves, E., Soriano, S., ... & Treangen, T. J.

(2022) Emu: species-level microbial community profiling of full-length 16S rRNA Oxford Nanopore

sequencing data. *Nature Methods*, 19(7), 845-853.

De-Bashan, L.E., Bashan, Y., Moreno, M., Lebsky, V.K., & Bustillos, J.J. (2002). Increased pigment

and lipid content, lipid variety, and cell and population size of the microalgae Chlorella spp. when co-

immobilized in alginate beads with the microalgae-growth-promoting bacterium Azospirillum

brasilense. *Canadian Journal of Microbiology*, 48(6), 514-521.

Deng, Y., Mauri, M., Vallet, M., Staudinger, M., Allen, R. J., & Pohnert, G. (2022) Dynamic diatom-

bacteria consortia in synthetic plankton communities. *Applied and Environmental Microbiology*,

88(22), e01619-22.

De Roy, K., Marzorati, M., Van den Abbeele, P., Van de Wiele, T., & Boon, N., 2014. Synthetic

microbial ecosystems: an exciting tool to understand and apply microbial

communities. *Environmental Microbiology*, 16(6), 1472-1481.

Dott, T., Culina, S., Chemali, R., Mansour, C. A., Dubois, F., Jagla, B., ... & Quintana-Murci, L.

(2024) Standardized high-dimensional spectral cytometry protocol and panels for whole blood

immune phenotyping in clinical and translational studies. *Cytometry Part A*, 105(2), 124-138.

Dubelaar, G. B., Groenewegen, A. C., Stokdijk, W., Van Den Engh, G. J., & Visser, J. W. (1989)

Optical plankton analyser: A flow cytometer for plankton analysis, II: Specifications. *Cytometry: The*

10(5), 529-539.

Dubelaar, G. B. J., & Gerritzen, P. L. (2000) CytoBuoy: a step forward towards using flow cytometry

in operational oceanography. *Scientia Marina*, 64(2), 255-265.

Falkowski, P.G.m & Oliver, M.J., 2007. Mix and match: how climate selects phytoplankton. *Nature*

Reviews Microbiology, 5(10), 813-819.

Fu, H., Uchimiya, M., Gore, J., & Moran, M. A. (2020) Ecological drivers of bacterial community

assembly in synthetic phycospheres. *Proceedings of the National Academy of Sciences USA*, 117(7),

3656-3662.

Futamura, K., Sekino, M., Hata, A., Ikebuchi, R., Nakanishi, Y., Egawa, G., ... & Tomura, M. (2015)

Novel full-spectral flow cytometry with multiple spectrally-adjacent fluorescent proteins and

fluorochromes and visualization of in vivo cellular movement. *Cytometry Part A*, 87(9), 830-842.

Geng, H., Sale, K. L., Tran-Gyamfi, M. B., Lane, T. W., & Yu, E. T. (2016) Longitudinal analysis of

microbiota in microalga *Nannochloropsis salina* cultures. *Microbial Ecology*, 72, 14-24.

Gorbunov, M. Y., Shirsin, E., Nikonova, E., Fadeev, V. V., & Falkowski, P. G. (2020) A multi-

spectral fluorescence induction and relaxation (FIRe) technique for physiological and taxonomic

analysis of phytoplankton communities. *Marine Ecology Progress Series*, 644, 1-13.

Hartwell, L. H., Hopfield, J. J., Leibler, S., & Murray, A. W. (1999) From molecular to modular cell

biology. *Nature*, 402(Suppl 6761), C47-C52.

Hochberg, E. J., & Atkinson, M. J. (2003) Capabilities of remote sensors to classify coral, algae, and

sand as pure and mixed spectra. *Remote Sensing of Environmen*t, 85(2), 174-189.

Jeppesen, E., Søndergaard, M., Jensen, J. P., Havens, K. E., Anneville, O., Carvalho, L., ... & Winder,

M. (2005) Lake responses to reduced nutrient loading–an analysis of contemporary long-term data

from 35 case studies. *Freshwater Biology*, 50(10), 1747-1771.

Kolter, R. (2024). Asking a question. *Journal of Bacteriology*, 206(6), 1-7.

Kramer, S.J., & Siegel, D.A. (2019). How can phytoplankton pigments be best used to characterize

surface ocean phytoplankton groups for ocean color remote sensing algorithms? *Journal of*

Geophysical Research: Oceans, 124(11), 557-7574.

Len, P., Meirkhanova, A., Nugumanova, G., Cestaro, A., Jeppesen, E., Vorobjev, I. A., ... &

Barteneva, N. S. (2023) Species-level classification provides new insights into the biogeographical

patterns of microbial communities in shallow saline lakes. *bioRxiv*, 2023-12.

Li, X., Zhou, Z., Li, W., Yan, Y., Shen, X., Wang, J., Sun, X. & Yuan, Q., 2022. Design of stable and

self-regulated microbial consortia for chemical synthesis. Nature Communications, 13(1), 1554.

Liu, J. Y., Zeng, L. H., & Ren, Z. H. (2021) The application of spectroscopy technology in the

monitoring of microalgae cells concentration. *Applied Spectroscopy Reviews*, 56(3), 171-192.

Peixoto, M. M., Soares-da-Silva, F., Schmutz, S., Mailhe, M. P., Novault, S., Cumano, A., & Ait-

Mansour, C. (2022) Identification of fetal liver stroma in spectral cytometry using the parameter

autofluorescence. *Cytometry Part A*, 101(11), 960-969.

Perera, I.A., Abinandan, S., Subashchandrabose, S.R., Venkateswarlu, K., Naidu, R., & Megharaj,

M., 2019. Advances in the technologies for studying consortia of bacteria and

cyanobacteria/microalgae in wastewaters. *Critical Reviews in Biotechnology*, 39(5),709-731.

Poryvkina, L., Babichenko, S., Kaitala, S., Kuosa, H., & Shalapjonok, A. (1994) Spectral fluorescence

signatures in the characterization of phytoplankton community composition. *Journal of Plankton*

Research, 16(10), 1315-1327.

Robinson, J. P. (2022) Flow cytometry: past and future. *Biotechniques*, 72(4), 159-169.

Rodríguez, Y. C., Gómez, J. D., Sánchez-Carnero, N., & Rodríguez-Pérez, D. (2017) A comparison

of spectral macroalgae taxa separability methods using an extensive spectral library. *Algal Research*,

26, 463-473.

Roy, S., Llewellyn, C.A., Egeland, E.S., & Johnsen, G. eds. (2011). *Phytoplankton pigments:*
characterization, chemotaxonomy and applications in oceanography. Cambridge University Press.

Segev, E., Wyche, T. P., Kim, K. H., Petersen, J., Ellebrandt, C., Vlamakis, H., ... & Kolter, R. (2016)
Dynamic metabolic exchange governs a marine algal-bacterial interaction. *elife*, 5, e17473.

Sieracki, C. K., Sieracki, M. E., & Yentsch, C. S. (1998) An imaging-in-flow system for automated
analysis of marine microplankton. *Marine Ecology Progress Series*, 168, 285-296.

Sommeria-Klein, G., Watteaux, R., Ibarbalz, F. M., Pierella Karlusich, J. J., Iudicone, D., Bowler, C.,
& Morlon, H. (2021) Global drivers of eukaryotic plankton biogeography in the sunlit ocean. *Science*,
374(6567), 594-599.

Sosik, H.M., Olson, R.J., & Armbrust, E.V. (2010). Flow cytometry in phytoplankton
research. *Chlorophyll a Fluorescence in aquatic sciences: methods and applications*, pp.171-185.

Swan, C. M., Vogt, M., Gruber, N., & Laufkoetter, C. (2016) A global seasonal surface ocean
climatology of phytoplankton types based on CHEMTAX analysis of HPLC pigments. *Deep Sea*
Research Part I: Oceanographic Research Papers, 109, 137-156.

Trask, B. J., Van den Engh, G. J., & Elgershuizen, J. H. B. W. (1982) Analysis of phytoplankton by
flow cytometry. *Cytometry*, 2(4), 258-264.

Wanner, N., Barnhart, J., Apostolakis, N., Zlojutro, V., & Asosingh, K. (2022) Using the
autofluorescence finder on the Sony ID7000TM spectral cell analyzer to identify and unmix multiple
highly autofluorescent murine lung populations. *Frontiers in Bioengineering and Biotechnology*, 10,
827987.

Xu, Y., Wang, Y., Yang, Y., & Zhou, D., 2016. The role of starvation in biomass harvesting and lipid
accumulation: Co-culture of microalgae–bacteria in synthetic wastewater. *Environmental Progress &*
Sustainable Energy, 35(1), 103-109.

Yentsch, C. M., Horan, P. K., Muirhead, K., Dortch, Q., Haugen, E., Legendre, L., ... & Zahuranec,

B. J. (1983) Flow cytometry and cell sorting: A technique for analysis and sorting of aquatic particles

1. *Limnology and Oceanography*, 28(6), 1275-1280.

647