A variational deep-learning approach to modeling memory T cell dynamics

Christiaan H. van Dorp[1,*], Joshua I. Gray[2,*], Daniel H. Paik[2], Donna L. Farber[2,†], and Andrew J. Yates[1,†]

[1]Department of Pathology and Cell Biology, Columbia University Irving Medical Center, New York City, USA
[2]Department of Microbiology and Immunology, Columbia University Irving Medical Center, New York City, USA
[*]Authors contributed equally
[†]Corresponding authors: `andrew.yates@columbia.edu`, `df2396@cumc.columbia.edu`

July 7, 2024

Abstract

Mechanistic models of dynamic, interacting cell populations have yielded many insights into the growth and resolution of immune responses. Historically these models have described the behavior of pre-defined cell types based on small numbers of phenotypic markers. The ubiquity of deep phenotyping therefore presents a new challenge; how do we confront tractable and interpretable mathematical models with high-dimensional data? To tackle this problem, we studied the development and persistence of lung-resident memory CD4 and CD8 T cells (T_{RM}) in mice infected with influenza virus. We developed an approach in which dynamical model parameters and the population structure are inferred simultaneously. This method uses deep learning and stochastic variational inference and is trained on the single-cell flow-cytometry data directly, rather than on the kinetics of pre-identified clusters. We show that during the resolution phase of the immune response, memory CD4 and CD8 T cells within the lung are phenotypically diverse, with subsets exhibiting highly distinct and time-dependent dynamics. T_{RM} heterogeneity is maintained long-term by ongoing differentiation of relatively persistent Bcl-2[hi] CD4 and CD8 T_{RM} subsets which resolve into distinct functional populations. Our approach yields new insights into the dynamics of tissue-localized immune memory, and is a novel basis for interpreting time series of high-dimensional data, broadly applicable to diverse biological systems.

Introduction

The widespread implementation of single-cell high-throughput assays in recent years has driven the development of a plethora of computational tools for generating lower-dimensional ('latent') representations of datasets that occupy high-dimensional 'feature' spaces, usually involving the identification of discrete clusters of cellular states or phenotypes[1,2]. Typically, inferences are derived from single snapshots of systems at steady state. However, following the response of a system to perturbations has the potential to reveal more about the components and interactions that underpin it. Mathematical models are used widely for formulating mechanistic descriptions of cellular dynamics, and drawing inferences from time-series of observations. Such models usually describe the directly-observed trajectories of small numbers of discrete, pre-defined cell types or states. An outstanding challenge, then, is to connect dynamical modeling tools to exploit the information contained with the burgeoning quantities of high dimensional data that are now available.

An acute immune response is an archetype of a dynamic, perturbed set of biological processes. In particular, T cells that respond to infectious challenge undergo clonal expansion and diversification, generating heterogeneous populations of memory-phenotype cells that are capable of responding rapidly and potently to re-exposure to the same infectious agent. These memory T cells can be broadly characterized as circulating or tissue-resident[3]. To date, the majority of modeling studies have focused on the dynamics of circulating memory subsets in both humans and mouse models, characterizing their ontogeny, diversity, kinetics, and persistence[4–8]. In contrast, the dynamics of tissue-resident memory T cells (T_{RM}), which patrol organs and barrier sites and provide indispensable front-line protection against a wide variety of pathogens[9], are relatively understudied[10]. Understanding how T_{RM} are established and maintained, and what determines the duration of protection that they offer, has the potential to revolutionize vaccine design[11], our understanding of autoimmune responses, and the control of tumor growth.

As with cell types in many areas of biology, T_{RM} are usually defined by small numbers of surface markers and are traditionally identified with flow cytometry using hierarchical binary gating strategies[11]. Lung $CD8^+$ T_{RM} are commonly associated with CD103, which binds to E-cadherin present on epithelial cells, supporting retention in barrier tissues, and CD69, which limits tissue egress by antagonizing S1P1-mediated extravasation[9,11]. Lung $CD4^+$ T_{RM} are often characterized by expression of CD69 and the chemokine receptor CXCR6[12,13]. However, the expression patterns of many of these markers do not yield objectively defined cutoff values to characterize positive and negative populations unambiguously, a problem compounded as the number of these markers grows[14]. The ontogeny and persistence of T_{RM} therefore provide an ideal setting to explore the application of dynamical modeling to high-dimensional phenotyping data.

An obvious approach is to perform initial unsupervised clustering step, which combines dimension reduction and phenotypic classification. One can then use sets of ordinary differential equations (ODEs) or other dynamical systems to describe the time evolution of the sizes of these clusters[15], some of which may represent phenotypes that are not canonically defined. This process typically yields estimates of rates of cell loss, self-renewal, differentiation, and cell killing. In prin-

ciple this sequential approach is highly tractable, because many off-the-shelf clustering methods are available, and standard tools can be applied to perform inference and model selection. Potential problems remain, however. First, clusters are usually not clearly separable and so in general the assignment of cells to clusters should be probabilistic. Second, as the phenotypic structure of a population that is out of equilibrium evolves, the uncertainty in cluster assignment may also be time dependent. These issues suggest that under some conditions it may be necessary to jointly model the distribution of the flow-cytometry data itself, as well as the underlying cellular dynamics.

To explore these issues, we used high dimensional flow cytometry to characterize the dynamics and phenotypic structure of T cell subsets within the lungs of mice that were infected with influenza A virus (IAV) and followed up to 57 days post-infection. To model the time evolution of these populations, we developed and compared two methodologies. The first, 'sequential' approach employs standard clustering methods applied to the processed flow cytometry data pooled across time points, and models the time evolution of the cluster sizes with ODE models. The second, 'integrated' method employs deep learning and stochastic variational inference[16] to simultaneously model the structure and dynamics of observed marker expression, via a lower dimensional representation of the data. With these approaches we find that both CD4 and CD8 T cells within the lung are phenotypically and dynamically heterogeneous. We also identified persistent populations within both subsets that may maintain this diversity over weeks to months through continuous differentiation. We discuss the benefits and limitations of the two approaches to modeling high dimensional time series, and conclude by highlighting areas for further development.

Results

The size and phenotypic composition of lung-localized T cell populations change dynamically following influenza infection in mice

The mouse model of influenza infection recapitulates the dynamics of virus morbidity and the immune response seen in human infection and results in the generation of lung CD4 and CD8 tissue resident memory T cells[17,18] (T_{RM}). We followed the T cell responses to influenza A virus (IAV) infection in mice, at timepoints between 6 and 57 days post-infection (DPI; Fig. 1A; see Methods). We restricted our analyses to antigen-experienced ($CD44^+ CD11a^+$) $CD8^+$ and $CD4^+$ T cells (which hereon we refer to as CD8 and CD4 T cells, for brevity) within the lung tissue, as defined by protection from *in vivo* antibody labeling as described[17,19] (Fig. S1). We previously determined that T cells in this lung resident niche represent the majority of influenza-specific memory T cells[19]. Both populations peaked in numbers between 8-10 DPI, shortly after the point of maximum weight loss as a measure of infection morbidity (Fig. S2A), and then declined with population half-lives of roughly 3 and 5 days respectively (Fig. 1B and C). Both CD8 and CD4 T cells approached stable numbers by 37 DPI, after contracting approximately 100-fold and 10-fold, respectively.

Figure 1: Studying the timecourse of the lung-localized CD8 and CD4 T cell responses to IAV infection.
A. Experimental design. Each cohort contained 4 or 5 mice and 42 mice were used in total. Eight mice were excluded due to lack of weight loss or i.v. label failure (Fig S2). **B-C.** Total numbers of antigen-experienced, protected CD8$^+$ and conventional CD4$^+$ T cells by day post infection (DPI), calculated by multiplying the estimated total number of lung cells with the fraction of protected CD44$^+$ CD11a$^+$ CD8$^+$ and conventional CD4$^+$ T cells. **D-E.** Frequencies of cells expressing combinations of the tissue-resident memory markers CD69 and CD103 within protected CD44$^+$ CD11a$^+$ CD8$^+$ T cells. **F-G.** Frequencies of cells expressing combinations of the tissue-resident memory markers CD69 and CXCR6 within protected CD44$^+$ CD11a$^+$ CD4$^+$ T cells. **H-I.** Marginal distributions of expression of a selection of markers, aggregated over mice and stratified by day post infection, and lineage (CD8 or CD4).

These smooth trajectories were highly characteristic of adaptive immune responses and of IAV responses in particular[20,21]. However, within them, we identified considerable heterogeneity with respect to surface markers of tissue residency[11,18,20]. The (relative) abundances of T cell subsets defined by combinations of the canonical markers CD69, CD103 and CXCR6 clearly shifted during the contraction phase of the response (Figs. 1D-G). For example, the relative abundance of CD8 T cells expressing both CD69 and CD103 increased with time, then decreased, while CD8 T cells expressing CD69 but not CD103 were rapidly lost in the second week post infection, and then stabilized (Fig. 1D-E). Around half of CD4 T cells expressed both CD69 and CXCR6 beyond 4 weeks post infection. CD4 T cells with low CXCR6 and high CD69 expression mostly disappeared around the peak of the immune response, and then very slowly grew in proportion (Fig. 1F-G). This (temporal) heterogeneity was manifested more clearly when phenotyping was expanded to include markers of cell survival (Bcl-2), access to lymphoid organs (CD62L, typically used to define circulating 'central' memory cells, T_{CM}), differentiation (KLRG1 and PD-1), a costimulatory receptor associated with cell survival (CD27), the chemokine receptor CX3CR1, the follicular helper T cell marker FR4, and the integrin CD49a. The marginal distributions of the expression of many of these markers showed clear variation with time (Fig. 1H and I). Thus, CD4 and CD8 T cells within the lung both exhibit complex and shifting phenotypic structures during the contraction and memory phases of IAV infection.

Sequential Approach

Standard unbiased clustering methods clearly delineate heterogeneity within CD8 T cells

To understand and quantify how these structures emerge and evolve, we began with a sequential approach to modeling the population dynamics of T cells within the lung from the peak of infection onward. For clarity, we first focus largely on the CD8 T cell response, and then present a more condensed description of the CD4 T cell response that highlights the key results.

We aggregated the flow cytometric data from all mice and all time points, and then assigned CD8 T cells to populations using the Leiden clustering method[22] and manual annotation. For each mouse, we tabulated the proportions of cells assigned to each population. Together with the measured total numbers of antigen-experienced protected CD8 T cells in the lung, this process yielded time series of the cluster sizes as a function of time post infection. We used Bayesian dynamical models and the leave-one-out information criterion, implemented in Stan[23,24], to infer biological parameters from these time series and assess the relative support for different models. The workflow of the sequential approach is summarized in Fig. 2A.

The Leiden algorithm is stochastic, and we found that both the number and marker expression profiles of the clusters varied from run to run. To address this uncertainty we performed a consensus clustering step. We ran the Leiden method 20 times, and manually annotated the clusters as cell types. Each cell's type was determined by its consensus assignment. We return to the issue of uncertainty in cluster assignment below.

Figure 2: Pre-processing flow cytometry data for the sequential approach. Results are based on data from $n = 27$ mice. **A.** Flow-chart of the sequential approach. **B.** Marker expression heatmap for selected markers and consensus T-cell populations. **C.** Uniform Manifold Approximation and Projection[25] embedding (UMAP) of the marker expression data, colored by annotation. **D.** UMAPs of marker expression data, split by day post infection (DPI). The color scale reflects cell density in UMAP space. **E.** Time series of the fraction of cells in each cluster. The lines show a linear interpolation on the log scale. The panels have the same y-scale for easy comparison and the inset axis shows the lower values of the DN T_{RM} population.

The Leiden clustering robustly identified eight distinct CD8 T cell populations (Fig. 2B and Fig. 2C). Central memory T cells (T_{CM}) were CD62L^{hi} Bcl-2^{hi}, and an effector-like population (T_{Eff}) was clearly defined by high levels of expression of KLRG1. We identified four T_{RM}-like populations based on expression of CD69 and CD103, together with CD49a, CXCR6, and PD-1. We denoted them DP T_{RM} (CD69^{hi} CD103^{hi}), CD69 SP T_{RM} (CD69^{hi} CD103^{lo}), CD103 SP T_{RM} (CD69^{lo} CD103^{hi}), and DN T_{RM} (CD69^{lo} CD103^{lo}), but note that these populations also vary in their expression of other markers. We also identified a population which we define as effector memory (T_{EM}) cells, which lacked expression of markers conventionally associated with T_{CM} and T_{RM}. A subset of these T_{EM} cells expressed CX3CR1 at high levels, which has been postulated to be a key marker for defining memory CD8 T cell subsets[26] and may indicate a reduced capacity for residency[27].

IAV nucleoprotein-specific T cells in lung cluster similarly to their polyclonal parent populations

To further validate our clustering step, we used a fluorescently labeled MHC class I tetramer to identify CD8 T cells specific for the NP$_{311-325}$ epitope of IAV, and examined their distribution relative to the cluster locations identified for the response in total (Fig. S3). The tetramer-positive (Tet$^+$) populations represented approximately 10% of CD8 T cells in bulk, and exhibited similar kinetics (Fig. S4), in concordance with previous findings[19]. The partitioning of the UMAP space into 8 sub-populations appeared to respect the density of Tet$^+$ cells (Fig. S3), although they were somewhat enriched within the CD69 SP T_{RM}, DN T_{RM} and both T_{EM} populations, suggesting some variation in phenotypic structure with epitope specificity, and/or that a proportion of CD8 T cells within the lung were not IAV-specific. Similar observations held for the NP$_{311-325}$-specific CD4 T cells, identified with an MHC class II tetramer, which represented roughly 5% of total CD4 and were slightly enriched for an effector-like population (Fig. S3). Overall, however, the congruent kinetics of epitope-specific and bulk populations, the fact that these bulk lung niche cells are likely IAV-specific[19], and the improved resolution provided by greater cell numbers, lead us to continue with our analyses of the bulk lung niche CD4 and CD8 T cells.

Dynamical modeling reveals evidence for ongoing differentiation and time-dependent loss rates of lung CD8 T cell populations

Having partitioned the CD8 T cells among the eight populations, we visualized their time evolution qualitatively by UMAP (Fig. 2D) and quantitatively, showing their abundances within the population as a whole (Fig. 2E). Overall, the dominant populations were DP T_{RM}, CD69 SP T_{RM} and T_{EM}, with a relatively stable but lower representation (5-10%) of T_{CM}. The T_{Eff} proportion quickly fell during the resolution phase to stabilize at around 1% by 16 DPI, while the DN T_{RM} fraction became almost undetectable over the same timescale. The dynamics of the DP T_{RM} and CD69 SP T_{RM} proportions resembled those derived by manual gating (Fig. 1D and E), serving as additional validation of the Leiden clustering approach.

To quantify the cellular dynamics underlying these shifts in phenotype, we fitted several models to the timecourses of cluster sizes (Fig. 3A). In the simplest model (I), we estimated a single net loss rate (the balance of loss due to cell death and/or egress from the lung, and self-renewal) for each population. In model II we allowed the net loss rates to be time-dependent, reflecting for example changes in inflammation level within the lung after the virus is cleared. In more flexible models we also allowed differentiation between populations, both without (model III) and with (model IV) time-dependent loss. To quantify the effect of migration of circulating T cells to the lung during the memory contraction phase, we performed a cell transfer experiment using CD90 congenic mice (Fig. S5 and SI Text A), which suggested that such ingress contributes minimally to the population dynamics after the infection is cleared. In all models we therefore make the simplifying assumption that ingress is negligible.

The loss of lung-resident CD8 T cells slows progressively following the resolution of IAV infection

From 8 DPI, the biphasic loss of CD8 T cells in the lung (Fig. 1C) could in principle be described by model I, in which two or more populations are lost at different but constant rates (SI Text B). A general feature of this model is that the trajectories of the subpopulation frequencies on a logarithmic scale are necessarily linear or concave, but not convex (SI Text C). The data were clearly incompatible with this behavior (Fig. 3B), and so we rejected it. In model II we allowed the rate of loss of each population to transition between initial and longer-term values at an exponential rate u. For simplicity, we assumed that this modulation in dynamics applied to all populations in tandem, potentially reflecting global changes in the lung environment as inflammation is resolved. This model yielded a markedly improved fit (Fig. 3B, second column), allowing for convexity in population frequencies (notably CD69 SP T_{RM}, T_{EM}, and T_{Eff}), and better capturing the biphasic decline in total cell numbers. The estimated timescale of transition between the early and long-term dynamics, $1/u$, was approximately 3 days, reflecting the large shifts in relative abundances of phenotypes between 8 and 14 DPI. Formal comparison of models I and II using LOO-IC[28] confirmed strong support for time-dependent loss rates (ΔLOO-IC $= 130 \pm 31$; see Table S1). The long-term loss rates λ_L yielded by model II, shown in Fig. 3C, are consistent with our prior estimates that influenza-specific CD8 T cells are lost from the lung with a population half life ($\ln(2)/\lambda_L$; reflecting the net effect of loss and any self-renewal) of 18-25 days[10].

Evidence for ongoing differentiation of CD8 T cell subsets within the lung

Model III extended model I to allow differentiation between CD8 phenotypes after 8 DPI. Allowing differentiation between every pair of populations risks overfitting, and so we penalized large values of Q_{ij} by regularizing them with an exponential prior distribution, similar to the lasso[29]. A key prediction of model III is that it allows subset frequencies to stabilize at late time points, as was observed for some populations (Fig. 3B, third column), because the balance of differentiation and loss can generate stable equilibria. Indeed we found substantial support for differentiation (model

Figure 3: Modeling lung resident CD8 T cell dynamics using the sequential approach. Results are based on data from $n = 27$ mice. **A.** Schematic of the mathematical models, illustrated with 3 populations. X_i is the number of cells in population i and $\lambda_i = \delta_i - \rho_i$ its (possibly time dependent) net loss rate. Q_{ji} is the *per capita* rate of differentiation from i to j. **B.** Top panels: numbers of CD8$^+$ T cells in the lung (black dots) and the model fits (blue lines). Remaining panels: observed and predicted subpopulation frequencies. Bands indicate 95% credible envelopes, bars indicate the 2.5 and 97.5 percentiles of the posterior predictive distributions. **C.** For model II, marginal posterior distributions of the parameters as violin plots (blue), with median (red) and 95% credible intervals (CrI; black). **D.** Differentiation rates in model IV. Diagonal; total egress rate by differentiation.

III) over the simplest model with constant loss rates, model I (ΔLOO-IC $= 82 \pm 29$). However, time-dependent loss only (model II) was still supported most strongly. We then explored model IV that combined the features of models II and III. This model was able to capture both the rapid early dynamics and long-term stability of subpopulation frequencies (Fig. 3B, fourth column), and it was very strongly supported statistically (ΔLOO-IC $= 29 \pm 6.8$, relative to model II). Indeed, after translating LOO-IC to Bayesian model weights[30], model IV had close to 100% support among the four candidates. However, it predicted differentiation between nearly all subpopulations (Fig. 3D), at rates with wide credible intervals (not shown). Further, the net loss rates had strong posterior correlations with the differentiation rates, indicating that the model can trade off loss with differentiation. For example, high rates of death and/or onward differentiation, balanced by influx from a precursor, are difficult to discriminate from a more isolated population that is terminally differentiated. This tension between quality of fit and interpretability highlights the limitations of relying on statistical support alone to select models.

Uncertainty in clustering highlights shortcomings of sequential approaches

Stratifying cells within the UMAP by timepoint allows one to visualize changes in the distribution of expression profiles within each cluster (Fig. 2D). Ideally, the distribution of cells within a subpopulation should be independent of time, but from 20 DPI onward the distribution within CD69 SP T_{RM} shifted towards T_{EM} and CD103 SP T_{RM}, T_{EM} moved towards to T_{CM}, and DP T_{RM} moved away from SP and DN T_{RM}. Related to these shifts, we found variation in the certainty with which cells could be assigned to each population. We quantified this uncertainty using the empirical entropy of these assignments (Fig. S3A). A low entropy corresponds to a cell that is always assigned to the same population, while high entropy indicates that different Leiden runs result in different assignments. Uncertainty of assignment also varied between populations (Fig. S3B). These discrepancies indicate that the Leiden clustering method may not capture important features of the data, and suggests that a more refined representation of the high-dimensional data is needed – one that takes into account the time evolution of the cell densities in phenotype space.

Integrated Approach

To deal with uncertainty in cluster assignment, we took inspiration from the single-cell modeling tools `scVI`[31] and `scANVI`[32]. We assumed that cell's flow cytometric phenotype, x, can be modeled by lower-dimensional latent vector z representing the cell's internal state, and that x can be reconstructed from z with a complex non-linear map, modeled with an artificial neural network (NN). The measured fluorescent intensities are noisy, and so z cannot be computed exactly; instead, for a given x we aim to generate a probability density for z. To avoid explicitly modeling this density for each cell, we make use of an amortized inference model, using the observations x to predict a parameterized distribution function for z. This predictor is also a NN model. In the language of

Figure 4: The integrated approach. A. Schematic of the VAE model. Using the fluorescence data x, we use a fully connected encoder network to infer the latent cell state (specifically, the variational parameters μ_z and σ_z for the posterior distribution of $z|x$; see Methods). The prior distribution of z is given by a Gaussian mixture model with time-dependent mixture weights determined by the dynamical model. The mixture components (multivariate normal likelihood functions) and mixture weights determine the categorical distribution of cluster membership of a cell. Using the decoder network, we compute parameters μ_x and σ_x for the distribution of $x|z$, thereby accounting for the measurement error in x. **B.** Flowchart indicating the logic behind the integrated approach. The model is fit directly to single-cell flow cytometry data and cell count data using stochastic variational inference (SVI). The posterior distribution of the model parameters (loss and differentiation rates, initial conditions, NN weights and biases, GMM parameters) are iteratively optimized. In each iteration we sample candidate parameters, and solve the system of ODEs. We then match the data with our predictions, and adjust the posterior to improve this match with the gradient ascent method.

variational auto-encoders (VAE), the amortization network is the 'encoder', and the network that generates observations x given a latent state z is the 'decoder' (Fig. 4A).

To model the population structure within the latent space, we used a multivariate Gaussian mixture model (GMM)[33,34]. The components and weights of the GMM correspond to the different T-cell subpopulations and their relative sizes, respectively. The ODE models predict the weights at any time point, and so the GMM describes a distribution of cell states that changes smoothly with time, parameterized by the rates of loss and differentiation, and the time-independent mixture component parameters. These mixtures overlap within the latent space, and so their time-dependent weights can capture shifting distributions of cells in both the latent and feature space. A schematic describing the time-dependent GMM is shown in Fig. 4A, where the latent space is represented by the 1-dimensional real line. A model predicting the size trajectories of four subpopulations is used to compute the weights of the mixture components (colored bell curves). The black curves show how the combined latent state distribution of all sub-populations evolves with time. The computational workflow for this method, which we refer to as the integrated approach, is illustrated in Fig. 4B. We implemented it in Pyro[35], and describe it in detail in Methods.

Dimension reduction using the integrated approach preserves information within the flow cytometry data

The sequential approach revealed evidence for both differentiation and time-dependent net loss rates, and so we proceeded with model IV. We first assessed our ability to find a latent representation consistent with our understanding of the marker expression data. We began with a 6-dimensional latent space representation of the 10-dimensional space of marker expression levels. Our results were similar with 5- and 7-dimensional latent spaces. Using a UMAP embedding of the latent space to visualize the learned representation, we saw that the VAE model was able to distinguish cells with different marker expression profiles (Fig. 5A) and yielded distinct populations (Fig. 5B). To visualize these sub-populations more clearly, we sampled latent vectors from each of the mixture components, mapped them the UMAP space, and calculated the contours of the 75% maximum density for each of the components. We also took the posterior samples of the location vectors of each of the components and projected them in UMAP space (indicated by colored dots in Fig. 5B). These location vectors can be interpreted as the typical cell state in a component. The contours and typical cell states match well with distinct regions in the latent space, indicating that the GMM is correctly capturing the distribution of latent cell states and, by extension, the distribution of the flow cytometry data. To account for possible batch effects, the batch information was included in the training data for the VAE. However, the trained models did not make use of large batch corrections (SI Text D), possibly due to the fact that cells from all mice were analyzed simultaneously.

The integrated approach yields finer-grained T cell phenotypes than the sequential approach

The GMM defines a distribution of cell states as a function of time, which we can match with data via the decoder NN. However, for each cell we can also use the encoder NN to compute its latent cell state, and then use the GMM to assign the cell to one of the mixture components. This is a probabilistic (or 'soft') cluster assignment; for each cell, the GMM gives a distribution of probabilities that the cell belongs to each component. By sampling from these distributions, we can assign to each cell a single component label (Fig. 5C). At this stage, we have effectively clustered the data, but here the key difference with the sequential approach is that this happens *after* we trained our dynamical model. The inference method does not rely at all on the fact that we assign cells to single clusters. Now that we have clustered the flow cytometry data, we can interpret the different mixture components as CD8 T cell sub-populations, and annotate them using the mean fluorescence intensities per cluster (Fig. 5D). We compared populations discovered by the two approaches using the Jaccard index (Fig. S6A). The integrated approach recovered 7 of 8 populations identified by the sequential method. Namely CD69 SP T_{RM}, DP T_{RM}, CD103 SP T_{RM}, which we denote TRM 1, 2, 3; T_{Eff} and T_{CM}; T_{EM} CX3CR1^{hi} and CX3CR1^{lo} T_{EM}, which we denote TEM 1 and 3. It did not identify the transient DN T_{RM} subset that expressed PD-1, CXCR6, and the early activation marker CD27, and was lost quickly early in the resolution phase (Fig. 2E). Instead, the integrated approach stratified two populations further; CX3CR1^{lo} T_{EM} were separated into Bcl-2^{lo} and Bcl-2^{hi} (TEM 1

Figure 5: Modeling the dynamics of lung CD8 T cells with the integrated approach. Results are based on data from $n = 27$ mice. **A.** For each marker, the fluorescence intensity of each cell is displayed in a UMAP representation of the latent space. **B.** Cell density and 95% confidence ellipsoids for GMM components projected into the same UMAP space. Dots are the GMM location vectors. **C.** Using the GMM, a population is assigned to each cell. **D.** Mean fluorescence intensity of each marker and population identified by the GMM. **E.** Calculated population frequencies per mouse and timepoint, with median trajectories predicted by the model and 95% CrI (bands). **F.** Measured numbers of $CD44^+$ $CD11a^+$ protected CD8 T cells in lung (points), trajectory of the median prediction (line), 95% CrI (dark band), 95% posterior predictive interval (light band). **G.** Marginal posterior distributions of net loss rates during the early and late stages, with medians and 95% CrIs. **H.** Differentiation matrix.

and 2), and CD69 SP T_{RM} became three subsets; Bcl-2^{lo} CX3CR1^{lo}, Bcl-2^{lo} CX3CR1^{hi}, and Bcl-2^{hi} CX3CR1^{lo} (TRM 1, 4, and 5 respectively). This partitioning could not be readily distinguished by UMAP inspection alone (Fig. 5A).

With the integrated approach, data and model are entwined

Having clustered the flow cytometry data, we could again construct time series, enumerating for each mouse the number of cells per population (points in Fig. 5E), and then compare their relative sizes to the trajectories predicted by the fitted model (model IV; curves and 95% credible envelopes in Fig. 5E). This model accurately captured many features of the distribution of phenotypes, as well as the total population size (Fig. 5F). The CD69^{hi} Bcl-2^{lo} T_{RM} (TRM 1) declined rapidly relative to the other populations between 8 and 12 DPI, recovered, but decreased again from around a month after infection, due to the continually increasing prevalence of CD69^{hi} Bcl-2^{hi} T_{RM} (TRM 5). This subset was present initially at a very low frequency, but come to comprise $> 30\%$ of lung CD8 T cells by 2 months post infection. A similar pattern can be seen for T_{EM}, which increased in frequency soon after the peak of infection but declined rapidly after 10 DPI. The Bcl-2^{hi} T_{EM} (TEM 2) population was initially very small, but quickly replaced Bcl-2^{lo} T_{EM} (TEM 1 and 3) and stabilized at 10% of CD8 T cells from 16 DPI. Therefore the integrated approach reveals that the Bcl-2^{hi} and Bcl-2^{lo} T_{RM} and T_{EM} populations, which are otherwise phenotypically similar, exhibit quite divergent dynamics. However, there is one subtlety with presenting the data as a timecourse of discrete observations. Calculating the probability that a cell is assigned to a cluster uses two pieces of information; the cell's marker expression, and the expected sizes of each of the clusters. The latter are dependent on time, and this time dependence is in turn learned from the flow cytometry data. Therefore, the data displayed in Fig. 5E are in part generated by the model, and as a result we expect the data points to follow the model trajectories more closely than in a standard fitting approach.

Bcl-2-expressing T_{RM} act to sustain lung CD8 T cell populations long-term

As with the sequential approach we found that up to around 14 DPI T_{Eff} are lost most rapidly, followed by TRM 1 and T_{CM}. At the same time, TRM 2 and 3, and TEM 2 and 3 were growing in numbers (Fig. 5G, left panel). From approximately 14 DPI onwards, all populations declined in number except for TRM 5, the Bcl-2^{hi} CD69^{hi} CD103^{lo} subset (Fig. 5G, right panel). Bcl-2^{lo} T_{EM} (TEM 1) were lost most rapidly, consistent with their lack of typical residency markers. Notably, at these late times T_{CM} and T_{Eff} were more persistent than canonical T_{RM}. Most strikingly, Bcl-2^{hi} CD69 SP T_{RM} (TRM 5) had a positive intrinsic net growth rate in the late memory phase (that is, we inferred that their rate of self-renewal was greater than their combined rates of death or egress from the lung), but their numbers declined due to their high rates of differentiation into DP T_{RM} (TRM 2), Bcl-2^{hi} T_{EM} (TEM 2), T_{CM}, and CD103 SP T_{RM} (TRM 3). Therefore, our analyses suggest that Bcl-2^{hi} T_{RM} help to maintain phenotypic diversity among CD8 T cells in the lung in the weeks following IAV infection.

Model validation with the integrated approach

In the integrated approach the generated and predicted cluster frequencies are interdependent to an extent. How then do we validate a model's utility or predictive capacity? In contrast to the sequential method, which predicts only the trajectories of the sizes of predefined clusters, the VAE model is generative – that is, it can be used to predict the trajectory of the flow cytometry data itself. Such a trajectory is shown in Fig. 6A. Each square shows a UMAP of the latent space at a particular time post infection. The squares highlighted in grey are UMAPs of the actual data, and the others are simulated by the model. This series of UMAPs acts as a visual posterior predictive check (i.e. a 'model fit'), analogous to a visual assessment of how well the curves in Fig. 3B capture the trends in the datapoints. In this case we can see the method describes the data well, and highlights time periods that may require denser sampling. For instance, the simulated data between 8 DPI and 12 DPI show rapid shifts in the distribution of phenotypes; CD69 SP T_{RM} rapidly disappear, DP T_{RM} increase in frequency, and then T_{EM} start to decline. This order of events is not directly observed, and might be resolved with intermediate samples.

Because the integrated approach models the joint density of the marker expression, we can easily marginalize to single cell surface markers and compare the model predictions with the data. This provides another posterior predictive check (Fig. 6B). Pseudo-observations are created by sampling from the GMM at the observation times, and using the decoder network to transform the latent space to the feature space. The marginal distributions of the pseudo observations (red density plots) largely coincide with the true data (black densities), but there are some discrepancies. Most notably, the VAE has difficulties capturing the bimodal shape of the CD49a density, as well as the shape of the Bcl-2 distribution at late time points. To investigate these discrepancies further, we made use of the auto-encoder architecture of the model. An auto-encoder finds a low-dimensional representation of the data, thereby removing noise and finding essential features[36]. We can test how well our auto-encoder has learned to represent the data by first encoding the marker expression data set, and then reconstructing the expression profiles from the latent cell states using the decoder network. For each marker, we plot the observed fluorescence intensities against the reconstructed values (Fig. 6C). A perfect reconstruction would yield a perfect correlation between input and output. For the markers CD69 and CD103, we find a very high correlation (R^2 = 0.92 and 0.98, respectively), while for CD49a and CD27, we find low correlations (R^2 = 0.18 and 0.15, respectively). Hence, it appears that the VAE discarded most of the information encoded by CD49a and CD27 expression, indicating that variation in these markers does not relate strongly to heterogeneity in the dynamics of CD8 T cell subsets in the lung.

The dynamics of lung-localized CD4 T cells following IAV infection

We then applied the two approaches to the CD4 T cell data. With Leiden clustering and consensus annotation, we identified eleven distinct populations (Figs. S7A and B). Seven clusters possessed a T_{RM} phenotype, delineated by varying expression levels of canonical T_{RM} markers CD69, CD49a, CXCR6, CD103. One of the T_{RM} subsets also expressed KLRG1, suggesting an effector phenotype.

Figure 6: Posterior predictive checks using simulated marker expression data. Results are based on data from $n = 27$ mice. **A.** UMAPs of marker expression data, either simulated using the fitted model (white background), or representing the observations (grey background). The location of the inferred clusters are highlighted in the top left panel (colors as in Figure 5B). The color (blue to yellow) shows the density of the points in UMAP space. The number of sampled cells is gradually decreased with DPI to match the sample size of the actual data. The marker expression data (simulated or real) is first mapped to the latent space using the trained VAE model. Then, a UMAP model is trained on the combined true data. The same UMAP model is used to map the simulated data to the plane. **B.** Marginal distributions of marker expression (cf. Fig. 1H). Data is shown in black, simulated data is shown in red. **C.** Input data and reconstruction using the auto-encoder model. The number in brackets is the coefficient of determination (R^2).

We also identified four populations lacking these residency markers: a CD62L^{hi} TCM-like subset, a subset expressing high levels of FR4 and PD-1, characteristics of T follicular helper (T$_{fh}$) cells in the lung[37], and two populations with low expression of all markers measured, distinguished only by the presence of Bcl-2, which we termed T$_{EM}$.

As we found for the CD8 T cells, the joint distribution of marker expression on CD4 T cells gradually shifted with time post infection (Fig. S7C). The peak of the response was dominated by Bcl-2^{lo} T$_{EM}$ and the CD69^{lo} CD49a^{lo} T$_{RM}$ population, which expressed high levels of PD-1 and CD27, markers upregulated during activation (Fig. S7D). By 3 weeks post infection, CD69^{hi} CD49a^{lo} and CD69^{hi} CD49a^{hi} T$_{RM}$ dominated, giving way in turn to two Bcl-2-expressing T$_{EM}$ and T$_{RM}$ subsets by 2 months post infection.

We fitted our four model variants to the timecourses of total CD4 T cell numbers (Fig. 1C) and the subset frequencies (Fig. S7D). Visual inspection and posterior predictive checks indicated that all four models fit the data reasonably well (Fig. S8A), but formal model comparisons indicated that time-dependent net loss alone (model II) or differentiation alone (model III) provided sufficient descriptions of the data (Table S1).

As was the case for CD8 T cells, all models exposed remarkable heterogeneity in the dynamics of CD4 T cell subsets in the lung after the resolution of IAV infection (Fig. S8B; using model II), particularly among the T$_{RM}$-like populations. The loss rates estimated with model III showed a similar, but slightly more homogeneous pattern (Fig. S8C). Again using model III, we found evidence for ongoing differentiation from CD69^{lo} CD49a^{lo} T$_{RM}$ to CD69^{hi} CD49a^{hi} T$_{RM}$ and CD69^{lo} CD49a^{hi} T$_{RM}$, and from Bcl-2^{hi} T$_{RM}$ to multiple T$_{EM}$ populations and effector cells (Fig. S8D). Finally, the data suggests a pathway from CD69^{hi} CD49a^{lo} T$_{RM}$ to T$_{EM}$. These data indicate the dynamically shifting T cell populations within the local environment, even after resolution of infection. high entropy scores within T$_{RM}$ clusters indicated substantial uncertainty in Leiden cluster assignment for CD4 T cells (Fig. S3D-E), underscoring the need for integrated analysis to dissect the subtle changes in subset composition.

Next, we used the integrated approach to refine the model fit for the CD4 data. We chose to continue with model III as it performed just as well as the full model and enabled us to further explore differentiation pathways. The VAE generated a latent representation that respected marker expression, and the location of the mixture components in UMAP space corresponded well with high density regions (Fig.7A-B). Using mean fluorescence intensity (Fig. 7C), the integrated approach identified T$_{CM}$ (CD62L^{hi}) and a KLRG1^{hi} subset that also expressed T$_{RM}$ markers (TRM 7) whose phenotypes were highly concordant with those identified using the sequential approach. In contrast, many of the T$_{RM}$ and T$_{EM}$ populations were distinct between the two approaches (Fig. S6B).

The integrated approach identified eight T$_{RM}$ clusters, defined by varying combinations of T$_{RM}$ markers. In contrast to the sequential approach, several of these clusters (TRM 1, 3, 6 and 7) expressed the T$_{fh}$ marker FR4, which was restricted to T$_{fh}$, T$_{EM}$, and T$_{CM}$ in the sequential approach. The remaining T$_{RM}$ clusters (TRM 2, 4 and 5), were presumably canonical T$_{h}$1-like cells characteristic of a typical IAV response. Again, these subpopulations exhibited diverse loss rates (Fig. 7D),

with the KLRG1-expressing TRM 7 lost at the highest rate, and TRM 4 (characterized by CD69, CD49a and Bcl-2, and low levels of FR4) showing an intrinsic net positive growth rate.

We found evidence for flows from FR4^{lo} to FR4^{hi} subsets but not *vice versa*, indicating fluxes towards populations with characteristics of T follicular helper cells (TEM 1 to TRM 7 and T$_{fh}$; TRM 2 to TRM 1, TRM 4 to TRM 7 and 8, TRM 5 to T$_{fh}$). However, these populations had relatively high loss rates and as the population declined as a whole (Fig. 7F) they were outcompeted by Bcl-2-expressing subsets (TRM 4 and 8, and TEM 2; Fig. 7G), akin to the long-lived T$_{RM}$ in the CD8 data.

As described for CD8 T cells, the generative nature of the VAE provided a visual predictive check (Fig. 7H), showing an excellent correspondence between the data and the fitted model. The observed and predicted marginal distributions for surface markers corresponded well (Fig. S9A), with the exception of CD49a and CD27. These discrepancies were also noticeable when we reconstructed data using the VAE (Fig. S9B) where CD49a and CD27 had the lowest coefficient of determination ($R^2 = 0.16$ and 0.12, respectively). In contrast, CD69, Bcl-2 and FR4 were almost perfectly reconstructed ($R^2 > 0.97$).

Figure 7: Modeling the dynamics of lung CD4 cells with the integrated approach. Results are based on data from $n = 27$ mice. **A.** For each marker, the fluorescence intensity of each cell is displayed in a UMAP representation of the latent space. **B.** Cell density and 95% confidence ellipsoids for GMM components projected into the same UMAP space. Dots are the GMM location vectors. **C.** Mean fluorescence intensity of each marker and population identified by the GMM. **D.** Marginal posterior distributions of net loss rates, with medians and 95% CrIs. **E.** Differentiation matrix. **F.** Measured numbers of $CD44^+$ $CD11a^+$ protected $CD4^+$ T cells in lung (points), trajectory of the median prediction (line), 95% CrI (dark band), 95% posterior predictive interval (light band). **G.** Calculated population frequencies per mouse and timepoint, with median trajectories predicted by the model and 95% CrI (bands). **H.** UMAPs of marker expression data, either simulated using the fitted model (white background), or representing the observations (gray background).

Discussion

Developing methods to connect time series of high dimensional data with mechanistic, dynamic models of cell behavior is a pressing challenge. Here we described a straightforward 'sequential' approach in which one clusters and then models the occupancy of these clusters over time. Intuitively, though, this approach may be problematic when phenotypic diversity within a population does not permit clear separation of discrete subpopulations. To address this issue we explored a more holistic approach which allows for time-varying uncertainty in the assignment of cells to clusters, simultaneously modeling the cellular dynamics and its low dimensional representation.

In the setting of the T cell response to influenza A virus infection, the sequential approach showed that the net loss rates of CD8 T cells within the lung are strongly dependent on their phenotype and on time, and also found evidence for ongoing differentiation between cell types. The integrated approach partitioned CD8 T cell phenotypes and their dynamics in a similar way, but also provided more resolution. Specifically, we found that lung-resident CD8 T cells appear to be maintained by a Bcl-2^{hi} CD69^{hi} subset that lacks expression of the canonical T_{RM} marker CD103. Expression of Bcl-2 may be important for counteracting pro-apoptotic signals arising from the nutrient-poor environment of the airway[38]. It has been suggested that CD69^{hi} CD103^{hi} T_{RM} in the airway are maintained by T_{RM} in the lung parenchyma and that migration is mediated by increased CXCR6 expression[12]. Although here we did not perform bronchoalveolar lavage (BAL) to study cells within the airway directly, we did find evidence for differentiation within the parenchyma from these Bcl-2^{hi} CD69^{hi} T_{RM} into CD69^{hi} CD103^{hi} (DP) T_{RM}, which also co-expressed CXCR6 at intermediate levels.

We found that CD4 T cells within the lung were also dynamically and phenotypically heterogeneous, but with subsets less well defined than their CD8 T cell counterparts. In contrast with the CD8 results, the CD4 T cell data could be explained by either time-dependent loss of independent populations, or constant loss of differentiating populations. As with CD8 T cells, however, we found that Bcl-2-expressing populations come to dominate the CD4 T cells within the lung at later time points, and that these cells might be a source for other populations, thereby sustaining heterogeneity. Notably, we also saw evidence for one-way differentiation from canonical T_{RM}, which were largely T_h1-like, to FR4-expressing cells that were more T_{fh}-like.

Our flow cytometry panel contained five markers commonly associated with tissue residency (CD69, CD103, CD49a, CXCR6, and PD1), and others that are typically not expressed on canonical T_{RM} (CD62L, CX3CR1, and KLRG1). Other markers have been used to classify T_{RM} (such as CD101, CXCR3 and CXCR4[14]), and adding these to the panel might reveal more fine-grained subpopulations, each with distinct dynamics. For instance, it has been shown that T_{RM} in the airway rapidly lose their CD11a expression[12]. Airway T_{RM} also show distinct dynamics as they are lost more rapidly than T_{RM} in the parenchyma[38]. Here, we used CD11a to pre-select antigen-experienced cells in our gating strategy. If data from BAL were included, adding CD11a to the training data instead could be beneficial. Our inferences could also benefit from inclusion of the Ki67 marker, which is expressed transiently after cell division. Using this marker could allow us to disentangle

net loss rates (λ) into division (ρ; Fig. 3A) and death or egress (δ). However, this would require more complex models, and will be left for future work.

A typical workflow may benefit from using both the sequential and integrated methods. The sequential approach makes it easy to visualize the data as a set of timeseries, and hence aids in the the model development phase, and may be sufficient if the subsets are very clearly defined (such as the T_{CM} populations we identified here). The integrated approach is particularly powerful when standard clustering approaches do not clearly delineate phenotypes. In any case it is well suited for fine tuning of the model, parameter estimates and the population structure. Here we used the sequential approach for model selection and explored the integrated approach only with the favored models for CD8 and CD4 T cell dynamics. As the integrated approach is likelihood-based it should also lend itself to model comparison, but performing this is difficult for variational methods and remains an open challenge[39]. Facilitating model selection within the integrated approach would provide significant advantages. In the sequential approach, it is not possible to apply standard selection criteria to establish the optimal number of clusters, because the clustering step occurs before model fitting, and so different numbers of clusters represent entirely distinct datasets. In contrast, the integrated approach could provide an objective measure of the number of subpopulations present in the data, because it deals with the likelihood of the observations themselves.

A related issue with the integrated method is that, because models are fitted to high-dimensional data, it is challenging to perform posterior predictive checks. We used two-dimensional UMAP projections to address this issue (Fig. 6A), comparing the density of the observations in UMAP space with the density of pseudo-observations generated by the model at intermediate time points. A recent analysis of dimension reduction methods argued that for high-dimensional single-cell data the density in UMAP space corresponds poorly with cell density in the feature space[40]. However, the dimensionality of our data is an order of magnitude lower than the data used in their analyses. Therefore we are confident that, in the realm we explored here, using UMAP for posterior predictive checks is a reasonable procedure.

In both approaches, the model equations take the same functional form for all populations. In general this constraint will be too stringent. For instance, during the expansion phase of the immune response diversifying subsets of T cells may exhibit highly distinct dynamics, which may need to be modeled with structurally different equations. Further, programmed bursts of division characteristic of clonal expansion phase require multiple equations to describe a single population, one for each round of division[41,42]. Structural heterogeneity within models is easily accommodated by sequential approaches, because one assigns phenotypes to each of the clusters before model fitting. It is more problematic for the integrated approach; because we do not know *a priori* which cells will end up in which mixture component, an equation may end up describing the wrong cluster of cells. A potential solution is to use semi-supervised clustering[32]. Here we assign to a small set of cells a phenotype that we want to model with a particular type of equation. We then add a term to the likelihood function that makes it much more likely that those selected cells end up in the appropriate cluster.

Our approach is similar to scVI[31] and scANVI[32] in its use of a VAE, although our method is currently designed for flow cytometry data instead of very high dimensional single-cell RNA sequence data, which requires more complex error models. The use of hierarchical Gaussian mixture models for the analysis of flow cytometry data has been explored elsewhere[43], but here we fit a GMM on a lower-dimensional state space after applying a non-linear transformation. The transformed data might be more easily described by a GMM than the raw flow cytemetry data, especially when the latter is high-dimensional. Autoencoders for single-cell data analysis were first introduced as a de-noising and imputation tool[44], but have evolved rapidy. Currently there exists a large collection of methods based on variational autoencoders[31] which can analyze scRNA-seq data and other modalities. These methods can roughly be grouped into 'exploratory' and 'guided'[45]. Exploratory methods do not use any covariates of interest to guide latent state recovery or community detection, while guided methods explicitly incorporate such data[46]. Our integrated method falls into the guided class of models, as we use time information directly in the data manifold reconstruction, as well as community detection steps.

Understanding the dynamical properties of immune responses is key to solving many fundamental problems, ranging from finding cures to chronic diseases such HIV-1 infection, designing vaccines that provide heterosubtypic immunity to respiratory viruses such as influenza, and immunotherapies targeting tumors. Continuing technological advances are yielding bountiful data at the single cell level, resolving phenotypes at unprecedented scales and promising to advance our abilities to tackle these problems. Meanwhile, the deep learning revolution is presenting us with an expanding suite of tools to process and visualize and interpret these data. Mathematical modeling has a long tradition of aiding the interpretation of immunological and virological data, but has largely relied on heavily curated datasets. The approaches we present in this paper, and others like them, bring dynamical models closer to the observations. This bridging will increase the utility of mathematical models for shaping and testing our intuitions regarding biological processes.

Methods

Experimental procedures

Mice   Mice were housed in specific pathogen–free conditions in the animal facilities at Columbia University Irving Medical Center (CUIMC). Eight week old female C57BL/6 were purchased from the Jackson Laboratory. Infections were performed in biosafety level 2 biocontainment animal facilities. All animal studies were approved by Columbia University Institutional Animal Care and Use Committee.

Influenza infection   C57BL/6 mice were briefly anesthetized using inhaled isoflurane and infected intranasally with X31 influenza A virus at a 50% tissue-culture infectious dose (TCID50) of 5000 in 30μl of PBS. Infected mice were weighed daily and animals that lost more than 30%

of their starting weight were humanely euthanized. The infections were staggered such that all samples were processed and analyzed on the same day, to mitigate batch effects.

Tissue preparation   Mice were intravenously injected with 5μg of Brilliant Violet (BV) 421- conjugated anti-CD45.2 antibody 5 minute before cervical dislocation. Spleen and mediastinal lymph nodes (MedLNs) were processed by mechanical disruption and passed through 70μm filters (Corning). Single-cell suspensions of lungs were prepared first be mechanical disruption followed by digestion with 1mg/ml collagenase and DNAse (Sigma) for 40 minutes at 37°C. Lungs were then passed through a 70μm filter. Red blood cells were lysed from the spleen and lungs using ACK lysis buffer (Gibco).

Flow cytometry   Single cell suspensions were washed in PBS before staining with Zombie NIR viability dye (1/1000 in 100μl PBS) for 30 minutes at 4°C. Cells were washed twice and stained with 0.5μl of PE-labeled IA^b/$NP_{311\text{-}325}$ (CD4) and 1μl of APC-labeled $H\text{-}2D^b$/$NP_{366\text{-}374}$ (CD8) tetramers (National Institute of Health (NIH) Tetramer Core Facility (NTCF)) in 50μl FACS buffer (PBS with 2% heat inactivated FBS + 0.05mM EDTA). For surface staining, fluorochrome-conjugated antibodies (STAR Methods) in FACS buffer were added to cell suspensions for 20 minutes at 4°C protected from light, followed by washing with FACS-buffer. Cells were then fixed and permeabilized with the FoxP3 fix/perm kit (Tonbo) for 30 minutes at 4°C. Following fixation, cells were washed with FACS buffer, and stained with antibodies for intracellular detection (STAR Methods) in Perm Buffer (Tonbo) for 20 minutes at 4°C, followed by washing and resuspension in FACS buffer. Stained cells were acquired using the Cytek Aurora flow cytometer and analyzed using FlowJo version 10 software (Tree Star).

Congenic transfer experiment   Three CD90.1 and twelve CD90.2 6-8 week old C57BL/6 male mice were infected with X31 influenza virus (TCID50 5000 in 30 μl). At day 14 of infection, mediastinal lymph nodes and spleen cells were pooled from the three CD90.1 infected mice. From these two groups of pooled cells, T cells were isolated using a Stemcell Technologies EasySepTM T cell isolation kit. 10^6 pooled T cells were transferred into the infected CD90.2 mice via intraperitoneal injection and tissue was analyzed at days 2, 4, and 6 post-transfer ($n = 3$ per time point). As a control, 10^6 pooled T cells from age- and cage-matched uninfected CD90.1 mice were transferred into D14 infected CD90.2 mice and tissue was isolated at 2 days post-transfer ($n = 3$).

Models

Sequential approach

The classical analysis pipeline for our data is as follows. First, the data is pooled and clustered using a heuristic clustering method. Second, for each mouse, the number of cells in each of the clusters is counted. As the samples were taken at different times post infection, this results in a time series of cell counts per cluster. Third, dynamical models are fitted to these time series. The resulting parameter estimates and model comparison can give clues about the dynamics of the different T-cell populations in the lung.

For the first step (clustering), we used the established Phenograph method[47], which first constructs a K nearest-neighbor graph, and then uses the Leiden algorithm[22] for community detection. We ran the algorithm with number of neighbors set to 10 and resolution parameter set to 0.8. We used the implementation from the `scanpy` package[48].

For the third step (fitting models), we used the probabilistic programming language Stan (cmdstan version 2.34)[23,24]. Bayesian inference was done with the NUTS algorithm, using 5 parallel chains, 500 warm-up iterations and 500 posterior samples. To prevent divergent transitions, we set the "adapt delta" parameter to 0.9.

ODE models. We developed a number of different models, each making different assumptions about T-cell dynamics and differentiation. The most general model is given in terms of the following system of ODEs

$$\frac{dX}{dt} = -\lambda(t) \circ X + QX \tag{1}$$

and initial condition $X(t_0) = X_0$. Here $X \in \mathbb{R}_+^d$ is a vector of population sizes for each of the d clusters. The vector λ contains the net loss rates for each of the clusters, and $\circ$ denotes the Hadamard product (element-wise multiplication). The matrix Q determines T-cell differentiation, and Q_{ij} is the per-capita rate at which cells in cluster j differentiate into cells in cluster i. The matrix Q is a stochastic matrix in the sense that the columns add up to zero. Hence, the diagonal elements $Q_{ii} \leq 0$ denote the total efflux from cluster i due to differentiation.

We consider two different forms of the net loss rate λ. The simplest form is time-independent loss $\lambda(t) = \lambda_E$. This means that the loss rates of each population stays constant over time. However, each population can have its own distinct loss rate. For instance, effector cell counts might decrease faster than T_{RM} or T_{CM} cells.

In addition to a time-homogeneous (or autonomous) model, we also consider a model in which the net loss rate decreases exponentially to a smaller loss rate.

$$\lambda_i(t) = \lambda_{E,i} e^{-u(t-t_0)} + \lambda_{L,i}(1 - e^{-u(t-t_0)}) \tag{2}$$

The indices E and L refer to 'early' and 'late' stage dynamics. The speed at which this switch happens (u) is shared between clusters. In both the homogeneous case and the inhomogeneous case, we can compare models in which we set $Q = 0$ (i.e. the clusters form independent populations) to models in which we estimate the off-diagonal coefficients of the matrix Q (i.e. allowing for differentiation between clusters).

As the system of ODEs is linear, the initial value problem in Eqn. 1 can be solved explicitly. This convenient for inference, as we will not have to use numerical integrators (and in particular sensitivity or adjoint equations). The solution is given by

$$X(t) = \exp\left(-\int_{t_0}^{t} \text{diag}(\lambda(s))ds + Q(t - t_0)\right) X_0,$$ (3)

where exp denotes the matrix exponential, and $\text{diag}(\lambda)$ denotes a square, diagonal matrix with vector λ on the main diagonal. The integral of λ is given by

$$\int_{t_0}^{t} \lambda_i(s)ds = \begin{cases} \frac{\lambda_{E,i} - \lambda_{L,i}}{u}(1 - e^{-u(t-t_0)}) + \lambda_{L,i}(t - t_0) & \text{if } u \neq 0 \\ \lambda_{E,i}(t - t_0) & \text{if } u = 0 \text{ and } \lambda_L = 0 \end{cases}$$ (4)

Observation (or error) model. In order to compare our dynamical model to the data, we need to specify a likelihood function. We have two data streams: For each sample (mouse) s, we have the number of cells per cluster i, denoted $K_{s,i}$. Additionally, we have an estimate of the total number of T cells in the tissue, denoted M_s. According to the model, the true cluster frequencies are given by $\pi_i(t_s) = X_i(t_s)/Y(t_s)$, where $Y = \sum_{j=1}^{d} X_j$ denotes the total number of T cells. We then assume that the the count vector K_s is sampled from a Dirichlet-multinomial distribution

$$K_s \sim \text{DirMult}(N_s, \pi(t_s)\tau_K)$$ (5)

where $N_s = \sum_{i=1}^{d} K_{s,i}$ is the total number of T cells in sample s, and τ_K is a dispersion parameter. Explicitly, the probability of sampling K_s is equal to $N_s B(N_s, \tau_K)/\prod_{i:K_{s,i}>0} K_{s,i} B(K_{s,i}, \tau_K \pi_i(t_s))$, where B is the Beta function. The dispersion parameter τ_K determines the over-dispersion of the Dirichlet-multinomial distribution, which converges to a multiniomial distribution as $\tau_K \to \infty$. We use an over-dispersed distribution because of the significant biological and technical variation between our samples[49].

We assume that the total number of cells in the tissue (M_s) has a log-normal distribution

$$M_s \sim \text{LogNorm}(\log(Y(t_s)), \sigma_M)$$ (6)

where Y is the total number of cells predicted by the dynamical model. The scale parameter σ_M is determined by the technical and biological variation between the experiments. In addition to a likelihood function, we have to specify prior distributions for the parameters, which are listed in Table S3.

Model comparison. For model comparison, we used the Pareto-smoothed importance sampling (PSIS) approximation of leave-one-out (LOO) cross validation[28]. We used the method implemented in the python package Arviz[50]. The PSIS approximation breaks down for highly influential observations, and Arviz package provides a diagnostic for when this happens. For these observations, we replaced the PSIS approximation with the true LOO-CV statistic by re-fitting the model.

Integrated approach

We estimate parameters of our ODE model and a set of "nuisance" parameters (NN weights and biases, GMM location vectors) simultaneously in a variational inference framework. We approximate the joint posterior distribution of the latent cell states, the ODE model parameters, and the GMM components with a so-called variational distribution, and optimize the variational parameters using stochastic gradient ascent. This procedure is outlined in Fig. 4B. The flow cytometry data is used together with the encoder network to generate a random sample of latent cell states z. At the same time, we also sample ODE parameters and GMM location parameters from the variational distribution. Using these parameters, we predict the trajectories of the T-cell population sizes at each sampling time. The total predicted population size is used to compute the likelihood of the total cell counts. The relative population sizes give us the weights of the GMM model, and allow us to compute the prior density for each latent cell state. The decoder network is used compute the likelihood of the flow cytometry data, given the sampled cell states. The prior density and likelihood are combined into the evidence lower bound (ELBO), the objective function of variational inference, which we explicitly define below. Using the gradient of the objective function, the variational parameters are perturbed in the direction of a larger ELBO at which point we can sample new parameters. We repeat this procedure until the ELBO converges.

In our integrated approach, we combine clustering with fitting of the dynamical model in a single machine learning algorithm. We use a variational auto-encoder (VAE) to infer a lower dimensional representation z of the marker expression data x. We then simultaneously fit a Gaussian mixture model (GMM) to the latent vectors z[33]. Each of the d components of this GMM represents a T-cell cluster. This means that for each data point x_n, we can compute the probabilities $p_{n,i}$ that cell n belongs to population i using Bayes rule

$$p_{n,i} = \frac{\pi_i(t_n)\psi_i(z_n)}{\pi_1(t_n)\psi_1(z_n) + \cdots + \pi_d(t_n)\psi_d(z_n)}. \tag{7}$$

In this equation, ψ_i denote the probability density functions of the Gaussian mixture components, each parameterized with a mean vector μ_i and covariance matrix Σ_i, and π_i represents the weight of mixture component i. In our method, π is a function of time and is determined by a dynamical model as in the sequential approach: $\pi(t) = X(t)/Y(t)$.

The VAE consists of a decoder network D and a encoder network E. The decoder network D : $(z, s) \mapsto (\mu_x, \sigma_x)$ takes as an argument a latent vector z and the one-hot-encoded batch information s, and returns vectors $\mu_x \in \mathbb{R}^m$ and $\sigma_x \in \mathbb{R}^m_+$ which we can use to sample a vector x in the feature

space.

$$x \sim \mathcal{N}(\mu_x, \sigma_x). \tag{8}$$

or compute the likelihood of an observed expression vector x, given latent vector z. The decoder network D has one hidden layer with 20 nodes and a softplus non-linearity. This hidden layer is shared by the two output layers for μ_x and σ_x. We use a softplus link function to ensure that the scale σ_x is positive. The encoder network $E : (x, s, t) \mapsto (\mu_z, \sigma_z)$ is similar in architecture to D, but takes as an input the expression vector x, the batch information s, and the sampling time t and returns a location and scale vector for the latent vector z, distributed as $z \sim \mathcal{N}(\mu_z, \sigma_z)$. The encoder depends on the sampling time, because it approximates the posterior distribution of z, which in turn depends on the time-dependent prior distribution (the GMM).

Target of optimization: ELBO. The goal of variational inference is to find an approximate posterior distribution q that is as close as possible to the true posterior, p. The Kullback-Leibler (KL) divergence provides a measure of their similarity;

$$D_{KL}(q_\phi(\theta) \| p_\phi(\theta|x)) = \mathbb{E}_{q_\phi(z,\theta)} \left[\log(q_\phi(\theta)) - \log(p_\phi(\theta|x)) \right]. \tag{9}$$

The posterior distribution is given by

$$p_\phi(\theta|x) = \frac{\pi_\phi(\theta) L(x|\theta, \phi)}{p_\phi(x)}, \tag{10}$$

where $p_\phi(x)$ is the evidence of the model, given the data. We would like to find a parameter vector ϕ that maximizes this evidence, but unfortunately $p_\phi(x) = \mathbb{E}_{\pi_\phi(\theta)}[L(x|\theta, \phi)]$ is generally intractable and difficult to approximate directly. This is where the variational distribution is useful. As the KL-divergence is positive, we have

$$\begin{aligned} \log(p_\phi(x)) &\geq \log(p_\phi(x)) - D_{KL}(q_\phi(\theta) \| p_\phi(\theta|x)) \\ &= \mathbb{E}_{q_\phi(\theta)}[\log(\pi_\phi(\theta) L(x|\theta, \phi)) - \log(q_\phi(\theta))]. \end{aligned} \tag{11}$$

This final quantity is called the evidence lower bound (ELBO). If we then maximize the ELBO, we both minimize the KL-divergence, and given that our variational distribution is close to the posterior, we also maximize the (log) evidence of the model.

To specify the ELBO for our model we have to define three quantities: the prior π_ϕ, the likelihood function L and the variational distribution q_ϕ. First, the prior factors into three components

$$\pi_\phi(\theta) = \pi_{\text{Dyn}}(\theta_{\text{Dyn}}) \times \pi_{\text{GMM}}(\theta_{\text{GMM}}) \times \pi_{\text{Lat}}(z|\theta_{\text{Dyn}}, \theta_{\text{GMM}}), \tag{12}$$

corresponding to the prior for the dynamical model parameters, the GMM parameters, and the prior for the latent vectors z, respectively. The prior and parameters for the dynamical model are specified in Table S3. Next, the parameters for the GMM consist of the location (or mean) vectors μ_i in the latent space of the Gaussian components, and their covariance matrices Σ_i. The component

weights are not free parameters, but are derived from the dynamical model. The location vectors have prior distributions $\mu_i \sim \mathcal{N}_k(0, I_k)$, where I_k is the k-dimensional identity matrix. To reduce identifiability issues, all components i have covariance matrix $\Sigma_i = I_k$.

Finally, the prior for the latent vectors z is given by the GMM with density

$$\pi_{\text{Lat}}(z_n|\theta_{\text{Dyn}}, \theta_{\text{GMM}}) = \sum_{i=1}^{d} \pi_i(t_n)(2\pi)^{-k/2} \det(\Sigma_i)^{-1/2} \exp\left(-\tfrac{1}{2}(z - \mu_i)\Sigma_i^{-1}(z - \mu_i)^T\right). \tag{13}$$

Here t_n is the time point that cell n was sampled. We assume that all latent vectors of cells z_n are independent. Hence, we ignore any potential dependencies due to common ancestry[51].

The likelihood function L consists of two factors

$$L(x, M|s, \theta, \phi) = L_{\text{Expr}}(x|s, \theta, \phi) \times L_{\text{Count}}(M|\theta, \phi) \tag{14}$$

which are the likelihood of marker expression data x_n, and the likelihood of cell counts M_n, respectively. For the likelihood of the expression data, we start with a latent vector z_n, and pass this through the decoder network D_ϕ to get parameters for the likelihood of x_n:

$$x_n \sim \mathcal{N}_m(\mu_x, \sigma_x), \quad (\mu_x, \sigma_x) = D_\phi(z_n, s_n). \tag{15}$$

The likelihood of the count data is given by

$$M_n \sim \text{LogNorm}(Y(t_n), \sigma_M). \tag{16}$$

Finally, in order to perform inference, we have to specify a variational distribution $q_\phi(\theta)$. We assume that the variational distribution can be factorized as

$$q_\phi(\theta) = q_{\text{Dyn}}(\theta_{\text{Dyn}}|\phi_{\text{Dyn}}) \times q_{\text{GMM}}(\theta_{\text{GMM}}|\phi_{\text{GMM}}) \times q_{\text{Lat}}(z|\phi_{\text{Lat}}, x), \tag{17}$$

with factors for the dynamical model parameters, the GMM parameters and the latent vectors, respectively. For the (possibly transformed) dynamical model parameter vector

$$\theta_{\text{IVP}} = \left(\rho_0, \eta, \log(u), \log(Q), \log(X_0), \mu_\rho, \log(\sigma_\rho), \mu_\eta, \log(\sigma_\eta), \mu_Q\right),$$

and σ_M, we assume a multivariate normal and log-normal distribution, respectively:

$$\theta_{\text{IVP}} \sim \mathcal{N}\left(\mu_{\text{IVP}}, \Sigma_{\text{IVP}}\right), \quad \sigma_M \sim \text{LogNorm}(\mu_{\sigma_M}, \sigma_{\sigma_M}). \tag{18}$$

Next, we use a multivariate normal distribution for the location vectors of the mixture components;

$$(\mu_1, \ldots, \mu_d) \sim \mathcal{N}_{kd}(\mu_{\text{GMM}}, \Sigma_{\text{GMM}}). \tag{19}$$

Finally, we have to specify a variational distribution for z. As we have a latent variable z_n for every observed marker expression vector x_n, we use the method of amortization; instead of specifying

variational parameter for each individual variable z_n, we define a function E that calculates variational parameters σ_z and μ_z, based on the data point x_n:

$$z_n \sim \mathcal{N}_k(\mu_z, \sigma_z) \,, \quad (\mu_z, \sigma_z) = E_{\phi_{\text{Lat}}}(x_n, s_n). \tag{20}$$

The function E is defined in terms of a neural network.

We implemented our method in the Pyro language for deep probabilistic programming[35]. We fitted the model to our data using stochastic variational inference (SVI). We used the Adam optimizer[52] with learning rate 10^{-3} and 5×10^5 epochs, and used 10 parallel particles to evaluate the ∇ELBO. The neural network weights and biases were L_2-regularized. All our code is available on `https://github.com/chvandorp/scdynsys.git`.

Acknowledgments

This work was funded by the National Institutes of Health (NIAID), U01 AI150680 and R01 AI093870.

References

1. Bouveyron C and Brunet-Saumard C. Model-based clustering of high-dimensional data: A review. Computational Statistics & Data Analysis 2014;71:52–78.

2. Karim MR, Beyan O, Zappa A, et al. Deep learning-based clustering approaches for bioinformatics. Briefings in Bioinformatics 2021;22:393–415.

3. Lam N, Lee Y, and Farber DL. A guide to adaptive immune memory. Nature Reviews Immunology 2024:1–20.

4. Antia R, Pilyugin SS, and Ahmed R. Models of immune memory: On the role of cross-reactive stimulation, competition, and homeostasis in maintaining immune memory. Proceedings of the National Academy of Sciences 1998;95:14926–31.

5. De Boer RJ, Homann D, and Perelson AS. Different Dynamics of CD4+ and CD8+ T Cell Responses During and After Acute Lymphocytic Choriomeningitis Virus Infection 1. The Journal of Immunology 2003;171:3928–35.

6. Amo PCd, Lahoz-Beneytez J, Boelen L, et al. Human TSCM cell dynamics in vivo are compatible with long-lived immunological memory and stemness. PLOS Biology 2018;16:e2005523.

7. Swain AC, Borghans JA, and Boer RJ de. Effect of cellular aging on memory T-cell homeostasis. Frontiers in Immunology 2022;13.

8. Gossel G, Hogan T, Cownden D, Seddon B, and Yates AJ. Memory CD4 T cell subsets are kinetically heterogeneous and replenished from naive T cells at high levels. eLife 2017;6. Ed. by Boer R de:e23013.

9. Paik DH and Farber DL. Anti-viral protective capacity of tissue resident memory T cells. Curr Opin Virol 2021;46:20–6.

10. Morris SE, Farber DL, and Yates AJ. Tissue-Resident Memory T Cells in Mice and Humans: Towards a Quantitative Ecology. The Journal of Immunology 2019;203:2561–9.

11. Szabo PA, Miron M, and Farber DL. Location, location, location: Tissue resident memory T cells in mice and humans. Science Immunology 2019;4:eaas9673.

12. Wein AN, McMaster SR, Takamura S, et al. CXCR6 regulates localization of tissue-resident memory CD8 T cells to the airways. Journal of Experimental Medicine 2019;216:2748–62.

13. Kumar BV, Ma W, Miron M, et al. Human Tissue-Resident Memory T Cells Are Defined by Core Transcriptional and Functional Signatures in Lymphoid and Mucosal Sites. Cell Reports 2017;20:2921–34.

14. Pizzolla A and Wakim LM. Memory T Cell Dynamics in the Lung during Influenza Virus Infection. J Immunol 2019;202:374–81.

15. Bachireddy P, Azizi E, Burdziak C, et al. Mapping the evolution of T cell states during response and resistance to adoptive cellular therapy. Cell Reports 2021;37:109992.

16. Kingma DP and Welling M. An Introduction to Variational Autoencoders. Foundations and Trends® in Machine Learning 2019;12:307–92.

17. Teijaro JR, Turner D, Pham Q, Wherry EJ, Lefrançois L, and Farber DL. Cutting Edge: Tissue-Retentive Lung Memory CD4 T Cells Mediate Optimal Protection to Respiratory Virus Infection. The Journal of Immunology 2011;187:5510–4.

18. Wu T, Hu Y, Lee YT, et al. Lung-resident memory CD8 T cells (TRM) are indispensable for optimal cross-protection against pulmonary virus infection. J Leukoc Biol 2014;95:215–24.

19. Turner DL, Bickham KL, Thome JJ, et al. Lung niches for the generation and maintenance of tissue-resident memory T cells. Mucosal Immunology 2014;7:501–10.

20. Pizzolla A, Nguyen THO, Smith JM, et al. Resident memory CD8$^+$ T cells in the upper respiratory tract prevent pulmonary influenza virus infection. Science Immunology 2017;2:eaam6970.

21. Fiege JK, Stone IA, Fay EJ, et al. The Impact of TCR Signal Strength on Resident Memory T Cell Formation during Influenza Virus Infection. J Immunol 2019;203:936–45.

22. Traag VA, Waltman L, and Eck NJ van. From Louvain to Leiden: guaranteeing well-connected communities. Sci Rep 2019;9:5233.

23. Carpenter B, Gelman A, Hoffman M, et al. Stan: A probabilistic programming language. J. Stat. Softw. 2017;76:1–32.

24. Team SD. Stan Modeling Language Users Guide and Reference Manual, version 2.34. 2024.

25. McInnes L, Healy J, and Melville J. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. arXiv:1802.03426 [cs, stat]. 2020.

26. Zwijnenburg AJ, Pokharel J, ė R, et al. Graded expression of the chemokine receptor CX3CR1 marks differentiation states of human and murine T cells and enables cross-species interpretation. Immunity 2023;56:1955–74.

27. Gerlach C, Moseman EA, Loughhead SM, et al. The Chemokine Receptor CX3CR1 Defines Three Antigen-Experienced CD8 T Cell Subsets with Distinct Roles in Immune Surveillance and Homeostasis. Immunity 2016;45:1270–84.

28. Vehtari A, Gelman A, and Gabry J. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. Statistics and Computing 2017;27:1413–32.

29. Tibshirani R. Regression Shrinkage and Selection via the Lasso. Journal of the Royal Statistical Society. Series B (Methodological) 1996;58:267–88.

30. Yao Y, Vehtari A, Simpson D, and Gelman A. Using Stacking to Average Bayesian Predictive Distributions (with Discussion). Bayesian Analysis 2018;13:917–1007.

31. Lopez R, Regier J, Cole MB, Jordan MI, and Yosef N. Deep generative modeling for single-cell transcriptomics. Nature Methods 2018;15:1053–8.

32. Xu C, Lopez R, Mehlman E, Regier J, Jordan MI, and Yosef N. Probabilistic harmonization and annotation of single-cell transcriptomics data with deep generative models. Mol Syst Biol 2021;17:e9620.

33. Jiang Z, Zheng Y, Tan H, Tang B, and Zhou H. Variational Deep Embedding: An Unsupervised and Generative Approach to Clustering. In: IJCAI'17. Place: Melbourne, Australia. AAAI Press, 2017:1965–72.

34. Kopf A, Fortuin V, Somnath VR, and Claassen M. Mixture-of-Experts Variational Autoencoder for clustering and generating from similarity-based representations on single cell data. PLoS Comput Biol 2021;17:e1009086.

35. Bingham E, Chen JP, Jankowiak M, et al. Pyro: Deep Universal Probabilistic Programming. Journal of Machine Learning Research 2018.

36. Goodfellow I, Bengio Y, and Courville A. Deep Learning. MIT Press, 2016.

37. Son YM, Cheon IS, Wu Y, et al. Tissue-resident CD4+ T helper cells assist the development of protective respiratory B and CD8+ T cell memory responses. Science Immunology 2021;6:eabb6852.

38. Hayward SL, Scharer CD, Cartwright EK, et al. Environmental cues regulate epigenetic reprogramming of airway-resident memory CD8+ T cells. Nature Immunology 2020;21:309–20.

39. Yao Y, Vehtari A, Simpson D, and Gelman A. Yes, but Did It Work?: Evaluating Variational Inference. In: *Proceedings of the 35th International Conference on Machine Learning*. ISSN: 2640-3498. PMLR, 2018:5581–90.

40. Chari T and Pachter L. The specious art of single-cell genomics. PLOS Computational Biology 2023;19:1–20.

41. Antia R, Bergstrom CT, Pilyugin SS, Kaech SM, and Ahmed R. Models of CD8+ Responses: 1. What is the Antigen-independent Proliferation Program. Journal of Theoretical Biology 2003;221:585–98.

42. Bresser K, Kok L, Swain AC, et al. Replicative history marks transcriptional and functional disparity in the CD8+ T cell memory pool. Nature Immunology 2022;23:791–801.

43. Johnsson K, Wallin J, and Fontes M. BayesFlow: latent modeling of flow cytometry cell populations. BMC Bioinformatics 2016;17:25.

44. Eraslan G, Simon LM, Mircea M, Mueller NS, and Theis FJ. Single-cell RNA-seq denoising using a deep count autoencoder. Nature Communications 2019;10:390.

45. Boyeau P, Hong J, Gayoso A, Jordan MI, Azizi E, and Yosef N. Deep generative modeling for quantifying sample-level heterogeneity in single-cell omics. Pages: 2022.10.04.510898 Section: New Results. 2022.

46. Zhao J, Jaffe A, Li H, et al. Detection of differentially abundant cell subpopulations in scRNA-seq data. Proc Natl Acad Sci U S A 2021;118.

47. Levine JH, Simonds EF, Bendall SC, et al. Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. Cell 2015;162:184–97.

48. Wolf FA, Angerer P, and Theis FJ. SCANPY: large-scale single-cell gene expression data analysis. Genome Biol 2018;19:15.

49. Bullock ME, Hogan T, Morris S, et al. Cell age, not chronological age, governs the dynamics and longevity of circulating CD4+ memory T cells. Pages: 2023.10.16.562650 Section: New Results. 2023.

50. Kumar R, Carroll C, Hartikainen A, and Martin O. ArviZ a unified library for exploratory analysis of Bayesian models in Python. Journal of Open Source Software 2019;4:1143.

51. Stadler T, Pybus OG, and Stumpf MPH. Phylodynamics for cell biologists. Science 2021;371.

52. Kingma DP and Ba J. Adam: A Method for Stochastic Optimization. arXiv:1412.6980 [cs]. 2017.

53. Perelson AS, Essunger P, Cao Y, et al. Decay characteristics of HIV-1-infected compartments during combination therapy. Nature 1997;387:188–91.

54. Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research 2011;12:2825–30.

Supplementary Information

Figure S1: Gating Strategy for CD8$^+$ and conventional CD4$^+$ antigen experienced T cells residing in the lung.

Figure S2: Weight curves of mice after IAV challenge and assessment of I.V. labeling. Each cohort contained 4 or 5 mice. In total data from 38 mice is presented. **A.** The curves indicate the percentage of the weight relative to that on the day of infection. The red curves correspond to mice that were excluded from further analysis due to lack of weight loss. **B.** Distribution of the IV label (CD45.2 IV) for each mouse. The red curves correspond to mice that were excluded from further analysis due to poor IV labeling.

Figure S3: Validation of population assignment in the sequential approach. Results are based on data from $n = 27$ mice. **A.** Entropy of CD8 T cell population assignment based on 20 Leiden clustering runs with different random seeds. Black dots corresponds to very certain assignments, yellow dots to highly uncertain assignments. The colored contours indicate the location of the different sub-populations in the UMAP. **B.** Entropy distribution per cluster. The bar plots show the median, IQR and 2.5 - 97.5 percentile range. The color of the bars and labels correspond to the contours in panel A. **C.** Distribution of IAV NP-specific CD8 T cells in the UMAP with contours indicating clusters. Gray dots indicate bulk antigen-experienced cells. **D-F.** Same as panels A-C, but for CD4 T cell data.

Figure S4: Timecourses of NP-specific T cells. Results are based on data from $n = 34$ mice. **A.** Number of NP-specific CD8 T cells, as a function of post-infection sampling time. The curve represents a spline fit to the log-transformed T cell counts, and the dashed lines represent the 95% confidence envelope (estimated by bootstrapping residuals). The red curve indicates the number of polyclonal CD8 T cells (cf. Fig. 1B). **B.** The Tet$^+$ fraction of bulk CD8 T cells in the lung niche. Splines are fitted on the logit scale. **C.** Number of NP-specific CD4 T cells. **D.** The Tet$^+$ fraction of bulk CD4 T cells.

Figure S5: Congenic transfer experiment to assess the amount of ingress during the memory phase. Results are based on data from $n = 12$ mice in total, with 3 mice per group. **A.** Design of the congenic transfer experiment. **B.** Numbers of protected and labeled, antigen-experienced CD8 and CD4 T cells from host and donors combined (upper panels) and donors only (lower panels). We fitted a log-linear model to the cell counts (solid line: ML estimate, dashed lines: 95% confidence envelope). The indicated λ the estimated net loss rate ($\pm$ standard error).

Figure S6: Comparison of results from the integrated and sequential approaches. Results are based on data from $n = 27$ mice. The sequential (SA) and integrated (IA) approaches both assign each cell to a subpopulation, but for any given cell this assignment may differ between the approaches. To quantify the similarity, we calculated the Jaccard index for each pair of sub-populations (one from the IA and the other from the SA). The Jaccard index is ratio of the number of cells that are in both clusters, and the number of cells that are in either one of the clusters. **A.** Results for CD8 T cell data. **B.** Results for CD4 T cell data.

Figure S7: Pre-processing CD4$^+$ T cell flow cytometry data for the sequential approach. Results are based on data from $n = 27$ mice. **A.** Marker expression heatmap for selected markers and consensus T-cell populations. **B.** UMAP of the marker expression data, colored by annotation. **C.** UMAPs of marker expression data, split by day post infection (DPI). The color scale reflects cell density in UMAP space. **D.** Time series of the fraction of cells in each cluster. The lines show a linear interpolation on the log scale.

Figure S8: Model fits to CD4 T cell timeseries using the sequential approach. Results are based on data from $n = 27$ mice. **A.** Data and predictions from fitted models. Top panels show total antigen-experienced CD4 T cell counts in the lung, other panels show relative population sizes of each of the sub-populations. **B.** Parameter estimates using model II. **C.** Parameter estimates using model III. **D.** Estimated differentiation matrix in model III.

Figure S9: Posterior predictive checks for the integrated approach with CD4 T cell data. Results are based on data from $n = 27$ mice. **A.** Marginal distributions of marker expression (cf. Fig. 1, panel I). Data is shown in black, simulated data is shown in red. **B.** Input data and reconstruction using the auto-encoder model. The number in brackets is the coefficient of determination (R^2).

Figure S10: Model fits to cell count data alone. Results are based on data from $n = 27$ mice. **A.** time-homogeneous model with a single compartment (i.e. a log-linear model) fit to CD8 T-cell count data (dots). The model fit is shown as a blue curve (posterior median), with 95% CrI as a dark-blue band. The light-blue band shows the posterior predictive interval (i.e. simulated observations). **B.** Fit of model with a single compartment, but with time-dependent net loss rates $\lambda(t)$. **C.** Fit of time-homogeneous model with two compartments. The population sizes (posterior median) of the two populations are shown as black curves. **D-F.** Fits of the three model to CD4 T cell counts.

Figure S11: The effect of batch correction in the integrated approach. Results are based on data from $n = 27$ mice. Shown is the entropy of the experimental batch distribution around each cell, using the latent vector z and its nearest neighbors. Values for batch-corrected latent vectors are shown in blue. Mock corrected values are shown in orange, and values for randomized batch information are shown in green. Panels A and B show CD8 and CD4 data, respectively. The distributions are capped at 1.5 as there was a very small number of cells with high relative entropy.

SI Text A   Ingress contributes minimally to the population dynamics of lung T$_{RM}$ during the memory phase

To inform the construction of models of cell dynamics within the lung, we aimed to quantify the extent to which they were supplemented by new immigrants following the peak of infection. Our approach is illustrated in Fig. S5A. To do this we simultaneously infected cohorts of CD90.1 and CD90.2 congenic mice with IAV, and at day 14 post infection, transferred 10^6 cells from the spleens and mediastinal lymph nodes of the CD90.1 (donor) mice to the CD90.2 (host) mice (see Methods). At 16, 18 and 20 days post infection (2, 4, and 6 days post transfer, respectively), lung tissue was collected from the host mice, and analyzed with flow cytometry. The kinetics of accumulation of any donor cells in the host lung tissue would then reflect the degree of ongoing recruitment of new T$_{RM}$ from circulating influenza-specific memory T cells.

Between 16 and 20 days post infection, the total numbers of CD4 and CD8 T cells (host+donor) fell by roughly 50% (Fig. S5B), consistent with the rates of decline observed in Fig. 1B-C. During the same time window, donor-derived cells were detectable in the lung but at low numbers that represented between 0.05-0.01% of transferred cells, and remained stable in number between 2-6 days post transfer. Between 16-18 DPI the rate of increase was 0.2 per day (95% CI $[-0.1, 0.4]$) for CD8$^+$ T cells and -0.2 per day for CD4$^+$ T cells (95% CI $[-0.5, 0.2]$). The number of donor cells in circulation (i.v. labeled cells from the lung sample) were consistently higher than cells in the tissue (CD8: 4.7 fold, $p = 2 \times 10^{-4}$; CD4: 3 fold, $p = 4 \times 10^{-3}$ paired t-test), while the total number of cells in circulation and in tissue were comparable (CD8: $p = 0.13$, CD4: $p = 0.1$). The number of donor cells in the lung tissue from uninfected donors was comparable to the number of cells from infected donors, indicating that the rate of ingress does not depend on IAV-specificity. We conclude that ingress of antigen-experienced T cells into the lung occurs at very low levels from at least 14 DPI onward.

SI Text B   A two-compartment, time-homogeneous model can explain the timecourse of CD8 T cell numbers

At first glance, the CD8 and CD4 T cell count data resemble typical biphasic exponential decay patterns[53]. Such time series can easily be modeled using two exponentially declining populations that decrease at two different rates. Initially, the more rapidly declining population is more abundant, but soon it is replaced (relatively) by the population that declines at a slower rate.

Because of this resemblance, it might be surprising that we need time-dependent loss rates or differentiation to model the T-cell populations in the lung. To make this initial expectation more rigorous, we fit three different models (denoted I-1, II and I-2) to the CD8 and CD4 T-cell count data alone. Model I-1 is time-homogeneous with only a single population (Fig S10A and D). This model can not describe the biphasic decay pattern seen in the data, and requires a large standard deviation for the error model (CD8: $\sigma_M = 0.86$, 95% CrI $[0.67, 1.17]$; CD4: $\sigma_M = 0.45$, 95% CrI $[0.35, 0.61]$).

Model II again has a single population, but now we include a time-dependent net loss rate $\lambda(t)$ as in Eqn. 2. This model fits the T-cell count data very well (Fig S10B and E, and requires a much smaller standard deviation for the error model (CD8: $\sigma_M = 0.34$, 95% CrI $[0.26, 0.46]$; CD4: $\sigma_M = 0.25$, 95% CrI $[0.19, 0.36]$). Finally, model I-2 has constant decay rates, but assumes that there are two distinct T-cell populations that decay at different rates. Judging from the posterior predictive check (Fig. S10C and E), this model fits the data as well as model II, and requires similar σ_M values as model II.

In terms of LOO-IC, model II and I-2 are indistinguishable (CD8: ΔLOO-IC $= 0.3 \pm 1.5$; CD4: ΔLOO-IC $= 0.3 \pm 0.5$), while model I-1 is significantly worse (CD8: ΔLOO-IC $= 24 \pm 3$; CD4: ΔLOO-IC $= 15 \pm 3$). This means that model II and I-2 describe the count data equally well, and hence using count data alone, it is not possible to distinguish between a model with two populations or a time-dependent net loss rate.

SI Text C The geometry of time-homogeneous loss of independent populations

The simplest model we considered has a geometric property that allows it to be easily tested against data. Suppose that the populations of T cells are independent (i.e. no differentiation, or $Q = 0$) and that their net loss rates are constant (i.e. time homogeneous). The model is then

$$\frac{dX}{dt} = -\lambda \circ X \tag{1}$$

with initial condition $X(t_0) = X_0$. This model admits the following solution

$$X(t) = X_0 \circ \exp(-\lambda(t - t_0)), \tag{2}$$

where the exponential function is taken element-wise. Again, we write $Y(t) = \sum_{i=1}^{d} X_i(t)$ for the total population size and $\pi_i(t) = X_i(t)/Y(t)$ for the population fractions.

When we look at the trajectories $\pi_i(t)$ on a logarithmic scale, a striking property is that they are all concave (Fig 3B and Fig. S8A, first column). We can see this mathematically as follows. A twice differentiable function is concave if the second derivative is non-positive. As the log of $\pi_i(t)$ is given by

$$\log \pi_i(t) = \log(X_{0,i}) + -\lambda_i t - \log Y(t), \tag{3}$$

the second derivative is given by

$$\frac{d^2}{dt^2} \log \pi_i(t) = -\frac{d^2}{dt^2} \log Y(t), \tag{4}$$

which does not depend on the population index i. The second derivative of $\log Y(t)$ is given by

$$\frac{d^2}{dt^2} \log Y(t) = \frac{Y(t)Y''(t) - Y'(t)^2}{Y(t)^2} \tag{5}$$

so its sign is determined by $Y(t)Y''(t) - Y'(t)^2$. We have $Y'(t) = -\sum_{i=1}^{d} \lambda_i X_i(t)$, and $Y''(t) = \sum_{i=1}^{d} \lambda_i^2 X_i(t)$. Now consider the vectors a and b given by $a_i = -\lambda_i \sqrt{X_i}$ and $b_i = \sqrt{X_i}$. By the Cauchy-Schwartz inequality, we have $\langle a, b \rangle^2 \leq \langle a, a \rangle \langle b, b \rangle$, and hence

$$Y'(t)^2 = \left(-\sum_{i=1}^{d} \lambda_i X_i \right)^2 \leq \sum_{i=1}^{d} \lambda_i^2 X_i(t) \sum_{i=1}^{d} X_i = Y''(t)Y(t) \tag{6}$$

which means that $\frac{d^2}{dt^2} \log Y(t) \geq 0$. Hence $\log \pi_i(t)$ is concave.

SI Text D   Batch correction

To measure how well batch correction performed in our integrated approach, we wanted to quantify the extent to which cells from different mice were well-mixed in phenotypic space[32]. To do this, we computed the K-nearest-neighbor graph of the latent representation z_i of the cells using the `scikit-learn` package[54]. For each cell i, we then counted the number of neighbors $k_{i,s}$ derived from mouse s (including the focal cell). If cells are well mixed, all of these counts should be distributed according to the sample sizes n_s of the animals s. A convenient measure of how well the distributions of the $k_{i,s}$ match is the relative entropy, given by

$$H_i = -\sum_s n_s/N \left(\log(k_{i,s}/K) - \log(n_s/N)\right),\qquad(7)$$

where $N = \sum_s n_s$ is the total number of cells from all animals. A lower relative entropy corresponds to more homogeneous mixing.

Fig. S11 shows the distribution of H_i as a blue histogram for the CD8 (panel A) and CD4 data (panel B). These distributions are difficult to interpret in isolation and so we compared them with two extremes. First, we computed the distribution of H_i under the assumption that all cells from all animals were homogeneously distributed. In this case we would have $\vec{k}_i \sim \text{Multinomial}(K, \vec{n}/N)$. We randomly generated samples for each cell i and the resulting distribution is shown as an green histogram in Fig. S11. The other extreme is the case when we switch off the effect of batch-correction. To accomplish this, we picked the animal s^* with the largest sample size n_{s^*} as a reference. We then used the encoder network to compute latent representations z_i from the pairs (x_i, s^*), instead of the usual pairs (x_i, s_i). This means that we simulate the scenario in which all cells came from the same animal s^*. The resulting distribution of H_i is shown as an orange histogram in Fig. S11.

Our relative entropy-based analysis shows that batch correction is effective at aligning the distributions of cells from different animals within the latent space. However, the relative entropy remains much higher than expected under a perfectly homogeneous distribution (green vs. orange histogram in Fig. S11). This is due to the fact that mice are sampled at different DPI, and the phenotype distribution is dependent on time, and possibly because all data was collected at the same time (Fig. 1), reducing batch effects.

Supporting Tables

Model	elpd_loo	p_loo	Δelpd	se Δelpd
CD8 T-cell data				
IV	-1219.1	28.7	-	-
II	-1254.4	24.8	35.3	8.8
III	-1287.7	18.7	68.6	12.5
I	-1372.8	22.3	153.8	29.9
CD4 T-cell data				
IV	-1807.0	38.5	-	-
II	-1813.2	33.4	6.2	4.9
III	-1816.0	33.8	9.0	4.7
I	-1889.9	25.3	82.9	21.2

Table S1: LOO-IC results for the sequential approach. Results are based on data from $n = 27$ mice. The models are ranked from best (top) to worst (bottom), using the "expected log predictive density" (elpd_loo) value. The p_loo value is a measure of the complexity of the model and generally increases with the number of parameters. The Δelpd value is the difference between the elpd_loo value and that of the best model. The se Δelpd is the standard error of the Δelpd.

Reagent or Resource	Source	Identifier
Antibodies		
Anti-Mouse Bcl2 Alexa Fluor 488	BD Biosciences	3F11; Cat. # 568426
Anti-Mouse CD103 BUV615	BD Biosciences	2E7; Cat. # 751631
Anti-Mouse CD11a PerCP-eFluor710	Thermofisher	M17/4; Cat. # 46-0111-82
Anti-Mouse CD19 BV480	BD Biosciences	1D3; Cat. # 566107
Anti-Mouse CD27 BV605	Biolegend	LG3A10; Cat. # 124249
Anti-Mouse CD3 BUV395	BD Biosciences	17A2; Cat. # 740268
Anti-Mouse CD4 APC-Cy/7	Biolegend	GK1.5; Cat. # 100412
Anti-Mouse CD44 BV786	Biolegend	IM7; Cat. # 103059
Anti-Mouse CD49a BUV737	BD Biosciences	Ha31/8; Cat. # 741776
Anti-Mouse CD62L BV570	Biolegend	MEL-14; Cat. # 104433
Anti-Mouse CD69 PE-Cy5	Biolegend	H1.2F3; Cat. # 104510
Anti-Mouse CD8 BUV805	BD Biosciences	53-6.7; Cat. # 612898
Anti-Mouse CX3CR1 APC-Fire 810	Biolegend	SA011F11; Cat. # 149053
Anti-Mouse CXCR5 BV650	Biolegend	L138D7; Cat. # 145517
Anti-Mouse CXCR6 BV711	Biolegend	SA051D1; Cat. # 151111
Anti-Mouse F4/80 BV480	BD Biosciences	T45-2342; Cat. # 565635
Anti-Mouse FoxP3 Pacific Blue	Biolegend	MF-14; Cat. # 126410
Anti-Mouse FR4 BUV563	BD Biosciences	12A5; Cat. # 748996
Anti-Mouse IA/IE (MHCII) BV480	BD Biosciences	M5/114.15.2; Cat. # 566088
Anti-Mouse Ki67 PE-Cy7	Biolegend	16A8; Cat. # 652426
Anti-Mouse KLRG1 PE-Fire 810	Biolegend	2F1/KLRG1; Cat. # 138437
Anti-Mouse PD-1 PE/Dazzle 594	Biolegend	RMP1-30; Cat. # 109115
Anti-Mouse Thy1.2 BV421	Biolegend	30-H12; Cat. # 105341
Chemicals and Reagents		
ACK Lysing buffer	Gibco	Ref. # A10492-01
Collagenase Millipore	Sigma	Cat. # 11088882001
DNAse Millipore	Sigma	Cat. # DN25-5G
DPBS	Corning	Cat. # 20-030-CV
EDTA	Corning	Cat. # 46-034-CI
Fetal Bovine Serum	GeminiBio	Cat. # 100-106
Fixable Viability Dye Zombie NIR	Biolegend	Cat. # 423106
FoxP3 Fix/Perm Concentrate	Cytek	Cat. # TNB-1020-L050
FoxP3 Fix/Perm Diluent	Cytek	Cat. # TNB-1022-L160
Flow Cytometry Perm Buffer	Cytek	Cat. # TNB-1213-L150
GolgiPlug	BD Biosciences	Cat # 555029
GolgiStop	BD Biosciences	Cat # 554724
IMDM	Gibco	Ref # 122440-053
Penicillin/Streptomycin/L-glutamine	GeminiBio	Cat# c400-110
Permeablization Buffer	Invitrogen	Cat # 00-8333-56
RPMI 1640	Corning	Cat #10-040-CM
Software and Algorithms		
FlowJo v 10.7 software	Tree Star	`https://www.flowjo.com`
Prism v 9.4.0 software	GraphPad	`www.graphpad.com`
CmdStan v 2.34 software	Stan	`www.mc-stan.org`
Pyro v 1.9.0 software	Pyro	`https://pyro.ai`

Table S2: Reagents and resources

Parameter	Description	Prior	Hyper-parameters
$\lambda_{E,i}$	Initial net loss rate of cluster i	$\lambda_{E,i} \sim \mathcal{N}(\mu_{\lambda_E}, \sigma_{\lambda_E})$	$\mu_{\lambda_E} \sim \mathcal{N}(0,1), \sigma_{\lambda_E} \sim \text{HalfNorm}(1)$
$\lambda_{L,i}$	Long-term net loss rate of cluster i	$\lambda_{L,i} \sim \mathcal{N}(\mu_{\lambda_L}, \sigma_{\lambda_L})$	$\mu_{\lambda_L} \sim \mathcal{N}(0,1), \sigma_{\lambda_L} \sim \text{HalfNorm}(1)$
u	Rate at which net loss rate λ changes from λ_E to λ_L	$u \sim \text{HalfNorm}(1)$	-
Q_{ij}	Rate of differentiation from cluster j to i	$Q_{ij} \sim \text{Exp}(\mu_Q^{-1})$	$\mu_Q \sim \text{Exp}(100)$
$X_{0,i}$	Initial size of cluster i	$X_{0,i} \sim \text{LogNorm}(0,10)$	-
σ_M	Scale parameter for likelihood of T cell numbers	$\sigma_M \sim \text{Exp}(1)$	-
τ_K	Dispersion parameter for likelihood of cluster size	$\tau_K^{-1} \sim \text{Exp}(10^3)$	-

Table S3: Prior distributions for the Bayesian model. Normal distributions are parameterized with scale parameters instead of variance. The half-normal distribution is denoted HalfNorm(σ), and the log-normal distribution LogNorm(μ, σ).