Early 2022 breakthrough infection sera from India target the conserved cryptic class 5 epitope to counteract immune escape by SARS-CoV-2 variants

Indrani Das Jana[1], Kawkab Kanjo[2], Subhanita Roy[1], Munmun Bhasin[2], Shatarupa Bhattacharya[5], Indranath Banerjee[3], Subhasis Jana[6], Arjun Chatterjee[7], Alok Chakraborty[8], Suman Chakraborty[4], Budhaditya Mukherjee[5], Raghavan Varadarajan[2*], Arindam Mondal[1*]

[1]Department of Bioscience and Biotechnology, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[2]Molecular Biophysics Unit (MBU), Indian Institute of Science, Bengaluru-560012, India
[3]B.C. Roy Technology Hospital, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[4]Department of Mechanical Engineering, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[5]School of Medical Science and Technology, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[6]Purba Medinipur District Hospital, Tamluk-721636, India
[7]Institute of Neuroscience, Kolkata-700017
[8]ICMR-National Institute of Cholera and Enteric Diseases, Kolkata-700010, India
[9]Lead contact: Arindam Mondal

Correspondence*: varadar@iisc.ac.in (R.V.), arindam.mondal@iitkgp.ac.in (A.M.)

Summary:

During the COVID-19 pandemic, the vast majority of epitope mapping studies have focused on sera from mRNA-vaccinated populations from high-income countries. In contrast, here we report an analysis of 164 serum samples isolated from breakthrough infection patients in India during early 2022 who received two doses of the ChAdOx viral vector vaccine. Sera were screened for neutralization breadth and potency against wildtype, Delta, and Omicron BA.1 viruses amongst which three were shortlisted for epitope mapping using charged scanning mutagenesis. These sera majorly targeted the recently identified class 5 cryptic epitope which is completely conserved across all SARS-CoV-2 variants and for different Sarbecoviruses. Consistent with these results, the sera show broad neutralizing activity against recent Omicron subvariants including XBB.1.5. Our study demonstrates that two vaccinations with the ChAdOx vaccine in a highly exposed population are sufficient to drive substantial neutralization breadth against emerging and future SARS-CoV-2 variants of concern.

Introduction

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has been circulating in the human population for over four years and rapidly evolving, causing the emergence of more than 50 different variants to date [1]. Multiple waves of infections were caused by the new variants despite massive vaccination programs, and their emergence was largely driven by the host immunity generated through natural infections and vaccination [2]. Detailed molecular-level understanding of the humoral immunity conferred by different vaccination regimens coupled with ongoing viral exposures is important both for the development of effective countermeasures against the current and future variants of SARS-CoV-2 and other Sarbecoviruses that may emerge in future.

More than 5.55 billion people, 72.3% of the world population, have received a single dose of COVID-19 vaccine [3], though in Africa, this number is less than 40% [4]. India ranks second highest after China in the number of COVID-19 vaccine doses administered (2.2 billion doses) [5]. The vaccination drive started in India on 16th January 2021, at the end of the first wave of infections caused mainly by the circulating B.1.1.7 (Alpha) and B.1.135 (Beta) variants. While the vaccination program was ongoing, the highly infectious B.1.617.2 (Delta) variant emerged and caused the second wave of the pandemic (March to October 2021), which resulted in the highest number of daily infections with high disease severity and mortality rate [6,7]. By January 2022, more than 77% of the Indian population were immunized; the new highly transmissible and heavily mutated B.1.1.529 (Omicron) variant emerged and led to the third wave of infection but was associated with milder symptoms and lower fatality rates [8]. A significant fraction of the vaccinated population in India got infected during the second and

third waves of the pandemic, leading to the large number of "breakthrough infections" [9]. These

breakthrough infections shaped the transmission dynamics and contributed to the overall

evolution of the virus [10]. After 2021, there was little vaccine uptake in the country and the

percentage of the population that received a third (booster) dose was only about 10%.

The SARS-CoV-2 spike protein (S) is the main surface glycoprotein that mediates the

interaction with the host angiotensin-converting enzyme 2 (ACE2) receptor and subsequent

fusion of the viral envelope with the host cell membrane that marks the initiation of the

infectious cycle [11,12]. The S protein is the primary target for neutralizing antibodies elicited

through natural infection as well as vaccination [13–15]. The spike constitutes a trimer where

individual protomers are comprised of two subunits, S1 and S2 [16]. S1 harbours the N-terminal

domain (NTD), the Receptor Binding Domain (RBD), and the S2 houses the fusion peptide,

the heptad repeats 1 and 2 (HR1, HR2), the transmembrane domain (TM) and the C-terminal

domain (CD). The majority of human neutralizing antibodies, isolated from convalescent or

vaccinated sera, target conformational epitopes in the RBD Receptor Binding Motif (RBM) or

in the conserved core of RBD[17–19] . In the context of the spike trimer, the individual RBDs can

be either in the up or down conformation, with only the former able to bind the ACE2 receptor.

The epitopes targeted by neutralizing antibodies can be classified into four major

classes (Class 1-4) [20]. Class 1 and class 2 epitopes overlap with the ACE2 binding site, and

antibodies targeting these sites competitively inhibit receptor binding. While class 1 epitopes

are accessible when RBD is present in the up conformation, class 2 epitopes are accessible

in both up and down conformations [21]. For certain class 2 epitope targeting antibodies (C144),

the long complementary determining region (CDR) of the heavy chain acts as a bridge

between adjacent RBDs from neighbouring protomers in the down conformation, thus locking

the S trimer into the prefusion closed conformation; an unusual mode of neutralization [21]. In

general, the antibodies targeting class 1 and class 2 epitopes have strong neutralizing activity.

However, different SARS-CoV-2 variants escape neutralization by these antibodies by

acquiring mutations in the RBM region that is part of the class 1 and 2 epitopes, compromising

their efficacy. Both class 3 and class 4 antibodies bind to the RBD core region away from the

ACE2 binding site. N343 glycan and the surrounding residues are accommodated in class 3

epitopes [22]. S309 antibody, a class 3 antibody isolated from a SARS-CoV infected individual,

exhibited broad cross-reactivity against SARS-CoV and SARS-CoV-2, although many SARS-

CoV-2 VOCs acquired mutations in class 3 epitopes to escape neutralization by these

antibodies [21]. The cryptic epitope around the RBD base (first described for CR3022 antibody)

constitutes the highly conserved class 4 epitope and is accessible only in the RBD up

conformation[23–26]. Despite their rare prevalence in convalescent sera, antibodies targeting

class 4 epitopes are suggested to protect against emerging VOCs and other sarbecoviruses.

Additionally, a highly conserved class 5 epitope has been recently identified and characterized

as a target for broadly neutralizing antibodies [27,28].

Mapping monoclonal antibody epitopes can inform about viral evolution and aid in the

development of epitope-based vaccines and therapeutics. However, mapping conformational

epitopes targeted by polyclonal sera remains challenging. Mapping these epitopes can

elucidate how escape mutations in the viral antigen impact the neutralization potency of the

human polyclonal sera harbouring a heterogeneous mixture of antibodies. In the present

study, we have conducted a comprehensive analysis of human sera collected from individuals

vaccinated with two doses of the ChAdOx1 nCoV-19 viral vectored vaccine and subsequently

had undergone breakthrough infections. Neutralization efficacies of 164 sera samples were

screened against SARS-CoV-2 pseudoviruses representing the parental strain WA.1 or VOCs

B.1.617.1, B.1.617.2 and B.1.1.529. Nine individuals whose serum samples exhibited broader

neutralization were subjected to longitudinal sampling and three of them were shortlisted for

epitope mapping. Utilizing charge scanning mutagenesis [29,30], a library of individual charged

substitutions at all the solvent-exposed positions of the wild-type RBD was generated and

displayed on the yeast surfaces to map the epitopes targeted by the polyclonal sera. A high

fraction of antibodies targeting either class 1 epitopes or the rare cryptic class 5 epitope, also

denoted as RBD8 [43], were observed in all three sera tested. The class 5/ RBD8 epitope

residues identified in this study are highly conserved across all SARS-CoV-2 variants,

consistent with the observed neutralizing potency of the selected serum samples against

heavily mutated Omicron BA.1, BA.5, and the recent XBB.1.5 variant. Our findings illustrate

how a combination of vaccinated and breakthrough infections in a highly exposed population

can elicit diverse antibody classes that confer potent neutralizing activity with broad cross-

reactivity, providing insights into why deaths and hospitalizations have remained low post-

2021 in virtually all low and middle-income countries (LMICs) even in the absence of variant

updated vaccine boosters.

Results

Study cohort

In contrast to vaccination alone, hybrid immunity resulting from SARS-CoV-2

breakthrough infections has been shown to boost neutralizing antibody titers while increasing

their breadth at the same time[31,32]. We shortlisted individuals who received two doses of the

ChAdOx1 nCoV-19 viral vector vaccine between April and November 2021 and then were

infected between November 2021 and January 2022, which marks the transition between the

second and third waves of the pandemic, in India caused by the B.1.617.2 (Delta) and

B.1.1.529 (Omicron) variants, respectively [33]. We hypothesized that such individuals would

have developed broadly cross-reactive neutralizing antibodies that are effective against

different variants of concern [31,32]. Blood samples, along with questionnaires about gender, age

and symptoms, were collected from 164 shortlisted individuals who were voluntarily recruited

by a separate consent for this study. All participants were confirmed positive by RT-PCR for

SARS-CoV-2 infection during the abovementioned time frame, which is within 2-4 months

post-second dose of vaccination (Figure 1A, B). The serum samples were collected from the

individuals within 4-6 weeks of their breakthrough infection. The study cohort consisted of 90

males and 74 females; the median age was 36 and 34 years, respectively (Figure 1D). Most

volunteers showed mild symptoms with 5.84% (10 individuals) population showing severe to

critical illness.

Neutralization profile of the polyclonal serum samples against SARS-CoV-2 variants.

We employed a high throughput neutralization assay based on reporter lentiviruses

pseudotyped with spike proteins from SARS-CoV-2 parental WA.1, B.1.617.1 (Kappa),

B.1.617.2 (Delta) and B.1.1.529 (Omicron) strains [34]. These pseudoviruses were incubated

with dilutions of serum samples ranging from 10^{-2} to 10^{-4} before infecting HEK-293T cells

overexpressing ACE2. The reporter signal was measured as a proxy of the infectivity, which

was the highest for the mock-treated cells and the lowest for a polyclonal antibody preparation

that binds the spike protein and has high neutralization activity (Novus Biologicals, NB100-

56578). The neutralization efficacies of 164 serum samples against WA.1 and different VOCs

was represented as a heatmap with colours ranging from red (high neutralization) to green

(low neutralization) (Figure 1C). Overall, the serum samples from this study exhibited

maximum neutralizing activity towards the Delta variant followed by the Kappa variant with a

geometric mean (GM) of 93% and 90%, respectively (Figure 1E). Reduced neutralizing activity

was observed against the parental WA.1 strain (GM-73%), which was further lowered against

the Omicron variant (GM-46%) (Figure 1E). These results suggest that majority of the

individuals in the study cohort had breakthrough infections caused by the Delta variant. Most

of the serum samples neutralized Delta and Kappa variants better than WA.1 (Figure 1C,

heatmap) due to the high similarity between the spikes of both variants [35]. Interestingly, certain

sera (BCRTH 29, 30, 33, 35, 45, 46) showed potent neutralization against WA.1 but failed to

neutralize other tested VOCs, especially at higher dilutions (Figure 1C). This suggests that

vaccination and subsequent infections have primed a strong immune response towards the

parental strain but failed to generate antibodies that are broadly cross reactive against the

VOCs, as also reported previously[36]. A few serum samples showed high neutralization activity

against the Omicron variant (Figure 1C). Limited neutralization activity against the heavily

mutated Omicron variant results from the high number of mutations conferring extensive

immune escape from vaccine and infection sera [37].

The breadth and cross-reactivity of the shortlisted polyclonal serum.

Serum samples with broadly cross-reactive neutralizing activity against WA.1, Kappa,

Delta and Omicron variants at 10^{-2} dilution were selected for further studies. Out of 164 serum

samples, 29 (17%) showed more than 65% neutralization against all four VOCs (Figure 1F &

Figure 1C, red diamonds) and hence, were considered as broadly neutralizing serum samples.

We selected 9 volunteers who have not been infected and had not received any vaccine

booster post the first round of blood collection and obtained their consent to participate in a

second round of blood sampling. The second set of serum samples (late) was collected from

the 9 selected individuals 4-6 weeks post the first set (early), and both early and late sets of

serum samples were subjected to further analysis.

Neutralization efficacies of the shortlisted serum samples were determined against

WA.1, Delta and the Omicron variants using the pseudovirus neutralization assay (Figure S1)

and neutralization titers were represented in the form of IC_{50} (Figure 2A, B, Early). Amongst

the nine serum samples collected at the early time point, four showed the highest IC_{50} values

for Delta, three for the parental WA.1 strain and two for the Omicron; a trend that matched

with the overall neutralizing activity of the larger cohort of 164 samples (Delta> WA.1>

Omicron; Figure 1E). Interestingly, this was not the case for the serum samples collected from

the same volunteers at the later time point (Figure S1 and Figure 2A, B Late). A moderate to

sharp decline in neutralizing activity (2-18 fold change in IC_{50}) was observed for most serum

samples between early and late time points (Figure 2 C, D). Interestingly, a few sera had an

increase or maintained similar variant-specific neutralization titers at both time points.

Compared to the early time point neutralization profile, BCRTH01 and INK41 showed a

significant decline in neutralization titer against WA.1 but a slight increase in neutralization

titer against the Omicron variant, at the late time point. This may reflect the higher durability

of the variant specific (Omicron or Delta) antibodies generated by the breakthrough infections

with respect to the parental strain (WA.1) specific antibodies elicited through vaccination.

Interestingly, BCRTH01, INK09, INK36 and INK41 sera maintained high neutralization titers

against all the pseudoviruses tested and at both time points, with INK36 being the most potent

neutralizing sera.

As the majority of the neutralizing antibodies target the receptor binding domain (RBD)

of the spike protein [17–19], we sought to test the binding activity of the four serum samples

(BCRTH01, INK09, INK36 and INK41) to the spike RBD. Enzyme-linked immunosorbent

assay (ELISA) was performed with recombinant purified spike RBD derived from the WA.1,

Omicron BA.1 and BA.5, which are used as coating antigens. As expected, all four serum

samples bound well to the RBD variants, indicating high abundance of RBD-targeted

antibodies in them (Figure S2 A). All sera samples showed highest IgG titres against WA.1

and marginally lower titres against BA.1 and BA.5 (2.2– 4.8 fold reduction) (Figure S2 B, C),

except for INK41, which showed 9.1 fold reduction for BA.1 compared to the WA.1. BA.5 RBD

exhibited higher sensitivity than BA.1, but lower sensitivity than WA.1 for binding to all four

serum samples tested. Three serum samples with high neutralizing activity against different

variants of SARS-CoV-2, namely BCRTH01, INK09 and INK36, were selected for further

studies.

Comprehensive mapping of RBD epitopes targeted by the shortlisted human serum.

We employed a charged scanning mutagenesis library of SARS-CoV-2 RBD displayed at the

yeast surface as a novel approach to identify the epitopes targeted by the antibodies present

in the polyclonal serum samples (Figure 3A). WA.1 RBD (332-532) was used as the template

for library construction and all RBD residues with solvent accessibility >15% were individually

mutated to negatively charged Aspartate, with the exception of negatively charged residues,

which were mutated to the positively charged Arginine. The generated RBD library was

displayed on the yeast surface and subjected to binding with a saturating concentration of the

polyclonal serum sample as determined by sera titration on the yeast surface (Figure S3 A).

Binding was detected using fluorescently labelled antibodies, and the yeast cell populations

were sorted into different bins through Fluorescence Activated Cell Sorting (FACS) (Figure S3

B-D). Yeast populations within these bins were subjected to deep sequencing to identify the

mutations responsible for abolishing binding to the serum. The deep sequencing reads were

analyzed as described in the methods section. Briefly, the binding and expression signals of

each mutant RBD were normalized with respect to the wildtype and the MFI ratio between the

normalized Mean Fluorescence Intensities for expression (MFI-expression) and binding (MFI-

binding) for each of the RBD variants was calculated.

Epitope mapping was performed with all three of the shortlisted serum samples

(BCRTH01, INK09 or INK36) and RBD residues whose mutation resulted in MFI_{Ratio} one

standard deviation higher than the mean MFI_{Ratio} were considered epitopes (Figure 3B, Table

S2). A high value of MFI_{Ratio} (darker shades of red) indicates that the charged substitution

significantly perturbed the binding of the serum samples with minimum or no effect on

expression. The MFI_{Ratio} of the identified epitope residues were mapped on the RBD structure

(Figure 3C). Overall, the identified epitopes (highlighted in lighter to dark shades of red) for all

three serum samples were clustered in two major groups; (i) epitopes that overlap with the

ACE2 binding motif (RBM) and (ii) epitopes that are away from the RBM. Interestingly, for

BCRTH01 serum, an enrichment in non-RBM over RBM epitope residues was observed.

However, this was not the case for INK36 or INK09, where epitope residues across different

RBD regions were enriched to similar levels.

To validate the results of the yeast-display screening and deep sequencing, we

selected three RBD residues identified as an epitope and another three residues identified as

non-epitope residues in the context of BCRTH01 serum and reintroduced these mutations to

the RBD individually followed by yeast surface display and BCRTH01 binding evaluation via

flow cytometry (Figure 3D). We selected the non-RBM epitope residues (T430, T437 and

R466) as they showed high values of MFI_{Ratio} compared to the wildtype. As shown in Figure

3D, charged substitution at the putative epitope residues significantly impacted the binding to

the serum while charged substitution at putative non-epitope residues had minimum or no

effect upon the binding (Figure S4 A, B & Figure 3D). This data validates the epitope residues

identified through the current study and establishes the robustness of the charged scanning

mutagenesis approach for the identification of epitope residues targeted by polyclonal human

serum.

Antibodies in the shortlisted sera target a cryptic epitope that is broadly conserved

across different SARS-CoV-2 variants.

Previous epitope mapping studies on convalescent polyclonal serum samples showed

enrichment of antibodies targeting the class-1, -2 and -3 epitopes and to a lesser extent the

class-4 epitope [37–41]. Hence, we analysed the different epitopes identified in the present study

and compared them to the epitopes reported in the literature. The major neutralizing epitopes

in the RBD are categorized into four major classes, class 1-4 (Figure 4 and Table S1). Out of

the 26 epitopes residues identified, eight were classified as Class 1, two as Class 2, two as

Class 3 and five as Class 4. Surprisingly, 11 out of the 26 (42%) epitope residues do not belong

to any of these major classes (except for K356 which is shared with class 3 and T470 which

is shared with class 2), but instead belong to a rare cryptic epitope [42,78], situated away from

the receptor binding motif of RBD (Figure 4A).

Residues constituting the cryptic epitope were recently designated as class 5 [27] or RBD8

class of epitope [43] (Figure S5). This rare cryptic epitope spans diagonally across the base of

the RBD (Figure 4A, S5) and is situated on the opposite surface of the ACE2 binding motif,

thereby remain buried and inaccessible in the prefusion closed conformation of the Spike

trimer (Figure 4B). Hence, binding of class 5 antibodies, such as BIOLS56, IMCAS74, WRAIR-

2063/ S2H97/ WRAIR-2057/ WRAIR-2134, to this rare cryptic epitope requires extensive

opening of the pre-fusion structure, thereby inducing rapid and premature transition to the

post-fusion conformation and S1 shedding [27,43–45]. These antibodies cross-reacted with all

SARS-CoV-2 VOCs, including Delta, Omicron BA.4/BA.5, BQ.1.1, and the recent highly

immune escaping variant XBB1[43,44]. Supporting this notion, class 5 epitope residues identified

in our analysis were highly conserved not only across all the different SARS-CoV-2 VOCs

(Figure 4C) but also across different Sarbecoviruses (Figure S6). Interestingly, the BCRTH01

serum had the highest number (total 10) of class 5 identified epitope residues compared to

the INK09 and INK36 (7 and 8 residues, respectively) (Figure 4A, Table S1), which explains

its breadth and high cross-reactivity against the different SARS-CoV-2 variants tested. In

contrast, INK36 has the highest number of class 1 and class 2 epitopes (8 and 2, respectively)

that are targeted by ACE2 competing antibodies, consistent with its highest neutralization

activity amongst the three shortlisted serum samples.

To evaluate the effect of charged substitutions on neutralizing activity of the polyclonal

sera, we selected four of the RBD-identified epitope residues and individually mutated them

to charged substitutions in the WA.1 backbone. This is followed by generation of

pseudoviruses harbouring these mutant spike proteins. Two mutations, R403D and G416D,

are situated within the RBM and are expected to perturb class 1 or class 1/ 4 epitope-specific

antibodies, respectively while the other two mutations, T430D and R466D, are situated away

from the RBM. T430D is expected to interfere with the binding of class 4 and class 5/ RBD8

epitope-specific antibodies, while R466D is expected to disturb the latter (Figure 4, Table S1).

We tested the sensitivity of WT and mutant pseudoviruses to neutralization by BCRTH01

serum. As expected (Figure 5 A, B), all the mutations reduced the neutralization activity of the

polyclonal serum in comparison to the WT pseudovirus, with R466D and R403D mutants

showing 132-fold and 48.9-fold reduction in the neutralization titer (IC_{50}), respectively. In

contrast, T430D and G416D mutations showed moderate and comparable effects with 3.55-

and 4.12-fold reduction in IC_{50} values, respectively. Together, our data suggests that the

polyclonal serum (BCRTH01) used in this study harbours the highest fraction of class 5/ RBD8

and class 1 antibodies, and a lower fraction of class 4 antibodies. This combination of ACE2

competing (class 1) and non-competing (class 5/ RBD8 and class 4) antibodies imparts high neutralization efficacy and broad cross reactivity to these serum samples, which render it effective against different variants of SARS-CoV-2, including the currently circulating ones.

Pre-XBB serum neutralizes the recent Omicron subvariant XBB.1.5.

The heavily mutated Omicron subvariant, XBB and XBB.1, emerged in late 2022 and soon, XBB and its descendants accounted for the majority of infections in the whole world [46,47]. The variant XBB.1.5 harbors a rare mutation, F486P, which demonstrates superior transmissibility and immune escape towards vaccines and monoclonal antibodies, accounting for its rapid spread in more than 100 counties [48–53]. Interestingly, the epitopes identified in this study remain completely conserved for XBB.1.5 (Figure 4C), suggesting that the shortlisted serum samples should neutralize this recent subvariant of SARS-CoV-2. Thus, we generated pseudoviruses harboring either WA.1, Omicron BA.1 or Omicron XBB.1.5 spike proteins at their surface and performed neutralization assays with the shortlisted serum samples (collected in early 2022, well before the emergence of XBB). As expected (Figure 5 C-F), all three serum samples showed neutralizing activities against Omicron BA.1 and XBB.1.5 comparable to the parental WA.1 strain, specifically for INK36 and INK09 serum. For BCRTH01, the neutralizing activity towards the XBB.1.5 is comparable to that of the Omicron BA.1 but 2.7 fold lower than for WA.1. This indicates that the shortlisted volunteers, who had breakthrough infections following vaccination during the Delta/ Omicron wave (December 2021) of the pandemic, generated humoral immunity that remained effective even against recently emerged subvariants (February-March 2023). This, in turn, establishes the potential importance of the identified epitopes as a target for developing vaccine candidates against future SARS-CoV-2 variants.

Discussion.

Breakthrough infections caused by waning immunity to SARS-CoV-2 VOCs have been

reported to induce enduring humoral immunity and generate broadly neutralizing antibodies

capable of combating different SARS-CoV-2 variants as well as SARS-CoV and other

Sarbecoviruses [32,54,55]. Human high-resolution epitope mapping studies following vaccination

and/or infection have been largely focused on sera from US or European populations that

were primarily immunized with mRNA vaccines [19,41,59,61] while a few studies focused on sera

from individuals vaccinated with CoronaVac whole virus inactivated vaccine with or without

breakthrough infection [52,63,65]. Little is known about the humoral response in LMICs and China,

which are home to ~85% of the world's population. In the present study, we describe epitope

mapping studies of sera from an Indian population of individuals who received two doses of

viral vectored vaccine followed by "breakthrough infections" within 4-6 weeks post-second

dose. The timing of the study was deliberately chosen at the intersection of the second and

third waves of the pandemic instigated by B.1.617.2 and B1.1.529, which, in theory, is

expected to enhance the breadth and potency of antibodies generated through vaccination

(parental strain WA.1) and subsequent infections by different VOCs[32]. Around 18% of the

serum samples exhibited > 65% neutralizing activity against the parental strain and different

VOCs including Kappa, Delta, and Omicron. Furthermore, via longitudinal sampling, three

serum samples were identified that not only maintained their broad neutralizing activity up to

12 weeks post-infection but also demonstrated effectiveness against the recent circulating

Omicron subvariant XBB.1.5. Although prior exposure to Omicron spike, including BA.1

bivalent vaccination was reported to generate XBB.1.5 specific antibodies [56], the neutralizing

activity remains low compared to the parental WA.1 or D614G strains [50,56–58]. Several studies

have demonstrated a substantial escape of XBB variants from neutralization by convalescent

and vaccinated sera, with more than 20-fold reduction in neutralization titres compared to

WA.1 [39,52,57,67]. Strikingly, the shortlisted serum samples used in this study showed less than

3-fold decrease in neutralizing titers against Omicron BA.1 and XBB.1.5 compared to the

parental strain, hence representing proof towards the role of hybrid immunity in generating

highly potent, durable, and broadly cross-reactive neutralizing antibodies.

We employed an aspartate scanning mutagenesis and yeast surface display coupled

with flow cytometry and deep sequencing to map epitope residues which, upon charged

substitution, severely impacted the binding of the serum antibodies without affecting the

expression. Charge scanning mutagenesis has been previously proven to be effective in

identifying residues buried at protein-protein interaction interface [60,62]. While Deep Mutational

Scanning using yeast surface display has been previously employed to identify serum escape

mutations on SARS-CoV-2 RBD [19,64], the relatively smaller size of the aspartate scanning

library, together with the unique barcoding strategy of the individual mutant ORFs employed

by us simplify library construction and downstream analysis. Using this method, we identified

a total of 26 surface-exposed amino acid residues whose mutation significantly impacted the

binding of the polyclonal sera and, hence, are designated as potential epitopes targeted by

the sera. The majority of these residues have been previously identified as part of epitopes for

different classes of RBD neutralizing monoclonal antibodies [66], thus supporting the robustness

of the novel screening approach in identifying epitopes targeted by polyclonal sera.

The epitopes identified using the proposed analysis can be broadly categorized into

two main groups: RBM epitopes overlapping with the ACE2 binding motif and non-RBM

epitopes. The RBM epitopes consist primarily of class 1 and class 2 epitopes which are

targeted by antibodies that competitively hinder receptor engagement and virus attachment to

host cells. While antibodies targeting these epitopes are responsible for the high neutralizing

activity of polyclonal serum used in this study [17,21,68–70], these regions are heavily mutated in

emerging VOCs, thereby limiting the breath and cross-reactivity of antibodies targeting these

epitopes [25,26,45,71]. For example, the R403K mutation, in the class 1 epitope, has been classified

as "vaccine escape mutation", leading to higher infectivity and lower neutralizing activity of

vaccine-elicited antibodies against the recent JN.1 subvariant of Omicron [72]. Interestingly, the

R403D mutation resulted in 48.9 fold reduction in neutralizing activity of the BCRTH01 serum

as evidenced in our study. Another identified class 2 epitope residue substitution L452R, was

associated with a massive expansion and immune escape of the Delta variant from vaccine-

elicited class1 and class 2 antibodies[41] during the second year of the pandemic [73]. Similarly,

the F456L mutation (class 1 epitope) is associated with heightened ACE2 binding, infectivity,

and immune evasion from the XBB.1.5 neutralizing antibodies [74] leading to the global spread

of EG.5 (Eris) subvariant [75] in the recent past. The K356T mutation in JN.1, in the class 3

epitope, drives resistance towards Sotrovimab, a class 3 antibody (S309) based antiviral drug

[76]. In contrast to classes 1-3, the class 4 epitopes targeted by the shortlisted sera described

above show high conservation (Figure 5B). These class 4 epitopes are situated in the RBD

core region, away from the RBM except for the G416, which resides within RBM. Class 4

antibodies targeting G416 (COVA1-16, C118 and C022) and surrounding residues sterically

hinder ACE2 binding [25,26] thereby resulted in high neutralizing activity.

A major fraction (11/26) of the epitope residues identified in the present study belong

to a rare cryptic epitope [42,77,78], rarely seen in the repertoire of antibodies from convalescent

sera [38–41]. Recently, this cryptic epitope has been categorized as class 5 or the RBD8 class of

epitopes [27,43]. These antibodies are reported to get enriched by breakthrough infections

caused by Omicron subvariants [52]. The amino acid residues constituting these epitopes are

clustered together at the opposite surface from the ACE2 binding motif of RBD and remain

inaccessible in the closed trimeric form of the spike (S) (Figure 4B, upper panel). Such

epitopes can only be accessible through breathing of the prefusion structure of the trimeric S

protein (Figure 4B, upper panel) whereby antibody binding ultimately leads to shedding of the

S1 subunit [43]. Additionally, it has been proposed that the binding of class 5 antibodies leads

to premature, receptor-independent conversion of S from pre to post-fusion conformation,

ultimately blocking viral entry and syncytium formation [79]. Interestingly, class 5 epitope

residues remain completely conserved across all SARS-CoV-2 variants and for different

Sarbecoviruses, indicating the cross-reactivity and increased breadth of class 5 antibodies

against these viruses. Consistent with this, the shortlisted serum samples, enriched for RBD8

epitope antibodies, effectively neutralized WA.1, Delta, Kappa, Omicron BA.1 and XBB.1.5.

Lately, antibodies targeting non-RBM epitopes in the spike protein have emerged as

prime candidates for developing broad-spectrum vaccines or therapeutics against SARS-CoV-

2 variants [28,80,81]. Some of these antibodies showed cross reactivity against different

Sarbecoviruses, thus further extending the prophylactic/therapeutic potential of such

antibodies against future emerging SARS-like coronaviruses. However, non-RBM antibodies

are limited by their weak neutralization activity. Rao et al., have shown that the RBD8 antibody

(BIOLS56) can rescue the immune escape caused by an RBM targeting antibody (IMCAS-

L4.65), together ensuring strong neutralizing activity and broad neutralizing efficacy against

different Omicron variants [43]. In this study, we show that an immune response elicited by hybrid

immunity and containing a combination of class-1 (RBM targeting) and class 5/ RBD8

antibodies can achieve neutralization potencies against different SARS-CoV-2 variants

including WA.1, Kappa, Delta and multiple Omicron subvariants including the recent XBB.1.5.

Furthermore, mutation of a single class5/ RBD8 epitope residue (R466D) resulted in 132 fold

reduction in the neutralization titer which is even higher than that observed in the case of

mutations of class 1 (R403D) epitope (48.9 fold), suggesting high abundance and comparable

neutralization efficacy of these two different classes of antibodies (class 5/ RBD8 and class 1)

in the shortlisted serum samples. This contrasts with the previously conducted epitope

mapping studies from the convalescent polyclonal serum samples, which led to the

identification of class 1-4 epitope targeting antibodies [38–41].

In conclusion, antibodies targeting different classes of epitopes may synergistically

contribute to achieving high neutralizing activity (Class 1+2 epitopes) and broader cross-

reactivity (Class 4 and RBD8) against all variants of SARS-CoV-2. Notably, RBD8 constitutes

a significant portion of the epitopes identified in all three serum samples, which differs from

the typical composition of the convalescent serum samples analysed previously [38–41]. In India,

after the Delta wave in mid-2021, there was little masking or isolation, and likely continued

extensive viral exposure. This, coupled with the relatively young population (median age 28.4

years) may have contributed to the generation of broadly cross reactive antibodies targeting

novel cryptic epitope presented here. These findings need to be confirmed in larger epitope

mapping studies. Understanding this adaptive strategy implemented by the host could pave

the way for the development of universal vaccines and therapeutics against future SARS-CoV-

2 variants and future sarbecovirus outbreaks.

Acknowledgement and Funding:

A.M. acknowledges the Indian Council for Medical Research (VIR/COVID-19/19/2021/ECD-I)

and the Science and Engineering Research Board (SERB-DST (CRG/2022/003628) for

providing financial support. R.V. is a JC Bose Fellow of DST. This work was funded by a grant

from the Bill and Melinda Gates Foundation (INV-042471) to RV. Funding for infrastructural

support was from DST FIST, MHRD, and the DBT IISc Partnership Program. I.D.J.

acknowledges the Science and Engineering Research Board (SERB) for the National Post

Doctoral Fellowship (PDF/2023/000284) awarded to her. K.K. acknowledges the support of

an MHRD-IISc doctoral fellowship. The authors acknowledge project staff Shyamal Manna for

his support in blood collection. The funders had no role in study design, data collection and

interpretation, or the decision to submit the work for publication.

Authors' Contributions:

Conceptualization—A.M., and R.V. Designing experiments—A.M., R.V., I.D.J. and K.K.

Performing experiments—I.D.J., K.K., and S.R. Manuscript writing—A.M., R.V., I.D.J. and

K.K. Fund acquisition—A.M., R.V., B.M., and S.C. Manuscript editing— A.M., R.V., I.D.J. and

K.K, B.M., S.C. and A.C. Deep Sequencing reads processing—M.B., Medical supervisors in

cohort identification and blood collection—S.J., A.C., and I.B.

Declaration of interests: The authors declare no competing financial interests or dispute in

the relationships that could have appeared to influence the work reported in this article.

Figure titles and legends

Figure 1. Study cohort and timeline. (A) Schematic representation of study design. (B)

Distribution of the study cohort as a function of the timeline of vaccination and breakthrough

infection in the context of the second and third wave of pandemics caused by different VOCs

of SARS-CoV-2. Red bars indicate the time points of serum sample collection for the entire

cohort (early) and subsequently for the shortlisted individuals (late). (C) Heatmap of

neutralizing activity of the 164 serum samples (at dilutions; 0.01, 0.001, 0.001) against the

parental strain and different VOCs of SARS-CoV-2 with green to red representing the lowest

to highest extent of neutralization (%). (D) Age distribution of male and female volunteers

included in the study cohort. (E) Comparison of the relative neuralization efficacies of the

serum samples (at 0.01 dilution) against parental strain and different VOCs. (F) Venn diagram

showing cross-reactivity of the shortlisted serum samples displaying more than 65%

neutralization (at 0.01 dilution) towards at least one of the pseudoviral variants tested in this

study. A total of 29 serum samples with >65% cross-reactivity towards all the VOCs tested in

this study were designated as broadly cross-reactive sera (red diamonds in 1C).

Figure 2. Neutralizing activity of the shortlisted serum samples collected at early and

late time points. (A) Comparative analysis of the neutralization titers (IC_{50}) of serum samples

isolated from the shortlisted 9 volunteers during 1st (early) and 2nd (late) batches of sample

collection, against WA.1, B.1.617.2 and BA.1. (B) Heatmap showing fold change in the IC_{50}

values for B.1.617.2 and BA.1 with respect to the parental WA.1 strain with red to white

indicating higher to lower neutralization titers. (C) Comparative analysis of the pseudovirus

neutralization titers of the serum samples isolated at early and late time points. (D) Heatmap

representing variant-specific fold change in IC_{50} values.

Figure 3. Epitope mapping through yeast display analysis. (A) Schematic representation

of the yeast display screening as described in the text. (B) The heat map represents the ratio

of MFI-expression to MFI-binding extracted from deep sequencing reads analysis. The higher

to lower ratios are represented in a red to blue gradient. The higher the ratio (*Red*), the more

sera escape at that position. The RBM is demarcated on the structure using a black line. (C)

The MFI-expression/ MFI-binding ratios are mapped on the RBD structure (PDB ID: 6M0J)

following the same color scheme. (D) The ratio of MFI-expression to MFI-binding of WT and

mutant RBDs w.r.t BCRTH01 serum, harboring mutations at the epitope and non-epitope

residues, are presented.

Figure 4. (A) Individual epitope residues, as identified for the BCRTH01, INK09 and INK36

serum samples, are mapped on the surface of the RBD (PDB ID: 6M0J) and are color-coded

based on their epitope class. (B) Class 5/ RBD8 epitopes identified in this study are mapped

on the pre-fusion trimeric structure of the spike in its closed (PDB ID: 6VXX) and open (PDB

ID: 6VYB) conformations. (C) Matrix representing the conservation of identified epitope

residues across all major SARS-CoV-2 variants. Mutations of a specific residue in a particular

variant are represented in red square. Epitope classes are designated using coloured

diamonds.

Figure 5. Mutation of epitope residues impacts the neutralizing activity of polyclonal

sera. (A) Neutralization curves for pseudoviruses with WT (WA.1) or mutant spike proteins,

harbouring specific mutations at the epitope residues with serial dilutions of the BCRTH-01

serum. (B) Neutralization titers (IC_{50}) for WT and mutant pseudoviruses. (C-E) Neutralization

curves for WA.1, BA.1 and XBB.1.5 pseudoviruses with serial dilutions of BCRTH-01, INK-09

and INK-36 sera, respectively. (F) IC_{50} fold changes with respect to the WA.1.

STAR Methods

Materials:

Cell line

Human embryonic kidney (HEK) 293T cells (ATCC #CRL-3216) were maintained in

Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum

(FBS) along with antibiotics like penicillin, streptomycin and anti mycotic agent (Gibco) at 37°C

and 5% CO_2.

Plasmids:

Plasmid pHAGE2-EF1aInt- ACE2-WT, expressing human ACE2 gene (BEI#NR-

52514) was used for overexpressing ACE2 in HEK 293T cells. Plasmid HDM-IDTSpike-fixK

(BEI#NR-52514) was used for expressing the codon-optimized spike protein from SARS-

CoV-2, Wuhan-Hu-1 strain (Genbank NC_045512). Plasmids pcDNA3.3-SARS2-B.1.617.1,

expressing spike protein of Kappa variant (addgene#172319), pcDNA3.3-SARS2-B.1.617.2

SARS-CoV-2, expressing spike protein of Delta variant (addgene#172320) [82] and SARS-

CoV-2 Omicron strain spike gene in Human codon_pcDNA3.1(+) expressing spike protein of

Omicron variant (B.1.1.529) (GenScript# MC_0101274) were utilized in the study [34]. XBB1.5

spike expression plasmid construct was a kind gift from Dr. Rajesh P. Ringe, Institute of

Microbial Technology, Council of Scientific and Industrial Research (CSIR), India.

Antibodies:

NB100-56578 is a polyclonal antibody raised in rabbits against SARS CoV-2 spike

protein and was obtained from Novus Biologicals (USA). Serum from the pre-COVID era

(Single donor human serum off the clot, ISERS2M-2019-33108-07) was obtained from

Innovative Research, Inc.

Methodology:

Bioinformatics and Structural Analysis:

Genome sequences of 47 different SARS-CoV-2 viruses that infect humans and 11

different sarbecovirus spike RBD sequences were obtained from the GISAID database.

Multiple sequence alignment was performed in Bioedit software using ClustalW multiple

alignment function. Further representation of the spike proteins' aligned RBD amino acid

sequences was performed using JALVIEW or the Multlin sequence alignment tool. Structural

alignment was performed in Pymol using specific PDB files of post-fusion spike protein

structures of SARS-CoV-2 viruses (PDB id: 6M0J).

Construction of SARS-CoV-2 S mutant plasmid constructs:

Plasmid HDM-IDTSpike-fixK expressing a codon-optimized spike protein from SARS-

CoV-2 was subjected to site-directed mutagenesis to generate R403D, G416D, T430D and

R466D, respectively, using the QuikChange II Site-directed mutagenesis kit (Agilent) according

to the manufacturer's instructions. Mutations generated in the SARS-CoV-2 S protein were

further verified using Sangers sequencing.

Generation of SARS-CoV-2 S pseudotyped lentiviruses:

Generation of SARS-CoV-2 lentivirus particles pseudotyped with either full-length WT,

different VoCs or mutation at SARS-CoV-2 spike protein (R403D, G416D, T430D and R466D)

were generated by transfecting HEK293T cells with the respective set of plasmids by using the

protocol as described previously by Crawford et al., 2020 [83]. Briefly, a lentiviral backbone

plasmid, pHAGE-CMV-Luc2-IRES-ZsGreen-W (BEI #NR-52516) bearing a CMV promoter to

express luciferase, was co-transfected with a set of other lentiviral plasmids like pHDM-

Hgpm2, expressing the HIV-1 gag and pol (BEI #NR-52517), pRC-CMV-rev1b, expressing the

HIV-1 rev (BEI #NR-52519), pHDM-tat1b, expressing the HIV-1 tat (BEI # NR- 52518), and

respective different WT or mutant spike protein-expressing plasmids. Post transfection, the

SARS CoV-2 pseudoviruses were collected by passing the supernatant through a 0.45 um

filter and used freshly.

Pseudovirus-based Serum Neutralization Assay:

Serum samples collected from the volunteers were inactivated at 65^0C for 30 minutes

prior to use. Serially diluted serum samples were incubated with either WT or different VoC/

mutant pseudoviruses at 37^0C for 1h. Serum-neutralized viral inoculum was then used to infect

hACE2-expressing HEK 293T cells at $37^{\circ}C$ for 1h in the CO_2 incubator in the presence of

polybrene till 18h. Successful neutralization of the pseudovirus results in a several fold

reduction in reporter virus activity, which were measured using a multimodal plate reader

(GloMax® Explorer Multimode Microplate Reader, Promega). Polyclonal SARS CoV-2

(NB100-56578) antibody served as a positive control, whereas sera from the pre-COVID era

were used as a negative control for the study.

Charged scanning library construction and yeast surface display:

Substitutions to Aspartate and Arginine are chosen for introduction in scanning

mutagenesis as analysis of existing DMS data suggests that these are the most disruptive for

binding. A charged scanning library of SARS CoV-2 spike RBD was generated using site-

directed mutagenesis. Exposed positions with accessibility >15% (total 125 residues) were

individually mutated to Aspartate or Arginine (for negatively charged residues), and each

mutant was fused by PCR to a unique 6 nucleotide barcode of known sequence to simplify

analysis and eliminate the need for long read sequencing. Upon mutagenesis by overlap PCR,

all the PCR products from each individual mutated position were pooled and transformed into

yeast via two fragment recombination by electroporation.

Human sera screening

Three human sera were selected for screening based on their high neutralization titres.

The sera were first titrated on yeast surface displaying WT-RBD and a saturating concentration

was chosen from the titration curve to be used for epitope mapping. The binding of these sera

with the aspartate library was analysed by flow cytometry, sorted into multiple gates and then

subjected to deep sequencing. Briefly, the pooled library of yeast cells expressing each of the

charged mutants was inoculated in 5 ml of SDCAA medium and grown overnight at 30°C, 250

rpm. The next day, a secondary inoculation from the overnight culture of the library was grown

at 30°C until an OD600 of 3–4, followed by induction in SGCAA medium at 20°C for 16 hr. A

total of 1× 10^7 cells of this library were washed twice with FACS buffer (1XPBS, pH 7.4 and

0.5% BSA) and then incubated at 4°C for 1 hr with a selected dilution of sera to probe for

binding and chicken anti-CMyc antibody (1:300 dilution) as a probe for the expression. The

cells were washed twice with FACS buffer and subsequently labelled with Alexa Fluor 633 goat

anti-human (1:1200 dilution) (catalogue no. A21091), plus Alexa Fluor 488 goat anti-chicken

(1:300 dilution), (catalogue no. A11039) for 20 min at room temperature in the dark. After

washing thrice with FACS buffer, cells were resuspended in 1 mL of FACS buffer, subjected to

flow cytometry, and sorted on an Aria-III cell sorter (BD Biosciences).

Deep sequencing and analysis

The sorted yeast cells from each gate were grown separately in liquid SDCAA medium

to saturation at 30°C for 24 hr, and the plasmids were extracted using EZ Yeast plasmid

miniprep kit (G Biosciences) according to the manufacturer's instructions and then subjected

to PCR. The PCR primers were designed to amplify the barcode region only (150 bp) and in a

way such that for both, the forward and reverse end reads, the first 3 bases are NNN followed

by 6 bases of unique sequence tag (MID) and the 21 bases of the primer sequence

complementary to the gene. Each forward MID sequence represents a particular serum, while

each reverse MID sequence represents a specific gate. The PCR for all the gates from three

sorted sera samples was carried out using Phusion polymerase for 15 cycles. Following

agarose gel electrophoresis and gel band purification, an equal amount (∼100 ng) of PCR

product from each sample was pooled and then gel-band purified, followed by sequencing on

the NovaSeq platform. The fraction of each mutant (*Xi*) distributed across all the bins was

calculated as in (Equation 1) and then the reconstructed MFI for each mutant was calculated

by the summation of the product obtained from multiplying the fraction of a particular mutant

in a particular gate (*Xi*) with the MFI of that gate that was obtained from the FACS experiment

(*Fi*), (Equation 2).

Fraction of mutant in each gate $\quad X_i = \dfrac{N_i}{\sum_1^n N_i}$ (Equation 1)

Ni: Number of reads of a specific mutant in bin i.

Reconstructed MFI= $\sum_1^n X_i * N_i$ (Equation 2).

The ratio of MFI expression/MFI binding for all positions was represented by a heat map using

GraphPad Prism software. The mutants with Ratio (Normalized MFI of expression: Normalized

MFI of binding), one standard deviation higher than the mean MFI were considered as epitope

residues.

Statistical analysis:

All experiments were performed in triplicates, and each data was repeated at least

three times except for the epitope mapping experiments, which were performed in

duplicates. Graphs are plotted using Graphpad Prism 10.1.2 and represented as mean

standard deviations (n = 3). Results were compared by performing a two-tailed Student's t-test.

Significance is defined as $P < 0.05$, and statistical significance is indicated with an asterisk (*).

*$P < 0.001$ were considered statistically significant.

Supplementary information titles and legends

Supplementary information.
Document S1. Figures S1-S6 and Table S1-S2

References:

1.   Ecdc.europa https://www.ecdc.europa.eu/en/covid-19/variants-concern.

2.   Zabidi, N.Z., Liew, H.L., Farouk, I.A., Puniyamurti, A., Yip, A.J., Wijesinghe, V.N., Low,
Z.Y., Tang, J.W., Chow, V.T.K., and Lal, S.K. (2023). Evolution of SARS-CoV-2
Variants: Implications on Immune Escape, Vaccination, Therapeutic and Diagnostic
Strategies, https://doi.org/10.3390/v15040944 https://doi.org/10.3390/v15040944.

3.   Our world in data https://ourworldindata.org/covid-vaccinations. covid 19 vaccinations
india.

4.   Doshi, R.H., Nsasiirwe, S., Dahlke, M., Atagbaza, A., Aluta, O.E., Tatsinkou, A.B.,
Dauda, E., Vilajeliu, A., Gurung, S., Tusiime, J., et al. (2024). COVID-19 Vaccination
Coverage - World Health Organization African Region, 2021-2023. MMWR. Morbidity
and mortality weekly report *73*, 307–311. https://doi.org/10.15585/mmwr.mm7314a3.

5.   Ministry of health and family welfare, G. of I. Ministry of Health and Family Welfare

672    https://www.mohfw.gov.in Govt. Of India: COVID-19 data.

673 6. Sarkar, A., Chakrabarti, A.K., and Dutta, S. (2021). Covid-19 Infection in India: A
674   Comparative Analysis of the Second Wave with the First Wave,
675   https://doi.org/10.3390/pathogens10091222
676   https://doi.org/10.3390/pathogens10091222.

677 7. Tareq, A.M., Emran, T. Bin, Dhama, K., Dhawan, M., and Tallei, T.E. (2021). Impact of
678   SARS-CoV-2 delta variant (B.1.617.2) in surging second wave of COVID-19 and
679   efficacy of vaccines in tackling the ongoing pandemic. Human Vaccines &
680   Immunotherapeutics *17*, 4126–4127.
681   https://doi.org/10.1080/21645515.2021.1963601.

682 8. Chavda, V.P., Balar, P., Vaghela, D., Solanki, H.K., Vaishnav, A., Hala, V., and Vora, L.
683   (2023). Omicron Variant of SARS-CoV-2: An Indian Perspective of Vaccination and
684   Management, https://doi.org/10.3390/vaccines11010160
685   https://doi.org/10.3390/vaccines11010160.

686 9. Krishna, B., Gupta, A., Meena, K., Gaba, A., Krishna, S., Jyoti, R., Aeron, N.,
687   Prashanth, S., Samriti, and Ganapathy, U. (2022). Prevalence, severity, and risk factor
688   of breakthrough infection after vaccination with either the Covaxin or the Covishield
689   among healthcare workers: A nationwide cross-sectional study. Journal of
690   anaesthesiology, clinical pharmacology *38*, S66–S78.
691   https://doi.org/10.4103/joacp.joacp_436_21.

692 10. Kavitha, C., Gowrisankar, A., and Banerjee, S. (2021). The second and third waves in
693   India: when will the pandemic be culminated? The European Physical Journal Plus
694   *136*, 596. https://doi.org/10.1140/epjp/s13360-021-01586-7.

695 11. Letko, M., Marzi, A., and Munster, V. (2020). Functional assessment of cell entry and
696   receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. Nature
697   microbiology *5*, 562–569. https://doi.org/10.1038/s41564-020-0688-y.

698 12. Jackson, C.B., Farzan, M., Chen, B., and Choe, H. (2022). Mechanisms of SARS-
699   CoV-2 entry into cells. Nature Reviews Molecular Cell Biology *23*, 3–20.
700   https://doi.org/10.1038/s41580-021-00418-x.

701 13. Xiaojie, S., Yu, L., lei, Y., Guang, Y., and Min, Q. (2021). Neutralizing antibodies
702   targeting SARS-CoV-2 spike protein. Stem Cell Research *50*, 102125.
703   https://doi.org/https://doi.org/10.1016/j.scr.2020.102125.

704 14. Xia, S., Liu, M., Wang, C., Xu, W., Lan, Q., Feng, S., Qi, F., Bao, L., Du, L., Liu, S., et
705   al. (2020). Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by a highly
706   potent pan-coronavirus fusion inhibitor targeting its spike protein that harbors a high
707   capacity to mediate membrane fusion. Cell Research *30*, 343–355.
708   https://doi.org/10.1038/s41422-020-0305-x.

709 15. Du, L., He, Y., Zhou, Y., Liu, S., Zheng, B.-J., and Jiang, S. (2009). The spike protein
710   of SARS-CoV — a target for vaccine and therapeutic development. Nature Reviews
711   Microbiology *7*, 226–236. https://doi.org/10.1038/nrmicro2090.

712 16. Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.L., Abiona, O., Graham,
713   B.S., and McLellan, J.S. (2020). Cryo-EM structure of the 2019-nCoV spike in the
714   prefusion conformation. Science *367*, 1260–1263.
715   https://doi.org/10.1126/science.aax0902.

716 17. Piccoli, L., Park, Y.-J., Tortorici, M.A., Czudnochowski, N., Walls, A.C., Beltramello, M.,
717   Silacci-Fregni, C., Pinto, D., Rosen, L.E., Bowen, J.E., et al. (2020). Mapping
718   Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding
719   Domain by Structure-Guided High-Resolution Serology. Cell *183*, 1024-1042.e21.
720   https://doi.org/10.1016/j.cell.2020.09.037.

721 18. Steffen, T.L., Stone, E.T., Hassert, M., Geerling, E., Grimberg, B.T., Espino, A.M.,
722   Pantoja, P., Climent, C., Hoft, D.F., George, S.L., et al. (2020). The receptor binding

domain of SARS-CoV-2 spike is the key target of neutralizing antibody in human
polyclonal sera. bioRxiv, 2020.08.21.261727.
https://doi.org/10.1101/2020.08.21.261727.

19. Greaney, A.J., Loes, A.N., Crawford, K.H.D., Starr, T.N., Malone, K.D., Chu, H.Y., and
Bloom, J.D. (2021). Comprehensive mapping of mutations in the SARS-CoV-2
receptor-binding domain that affect recognition by polyclonal human plasma
antibodies. Cell Host & Microbe *29*, 463-476.e6.
https://doi.org/https://doi.org/10.1016/j.chom.2021.02.003.

20. Chen, Y., Zhao, X., Zhou, H., Zhu, H., Jiang, S., and Wang, P. (2023). Broadly
neutralizing antibodies to SARS-CoV-2 and other human coronaviruses. Nature
Reviews Immunology *23*, 189–199. https://doi.org/10.1038/s41577-022-00784-3.

21. Barnes, C.O., Jette, C.A., Abernathy, M.E., Dam, K.-M.A., Esswein, S.R., Gristick,
H.B., Malyutin, A.G., Sharaf, N.G., Huey-Tubman, K.E., Lee, Y.E., et al. (2020).
SARS-CoV-2 neutralizing antibody structures inform therapeutic strategies. Nature
588, 682–687. https://doi.org/10.1038/s41586-020-2852-1.

22. Pinto, D., Park, Y.-J., Beltramello, M., Walls, A.C., Tortorici, M.A., Bianchi, S., Jaconi,
S., Culap, K., Zatta, F., De Marco, A., et al. (2020). Cross-neutralization of SARS-
CoV-2 by a human monoclonal SARS-CoV antibody. Nature *583*, 290–295.
https://doi.org/10.1038/s41586-020-2349-y.

23. Yuan, M., Wu, N.C., Zhu, X., Lee, C.-C.D., So, R.T.Y., Lv, H., Mok, C.K.P., and Wilson,
I.A. (2020). A highly conserved cryptic epitope in the receptor binding domains of
SARS-CoV-2 and SARS-CoV. Science *368*, 630–633.
https://doi.org/10.1126/science.abb7269.

24. Huang, Y., Yang, C., Xu, X. feng, Xu, W., and Liu, S. wen (2020). Structural and
functional properties of SARS-CoV-2 spike protein: potential antivirus drug
development for COVID-19. Acta Pharmacologica Sinica *41*, 1141–1149.
https://doi.org/10.1038/s41401-020-0485-4.

25. Liu, H., Wu, N.C., Yuan, M., Bangaru, S., Torres, J.L., Caniels, T.G., van Schooten, J.,
Zhu, X., Lee, C.-C.D., Brouwer, P.J.M., et al. (2020). Cross-Neutralization of a SARS-
CoV-2 Antibody to a Functionally Conserved Site Is Mediated by Avidity. Immunity *53*,
1272-1280.e5. https://doi.org/https://doi.org/10.1016/j.immuni.2020.10.023.

26. Jette, C.A., Cohen, A.A., Gnanapragasam, P.N.P., Mueecksch, F., Lee, Y.E., Huey-
Tubman, K.E., Schmidt, F., Hatziioannou, T., Bieniasz, P.D., Nussenzweig, M.C., et al.
(2021). Broad cross-reactivity across sarbecoviruses exhibited by a subset of COVID-
19 donor-derived neutralizing antibodies. Cell Reports *36*, 109760.
https://doi.org/https://doi.org/10.1016/j.celrep.2021.109760.

27. Jensen, J.L., Sankhala, R.S., Dussupt, V., Bai, H., Hajduczki, A., Lal, K.G., Chang,
W.C., Martinez, E.J., Peterson, C.E., Golub, E.S., et al. (2023). Targeting the Spike
Receptor Binding Domain Class V Cryptic Epitope by an Antibody with Pan-
Sarbecovirus Activity. Journal of virology *97*, e0159622.
https://doi.org/10.1128/jvi.01596-22.

28. Rouet, R., Henry, J.Y., Johansen, M.D., Sobti, M., Balachandran, H., Langley, D.B.,
Walker, G.J., Lenthall, H., Jackson, J., Ubiparipovic, S., et al. (2023). Broadly
neutralizing SARS-CoV-2 antibodies through epitope-based selection from
convalescent patients. Nature Communications *14*, 687.
https://doi.org/10.1038/s41467-023-36295-5.

29. Chandra, S., Manjunath, K., Asok, A., and Varadarajan, R. (2023). Mutational scan
inferred binding energetics and structure in intrinsically disordered protein CcdA.
Protein science : a publication of the Protein Society *32*, e4580.
https://doi.org/10.1002/pro.4580.

30. Chandra, S., Chattopadhyay, G., and Varadarajan, R. (2021). Rapid Identification of
Secondary Structure and Binding Site Residues in an Intrinsically Disordered Protein

Segment. Frontiers in genetics *12*, 755292.
https://doi.org/10.3389/fgene.2021.755292.

31.   Bates, T.A., McBride, S.K., Leier, H.C., Guzman, G., Lyski, Z.L., Schoen, D., Winders,
B., Lee, J.-Y., Lee, D.X., Messer, W.B., et al. (2022). Vaccination before or after
SARS-CoV-2 infection leads to robust humoral response and antibodies that
effectively neutralize variants. Science immunology *7*, eabn8014.
https://doi.org/10.1126/sciimmunol.abn8014.

32.   Walls, A.C., Sprouse, K.R., Bowen, J.E., Joshi, A., Franko, N., Navarro, M.J., Stewart,
C., Cameroni, E., McCallum, M., Goecker, E.A., et al. (2022). SARS-CoV-2
breakthrough infections elicit potent, broad, and durable neutralizing antibody
responses. Cell *185*, 872-880.e3.
https://doi.org/https://doi.org/10.1016/j.cell.2022.01.011.

33.   Swami, A., Mohanty, A., Jamwal, A., Turbadkar, D., Baveja, S., Shastri, J., and
Chitalia, V. (2023). A comparative analysis of the second and third wave of the Covid-
19 pandemic: an experience from a tertiary care hospital in Western India. Journal of
medical microbiology *72*. https://doi.org/10.1099/jmm.0.001685.

34.   Jana, I. Das, Bhattacharya, P., Mayilsamy, K., Banerjee, S., Bhattacharje, G., Das, S.,
Aditya, S., Ghosh, A., McGill, A.R., Srikrishnan, S., et al. (2022). Targeting an
evolutionarily conserved "e-L-L" motif in spike protein to identify a small molecule
fusion inhibitor against SARS-CoV-2. PNAS Nexus *1*, 1–14.
https://doi.org/10.1093/pnasnexus/pgac198.

35.   Wenlin, R., Xiaohui, J., Mingli, G., Jun, L., Yanying, Y., Quanxin, L., J., K.D., K., O.A.,
Yu, Z., Jin, Z., et al. (2022). Characterization of SARS-CoV-2 Variants B.1.617.1
(Kappa), B.1.617.2 (Delta), and B.1.618 by Cell Entry and Immune Evasion. mBio *13*,
e00099-22. https://doi.org/10.1128/mbio.00099-22.

36.   Servellita, V., Syed, A.M., Morris, M.K., Brazer, N., Saldhi, P., Garcia-Knight, M.,
Sreekumar, B., Khalid, M.M., Ciling, A., Chen, P.-Y., et al. (2022). Neutralizing
immunity in vaccine breakthrough infections from the SARS-CoV-2 Omicron and Delta
variants. Cell *185*, 1539-1548.e5.
https://doi.org/https://doi.org/10.1016/j.cell.2022.03.019.

37.   Ai, J., Zhang, H., Zhang, Y., Lin, K., Zhang, Y., Wu, J., Wan, Y., Huang, Y., Song, J.,
Fu, Z., et al. (2022). Omicron variant showed lower neutralizing sensitivity than other
SARS-CoV-2 variants to immune sera elicited by vaccines after boost. Emerging
Microbes & Infections *11*, 337–343. https://doi.org/10.1080/22221751.2021.2022440.

38.   Greaney, A.J., Starr, T.N., Barnes, C.O., Weisblum, Y., Schmidt, F., Caskey, M.,
Gaebler, C., Cho, A., Agudelo, M., Finkin, S., et al. (2021). Mapping mutations to the
SARS-CoV-2 RBD that escape binding by different classes of antibodies. Nature
Communications *12*, 4196. https://doi.org/10.1038/s41467-021-24435-8.

39.   Wang, Q., Iketani, S., Li, Z., Liu, L., Guo, Y., Huang, Y., Bowen, A.D., Liu, M., Wang,
M., Yu, J., et al. (2023). Alarming antibody evasion properties of rising SARS-CoV-2
BQ and XBB subvariants. Cell *186*, 279-286.e8.
https://doi.org/10.1016/j.cell.2022.12.018.

40.   Kugathasan, R., Sukhova, K., Moshe, M., Kellam, P., and Barclay, W. (2023). Deep
mutagenesis scanning using whole trimeric SARS-CoV-2 spike highlights the
importance of NTD-RBD interactions in determining spike phenotype. PLoS
pathogens *19*, e1011545. https://doi.org/10.1371/journal.ppat.1011545.

41.   Greaney, A.J., Eguia, R.T., Starr, T.N., Khan, K., Franko, N., Logue, J.K., Lord, S.M.,
Speake, C., Chu, H.Y., Sigal, A., et al. (2022). The SARS-CoV-2 Delta variant induces
an antibody response largely focused on class 1 and 2 antibody epitopes. PLOS
Pathogens *18*, e1010592.

42.   Tanaka, S., Olson, C.A., Barnes, C.O., Higashide, W., Gonzalez, M., Taft, J.,
Richardson, A., Martin-Fernandez, M., Bogunovic, D., Gnanapragasam, P.N.P., et al.

(2022). Rapid identification of neutralizing antibodies against SARS-CoV-2 variants by
mRNA display. Cell Reports *38*, 110348.
https://doi.org/https://doi.org/10.1016/j.celrep.2022.110348.

43. Rao, X., Zhao, R., Tong, Z., Guo, S., Peng, W., Liu, K., Li, S., Wu, L., Tong, J., Chai,
Y., et al. (2023). Defining a de novo non-RBM antibody as RBD-8 and its synergistic
rescue of immune-evaded antibodies to neutralize Omicron SARS-CoV-2.
Proceedings of the National Academy of Sciences *120*, e2314193120.
https://doi.org/10.1073/pnas.2314193120.

44. Sankhala, R.S., Dussupt, V., Chen, W.-H., Bai, H., Martinez, E.J., Jensen, J.L., Rees,
P.A., Hajduczki, A., Chang, W.C., Choe, M., et al. (2024). Antibody targeting of
conserved sites of vulnerability on the SARS-CoV-2 spike receptor-binding domain.
Structure (London, England : 1993) *32*, 131-147.e7.
https://doi.org/10.1016/j.str.2023.11.015.

45. Starr, T.N., Czudnochowski, N., Liu, Z., Zatta, F., Park, Y.-J., Addetia, A., Pinto, D.,
Beltramello, M., Hernandez, P., Greaney, A.J., et al. (2021). SARS-CoV-2 RBD
antibodies that maximize breadth and resistance to escape. Nature *597*, 97–102.
https://doi.org/10.1038/s41586-021-03807-6.

46. Hotez, P. (2023). XBB.1.5 emerges in the Americas: what it means to the region.
Lancet regional health. Americas *18*, 100433.
https://doi.org/10.1016/j.lana.2023.100433.

47. Callaway, E. (2023). Coronavirus variant XBB.1.5 rises in the United States - is it a
global threat?, https://doi.org/10.1038/d41586-023-00014-3
https://doi.org/10.1038/d41586-023-00014-3.

48. Ao, D., He, X., Hong, W., and Wei, X. (2023). The rapid rise of SARS-CoV-2 Omicron
subvariants with immune evasion properties: XBB.1.5 and BQ.1.1 subvariants.
MedComm *4*, e239. https://doi.org/10.1002/mco2.239.

49. Parums, D. V (2023). Editorial: The XBB.1.5 ('Kraken') Subvariant of Omicron SARS-
CoV-2 and its Rapid Global Spread., https://doi.org/10.12659/MSM.939580
https://doi.org/10.12659/MSM.939580.

50. Machado, R.R.G., Candido, É.D., Aguiar, A.S., Chalup, V.N., Sanches, P.R., Dorlass,
E.G., Amgarten, D.E., Pinho, J.R.R., Durigon, E.L., and Oliveira, D.B.L. (2024).
Immune Evasion of SARS-CoV-2 Omicron Subvariants XBB.1.5, XBB.1.16 and
EG.5.1 in a Cohort of Older Adults after ChAdOx1-S Vaccination and BA.4/5 Bivalent
Booster. Vaccines *12*. https://doi.org/10.3390/vaccines12020144.

51. Chen, J.-J., Li, L.-B., Peng, H.-H., Tian, S., Ji, B., Shi, C., Qian, C., Jiang, W.-G., Liu,
M.-C., Li, T.-T., et al. (2023). Neutralization against XBB.1 and XBB.1.5 after omicron
subvariants breakthrough infection or reinfection. The Lancet regional health.
Western Pacific *33*, 100759. https://doi.org/10.1016/j.lanwpc.2023.100759.

52. Cao, Y., Jian, F., Wang, J., Yu, Y., Song, W., Yisimayi, A., Wang, J., An, R., Chen, X.,
Zhang, N., et al. (2023). Imprinted SARS-CoV-2 humoral immunity induces
convergent Omicron RBD evolution. Nature *614*, 521–529.
https://doi.org/10.1038/s41586-022-05644-7.

53. Jian, F., Feng, L., Yang, S., Yu, Y., Wang, L., Song, W., Yisimayi, A., Chen, X., Xu, Y.,
Wang, P., et al. (2023). Convergent evolution of SARS-CoV-2 XBB lineages on
receptor-binding domain 455-456 synergistically enhances antibody evasion and
ACE2 binding. PLoS pathogens *19*, e1011868.
https://doi.org/10.1371/journal.ppat.1011868.

54. Kitchin, D., Richardson, S.I., van der Mescht, M.A., Motlou, T., Mzindle, N., Moyo-
Gwete, T., Makhado, Z., Ayres, F., Manamela, N.P., Spencer, H., et al. (2022).
Ad26.COV2.S breakthrough infections induce high titers of neutralizing antibodies
against Omicron and other SARS-CoV-2 variants of concern. Cell Reports Medicine *3*,
100535. https://doi.org/https://doi.org/10.1016/j.xcrm.2022.100535.

55. Nutalai, R., Zhou, D., Tuekprakhon, A., Ginn, H.M., Supasa, P., Liu, C., Huo, J.,
Mentzer, A.J., Duyvesteyn, H.M.E., Dijokaite-Guraliuc, A., et al. (2022). Potent cross-
reactive antibodies following Omicron breakthrough in vaccinees. Cell *185*, 2116-
2131.e18. https://doi.org/https://doi.org/10.1016/j.cell.2022.05.014.

56. Carr, E.J., Wu, M.Y., Gahir, J., Harvey, R., Townsley, H., Bailey, C., Fowler, A.S.,
Dowgier, G., Hobbs, A., Herman, L., et al. (2023). Neutralising immunity to omicron
sublineages BQ.1.1, XBB, and XBB.1.5 in healthy adults is boosted by bivalent BA.1-
containing mRNA vaccination and previous Omicron infection.,
https://doi.org/10.1016/S1473-3099(23)00289-X https://doi.org/10.1016/S1473-
3099(23)00289-X.

57. Nham, E., Kim, J., Lee, J., Park, H., Kim, J., Lee, S., Choi, J., Kim, K.T., Yoon, J.G.,
Hwang, S.Y., et al. (2023). Low Neutralizing Activities to the Omicron Subvariants
BN.1 and XBB.1.5 of Sera From the Individuals Vaccinated With a BA.4/5-Containing
Bivalent mRNA Vaccine. Immune network *23*, e43.
https://doi.org/10.4110/in.2023.23.e43.

58. Qian Wang , Yicheng Guo , Anthony Bowen , Ian A Mellis , Riccardo Valdez , Carmen
Gherasim , Abree Gordon , Lihong Liu, D.D.H. (2024). XBB.1.5 monovalent mRNA
vaccine booster elicits robust neutralizing antibodies against XBB subvariants and
JN.1. Cell host & microbe *32*, 315-321.e3.
https://doi.org/10.1016/j.chom.2024.01.014.

59. Greaney, A.J., Starr, T.N., Gilchuk, P., Zost, S.J., Binshtein, E., Loes, A.N., Hilton,
S.K., Huddleston, J., Eguia, R., Crawford, K.H.D., et al. (2021). Complete Mapping of
Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape Antibody
Recognition. Cell host & microbe *29*, 44-57.e9.
https://doi.org/10.1016/j.chom.2020.11.007.

60. Raksha, D., Rohini, D., and Raghavan, V. (2020). Probing the Structure of the HIV-1
Envelope Trimer Using Aspartate Scanning Mutagenesis. Journal of Virology *94*,
10.1128/jvi.01426-20. https://doi.org/10.1128/jvi.01426-20.

61. Greaney, A.J., Loes, A.N., Gentles, L.E., Crawford, K.H.D., Starr, T.N., Malone, K.D.,
Chu, H.Y., and Bloom, J.D. (2021). Antibodies elicited by mRNA-1273 vaccination
bind more broadly to the receptor binding domain than do those from SARS-CoV-2
infection. Science translational medicine *13*.
https://doi.org/10.1126/scitranslmed.abi9915.

62. Bajaj, K., Chakrabarti, P., and Varadarajan, R. (2005). Mutagenesis-based definitions
and probes of residue burial in proteins. Proceedings of the National Academy of
Sciences *102*, 16221–16226. https://doi.org/10.1073/pnas.0505089102.

63. Cao, Y., Yisimayi, A., Jian, F., Song, W., Xiao, T., Wang, L., Du, S., Wang, J., Li, Q.,
Chen, X., et al. (2022). BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by
Omicron infection. Nature *608*, 593–602. https://doi.org/10.1038/s41586-022-04980-y.

64. Rappazzo, C.G., Tse, L. V, Kaku, C.I., Wrapp, D., Sakharkar, M., Huang, D., Deveau,
L.M., Yockachonis, T.J., Herbert, A.S., Battles, M.B., et al. (2021). Broad and potent
activity against SARS-like viruses by an engineered human monoclonal antibody.
Science (New York, N.Y.) *371*, 823–829. https://doi.org/10.1126/science.abf4830.

65. Cao, Y., Wang, J., Jian, F., Xiao, T., Song, W., Yisimayi, A., Huang, W., Li, Q., Wang,
P., An, R., et al. (2022). Omicron escapes the majority of existing SARS-CoV-2
neutralizing antibodies. Nature *602*, 657–663. https://doi.org/10.1038/s41586-021-
04385-3.

66. Joseph, J., Darpe, S., Kulkarni, G., and Raj, V.S. (2023). SARS-CoV-2 neutralizing
antibody epitopes are overlapping and highly mutated which raises the chances of
escape variants and requires development of broadly reactive vaccines. Proteins *91*,
1021–1031. https://doi.org/10.1002/prot.26488.

67. Miller, J., Hachmann, N.P., Collier, A.-R.Y., Lasrado, N., Mazurek, C.R., Patio, R.C.,

Powers, O., Surve, N., Theiler, J., Korber, B., et al. (2023). Substantial Neutralization
Escape by SARS-CoV-2 Omicron Variants BQ.1.1 and XBB.1.,
https://doi.org/10.1056/NEJMc2214314 https://doi.org/10.1056/NEJMc2214314.

68. Dejnirattisai, W., Zhou, D., Ginn, H.M., Duyvesteyn, H.M.E., Supasa, P., Case, J.B.,
Zhao, Y., Walter, T.S., Mentzer, A.J., Liu, C., et al. (2021). The antigenic anatomy of
SARS-CoV-2 receptor binding domain. Cell *184*, 2183-2200.e22.
https://doi.org/https://doi.org/10.1016/j.cell.2021.02.032.

69. Huang, K.-Y.A., Tan, T.K., Chen, T.-H., Huang, C.-G., Harvey, R., Hussain, S., Chen,
C.-P., Harding, A., Gilbert-Jaramillo, J., Liu, X., et al. (2021). Correction: Breadth and
function of antibody response to acute SARS-CoV-2 infection in humans. PLOS
Pathogens *17*, e1010148.

70. Tortorici, M.A., Beltramello, M., Lempp, F.A., Pinto, D., Dang, H. V, Rosen, L.E.,
McCallum, M., Bowen, J., Minola, A., Jaconi, S., et al. (2020). Ultrapotent human
antibodies protect against SARS-CoV-2 challenge via multiple mechanisms. Science
370, 950–957. https://doi.org/10.1126/science.abe3354.

71. Tortorici, M.A., Czudnochowski, N., Starr, T.N., Marzi, R., Walls, A.C., Zatta, F.,
Bowen, J.E., Jaconi, S., Di Iulio, J., Wang, Z., et al. (2021). Broad sarbecovirus
neutralization by a human monoclonal antibody. Nature *597*, 103–108.
https://doi.org/10.1038/s41586-021-03817-4.

72. Wang, R., Chen, J., Gao, K., and Wei, G.-W. (2021). Vaccine-escape and fast-growing
mutations in the United Kingdom, the United States, Singapore, Spain, India, and
other COVID-19-devastated countries. Genomics *113*, 2158–2170.
https://doi.org/https://doi.org/10.1016/j.ygeno.2021.05.006.

73. Veronika, T., Hemantha, K., Lydia, L., Victoria, B., Elena, R., Dagmara, K., Yulia, S.,
Debarati, C., Iryna, M., Kai, D., et al. (2021). Acquisition of the L452R Mutation in the
ACE2-Binding Interface of Spike Protein Triggers Recent Massive Expansion of
SARS-CoV-2 Variants. Journal of Clinical Microbiology *59*, 10.1128/jcm.00921-21.
https://doi.org/10.1128/jcm.00921-21.

74. Kurhade, C., Zou, J., Xia, H., Liu, M., Chang, H.C., Ren, P., Xie, X., and Shi, P.
(2023). Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1 and XBB.1 by
parental mRNA vaccine or a BA.5 bivalent booster. Nature Medicine *29*, 344–347.
https://doi.org/10.1038/s41591-022-02162-x.

75. Parums, D. V (2023). Editorial: A Rapid Global Increase in COVID-19 is Due to the
Emergence of the EG.5 (Eris) Subvariant of Omicron SARS-CoV-2.,
https://doi.org/10.12659/MSM.942244 https://doi.org/10.12659/MSM.942244.

76. Vellas, C., Trémeaux, P., Del Bello, A., Latour, J., Jeanne, N., Ranger, N., Danet, C.,
Martin-Blondel, G., Delobel, P., Kamar, N., et al. (2022). Resistance mutations in
SARS-CoV-2 omicron variant in patients treated with sotrovimab. Clinical Microbiology
and Infection *28*, 1297–1299. https://doi.org/https://doi.org/10.1016/j.cmi.2022.05.002.

77. Xiong, H., Sun, H., Wang, S., Yuan, L., Liu, L., Zhu, Y., Zhang, J., Huang, Y., Qi, R.,
Jiang, Y., et al. (2022). The neutralizing breadth of antibodies targeting diverse
conserved epitopes between SARS-CoV and SARS-CoV-2. Proceedings of the
National Academy of Sciences *119*, e2204256119.
https://doi.org/10.1073/pnas.2204256119.

78. Li, T., Xue, W., Zheng, Q., Song, S., Yang, C., Xiong, H., Zhang, S., Hong, M., Zhang,
Y., Yu, H., et al. (2021). Cross-neutralizing antibodies bind a SARS-CoV-2 cryptic site
and resist circulating variants. Nature Communications *12*, 5652.
https://doi.org/10.1038/s41467-021-25997-3.

79. Starr, T.N., Greaney, A.J., Dingens, A.S., and Bloom, J.D. (2021). Complete map of
SARS-CoV-2 RBD mutations that escape the monoclonal antibody LY-CoV555 and its
cocktail with LY-CoV016. Cell Reports Medicine *2*, 100255.
https://doi.org/https://doi.org/10.1016/j.xcrm.2021.100255.

80. Hastie, K.M., Li, H., Bedinger, D., Schendel, S.L., Dennison, S.M., Li, K., Rayaprolu,
V., Yu, X., Mann, C., Zandonatti, M., et al. (2021). Defining variant-resistant epitopes
targeted by SARS-CoV-2 antibodies: A global consortium study. Science *374*, 472–
478. https://doi.org/10.1126/science.abh2315.

81. Martinez, D.R., Schäfer, A., Gobeil, S., Li, D., De la Cruz, G., Parks, R., Lu, X., Barr,
M., Stalls, V., Janowska, K., et al. (2024). A broadly cross-reactive antibody
neutralizes and protects against sarbecovirus challenge in mice. Science
Translational Medicine *14*, eabj7125. https://doi.org/10.1126/scitranslmed.abj7125.

82. Cho, H., Gonzales-Wartz, K.K., Huang, D., Yuan, M., Peterson, M., Liang, J., Beutler,
N., Torres, J.L., Cong, Y., Postnikova, E., et al. (2024). Bispecific antibodies targeting
distinct regions of the spike protein potently neutralize SARS-CoV-2 variants of
concern. Science Translational Medicine *13*, eabj5413.
https://doi.org/10.1126/scitranslmed.abj5413.

83. Crawford, K.H.D., Eguia, R., Dingens, A.S., Loes, A.N., Malone, K.D., Wolf, C.R., Chu,
H.Y., Tortorici, M.A., Veesler, D., Murphy, M., et al. (2020). Protocol and reagents for
pseudotyping lentiviral particles with SARS-CoV-2 spike protein for neutralization
assays. Viruses *12*. https://doi.org/10.3390/v12050513.

84. Li, T., Cai, H., Zhao, Y., Li, Y., Lai, Y., Yao, H., Liu, L.D., Sun, Z., van Vlissingen, M.F.,
Kuiken, T., et al. (2021). Uncovering a conserved vulnerability site in SARS-CoV-2 by
a human antibody. EMBO molecular medicine *13*, e14544.
https://doi.org/10.15252/emmm.202114544.

85. Bullen, G., Galson, J.D., Hall, G., Villar, P., Moreels, L., Ledsgaard, L., Mattiuzzo, G.,
Bentley, E.M., Masters, E.W., Tang, D., et al. (2021). Cross-Reactive SARS-CoV-2
Neutralizing Antibodies From Deep Mining of Early Patient Responses. Frontiers in
immunology *12*, 678570. https://doi.org/10.3389/fimmu.2021.678570.

86. Wu, N.C., Yuan, M., Bangaru, S., Huang, D., Zhu, X., Lee, C.-C.D., Turner, H.L.,
Peng, L., Yang, L., Burton, D.R., et al. (2020). A natural mutation between SARS-CoV-
2 and SARS-CoV determines neutralization by a cross-reactive antibody. PLOS
Pathogens *16*, e1009089.

87. Barnes, C.O., West, A.P.J., Huey-Tubman, K.E., Hoffmann, M.A.G., Sharaf, N.G.,
Hoffman, P.R., Koranda, N., Gristick, H.B., Gaebler, C., Muecksch, F., et al. (2020).
Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal Common
Epitopes and Recurrent Features of Antibodies. Cell *182*, 828-842.e16.
https://doi.org/10.1016/j.cell.2020.06.025.

1017

1018

1019

1020
1021

1022

1023

1024

1025

1026

1027

1028

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Early 2022 breakthrough infection sera from India target the conserved cryptic class 5 epitope to counteract immune escape by SARS-CoV-2 variants

Indrani Das Jana[1], Kawkab Kanjo[2], Subhanita Roy[1], Munmun Bhasin[2], Shatarupa Bhattacharya[5], Indranath Banerjee[3], Subhasis Jana[6], Arjun Chatterjee[7], Alok Chakraborty[8], Suman Chakraborty[4], Budhaditya Mukherjee[5], Raghavan Varadarajan[2*], Arindam Mondal[1*]

[1]Department of Bioscience and Biotechnology, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[2]Molecular Biophysics Unit (MBU), Indian Institute of Science, Bengaluru-560012, India
[3]B.C. Roy Technology Hospital, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[4]Department of Mechanical Engineering, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[5]School of Medical Science and Technology, Indian Institute of Technology Kharagpur, Kharagpur-721302, India
[6]Purba Medinipur District Hospital, *Tamluk-721636*, India
[7]Institute of Neuroscience, *Kolkata*-700017
[8]ICMR-National Institute of Cholera and Enteric Diseases, Kolkata-700010, India

Correspondence*: varadar@iisc.ac.in (R.V.), arindam.mondal@iitkgp.ac.in (A.M.)

39 **Supplementary information**

41 **Figure S1. Neutralizing activity of serum samples collected at early and late time points.**
42 Neutralization of the WA.1, B.1.617.2 and BA.1 pseudoviruses with nine serum samples collected
43 from the shortlisted volunteers during 1st and 2nd batches of sample collections, designated as (A)
44 early and (B) late time points.

45
Figure S2. The serum samples show high binding affinity towards spike RBD. (A) ELISA was
performed to determine the binding affinity of the polyclonal serum samples towards spike RBD derived
from WA.1 and Omicron subvariants BA.1 and BA.5. (B) Comparison between different serum samples
in terms of their binding affinity towards different RBD variants. (C) Showing fold change in IC50 values
for the BA.1 and BA.5 variants with respect to the WA.1.

Figure S3. (A) The binding profile of the shortlisted serum samples to wild type RBD displayed on the
yeast surface. (B, C and D) Histograms showing yeast cells binding to the indicated serum samples,

were sorted into eight gates based upon low to high binding affinity and the frequency of each
variant's barcode in individual gates was determined by Illumina sequencing.

56
57
58   **Figure S4**. Validation of epitope and non-epitope residues identified by deep sequencing analysis. (A)
59   The validation was carried out using FACS by quantifying the binding of WT and the selected RBD
60   mutants to BCRTH-01 serum (B) MFI-Expression and MFI-binding of WT and mutant RBDs,
61   harbouring mutations at the epitope and non-epitope residues are presented individually.
62

63
64
65   **Figure S5:** Surface representation of the RBD structure (PDB ID: 6M0J) with specific epitope residues
66   color coded based on their classification. Epitope residues of RBD8 (colored in yellow), class 5 (colored
67   in pale pink) and the rare cryptic epitope identified in this study (colored in orange) are individually
68   depicted with separate colors.

70
71 **Figure S6.** Multiple sequence alignment of different sarbecovirus spike RBD sequences. Black
72 triangles mark the Class5/ RBD epitope residues.
73
74 **Table S1. Classification of epitopes.**
75

Residue No.	Epitopes for serum samples			Epitope class	Antibody	Reference
	BCRTH -01	INK -09	INK -36			
351	+	+	+	Class 5/ RBD8	N-612-056; BIOLS56/ IMCAS74/ WRAIR-2057/ WRAIR-2134	[42–44,78]
355	+	+	+	Class 5/ RBD8	N-612-056/ BIOLS56/ IMCAS74/ WRAIR-2063/ S2H97/ FD20/WRAIR-2057/WRAIR-2134	[27,42–44,78,84,85]
356	+	-	-	Class3/ Class 5/ RBD8	S309/ N-612-056/ IMCAS74/ WRAIR-2134	[22,42–44]
359	-	-	+	Class3	S309	[22]
364	+	-	-	Unclassified	-	-
380	-	+	+	Class4	COVA1-16, CR3022	[25,86]
381	+	-	-	Class4	COVA1-16, CR3022	[25,86]
387	+	-	-	Unclassified		-
403	+	+	+	Class1	C002, C102, C121	[21]
416	+	+	+	Class1/ Class4	C105, COVA1-16, C118, C022,	[25,26,87]
420	+	-	+	Class1	C105, LY-CoV-016	[38,79,87]
421	+	+	+	Class1	C102	[21]
424	-	+	+	Class 5/ RBD8	BIOLS56/ IMCAS74/ WRAIR-2057/ WRAIR-2134	[43,44]
430	+	-	-	Class4/ Class 5/ RBD8	C022/ BIOLS56/ IMCAS74/ WRAIR-2057/ WRAIR-2134	[26,43,44]
437	+	+	+	Class 4	C118	[26]
452	-	-	+	Class2	C144, C002	[21]
456	+	+	+	Class1	C002, C102, C121	[21]
457	+	+	-	Class 5	N-612-056, 6D6; WRAIR-2057/WRAIR-2134	[42,44]

464	+	+	+	Class 5/ RBD8	N-612-056/ 6D6 & 7D6/ BIOLS56/ IMCAS74/ WRAIR-2063/ S2H97/ WRAIR-2057/ WRAIR-2134	27,42–44,78,84,85
465	+	-	+	Class 5/ RBD8	N-612-056/ 6D6 & 7D6/ BIOLS56/ IMCAS74/ WRAIR-2063/ S2H97/ WRAIR-2057/ WRAIR-2134	27,42–44,78,85
466	+	+	+	Class 5/ RBD8	N-612-056/ 6D6 & 7D6/ BIOLS56/ IMCAS74/ WRAIR-2063/ S2H97/ WRAIR-2057/ WRAIR-2134	42,43,78
467	+	+	+	Class 5/ RBD8	N-612-056; IMCAS74/ WRAIR-2063/ S2H97/ WRAIR-2057/ WRAIR-2134	42,43
470	+	-	+	Class2/ Class 5/ RBD8	C144/ 7D6, 6D6/ WRAIR-2057	21,78
473	+	+	+	Class1	C102	21
475	-	-	+	Class1	C102	21
489	-	+	+	Class1	C102	21

78
79 **Table S2. MFI ratios for the potential epitope residues**
80
81

SI No.	Residue	MFI-expression/ MFI-binding		
		BCRTH-01	INK-09	INK-36
1	351	1.295788	1.273718	1.157062
2	355	1.219729	1.239262	1.197349
3	356	1.17089	-	-
4	359	-	-	1.081459
5	364	1.182295	-	-
6	380	-	1.148814	1.077429
7	381	1.260786	-	-
8	387	1.230637	-	-
9	403	1.292267	1.172926	1.120017
10	416	1.329536	1.175896	1.101252
11	420	1.163761	-	1.072276
12	421	1.218064	1.168982	1.115263
13	424	-	1.192174	1.081695
14	430	1.270088	-	-
15	437	1.301646	1.107961	1.117455
16	452	-	-	1.076026
17	456	1.185845	1.252503	1.192579
18	457	1.225249	1.222689	-

19	464	1.246659	1.128143	1.087599
20	465	1.225858	-	1.080284
21	466	1.341557	1.187018	1.116536
22	467	1.242733	1.221766	1.118558
23	470	1.178698	-	1.063863
24	473	1.169721	1.145745	1.100281
25	475	-	-	1.102217
26	489	-	1.234014	1.144957

82
83
84
85