Membrane Binding Controls the ATPase Cycle and Localization of

MinD in *Bacillus subtilis*

Helge Feddersen[1], Charlotte Dyckmanns[1] and Marc Bramkamp[1,]*

[1] *Institute for General Microbiology, Christian-Albrechts-University Kiel, Am Botanischen*

Garten 1-9, 24118 Kiel, Germany

Running title: MinD dynamics in *B. subtilis*

*To whom correspondence should be addressed: Marc Bramkamp, Christian-Albrechts-
University Kiel, Institute for General Microbiology, Am Botanischen Garten 1-9, 24118 Kiel,
Germany, Email: bramkamp@ifam.uni-kiel.de; Phone: +49 (0)431-880-4341; Telefax:
+49(0)431-880-2198
Bluesky: @bramkamplab.bsky.social

Abstract

Bacteria precisely regulate the place and timing of their cell division. One of the best-understood systems for division site selection is the Min system in *Escherichia coli*. In *E. coli*, the Min system displays remarkable pole-to-pole oscillation, creating a time-averaged minimum at the cell's geometric center, which marks the future division site. Interestingly, the Gram-positive model species *Bacillus subtilis* also encodes homologous proteins: the cell division inhibitor MinC and the Walker-ATPase MinD. However, *B. subtilis* lacks the activating protein MinE, which is essential for Min dynamics in *E. coli*. We have shown before that the *B. subtilis* Min system is highly dynamic and quickly relocalizes to active sites of division. This raised questions about how Min protein dynamics are regulated on a molecular level in *B. subtilis*. Here, we show with a combination of *in vitro* experiments and *in vivo* single-molecule imaging that the ATPase activity of *B. subtilis* MinD is activated solely by membrane binding. Additionally, both monomeric and dimeric MinD bind to the membrane, and binding of ATP to MinD is a prerequisite for fast membrane detachment. Single-molecule localization microscopy data confirm membrane binding of monomeric MinD variants. However, only wild type MinD enriches at cell poles and sites of ongoing division, likely due to interaction with MinJ. Monomeric MinD variants and locked dimers remain distributed along the membrane and lack the characteristic pattern formation. Single-molecule tracking data further support that MinD has a freely diffusive population, which is increased in the monomeric variants and a membrane binding defective mutant. Thus, MinD dynamics in *B. subtilis* do not require any unknown protein component and can be fully explained by MinD's binding and unbinding kinetics with the membrane. The spatial organization of MinD relies on the short-lived temporal residence of MinD dimers at the membrane.

Keywords

B. subtilis, Min system, cell division, ATPase, membrane binding, protein patterns, reaction

diffusion, self-organization, SMLM, SMT

Introduction

Accurate selection of the division site is essential in bacteria for maintaining cell size, ensuring proper chromosome segregation, and preventing the formation of anucleate minicells. Cell division is governed by a complex protein machine known as the divisome (3-5). The spatiotemporal organization of this protein complex is directed by the dynamic arrangement of a bacterial tubulin homolog, FtsZ (6, 7). FtsZ forms treadmilling protofilaments that condense into a ring-like structure at the division site (8-10), recruiting several cell wall synthesis proteins, such as division-specific transglycosylases and transpeptidases (11, 12). Together, these proteins synthesize the bacterial peptidoglycan cell wall. To ensure that FtsZ polymerizes only at the correct location, many bacterial species employ the Min system, a conserved pathway that prevents aberrant polar septation events.

The Min system was identified through the formation of small, round, anucleate cells that divide close to the cell poles (27). The gene locus responsible for the "*min*iature" cell phenotype in *E. coli* was termed *minB* (28, 29), later shown to encode the three structural genes *minC*, *minD* and *minE* (30). MinC is the actual inhibitor of FtsZ-polymer formation (31, 32). The MinC protein contains an N- and C-terminal domain (33), both of which contribute to FtsZ inhibition. The C-terminal domain binds to the C-terminal tail of FtsZ, whereas the N-terminal domain, which also mediates MinC dimerization, binds at the interface of two FtsZ subunits (34-36). MinC-mediated inhibition of FtsZ has also been demonstrated in *B. subtilis* (37-39), suggesting a conserved mechanism across species.

The spatial localization of MinC is regulated by MinD, a Walker-type ATPase that belongs to the ParA/MinD family of proteins (37, 40, 41). These partitioning ATPases are involved in positioning cargo within bacterial cells (42, 43). While the molecular details of partitioning

differ among ParA-like proteins, a common feature is their use of binding to a large

intracellular surface as a spatial template (42, 43). ParA proteins typically bind non-

specifically to DNA, covering the nucleoid and exploiting it as a track for cargo positioning

(44). In contrast, MinD proteins bind to the membrane via an amphipathic helix (45, 46). In *E.*

coli, membrane binding of MinD strictly depends on ATP binding and subsequent

dimerization (40). Dimerized MinD recruits MinC and thereby activates MinC to inhibit FtsZ

oligomerization. In *E. coli*, MinE further regulates MinD localization by binding to the

membrane and interacting with MinD, stimulating ATP hydrolysis and triggering MinD

release from the membrane. Slow nucleotide exchange in cytosolic MinD leads to an

oscillatory cycle of membrane attachment and detachment, resulting in pole-to-pole

oscillation of MinCD (48-50). This oscillatory behavior generates a temporal minimum of

MinC at midcell, such that Z-ring assembly is restricted exclusively to this position. The

oscillation can be reconstituted *in vitro* on planar membranes, demonstrating the robustness

of this self-organizing system and a prime example of pattern formation. (51).

The Gram-positive model bacterium *B. subtilis* likewise encodes the *minC* and *minD* genes

(52, 53), and loss of either results in the characteristic minicell phenotype. Thus, a role for

the Min system analogous to that in *E. coli* has been proposed (37, 54). However, *B. subtilis*

lacks a MinE homolog, and the Min proteins do not oscillate from pole to pole (37, 54).

Instead, static imaging shows that MinD accumulates at the cell poles and at active division

sites. Initial genetic and cell biological studies demonstrated that the polar scaffold protein

DivIVA is essential for polar MinD localization (37). The interaction between DivIVA and

MinD is mediated by the integral membrane protein MinJ (55, 56). MinJ is also part of the

divisome and interacts with multiple division proteins (56). Loss of MinJ causes defects in

divisome disassembly, as proteins such as FtsL and PBP2b remain associated with the

nascent cell pole after division in Δ*minJ* cells (57). This function is supported by recent

findings that MinD promotes FtsZ polymer disassembly at the cell poles, and that deletion of

minD inhibits polar peptidoglycan remodeling (58).

We have recently shown that all components of the *B. subtilis* Min system, including DivIVA

and MinJ, dynamically localize within the bacterial cell (57, 59-61). Interestingly, even the

scaffold protein DivIVA, although highly conserved in many Gram-positive phyla, is not

statically anchored at the poles but instead redistributes along the membrane from poles to

septa (59-61). In contrast, DivIVA homologs in actinobacteria show greatly reduced mobility,

consistent with their role in apical growth (60).

Despite these advances, the mechanism by which MinD activity and localization are

regulated in the absence of MinE has remained unclear, raising the possibility of additional,

unidentified regulatory factors or alternative modes of ATPase control. In this study, we

address these long-standing questions by combining *in vitro* biochemical analyses with

single-molecule imaging *in vivo*. We define the ATPase and membrane-binding properties of

B. subtilis MinD and examine how its dynamics are shaped by interactions with MinJ and

MinC. Our findings reveal an unexpected mechanism of MinD regulation that operates

independently of MinE-like factors, providing a revised understanding of how the non-

oscillatory *B. subtilis* Min system robustly controls cytokinesis and prevents aberrant

septation events.

Results

Purification and gel filtration of His-MinD

Since localization and function of MinD appear to be strongly tied to its ATPase activity, the first logical step in describing and understanding details of the ATPase cycle was identification of the relevant biochemical characteristics, to eventually be able to correlate these properties to *in vivo* observations. To this end, *B. subtilis minD* was heterologously expressed in *E. coli*, yielding an N-terminally His-tagged fusion protein (Fig. S 1), as C-terminal MinD fusions have been shown to interfere with membrane binding (62). Since MinD is known for being membrane-associated, a well-established *E. coli* MinD purification protocol was adapted (63), where buffers were supplied with excess Mg^{2+}-ADP to potentially reduce MinD self- and membrane interaction. Based on the respective western blot and gel filtration chromatogram (Fig. S 1 and Fig. S 2), His-MinD is not able to dimerise when purified in presence of excess Mg^{2+}-ADP, as evidenced by a main peak at the expected sizes of the monomeric form (~31 kDa, 16.11 ml) and the absence of a peak at the expected size of the dimeric form (~62 kDa, 14.65 ml, Fig. S 2). However, Western blotting of Ni-NTA and size-exclusion chromatography (SEC) fractions indicates the presence of multiple His-tagged degradation products alongside full-length His-MinD (Fig. S 1). Because these fragments retain the His-tag, they co-purify during Ni-NTA and can co-elute or form non-covalent assemblies with full-length MinD, increasing the heterogeneity of the sample and likely producing the minor shoulder observed at an apparent molecular weight of ~44 kDa (15.4 ml, Fig. S 2). Unfortunately, the Mg^{2+}-ADP excess in the purification buffers did not seem to stop His-MinD membrane interaction, as a significant fraction of the recombinant protein

were later detected to be lost in the cell pellet. However, sufficiently high His-MinD

concentrations could be obtained to proceed with biochemical analysis.

B. subtilis MinD displays full ATPase activity when incubated with ATP and liposomes.

Being able to purify His-MinD allowed us to investigate the biochemical conditions affecting

the MinD ATPase cycle next. For this purpose, we choose to use the commercially available

enzyme coupled EnzChek™ phosphate assay kit (Thermo Fisher Scientific) to measure

ATPase activity. The activity was determined by measuring the P_i production rate (in μM

min^{-1}) as a function of His-MinD concentration. We first tested His-MinD ATPase activity at

protein concentrations between 0.5 μM and 16 μM. Under these conditions, only baseline

ATPase activity below the detection limit was observed (consistent with Fig. 1 D, 0 mg ml^{-1}

liposomes condition). Since MinD interacts with the membrane and it was expected that the

ATPase cycle is involved in controlling this interaction, we next measured ATPase activity in

the presence of liposomes extruded from _E. coli_ total lipids (Avanti). Remarkably, this led to

a robust increase in ATPase activity of His-MinD, appearing to be linearly dependent on

protein concentration (Fig. 1 A & B). To be able to compare the activity level to the published

E. coli MinD _in vitro_ ATPase data (64-66), the specific activity is shown as a specific activity in

nmol mg^{-1} min^{-1} (Fig. 1 B). Interestingly, activity levels of recombinant _B. subtilis_ MinD were

found to be on par with fully active _E. coli_ MinD in conjunction with MinE and phospholipids

(64). Moreover, the ATPase activity did not show cooperative behavior compared to _E. coli_

MinD at lower concentrations (64). We further measured that ATP hydrolysis can be titrated

with both ATP and liposomes and is described by Michaelis-Menten kinetics (Fig. 1 C & D).

Again, the hereby-obtained V_{max} values (19.50 nmol mg^{-1} min^{-1} and 22.05 nmol mg^{-1} min^{-1},

respectively) indicate similar ATPase activity levels between MinD from _E. coli_ and _B. subtilis_.

158   MinC and the PDZ domain of MinJ have different effects on MinD ATPase activity

Even though ATPase activity levels were similar, proteins that directly interact with MinD

could potentially modulate this activity *in vivo* as seen for MinD:MinE in *E. coli* (64-66). To

get the complete picture of MinD activity, investigating the potential effects of both known

B. subtilis MinD interactors MinC and MinJ was therefore necessary. For this purpose, both

proteins were subjected to purification attempts. While purification of the cytosolic His-

MinC could be successfully established using affinity chromatography followed by SEC (Fig. S

3), the intermembrane protein His-MinJ proved more challenging. Since several different

purification attempts of the full-length protein yielded no stable product, a truncated form

containing only the soluble C-terminal PDZ domain of MinJ fused to an N-terminal His-tag

(His-PDZ) via a flexible linker was expressed instead (Fig. S 4). PDZ domains are often found

in membrane-associated proteins and known to selectively interact with the C-terminus of

partner peptides or proteins to sequester them to highly specific sites of the cell, both in

pro- and eukaryotes (67). Although His-PDZ could be isolated via Ni-NTA affinity

chromatography, it was unstable during SEC and consistently lost from the column,

precluding further purification by this method. Hence, further experiments were performed

using His-PDZ directly after affinity chromatography.

To test the effect of His-MinC and His-PDZ on His-MinD ATPase activity, the previously

established enzymatic assays were repeated in the presence of either His-MinC or His-PDZ

(Fig. S 5). Addition of His-MinC resulted in a very small but statistically significant decrease in

MinD ATPase activity (~5%). Given the small sample size this effect should be interpreted

cautiously. However, it could be caused by the experimental set-up, since the high amounts

of MinC (6 µM) that have been used might sequester a small fraction of MinD away from

membrane surfaces that are required for efficient ATP hydrolysis, due to direct interaction of

MinC and MinD.

In contrast, addition of His-PDZ led to a slight but reproducible increase in MinD ATPase

activity, which prompted us to repeat the experiment at different His-PDZ concentrations

(Fig. S 5). All His-PDZ concentrations significantly increased His-MinD ATPase activity

compared to the control (1 µM His-PDZ = 25% ± 6.8%, 3 µM His-PDZ = 37 ± 12.5 %; 5 µM His-

PDZ = 22% ± 2.8 %). We speculate that the PDZ domain may promote MinD:MinD

encounters, facilitating dimerization rather than directly enhancing catalytic turnover.

Because only the soluble PDZ domain could be purified, it remains unclear how full-length,

membrane-anchored MinJ would influence MinD activity in vivo. The magnitude of the

effect might differ when the PDZ domain is present in its native membrane-associated

context.

MinD ATPase activity is triggered by membrane binding

To determine whether interaction with the membrane is sufficient to stimulate MinD ATPase

activity, a MinD mutant that is defective in membrane binding was required. MinD interacts

with the membrane through an amphipathic α-helix located at the N-terminus, thus being

the membrane targeting sequence (MTS) (Fig. 2 A). Exchanging the hydrophobic isoleucine

residue in the apolar core of the MTS with a charged glutamate (I260E) has been shown to

abolish membrane binding of MinD by perturbing the amphipathic properties, demonstrated

through fluorescence microscopy (68). Accordingly, a His-MinD-I260E variant was

constructed and purified via the established purification protocol. The yield of the

purification was extensively higher compared to the wild type protein (Fig. S 6), hinting

towards a successful abolishment of membrane interaction.

We next repeated ATPase activity assays with the purified, recombinant I260E mutant, and

remarkably, activity remained below the detection limit of the assay, while a positive control

of His-MinD wild type protein displayed regular activity (Fig. S 7). In summary, these data

collectively suggest that membrane binding is necessary and the key trigger for ATP

hydrolysis of *B. subtilis* MinD, without the apparent need for another stimulus.

Next, we wanted to compare *in vivo* and *in vitro* data. Therefore, a *B. subtilis* strain for

microscopy was constructed, where the native copy of MinD was replaced by a HaloTag-

MinD fusion protein (Fig. 2 B, HaloTag will henceforth be referred to as Halo). We then

introduced the I260E mutation, resulting in a strain producing Halo-MinD-I260E as the only

copy. Microscopic analysis of a Halo-MinD-I260E with TMR dye confirmed drastically

reduced membrane binding through its almost exclusive cytosolic localization (Fig. 2 B).

Furthermore, cell length and the number of minicells increased drastically, comparable to a

minD knockout (Tab. S 1).

Construction of catalytic MinD mutants

To determine the role of ATP-mediated dimerization in membrane binding, we explored

further mutations that affect the ATP binding region and thus the dimerization process. Due

to the conserved nature of Walker-type MinD/ParA-like ATPases, residues involved in

catalytic activity have been well characterized (69-71). We therefore constructed expression

plasmids encoding three different MinD variants (illustrated in Fig. 2 A): This first mutation,

MinD-G12V, produces a steric clash within the active site, so that ATP can still be bound, but

dimers cannot form (69, 70). The second mutation, K16A, completely disrupts ATP binding

and hence results in monomeric protein (69). The third variant encodes the mutation D40A,

which should interfere with ATP hydrolysis (70, 71). Therefore, it was expected that MinD-

D40A forms a trapped nucleotide sandwich dimer similar to Soj or *E. coli* MinD (70, 71). All

MinD variants were successfully expressed and purified and resulted in similar protein yields

when compared to the wild type protein (Fig. S 1, Fig. S 8, Fig. S 9 and Fig. S 10). When

evaluating the gel filtration chromatograms, His-MinD-G12V elute in a single peak, which can

be interpreted as a monomer when compared to the wild type His-MinD elution peak (Fig. S

2). His-MinD-D40A on the other hand also eluted in a single peak, albeit being wider and

eluting earlier, indicating a mainly dimeric conformation. It eluted between the expected

dimer and monomer positions during SEC as a wide peak (Fig. S 2). As His-MinD-D40A cannot

hydrolyze ATP, it should be dimeric and likely also form small oligomers or transient

multimers. A mixture of monomer/dimer/oligomer likely produced the broad, left-shifted

peak whose center is between monomer and dimer. Only His-MinD-K16A reproducibly

formed a rather unexpected shoulder in its main elution peak, which could hint towards

further interaction with the gel filtration matrix or other potential self-interactions. All of the

recombinant His-MinD mutants displayed negligible ATPase activity when tested using the

EnzChek phosphate assay kit (Fig. S 7 C). This was expected, as all of these MinD variants

(G12V, K16A and D40A) were mutated in residues which significantly alter the ATP binding

site. In all cases, activity levels remained consistently near the detection limit, precluding

reliable quantitative interpretation.

To investigate *in vivo* effects of these variants, the same mutations were introduced into the

strain expressing Halo-MinD in *B. subtilis*, respectively, and subsequently imaged via wide-

field microscopy (Fig. 2 B). All strains producing MinD mutants displayed an increased cell

length and minicell production comparable to Δ*minD* (Tab. S 1), as it was expected in cells

containing dysfunctional Min proteins. Surprisingly, both monomeric MinD variants (G12V

and K16A) were still observed to be membrane associated, however resulting in the

apparent loss of the polar-septal MinD localization (Fig. 2 B). Finally, the locked dimer D40A

mutant appeared to be mainly membrane associated, thereby forming large foci indicating

accumulation of MinD, but much less dispersed when compared to G12V or K16A (Fig. 2 B).

In summary, the time-resolved MinD gradient appears to be lost in all MinD mutants,

including D40A. This suggests that the ATPase cycle and thus quick cycling of MinD is indeed

necessary for formation of the observed dynamic gradient, as it has been previously

suggested (61).

Monomeric MinD binds the membrane with high efficiency.

Membrane binding appeared to be a key factor of the ATPase cycle and since the

microscopy-data indicated binding of MinD monomers to the membrane, we next aimed to

determine biochemically, if the respective MinD variants are capable of binding to

membrane and if so, to characterize their respective binding kinetics. To this end, we

employed Bio-layer interferometry (BLI), a technique capable of quantifying binding strength

and protein interaction, as well as identifying reaction kinetics. Liposomes were immobilized

on a biosensor to create a membrane-like surface for subsequent MinD binding assays. After

successful saturation of the sensor with liposomes, MinD-liposome association and

dissociation was measured, a scheme of the experimental set-up is depicted in Fig. 3 A. As a

negative control for unspecific binding, MinD-sensor interaction was tested on empty SA

sensors. Furthermore, unspecific binding to liposome-saturated sensors was tested using

bovine serum albumin (BSA). Both resulted in negligible, diffuse signal (data not shown).

Kinetics of His-MinD binding was tested first, using different protein concentrations between

1 µM and 16 µM in presence of ATP. Binding to the sensor-bound liposomes scaled with

increasing protein concentration, resulting in an equilibrium dissociation constant $K_D = 7.17$

μM and an association rate constant k_{on} = 3.03 x 10^3 $M^{-1}s^{-1}$ as a reference point for the wild

type protein (Fig. 3 B, Tab. 1**Error! Reference source not found.**). Unsurprisingly, His-MinD-

I260E showed a more than 10-fold reduced affinity to the membrane with a K_D = 74.36 μM

(Tab. 1**Error! Reference source not found.**), which is also apparent in the binding graph (Fig.

3 B). The strong increase in K_D is based on the over 16-fold lowered association k_{on} = 0.18 x

10^3 $M^{-1}s^{-1}$, while the dissociation k_{off} = 13.45 x 10^{-3} s^{-1} was also slightly lower when compared

to wild type MinD. Next, both monomeric mutants G12V and K16A were probed, and to our

surprise, capable of membrane interaction (Fig. 3 C & D, **Error! Reference source not**

found.). This is in stark contrast to *E. coli* MinD, where dimerization needs to precede

membrane binding to achieve sufficient affinity (45, 46, 65, 71, 72). Similar to I260E, the

G12V mutant displayed an around 10-fold reduced membrane affinity equilibrium with K_D =

75.38 μM (Tab. 1**Error! Reference source not found.**). However, G12V only exhibited a ~5

fold reduced binding affinity with k_{on} = 0.63 x 10^3 M^{-1} s^{-1} when compared to the wild type

MinD, while the k_{off} = 47.8 x 10^{-3} s^{-1} represents the fastest dissociation rate constant

between all MinD variants and more than 2-fold increase compared to wild type MinD (Tab.

1**Error! Reference source not found.**). This indicates a higher turnover and thus a quicker

release of the protein from the membrane. Markedly different from this, K16A displayed a

K_D of 1.36 μM, but not much lower binding rates (k_{on} = 2.03 x 10^3 M^{-1} s^{-1}, Tab. 1**Error!**

Reference source not found. & Fig. 3 C). Instead, the low K_D can be mostly attributed to the

reduced dissociation (k_{off} = 2.77 x 10^{-3} s^{-1}), implying a much longer retention of His-MinD-

K16A at the membrane (Fig. 3 D). Since G12V can bind ATP while K16A cannot, these findings

reveal that presence of ATP in the binding pocket affects association and dissociation of

MinD.

Last, the His-MinD-D40A mutant protein was probed, resulting in an almost 10-fold reduced

dissociation equilibrium K_D = 0.74 µM (Tab. 1Error! Reference source not found., Fig. 3 E).

Here, the strongly reduced membrane dissociation (k_{off} = 1.76 x 10^{-3} s^{-1}) causes the drastic

change in K_D, which we expected for the trapped dimer, as it could position the two

amphipathic helices into a more optimal orientation of the apolar residues towards the

membrane, resulting in stronger binding and no capability of hydrolyzing ATP and thus

relieving the dimer (68, 70, 71).

In summary, these BLI data demonstrate that *B. subtilis* MinD is capable of binding

membrane in both monomeric and dimeric form, while the presence of ATP in the binding

pocket appears to affect both membrane binding and dissociation.

MinC and the PDZ-domain of MinJ have no significant impact on MinD membrane kinetics.

In vivo, the integral membrane protein MinJ recruits MinD to poles and septum, which in

turn recruits MinC to form a functional complex that inhibits FtsZ-ring assembly. However,

the mechanistic details of how MinJ influences membrane association of MinD remain

unclear. Does MinJ actively recruit MinD from the cytoplasm, or does it primarily sequester

MinD already bound to the membrane? Furthermore, does MinJ promote MinD dimerization

or oligomerization, potentially enhancing ATPase activity and triggering membrane

dissociation? Similarly, MinC binding to MinD may influence the stability or turnover of MinD

at the membrane, yet its impact on association/dissociation kinetics is unknown. To address

these questions directly, we employed BLI to test the effect of His-MinC and His-PDZ on His-

MinD membrane kinetics.

After validation of the experimental setup (see Material and Methods), we next tested how

His-MinC and His-PDZ affect MinD-liposome binding kinetics by subjecting a liposome coated

sensors to solutions containing 8 µM His-MinD and varying concentrations [1, 2, 4, 8 and 16

µM] of His-MinC or His-PDZ, respectively (Fig. S 11, Tab. 2). MinD controls appear twice in

the table, because the experiments were performed using two independent purification

batches. The corresponding His-MinC and His-PDZ samples were each paired with the His-

MinD batch they contained. Due to batch-specific differences in His-MinD stability and

degradation, we did not directly compare measurements across batches. Interestingly,

neither His-MinC nor the His-PDZ domain significantly altered the membrane binding

kinetics of His-MinD in BLI experiments. The extracted K_D, k_{on}, and k_{off} values differed only

marginally between conditions (Tab. 2**Error! Reference source not found.**), and these

apparent shifts fell within the fitting uncertainty and typical inter-run variability of BLI. This

contrasts with the slight increase in His-MinD ATPase activity observed in the presence of

His-PDZ (Fig. S 5), which had led to the hypothesis that PDZ might promote MinD

oligomerization at the membrane, enhancing ATP hydrolysis. However, the discrepancy

between the two systems and the high lipid concentration (2 mg/ml) in the ATPase assay

versus a liposome-coated sensor in BLI may account for the different outcomes. In the BLI

experiment, limited surface area and distinct spatial organization of liposomes may restrict

the ability of the soluble PDZ domain to effectively cluster MinD, unlike full-length MinJ,

which is membrane-anchored via its transmembrane helices. Thus, while the PDZ domain

alone appears insufficient to modulate membrane binding kinetics of MinD, it remains

plausible that the full-length MinJ could exert a more pronounced effect through spatial

coordination at the membrane.

Single-molecule localization microscopy confirms *in vitro* data.

To validate the biochemical *in vitro* results, we decided to utilize single-molecule localization

microscopy (SMLM) with focus on single-molecule tracking (SMT), a technique allowing to

observe and evaluate MinD dynamics *in vivo* inside *B. subtilis* cells. We began SMLM analysis

by plotting a heat-map of the intracellular localization of all individually recorded Halo-MinD

molecules as well as the mutated variants, respectively, on normalized cells (Fig. 4 A). This

representation did not only provide a robust internal control for the imaging and post-

processing process for the well characterized localization of MinD, but also revealed

systemic differences in localization caused by the respective mutations. The first obvious

difference is reduced MinD enrichment at the cell center and poles in the G12V or K16A

variants. (Fig. 4 A). Here, the membrane-bound MinD fraction seems to be much more

evenly distributed along the membrane, similar to what we observed during wide-field

microscopy (Fig. 2 B). This result does not only confirm the ability of MinD monomers to bind

the membrane (Fig. 3 C & D), but also fuels the speculation that MinJ might be unable to

recruit MinD in its monomeric form, so that polar and septal recruitment by MinJ requires

dimeric MinD. In combination with the information obtained from BLI results (Fig. 3), this

could mean that the membrane itself serves as proxy for MinD dimerisation and subsequent

recruitment, quickly followed by ATP hydrolysis and membrane detachment of MinD. In

contrast, but in agreement with wide-field imaging (Fig. 2 B), the Halo-MinD-D40A mutant

does appear to frequently form foci and larger local accumulations, with little protein in the

cytosolic fraction (Fig. 4 A). This, again, could be caused by local MinJ interaction and

recruitment in combination with the missing capability of membrane detachment or with

increased lateral MinD interactions (73). Finally, the I260E mutant appears evenly distributed

throughout the cell (Fig. 4 A), as it was expected.

Independently, all protein trajectories were subjected to a stationary localization analysis

(SLA) (Fig. 4 B). This was done by first grouping all trajectories into an either mobile or static

population by testing if the molecule left a circle with a diameter of 97 nm (equaling the

pixel size of the imaging setup) within at least five consecutive frames. Trajectories that were

classified as mobile were further sub-grouped into mixed and free trajectories, where only

free trajectories never presented a confinement event (74). In agreement with previous

data, the D40A mutant displayed the highest fraction of static tracks (Fig. 4 B), and by far the

lowest proportion of truly free trajectories (48.5 %). In contrast, more than 90% of Halo-

MinD-I260E molecules were classified as free with only few trajectories classifying as static

(Fig. 4 B), accentuating the absence of interaction with the membrane or other structures.

When comparing Halo-MinD to the monomeric mutants G12V and K16A, the number of

static tracks was visibly reduced in the mutants. Since both monomers are unable to

dimerise, it is less likely that they can form larger clusters, which we observed in a previous

study (61). Between both monomeric mutants, K16A displayed a slightly larger number of

static trajectories (Fig. 4 B), which is in alignment with BLI data, where the K16A mutant

dissociates much less effective from the membrane (Fig. 2 C & D, Tab. 1**Error! Reference**

source not found.).

Next, we examined the average intracellular mobility of Halo-MinD compared with the

mutant variants by analyzing the mean-squared displacement (MSD, Fig. 4 C). In MSD

analysis, the traveled distances of recorded trajectories are plotted against a given time lag

(Δt) as a function of Δt, here one imaging frame (24 ms), where the slope can indicate the

speed (75-77). As expected, the I260E mutant displayed the fastest average mobility, since it

does not bind the membrane effectively anymore (Fig. 4 C). In contrast, the D40A mutant

was the slowest recorded protein, indicated by the only mild incline (Fig. 4 C). Since we

expect most of the proteins' population to be membrane bound, and furthermore observed

larger foci that suggest cluster formation, this was expected. Halo-MinD wild type protein

appeared to be more mobile in comparison (Fig. 4 C), as it should exist in both monomeric

and dimeric conformation and either freely diffusive in the cytosol or interact with the

membrane. Last, the two monomeric mutants G12V and K16A displayed extremely similar

average speeds that were both faster than the wild type protein (Fig. 4 C). Since these MinD

variants cannot form dimers, general diffusion should be faster, as there should be no MinD

dimers or oligomers, and possibly no recruitment by MinJ.

If the underlying movement in MSD analysis can be described by simple Brownian motion,

the gradient of the curve is proportional to the diffusion coefficient D (77). We know,

however, that MinD interacts at least with MinC, MinJ, the membrane and itself, in

combination with the fact that bacterial cells exhibit natural confinement. With these

information one cannot assume simple Brownian motion, therefore we have chosen jump

distance (JD) analysis as a more accurate assessment to describe Halo-MinD dynamics (Fig. 4

D & E). In JD analysis, the distances molecules travel between subsequent frames are plotted

in a probability distribution (Fig. 4 E), and diffusion coefficients are obtained by curve fitting

(77), in this case with a three-component model separating fast, slow and confined

molecules (Fig. 4 D & E). To accurately compare population sizes, the simultaneous option

was chosen in the SMTracker software, fixing the diffusion coefficients for every MinD

variant (74). To also obtain individual, variant specific diffusion coefficients, the same data

was additionally fitted individually (non-simultaneous SQD analysis in SMTracker2, Tab.

3**Error! Reference source not found.**).

Generally, the results of JD analysis correlated well with our *in vitro* data. When dissecting

the different populations in the wild type, more than a third of all proteins appear to be

confined (36.6%, Fig. 4 D), which likely comprises membrane bound protein that is further

stabilized by protein-protein interactions via lateral interaction of MinD with itself, and

possibly MinC and MinJ. The slow mobile population was the largest (45.7%, Fig. 4 D), and

likely summarizes different conformations of MinD that interact with the membrane, but are

still diffusive and not captured or stabilized through further interactions. The third, fast

diffusive population (17.7%, Fig. 4 D) likely reflects cytosolic MinD. These results stress the

importance of the MinD ATPase cycle for its localization and dynamics, as these are severely

altered in the MinD mutants.

In detail, the probability distributions of Halo-MinD-D40A and I260E (Fig. 4 E) illustrate the

strong shift between MinD molecules in a confined (D40A) and a diffusive state (I260E) best.

When examining the different populations, most Halo-MinD-D40A molecules were classified

as confined (54.2%) and slow mobile (40.2%), whereas I260E only displayed ~9% of confined

molecules (Fig. 4 D), matching prior results and expectations of the membrane binding

mutant. When comparing D40A to wild type Halo-MinD, the largest shift is observed

between the fast mobile and confined populations, shifting towards a distinctively more

confined regiment in the D40A mutant (36.6% WT, 52.2% D40A; Fig. 4 D) and visible

disappearance of the fast population in the probability distribution (Fig. 4 E). These results

are in line with the previous observations, as a trapped dimer of MinD will remain at the

membrane at much higher efficiency (Fig. 3 E & Tab. 1<mark>Error! Reference source not found.</mark>),

and appeared to form foci and clusters (Fig. 2 B, Fig. 4 A). G12V and K16A mutants on the

other hand showed an increase in fast mobile populations (WT 17.7%, G12V 25.8%, K16A

26.6%; Fig. 4 D E), which could be attributed to a reduced interaction with MinD interaction

partners (MinC, MinD, MinJ and plasma membrane). This notion is supported by the

decrease in confined molecules for both G12V and K16A mutants compared to the wild type

(WT 36.6%, G12V 23.7%, K16A 29%; Fig. 4 D & E), which are expected to be membrane

bound and likely stabilized through other protein-protein interaction, which may be reduced

or absent in the monomeric conformation. This notion is also supported by the absence of a

MinD gradient in epifluorescence images of G12V and K16A mutants (Fig. 2 B), a pattern

usually formed through sequestering via MinJ. Finally, when comparing both monomeric

variants, the K16A mutant displays a noticeable shift between the slow diffusive (G12V

50.5%, K16A 44.3%; Fig. 4 D) and confined populations (23.7% G12V, 29% K16A; Fig. 4 D).

The larger confined population of Halo-MinD-K16A is consistent with the BLI results (Fig. 3 D,

Tab. 1<mark>Error! Reference source not found.</mark>), where K16A appears to have a lower dissociation

from the membrane, increasing its likelihood of confinement, possibly caused by the inability

to bind ATP.

In summary, these results emphasize the importance of the MinD ATPase cycle for

functionality of the Min system in *B. subtilis*. Even though *B. subtilis* MinD is not capable of

oscillation, its intracellular dynamics are highly dependent on the functionality of the MinD

ATPase domain and its catalytic site, which is reflected by the variety of localization and

interaction defects displayed by the MinD mutants on a molecular level, that all result in

severe division anomalies.

MinC and MinJ have different effects on MinD protein dynamics in SMLM.

MinC and MinJ interact with MinD *in vivo*, suggesting protein dynamics should be affected

by this. However, in our ATPase and BLI experiments, MinC had only minor effects on ATPase

activity and membrane binding kinetics of MinD, while the PDZ domain slightly enhanced

ATPase activity without altering membrane binding kinetics of MinD. To validate these

observations *in vivo* and to identify differences between MinD interacting with a cytosolic

PDZ-domain or a full-length MinJ, we performed the same SMLM workflow and analysis

pipeline described in the previous section on cells expressing Halo-MinD in Δ*minC* and Δ*minJ*

backgrounds and compared them to the wild type background (Fig. S 12, Tab. 4).

When comparing the average Halo-MinD distribution between the respective strains, the

Δ*minC* mutant exhibited a similar intracellular localization pattern as the wild type, with

strong membrane enrichment and highest concentrations at midcell and poles (Fig. S 12 A).

However, Δ*minC* cells interestingly exhibited a slightly reduced cytosolic signal and a

corresponding increased membrane association. As expected, Δ*minJ* cells lacked the

characteristic septal accumulation. Instead, Halo-MinD was distributed relatively uniformly

along the membrane, consistent with MinJ being required for efficient sequestering of MinD

to the future division site and cell poles.

In the stationary localization analysis (Fig. S 12 B), Halo-MinD in the Δ*minC* strain displayed

only a modest increase in the static fraction compared to wild type (23.4 % from 20.1 %).

This shift can be attributed to a reduction in the mixed, likely membrane-interacting

population (from 19.0 % to 14.7 %), whereas the freely mobile fraction remained essentially

unchanged (61 % to 62 %). These results indicate that in the absence of MinC, less MinD

moves along the membrane and a slightly larger proportion becomes immobilized. Although

the effect size is small, it suggests that MinD may form larger or slightly more stable

membrane-associated assemblies when MinC is absent *in vivo*. In the Δ*minJ* background, the

static Halo-MinD population surprisingly increased (20.1 % to 28.2 %), together with

respective reductions in both the freely diffusing cytosolic pool (61.0 % to 56.6 %) and the

mixed, likely membrane-interacting population (19.0 % to 15.1 %). This redistribution

suggests that MinJ contributes to efficient MinD turnover on the membrane, in line with

MinD ATPase data with His-PDZ (Fig. S 12). In the absence of MinJ, MinD appears to remain

membrane-bound for longer, consistent with impaired detachment or reduced cycling

between membrane-associated states. One possible explanation is that MinD may form

more stable or aggregation-prone assemblies when MinJ is absent, whereas the presence of

MinJ could promote more dynamic membrane association and hence sharper local

accumulations and cycling of MinD, necessary for enriched polar and septal localization.

Jump-distance analysis with fixed diffusion coefficients corroborated these findings (Fig. S 12

C & D), showing that Δ*minJ* cells contained a substantially enlarged confined population of

MinD (27.9 % vs. 44.9 %), together with a large reduction of the slow diffusive population

(50.9 % vs. 37.8 %). Fitting the data without fixed diffusion coefficients produced the same

trend (Tab. 4), indicating that the shift reflects MinD spending more time in the respective

state rather than actual changes in the underlying diffusive coefficients.

In the Δ*minC* strain, Halo-MinD shows a slight redistribution toward the confined state, with

an increased fraction of very slow or immobile molecules (27.9 % vs. 36.8 %) and a

corresponding decrease in both the fast and intermediate populations (50.9 % vs. 45.8 % and

21.2 % vs. 17.4 %, respectively, Fig. S 12 C & D). SQD analysis without fixed diffusion

coefficient (Tab. 4) shows that the diffusion coefficients of the fast and intermediate

populations are reduced compared with wild-type, indicating that the mobile molecules

themselves move more slowly in average. Combined, this could suggest that in the absence

of MinC, membrane-bound MinD is more prone to forming clusters or larger aggregates,

which restrict mobility and reduce the efficiency of its membrane cycling.

Together, these data suggest that MinJ promotes productive membrane cycling of MinD,

and in its absence, MinD accumulates more often in a confined, likely non-productive

membrane-associated state, whereas MinC plays a comparatively minor regulatory role,

where it passively contributes to MinD mobility by preventing excessive clustering and

facilitating proper membrane-associated dynamics.

Discussion

In bacteria, the oscillation of MinCDE in *E. coli* is one of the best studied examples of a reaction-diffusion system, and the two core components MinD and MinE cycle between membrane and cytosol in an ATP-dependent manner, generating a standing wave with a time-resolved minimum at midcell (15, 83). Here, binding and unbinding of the components to the membrane surface influences the diffusion coefficients and controls the dynamic activity of MinC (84). While the *E. coli* MinCDE system is one of the best-studied examples of intracellular pattern formation in bacteria (15, 84-86), the Min system in *B. subtilis* remains less well characterized. In a previous study (61), we suggested that gradient formation depends on ATPase-driven MinD dynamics, backed by PALM and FRAP analysis as a basis for a mathematical model. However, due to a lack of biochemical data, we had to assume that the *B. subtilis* MinD cycles between membrane-bound and cytosolic localizations based on ATP-binding and hydrolysis, similar to *E. coli* MinD (61). A further gap in our knowledge of the *B. subtilis* Min system concerned the activation mechanism of MinD in absence of a MinE homolog.

To address this gap, we first analyzed *B. subtilis* MinD biochemically *in vitro*. Purified MinD showed basal ATPase activity that drastically increased upon liposome addition. Although *E. coli* MinD similarly requires lipids, it additionally depends on MinE for full activation (64). The reported maximum ATPase rate, around 20 nmol mg^{-1} min^{-1} (64, 65), was nearly identical to that of *B. subtilis* MinD, indicating that membrane binding alone is sufficient to trigger robust ATP hydrolysis in the *Bacillus* system. When comparing to other MinD/ParA ATPases that have been biochemically characterized, the activity level of *B subtilis* MinD is on the lower end of the spectrum with a k_{cat} of 36.27 h^{-1}. Soj, the ParA homologue in *B. subtilis*, reaches

more than 2-fold higher catalytic activity (83.6 h^{-1}) when fully activated by supplementing

SpoOJ (ParB) and DNA (87), similarly to Soj in *Thermus thermophiles* combined with SpoOJ

and *parS* containing DNA (70). Studies investigating the activity levels of ParA in *Caulobacter*

crescentus, purified in presence of ATP, determined a k_{cat} of 0.79 min^{-1} (47.4 h^{-1}) in absence

of ParB, increasing almost 5-fold to 3.7 min^{-1} (222 h^{-1}) in presence of ParB (88). A more

recent study found the *C. crescentus* ParA k_{cat} increase from 5.8 hr^{-1} to 120 hr^{-1} upon

addition of ParB (89).

These findings suggest that membrane association is the primary trigger for MinD ATPase

activity in *B. subtilis*. Specifically, the comparison of maximum velocities, activity stimulation

factors, and K_M values between the *E. coli* and *B. subtilis* MinD proteins suggests that a

functional homolog of the MinE protein is not necessary in *B. subtilis*. At the same time, our

biochemical data indicate that MinJ/PDZ increases MinD ATP hydrolysis rates at least to a

modest extent, (between 22% and 37%, Fig. S 5 B), pointing to a broader regulatory

influence of MinJ on MinD that becomes evident from additional mechanistic analyses

discussed below. In contrast, the presence of MinC led only to a slight decrease in MinD

ATPase activity in our APTase assays (~ 5%, Fig. S 5 A). This modest inhibitory effect suggests

that MinC does neither directly stimulate MinD turnover nor inhibit it, and may instead

stabilize MinD in a conformation or assembly state with mildly reduced catalytic activity, also

discussed below.

With the goal of testing binding of isolated MinD to the membrane, we designed a Bio-layer

interferometry experiment in which we immobilized liposomes and measured binding of

purified MinD proteins. Importantly, without addition of ATP (and ADP present from

purification) we observed only very weak MinD binding to the liposomes (Fig. S 13),

indicating that MinD membrane binding and dissociation is coupled to the ATP hydrolysis

cycle or the nucleotide binding state. In presence of ATP, MinD readily binds to the

liposomes with high affinity. The ParA/MinD family is well studied, and mutations producing

strictly monomeric (G12V, K16A), dimeric (D40A), or membrane-binding-deficient (I260E)

MinD variants are well established (68-70). The most intriguing observation of the binding

experiments is that MinD monomers can apparently bind to the liposome membrane *in*

vitro. Both monomeric mutants (G12V and K16A) are readily binding to the membrane, but

the release of the proteins from the membrane surface differs drastically. While the K16A

mutant, unable to bind ATP, has a slow membrane release rate (k_{off} rate of only $2.77*10^{-3}$ s^{-}

^{1}), the G12V monomeric mutant, which can still bind ATP, is released fast from the

membrane (k_{off} rate $47.8*10^{-3}$ s^{-1}). The D40A mutant of MinD, which is thought to be a stable

ATP-bound dimer, releases with very slow kinetics (k_{off} rate $1.76*10^{-3}$ s^{-1}). This indicates that

ATP hydrolysis accelerates MinD release from the membrane and that ATP binding itself

influences MinD association and dissociation kinetics, although we cannot entirely exclude

that the mutations introduce subtle structural changes that affect membrane affinity. These

findings crucially support a reaction diffusion model in which the MinD ATPase cycle is

essential for correct pattern formation of *B. subtilis* MinD. When testing the impact of MinC

on MinD membrane-binding kinetics, no significant differences were observed, suggesting

that MinC does not substantially influence the binding dynamics of MinD, in-line with the

collective view of MinC as a passenger rather than a driver of MinD dynamics (84). It would

have been desirable to study the interaction of MinD with reconstituted full-length MinJ

proteins to specifically investigate the influence that MinJ has on membrane interaction of

MinD. Unfortunately, we did not succeed in the purification of a full length MinJ protein so

far. MinJ contains seven predicted transmembrane helices and displays a pronounced

instability, rendering purification difficult. While the binding kinetics were unaffected during

BLI, our *in vitro* ATPase experiments combining the soluble PDZ domain of MinJ with MinD

already demonstrate that MinJ possibly modulates MinD:membrane interactions to some

extent through an effect on the hydrolysis efficiency.

To clarify this point and to better understand the isolated MinD membrane interaction under

in vivo conditions, we turned to state-of-the-art microscopy (SMLM) (92, 93).

SMLM data comparing the behavior of MinD in wild type vs. Δ*minJ* background (Fig. S 12)

collectively suggest that MinJ stabilizes correct pattern formation of MinD. By promoting

efficient membrane detachment and maintaining MinD in a dynamically cycling state, MinJ

appears to ensure that MinD remains properly redistributed throughout the cell cycle rather

than becoming sequestered in non-productive clusters. This organizational role likely

underlies the ability of MinJ to position MinD, and thus MinC, at the appropriate cellular

sites (55, 56), thereby supporting robust spatial regulation of division. However, a short-lived

MinJ mediated stabilization of MinD-dimers at the membrane would be comparable to the

stabilizing effect that MinE has on MinD oligomers in *E. coli* (90, 91), and was previously

observed in our *in vivo* FRAP studies (61). In contrast to MinJ, MinC did not seem to have a

major impact on MinD dynamics *in vivo*. When observing MinD in absence of *minC*, the main

effect was a subtle reduction in MinD cycling activity, shifting a small fraction of MinD into

more static or confined membrane-associated states. In the wild type situation, MinC may

passively restrict uncontrolled aggregation by capping or destabilizing certain interfaces, or

conversely transiently stabilize short-lived assemblies required for regulation.

The locked MinD dimers (D40A mutant) frequently nucleated into larger membrane clusters.

This cluster formation is underlined by the drastic increase of the confined/immobile

population (54.2%) of D40A in SMLM. This increase in the static population comes to the

expense of freely diffusive protein (~5.6 %), indicating that the vast majority of MinD D40A is

membrane associated. We observed a clear enrichment of these MinD clusters at the cell

poles and septa, lending support to the notion that MinD dimers are specifically

stabilized/recruited by MinJ. This interpretation is supported by previous data showing that

in absence of MinJ, MinD proteins are randomly localized along the membrane in clusters,

but not enriched at poles or (in a *minJ* mutant background rare) septa (57). In line with this

idea, the monomeric MinD variants (G12V, K16A) are evenly bound to the membrane

surface along the entire cell length.

The fact that most monomeric MinD appears membrane associated *in vivo* also supports our

in vitro measurements in which monomeric MinD variants efficiently bind the membrane, a

notion strongly supported by a recent report that examined localization of the same

monomeric variants in epifluorescence microscopy (94). The time-resolved MinD gradient is

lost in all mutants - including D40A - which should still be recruited by MinJ, suggesting the

ATPase cycle and thus MinD cycling is indeed necessary for proper formation of a gradient,

as suggested before (61). Our data are in perfect agreement with recent *in vitro* data

collected with the *E. coli* MinD using mass-sensitive particle tracking (MSPT) (73). MSPT

analysis revealed that *E. coli* MinD forms lateral interactions, leading to higher order

oligomers (trimer, tetramer and even higher oligomers). It was speculated that dimerization

of *E. coli* MinD at the membrane leads to conformational changes opening an additional

interaction surface of MinD for lateral interaction (73). These lateral MinD interactions were

thought to recruit further monomeric MinD from solution to the membrane, thereby

forming a nucleation for larger clusters. The MinD cluster formation that we observed in *B.*

subtilis using SMLM (61) are likely *in vivo* evidence for this cooperative membrane binding. It

should be noted that the MSPT analysis was not sensitive enough to detect monomer

binding to the membrane, but this was explicitly not ruled out (73). Addition of a second

amphipathic helix to the *E. coli* MinD however, was shown to promote monomer binding of

E. coli MinD to the membrane (66). Thus, the difference between the *E. coli* and *B. subtilis*

MinD is likely the higher membrane binding capacity of the *Bacillus* protein. The observed

cooperativity in membrane binding and thus nucleation, as shown for the *E. coli* MinD in

vitro (66, 73) and here for the *B. subtilis* MinD *in vitro* and *in vivo*, is required for pattern

formation by conferring nonlinearity (61, 95, 96).

Consistent with this model, our SMT data revealed three major dynamic states of MinD: fast

cytosolic diffusion, a slower membrane-associated pool, and a confined fraction

corresponding to clustered or multimeric assemblies. Most of the membrane-binding-

deficient mutant I260A shifted drastically into the fast, cytosolic population, validating this

assignment. Conversely, the locked-dimer (D40A) mutant showed a pronounced increase in

confined, slow-moving molecules, in line with its strong tendency to nucleate clusters.

Monomeric variants (G12V, K16A) displayed reduced confinement, supporting the view that

dimerization is required for higher-order assembly or recruitment. Together, these

observations indicate that MinD dynamically cycles between cytosol and membrane in an

ATP-dependent manner. Thus, we conclude that membrane bound dimers of MinD have a

much stronger tendency to interact with and recruit binding partners (including self-

nucleation). This finding is further supported by a recent report that analyzed MinD

dynamics *in vivo* using diffraction limited microscopy (94). In this study, a MinD D40A mutant

hyper-recruited MinC, leading to cluster formation of MinCD at the membrane. Deletion of

MinC restored the MinD gradient formation, indicating that MinC might indeed be part of a

MinC:MinD co-polymer, as suggested for the *E. coli* system (97). This assumption is in good

agreement with our *in vivo* findings, where self-nucleation seems to increase drastically for

the D40A mutant or in absence of MinJ. In contrast, monomeric G12V and K16A mutants do

not display this behavior, in line with monomeric MinD variants not being able to interact

with MinC, which was similarly observed through widefield microscopy in the same study

(94).

Collectively, our data reveal that the MinD ATPase cycle is essential for regulated membrane

binding and release, which is crucial for the formation of a sharp MinD gradient at the

septum (61). However, in contrast to *E. coli* MinD, monomeric *B. subtilis* MinD can bind the

membrane with higher affinity, suggesting that MinD also dimerizes at the membrane

surface. Dimer formation is essential for ATPase activity, as indicated by our *in vitro*

measurements, and is necessary for efficient membrane detachment. The main

characteristics of the *E. coli* MinD with respect to membrane binding, self-interaction leading

to nucleation, and membrane detachment are therefore shared in the *B. subtilis* protein. The

difference, that ATP hydrolysis of the *B. subtilis* MinD does not require a trigger other than

membrane binding, makes sense in a system that does not need to oscillate, but rather

accumulate and sustain a sharp gradient at the current site of division. However, the balance

of a dynamic membrane cycle of MinD seems to be fine-tuned by the interaction with MinC

and especially MinJ, which appears to keep MinD in a productive state and protects it from

aberrant aggregation, while recruiting it towards poles and septum, the intended site of

action of MinC.

Materials and Methods

Bacterial strains, plasmids and oligonucleotides

Plasmid and strain construction of all relevant constructs is described in the supplementary information. Oligonucleotides, plasmids and strains used in this study are listed in Tab. S 2, Tab. S 3 and Tab. S 4, respectively.

Media and growth conditions

E. coli was grown on lysogeny broth (LB) agar plates containing 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract, 10 g L^{-1} NaCl and 1.5% (w/v) agar at 37°C overnight using appropriate selection (kanamycin 50 µg ml^{-1}, chloramphenicol 35 µg ml^{-1}).

B. subtilis was grown on nutrient agar plates using commercial nutrient broth and 1.5% (w/v) agar at 37°C overnight. To reduce inhibitory effects, antibiotics were only used for transformations and when indicated, since allelic replacement is stable after integration (kanamycin 5 µg ml^{-1}, spectinomycin 100 µg ml^{-1}).

For microscopy, *B. subtilis* was inoculated to an OD_{600} 0.05 from a fresh overnight culture and grown in MD medium - a modified version of Spizizen Minimal Medium (98) – at 37°C with aeration in baffled shaking flasks (200 rpm) to OD_{600} 0.5-0.8. MD medium contains 10.7 mg ml^{-1} K_2HPO_4, 6 mg ml^{-1} KH_2PO_4, 1 mg ml^{-1} Na_3 citrate, 20 mg ml^{-1} glucose, 50 µg ml^{-1} L-tryptophan, 11 µg ml^{-1} ferric ammonium citrate, 2.5 mg ml^{-1} L-aspartate and 0.36 mg ml^{-1} $MgSO_4$ and was always supplemented with 1 mg ml^{-1} casamino acids. Subsequently, cultures were diluted to OD_{600} 0.05 in fresh, pre-warmed MD medium and grown to OD_{600} 0.5 (exponential phase).

Purification of His-MinC, His-PDZ, His-MinD and variants

For heterologous expression of His-MinC, His-PDZ and His-MinD and its variants, freshly

transformed BL21(DE3) cells carrying the respective pET-28a expression vector (Tab. S 3) as

well as pLysS (His-MinC, His MinD and variants only) were transferred to LB medium with

appropriate selection (kanamycin 50 µg ml^{-1}; chloramphenicol 35 µg ml^{-1} in presence of

pLysS) and grown overnight (37°C, 200 rpm). The next day, fresh LB medium was inoculated

1:150 from the overnight culture and grown to an OD$_{600}$ of 0.5 at 37 °C and 200 rpm. Since

His-PDZ tended to form inclusion bodies under these conditions, its expression culture was

instead grown at 18 °C and 120 rpm without IPTG induction. For all other constructs, protein

expression was induced with 1 mM IPTG. Cells expressing His-MinC and all His-MinD variants

were harvested after 3 h of induction, whereas His-PDZ cultures were collected after

overnight incubation. Cells were harvested by centrifugation (4 °C, 5000 × g, 15 min) and

stored at -80°C. For purification, a protocol of *E. coli* MinD purification was adapted from

(63): the cell pellet was resuspended on ice in lysis buffer (50 mM NaH$_2$PO$_4$, 500 mM NaCl

and 10 mM imidazole; pH 8.0), supplemented with protease inhibitor cocktail (Roche), 0.2

mM Mg^{2+}-ADP and 0.4 mM TCEP (His-MinD and variants only) and ~250 U mL^{-1} DNase I

(Roche), and subsequently lysed in a pre-cooled (4°C) French pressure cell (Amico; 3 x 1,200

psi). The debris was then removed via centrifugation (10,000 g, 30 min, 4°C) and the

supernatant was passed through a 1 ml Ni-NTA column (Protino, Macherey-Nagel) utilizing

liquid chromatography (ÄKTApurifier). Here, the resin was washed with 10 ml binding buffer

A (50 mM NaH$_2$PO$_4$, 500 mM NaCl, 20 mM imidazol; pH 8) before elution buffer B (50 mM

NaH$_2$PO$_4$, 500 mM NaCl, 250 mM imidazol; pH 8, (and 0.2 mM Mg^{2+}-ADP and 0.4 mM TCEP

for His-MinD and variants)) was added sequentially at concentrations of 5% (10 ml) 10% (20

ml), and finally 100% (10 ml). Recombinant protein eluate was collected in 0.5 ml fractions.

Fractions containing high concentrations of protein (determined by chromatogram analysis,

sodium dodecyl-sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting

using Penta-His antibody (QIAGEN)) were pooled and concentrated using Amicon filter

devices with 10 kDa molecular weight cut-off (Millipore; Merck). Additionally, protein

aggregates were removed by pelleting them through centrifugation (10,000 g, 10 min, 4°C).

Samples were further purified by size-exclusion chromatography through a Superdex200

Increase 10/300 GL column (Cytiva) at a flow rate of 0.75 ml min^{-1} using gel filtration buffer C

(50 mM HEPES-KOH, 150 mM KCl, 10% (v/v) glycerol, 0.1 mM EDTA, pH 7.2). For His-MinD

and its variants, buffer C was supplemented with 0.4 mM TCEP and 0.2 mM Mg^{2+}-ADP

immediately before use. His-PDZ was not subjected to size-exclusion chromatography

because it could not be recovered after SEC. Instead, it was used directly following affinity

purification. To calibrate the column and allow identification of relevant fractions, a gel

filtration standard (Bio-Rad) was run according to the manufacturers protocol and analyzed

using the same conditions, respectively. Fractions containing recombinant protein were

pooled and either directly used for the respective assays, stored at 4°C for a maximum of

one week or flash-frozen and stored at -80 °C. Final protein concentration was determined

using Bradford assay with bovine serum albumin (BSA) as standards. Protein purity and

stability was confirmed via SDS-PAGE using 12% polyacrylamide gels and Western blotting

using Penta-His antibody (QIAGEN) (Fig. S 1, Fig. S 3, Fig. S 4, Fig. S 6 Fig. S 8, Fig. S 9 and Fig.

S 10).

ATP hydrolysis assays

ATP hydrolysis was analyzed in continuous reactions using the EnzChek Phosphate Assay Kit

(Thermo Fisher Scientific), according to the manufacturer's manual. Reaction volumes were

reduced to 100 µl and assayed in flatbottom 96-well plates (Greiner-UV-Star 96-well plates).

Reactions were measured at 37 °C continuously every minute over a time-course of 3 h in a

Tecan Infinite 200 Pro with the Software Tecan i-control v.2.0. If not indicated differently,

the reactions contained either 0-16 µM His-MinD or the respective variant/protein

combination as well as 2 mM Mg^{2+}-ATP and 0.2 mg ml^{-1} liposomes. Liposomes were made

from *E. coli* total extract (Avanti) by first drying and then resuspending and rehydrating lipids

in gel filtration buffer C (50 mM HEPES-KOH, 150 mM KCl, 10% (v/v) glycerol, 0.1 mM EDTA).

Next, the lipids were extruded 40 times through 100 nm polycarbonate membranes

(Nuclepore™ Track-Etched Membrane 0.1 µm, Whatman). Before starting the reactions with

2 mM Mg^{2+}-ATP, the plate was preincubated for 10 min in order to eliminate phosphate

contamination.

Data were first analyzed with Excel (normalized, subtraction of ATP auto hydrolysis and

subtraction of no substrate control), a linear regression was used to determine the

hydrolysis rate per minute. Data visualization and further analysis was performed using

GraphPad Prism version 9.4.0 for Windows, (GraphPad Software). Statistical comparisons

between His-MinD alone (n = 2) and His-MinC/His-PDZ conditions (n = 3 each) were

performed using Welch's t-test (two-tailed), which is appropriate for unequal variances and

unequal sample sizes.

AlphaFold prediction and structure-based analysis

A structural model of *B. subtilis* MinD was calculated using AlphaFold2 (99) installed on a

local computer (Fig. 2 A), where the structural homology search was performed against the

full AlphaFold database as of 11.02.2024 (99).

A structural model of *B. subtilis* MinJ was calculated using AlphaFold3 (100), where the

structural homology search was performed against the full AlphaFold database as of

01.02.2025 (100).

Bio-layer Interferometry

Measurements were performed on the BLItz platform (Sartorius) using Streptavidin (SA)

Biosensors (Sartorius, 18-5019). The BLItz device was operated with the Software BLItz Pro

(version 1.3.1.3) using the advanced kinetics protocol modified as shown in Table S5 and

essentially as described before (101-103).

Beforehand, Liposomes were made from a 33:1 mixture of *E. coli* total extract (Avanti) and

DSPE-PEG$_{(2000)}$-biotin (Avanti) by first drying and then resuspending and rehydrating lipids in

gel filtration buffer C (50 mM HEPES-KOH, 150 mM KCl, 10% (v/v) glycerol, 0.1 mM EDTA).

Next, the lipids were extruded 40 times through 100 nm polycarbonate membranes

(Nuclepore™ Track-Etched Membrane 0.1 μm, Whatman). Biosensors were hydrated for at

least 10 min in a 96-well plate in 200 μl gel filtration buffer C.

As a starting point, unspecific binding was tested. For this, non-biotinylated His-MinD was

used to be loaded onto the Streptavidin biosensor. For the binding assays, all measurements

were performed in black reaction tubes with 200 μl gel filtration buffer C or the indicated

solution (Tab. S 5). After measuring the initial baseline, biotinylated liposomes were allowed

to bind the biosensor. Another baseline was measured in the same tube as before. For

association, 200 μl solutions of His-MinD and variants in different concentrations and/or in

combination with His-MinC or His-PDZ, respectively, were exposed to the sensor, first

without and later with addition of 2 mM Mg-$^{2+}$ATP (final concentration) just before use.

Furthermore, unspecific binding to liposome-saturated sensors was tested using bovine

serum albumin (BSA, 16 µM). For dissociation, the sensor was exposed to buffer again. Each

measurement was performed with a fresh biosensor.

For BLI experiments involving His-MinD and one of its direct interactors, we first assessed

whether His-MinC or His-PDZ can directly interact with His-MinD *in vitro*. To this end, a

liposome-coated BLI sensor was first dipped into a 8 µM His-MinD solution. Subsequently,

the sensor was exposed to solutions containing either 8 µM His-MinD together with 8 µM

His-MinC or 8 µM His-PDZ, respectively. As a negative control, a sample containing 8 µM

His-MinD and 8 µM BSA was used. Both His-MinC and His-PDZ induced a significant increase

in the equilibrium binding signal upon interaction with immobilized His-MinD, consistent

with specific complex formation. In contrast, the BSA control showed only a minimal signal

increase, confirming the specificity of the observed binding. To further validate the sensor

specificity, individual proteins were tested. Only His-MinD exhibited robust binding to the

liposome-coated sensor, while His-MinC and His-PDZ showed negligible signal, indicating

non-specific interactions (Fig. S11). These initial tests support the expected specificity of the

MinD:MinC and MinD:PDZ interactions.

Kinetic analysis and fitting were performed by the BLItz Pro Software (version 1.3.1.3). Data

was exported and visualized using GraphPad Prism version 9.4.0 for Windows, (GraphPad

Software).

The BLItz Pro Software analysis follows a 1:1 binding model, where:

$$A + B \underset{k_{\text{off}}}{\overset{k_{\text{on}}}{\longleftrightarrow}} AB$$

806   with k_{on} = rate of association or "on-rate" and k_{off} = rate of dissociation or "off-rate".

807   Formula for fitting the association (a = slope):

$$Y = Y_0 + a\ (1 - e^{-k_{\mathrm{obs}} \times t)}$$

$$k_{\mathrm{obs}} = \ k_{\mathrm{on}} \times [A(M)] + k_{\mathrm{off}}$$

808   Formula for fitting the dissociation:

$$Y = \ Y_0 + a\ \times e^{-k_{\mathrm{off}} \times t}$$

809   Resulting in the equilibrium dissociation constant K_{D}:

$$K_{\mathrm{D}} = \frac{k_{\mathrm{off}}}{k_{\mathrm{on}}}$$

Fluorescence microscopy

Microscopy slide preparation

For epifluorescence imaging, *B. subtilis* cells were mounted on 1.5% MD agarose pads using

1.5 x 1.6 cm "Gene Frames" (Thermo Scientific), and incubated 10 min at 37°C before

microscopic analysis.

For SMLM, slides and coverslips were first cleaned by overnight storage in 1 M KOH,

carefully rinsed with ddH2O, and subsequently dried with pressurized air. Next, 1.5% (w/v)

low melting agarose (Sigma-Aldrich) was dissolved in MD medium. Medium was sterile

filtered (0.2 μm pore size) shortly before being used to remove particles. To produce flat,

uniform, and reproducible agarose pads, 1.5 x 1.6 cm "Gene Frames" (Thermo Fisher) were

utilized, and pads were allowed to solidify for 1 h at room temperature to be used within the

next 3 h.

Staining of HaloTag

For staining, 1 mL of bacterial culture was moved to a 2 mL microcentrifuge tube and incubated with 5 nM (SMLM) or 50 nM (epifluorescence) HaloTag TMR ligand at 37 ⬚C for 10 min. Cells were harvested (4000 g, 3 min, 37 °C) and washed four times in pre-warmed MD medium, before being mounted on the previously described microscopy slides.

Epifluorescence imaging

For strain characterization (Fig. 2 B), microscopy images were taken on a Axio Observer 7 microscope (Zeiss) equipped with a Colibri 7 LED light source (Zeiss) and an OrcaR[2] camera (Hamamatsu) using a Plan-Apochromat 100×/1.4 oil Ph3 objective (Zeiss). TMR fluorescence was visualized with a 90 HE LED filter set (QBP 425/30 + 514/30 + 592/30 + 709/100) (Zeiss) and excited at 555 nm (100% intensity of Colibri 7 LED) for 400 ms. The microscope was equipped with an environmental chamber set to 37 °C. Digital images were acquired with Zen Blue (Zeiss) and analyzed and edited with Fiji (104) ImageJ2 (105). For measurement of cell length and minicells, cells were stained with NileRed (1 µg/ml final concentration) at OD_{600} 0.5 and incubated for further 20 minutes (200 rpm, 37°C) before mounting them on an agarose pad. To minimize bias during analysis, cell lengths and number of minicells were identified using the Fiji (104) plugin MicrobeJ 5.11c (106) on phase contrast images with manual correction by screening the results against the simultaneously acquired membrane stain images and sorting out mislabeled cells. For cell length measurements, MicrobeJ modules *options*, *exclude on edges*, *shape descriptor*, *segmentation* and *chain* were used with default settings, except for following parameters in *options* (*area* 1-max; *length* 1-max; *width* 0.6-1; *sinuosity* 1-max; *angularity* 0-0.5) and *chain* (*tolerance* 0.1; *area* 1.5-max). To identify minicells, MicrobeJ modules *options*, *exclude on edges*, *shape descriptor* and

segmentation were used with default settings, except for following parameters in *options*

(*area* 0.15-1; *length* 0.5-1.2; *width* 0.3-1; *circularity* 0.95-max).

SMLM imaging

SMLM imaging was performed with an Elyra 7 (Zeiss) inverted microscope equipped with

two pco.edge sCMOS 4.2 CL HS cameras (PCO AG), connected through a DuoLink (Zeiss), only

one of which was used in this study. Cells were observed through an alpha Plan-Apochromat

63x/1.46 Oil Korr M27 Var2 objective, yielding a final pixel size of 97 nm. During image

acquisition, the focus was maintained with the help of a Definite Focus.2 system (Zeiss).

Fluorescence was excited with a 561 nm (100 mW) laser, and signals were observed through

a multiple beam splitter (405/488/561/641 nm) and laser block filters (405/488/561/641

nm) followed by a Duolink SR QUAD (Zeiss) filter module (secondary beam splitter: LP 560,

emission filters: EF BP420-480 + BP495-550).

Cells were illuminated with the 561 nm laser (50% intensity) in TIRF mode (62° angle). For

each time lapse series 10,000 frames were taken with 20 ms exposure time (~24 ms with

transfer time included) and 50% 561 nm intensity laser.

SMLM analysis

For average localization analysis (Fig. 4 A), a single-molecule localization table was

constructed from a 2D Gaussian fitting in the ZEN 3.0 SR (black) software (Zeiss) with a peak

mask size of 9 and a peak intensity-to-noise ratio of 6, and further allowing molecules to

overlap. Filtering was used to minimize noise, background, and out-of-focus emitters,

discarding events that did not emit within a Point spread function size (PSF) of 100-200 nm

and did not display a photon count of 50-600. Further analysis was performed using

individual Fiji (104) scripts for the cell outlines and R (4.2.2) scripts in R Studio

(2023.12.1+402) for the corresponding localization events (107, 108). Here, the remaining

events were transformed to align the orientation of all cells in which the signals were

detected, to be able to summarize all recorded localizations in one normalized cell per strain.

The localization frequency is displayed as a heatmap using the gradual viridis scaling from

the ggplot2 R package (109), where the highest frequency (normalized frequency 1) is

displayed in yellow and the lowest frequency (normalized frequency 0) in blue (Fig. 4 A).

For single-molecule tracking, spots were identified with the LoG Detector of TrackMate

v6.0.1 (110) implemented in Fiji 1.53 g (104), an estimated diameter of 0.5 µm, and median

filter and sub-pixel localization activated. The signal-to-noise threshold for the identification

of the spots was set at 7. To limit the detection of particles to single-molecules, frames

belonging to the bleaching phase of TMR were removed individually from the time lapses

prior to the identification of spots. Spots were merged into tracks via the "Simple LAP

Tracker" of TrackMate, with a maximum linking distance of 500 nm and no frame gaps

allowed. Only tracks with a minimum length of 5 frames were used for further analysis,

yielding a minimum number of total tracks per sample of 2596.

To identify differences in protein mobility and/or behavior, the resulting tracks were

subjected to stationary localization analysis (SLA), mean-squared-displacement (MSD) and

square displacement (SQD) analysis in SMTracker 2.0 (74). SLA analysis was performed with

a confinement circle radius of 97 nm and a minimum number of 5 steps to be considered

static. The average MSD was calculated for four separate time points per strain (exposure of

20 ms - τ = 24, 48, 72 and 96 ms), followed by fitting of the data to a linear equation. The last

time point of each track was excluded to avoid track-ending-related artifacts. The cumulative

probability distribution of the square displacements (SQD) was used to estimate the

diffusion constants and relative fractions of up to three diffusive states. Diffusion constants

were determined simultaneously for the compared conditions, therefore allowing for a more

direct population fractions comparison. Additionally, the analysis was performed for each

strain non-comparatively, yielding individual diffusion coefficients, displayed in Tab. 3**Error!**

Reference source not found..

Acknowledgments

This research was supported by the Deutsche Forschungsgemeinschaft (DFG) within the Transregio Collaborative Research Center (TRR 174) "Spatiotemporal Dynamics of Bacterial Cells". We thank Leendert Hamoen (Amsterdam) and Henrik Strahl (Newcastle) for communicating unpublished results prior to publication. We are also grateful to Anna Becker and Vivian Hennig (Kiel) for help in the initial stages of this project as part of their BSc theses.

Author Contributions

M.B. and H.F. conceived the study, H.F. constructed the strains, performed the *in vivo* and *in vitro* experiments incl. microscopy. C.D. performed *in vitro* experiments. H.F., C.D. and M.B. analyzed the data, H.F. and M.B. wrote the manuscript.

Declaration of Interests

The authors declare no competing interests.

 # Figure legends

Fig. 1: Biochemical analysis of *B. subtilis* MinD ATPase cycle.

(A) Phosphate release plotted against different MinD concentrations, fitted with a simple linear regression (R^2 = 0.97). Phosphate release was measured using the EnzChek™ phosphate assay kit. Samples contained 0.2 mg ml^{-1} liposomes and were pre-incubated for 10 minutes before addition of 2 mM Mg^{2+}-ATP; n ≥ 3.

(B) Specific activity of the MinD ATPase from (A) measured as a function of MinD concentration.

(C) Specific activity of MinD measured as a function of ATP concentration, determined as in (B) with fixed MinD concentration of 10 µM. Fitting the Michaelis–Menten equation (black lines) gives k_{cat} = 36.27 h^{-1}, V_{max} = 19.50 nmol mg^{-1} min^{-1} and K_M = 0.173 mM.

(D) Specific activity of MinD measured as a function of liposome concentration, determined as in (B) with fixed MinD concentration of 6 µM. Fitting the Michaelis–Menten equation (black lines) gives k_{cat} = 41.01 h^{-1}, V_{max} = 22.05 nmol mg^{-1} min^{-1} and K_M = 0.0437 mg ml^{-1}.

Fig. 2: Cartoon of MinD AlphaFold model with highlighted key amino acids and the effects

of their mutagenesis.

(A) Left: Rendering of MinD AlphaFold model with ATP binding region highlighted in light

blue, membrane targeting sequence (MTS) in turquoise and key residues in orange; Box:

explanation of mutagenesis effects. Right: Close-up on hydrophobic region of amphipathic

helix.

(B) Representative wide-field microscopy images of *B. subtilis* expressing indicated Halo-

MinD variants as the only MinD copy from the native locus, stained with TMR ligand. Left:

Phase contrast; scale bar 5 μm. Right: fluorescence.

Fig. 3: BLI analysis of His-MinD and different mutants.

(A) Left: Cartoon representation of the different BLI steps, starting with (1) establishing a

baseline in protein buffer, (2) binding of liposomes through the biotinylated DSPE-PEG(2000)

phospholipid, (3) establishing a new baseline, (4) binding of MinD and finally (5) dissociation

of MinD. Right: Exemplary graph resulting from sensor readouts of steps (1) - (5). Scheme

adapted from (102).

(B) MinD binding plotted against time through association and dissociation phase (steps 4-5)

at different protein concentrations, including the His-MinD-I260E mutant.

(C) – (E) Same as (B) using the indicated respective mutant of His-MinD (G12V, K16A and

D40A).

Fig. 4: Single-molecule localization microscopy analysis of Halo-MinD and mutants expressed in *B. subtilis*.

Exponentially growing *B. subtilis* cells expressing Halo-MinD and variants (n ≥ 48 cells, respectively) were stained with TMR ligand and subsequently imaged. Individual protein trajectories were recorded using SMLM and analyzed with Zen blue (Zeiss), Trackmate, the SMTracker 2.0 software package and manual scripts in R. Minimum track-length 4 frames of 24 ms, with at least 2596 trajectories per strain.

(A) Heat map representation of intracellular localization of individual molecules of Halo-MinD and variants, respectively, plotted on normalized cells. Brighter colors indicate higher abundance.

(B) Barplot of stationary localization analysis (SLA), comparing different track types within the protein population. Tracks were considered static when not leaving a circular area of 97 nm diameter within 5+ frames. Mobile populations were further divided into free and mixed tracks, where mixed tracks displayed a switch between free and confined movement.

(C) Plot of the mean-squared displacement of Halo-MinD and variants over time, fitted with a linear fit excluding the last timelag.

(D) Bubble plot displaying single-molecule diffusion rates of the indicated MinD fusions. Populations were determined by fitting the probability distributions of the frame-to-frame displacement (jump distance) data of all respective tracks to a three components model (fast mobile, slow mobile and confined protein populations).

(E) Probability distributions of jump distances of Halo-MinD and variants. Data was fit with a three component model, indicating confined, slow and fast tracks.

 # Tables

977   **Tab. 1: Kinetic constants of His-MinD and variants obtained via BLI.**

Protein	K_D [μM]	k_{on} [M^{-1} s^{-1}]	$k_{on\ error}$	k_{off} [s^{-1}]	$k_{off\ error}$	X^2	R^2
MinD	7.17	3.03 x 10^3	0.13 x 10^3	21.7 x 10^{-3}	0.33 x 10^{-3}	16.07	0.95
MinD-G12V	75.38	0.63 x 10^3	0.19 x 10^3	47.8 x 10^{-3}	1.13 x 10^{-3}	13.89	0.95
MinD-K16A	1.36	2.03 x 10^3	0.062 x 10^3	2.77 x 10^{-3}	0.088 x 10^{-3}	6.97	0.95
MinD-D40A	0.74	2.39 x 10^3	0.053 x 10^3	1.76 x 10^{-3}	0.069 x 10^{-3}	6.01	0.98
MinD-I260E	74.36	0.18 x 10^3	0.19 x 10^3	13.45 x 10^{-3}	0.38 x 10^{-3}	3.61	0.95

978   K_D = equilibrium dissociation constant; k_{on} = association rate constant; k_{off} = dissociation rate
979   constant

980 **Tab. 2: Kinetic constants of His-MinD in combination with His-MinC or His-PDZ.**

Protein	K_D [µM]	k_{on} [M^{-1} s^{-1}]	$k_{on \text{ error}}$	k_{off} [s^{-1}]	$k_{off \text{ error}}$	X^2	R^2
MinD*	31.7	0.27×10^3	0.21×10^3	8.65×10^{-3}	0.26×10^{-3}	4.10	0.957
+ MinC	33.4	0.35×10^3	0.084×10^3	11.7×10^{-3}	0.16×10^{-3}	8.07	0.979
MinD*	51.1	0.34×10^3	0.12×10^3	17.5×10^{-3}	0.24×10^{-3}	2.40	0.986
+ PDZ	47.5	0.35×10^3	0.14×10^3	16.6×10^{-3}	0.25×10^{-3}	8.98	0.973

K_D = equilibrium dissociation constant; k_{on} = association rate constant; k_{off} = dissociation rate
constant. *His-MinD control appears twice, from independent purifications. Since batches were not
directly comparable, controls are only compared to samples containing the same batch.
Concentrations: His-MinD 8 µM, His-MinC 1-16 µM [1, 2, 4, 8 & 16], His-PDZ 1-16 µM [1, 2, 4, 8 & 16],
see Fig. S 11.

Tab. 3: Diffusion coefficients obtained from non-comparative SQD analysis of Halo-MinD
and variants fitted with three populations.

Strain	Fast diffusive [$\mu m^2 s^{-1}$]	Slow diffusive [$\mu m^2 s^{-1}$]	Confined [$\mu m^2 s^{-1}$]
MinD WT	0.60 (22.3 %)	0.084 (50.4 %)	0.020 (27.3 %)
MinD-G12V	0.68 (28.5 %)	0.10 (47.5 %)	0.026 (24.0 %)
MinD-K16A	0.70 (27.3 %)	0.11 (43.9 %)	0.024 (28.8 %)
MinD-D40A	0.44 (13.7 %)	0.063 (50.0 %)	0.021 (36.3 %)
MinD-I260E	0.76 (73.4 %)	0.095 (20.3 %)	0.022 (6.3 %)

989 Percentages indicate relative population size

990 **Tab. 4: Diffusion coefficients obtained from non-comparative SQD analysis of Halo-MinD in**
991 **different genetic backgrounds fitted with three populations.**

Strain	Fast diffusive [$\mu m^2\,s^{-1}$]	Slow diffusive [$\mu m^2\,s^{-1}$]	Confined [$\mu m^2\,s^{-1}$]
WT	0.57 (22.1 %)	0.083 (50.2 %)	0.019 (27.7 %)
Δ*minC*	0.31 (29.3 %)	0.059 (44.3 %)	0.01 (26.4 %)
Δ*minJ*	0.6 (19 %)	0.069 (41.2 %)	0.018 (39.7 %)

992 Percentages indicate relative population size

References

1. Kotak S, Gonczy P. 2013. Mechanisms of spindle positioning: cortical force generators in the limelight. Curr Opin Cell Biol 25:741-8.

2. Pollard TD, O'Shaughnessy B. 2019. Molecular Mechanism of Cytokinesis. Annu Rev Biochem 88:661-689.

3. Cameron TA, Margolin W. 2024. Insights into the assembly and regulation of the bacterial divisome. Nat Rev Microbiol 22:33-45.

4. Mahone CR, Goley ED. 2020. Bacterial cell division at a glance. J Cell Sci 133.

5. den Blaauwen T, Hamoen LW, Levin PA. 2017. The divisome at 25: the road ahead. Curr Opin Microbiol 36:85-94.

6. Bi EF, Lutkenhaus J. 1991. FtsZ ring structure associated with division in Escherichia coli. Nature 354:161-4.

7. Lowe J, Amos LA. 1998. Crystal structure of the bacterial cell-division protein FtsZ. Nature 391:203-6.

8. Schaper S, Brito AD, Saraiva BM, Squyres GR, Holmes MJ, Garner EC, Hensel Z, Henriques R, Pinho MG. 2024. Cell constriction requires processive septal peptidoglycan synthase movement independent of FtsZ treadmilling in Staphylococcus aureus. Nat Microbiol 9:1049-1063.

9. Whitley KD, Jukes C, Tregidgo N, Karinou E, Almada P, Cesbron Y, Henriques R, Dekker C, Holden S. 2021. FtsZ treadmilling is essential for Z-ring condensation and septal constriction initiation in Bacillus subtilis cell division. Nat Commun 12:2448.

10. Squyres GR, Holmes MJ, Barger SR, Pennycook BR, Ryan J, Yan VT, Garner EC. 2021. Single-molecule imaging reveals that Z-ring condensation is essential for cell division in Bacillus subtilis. Nat Microbiol 6:553-562.

11. Egan AJF, Errington J, Vollmer W. 2020. Regulation of peptidoglycan synthesis and remodelling. Nat Rev Microbiol 18:446-460.

12. Adams DW, Errington J. 2009. Bacterial cell division: assembly, maintenance and disassembly of the Z ring. Nat Rev Microbiol 7:642-53.

13. Chan H, Mohamed AMT, Grainge I, Rodrigues CDA. 2022. FtsK and SpoIIIE, coordinators of chromosome segregation and envelope remodeling in bacteria. Trends Microbiol 30:480-494.

14. Reyes-Lamothe R, Sherratt DJ. 2019. The bacterial cell cycle, chromosome inheritance and cell growth. Nat Rev Microbiol 17:467-478.

15. Bramkamp M, van Baarle S. 2009. Division site selection in rod-shaped bacteria. Curr Opin Microbiol 12:683-8.

16. Szwedziak P, Ghosal D. 2017. FtsZ-ring Architecture and Its Control by MinCD. Subcell Biochem 84:213-244.

17. Ortiz C, Natale P, Cueto L, Vicente M. 2016. The keepers of the ring: regulators of FtsZ assembly. FEMS Microbiol Rev 40:57-67.

18. Rodrigues CD, Harry EJ. 2012. The Min system and nucleoid occlusion are not required for identifying the division site in *Bacillus subtilis* but ensure its efficient utilization. PLoS Genet 8:e1002561.

19. Wu LJ, Errington J. 2011. Nucleoid occlusion and bacterial cell division. Nat Rev Microbiol 10:8-12.

20. Bernhardt TG, de Boer PA. 2005. SlmA, a nucleoid-associated, FtsZ binding protein required for blocking septal ring assembly over Chromosomes in E. coli. Mol Cell 18:555-64.

21. Cho H, McManus HR, Dove SL, Bernhardt TG. 2011. Nucleoid occlusion factor SlmA is a DNA-activated FtsZ polymerization antagonist. Proc Natl Acad Sci U S A 108:3773-8.

22. Tonthat NK, Arold ST, Pickering BF, Van Dyke MW, Liang S, Lu Y, Beuria TK, Margolin W,
Schumacher MA. 2011. Molecular mechanism by which the nucleoid occlusion factor, SlmA,
keeps cytokinesis in check. EMBO J 30:154-64.
23. Wu LJ, Errington J. 2004. Coordination of cell division and chromosome segregation by a
nucleoid occlusion protein in Bacillus subtilis. Cell 117:915-25.
24. Wu LJ, Ishikawa S, Kawai Y, Oshima T, Ogasawara N, Errington J. 2009. Noc protein binds to
specific DNA sequences to coordinate cell division with chromosome segregation. EMBO J
28:1940-52.
25. Jalal ASB, Tran NT, Wu LJ, Ramakrishnan K, Rejzek M, Gobbato G, Stevenson CEM, Lawson
DM, Errington J, Le TBK. 2021. CTP regulates membrane-binding activity of the nucleoid
occlusion protein Noc. Mol Cell 81:3623-3636 e6.
26. Adams DW, Wu LJ, Errington J. 2015. Nucleoid occlusion protein Noc recruits DNA to the
bacterial cell membrane. EMBO J 34:491-501.
27. Adler HI, Fisher WD, Cohen A, Hardigree AA. 1967. MINIATURE escherichia coli CELLS
DEFICIENT IN DNA. Proc Natl Acad Sci U S A 57:321-6.
28. Schaumberg TH, Kuempel PL. 1983. Genetic mapping of the minB locus in Escherichia coli K-
12. J Bacteriol 153:1063-5.
29. Davie E, Sydnor K, Rothfield LI. 1984. Genetic basis of minicell formation in Escherichia coli K-
12. J Bacteriol 158:1202-3.
30. de Boer PA, Crossley RE, Rothfield LI. 1989. A division inhibitor and a topological specificity
factor coded for by the minicell locus determine proper placement of the division septum in
E. coli. Cell 56:641-9.
31. de Boer PA, Crossley RE, Rothfield LI. 1992. Roles of MinC and MinD in the site-specific
septation block mediated by the MinCDE system of Escherichia coli. J Bacteriol 174:63-70.
32. Hu Z, Mukherjee A, Pichoff S, Lutkenhaus J. 1999. The MinC component of the division site
selection system in Escherichia coli interacts with FtsZ to prevent polymerization. Proc Natl
Acad Sci U S A 96:14819-24.
33. Cordell SC, Anderson RE, Lowe J. 2001. Crystal structure of the bacterial cell division inhibitor
MinC. EMBO J 20:2454-61.
34. Hu Z, Lutkenhaus J. 2000. Analysis of MinC reveals two independent domains involved in
interaction with MinD and FtsZ. J Bacteriol 182:3965-71.
35. Shen B, Lutkenhaus J. 2009. The conserved C-terminal tail of FtsZ is required for the septal
localization and division inhibitory activity of MinC(C)/MinD. Mol Microbiol 72:410-24.
36. Shen B, Lutkenhaus J. 2010. Examination of the interaction between FtsZ and MinCN in E. coli
suggests how MinC disrupts Z rings. Mol Microbiol 75:1285-98.
37. Marston AL, Errington J. 1999. Selection of the midcell division site in Bacillus subtilis through
MinD-dependent polar localization and activation of MinC. Mol Microbiol 33:84-96.
38. Gregory JA, Becker EC, Pogliano K. 2008. Bacillus subtilis MinC destabilizes FtsZ-rings at new
cell poles and contributes to the timing of cell division. Genes Dev 22:3475-88.
39. Scheffers DJ. 2008. The effect of MinC on FtsZ polymerization is pH dependent and can be
counteracted by ZapA. FEBS Lett 582:2601-8.
40. de Boer PA, Crossley RE, Hand AR, Rothfield LI. 1991. The MinD protein is a membrane
ATPase required for the correct placement of the Escherichia coli division site. EMBO J
10:4371-80.
41. Marston AL, Thomaides HB, Edwards DH, Sharpe ME, Errington J. 1998. Polar localization of
the MinD protein of Bacillus subtilis and its role in selection of the mid-cell division site.
Genes Dev 12:3419-30.
42. Pulianmackal LT, Vecchiarelli AG. 2024. Positioning of cellular components by the ParA/MinD
family of ATPases. Curr Opin Microbiol 79:102485.
43. Lutkenhaus J. 2012. The ParA/MinD family puts things in their place. Trends Microbiol
20:411-8.

44.   Szardenings F, Guymer D, Gerdes K. 2011. ParA ATPases can move and position DNA and
subcellular structures. Curr Opin Microbiol 14:712-8.
45.   Szeto TH, Rowland SL, Habrukowich CL, King GF. 2003. The MinD membrane targeting
sequence is a transplantable lipid-binding helix. J Biol Chem 278:40050-6.
46.   Szeto TH, Rowland SL, Rothfield LI, King GF. 2002. Membrane localization of MinD is
mediated by a C-terminal motif that is conserved across eubacteria, archaea, and
chloroplasts. Proc Natl Acad Sci U S A 99:15693-8.
47.   Zhou H, Lutkenhaus J. 2004. The switch I and II regions of MinD are required for binding and
activating MinC. J Bacteriol 186:1546-55.
48.   Raskin DM, de Boer PA. 1999. Rapid pole-to-pole oscillation of a protein required for
directing division to the middle of Escherichia coli. Proc Natl Acad Sci U S A 96:4971-6.
49.   Hale CA, Meinhardt H, de Boer PA. 2001. Dynamic localization cycle of the cell division
regulator MinE in Escherichia coli. EMBO J 20:1563-72.
50.   Hu Z, Lutkenhaus J. 1999. Topological regulation of cell division in Escherichia coli involves
rapid pole to pole oscillation of the division inhibitor MinC under the control of MinD and
MinE. Mol Microbiol 34:82-90.
51.   Loose M, Fischer-Friedrich E, Ries J, Kruse K, Schwille P. 2008. Spatial regulators for bacterial
cell division self-organize into surface waves in vitro. Science 320:789-92.
52.   Varley AW, Stewart GC. 1992. The divIVB region of the Bacillus subtilis chromosome encodes
homologs of Escherichia coli septum placement (minCD) and cell shape (mreBCD)
determinants. J Bacteriol 174:6729-42.
53.   Levin PA, Margolis PS, Setlow P, Losick R, Sun D. 1992. Identification of Bacillus subtilis genes
for septum placement and shape determination. J Bacteriol 174:6717-28.
54.   Lee S, Price CW. 1993. The minCD locus of Bacillus subtilis lacks the minE determinant that
provides topological specificity to cell division. Mol Microbiol 7:601-10.
55.   Patrick JE, Kearns DB. 2008. MinJ (YvjD) is a topological determinant of cell division in Bacillus
subtilis. Mol Microbiol 70:1166-79.
56.   Bramkamp M, Emmins R, Weston L, Donovan C, Daniel RA, Errington J. 2008. A novel
component of the division-site selection system of Bacillus subtilis and a new mode of action
for the division inhibitor MinCD. Mol Microbiol 70:1556-69.
57.   van Baarle S, Bramkamp M. 2010. The MinCDJ system in Bacillus subtilis prevents minicell
formation by promoting divisome disassembly. PLoS One 5:e9850.
58.   Yu Y, Zhou J, Dempwolff F, Baker JD, Kearns DB, Jacobson SC. 2020. The Min System
Disassembles FtsZ Foci and Inhibits Polar Peptidoglycan Remodeling in Bacillus subtilis. mBio
11.
59.   Bach JN, Albrecht N, Bramkamp M. 2014. Imaging DivIVA dynamics using photo-convertible
and activatable fluorophores in Bacillus subtilis. Front Microbiol 5:59.
60.   Giacomelli G, Feddersen H, Peng F, Martins GB, Grafemeyer M, Meyer F, Mayer B, Graumann
PL, Bramkamp M. 2022. Subcellular Dynamics of a Conserved Bacterial Polar Scaffold Protein.
Genes (Basel) 13.
61.   Feddersen H, Wurthner L, Frey E, Bramkamp M. 2021. Dynamics of the Bacillus subtilis Min
System. mBio 12.
62.   Ishikawa K, Matsuoka S, Hara H, Matsumoto K. 2017. Septal membrane localization by C-
terminal amphipathic alpha-helices of MinD in Bacillus subtilis mutant cells lacking MinJ or
DivIVA. Genes Genet Syst 92:81-98.
63.   Ramm B, Glock P, Schwille P. 2018. In Vitro Reconstitution of Self-Organizing Protein Patterns
on Supported Lipid Bilayers. J Vis Exp doi:10.3791/58139.
64.   Hu Z, Lutkenhaus J. 2001. Topological regulation of cell division in E. coli: Spatiotemporal
oscillation of MinD requires stimulation of its ATPase by MinE and phospholipid. Mol Cell
7:1337-43.
65.   Hu Z, Lutkenhaus J. 2003. A conserved sequence at the C-terminus of MinD is required for
binding to the membrane and targeting MinC to the septum. Mol Microbiol 47:345-55.

66.   Heermann T, Ramm B, Glaser S, Schwille P. 2020. Local Self-Enhancement of MinD
Membrane Binding in Min Protein Pattern Formation. J Mol Biol 432:3191-3204.
67.   Ponting CP, Phillips C, Davies KE, Blake DJ. 1997. PDZ domains: targeting signalling molecules
to sub-membranous sites. Bioessays 19:469-79.
68.   Strahl H, Hamoen LW. 2010. Membrane potential is important for bacterial cell division. Proc
Natl Acad Sci U S A 107:12281-6.
69.   Karoui ME, Errington J. 2001. Isolation and characterization of topological specificity mutants
of minD in Bacillus subtilis. Mol Microbiol 42:1211-21.
70.   Leonard TA, Butler PJ, Lowe J. 2005. Bacterial chromosome segregation: structure and DNA
binding of the Soj dimer--a conserved biological switch. EMBO J 24:270-82.
71.   Wu W, Park KT, Holyoak T, Lutkenhaus J. 2011. Determination of the structure of the MinD-
ATP complex reveals the orientation of MinD on the membrane and the relative location of
the binding sites for MinE and MinC. Mol Microbiol 79:1515-28.
72.   Lackner LL, Raskin DM, de Boer PA. 2003. ATP-dependent interactions between Escherichia
coli Min proteins and the phospholipid membrane in vitro. J Bacteriol 185:735-49.
73.   Heermann T, Steiert F, Ramm B, Hundt N, Schwille P. 2021. Mass-sensitive particle tracking
to elucidate the membrane-associated MinDE reaction cycle. Nat Methods 18:1239-1246.
74.   Oviedo-Bocanegra LM, Hinrichs R, Rotter DAO, Dersch S, Graumann PL. 2021. Single
molecule/particle tracking analysis program SMTracker 2.0 reveals different dynamics of
proteins within the RNA degradosome complex in Bacillus subtilis. Nucleic Acids Res 49:e112.
75.   Barak LS, Webb WW. 1982. Diffusion of low density lipoprotein-receptor complex on human
fibroblasts. J Cell Biol 95:846-52.
76.   Goulian M, Simon SM. 2000. Tracking single proteins within cells. Biophys J 79:2188-98.
77.   Weimann L, Ganzinger KA, McColl J, Irvine KL, Davis SJ, Gay NJ, Bryant CE, Klenerman D.
2013. A quantitative comparison of single-dye tracking analysis tools using Monte Carlo
simulations. PLoS One 8:e64287.
78.   Landge AN, Jordan BM, Diego X, Muller P. 2020. Pattern formation mechanisms of self-
organizing reaction-diffusion systems. Dev Biol 460:2-11.
79.   St Johnston D, Nusslein-Volhard C. 1992. The origin of pattern and polarity in the Drosophila
embryo. Cell 68:201-19.
80.   Negrete J, Jr., Oates AC. 2021. Towards a physical understanding of developmental
patterning. Nat Rev Genet 22:518-531.
81.   Turing AM. 1990. The chemical basis of morphogenesis. 1953. Bull Math Biol 52:153-97;
discussion 119-52.
82.   Meinhardt H, Gierer A. 2000. Pattern formation by local self-activation and lateral inhibition.
Bioessays 22:753-60.
83.   Rowlett VW, Margolin W. 2013. The bacterial Min system. Curr Biol 23:R553-6.
84.   Ramm B, Heermann T, Schwille P. 2019. The E. coli MinCDE system in the regulation of
protein patterns and gradients. Cell Mol Life Sci 76:4245-4273.
85.   Kretschmer S, Schwille P. 2016. Pattern formation on membranes and its role in bacterial cell
division. Curr Opin Cell Biol 38:52-9.
86.   Halatek J, Brauns F, Frey E. 2018. Self-organization principles of intracellular pattern
formation. Philos Trans R Soc Lond B Biol Sci 373.
87.   Scholefield G, Whiting R, Errington J, Murray H. 2011. Spo0J regulates the oligomeric state of
Soj to trigger its switch from an activator to an inhibitor of DNA replication initiation. Mol
Microbiol 79:1089-100.
88.   Easter J, Jr., Gober JW. 2002. ParB-stimulated nucleotide exchange regulates a switch in
functionally distinct ParA activities. Mol Cell 10:427-34.
89.   Lim HC, Surovtsev IV, Beltran BG, Huang F, Bewersdorf J, Jacobs-Wagner C. 2014. Evidence
for a DNA-relay mechanism in ParABS-mediated chromosome segregation. Elife 3:e02758.

90.   Loose M, Fischer-Friedrich E, Herold C, Kruse K, Schwille P. 2011. Min protein patterns
emerge from rapid rebinding and membrane interaction of MinE. Nat Struct Mol Biol 18:577-
83.
91.   Vecchiarelli AG, Li M, Mizuuchi M, Hwang LC, Seol Y, Neuman KC, Mizuuchi K. 2016.
Membrane-bound MinDE complex acts as a toggle switch that drives Min oscillation coupled
to cytoplasmic depletion of MinD. Proc Natl Acad Sci U S A 113:E1479-88.
92.   Turkowyd B, Virant D, Endesfelder U. 2016. From single molecules to life: microscopy at the
nanoscale. Anal Bioanal Chem 408:6885-911.
93.   Yao Z, Carballido-Lopez R. 2014. Fluorescence imaging for bacterial cell biology: from
localization to dynamics, from ensembles to single molecules. Annu Rev Microbiol 68:459-76.
94.   Bohorquez L, Strahl H, Marenduzzo D, Thiele M, Bürmann F, Hamoen L. 2024. Cyclical MinD
membrane affinity differences are not necessary for MinD gradient formation in *Bacillus*
subtilis. doi:10.1101/2024.06.11.598461. bioRxiv.
95.   Miyagi A, Ramm B, Schwille P, Scheuring S. 2018. High-Speed Atomic Force Microscopy
Reveals the Inner Workings of the MinDE Protein Oscillator. Nano Lett 18:288-296.
96.   Halatek J, Frey E. 2012. Highly canalized MinD transfer and MinE sequestration explain the
origin of robust MinCDE-protein dynamics. Cell Rep 1:741-52.
97.   Ghosal D, Trambaiolo D, Amos LA, Lowe J. 2014. MinCD cell division proteins form alternating
copolymeric cytomotive filaments. Nat Commun 5:5341.
98.   Anagnostopoulos C, Spizizen J. 1961. Requirements for Transformation in *Bacillus subtilis*. J
Bacteriol 81:741-6.
99.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Zidek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-
Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M,
Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,
Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature 596:583-589.
100.   Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard
AJ, Bambrick J, Bodenstein SW, Evans DA, Hung CC, O'Neill M, Reiman D, Tunyasuvunakool K,
Wu Z, Zemgulyte A, Arvaniti E, Beattie C, Bertolli O, Bridgland A, Cherepanov A, Congreve M,
Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR,
Perlin K, Potapenko A, Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S,
Zhong ED, Zielinski M, Zidek A, Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024.
Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630:493-
500.
101.   Weiss M, Giacomelli G, Assaya MB, Grundt F, Haouz A, Peng F, Petrella S, Wehenkel AM,
Bramkamp M. 2023. The MksG nuclease is the executing part of the bacterial plasmid
defense system MksBEFG. Nucleic Acids Res 51:3288-3306.
102.   Maximova K, Reuter N, Trylska J. 2019. Peptidomimetic inhibitors targeting the membrane-
binding site of the neutrophil proteinase 3. Biochim Biophys Acta Biomembr 1861:1502-
1509.
103.   Wallner J, Lhota G, Schosserer M, Vorauer-Uhl K. 2017. An approach for liposome
immobilization using sterically stabilized micelles (SSMs) as a precursor for bio-layer
interferometry-based interaction studies. Colloids Surf B Biointerfaces 154:186-194.
104.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona
A. 2012. Fiji: an open-source platform for biological-image analysis. Nature Methods 9:676-
682.
105.   Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri KW. 2017.
ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics 18:529.
106.   Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput bacterial cell
detection and quantitative analysis. Nat Microbiol 1:16077.

107.   R Core Team. 2022. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing:Vienna, Austria. https://www.R-project.org/.
108.   Posit team. 2024. RStudio: Integrated Development Environment for R. Posit Software, PBC;
Boston, MA; http://wwwpositco/.
109.   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York:ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.
110.   Tinevez JY, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, Bednarek SY, Shorte
SL, Eliceiri KW. 2017. TrackMate: An open and extensible platform for single-particle tracking.
Methods 115:80-90.

1256

A

MinD (AlphaFold)

ASP 40

GLY 12   LYS 16

ATP binding region

Mutants
G12V: ATP binding, monomeric
K16A: No ATP binding, monomeric
D40A: No ATP hydrolysis, dimeric
I260E: No membrane binding

Wild type

ILE 260

GLU 260

Amphipathic α-helix

Membrane-binding deficient

B

Halo-MinD

Halo-MinD-G12V

Halo-MinD-K16A

Halo-MinD-D40A

Halo-MinD-I260E

A

WT G12V K16A D40A I260E

B

Comparison of track types by strain

Track Type
- static
- mobile (free)
- mobile (mixed)

Percentage [%]

Strain: WT, G12V, K16A, D40A, I260E

C

Mean-squared displacement of Halo-MinD

MSD [μm²] vs time lag [s]

- WT
- G12V
- K16A
- D40A
- I260E

D

Bubbleplot of Halo-MinD diffusive populations

Diffusion [μm²s⁻¹] vs Strain

	WT	G12V	K16A	D40A	I260E
	17.7%	25.8%	26.6%	5.61%	71.4%
	45.7%	50.5%	44.3%	40.2%	19.6%
	36.6%	23.7%	29%	54.2%	8.99%

E

WT G12V K16A D40A I260E

Probability vs Jump distance [μm]

- Confined
- Slow
- Fast