Disentangling the Roles of Distinct Cell Classes with Cell-Type Dynamical Systems

Aditi Jha
Electrical and Computer Engineering
Princeton University
aditijha@princeton.edu

Diksha Gupta
Sainsbury Wellcome Center
University College London
diksha.gupta@ucl.ac.uk

Carlos D. Brody
Princeton Neuroscience Institute
Princeton University
brody@princeton.edu

Jonathan W. Pillow
Princeton Neuroscience Institute
Princeton University
pillow@princeton.edu

Abstract

Latent dynamical systems have been widely used to characterize the dynamics of neural population activity in the brain. However, these models typically ignore the fact that the brain contains multiple cell types. This limits their ability to capture the functional roles of distinct cell classes, or to accurately predict the effects of cell-specific optogenetic perturbations on neural activity or behavior. To overcome these limitations, we introduce the "cell-type dynamical systems" (CTDS) model. This model extends latent linear dynamical systems to contain distinct latent variables for each cell class, with biologically inspired constraints on both dynamics and emissions. To illustrate our approach, we consider neural recordings with distinct excitatory (E) and inhibitory (I) populations. The CTDS model defines separate latents for E and I cells, and constrains the dynamics so that E (I) latents have a strictly positive (negative) effects on other latents. We applied CTDS to recordings from rat frontal orienting fields (FOF) and anterior dorsal striatum (ADS) during an auditory decision-making task. The model achieved higher accuracy than a standard linear dynamical system (LDS), and revealed that both E and I latents could be used to decode the animal's choice, showing that choice-related information is not restricted to a single cell class. We also performed in-silico optogenetic perturbation experiments in the FOF and ADS, and found that CTDS was able to replicate the causal effects of different perturbations on behavior, whereas a standard LDS model—which lacks the ability to capture cell-specific perturbations—did not. Crucially, our model allowed us to understand the effects of these perturbations by revealing the dynamics of different cell-specific latents. Finally, CTDS can also be used to identify cell types for neurons whose class labels are unknown in electrophysiological recordings. These results illustrate the power of the CTDS model to provide more accurate and more biologically interpretable descriptions of neural population dynamics and their relationship to behavior.

1   Introduction

Advancements in neural recording technologies have made it possible to record from hundreds of neurons simultaneously [30–32]. Understanding the dynamics of these high-dimensional populations and their relationship to complex behavior is a fundamental goal of computational neuroscience. Dynamical systems have proven to be extremely useful in this pursuit [18, 3, 10, 11]. Latent dynamical

systems model neural activity as arising from a low-dimensional latent state, and describe how the activity evolves as a function of time in this low-dimensional space.

However, while latent dynamical systems provide parsimonious descriptions of neural activity, they fail to capture key functional and biological properties of real neural circuits. In particular, they ignore the fact that neural circuits consist of multiple cell types. Standard models describe the activity of all neurons as arising from the same set of latent variables, without any differences among cell types or constraints on the interactions between them. This prevents latent dynamical systems from shedding light on the functional roles of distinct cell classes [25, 34, 26, 24]. Furthermore, it prevents them from accurately describing optogenetic perturbation experiments, in which specific cell classes within a neural circuit are perturbed. This severely limits the usefulness of classic latent dynamical systems models for identifying the causal effects of neural activity on behavior [14, 22].

To overcome these limitations, we introduce the "cell-type dynamical systems" (CTDS) model. This model is an extension of latent linear dynamical system (LDS) models in which a distinct set of latent variables governs the activity of each cell class, with constraints on the model parameters governing the sign of the interactions between different cell types. Here we focus on networks with two distinct cell types: excitatory (E) and inhibitory (I) neurons. Standard analyses would characterize the dominant modes of network activity without considering the distinct roles played by E and I cells. By contrast, the CTDS model defines separate latents for E and I cells, and constrains the dynamics matrix so that E and I latents have a positive or negative (respectively) effect on the other latents. Our model also easily extends to multi-region settings, and allows us to explicitly constrain the types of cells (e.g., E cells) making long range projections.

We first demonstrate an equivalence between CTDS models with E and I cells and low-rank recurrent neural network models composed of E and I neurons (EI-RNN), both theoretically and using simulated experiments. EI-RNNs [12, 29, 27] are widely studied as proxies of neural circuits, thus this equivalence validates the utility of our model. Next, we apply CTDS to recordings from rat frontal orienting fields (FOF) and anterior dorsal striatum (ADS) during an auditory decision-making task [4]. We show that CTDS predicts neural activity better on held out trials than a standard LDS model; a multi-region CTDS with distinct latents for FOF and ADS furthermore outperforms CTDS with no regional constraints. We show that the latents extracted from both E and I populations encode the animal's choice, revealing that choice information is not restricted to a single cell class, consistent with recent findings [21]. Additionally, we show that a classifier trained on the inferred latent states during training trials predicted the animal's behavioral choice well during test trials.

Next, we use CTDS to perform in-silico optogenetic perturbation experiments modeled after those performed in vivo. In these experiments, one class of neurons (E or I) in a single brain region is perturbed during a portion of each trial. Remarkably, we find that the CTDS model—despite not being trained on perturbation data—can accurately predict the effects of different perturbations on behavior. Moreover, CTDS allows us to visualize the effects of these perturbations on the underlying dynamics of different regions, allowing us to obtain accurate roles of distinct regions and cell-types. A standard LDS, however, is unable to replicate experimental findings during the same set of perturbations due to the lack of cell information and appropriate structural constraints. Finally, we show that the CTDS model can be used to identify the cell class of neurons whose type is unknown in experimental recordings. While in previous analyses we assumed that the identity of neurons were known, cell-type information is often not known in neurophysiology experiments. We developed an approach to infer the indentities of unknown cells using CTDS, and find that CTDS is indeed able to infer cell-types of upto 50% of unidentified neurons using recordings from the FOF and ADS. This an exciting application of our model as tagging cell-types can be challenging during the course of real-world experiments. Overall, our finding underscores that CTDS provides a versatile tool to study the effects of optogenetic perturbations on neural dynamics, and to infer cell identities.

2   Cell-type latent linear dynamical systems (CTDS)

2.1   Background: LDS models

A latent linear dynamical system (LDS) model describes the firing activity of a population of N neurons at time t, $\mathbf{y}_t \in \mathbb{R}^N$, as arising from a low-dimensional latent state $\mathbf{x}_t \in \mathbb{R}^D$. This state

Figure 1: **A.** Graphical model of a Cell-Type latent Dynamical System (CTDS). Red and blue circles represent the observed activity, $\mathbf{y}_t$, of excitatory and inhibitory neurons respectively at time t. The color of the latents $\mathbf{x}_t$ reflects the cell-type they govern. $\mathbf{u}_t$ is any input to the system at the time t. **B.** Latent dynamics, A, obeys Dale's law, such that the columns of A are either non-negative or non-positive corresponding to E and I latents. **C.** Distinct sets of latents govern the activity of E and I neurons. The emission matrix, C, is also constrained to be non-negative such that the Dale's law constraint in the latent space is reflected to the neural activity space.

evolves according to linear dynamics and any under the influence of any external input $\mathbf{u}_t \in \mathbb{R}^M$:

$$\mathbf{x}_{t+1} = A\mathbf{x}_t + B\mathbf{u}_t + \mathbf{w}_t; \ \mathbf{w}_t \in \mathcal{N}(\mathbf{0}, Q) \tag{1}$$

Here, $A \in \mathbb{R}^{D \times D}$ captures dynamics in the latent space, $B \in \mathbb{R}^{D \times M}$ captures the influence of the external input, while $Q \in \mathbb{R}^{D \times D}$ is the noise covariance. The observed activity is then a linear transformation of this latent state such that

$$\mathbf{y}_t = C\mathbf{x}_t + \mathbf{v}_t, \ \mathbf{v}_t \in \mathcal{N}(\mathbf{0}, R) \tag{2}$$

Here, $C \in \mathbb{R}^{N \times D}$ is called the emission matrix, and $R \in \mathbb{R}^{N \times N}$ is the observation noise covariance.

2.2 The CTDS model

To disentangle the roles of distinct cell types in a population, we modify the standard LDS model in two ways: (1) we define a distinct set of latent variables for the activity of each cell class; and (2) we constrain the dynamics to obey functional properties of these cell types. We refer to the resulting models as Cell-Type latent linear Dynamical Systems (CTDS). Here we focus on the setting of excitatory (E) and inhibitory (I) cell types, as schematized in Fig. 1. We use distinct sets of latents for E and I neurons, $\mathbf{x}_t^e \in \mathbb{R}^{D_e}$ and $\mathbf{x}_t^i \in \mathbb{R}^{D_i}$ respectively, such that $D_e + D_i = D$ (shown in Fig. 1C). As a result, the emission matrix C is block-diagonal with blocks of shape $N_e \times D_e$ and $N_i \times D_i$, where the first block maps the E latents to the N_e excitatory neurons, and the second block maps the I latents to the N_i inhibitory neurons. Note that we constrain these blocks to be non-negative, thus ensuring that an increase or decrease in a given latent maps to an increased or decreased firing rate (respectively) in neurons of the associated cell type.

Next, to ensure consistency with Dale's law—E neurons excite and I neurons inhibit—we constrain the sign of the columns of the dynamics matrix A so that columns associated with E latents are positive and columns associated with I latents are negative. (Fig. 1B). (Note we do not apply this constraint to the diagonal elements of A; this allows individual latents to have positive auto-correlation and evolve smoothly in time, regardless of cell type).

2.3 Multi-region CTDS

Given simultaneously recorded neural activity from multiple brain regions, we often want to understand the computations performed by each region as well as the interactions between regions.

To do so, following existing work on multi-region modeling [11], we extend our model to multiple regions such that each region has its own set of latents. For example, if we have K regions such that $\mathbf{x}_t^k \in \mathbb{R}^{D_k}$ is the latent vector for region k, this latent state evolves as follows:

$$\mathbf{x}_t^k = A_{kk}\mathbf{x}_{t-1}^k + \sum_{j \neq k} A_{jk}\mathbf{x}_{t-1}^j + B\mathbf{u}_t + \epsilon_t, \tag{3}$$

where $A_{kk} \in \mathbb{R}^{D_k \times D_k}$ captures the within-region dynamics of the kth region, while $A_{jk} \in \mathbb{R}^{D_j \times D_k}$ models the inter-region communications from region j to region k. Since we are interested in cell-type specific information, we further assume separate latents for distinct cell types within a region such that $\mathbf{x}_t^k = [\mathbf{x}_t^{k,e}, \mathbf{x}_t^{k,i}]$. The within-region dynamics A_{kk} obeys Dale's law, similar to the single-region model. Furthermore, since long range connections in the cortex are generally known to be excitatory [28], we can also constrain the inter-region communication matrix A_{jk} to have non-negative columns corresponding to the E latents of region j, and zero for the I latents. Finally, to ensure that cell-type specific and region-specific latents only control their respective neurons, the emission matrix $C \in \mathbb{R}^{N \times KD}$ is non-negative and block diagonal. Mathematically, we can write the activity of E and I cell classes in region k as follows:

$$\mathbf{y}_t^{k,e} = C^{k,e}\mathbf{x}_t^{k,e} + \mathbf{v}_t^k, \ \mathbf{v}_t^k \in \mathcal{N}(\mathbf{0}, R^k) \tag{4}$$

$$\mathbf{y}_t^{k,i} = C^{k,i}\mathbf{x}_t^{k,i} + \mathbf{v}_t^{k,i}, \ \mathbf{v}_t^k \in \mathcal{N}(\mathbf{0}, R^k). \tag{5}$$

2.4 Inference procedure

In order to infer the parameters of a CTDS model, we maximize the expected log-likelihood of observed data under the model using the Expectation Maximization (EM) algorithm [5]. We will describe inference for the single-region model for simplicity, which can easily be extended to the multi-region variant. We want to infer the model parameters, $\Theta = \{A, B, C, Q, R\}$, given N trials of observed data $\{\mathbf{y}_{1:T_n}^n\}_{n=1}^N$ where n represents the trial index.

In the Expectation step, we perform standard Kalman filtering and smoothing, to learn posterior distributions of the latent states $P(\mathbf{x}_t^n \mid \mathbf{y}^n, \Theta)$, and $P(\mathbf{x}_t^n, \mathbf{x}_{t+1}^n \mid \mathbf{y}_{1:T_n}^n, \Theta) \ \forall t \in \{1, T_n\}$. In the Maximization step, we learn the model parameters given the computed expectations of the latent states. Specifically, we optimize the following expression

$$\mathcal{L}_{CD}(\Theta) = -\frac{1}{2}\sum_{n=1}^N \left(\mathbb{E}\Big[\sum_{t=1}^{T_n-1} \log \mathcal{N}(\mathbf{x}_{t+1}^n; A\mathbf{x}_t^n + B\mathbf{u}_t^n, Q) + \sum_{t=1}^{T_n} \log \mathcal{N}(\mathbf{y}_t^n; C\mathbf{x}_t^n, R)\Big]\right) \tag{6}$$

for $\{A, B, C, Q, R\}$, such that A and C obey their respective structural constraints. As a result, unlike a standard LDS, we no more have closed form updates for the model parameters. We instead solve two quadratic program under the constraints on A and C separately:

$$\max_{A,B} -\frac{1}{2}\sum_{n=1}^N \left(\mathbb{E}\Big[\sum_{t=1}^{T_n-1} \log \mathcal{N}(\mathbf{x}_{t+1}^n; A\mathbf{x}_t^n + B\mathbf{u}_t^n, Q)\Big]\right) \ \text{s.t. } A \text{ obeys Dale's law} \tag{7}$$

$$\max_{C} -\frac{1}{2}\sum_{n=1}^N \left(\mathbb{E}\Big[\sum_{t=1}^{T_n} \log \mathcal{N}(\mathbf{y}_t^n; C\mathbf{x}_t^n, R)\Big]\right) \ \text{s.t. } C >= 0 \text{ and block-diagonal} \tag{8}$$

We alternate between optimizing A, B, C and the noise matrices $\{Q, R\}$, which have closed form expressions once $\{A, B, C\}$ are fixed. Overall, we alternate between the expectation and maximization steps until the log-likelihood converges. We use the MOSEK solver [2] with CVXPY [6, 1] to perform the quadratic optimizations for the constrained matrices $\{A, C\}$.

3 Relationship to E-I recurrent neural networks

Recurrent neural networks (RNNs) are widely used to study neural dynamics in the brain [19, 23, 7, 12, 29]. An important line of work has focused on RNNs with excitatory and inhibitory units, which mirror the functional organization of real neural circuits [9, 12, 22, 29, 27]. However, such models are often hard to analyze or interpret. Here we show that CTDS provides a natural bridge between EI-RNNs and latent dynamical systems, in which neural activity is explained in terms of

a low-dimensional latent variable. Our work thus also provides an explicit connection to low-rank RNNs, which have been shown to account for a wide variety of tasks and datasets [20, 7, 33], but which have not to our knowledge been equipped with distinct cell types. Specifically, we show that under certain conditions a low-rank EI-RNN is formally equivalent to a CTDS, and we derive the mapping from one model class to the other.

Let $\mathbf{y}_t \in \mathbb{R}^N$ be the activity generated by a linear RNN containing N neurons, such that:

$$\mathbf{y}_{t+1} = J\mathbf{y}_t + \eta_t, \quad \eta_t \sim \mathcal{N}(0, P), \tag{9}$$

where $J \in \mathbb{R}^{N \times N}$ obeys Dale's law, with non-negative columns for E cells and non-positive columns for I cells, and $P \in \mathbb{R}^{N \times N}$ is the noise covariance. Let's further assume N_e to be the number of E cells, and N_i as the number of I cells. For simplicity, we will focus here on the noiseless case, $P = \mathbf{0}$; we discuss the noisy case in the supplement.

We can further write the activity of the E and I cells separately as follows:

$$\mathbf{y}_{t+1}^e = J_{ee}\mathbf{y}_t^e + J_{ei}\mathbf{y}_t^i, \quad \mathbf{y}_{t+1}^i = J_{ii}\mathbf{y}_t^i + J_{ie}\mathbf{y}_t^e \tag{10}$$

where J_{ee} and J_{ie} represent blocks of J that contain outgoing weights from E cells and are thus non-negative, J_{ii} and J_{ei} are blocks of outgoing weights from I cells and contain non-positive elements. Let's define a new matrix J^+ which contains the absolute values of J such that:

$$J^+ = \begin{bmatrix} J_{ee} & J_{ie}^+ \\ J_{ei} & J_{ii}^+ \end{bmatrix} \tag{11}$$

Here J_{ii}^+ and J_{ie}^+ contain absolute values of the all-negative matrices J_{ii} and J_{ie} respectively. Mapping this RNN to a CTDS with E and I cells requires that the top and bottom sub-matrices of J^+ have low-rank non-negative matrix factorization (NNMF) solutions:

$$\begin{bmatrix} J_{ee} & J_{ie}^+ \end{bmatrix} = U_1 V_1^\top; \quad \begin{bmatrix} J_{ei} & J_{ii}^+ \end{bmatrix} = U_2 V_2^\top \tag{12}$$

where $U_1 \in \mathbb{R}^{N_e \times K_1}$, $V_1 \in \mathbb{R}^{N \times K_1}$ and $U_2 \in \mathbb{R}^{N_i \times K_2}$, $V_2 \in \mathbb{R}^{N \times K_2}$. Thus, J^+ can be written as:

$$J^+ = UV^\top = \begin{bmatrix} U_1 & \mathbf{0} \\ \mathbf{0} & U_2 \end{bmatrix} \begin{bmatrix} V_1^\top \\ V_2^\top \end{bmatrix} \tag{13}$$

where $U, V \in \mathbb{R}^{N \times (K_1 + K_2)}$ are non-negative matrices. If we make the rows of V corresponding to I cells negative, and call this new matrix V_{dale}, we obtain $J = UV_{\text{dale}}^\top$. This EI-RNN can then be equivalently be written as a CTDS of rank at most $K_1 + K_2$ as follows [33]:

$$\mathbf{x}_t = V_{\text{dale}}^\top U \mathbf{x}_{t-1} \tag{14}$$

$$\mathbf{y}_t = U \mathbf{x}_t, \tag{15}$$

with the emissions $C = U$, and latent dynamics $A = V_{\text{dale}}^\top U$. Both C and A obey the required constraints: C is non-negative and block diagonal, and A obeys Dale's law. Hence, in summary, a linear EI-RNN can be mapped to a CTDS if the excitatory and inhibitory rows of the absolute connectivity matrix, J^+, have low-rank NNMF solutions.

In the case of non-zero noise in the EI-RNN, it is still possible to find a mapping to a cell-type LDS model, however that requires further constraints on the connectivity and noise structure of the RNN (see sec. A1). We also generated activity from E-I networks whose connectivity J obeys the above constraints, and show that CTDS models recover the connectivity accurately, and with smaller error than a standard LDS using the same amount of data (see sec. A2). Finally, when fitting CTDS models to real data, we use insights from this equivalence to initialize them. In particular, we first learn a J matrix given data $\{\mathbf{y}_{1:T}\}$ that obeys Dale's law by solving the following constrained regression:

$$\mathbf{y}_{t+1} = J\mathbf{y}_t + \mathbf{v}_t; \quad \text{s.t } J \text{ obeys Dale's law} \tag{16}$$

We apply NNMF as discussed earlier on the absolute values of the learned J per eq. 13, and thus initialize $C = U$, $A = V_{\text{dale}}^\top U$.

4 Application to rodent decision-making data

Next, we applied the CTDS model to neural data collected from rats trained to perform an auditory decision-making task [4]. On each trial, animals heard randomly timed clicks from the left and the right, and were rewarded for selecting the side with more clicks (Fig. 2A). Neuropixel probes were used to record simultaneous neural activity from two regions known to be involved in evidence accumulation: the frontal orienting fields (FOF) and the anterior dorsal striatum (ADS) [13, 35]. Previous work has shown that individual neurons in FOF and ADS exhibit side selectivity in this task, with activity ramping upwards for stimuli on the cell's preferred side [13, 35]. However, to carefully understand the causal role of each region during evidence accumulation (e.g., whether regions carry out different aspects of the task), it is important to disentangle the roles of different cell types in each region. We point out that prior work has analyzed the two regions independently, while here we also study communication between them.

4.1 Model fitting details

We fitted a standard LDS and the CTDS model to zero-centered firing rates of 109 active neurons recorded in FOF and ADS regions of a single rat. These neurons were filtered to have a minimum firing rate of 1Hz during the (~1s) stimulus period. We used 50ms bins to obtain firing rates from the spiking activity of neurons, and zero-centered the response of each neuron across trials. We had access to 353 trials for this animal. To fit CTDS models, we labeled neurons as E or I using both anatomical information about the two regions as well a clustering of spike width histograms (see sec. A3). Additionally, since we had data from two regions, we also used region identity to fit multi-region CTDS models. Since ADS is a sub-cortical region, we did not restrict across-region communication between ADS and FOF to be only excitatory. However, we did restrict communication from FOF to ADS to be excitatory (as discussed in subsec. 2.3).

We used 80% of 353 trials to fit our models, the remaining 20% trials were held out. The models also received a 2-dimensional input at every time point, containing the number of left and right clicks played between the previous and current time points. We initialized CTDS models using the NNMF initialization process described in sec. 3. For all models, we used 10 initialization seeds and picked the best seed based on training log-likelihood. Finally, we varied the number of dimensions in the latent space of the models, and for simplicity used the same number of latents for each cell type.

4.2 Both cell types encode choice information

We found that CTDS and its multi-region variant outperformed LDS models in log-likelihood of held-out trials. Fig. 2B shows test log-likelihood as a function of the number of latents available to each cell-type. This result confirms that adding E-I structure to latent linear dynamical systems allows the model to capture neural data well, while also providing additional interpretability. For each model class, we also trained a logistic regression classifier to predict the animal's choice from the last time-step of the latent variable. Fig. 2C shows the choice prediction accuracy of the classifier corresponding to each model on test trials (for $D = 4$ per population). We observe that all models are able to predict choice well (and far above chance, i.e. 50%; error bars show one standard deviation across 10 different sampled latent states).

Next, we analyzed a fitted multi-region CTDS model to draw scientific insights about the roles of the two regions and cell types. We chose the model with 4 dimensions per cell-type for interpretability, resulting in overall 16 latents across the two regions and the two cell-types. In Fig. 2D, we show the recovered dynamics matrix—our results reveal cross-region communications between FOF and ADS in both directions suggesting that they are recurrently connected during evidence accumulation tasks. We also plotted the magnitude of the eigenvalues corresponding to the within-region dynamics in both regions, in Fig. 2, and found that both regions have distinct eigenvalues suggesting distinct within-region dynamics.

Finally, we plotted the top two PCs of the inferred latent states for FOF and ADS in Fig. 2E, with left and right choice trials colored distinctly. We found that both E and I latents in the FOF encoded the animal's choice, with their trajectories separating out for left and right choices in opposite directions. This is consistent with recent work from Najafi et al. [21], which showed that both excitatory and

Figure 2: CTDS applied to decision-making task in rodents. **A**. "Poisson clicks" auditory decision-making task [4] **B**. Log-likelihood on held out trials as a function of the number of latents per cell-type, for three different model classes. LDS (gray), CTDS with no region structure (dashed black), multi-region CTDS (solid black). **C**. Choice accuracy (between 0–1) on test trials using the three models with four latents per cell-type. Dotted line represents chance performance. **D**. (Left) Recovered dynamics matrix, A, using a multi-region CTDS with 4 latents per population. (Right) Real parts of eigenvalues of the top left and bottom right blocks of A, corresponding to within-region dynamics in FOF and ADS, respectively. **E**. Latent state trajectories plotted along their top two PCs colored distinctly for left (coral) and right (teal) choice trials. (Left) E and I latents in FOF, and (right) I latents in ADS.

inhibitory neurons are equally selective to choice. Thus, CTDS enabled us to distentangle and understand the dynamics underlying distinct classes of cells and regions.

4.3 In-silico optogenetic perturbation experiments

By dissociating dynamics underlying each cell-type, CTDS models allows us to study cell-specific optogenetic perturbations. These experiments involve activating or silencing a targeted class of neurons in the brain to causally establish links between neural circuits and observed behavior. We can perform in-silico perturbation experiments in CTDS models by silencing or exciting latents corresponding to a particular cell class, and thus study the effects on both neural population dynamics and behavior of the animal.

Previous work by Hanks et al. [13] has shown that during the clicks task, silencing excitatory neurons in FOF during the the first half of the trial does not affect the animal's behavior. However inactivation during the second half produces biased choices and reduced task accuracy. Furthermore, Yartsev et al. [35] have shown that inactivating inhibitory ADS neurons results in behavior deficits during both early and late half of the trial.

To investigate the circuit-level origin of these effects, we performed in-silico perturbations using our fitted multi-region CTDS model from Fig. 2 ($D = 4$ latents per cell-type). Note that the model was fit using unperturbed data, and never saw perturbed trials during training. We provided the model with inputs from the held-out trials, and generated data while clamping cell-specific latents in either region to a negative value during the first or second half of the trial (Fig. 3A & B). This effectively clamped the activity of the associated cell class (we chose the inactivation strength of cell-specific latents via a parameter search, but our results are qualitatively robust; see Sec. A4). We then predicted choice on these trials using the classifier previously trained on unperturbed training data.

Remarkably, we found that the perturbed CTDS model produced the same pattern of deficits in FOF as observed in previous work [13](Fig. 3C, solid bars). Early-half perturbations produced low choice error rates in both the CTDS model and in real animal experiments. On the contrary, late-half perturbations resulted in biased animal behavior, and in high error rates in the CTDS model. We also

Figure 3: In-silico perturbation experiments. **A**. (*Left*) Latent trajectories of FOF during a sample trial when unperturbed, (*center*) perturbation of E latents in early half of the same trial (using same sets of inputs), (*right*) perturbed latents during later half of the trial. Red and blue represent E and I latents, respectively. **B**. Latent trajectories in ADS under no perturbation, early half perturbation, later half perturbation during an example trial (*from left to right*). **C**. Inactivating E cells in FOF. (*Right*) Experimental data from [13] showing that behavioral deficit (measured through ipsilateral bias) is only observed when inactivating FOF during the later half of the trial. (*Left*) In-silico perturbation experiments when inactivating E latents from FOF in a fitted CTDS replicate these findings. Gray represents error (0–1) between predicted and true choices during unperturbed trials. Yellow shows classifier's error for perturbed trials, inactivations were performing during the first half of trials. Green represents the same for inactivations during later half of trials. Hatched bars show the same for a standard multi-region LDS. **D**. Inactivating I cells in ADS. (*Right*) Experimental data from [35] showing behavioral deficits when inactivating ADS during both start and later half of trials. (*Left*) In-silico perturbation experiments when inactivating I latents from ADS in a fitted CTDS replicate these findings. Gray, yellow and green bars correspond to no perturbation, early stage and late stage perturbation respectively. Solid and hatched bars represent CTDS and standard LDS respectively.

performed in-silico perturbations in the ADS by inactivating I latents during either early half or later half of the trial. Excitingly, we were again able to replicate experimental findings (Fig. 3D, solid bars). Choice error was high when inactivating ADS during both early half and late half, suggesting that animal behavior changes due to perturbations during both halves of a trial.

These results show that CTDS is able to capture the findings of optogenetic inactivation experiments. This is exciting because the model was never trained on perturbed trials. Additionally, CTDS allows us to visualize the changes in underlying dynamics to understand the differential effects of inactivations in FOF and ADS. We visualized the unperturbed latent states for an example trial and compared them with perturbed latent states during early and later half of the trial. We see in Fig. 3A that FOF latents recovered quickly after inactivations during early half of the trial (presumably using information from ADS, see sec. A4), and as a result we did not observe behavior deficits in this case. However, silencing FOF during later half of the trial caused the latent states of FOF to shrink, resulting in the observed behavioral deficits. However, when perturbing ADS (Fig. 3B), the latent dynamics changed significantly during both early and late stage inactivations explaining the behavioral deficits observed

A cell-type identification

B accuracy of cell-types

Figure 4: Cell-type identification. (**A**) Schematic of unknown cell type problem. (**B**) Accuracy of cell-type inference as a function of the number of masked neurons. Error bars show one standard error over 10 initialization seeds. Dotted line represents random performance.

during inactivation in ADS. Thus, these results suggest that ADS is crucially involved during the task throughout, and perturbing this region prevents the animal from performing the task accurately.

For comparison, we also trained a multi-region LDS model on the same dataset, again with $D = 4$ latents per region. This model had no explicit representation of cell-types, but it did contain distinct latents for each brain region. We attempted to replicate the perturbation experiments in this model by clamping two of the four latents in each region to a negative value (just as we did with CTDS, although in this case the latents had no explicit assignment to distinct cell types). As before, we then used the perturbed model activity to predict choice using a classifier trained on unperturbed trials. This model accurately captured behavior on unperturbed trials (Fig. 3 C & D, hatched gray bar shows error between model predictions and animal behavior). However, it notably failed to capture the pattern of deficits observed in perturbed trials (Fig. 3 C & D, hatched yellow and green bars). In particular, perturbations in either region, whether early or late, produced behavior similar to that when no perturbations were present. This shows that standard LDS did not accurately capture the functional contributions of different cell-types to neural population dynamics (see sec. A5). These results underscore the importance of cell-type models for understanding perturbation experiments, and for uncovering the contributions of different cell types to population dynamics.

5 Inferring cell-type information of unknown neurons

Classifying neurons into cell-types facilitates scientific discovery, provides a nuanced view of neural circuits engaged in tasks, and is also crucial for analyses of diseases [36]. However, identifying cell-types in challenging. For example, spike width is commonly used to classify putative excitatory or inhibitory neurons, but many neurons exhibit intermediate spike widths, and thus cannot be confidently assigned to one class or the other. Here we describe a method for using the CTDS model to infer cell-types from neural activity for a subset of neurons in a recording.

Our proposed method involves first fitting the CTDS model using only neurons with known cell type. Then, we run an altered version of the M-step, in which we select for each neuron which set of latents best describe its activity (e.g., E latents or I latents). Specifically, for an unknown k-th neuron, assuming diagonal observation noise R, we solve the following two regression problems:

$$\min_{C^k} \sum_{n=1}^{N} \mathbb{E}\big[\mathbf{y}_t^{n,k} - C^k \mathbf{x}_t^e - \mathbf{v}_t^k\big], \quad \min_{C^k} \sum_{n=1}^{N} \mathbb{E}\big[\mathbf{y}_t^{n,k} - C^k \mathbf{x}_t^i - \mathbf{v}_t^k\big] \ \text{ s.t } C^k >= 0 \qquad (17)$$

Here $C^k \in \mathbb{R}^D$, where D is the latent dimensionality of each cell-type. This step was repeated within each M step during fitting for each unknown neuron. We then picked the C^k that resulted in minimum regression error in the end, thus obtaining the identity of the neuron.

To test our method, we masked between $5 - 50$ neurons (equal number of E and I cells) from the FOF-ADS data, and inferred their identities while fitting a multi-region CTDS ($D = 4$ latents per cell-type). Fig. 4 shows that we were indeed able to infer cell identities well above chance, thus demonstrating the usefulness of CTDS for inferring of cell identities.

6 Conclusions and discussion

We have developed a novel framework for disentangling the roles of distinct cell-types in neural circuits. In particular, we extended linear dynamical systems to incorporate cell-type specific information. We focused on excitatory and inhibitory cell classes, with latents constrained to interact in accordance with Dale's law.

We have also derived a theoretical equivalence between linear RNNs with an E-I structure and our model. Cell-type dynamical systems thus bridge a gap between interpretable state-space models and mechanistic models of recurrent computation. We extended CTDS to incorporate multiple regions, resulting in multi-region CTDS models. Application of our model to decision-making data from rodents revealed that including E-I structure in latent linear dynamical systems improved their ability to capture neural activity, and allowed us to understand roles of distinct cell classes, In particular, in line with Najafi et al. [21], we found that both E and I neurons encode choice information.

Crucially, CTDS allowed us to replicate optogenetic perturbation experiments in the FOF and ADS [13, 35], due to the separation of latents based on cell types. Our model predicted the same changes in behavior as observed in experimental studies, and also allowed us to visualize and understand the effects of these perturbation on the dynamics of different cell-typess. This is particularly exciting as a standard LDS was unable to capture the same causal effects. We also developed an approach using CTDS to infer the cell-type information of neurons using their activity. While we assumed linear dynamics in our current setup, the core idea of separating latents into cell types is applicable to non-linear systems as well [15, 16]. Overall, we believe that CTDS can be broadly useful for several applications, and can serve as a helpful tool to obtain a nuanced view of the dynamics underlying different populations of neurons in the brain.

References

[1] A. Agrawal, R. Verschueren, S. Diamond, and S. Boyd. A rewriting system for convex optimization problems. *Journal of Control and Decision*, 5(1):42–60, 2018.

[2] M. ApS. *MOSEK version 10.1*, 2024. URL https://www.mosek.com.

[3] E. W. Archer, U. Koster, J. W. Pillow, and J. H. Macke. Low-dimensional models of neural population activity in sensory cortical circuits. In *Advances in Neural Information Processing Systems*, volume 27. Curran Associates, Inc., 2014. URL https://proceedings.neurips.cc/paper/2014/hash/cb70ab375662576bd1ac5aaf16b3fca4-Abstract.html.

[4] B. W. Brunton, M. M. Botvinick, and C. D. Brody. Rats and humans can optimally accumulate evidence for decision-making. *Science (New York, N.Y.)*, 340(6128):95–98, Apr. 2013. ISSN 1095-9203. doi: 10.1126/science.1233912.

[5] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1): 1–22, Sept. 1977. ISSN 0035-9246, 2517-6161. doi: 10.1111/j.2517-6161.1977.tb01600.x.

[6] S. Diamond and S. Boyd. CVXPY: A Python-embedded modeling language for convex optimization. *Journal of Machine Learning Research*, 17(83):1–5, 2016.

[7] A. Dubreuil, A. Valente, M. Beiran, F. Mastrogiuseppe, and S. Ostojic. The role of population structure in computations through neural dynamics. *Nature Neuroscience*, 25(6):783–794, June 2022. ISSN 1546-1726. doi: 10.1038/s41593-022-01088-4.

[8] J. T. Dudman and C. R. Gerfen. *Chapter 17 - The Basal Ganglia*, page 391–440. Academic Press, San Diego, Jan. 2015. ISBN 978-0-12-374245-2. doi: 10.1016/B978-0-12-374245-2.00017-6. URL https://www.sciencedirect.com/science/article/pii/B9780123742452000176.

[9] D. Fisher, I. Olasagasti, D. W. Tank, E. R. F. Aksay, and M. S. Goldman. A modeling framework for deriving the structural and functional architecture of a short-term memory microcircuit. *Neuron*, 79(5):987–1000, Sept. 2013. ISSN 0896-6273. doi: 10.1016/j.neuron.2013.06.041.

[10] Y. Gao, E. W. Archer, L. Paninski, and J. P. Cunningham. Linear dynamical neural population models through nonlinear embeddings. In *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016. URL https://proceedings.neurips.cc/paper/2016/hash/76dc611d6ebaafc66cc0879c71b5db5c-Abstract.html.

[11] J. Glaser, M. Whiteway, J. P. Cunningham, L. Paninski, and S. Linderman. Recurrent switching dynamical systems models for multiple interacting neural populations. In *Advances in Neural Information Processing Systems*, volume 33, page 14867–14878. Curran

Associates, Inc., 2020. URL `https://proceedings.neurips.cc/paper/2020/hash/aa1f5f73327ba40d47ebce155e785aaf-Abstract.html`.

[12] A. Haber and E. Schneidman. The computational and learning benefits of daleian neural networks. Oct. 2022. URL `https://openreview.net/forum?id=ckQvYXizgd1`.

[13] T. D. Hanks, C. D. Kopec, B. W. Brunton, C. A. Duan, J. C. Erlich, and C. D. Brody. Distinct relationships of parietal and prefrontal cortices to evidence accumulation. *Nature*, 520 (75467546):220–223, Apr. 2015. ISSN 1476-4687. doi: 10.1038/nature14066.

[14] N. Li, K. Daie, K. Svoboda, and S. Druckmann. Robust neuronal dynamics in premotor cortex during motor planning. *Nature*, 532(76007600):459–464, Apr 2016. ISSN 1476-4687. doi: 10.1038/nature17643.

[15] S. Linderman, M. Johnson, A. Miller, R. Adams, D. Blei, and L. Paninski. Bayesian Learning and Inference in Recurrent Switching Linear Dynamical Systems. In A. Singh and J. Zhu, editors, *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*, pages 914–922. PMLR, 20–22 Apr 2017. URL `https://proceedings.mlr.press/v54/linderman17a.html`.

[16] S. Linderman, A. Nichols, D. Blei, M. Zimmer, and L. Paninski. Hierarchical recurrent state space models reveal discrete and continuous dynamics of neural activity in c. elegans. page 621540, Apr. 2019. doi: 10.1101/621540. URL `https://www.biorxiv.org/content/10.1101/621540v1`.

[17] S. Linderman, B. Antin, and D. Zoltowski. SSM: Bayesian learning and inference for state space models, 2020. URL `http://github.com/slinderman/ssm`.

[18] J. H. Macke, L. Buesing, J. P. Cunningham, B. M. Yu, K. V. Shenoy, and M. Sahani. Empirical models of spiking in neural populations. In *Advances in Neural Information Processing Systems*, volume 24. Curran Associates, Inc., 2011. URL `https://papers.nips.cc/paper_files/paper/2011/hash/7143d7fbadfa4693b9eec507d9d37443-Abstract.html`.

[19] V. Mante, D. Sussillo, K. V. Shenoy, and W. T. Newsome. Context-dependent computation by recurrent dynamics in prefrontal cortex. *Nature*, 503(7474):78–84, Nov. 2013. ISSN 1476-4687. doi: 10.1038/nature12742.

[20] F. Mastrogiuseppe and S. Ostojic. Linking connectivity, dynamics, and computations in low-rank recurrent neural networks. *Neuron*, 99(3):609–623.e29, Aug. 2018. ISSN 08966273. doi: 10.1016/j.neuron.2018.07.003.

[21] S. F. Najafi, G. F. Elsayed, R. Cao, E. Pnevmatikakis, P. E. Latham, J. P. Cunningham, and A. K. Churchland. Excitatory and inhibitory subnetworks are equally selective during decision-making and emerge simultaneously during learning. *Neuron*, 105(1):165–179.e8, Jan 2020. ISSN 0896-6273. doi: 10.1016/j.neuron.2019.09.045.

[22] D. J. O'Shea, L. Duncker, W. Goo, X. Sun, S. Vyas, E. M. Trautmann, I. Diester, C. Ramakrishnan, K. Deisseroth, M. Sahani, and K. V. Shenoy. Direct neural perturbations reveal a dynamical mechanism for robust computation. page 2022.12.16.520768, Dec 2022.

[23] C. Pandarinath, D. J. O'Shea, J. Collins, R. Jozefowicz, S. D. Stavisky, J. C. Kao, E. M. Trautmann, M. T. Kaufman, S. I. Ryu, L. R. Hochberg, J. M. Henderson, K. V. Shenoy, L. F. Abbott, and D. Sussillo. Inferring single-trial neural population dynamics using sequential auto-encoders. *Nature Methods*, 15(10):805–815, Oct. 2018. ISSN 1548-7105. doi: 10.1038/s41592-018-0109-9.

[24] W. F. Podlaski and C. K. Machens. Approximating nonlinear functions with latent boundaries in low-rank excitatory-inhibitory spiking networks. *Neural Computation*, 36(5):803–857, Apr. 2024. ISSN 0899-7667. doi: 10.1162/neco_a_01658.

[25] J. P. Roach, A. K. Churchland, and T. A. Engel. Choice selective inhibition drives stability and competition in decision circuits. *Nature Communications*, 14(11):147, Jan. 2023. ISSN 2041-1723. doi: 10.1038/s41467-023-35822-8.

[26] S. Sanghavi and K. Kar. Distinct roles of putative excitatory and inhibitory neurons in the macaque inferior temporal cortex in core object recognition behavior. page 2023.08.01.551579, Aug. 2023. doi: 10.1101/2023.08.01.551579. URL https://www.biorxiv.org/content/10.1101/2023.08.01.551579v1.

[27] M. Schimel. *Models of neural circuits as optimally driven dynamical systems*. PhD thesis, Apollo - University of Cambridge Repository, 2024. URL https://www.repository.cam.ac.uk/handle/1810/365584.

[28] B. Schuman, S. Dellal, A. Prönneke, R. Machold, and B. Rudy. Neocortical layer 1: An elegant solution to top-down and bottom-up integration. *Annual Review of Neuroscience*, 44(Volume 44, 2021):221–252, July 2021. ISSN 0147-006X, 1545-4126. doi: 10.1146/annurev-neuro-100520-012117.

[29] Y. Shao and S. Ostojic. Relating local connectivity and global dynamics in recurrent excitatory-inhibitory networks. *PLOS Computational Biology*, 19(1):e1010855, Jan. 2023. ISSN 1553-7358. doi: 10.1371/journal.pcbi.1010855.

[30] N. J. Sofroniew, D. Flickinger, J. King, and K. Svoboda. A large field of view two-photon mesoscope with subcellular resolution for in vivo imaging. *eLife*, 5:e14472, June 2016. ISSN 2050-084X. doi: 10.7554/eLife.14472.

[31] A. Song, A. S. Charles, S. A. Koay, J. L. Gauthier, S. Y. Thiberge, J. W. Pillow, and D. W. Tank. Volumetric two-photon imaging of neurons using stereoscopy (vtwins). *Nature Methods*, 14(4): 420–426, Apr. 2017. ISSN 1548-7105. doi: 10.1038/nmeth.4226.

[32] N. A. Steinmetz, C. Aydin, A. Lebedeva, M. Okun, M. Pachitariu, M. Bauza, M. Beau, J. Bhagat, C. Böhm, M. Broux, S. Chen, J. Colonell, R. J. Gardner, B. Karsh, F. Kloosterman, D. Kostadinov, C. Mora-Lopez, J. O'Callaghan, J. Park, J. Putzeys, B. Sauerbrei, R. J. J. van Daal, A. Z. Vollan, S. Wang, M. Welkenhuysen, Z. Ye, J. T. Dudman, B. Dutta, A. W. Hantman, K. D. Harris, A. K. Lee, E. I. Moser, J. O'Keefe, A. Renart, K. Svoboda, M. Häusser, S. Haesler, M. Carandini, and T. D. Harris. Neuropixels 2.0: A miniaturized high-density probe for stable, long-term brain recordings. *Science (New York, N.Y.)*, 372(6539):eabf4588, Apr. 2021. ISSN 1095-9203. doi: 10.1126/science.abf4588.

[33] A. Valente, S. Ostojic, and J. W. Pillow. Probing the relationship between latent linear dynamical systems and low-rank recurrent neural network models. *Neural Computation*, 34(9):1871–1892, 08 2022. ISSN 0899-7667. doi: 10.1162/neco_a_01522. URL https://doi.org/10.1162/neco_a_01522.

[34] J. C. R. Whittington, W. Dorrell, S. Ganguli, and T. Behrens. Disentanglement with biological constraints: A theory of functional cell types. Feb. 2023. URL https://openreview.net/forum?id=9Z_GfhZnGH.

[35] M. M. Yartsev, T. D. Hanks, A. M. Yoon, and C. D. Brody. Causal contribution and dynamical encoding in the striatum during evidence accumulation. *eLife*, 7:e34929, Aug. 2018. ISSN 2050-084X. doi: 10.7554/eLife.34929.

[36] H. Zeng and J. R. Sanes. Neuronal cell-type classification: challenges, opportunities and the path forward. *Nature Reviews Neuroscience*, 18(9):530–546, Sept. 2017. ISSN 1471-0048. doi: 10.1038/nrn.2017.85.

A Appendix

A1 Relationship between a noisy EI-RNN and a CTDS

In sec. 3, we showed that in the case of no noise, it is possible to map a linear RNN with E-I structure to a CTDS under certain conditions on the connectivity of the RNN. Here, we discuss the conditions under which a noisy EI-RNN maps to a CTDS.

Following sec. 3, let us assume the following linear RNN with N neurons:

$$\mathbf{y}_{t+1} = J\mathbf{y}_t + \eta_t; \eta_t \sim \mathcal{N}(0, P) \tag{A1}$$

We know from sec. 3 that for this RNN to map to a CTDS, $J = UV^\top$, where U and V are low rank matrices such that U is non-negative block diagonal and V obeys Dale's law. As a result, in the noiseless case, the CTDS that captures this model will have emission $C = U$, and the dynamics $A = V^\top U$.

In the noisy case, let's look at the joint distribution of the first two consecutive observations in a linear RNN model:

$$\begin{bmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \end{bmatrix} = \mathcal{N}\left(\begin{bmatrix} J\mathbf{y}_0 \\ J^2\mathbf{y}_0 \end{bmatrix}; \begin{bmatrix} P & JP \\ PJ^\top & JPJ^\top + P \end{bmatrix} \right) \tag{A2}$$

Next, we can also write the joint distribution of observations in any LDS model:

$$\begin{bmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \end{bmatrix} = \mathcal{N}\left(\begin{bmatrix} CA\mathbf{x}_0 \\ CA^2\mathbf{x}_0 \end{bmatrix}; \begin{bmatrix} CQC^\top + R & CAQC^\top \\ CQAC^\top & CAQA^\top C^\top + CQC^\top + R \end{bmatrix} \right) \tag{A3}$$

Now, if $C = U$ and $A = V^\top U$, the means of the observations in both settings match already. However, we need to compute the noise terms Q and R, so that the resultant CTDS maps to a noisy EI-RNN.

Following the joint distributions of observations in an RNN and an LDS, the following three expressions should hold true:

$$P = CQC^\top + R \tag{A4}$$

$$JP = CAQC^\top \tag{A5}$$

$$JPJ^\top + P = CAQA^\top C^\top + CQC^\top + R \tag{A6}$$

$$\tag{A7}$$

Let's start with the second expression: $JP = CAQC^\top$. Since $C = U$ and $A = V^\top U$, we obtain:

$$JP = UV^\top UQU^\top \tag{A8}$$

$$UV^\top P = UV^\top UQU^\top \tag{A9}$$

$$Q = U^\dagger P U^{\dagger\top} \tag{A10}$$

Since U is low-rank, not all of the covariance of P will be captured by Q. Thus, R should capture the remaining noise:

$$R = (I - UU^\top)(I - UU^\top)^\top P(I - UU^\top)(I - UU^\top)^\top \tag{A11}$$

Thus, $JP = CAQC^\top$ is now satisfied.

However, since we also want:

$$P = CQC^\top + R = UQU^\top + R \tag{A12}$$

$$\tag{A13}$$

For a linear RNN to be perfectly captured by a CTDS, the eigenvectors of P should either be aligned to the space spanned by U or to the space orthogonal to it.

Figure A1: Simulations with an E-I network. **A** Schematic of an E-I RNN, with red units representing excitatory cells and blue units representing inhibitory cells. Example generated activity from 10 randomly selected neurons. **B**. Top shows the true connectivity matrix J, while the bottom plot shows the J recovered using a CTDS. **C**. Root mean squared error between true and recovered J using a standard LDS and a CTDS model, each of latent state dimensionality 4. We fitted them to data from two E-I networks with 100 and 200 units each. The shaded region represents 95% confidence interval.

Finally, the third remaining equation has stricter implications:

$$JPJ^\top = CAQA^\top C^\top \tag{A14}$$

$$(UV^\top)P(VU^\top) = UV^\top UQU^\top VU^\top \tag{A15}$$

$UQU^\top$ is low-rank and thus cannot capture P entirely—it only captures covariance in the subspace spanned by U. Hence, for the LHS and RHS to be equal, we need U and $V^\top$ to have aligned subspaces so that the effect of projecting P on $V^\top$ results in the same covariance as that on the RHS.

So, in summary, we are able to find a restricted linear low-rank RNN that can be perfectly captured by a CTDS. Specifically, we want the connectivity J to be expressible as $UV^\top$, with rank D. Here, V should be in the column-space of U and obey Dale's law, while U should be positive block diagonal. Finally, we require that the noise covariance P has eigenvectors either completely aligned with the U subspace or entirely orthogonal to this subspace. In such a case, we have an CTDS that perfectly captures the activity of the RNN with the following parameters:

$$C = U, \ A = V^\top U \tag{A16}$$

$$Q = U^\dagger P U^{\dagger\top} \tag{A17}$$

$$R = (I - UU^\top)(I - UU^\top)^\top P(I - UU^\top)(I - UU^\top)^\top \tag{A18}$$

A2 Simulations with an E-I RNN

In order to illustrate the mapping between an EI-RNN and a CTDS, we generated simulated activity from two E-I networks with 100 and 200 units respectively. The connectivity matrix, J, had a non-negative rank of 2 for each of the sub-matrices formed using the excitatory and inhibitory rows. In each case, the connectivity matrix J was constrained to obey Dale's law, and was normalized to have eigenvalues less than 1. We set the noise matrix P in accordance with sec. A1.

We then generated 10 trials of 1000 time steps each from each network (Fig. A1 A), and fitted both standard LDS and CTDS models of latent-space dimensionality 4 to the generated activity.

We initialized both models randomly, and computed recovered connectivity matrices post-fitting as follows:

$$J = CA\Sigma_\infty C^\top \left(C\Sigma_\infty C^\top + R\right)^{-1} \tag{A19}$$

We solve the lyapunov equation $\Sigma_t = A\Sigma_{t-1}A^\top + Q$ to obtain Σ_∞. As shown in Fig. A1C, we found that the CTDS model recovers the connectivity matrix in both settings accurately, while a standard LDS does much worse in terms of root mean squared error. Fig. A1B shows the true and recovered connectivity matrices.

Figure A2: Histogram of the spike widths of FOF neurons. We see a bimodal distribution.

A3 Assigning cell-types to neurons in FOF and ADS

In this section, we discuss how we assigned cell-types to neurons in ADS and FOF. The ADS is a part of the striatum which is known to have primarily inhibitory neurons [8], hence we assumed all ADS neurons to be I cells. To identify cell types in the FOF, we constructed a histogram of the average spike widths of neurons in FOF. We found this to be a bimodal distribution (Fig. A2, and thus we labeled the neurons that had spike width less than 0.4ms as inhibitory and the remaining as excitatory. This is also consistent with the known distribution of neuronal cell types in this region (20% inhibitory, 80% excitatory).

A4 Inactivation strength during in-silico perturbation experiments

Figure A3: In-silico perturbation experiments with varying inactivation strengths. **A**. Perturbations in FOF at inactivation strengths of -2, -4 and -4. Solid bars show choice error using a multi-region CTDS, hatched bars show the same for a standard multi-region LDS. Gray shows error between the classifier's predictions and the animal's true behavior during no perturbation, yellow shows classifier's error on perturbed trials, when inactivating during early-half of trials, and green shows the same for late-half perturbations. **B**. Perturbations in ADS at inactivation strengths of -4, -8 and -12. Again, solid and hatched bars correspond to a multi-region CTDS and standard mutli-region LDS respectively. Similar to panel A, gray represents no perturbation, yellow shows early-half perturbation and green represents late-half perturbation.

As discussed in sec. 4.3, we performed in-silico optogenetic perturbation experiments in a fitted CTDS model by clamping latents corresponding to E cells in FOF, and those corresponding to I cells in ADS. In order to choose the inactivation strength for clamping the latent states, we varied the magnitude of the latents between $[-2, 20]$ during the perturbation time steps (either the first half of a trial, or the later half of trial). Since the model was fit on neural data that was processed to have a mean firing rate of 0Hz for every neuron, to ensure that clamping of latents indeed resulted in the inactivation of the corresponding neurons, we wanted their median firing rate to be negative. Hence, we chose the inactivation magnitude that resulted in a median firing rate of at least -1Hz across all neurons being perturbed. This resulted in an inactivation strength of -4 for FOF E latents, and -12 for ADS I latents (Fig. 3 shows results for these values).

However, our results are robust to variations in the inactivation strength. In Fig. A3A, solids bars in the center and right figures show choice errors when perturbing FOF E latents at strengths -4 and -8 respectively. We found that the experimental findings held true across these strengths, and that choice error was high only during perturbations in later half of the trial. The leftmost figure shows choice errors for inactivation strength of -2, which was presumably too small for the behavior to change. At the inactivation of -4, we obtained a median firing rate of -1 Hz across FOF E cells.

Similarly, in Fig. A3B, we saw the same perturbation trends for ADS across three values of inactivation strength ($-8, -12, -16$)—choice error was high during perturbations in either early or late half of the trial. We chose -12 as it resulted in a median firing rate of -1Hz. Finally, we also saw that across perturbation strengths, a standard multi-region LDS (Fig. A3 A & B, hatched bars) had low choice errors during perturbations in both regions, and they did not resemble experimental findings.

A5 Latent space visualization during in-silico perturbations

Figure A4: Latent state trajectories upon perturbations in FOF. **A**. This is the same as fig. 3A. Left shows unperturbed latents in FOF on an example trial, center shows latent trajectories when inactivating FOF latents during start of the same trial, and right show latent trajectories when inactivating FOF latents during end of the same trial. **B**. Effects of FOF perturbation on latents in ADS. Left shows unperturbed ADS latents during the same trial as **A**. Center shows ADS latents in response to early-half perturbations in FOF. Right shows ADS latents in response to late-half perturbations in FOF.

In Fig. 3A, we visualized the E and I latent states for FOF during an example trial—unperturbed, when perturbing FOF E latents during the early half of the trial, and the same during the later half of trial. Here, in Fig. A4B, we visualized the effects of FOF perturbations on latent states in ADS in the fitted CTDS model. We noticed that the trajectory of ADS latents did not suffer much upon early-half perturbations in FOF (compare Fig. A4B center to the leftmost plot). Thus, presumably

Figure A5: Latent state trajectories in a CTDS and a standard LDS. **A**. This is the same as fig. 3B. Left shows unperturbed latents in ADS on an example trial, center shows latent trajectories when inactivating ADS latents during start of the same trial, and right show latent trajectories when inactivating ADS latents during end of the same trial. **B** Same for a multi-region LDS with no cell-type constraints. We can see that the latent trajectories recover quickly after perturbations in a typical LDS. This is not the case with a CTDS as shown in **A**.

FOF latents recovered after early-half perturbations using the information coming in from the ADS. This is also supported by findings in Fig. 2D, where the cross-region dynamics between ADS and FOF is non-zero, indicating that information indeed flows from ADS to FOF.

Finally, we also visualized the latent trajectories of a standard multi-region LDS model when performing in-silico perturbations (Fig. A5). Here, we randomly inactivated 2 out of 4 latents per region during either early half or late half of trials. While in a multi-region CTDS model, inactivations in ADS derailed the latent dynamics entirely (Fig. A5A), we found that in a standard LDS the latent trajectories recovered quickly after perturbations (Fig. A5B). This is likely because CTDS constrained the connectivity between neurons and regions based on structural constraints in the brain, while the LDS did not have any such constraints. As a result, CTDS is able to accurately capture the dynamical effects of perturbation experiments, while LDS is unable to provide an accurate representation of underlying brain dynamics.

A6 Compute requirements and code

Finally, we trained these models on 2.8 GHz Intel Cascade Lake with 8 CPU cores, and developed our code on top of [17].