Goal-directed action preparation in humans entails a mixture of corticospinal neural computations

Corey G. Wadsley[1,2], Thuan Nguyen[3], Chris Horton[1], Ian Greenhouse[1,2,*]

[1]Action Control Lab, Department of Human Physiology, University of Oregon, Eugene, Oregon, USA

[2]Institute of Neuroscience, University of Oregon, Eugene, Oregon, USA

[3]School of Public Health, Portland State University-Oregon Health and Science University, Portland, Oregon, USA

Corresponding author: Ian Greenhouse (img@uoregon.edu)

Department of Human Physiology

181 Esslinger Hall,

1525 University St.

Eugene, OR 97403

Running title: Human action preparation entails corticospinal neural computations

Keywords: Human, Motor cortex, Action preparation, Gain modulation, Noninvasive brain stimulation

Key points:

- Neural computations determine what information is transmitted through brain circuits.

- We investigated whether the motor system uses computations similar to those observed in sensory systems by noninvasively stimulating the corticospinal pathway in humans during goal-directed action preparation.

- We discovered physiological evidence that corticospinal projections to behaviorally relevant muscles exhibit nonlinear gain computations, while projections to behaviorally irrelevant muscles exhibit linear suppression.

- Our findings suggest that certain computational principles generalize to the human motor system and serve to enhance the contrast between relevant and background neural activity.

- These results indicate that neural computations during goal-directed action preparation may support motor control by increasing signal-to-noise within the corticospinal pathway.

Abstract: The seemingly effortless ability of humans to transition from thinking about actions to initiating
them relies on sculpting corticospinal output from primary motor cortex. This study tested whether canonical
additive and multiplicative neural computations, well-described in sensory systems, generalize to the
corticospinal pathway during human action preparation. We used non-invasive brain stimulation to measure
corticospinal input-output across varying action preparation contexts during instructed-delay finger
response tasks. Goal-directed action preparation was marked by increased multiplicative gain of

corticospinal projections to task-relevant muscles and additive suppression of corticospinal projections to

non-selected and task-irrelevant muscles. Individuals who modulated corticospinal gain to a greater extent

were faster to initiate prepared responses. Our findings provide physiological evidence of combined additive

suppression and gain modulation in the human motor system. We propose these computations support

action preparation by enhancing the contrast between selected motor representations and surrounding

background activity to facilitate response selection and execution.

Introduction

Humans are experts at using their fingers for goal-directed actions to interact with their

environment effectively (Xu *et al.*, 2024). This behavioral repertoire is supported by a specially evolved

neural architecture comprising excitatory and inhibitory mechanisms that sculpt corticospinal (CS)

pathway output during goal-directed action preparation (Prut & Fetz, 1999; Duque *et al.*, 2017; Bundt &

Huster, 2024). The anatomy of the distributed cortical and spinal circuits that modulate CS output during

action preparation and execution is well characterized (Wang *et al.*, 2017; Derosiere & Duque, 2020).

However, the precise neural computations that unfold throughout the transition from action preparation to

execution are not (Churchland & Shenoy, 2024). Candidate computations have been proposed but are

based on neuronal circuit modeling and experimental evidence from investigations of non-motor neural

systems and behaviors (Greenhouse, 2022). Thus, it is unclear whether computational principles from

non-motor circuits generalize to the human motor system and account for the top-down modulation of CS

output during action preparation.

Influential models of goal-directed action preparation postulate that inhibition of the CS pathway

acts as a functional gate, with the removal of inhibition allowing selective excitatory signals to pass

through to the musculature and produce intended actions (Zagha *et al.*, 2015; Soteropoulos, 2018). Yet,

physiological recordings from non-human primates have provided mixed evidence for gating mechanisms

within the CS pathway (Merchant *et al.*, 2008; Kaufman *et al.*, 2010, 2013). Indeed, (dis)inhibition

supports a variety of computations other than gating in non-motor neural circuits (Katzner *et al.*, 2011;

Lee *et al.*, 2012; Mischler *et al.*, 2023). Multifaceted computations are also likely required in the motor

system since movements are not static but instead defined by various kinetic and kinematic features that

characterize a given action. Therefore, alternative interpretations to the functional gating account should

be considered to better explain how preparatory activity determines CS pathway excitability to shape

ensuing actions.

The selection and timing of goal-directed actions may be determined by neural computations

similar to, or bound to, those of non-motor circuits (Beste *et al.*, 2023). Such computations, responsible

for sculpting descending signals, are expected to manifest in distinct patterns of CS input-output during

goal-directed action preparation. One candidate computation is gain modulation, defined in studies of

sensory processing as a multiplicative (non-linear) change in a system's input-output relationship and a

proposed feature of canonical normalization processes within the brain (Hillyard *et al.*, 1998; Salinas &

Thier, 2000; Pi *et al.*, 2013). Computational models indicate gain modulation is important in determining

primary motor cortex (M1) output (Pouget & Snyder, 2000; Stroud *et al.*, 2018), the predominant source of

descending signals of the human CS tract (Welniarz *et al.*, 2017). Additive modulation, defined by a linear

shift across an input-output spectrum (Silver, 2010), is an alternative candidate computation that may

account for CS input-output adjustments during action preparation. Computational models of additive

modulation indicate effective changes in an input-output threshold (Mitchell & Silver, 2003), whereby

widespread suppression can stabilize a neuronal population by reducing the preponderance of

spontaneous firing (Ferguson & Cardin, 2020). During action preparation, additive modulation may

function as the 'quiet before the storm' to prepare the CS pathway for impending descending commands.

A third alternative is that multiplicative and additive computations are concurrent and

complementary. A precedent for synergistic computations can be found in the vision system. Extending a

divisive normalization framework (Carandini & Heeger, 2012) to the motor system, the activity of a motor

representation selected for action can be divided by the summed activity of alternative non-selected and

bystander motor representations, producing enhanced signal-to-noise (Georgopoulos & Stefanis, 2007;

Greenhouse, 2022). A combination of multiplicative and additive computations could serve to separate

figure from ground in a manner resembling neural sensory processing and would be consistent with a

general neurocomputational principle throughout the human nervous system akin to exponentiation, linear

filtering, and divisive normalization (Carandini & Heeger, 2012).

61  In this study, we tested the presence of multiplicative and additive computations in human CS

62 input-output during the behavioral state change from rest to action preparation of goal-directed finger

63 movements (Figure 1a). We hypothesized that goal-directed action preparation involves a concurrent gain

64 increase and additive suppression to enhance the contrast between selected and background motor

65 representations along the CS pathway.

66 **Methods**

67 *Participants*

68  Forty-five young, neurologically healthy adults volunteered to participate. The study was

69 approved by the institutional review board of the University of Oregon and performed with each

70 participant's understanding and written consent. All participants self-reported as right-handed and were

71 screened for contraindications to TMS before data collection. Five participants were excluded from data

72 collection due to a high resting motor threshold ($\geq$ 60% maximum stimulator output), making the TMS

73 input-output protocol unfeasible. An additional participant was excluded after data collection due to

74 noncompliance with task instructions. The remaining 39 participants completed the entire study and were

75 available for all statistical analyses (20 females, 19 males; mean age 20.3 yrs., range 18 to 30 yrs.).

76 *Task protocol*

77  Goal-directed action preparation was assessed with an instructed-delay two-choice reaction time

78 task (Figure 1b). Participants were seated comfortably in front of a computer monitor (59 Hz refresh rate,

79 ~60 cm viewing distance) with their hands palm-down and shoulder-width apart on a table surface. The

80 task was designed using Psychtoolbox-3 integrated with MATLAB (R2019b; The MathWorks) and a

81 custom response board (Makey Makey v.1.2; Joylabz) for accurate timing. Stimuli were synchronized with

82 task equipment through an analog-to-digital data-acquisition module (PCI-6620; National Instruments)

83 and verified during later offline analyses using photodiode recordings. The default task display consisted

84 of a blank grey background at trial onset. A fixation cue (centered white rectangle, 20 x 20 pixels)

85 appeared 0.3 s after trial onset. A warning cue appeared 1 to 1.5 s after trial onset (randomly drawn from

86 a uniform distribution) and indicated the forthcoming response (hollow leftward or rightward pointed

triangle, 140 x 150 pixels). An imperative stimulus (triangle filling white) appeared 0.9 s after the warning

cue and lasted until a button press was recorded or 0.8 s elapsed. Each trial was followed by an intertrial

interval (blank grey background) after the imperative stimulus and lasted until 7.6 s had elapsed from the

trial onset.

Figure 1: Assessment of human CS input-output during goal-directed action preparation. **a**, Corticospinal (CS) input-output
can exhibit additive and/or multiplicative modulation from baseline rest to a preparatory state. **b**, Goal-directed action preparation was
assessed with an instructed-delay two-choice reaction time task requiring left (L) or right (R) responses. Most trials were go trials
during which the imperative stimulus appeared until a reaction time (RT) was recorded or 0.8 s elapsed. A subset of catch trials (~8%)
was included to prevent participants from predicting imperative stimulus onset. **c**, The experiment was split into between-hand and
within-hand tasks, enabling the assessment of CS output from the left index finger with single-pulse transcranial magnetic stimulation
(TMS) when the targeted motor representation was at baseline and when it was selected, nonselected, or irrelevant for action
preparation. Here, input was TMS intensity relative to a participant's resting motor threshold (%RMT), and output was the amplitude
of ensuing motor evoked potentials (MEPs). **d**, CS input-output curves derived from pre-task resting state MEP data and fit with a
population-based nonlinear mixed effects model using a three-parameter Boltzmann function. Green dots and error bars indicate
mean MEP amplitude ± SEM at each TMS intensity. Dashed grey lines illustrate parameter estimates.

The experiment was split into between-hand and within-hand choice tasks (Figure 1c). Both tasks

were performed in the same session with a counterbalanced order across participants. Participants were

provided a short break in between tasks. Left and right imperative stimuli required button presses from

the left and right index fingers during the between-hand choice task and button presses from the right

index and pinky fingers during the within-hand task. Participants were instructed to relax during the

fixation period, to prepare but not execute the forthcoming response during the cue period, and to

execute the cued response as fast as possible when the imperative stimulus was presented. A subset of

no-go catch trials (i.e., no imperative stimulus; 8% of trials) was included to discourage participants from

responding prematurely in anticipation of the imperative stimulus. For each task, there was a total of 238

trials evenly distributed across left and right imperative stimuli and split across 4 blocks alternating

between 59 and 60 trials. The entire experiment lasted 2.5 hours on average.

Electromyography

Electrophysiological muscle activity was assessed with surface electromyography (EMG)

recorded using bipolar Ag-AgCl surface electrodes (Delsys Incorporated). EMG was collected from the

left and right FDI and the right *abductor digiti minimi* (ADM) hand muscles, which corresponded to the

primary agonist muscles for between-hand$_{left}$ (left FDI), between-hand$_{right}$ and within-hand$_{left}$ (right FDI),

and within-hand$_{right}$ (right ADM) choice alternatives. An additional EMG electrode was positioned over the

C3 vertebra of the neck to record TMS artifacts, and a shared ground electrode was positioned on the left

ulnar styloid process. EMG activity was amplified (×1000), bandpass filtered (20 – 450 Hz), and sampled

at 5000 Hz with a Bagnoli-8 amplifier (Delsys Incorporated) connected to a BNC-2090A terminal block

(National Instruments). EMG data were recorded on each trial for 4 s from onset using the MATLAB-

based VETA toolbox (Jackson & Greenhouse, 2019) and stored for later offline analyses.

Transcranial Magnetic Stimulation

Physiological changes in CS input-output during goal-directed action preparation were assessed

with single-pulse TMS delivered using a 70 mm figure-of-eight coil connected to a MagStim 200^2

stimulator (MagStim Ltd.). The coil was oriented to induce a posterior-to-anterior flowing current in the

underlying cortical tissue. The optimal coil position ('motor hotspot') over right M1 for eliciting a motor

evoked potential (MEP) in the left FDI was then assessed and marked on the scalp. The resting motor

threshold (RMT) of left FDI was defined as the minimum stimulator output required to elicit an MEP

amplitude of at least 0.05 mV at rest using a maximum-likelihood parameter estimation by sequential

testing strategy (Silbert *et al.*, 2013). TMS was delivered across a range of stimulation intensities relative

to a participant's RMT (90%, 110%, 130%, 150%, and 170%; 14 stimuli per intensity) to derive CS input-

output curves, with MEP peak-to-peak amplitude used as an index of CS output (Bestmann & Duque,

2016). The TMS input-output protocol was performed at baseline (start of fixation period) and during

action preparation (100 ms before the imperative stimulus) in the between-hand and within-hand tasks.

The task design enabled assessment of CS output to the left FDI muscle during baseline and during

action preparation when the muscle was selected (i.e., the agonist for prepared action), nonselected, or

task-irrelevant (Figure 1c). Nonselected refers to the context in which the left index finger was a choice

alternative in the task but was not cued to respond on a given trial, while task-irrelevant refers to the

context in which the left index finger was never a choice alternative in the task.

Dependent measures

Data were preprocessed using the VETA toolbox in two stages. In the first stage, EMG bursts,

MEPs, and TMS artifacts were marked with the automated detection *findEMG.m* function. In the second

stage, data was manually reviewed using the *visualizeEMG.m* function to ensure the accuracy of EMG

measures. Here, EMG data contaminated by artifacts were rejected, and EMG burst onset timings were

adjusted when required. Dependent measures were then extracted with custom scripts in MATLAB.

Behavioral measures were analyzed from non-TMS trials to avoid the influence of TMS on response

times. Go success was calculated as the percentage of go trials where a correct response was made

within 0.8 s from imperative stimulus onset. Catch trial success was calculated as the percentage of no-

go trials where the cued response was withheld correctly. Trials with button press RTs less than 100 ms

or EMG RTs less than 50 ms were then marked as premature responses and excluded from subsequent

analyses (M = 2.2 % of all trials). There was a strong positive correlation between button press and EMG

RT (r = 0.68, P < 0.001), as such, we report only EMG-based RTs for subsequent analyses since it is free

from electromechanical delays.

Dependent measures from stimulated trials during the between-hand and within-hand choice

tasks included pre-trigger root mean square EMG and MEP amplitudes. Trials with TMS were excluded

when pre-trigger root mean square EMG activity exceeded 30 µV in a -30 to -5 ms pre-TMS window or

when outlier MEP amplitudes were identified, defined as values more than three times the scaled median

absolute deviation from the median (M = 11.4 % of stimulated trials). The mean pre-TMS root mean

square EMG activity of the left FDI was calculated from the remaining trials for each preparation context,

collapsed across stimulation intensities. CS output of the left FDI was calculated as the mean MEP peak-

to-peak amplitude for each preparation context and stimulation intensity separately.

Statistical analyses

Statistical analyses were performed using R software (Version 4.3.0). The normality of data and

model-averaged residual plots were visually inspected where appropriate. Logarithmic transformations

were used in the case of non-normally distributed data. The alpha level was set to 0.05 for statistical

significance. P-values for pairwise post-hoc comparisons were adjusted for multiple comparisons using

the Benjamini-Hochberg method to control the false discovery rate at 5%. In-text data are presented as

nontransformed mean ± standard deviation unless otherwise specified.

Go and catch trial success rates were analyzed using an omnibus Friedman test with the factor of

Response (within-hand$_{left}$, within-hand$_{right}$, between-hand$_{left}$, between-hand$_{right}$) to verify task performance

was high and consistent across response alternatives. EMG RT was analyzed using a linear mixed-

effects model with fixed effects of Task (within-hand, between-hand) and Side (left, right) and random

effect of participant specificity to verify participants were responding fast and to determine any differences

between tasks and response alternatives.

We employ a population-based nonlinear mixed-effects model to determine the influence of

action preparation context on the CS input-output relationship. A modified three-parameter Boltzmann

sigmoid function (Heusser *et al.*, 2021) was used to model the nonlinear sigmoidal input-output

relationship between TMS intensity and MEP amplitude (Kukke *et al.*, 2014):

$$MEP(\%RMT) = 0 + Max/\{1 + exp[4 * Slope * (Midpoint - \%RMT)/Max]\}$$

where MEP(%RMT) is the predicted MEP peak-to-peak amplitude at a given TMS intensity, calculated as

a function of Max (upper plateau), Midpoint (inflection point), and Slope (maximum rate of change) input-

output parameter estimates. Here, the Midpoint and Slope parameters indicate additive and multiplicative

computations, respectively (Figure 1d). The lower plateau of CS input-output was constrained to 0 mV for

greater biological plausibility (i.e., MEP amplitude cannot be negative and should always be 0 mV at a 0%

TMS intensity) and provided a better fit of the data compared to an unconstrained model (constrained

AIC: 1307.5, unconstrained AIC: 1317.0).

Initial parameter estimates for the nonlinear mixed-effects models were derived from pre-task

resting state data to ensure independence between model optimization and statistical analyses.

Specifically, initial estimates were set to mean MEP amplitude at 170% RMT for Max, median stimulation

intensity for Midpoint (i.e., 130% RMT), and the linear gradient between 110% and 130% RMT for Slope

based on visual inspection of averaged MEP amplitude data. We then applied nonlinear least squares

modeling (*nls* function in R) with these initial parameter estimates to obtain corresponding optimal

parameter estimates of 2.34 mV for Max, 125.6 %RMT for Midpoint, and 0.055 mV/%RMT for Slope with

an iterative numerical integration procedure. These estimates were subsequently applied to a model of in-

task TMS data (*nlme* function in R). Max, Midpoint, and Slope parameters were estimated from this

random intercept model to account for participant variability. The model was then updated with the fixed

effect of Context (baseline, irrelevant, nonselected, selected) to examine the influence of action

preparation context on CS input-output. Updated models with the fixed effect of Task Order (within-hand

first, between-hand first) or Sex (male, female) were also included to determine any effects of task order

or participant sex on CS input-output parameters.

A linear mixed-effects model with a fixed effect of Context (baseline, irrelevant, nonselected,

selected) was used to confirm modulation of CS input-output was not attributable to shifts in pre-TMS root

mean square EMG activity.

We estimated a Gain Index *post priori* for each participant to capture interindividual differences in

CS input-output Slopes and evaluate whether observed context-dependent gain adjustments correlated

with task behavior. Direct parameter estimations could not be used from the population-based model

since variance is shared across participants and, thus, interindividual differences between contexts

derived from the population-based model are not meaningful. As such, we estimated the Gain Index by 1)

extracting the estimated Midpoint for each context and participant separately, 2) determining the two

neighboring input TMS intensities (e.g., 110% and 130% RMT if the estimated Midpoint was within these

bounds), 3) estimating the linear gradient (slope) between the mean MEP amplitudes from their individual

data corresponding to the selected input TMS intensities, and 4) calculating the estimated difference

between task-relevant contexts (average of selected and nonselected contexts) and task-irrelevant

contexts, normalized to baseline, i.e. $(\text{slope}_{relevant} - \text{slope}_{irrelevant})/\text{slope}_{baseline}$. Thus, a greater Gain Index

reflects a larger difference in the Slope of CS input-output between task-relevant and task-irrelevant

preparation contexts. The association between mean EMG RT, collapsed across all response types

(average non-stimulated trial number per response type: between-hand$_{left}$ 10.2 ± 2.0; between-hand$_{right}$

10.7 ± 1.8; within-hand$_{left}$ 10.4 ± 2.3; within-hand$_{right}$ 10.9 ± 2.0), and the Gain Index was then tested using

Pearson's correlation.

Results

CS input-output parameters changed during goal-directed action preparation in a context-

dependent manner (Figure 2a). The Context model provided a better fit of the data than the Null model

(Context AIC: 1264.3, Null AIC: 1312.9). Importantly, there was no effect of Context on pre-TMS muscle

activity ($F_{3,144} = 0.58$, $P = 0.629$), which was below 10 μV in all contexts (baseline 5.5 ± 2.5 μV; irrelevant

5.8 ± 4.0 μV; nonselected 5.5 ± 2.7 μV; selected 5.8 ± 2.5 μV). Thus, participants' muscles rested during

action preparation, and differences in alpha motoneuron depolarization did not explain the modulation of

CS input-output (Devanne *et al.*, 1997).

CS gain increases during response preparation for task-relevant representations

The Slope of CS input-output differed significantly across Context ($F_{3,730} = 4.26$, $P = 0.005$).

Estimated Slope was greater in the selected context compared to baseline ($F_{1,730} = 5.42$, $P = 0.040$) and

the irrelevant context ($F_{1,730} = 9.76$, $P = 0.011$). Estimated Slope in the nonselected context was also

greater than the irrelevant context ($F_{1,730} = 5.87$, $P = 0.040$) but not baseline ($F_{1,730} = 2.79$, $P = 0.143$).

There were no significant differences between baseline and irrelevant ($F_{1,730} = 0.95$, $P = 0.397$) as well as

between nonselected and selected contexts ($F_{1,730} = 0.23$, $P = 0.628$). Thus, context-dependent

multiplicative modulation of CS input-output was explained by an increase of Slope when the stimulated

motor representation was selected or nonselected for forthcoming action (i.e., task-relevant) compared to

baseline or when it was task-irrelevant (Figure 2b). Such nonlinear increases in sensitivity to excitatory

inputs are associated with response potentiation across sensory systems (Wood *et al.*, 2017).

Multiplicative gain modulation may be important for maximizing signal-to-noise and, computationally, can

extend the dynamic range of neuronal operations (Mitchell & Silver, 2003; Shine *et al.*, 2018). Of note is

that increased Slope was observed in both the selected and nonselected preparation contexts, indicating

a general role of gain modulation in potentiating task-relevant motor representations rather than

promoting effector-specific action selection. Overall, gain modulation may be pertinent for establishing the

pool of candidate actions.

The Max of CS input-output also differed significantly across Context ($F_{3,730} = 24.04$, $P < 0.001$).

Estimated Max was greater in the selected context compared to irrelevant ($F_{1,730} = 14.62$, $P = 0.001$) and

nonselected ($F_{1,730} = 6.44$, $P = 0.034$) contexts, which did not differ from each other ($F_{1,730} = 1.89$, $P =$

0.204). There were no differences between baseline and other contexts (all $P > 0.050$). Thus, CS output

maximum was greater during selected than nonselected and irrelevant contexts, but not baseline (Figure

2c). This pattern differed from that observed for the Slope and indicates the dynamic range of excitability

in the CS pathway is influenced by more than gain adjustments alone.

Additive suppression of CS input-output in nonselected and task-irrelevant preparatory contexts

The Midpoint of CS input-output differed significantly across Context ($F_{3,730} = 11.48$, $P < 0.001$).

Estimated Midpoint was greater in the irrelevant context than baseline ($F_{1,730} = 7.00$, $P = 0.012$) and the

selected context ($F_{1,730} = 18.54$, $P < 0.001$). Estimated Midpoint in the nonselected context was also

greater than baseline ($F_{1,730} = 9.65$, $P = 0.004$) and the selected context ($F_{1,730} = 25.32$, $P < 0.001$). There

was no significant difference between nonselected and irrelevant ($F_{1,730} = 0.05$, $P = 0.828$) and between

baseline and selected contexts ($F_{1,730} = 2.50$, $P = 0.137$). Thus, context-dependent modulation of CS

input-output was explained by an additive rightward shift of the Midpoint when the stimulated motor

representation was task-irrelevant or nonselected for the forthcoming action compared to when it was

selected (Figure 2d). The additive increase of the Midpoint during nonselected and irrelevant contexts

was also greater than baseline, consistent with inhibition of CS input-output relative to intertrial rest

(Derosiere & Duque, 2020). This nonspecific and apparently uniform suppression of irrelevant and

nonselected motor representations may reflect widespread noise reduction within the CS pathway

(Greenhouse *et al.*, 2015). These findings extend previous studies of preparatory inhibition and are

consistent with a mechanism for enhancing the contrast of selected motor representations through the

suppression of background activity.

Figure 2: Modulation of CS input-output across goal-directed action preparation contexts. **a**, CS input-output curves across
action preparation contexts. Insets indicate the estimated Max, Slope, and Midpoint parameters. **b**, Estimated Slope was greater in
the selected context compared to baseline and irrelevant, greater in the nonselected context than irrelevant, with no significant
differences between baseline and irrelevant or between nonselected and selected contexts. **c**, Estimated Max was greater in the
selected context compared to the irrelevant and nonselected contexts, which did not differ from each other, with no differences
between baseline and any other contexts. **d**, Estimated Midpoint was greater in the irrelevant and nonselected contexts compared to
baseline and the selected context, with no significant differences between nonselected and irrelevant or between baseline and
selected contexts. Data ranges represent mean ± SEM of the model-derived parameters in panels b – d, where the standard error
reflects the precision of the estimated coefficients.

Preparatory suppression during goal-directed action preparation

Preparatory suppression, expressed as a reduced MEP amplitude relative to baseline, was

observed during the irrelevant context across all stimulation intensities (110: $W = 615$, $P = 0.004$; 130: W

$= 573$, $P = 0.029$; 150: $W = 608$, $P = 0.005$; 170: $W = 616$, $P = 0.004$), only at the 110 %RMT stimulation

intensity during the nonselected context (110: $W = 644$, $P = 0.001$; 130: $W = 501$, $P = 0.372$; 150: $W =$

411, $P = 1.000$; 170: $W = 446$, $P = 1.000$), but at no stimulation intensity during the selected context (110:

$W = 393$, $P = 1.000$; 130: $W = 238$, $P = 0.100$; 150: $W = 271$, $P = 0.296$; 170: $W = 280$, $P = 0.382$). Thus,

preparatory suppression was observed across the entire input spectrum in the irrelevant context, only at

the lower end of the input spectrum in the nonselected context and at no part of the input spectrum in the

selected context (Figure 3a).

Figure 3: Preparatory suppression during goal-directed
action preparation. **a**, Preparatory suppression, expressed
as a reduced MEP amplitude relative to baseline, was
observed during the irrelevant context across all stimulation
intensities, only at the 110 %RMT stimulation intensity
during the nonselected context, and at no stimulation
intensity during the selected context. Crossbar boxes
indicate median hilows and the dashed horizontal line
indicates baseline MEP amplitude. Statistical results were
obtained from paired Wilcoxon-signed rank tests of mean
MEP amplitude against baseline. **b**, Conceptual overview of
how CS input-output modulation can reflect preparatory
suppression or enhancement depending on CS input and
modulation. Here, assuming fixed alternative parameters, a
multiplicative increase would be captured as preparatory
suppression or enhancement at input intensities below or
above the midpoint, respectively. In contrast, an additive
increase (i.e., rightward shift) would be captured as
preparatory suppression irrespective of the input intensity
relative to the midpoint.

Gain modulation of CS input-output across preparatory contexts is associated with reaction time

Participants responded accurately and quickly during the behavioral tasks. There were no

significant differences in accuracy across the response options during either go (Go: $X^2_3 = 5.08$, $P =$

0.166) or catch ($X_2^3 = 3.93$, $P = 0.269$) trials, with a high average trial success indicating that participants

performed the tasks correctly (Figure 4a). Reaction time (RT), calculated based on EMG burst onset, was

measured during task trials without TMS to avoid the direct influence of stimulation on motor output

(Foltys *et al.*, 2001). For EMG RT during non-stimulated trials, there was a significant fixed effect of Task

($F_{1,114}$ = 10.18, P = 0.002), with faster responses in the within-hand compared to the between-hand task.

There was no fixed effect of Side ($F_{1,114}$ = 0.17, P = 0.684) or a Task × Side interaction ($F_{1,114}$ = 2.88, P =

0.092). Thus, responses were faster when the prepared action involved a within-hand choice between the

fingers of the dominant hand compared to a between-hand choice. However, RTs remained fast across

all response alternatives (Figure 4b). Importantly, RTs were correlated between the two tasks (R = 0.38,

P = 0.001), indicating individuals differed reliably in their RTs. There was a significant negative Pearson

correlation between the Gain Index and EMG RT from non-TMS trials (R = -0.32, P = 0.045), whereby

participants with a larger gain modulation index had faster RTs (Figure 4c). That is, participants who

exhibited greater gain adjustments in the form of slope differences between task-relevant and task-

irrelevant behavioral contexts were faster to initiate responses.

Figure 4: Behavioral performance and relationship to CS input-output gain modulation. **a**, Accuracy during go (left) and catch
(right) trials for which there was no significant effect of Response type on either measure. **b**, Response initiation times based on EMG
were faster during the within-hand compared to the between-hand task. **c**, There was a significant negative Pearson correlation
between Gain Index and EMG RT from non-TMS trials whereby participants with greater gain in task-relevant than task-irrelevant
action preparation contexts were faster responders. Crossbar boxes indicate median hilows in all panels.

CS input-output is not affected by participant sex or task order

CS input-output curves derived from the Sex model did not provide a better fit of the data than the

null model (Sex AIC: 1319.1, Null AIC: 1312.9). There were no significant effects of Sex on estimated

Max ($F_{1,736}$ = 0.61, P = 0.434), Slope ($F_{1,736}$ = 0.27, P = 0.605), or Midpoint ($F_{1,736}$ = 2.85, P = 0.092) CS

input-output parameters. CS input-output curves derived from the Order model also did not fit the data

better than the null model (Order AIC: 1320.1). There were no significant effects of Order on estimated

Max ($F_{1,736}$ = 0.04, P = 0.844), Slope ($F_{1,736}$ < 0.01, P = 0.944), or Midpoint ($F_{1,736}$ = 0.02, P = 0.876) CS

input-output parameters. Thus, neither participant Sex nor task order influenced CS input-output during

goal-directed action preparation (Figure 5).

Figure 5: Modulation of CS input-output across participant sex and task order. **a**, CS input-out curves derived from the Sex
model. **b**, There were no significant effects of Sex on estimated Max, Slope, or Midpoint CS input-output parameters. **c**, CS input-
output curves derived from the Order model. **d**, Task order (WH: within-hand first; BH: between-hand first) had no significant effects
on estimated Max, Slope, or Midpoint CS input-output parameters. **e**, Extracted participant-wise estimates of Max, Slope, and Midpoint
parameters for between-subjects effects of Sex (*left*) and Order (*right*). Insets in subpanels a and c indicate estimated population
means for the Max, Slope, and Midpoint parameters. Data ranges in subpanels b and d represent mean ± SEM of the model-derived
parameters, where the standard error reflects the precision of the estimated coefficients.

Discussion

The present study provides novel physiological evidence of multiplicative and additive neural

computations within the human CS pathway during goal-directed action preparation. Specifically, we

showed that the gain of behaviorally relevant motor representations (selected and non-selected)

increases against a background of additive suppression of nonselected and task-irrelevant motor

representations during goal-directed action preparation. Furthermore, the magnitude of gain modulation

was associated with behavioral output, such that stronger modulators were faster responders. Below, we

discuss the implications of these findings for understanding state-dependent computations within the

human CS pathway during motor control.

Computational parallels between sensory and motor systems during action preparation

First, the findings indicate canonical computations well-described in sensory cortices (Salinas &

Sejnowski, 2001; Mitchell & Silver, 2003; Silver, 2010) generalize to the human motor system. Previous

work has established feature-based attention modulates gain within visual cortices (Treue & Trujillo,

1999), and normalization in the human visual system further supports attentional processes by

theoretically separating figures from the ground (Bloem & Ling, 2019). Similar gain adjustments have

been observed in the auditory cortex to increase spectrotemporal contrast (Rabinowitz *et al.*, 2011).

Computational models of auditory (Mischler *et al.*, 2023) and visual (Polack *et al.*, 2013) sensory systems

further indicate specific roles of gain for normalization and may mirror the contrast enhancement

mechanism we propose operates within the CS pathway during action preparation. Such state-dependent

gain adjustments for contrast enhancement could facilitate the transmission of descending signals along

the CS pathway, similar to models of basal ganglia output for action selection (Nambu *et al.*, 2002).

Importantly, while sensory and motor systems differ in their representational content and anatomy, they

share critical functional similarities for guiding behavior (Hatsopoulos & Suminski, 2011). Therefore,

contrast enhancement via simultaneously increased gain of behaviorally relevant and suppression of

background neural representations may be a canonical motif for sculpting cortical circuit output.

The current findings also align with recurrent neural network models of the non-human primate

motor cortex that indicate a specific role of gain modulation in moderating movement features such as

speed (Stroud *et al.*, 2018). Such models are typically based on electrophysiological recordings from a

neuronal subpopulation in a target region acquired without experimental control over the input strength to

the efferent motor pathway. Here, by controlling the input intensity and tailoring it to each participant, we

provide corroborative physiological evidence for these models by demonstrating that gain modulation

along the human CS pathway during a preparatory state facilitates the execution of subsequent actions.

Based on our data, we propose that increased gain of task-relevant motor representations occurs against

a backdrop of suppressed nonselected and irrelevant motor representations to enhance the contrast

between selected and bystander motor representations (Greenhouse *et al.*, 2015). We theorize this

extension to existing models further promotes the initiation of prepared responses by increasing signal-to-

noise within the CS pathway via divisive normalization (Figure 6). Our data can also inform computational

models of the motor cortex by providing useful information about the dynamic range of physiological

outputs and behavioral relationships in a human model. Translation of the context-dependent approach to

other model systems can help evaluate whether CS input-output adjustments during an output-null

(preparatory) motor state hasten the transition to an output-potent (execution) motor state (Hannah *et al.*,

2018; Churchland & Shenoy, 2024).

Figure 6: Theorized role of divisive normalization during goal-directed action preparation. CS input-output during action
preparation depends upon task contexts. Schematized CS input-output plots represent the multiplicative increase in the selected and
nonselected contexts and the additive increase in nonselected and task-irrelevant contexts. Speculatively, complementary
multiplicative and additive changes can enhance the signal-to-noise ratio (SNR) between selected and bystander motor
representations through divisive normalization to improve the fidelity of descending CS commands during action preparation.

Context-dependent computations with common coding frameworks

Context-dependent computations during goal-directed action preparation may fit within cognitive

'common coding' frameworks. These frameworks posit that human action control tasks are supported by

internal representations ('event files') of stimulus-response relationships and their sensory effects

(Hommel, 2009). Accordingly, action preparation entails retrieving an event file that triggers integrated

neuronal processing across nonmotor and motor brain regions to prime action execution (Beste *et al.*,

2023). We observed neurophysiological evidence of corticospinal input-output computations recapitulated

from sensory systems during action preparation, which may suggest a perception-action common code

between motor and nonmotor systems. The current results, conceptualized within a common code

framework, indicate that event files may have widespread motoric effects on not only the CS pathway to

effectors involved directly in stimulus-response bindings but also suppression of task-irrelevant CS

effector representations. In a somewhat homologous manner, modulation of sensory feedback during

action control occurs at both spinal and supraspinal levels and can include a mixture of gain and

suppression computations (Azim & Seki, 2019). Binding the motor command to anticipated perceptual

feedback may occur via common gain modulation and suppression processes, with up-or-down regulation

of neural pathways contingent upon their behavioral relevance (Frings *et al.*, 2020). Event file bindings

are likely strong in the instructed-delay response task we employed; however, it is important to note that

we did not set out to test event files explicitly and did not manipulate binding versus retrieval processes.

Nevertheless, our findings raise the interesting possibility that gain increases in CS output during goal-

directed action preparation are reciprocal to gain decreases in cutaneous feedback (Seki & Fetz, 2012)

but coupled to gain increases in proprioceptive feedback during action execution (Confais *et al.*, 2017).

Future work can investigate whether such changes arise from a common descending command.

Reevaluating context-dependent preparatory suppression across the CS input spectrum

An interesting observation from the current study was the lack of MEP suppression in the

selected context. Previous TMS studies have observed *preparatory suppression* (i.e., reduced MEP

amplitude below baseline during action preparation) across selected, nonselected, and task-irrelevant

contexts, albeit variably (Bundt & Huster, 2024). Our finding of enhanced CS gain in task-relevant

contexts may appear in conflict with inhibitory accounts (Ibáñez *et al.*, 2020). However, previous studies

often used a TMS intensity of ≤ 120 %RMT and, thus, only captured CS output at the lower end of the

input spectrum. By using a range of TMS intensities, we could model the entire CS input-output function,

providing a more complete picture of the underlying processes. Mathematically, both additive and

multiplicative changes in a sigmoidal input-output function would have complex effects on whether

context-dependent modulation is captured as suppression or enhancement when input intensity is below

or above the Midpoint (Figure 3b). We speculate that previous evidence of preparatory suppression of

task-relevant motor representations may result from TMS intensities below the CS input-output Midpoint.

Importantly, TMS may not mimic endogenous input to the CS pathway when transitioning from action

preparation to execution, which remains open for further investigation.

Neural basis of context-dependent gain adjustments along the CS pathway

Simultaneous gain adjustments and the suppression of background activity along the entire CS

pathway may rely on cooperative cortical, subcortical, and spinal mechanisms. For example, subcortical-

cortical projections from the thalamus to the prefrontal cortex selectively influence signal (figure) and

noise (ground) to guide cued behavior in the context of uncertainty via two distinct intracortical

mechanisms (Mukherjee *et al.*, 2021). Similar motifs may exist in thalamic projections to M1 to modulate

CS output (Halassa & Sherman, 2019). According to this framework, both increased gain of selected

motor representations and suppressed background activity would arise from a modulatory influence of

intracortical inhibitory circuits (Ferguson & Cardin, 2020). Alternatively, inhibition of background activity

may reflect the engagement of CS projections to spinal inhibitory neurons to suppress muscles that, if

activated, would interfere with the execution of the prepared action (Griffin & Strick, 2020). Computations

may also be mediated directly by spinal motoneurons, as MEPs measured with surface EMG exhibit

sigmoidal input-output properties, but individual spinal motor unit discharge rates exhibit linear input-

output properties (Devanne *et al.*, 1997). Summations across pools of spinal motoneurons operating

around the subliminal fringe may contribute to nonlinear transformations at the population level (Di

Lazzaro *et al.*, 1999), producing the sigmoidal shape modeled in our data. This account indicates that

population-level activity of spinal motoneurons operating more closely to an activation threshold during

action preparation could be responsible for the context-dependent gain changes we observed,

independent of cortical mechanisms. However, evidence of altered corticospinal input-output properties,

such as elevated gain in Parkinson's disease (Bologna *et al.*, 2018) and reduced maximum MEP output

after stroke (Buetefisch *et al.*, 2018), underscores the importance of cortico-subcortical mechanisms.

Therefore, it is likely that the observed CS input-output computations depend on cooperative cortical and

spinal mechanisms rather than a single stage in the pathway.

Limitations & Future Directions

The present study has some limitations. Using motor-evoked potentials (MEPs) elicited by TMS

provides an accurate assay of CS output. However, when probed with single-pulse TMS, MEP

modulation can arise from various mechanisms along the entire CS pathway, including intracortical,

transcortical, subcortical, and spinal mechanisms (Duque *et al.*, 2017), which may have unique roles in

action preparation (Derosiere & Duque, 2020). Thus, while we demonstrate that various computations

manifest in CS output, we cannot disentangle the level of the motor system in which they arise. Future

experiments will be able to determine at which level within the CS pathway these computations occur by

leveraging multimodal approaches, such as paired-pulse TMS, to characterize better the multifaceted

neuronal response to noninvasive brain stimulation and by studying clinical populations known to have

CS input-output irregularities. Another limitation relates to in-task TMS only being delivered at baseline or

a fixed time point late during action preparation. Limiting the temporal sampling was an important

methodological choice to collect a reliable number of trials for accurate CS input-output modeling across

various contexts. However, the time course of the observed computations during the transition from

action preparation to execution remains unclear. Future experiments may employ methods to sample CS

input-output effectively to enable a greater temporal resolution during action preparation (e.g., Alavi et al.,

2019).

Conclusion

The present study aimed to test the presence of multiplicative and additive computations in

human CS input-output during goal-directed action preparation. We provided physiological evidence that

goal-directed action preparation of finger responses entails a mixture of input-output computations within

the human motor system. These computations are context-dependent, characterized by gain increases in

task-relevant motor representations against a backdrop of suppression in bystander motor

representations, potentially facilitating action initiation through contrast enhancement. The recapitulation

of canonical neural computations across motor and non-motor neuronal circuits suggests that these

processes may generalize throughout the human nervous system.

Additional information

Data availability statement: The data and reproducible code support this study's findings will be made

publicly available in OSF upon publication, reference number osf.io/837ms.

Acknowledgments: The authors thank Kate Bakken, Mitch Fisher, Charlie Lewkowitz, Hayami Nishio,

Michelle Ortman, and Tania Sarabia for their help with data collection and preprocessing. The authors

also thank Luca Mazzucato and Richard Ivry for their helpful feedback on this manuscript.

Author contributions: Conceptualization, C.G.W and I.G.; Methodology, C.G.W, T.N., and I.G.; Formal

Analysis, C.G.W. and T.N.; Investigation, C.G.W. and C.H.; Writing – Original Draft, C.G.W. and I.G.;

Writing – Reviewing & Editing, C.G.W, T.N., C.H., and I.G.; Visualization, C.G.W.; Supervision, I.G.;

Funding acquisition, I.G.

Competing interests: The authors declare no competing interests.

Funding: This research was supported by NIH grant NINDS R01NS123115.

References

Alavi SMM, Goetz SM & Peterchev A V. (2019). Optimal Estimation of Neural Recruitment Curves Using

Fisher Information: Application to Transcranial Magnetic Stimulation. *IEEE Transactions on Neural*

Systems and Rehabilitation Engineering **27,** 1320–1330.

Azim E & Seki K (2019). Gain control in the sensorimotor system. *Curr Opin Physiol* **8,** 177–187.

Beste C, Münchau A & Frings C (2023). Towards a systematization of brain oscillatory activity in actions.

Commun Biol **6,** 137.

Bestmann S & Duque J (2016). Transcranial Magnetic Stimulation. *The Neuroscientist* **22,** 392–405.

Bloem IM & Ling S (2019). Normalization governs attentional modulation within human visual cortex. *Nat*

Commun **10,** 5660.

Bologna M, Guerra A, Paparella G, Giordo L, Alunni Fegatelli D, Vestri AR, Rothwell JC & Berardelli A

(2018). Neurophysiological correlates of bradykinesia in Parkinson's disease. *Brain* **141,** 2432–

2444.

Buetefisch CM, Revill KP, Haut MW, Kowalski GM, Wischnewski M, Pifer M, Belagaje SR, Nahab F,

Cobia DJ, Hu X, Drake D & Hobbs G (2018). Abnormally reduced primary motor cortex output is

related to impaired hand function in chronic stroke. *J Neurophysiol* **120,** 1680–1694.

Bundt C & Huster RJ (2024). Corticospinal excitability reductions during action preparation and action

stopping in humans: Different sides of the same inhibitory coin? *Neuropsychologia* **195,** 108799.

Carandini M & Heeger DJ (2012). Normalization as a canonical neural computation. *Nat Rev Neurosci* **13,**

51–62.

Churchland MM & Shenoy K V. (2024). Preparatory activity and the expansive null-space. *Nat Rev*

Neurosci **25,** 213–236.

Confais J, Kim G, Tomatsu S, Takei T & Seki K (2017). Nerve-Specific Input Modulation to Spinal

Neurons during a Motor Task in the Monkey. *The Journal of Neuroscience* **37,** 2612–2626.

Derosiere G & Duque J (2020). Tuning the Corticospinal System: How Distributed Brain Circuits Shape

Human Actions. *The Neuroscientist* **26,** 359–379.

Devanne H, Lavoie BA & Capaday C (1997). Input-output properties and gain changes in the human

corticospinal pathway. *Exp Brain Res* **114,** 329–338.

Duque J, Greenhouse I, Labruna L & Ivry RB (2017). Physiological Markers of Motor Inhibition during

Human Behavior. *Trends Neurosci* **40,** 219–236.

Ferguson KA & Cardin JA (2020). Mechanisms underlying gain modulation in the cortex. *Nat Rev*

Neurosci **21,** 80–92.

Foltys H, Sparing R, Boroojerdi B, Krings T, Meister IG, Mottaghy FM & Töpper R (2001). Motor control in

simple bimanual movements: a transcranial magnetic stimulation and reaction time study. *Clinical*

Neurophysiology **112,** 265–274.

Frings C, Hommel B, Koch I, Rothermund K, Dignath D, Giesen C, Kiesel A, Kunde W, Mayr S, Moeller B,

Möller M, Pfister R & Philipp A (2020). Binding and Retrieval in Action Control (BRAC). *Trends Cogn*

Sci **24,** 375–387.

Georgopoulos AP & Stefanis CN (2007). Local shaping of function in the motor cortex: Motor contrast,

directional tuning. *Brain Res Rev* **55,** 383–389.

Greenhouse I (2022). Inhibition for gain modulation in the motor system. *Exp Brain Res* **240,** 1295–1302.

Greenhouse I, Sias A, Labruna L & Ivry RB (2015). Nonspecific Inhibition of the Motor System during

Response Preparation. *Journal of Neuroscience* **35,** 10675–10684.

Griffin DM & Strick PL (2020). The motor cortex uses active suppression to sculpt movement. *Sci Adv*;

DOI: 10.1126/sciadv.abb8395.

Halassa MM & Sherman SM (2019). Thalamocortical Circuit Motifs: A General Framework. *Neuron* **103,**

762–770.

Hannah R, Cavanagh SE, Tremblay S, Simeoni S & Rothwell JC (2018). Selective Suppression of Local

Interneuron Circuits in Human Motor Cortex Contributes to Movement Preparation. *The Journal of*

Neuroscience **38,** 1264–1276.

Hatsopoulos NG & Suminski AJ (2011). Sensing with the Motor Cortex. *Neuron* **72,** 477–487.

Heusser K, Heusser R, Jordan J, Urechie V, Diedrich A & Tank J (2021). Baroreflex Curve Fitting Using a

WYSIWYG Boltzmann Sigmoidal Equation. *Front Neurosci*; DOI: 10.3389/fnins.2021.697582.

Hillyard SA, Vogel EK & Luck SJ (1998). Sensory gain control (amplification) as a mechanism of selective

attention: electrophysiological and neuroimaging evidence. *Philos Trans R Soc Lond B Biol Sci* **353,**

1257–1270.

Hommel B (2009). Action control according to TEC (theory of event coding). *Psychological Research*

Psychologische Forschung **73,** 512–526.

Ibáñez J, Hannah R, Rocchi L & Rothwell JC (2020). Premovement Suppression of Corticospinal

Excitability may be a Necessary Part of Movement Preparation. *Cerebral Cortex* **30,** 2910–2923.

Jackson N & Greenhouse I (2019). VETA: An Open-Source Matlab-Based Toolbox for the Collection and

Analysis of Electromyography Combined With Transcranial Magnetic Stimulation. *Front Neurosci*;

DOI: 10.3389/fnins.2019.00975.

Katzner S, Busse L & Carandini M (2011). GABA $_A$ Inhibition Controls Response Gain in Visual Cortex.

The Journal of Neuroscience **31,** 5931–5941.

Kaufman MT, Churchland MM, Santhanam G, Yu BM, Afshar A, Ryu SI & Shenoy K V. (2010). Roles of

Monkey Premotor Neuron Classes in Movement Preparation and Execution. *J Neurophysiol* **104,**

799–810.

Kaufman MT, Churchland MM & Shenoy K V. (2013). The roles of monkey M1 neuron classes in

movement preparation and execution. *J Neurophysiol* **110,** 817–825.

Kukke SN, Paine RW, Chao C-C, de Campos AC & Hallett M (2014). Efficient and Reliable

Characterization of the Corticospinal System Using Transcranial Magnetic Stimulation. *Journal of*

Clinical Neurophysiology **31,** 246–252.

Di Lazzaro V, Oliviero A, Profice P, Insola A, Mazzone P, Tonali P & Rothwell JC (1999). Effects of

voluntary contraction on descending volleys evoked by transcranial electrical stimulation over the

motor cortex hand area in conscious humans. *Exp Brain Res* **124,** 525–528.

Lee S-H, Kwan AC, Zhang S, Phoumthipphavong V, Flannery JG, Masmanidis SC, Taniguchi H, Huang

ZJ, Zhang F, Boyden ES, Deisseroth K & Dan Y (2012). Activation of specific interneurons improves

V1 feature selectivity and visual perception. *Nature* **488,** 379–383.

Merchant H, Naselaris T & Georgopoulos AP (2008). Dynamic Sculpting of Directional Tuning in the

Primate Motor Cortex during Three-Dimensional Reaching. *The Journal of Neuroscience* **28,** 9164–

9172.

Mischler G, Keshishian M, Bickel S, Mehta AD & Mesgarani N (2023). Deep neural networks effectively

model neural adaptation to changing background noise and suggest nonlinear noise filtering

methods in auditory cortex. *Neuroimage* **266,** 119819.

Mitchell SJ & Silver RA (2003). Shunting Inhibition Modulates Neuronal Gain during Synaptic Excitation.

Neuron **38,** 433–445.

Mukherjee A, Lam NH, Wimmer RD & Halassa MM (2021). Thalamic circuits for independent control of
prefrontal signal and noise. *Nature* **600,** 100–104.

Nambu A, Tokuno H & Takada M (2002). Functional significance of the cortico–subthalamo–pallidal
'hyperdirect' pathway. *Neurosci Res* **43,** 111–117.

Pi H-J, Hangya B, Kvitsiani D, Sanders JI, Huang ZJ & Kepecs A (2013). Cortical interneurons that
specialize in disinhibitory control. *Nature* **503,** 521–524.

Polack P-O, Friedman J & Golshani P (2013). Cellular mechanisms of brain state–dependent gain
modulation in visual cortex. *Nat Neurosci* **16,** 1331–1339.

Pouget A & Snyder LH (2000). Computational approaches to sensorimotor transformations. *Nat Neurosci*
3, 1192–1198.

Prut Y & Fetz EE (1999). Primate spinal interneurons show pre-movement instructed delay activity.
Nature **401,** 590–594.

Rabinowitz NC, Willmore BDB, Schnupp JWH & King AJ (2011). Contrast Gain Control in Auditory
Cortex. *Neuron* **70,** 1178–1191.

Salinas E & Sejnowski TJ (2001). Book Review: Gain Modulation in the Central Nervous System: Where
Behavior, Neurophysiology, and Computation Meet. *The Neuroscientist* **7,** 430–440.

Salinas E & Thier P (2000). Gain Modulation. *Neuron* **27,** 15–21.

Seki K & Fetz EE (2012). Gating of Sensory Input at Spinal and Cortical Levels during Preparation and
Execution of Voluntary Movement. *The Journal of Neuroscience* **32,** 890–902.

Shine JM, Aburn MJ, Breakspear M & Poldrack RA (2018). The modulation of neural gain facilitates a
transition between functional segregation and integration in the brain. *Elife*; DOI:
10.7554/eLife.31130.

Silbert BI, Patterson HI, Pevcic DD, Windnagel KA & Thickbroom GW (2013). A comparison of relative-

frequency and threshold-hunting methods to determine stimulus intensity in transcranial magnetic

stimulation. *Clinical Neurophysiology* **124,** 708–712.

Silver RA (2010). Neuronal arithmetic. *Nat Rev Neurosci* **11,** 474–489.

Soteropoulos DS (2018). Corticospinal gating during action preparation and movement in the primate

motor cortex. *J Neurophysiol* **119,** 1538–1555.

Stroud JP, Porter MA, Hennequin G & Vogels TP (2018). Motor primitives in space and time via targeted

gain modulation in cortical networks. *Nat Neurosci* **21,** 1774–1783.

Treue S & Trujillo JCM (1999). Feature-based attention influences motion processing gain in macaque

visual cortex. *Nature* **399,** 575–579.

Wang X, Liu Y, Li X, Zhang Z, Yang H, Zhang Y, Williams PR, Alwahab NSA, Kapur K, Yu B, Zhang Y,

Chen M, Ding H, Gerfen CR, Wang KH & He Z (2017). Deconstruction of Corticospinal Circuits for

Goal-Directed Motor Skills. *Cell* **171,** 440-455.e14.

Welniarz Q, Dusart I & Roze E (2017). The corticospinal tract: Evolution, development, and human

disorders. *Dev Neurobiol* **77,** 810–829.

Wood KC, Blackwell JM & Geffen MN (2017). Cortical inhibitory interneurons control sensory processing.

Curr Opin Neurobiol **46,** 200–207.

Xu J, Mawase F & Schieber MH (2024). Evolution, biomechanics, and neurobiology converge to explain

selective finger motor control. *Physiol Rev* **104,** 983–1020.

Zagha E, Ge X & McCormick DA (2015). Competing Neural Ensembles in Motor Cortex Gate Goal-

Directed Motor Output. *Neuron* **88,** 565–577.