Title: Smooth muscle expression of RNA editing enzyme *ADAR1* controls activation of RNA sensor MDA5 in atherosclerosis

Authors: Chad S. Weldy[1,2,3,*], Qin Li[4], João P. Monteiro[1], Hongchao Guo[5], Drew Galls[6], Wenduo Gu[1], Paul P. Cheng[1,3], Markus Ramste[1], Daniel Li[1], Brian T. Palmisano[1], Disha Sharma[1], Matthew D. Worssam[1], Quanyi Zhao[1], Amruta Bhate[6], Ramendra K. Kundu[1], Trieu Nguyen[1], Jin Billy Li[3,6,†], Thomas Quertermous[1,3,6,†,*]

Affiliations:
[1]Department of Medicine, Division of Cardiovascular Medicine, Stanford University
[2]Stanford Center for Inherited Cardiovascular Disease, Stanford University
[3]Stanford Cardiovascular Institute
[4]Department of Genetics, University of Pennsylvania
[5]University of Utah, Cardiovascular Research and Training Institute
[6]Department of Genetics, Stanford University
[†]Co-senior authorship

*Corresponding Authors:
    Chad S. Weldy, MD, PhD
    Instructor of Medicine
    300 Pasteur Drive, Falk CVRC
    Stanford, CA 94305
    360-305-8151
    weldyc@stanford.edu

    Thomas Quertermous, MD
    William G. Irwin Professor of Medicine
    300 Pasteur Drive, Falk CVRC
    Stanford, CA 94305
    650-498-5794
    tomq1@stanford.edu

Short title: ADAR1 as a causal mechanism of atherosclerosis

One Sentence Summary: Smooth muscle expression of RNA editing enzyme ADAR1 regulates activation of double strand RNA sensor MDA5 in novel mechanism of atherosclerosis.

Abstract:

Mapping the genomic architecture of complex disease has been predicated on the understanding that genetic variants influence disease risk through modifying gene expression. However, recent discoveries have revealed that a significant burden of disease heritability in common autoinflammatory disorders and coronary artery disease is mediated through genetic variation modifying post-transcriptional modification of RNA through adenosine-to-inosine (A-to-I) RNA editing. This common RNA modification is catalyzed by ADAR enzymes, where ADAR1 edits specific immunogenic double stranded RNA (dsRNA) to prevent activation of the double strand RNA (dsRNA) sensor MDA5 (*IFIH1*) and stimulation of an interferon stimulated gene (ISG) response. Multiple lines of human genetic data indicate impaired RNA editing and increased dsRNA sensing by MDA5 to be an important mechanism of coronary artery disease (CAD) risk. Here, we provide a crucial link between observations in human genetics and mechanistic cell biology leading to progression of CAD. Through analysis of human atherosclerotic plaque, we implicate the vascular smooth muscle cell (SMC) to have a unique requirement for RNA editing, and that ISG induction occurs in SMC phenotypic modulation, implicating MDA5 activation. Through culture of human coronary artery SMCs, generation of a conditional SMC specific *Adar1* deletion mouse model on a pro-atherosclerosis background with additional constitutive deletion of MDA5 (*Ifih1*), and with incorporation of single cell RNA sequencing cellular profiling, we further show that Adar1 controls SMC phenotypic state by regulating Mda5 activation, is required to maintain vascular integrity, and controls progression of atherosclerosis and vascular calcification. Through this work, we describe a fundamental mechanism of CAD, where cell type and context specific RNA editing and sensing of dsRNA mediates disease progression, bridging our understanding of human genetics and disease causality.

Main Text:

INTRODUCTION

As coronary artery disease (CAD) is the worldwide leading cause of death[1], there is enormous interest to map the genomic mechanisms of disease, establish biological causality, and accelerate therapeutic drug development. Despite the discovery of hundreds of genomic loci associated with CAD[2-5] the promise of genome wide association studies (GWAS) to identify new mechanisms for treatment outside of lipid lowering therapy has not been fulfilled. Quantitative trait loci (QTL) studies mapping gene expression QTLs (eQTLs) have helped bridge GWAS variants to their molecular mechanisms[6,7]. However, post-transcriptional processes, such as RNA editing, have critical functions in health and disease and genetic determinants of RNA modification are important and poorly appreciated disease mechanisms[8-10]. Catalyzed by adenosine deaminase acting on RNA enzymes (ADARs), adenosine-to-inosine (A-to-I) RNA editing is a highly frequent RNA modification that is evolutionarily conserved across all metazoan species[8,11]. We have recently reported that common genetic variants that modify the frequency of A-to-I RNA editing, loci termed 'editing-QTLs' (edQTL), significantly increase CAD risk as well as numerous other autoinflammatory disorders[12]. This discovery raises the important possibility that genetically determined RNA editing function is a causal mechanism of CAD, yet the biological underpinning of this relationship is unclear.

Foundational work has revealed that ADAR1 (gene symbol *ADAR*) mediated RNA editing is a novel mechanism of inflammation and disease[13-16]. A principle function of ADAR1 is to bind double-stranded RNA (dsRNA) in non-coding repetitive elements and convert adenosines to inosines to facilitate the clearance of dsRNA molecules[17]. This is in contrast to ADAR2 (gene symbol *ADARB1*) that serves to edit coding region of transcripts to modify transcript function[11,18]. ADAR1 is ubiquitously expressed across human tissues and its function to edit dsRNA suppresses the sensing of "non-self" dsRNA that is mediated by the cytosolic sensor of dsRNA, MDA5 (gene symbol *IFIH1*) (Fig 1A)[14,19,20]. Mice deficient in Adar1 editing are embryonic lethal due to activation of an interferon-stimulated gene (ISG) response, which is rescued to full life span when Mda5 (*Ifih1*) is also knocked out[14]. In humans, rare variants resulting in loss of function

in ADAR1 or gain of function in MDA5 result in two severe and clinically overlapping

Mendelian interferon mediated disorders, Acardi Goutières Syndrome and Singleton

Merton Syndrome, both of which are characterized by a profound early onset vascular

calcification phenotype demonstrating a link between RNA editing and dsRNA sensing

in vascular disease[21,22].

Common genetic variation that decreases RNA editing and increases CAD risk

have similar effects on risk for autoinflammatory disorders such as inflammatory bowel

disease and lupus[12]. To further link RNA editing to ISG response in human coronary

arteries, risk allele-specific RNA editing was found to be lower and closely predicted an

increased interferon gene response (implicating MDA5 activation) in the coronary

arteries of patients with CAD[12]. Although prior work has identified ADAR1 as mediating

atherosclerotic disease risk through RNA editing of specific RNA transcripts such as

NEAT1 and *CTSS*[23,24], human genetic data corroborates a separate 'axis' of global

ADAR1 RNA editing of immunogenic dsRNA to regulate MDA5 activation (ADAR1-

dsRNA-MDA5 axis) as a mechanism of atherogenesis (Fig 1A). The relationship of this

axis to CAD is further supported by the notable discovery that loss of function variants in

IFIH1 (MDA5) are protective against CAD[25] and that the *IFIH1* locus meets genome

wide level significant for association to CAD in GWAS[5].

Here, by analyzing human atherosclerotic plaque single cell RNA sequencing

(scRNAseq) data, we identified that vascular smooth muscle cells in atherosclerosis

have a uniquely high expression of RNA that require ADAR1 mediated RNA editing to

evade MDA5 activation. Through *in vitro* culture of primary human coronary artery

smooth muscle cells (HCASMCs) in relevant models, generation of genetic mouse

models with conditional deletion of *Adar1* in vascular smooth muscle cells (SMCs) in

homozygous loss of function and haploinsufficiency in atherosclerosis, extensive single

cell RNA sequencing and histological characterization, with concurrent deletion of *Ifih1*

(Mda5), we provide crucial data to support the causality behind the genetic

determinants of impaired RNA editing and CAD through MDA5 activation. We identify

SMC as a unique mediator of RNA editing and vascular disease and by doing so create

a new paradigm linking human genetics, epigenomics, and vascular biology to CAD.

RESULTS

Distinct expression of immunogenic RNA within vascular SMCs in human atherosclerotic plaque

Activation of MDA5 is dependent on the concentration of specific RNA molecules that are prone to the formation dsRNA structures. By combining our data from experimental *in vitro* models[26] along with human genetic edQTL analysis[12], we generated a putative list of high probability human RNA molecules that are 'immunogenic' and require RNA editing to evade sensing by MDA5. Given the relationship between edQTLs and CAD, we evaluated human scRNAseq data from carotid atherosclerotic plaque[27] to determine if there is a cell type specific expression of these immunogenic RNA. Given that edQTLs that decrease RNA editing increase the risk of CAD, we hypothesized that cells with a higher expression of 'immunogenic' RNAs would be more prone to MDA5 activation if RNA editing was impaired. We evaluated a scRNAseq dataset from human carotid atherosclerotic plaque[27] to determine cell clusters and cell types (Fig 1B). We generated a cell module 'score' that summates the expression of all 'immunogenic' RNA and discovered that human SMCs expressed immunogenic RNA at a markedly higher level compared to all other cell types (Fig 1C). This observation suggests a unique role for SMCs in mediating RNA editing within the vessel wall.

SMC phenotypic modulation and ISG induction in atherosclerosis

The formation of atherosclerotic plaque and vascular calcification involves SMCs undergoing a process of epigenetic reprogramming, termed "phenotypic modulation"[28,29]. In this process, SMCs dedifferentiate, down-regulate lineage marker expression, and migrate into the atheroma where these cells then proliferate and differentiate into different progenies that affect the biology of the plaque. Using SMC lineage tracing mouse models of atherosclerosis with single cell RNA and ATAC sequencing of the aortic root, we have previously shown that numerous CAD risk genes mediate disease risk through modification of this phenotypic modulation, including TCF21, ZEB2, AHR, PDGFD and SMAD3[30-34]. Mature SMCs will undergo phenotypic modulation to a more 'fibroblast like' fibromyocyte (FMC) and then into a calcification promoting chondromyocyte (CMC)[32] phenotype in the setting of disease.

We hypothesized that given SMCs express immunogenic RNA at markedly

higher levels than other cell types, that insufficient RNA editing and dsRNA sensing,

could be identified as expression of key ISGs in the SMC population. To evaluate if

there is ISG induction with SMC phenotypic modulation, we performed a subset

analysis of the SMC population from a human carotid atherosclerotic plaque dataset[27].

Clustering analysis revealed 4 major clusters (cluster '0', '1', '2', and '3') (Fig 1D), where

SMCs undergo a process of downregulation of the mature SMC marker *MYH11* (Fig 1E)

with upregulation of FMC/CMC markers including *TNFRSF11B*, *HAPLN1* and *LUM* (ex.

TNFRSF11B Fig 1F) as SMCs transition from mature SMC cluster '2' to FMC/CMC

cluster '0' — a finding we have previously reported[32]. However, findmarker analysis of

these clusters identified numerous ISGs as top markers of the FMC/CMC cluster '0',

including *ISG15*, *IFI35*, *IFI16*, *IFI27*, *IFIT3* (ex. *ISG15*, *IFI35*, *IFI16* Fig 1G-I). This

finding is suggestive of dsRNA sensing by MDA5 with phenotypic modulation. Top gene

ontology (GO) terms of each cluster highlighted clusters '2' and '1' to be defined by

'smooth muscle contraction', 'muscle system process', and 'actomyosin structure

organization', while cluster '0' is marked by immune mediated processes including

'complement activation' and 'humoral immune response' (Fig 1J).

To investigate if ISG induction with SMC phenotypic modulation is conserved in

mouse models of atherosclerosis, we evaluated our previously published

comprehensive integrated dataset of multiple single cell transcriptomic experiments[35]. In

this dataset, we evaluated transcriptomic profiles in the aortic root of lineage traced

SMCs in hyperlipidemic models of atherosclerosis from multiple publicly available

datasets[35]. We observed that in comparing between 0 and 16 weeks of high fat diet,

SMCs underwent significant phenotypic modulation, marked by downregulation of SMC

contractile marker *Myh11* (Fig 1K), with upregulation of *Col2a1* (Fig 1L), a marker of

chondromyocyte formation. With this phenotypic transition, we observed significant

upregulation of ISGs including *Isg15* (Fig 1M) and *Ifit3* (Fig 1N) in these phenotypically

modified SMCs. These cells lie at the juncture between FMC and CMC formation and

expression was absent in mature SMCs prior to phenotypic modulation.

SMC ADAR1 and MDA5 expression increases in atherosclerosis in mice

To understand the expression patterns of *ADAR* and *IFIH1* (MDA5) in

atherosclerosis SMC phenotypic modulation, we first evaluated the expression of *ADAR*

and *IFIH1* in this human carotid atherosclerosis dataset where we observed minimal

differences in the expression of either *ADAR* or *IFIH1* across the different SMC subsets

(Fig S1A-B). However, upon evaluation of our comprehensive SMC lineage traced

mouse dataset comparing week 0 and week 16, we observe that both *Adar* and *Ifih1*

increase significantly at 16 weeks compared to the 0-week baseline (Fig S1C-E). At the

16-week timepoint, *Adar* expression appears largely consistent across SMC populations

with phenotypic modulation, however *Ifih1* (Mda5) increases significantly at the

FMC/CMC junction as observed with other ISGs (Fig S1D-E).

ADAR1 controls global RNA editing and ISG response in human coronary artery

smooth muscle cells

To investigate ADAR1 in human SMCs, we used an *in vitro* model of phenotypic

modulation with primary human coronary artery smooth muscle cells (HCASMCs) and

knocked down (KD) *ADAR* (ADAR1) and *IFIH1* (MDA5) with siRNA transfection paired

with bulk RNA sequencing (RNAseq). In this model (Fig 2A), we cultured HCASMCs to

confluency, treated with siRNA for 48 hours, and then transitioned to serum free

medium (serum starvation) to induce a mature SMC phenotype marked by high

expression of SMC markers (*e.g. ACTA2*, *CNN1*). Following 48 hours of serum

starvation, cells were stimulated with either serum replete medium or maintained in a

serum free medium for 48 hours. With serum stimulation, HCASMCs subsequently

downregulated expression of mature SMC markers, a phenotype akin to phenotypic

modulation in the SMC to FMC transition in atherosclerosis[32].

With RNAseq data we inferred total RNA editing by detection of G base

sequences at RNA editing sites as we have done previously (Fig 2B)[12,36]. We observed

a notably strong linear relationship between *ADAR* expression and global RNA editing

frequency (r^2=0.9509, *p<0.0001*) (Fig 2B&C). Intriguingly, RNA editing frequency

significantly decreased with serum stimulation alone (Fig 2C).

ADAR siRNA transfection in serum starved conditions induced a significant gene

expression response with 6,351 differentially expressed genes (*FDR <0.05*) (Fig 2D).

Top upregulated genes were interferon stimulated genes (ISGs) including *ISG15, IFI6,*

IFI44L, *IFIT2*, and *IFIT1* (Fig 2D). Importantly, classical interferon genes interferon α, β,

and γ (i.e. *IFNA1*, *IFNB1, IFNG*) were either not expressed (*IFNA1* & *IFNG*), or

minimally affected by *ADAR* KD in an MDA5 dependent mechanism (*IFNB1*) (Fig S2A-

C) with similar minimal effect on the key IFN receptor, *IFNAR1* (Fig S2D).

Top KEGG pathways of upregulated genes indicated a global viral sensing and

inflammatory response, including "TNF signaling pathway", "Influenza A", "NOD-like

receptor pathway", and "EBV infection", as well as "Lipid and atherosclerosis" (Fig S2E).

GO biological processes similarly revealed "Defense Response to Virus" as the top

process (Fig S2F). Downregulated pathways highlighted "Ribosome", "Focal Adhesion",

and "ECM-receptor interaction" (Fig S2G) and GO terms suggested downregulation of

RNA and protein processing with "Cytoplasmic translation" and related terms (Fig S2H).

With transfection of both *ADAR* (ADAR1) and *IFIH1* (MDA5) siRNAs we observed that

co-KD of *IFIH1* blocked the majority of the gene expression response (Fig 2E).

With serum stimulation (representing phenotypic modulation) we observed *ADAR*

KD induced a much greater effect than seen with serum starvation with 12,747

differentially expressed genes (*FDR <0.05*) (Fig 2F). This effect was characterized with

a dominant ISG response and appeared nearly entirely dependent on activation of

MDA5 (Fig 2G). When using principal component analysis (PCA) of bulk RNAseq data

for each treatment group, we observed that the two dominant principal components

tracked with MDA5 activation (PC1) and serum stimulation (PC2), where serum starved

cells with *ADAR* KD produced an outwards shift on PC1, and then returned to near

baseline with co-KD of *IFIH1* (Fig 2H). With serum stimulation, there was an upwards

shift on PC2, and an even greater shift on PC1 with *ADAR* KD, and a return to baseline

with co-KD of *IFIH1* (Fig 2I). These findings highlight that the dominant effect of

impaired RNA editing in HCASMCs is driven by activation of MDA5 and SMC cell state.

ADAR1 and MDA5 regulate SMC maturation and key CAD transcription factors in

phenotypic transition

In this HCASMC model of phenotypic transition, we observed that serum stimulation

downregulated mature SMC markers including *ACTA2* and *CNN1* (Fig 2I&J). However,

we also observed that *ADAR* KD alone downregulated these SMC markers, an effect

that was accentuated with serum stimulation (Fig 2I&J). Using *ISG15* expression as a

marker of ISG activation, *ADAR* KD induced an increase in *ISG15* expression (2K), an

effect that was enhanced with serum stimulation and entirely blocked with co-KD of

IFIH1 (MDA5) (2K). Interestingly, numerous other key transcription factors with known

functions in SMC maturation and risk of CAD (*i.e. KLF4, EGR1*, and *ATF3*)[37] were

similarly affected with *ADAR* KD (Fig 2L-2M).

Global RNA editing is decreased in HCASMC following phenotypic modulation in

vitro

In the evaluation of global RNA editing frequency in HCASMC with phenotypic

modulation (serum stimulation) with control, *ADAR,* or *ADAR* + *IFIH1* KD, we observed

that serum stimulation alone appeared to decrease total RNA editing (Fig 2C). Given we

see serum stimulation to increase MDA5 activation with *ADAR* KD, this suggests a

potential interaction between phenotypic modulation, decreased RNA editing, and

dsRNA formation. To better understand this relationship, we performed a genome wide

RNA editing site specific comparison in serum starved and serum stimulated HCASMC

(Fig S3). In comparing the mean RNA editing level for each RNA editing site in control

siRNA treated cells in serum starved and serum stimulated conditions, we observed a

significant shift where serum stimulated cells have an overall decrease in RNA editing

(Fig S3A). Although no specific RNA editing site met genome wide level significance,

there was a significant overall net effect to decrease RNA editing frequency across

hundreds of RNA editing sites. To evaluate the effect of *ADAR* KD we performed this

editing site specific analysis in the *ADAR* KD with concurrent *IFIH1* KD (minimize the

effect of MDA5 activation on ADAR1 editing) that revealed a massive reduction in RNA

editing with *ADAR* KD in both serum starved and serum stimulated conditions (Fig

S3B&C). From this analysis, we then identified 1,101 differentially edited sites (DES)

across 219 genes (Fig S3D&E) that are ADAR1 dependent. The majority of these DES

are shared between serum starved and serum stimulated conditions (Fig S3D&E),

however we identified 226 sites within 38 genes that are distinct to serum stimulated

conditions, and 357 sites within 49 genes that are distinct to serum starved conditions.

Evaluation of these genes show no consistent biological pattern where gene ontology

enrichment analysis shows no significant pathways (data not shown), however several

genes (*TCTA, FN1, GGCX*) have been identified to reside at loci that meet genome

wide level significance for CAD[5]. These data implicate specific genes that may be

responsible for this enhanced MDA5 activation seen with serum stimulation.

TGFβ upregulates ADAR1 and blunts MDA5 activation in HCASMC

As we observed *ADAR* and *IFIH1* (MDA5) to have distinct effects on HCASMC

response to serum stimulation (a "synthetic" response), we aimed to further understand

how *ADAR* may regulate alternative SMC programming and investigated how *ADAR*

and *IFIH1* may regulate HCASMC response to TGFβ as a 'contractile' response. We

cultured HCASMC to confluency, treated with siRNA for 48 hours, and transitioned cells

to a serum free condition for 48 hours. We then stimulated cells with TGFβ (10ng/mL) or

control for 72 hours before we isolated RNA and performed quantitative PCR. In

comparing the effect of serum stimulation or TGFβ treatment on *ADAR* expression, we

observed that serum stimulation induced a small increase in *ADAR* expression (~1.2

fold) (similar to what we observed in vivo during atherosclerosis, Fig S1), however,

TGFβ treatment produced a greater than 2-fold upregulation of *ADAR* (Fig S4A&B). At

this timepoint, 7 days after siRNA treatment, *ADAR* expression in the *ADAR* KD

treatment was found to have normalized in a manner dependent on *IFIH1* (MDA5),

indicating that MDA5 activation is stimulating a compensatory upregulation of *ADAR*

(Fig S4B). However, when evaluating expression of key ISGs such as *ISG15*, we

observed that TGFβ treatment significantly blunted the effect of *ADAR* KD, where a 25-

fold upregulation of *ISG15* with *ADAR* KD in serum starved conditions was limited to

less than 10-fold upregulation with TGFβ treatment (Fig S4C). In our serum stimulation

study, *ADAR* KD induced a marked upregulation of *IFIH1* (MDA5) that was exacerbated

with serum stimulation in a pattern that was consistent with all other ISGs (Fig S4D).

However, TGFβ treatment markedly reduced this effect of *ADAR* KD (Fig S4E). We

further evaluated expression of *PKR*, a key ISG that acts to control cellular signaling

following MDA5 activation[38]. Here, we found *PKR* to similarly be upregulated by *ADAR*

KD, however this effect is significantly reduced with TGFβ treatment (Fig S4F). This

finding suggests that TGFβ treatment regulates expression of *ADAR* and perhaps

implicates a mechanism of TGFβ to control dsRNA editing by ADAR1 and dsRNA

sensing by MDA5.

ADAR1 and MDA5 regulate the transcriptomic and inflammatory response to

calcification in vitro

To model SMC calcification *in vitro*, we cultured an immortalized HCASMC line under

high calcium/phosphate culture conditions as has been previously reported[39]. In this

model, SMCs begin to form deposits that stain with Alizarin red, consistent with calcium

crystals and has been proposed to represent vascular calcification[39]. Following siRNA

transfection for 48 hours, we cultured cells for 7 days with pro-calcification medium

under low serum (1%) conditions and performed bulk RNAseq. After 7 days,

calcification deposits became visible, however with *ADAR* KD, there was a significant

phenotypic change with increased cell death and increased density of calcium deposits

— an effect that was blocked with co-KD of *IFIH1* (Fig S5A&B). Calcification medium

under scramble siRNA treatment promoted a significant gene expression response with

584 differentially expressed genes (*FDR <0.05*) (Fig S3C). Top upregulated genes

include numerous BMP related signaling genes including *FBLN5*, *FMOD*, and

RANBP3L (Fig S5D-F), effects that have been previously reported[39]. However, with

ADAR KD this effect was blunted in an MDA5 dependent mechanism (Fig S5D-F).

Calcification did not overtly activate MDA5 as measured by expression of *ISG15*

(Fig S5G&H), however, *ADAR* KD modified the HCASMC inflammatory response to

calcification. Calcification had minimal effect on expression of NFκB (*NFKB1*), however

with *ADAR* KD *NFKB1* is modestly upregulated and is further upregulated with

calcification in *ADAR* KD cells, an effect blocked with co-KD of *IFIH1* (Fig S5I). This

observation revealed a unique relationship where calcification induced a distinct

proinflammatory effect only with concurrent *ADAR* KD and MDA5 activation. Similar

effects were observed for numerous proinflammatory signaling and matrix proteins

including *JUND*, *MMP1*, and *MMP10* (Fig S5J-L).

SMC specific Adar1 maintains vascular integrity and survival in vivo

Although prior reports have implicated SMC specific RNA editing in mediating

phenotypic response[40,41], the relationship between deficient SMC RNA editing, MDA5

activation, vascular inflammation, and atherosclerosis has not been investigated. To

understand how homozygous deletion of *Adar1* within SMCs influence vascular

homeostasis, we bred a mouse model with four distinct alleles, including *Adar1*[fl/fl] (exon

7-9), a tamoxifen inducible SMC specific Cre-recombinase (*Myh11*[CreERT2]), a lineage

tracing fluorophore (*ROSA*[tdTomato]), on a hyperlipidemic pro-atherosclerosis genetic

background (*ApoE*[-/-]) (Fig 3A). A total of 56 mice were treated, comparing *SMC-Adar1*[-/-],

SMC-Adar1[-/+] (*Adar1*[fl/WT], *Myh11*[CreERT2], *ROSA*[tdTomato], *ApoE*[-/-]), and control (*Adar1*[WT/WT],

Myh11[CreERT2], *ROSA*[tdTomato], *ApoE*[-/-]) genotypes. We observed that at day 4, *SMC-*

Adar1[-/-] mice became notably ill as characterized by weight loss, piloerection, and a

hunched posture, a phenotype that was not present in *SMC-Adar1*[-/+] or control genotype

animals. We observed mortality in the *SMC-Adar1*[-/-] mice at this time and by 14 days,

10/20 (50%) mice had died (Fig 3B). A recent report while this manuscript was in review

has further supported this role of SMC specific *Adar1* to maintain survival and vascular

integrity[42]. Gross dissection and visual inspection of *SMC-Adar1*[-/-] mice did not reveal

prominent findings of the lungs, heart, or intestines however visual appearance of the

aorta suggested petechial hemorrhage and erythema (Fig 3C). H&E staining (Fig

3D&E), fluorescent imaging (Fig 3F&G), bright field imaging (Fig 3H), Cd68

immunostaining (Fig 3I), and Masson's trichrome staining (Fig 3J) of ascending and

descending aortic tissue in 6 *SMC-Adar1*[-/-] mice revealed a loss of elastin organization

(present in 6/6 mice), dissociation of the matrix at the adventitial aspect with associated

'ballooning' out and separation of layers (present in 3/6 mice), with evidence of

intramural hemorrhage (present in 2/6 mice) (Fig 3D-H, Fig S6). These sites of

ballooning occurred with complete loss of tdTomato SMCs at the adventitial aspect (Fig

3F&G). H&E staining of the aorta confirmed severe loss of organization of the elastin

matrix and evidence of intravascular hemorrhage with further discovery of dense cellular

infiltrate into the adventitia (Fig 3D&E). These patterns were not observed in control

mice (Fig 3E). Immunostaining for CD68 revealed macrophage cell infiltration from the

luminal as well as the adventitial aspect of the vessel wall (Fig 3I). Additional Masson's

trichrome stain to evaluate for fibrosis revealed evidence of intra-medial collagen

deposition (Fig 3J).

In this model, there was a rapid and early inflammatory response that resulted in

profound weight loss and mortality. Although the cause of death was not immediately

apparent, as gross examination and H&E sections of liver, lung, and heart were

nonrevealing, the acute systemic inflammatory response is likely eliciting death through

a multifactorial mechanism akin to a systemic inflammatory response. There was no

evidence of aortic rupture as a cause of death. However, mice that survived this period

remained underweight and ill in appearance until ~7 months of age. Importantly, no

mice that were taken past this 2-week timepoint regained weight or lived a normal

lifespan. Dissection of mice at this 7-month timepoint demonstrated dense fecal

impaction of the large intestine, consistent with bowel obstruction, a finding not seen at

earlier timepoints.

Haploinsufficiency of MDA5 is sufficient to protect against mortality with SMC

Adar1$^{-/-}$ mice

Our *in vitro* work indicates MDA5 to be the principal mediator of downstream signaling

with ADAR1 KD in SMC, however it remains unclear if this relationship is true *in vivo*. In

addition, it is unclear if only partial inhibition of MDA5 can prevent overt ISG activation,

a model more akin to what might be achieved with future pharmacologic inhibition. To

test this, we bred the SMC *Adar1* KO mouse onto a constitutive KO of *Ifih1* (Mda5) on

both a heterozygous and homozygous background generating mice that are *SMC-*

Adar1$^{-/-}$ + Mda5$^{-/+}$ (*Adar1$^{fl/fl}$, Ifih1$^{-/+}$, Myh11^{CreERT2}, ROSAtdTomato, ApoE$^{-/-}$*) and *SMC-*

Adar1$^{-/-}$ + Mda5$^{-/-}$ (*Adar1$^{fl/fl}$, Ifih1$^{-/-}$, Myh11^{CreERT2}, ROSAtdTomato, ApoE$^{-/-}$*). Following

the same model as shown previously (Fig 3), we performed an additional experiment

using litter-mate controlled mice with tamoxifen treatment for a total of 32 mice (12

SMC-*Adar1$^{-/-}$*, 8 *SMC-Adar1$^{-/-}$ + Mda5$^{-/+}$*, and 12 *SMC-Adar1$^{-/-}$ + Mda5$^{-/-}$*). Mice that were

homozygous SMC *Adar1* KO with MDA5 WT had severe weight loss and significant

mortality, with 10/12 (83%) *SMC-Adar1$^{-/-}$* mice dying prior to the 14-day timepoint and

greater than 30% loss of total body weight by 7 days for the 6 mice that survived to the

7-day timepoint (Fig 4A-C). Both *SMC-Adar1$^{-/-}$ + Mda5$^{-/+}$* and *SMC-Adar1$^{-/-}$ + Mda5$^{-/-}$*

mice showed minimal weight loss and no mortality, and visually appeared largely

unaffected by *Adar1* KO at this 2-week timepoint (Fig 4A-C). *SMC-Adar1*[-/-] *+ Mda5*[-/+]

mice have been taken out to 24 weeks following tamoxifen dosing and continue to show

minimal phenotype, demonstrating that MDA5 is required to initiate this severe

phenotype with SMC *Adar1* KO (Fig 4D). Histological evaluation of aortic tissue from

SMC-Adar1[-/-] *+ Mda5*[-/+] mice show that aortic cross section, aortic media, and elastin

structures are indistinguishable from control mice (Fig S7). Intriguingly, a portion of both

SMC-Adar1[-/-] *+ Mda5*[-/+] and *SMC-Adar1*[-/-] *+ Mda5*[-/-] mice (2/8 and 4/12, respectively) did

lose significant weight within the first week, however had significant weight recovery by

week 2 (Fig 4C). This suggests that although MDA5 deletion protects against mortality

and any major phenotype, that alternative dsRNA receptors such as RIG-I may be

mediating a partial inflammatory response.

Single cell RNA sequencing reveals SMC Adar1 controls ISG activation,

phenotypic response, and macrophage infiltration into the vessel wall

To further characterize the vascular response of SMC specific *Adar1* KO in mouse, we

performed single cell RNA sequencing (scRNAseq) of the aortic root and ascending

aorta of *SMC-Adar1*[-/-] mice at day 14 following tamoxifen dosing. With a total of 10

individual scRNAseq captures, we characterized over 50,000 cells with high sequencing

depth (>50K reads per cell). Using dimensionality reduction with UMAP as we have

done previously[43,44], we observed *SMC-Adar1*[-/-] aortic cells to have a dramatic

transcriptomic change in comparison to control genotype (Fig 5A). Cluster analysis

identified 13 main clusters, including 3 SMC clusters — a control SMC cluster marked

by mature contractile markers (i.e. *Myh11* and *Cnn1*), and two SMC clusters that

represented cells with distinct transcriptomic signatures secondary to SMC *Adar1* KO

that are marked by tdTomato (clusters SMC_cAdar1_KD1; SMC_cAdar1_KD2) (Fig

5B&C). Genes expressed by cells in these two clusters were found to show gene

ontology (GO) enrichment for terms related to 'response to interferon beta', 'response to

virus', 'defense response to virus' as top GO terms (Fig S8). Using *Isg15* as a marker of

MDA5 activation, clusters SMC_cAdar1_KD1 and SMC_cAdar1_KD2 were marked by

notably high expression of *Isg15* (Fig 5D). To validate this finding and characterize the

location of *Isg15* expression, we performed RNAscope on aortic tissue from *SMC-*

Adar1^{-/-} mice. We observed robust *Isg15* expression within the vessel wall that was

highest at the luminal aspect of the media, with additional expression within the

adventitia (Fig 5E).

Within tdTomato lineage traced SMCs, conditional *Adar1* KO produced marked

gene expression changes with 2,382 differentially expressed genes (*FDR<0.05,*

log2FC>0.1), with notable increase in ISGs. In comparing clusters SMC_cAdar1_KD1

and SMC_cAdar1_KD2, SMC_cAdar1_KD2 was marked by higher expression of

numerous inflammatory signaling genes (i.e. *Isg15*, *Ccl5*, *Cxcl10*) (Fig S9A).

Importantly, consistent with our *in vitro* findings (Fig S2A-D), there was only

modest activation of classical interferon genes with no observed expression of *Ifna1* and

minor activation of *Ifnb1* (Fig 5F). Similarly, we observed minimal expression of IL6

family cytokines (i.e. *Il6*) with *SMC-Adar1*^{-/-} (Fig S8B). The minimal expression of these

markers appeared limited to the more proinflammatory SMC_cAdar1_KD2 cluster. In

addition, consistent with our *in vitro* findings, we observed that SMC *Adar1* KO led to

downregulation of mature SMC markers (Fig S9C). By comparing the three major SMC

clusters, expression of *Cnn1* and *Myh11* decreased in a pattern consistent with

increasing MDA5 activation (Fig S9C).

There was a new distinct fibroblast population seen only with SMC *Adar1*^{-/-} and is

marked by ISGs and high expression of *Ly6a* (Sca1) (Fig 5B, Fig S9D), consistent with

a proinflammatory adventitial fibroblast population[45]. Similarly, in SMC *Adar1*^{-/-} there

was a loss of cells within the 'Fibro_1' cluster that was only seen in controls (Fig 5B),

suggesting a dynamic shift in gene expression from resident fibroblasts to this new

reparative and inflammatory cell cluster. In addition, there was a large infiltrating

macrophage population marked by high *Cd68* (resident macrophage cluster – 'Mac',

infiltrating macrophage cluster – 'Mac_2') (Fig 5B, Fig S9E) with an additional small T

cell population with SMC *Adar1*^{-/-} marked by high *Gzma*, *Cd3g*, and *Cd52*, suggestive of

a cytotoxic T lymphocyte (CTL) population (Fig 5B, Fig S9F). These data indicate that

with loss of *Adar1* within SMCs, a profound vascular response occurs throughout the

vessel wall that is coordinated by SMCs.

Receptor-ligand interaction modeling implicates CCL5:CCR5 to coordinate

macrophage infiltration in SMC-Adar1$^{-/-}$ mice

To better understand the cell-cell communications that mediate this response, we performed an inference analysis of cell-cell communication focused on receptor-ligand interactions (CellChat)[46]. This analysis revealed that with SMC *Adar1$^{-/-}$*, SMC clusters SMC_cAdar1_KD1 and SMC_cAdar1_KD2 are notably high outward senders of CCL signaling markers (Fig 5G). When interrogating the specific signaling pairs, Ccl5:Ccr5 was the dominant interaction (Fig 5H). A circos plot depicting Ccl5:Ccr5 interaction strength revealed dominant outward interaction arising from SMC_cAdar1_KD1 and SMC_cAdar1_KD2 clusters with key receiving clusters being resident macrophage, infiltrating macrophage, and T cell populations (Fig 5I). Feature plots of Ccl5 and Ccr5 further revealed the distinct expression of Ccl5 within SMCs following *Adar1* KD and Ccr5 within macrophage populations (Fig 5J&K). Evaluation of the next top 3 interaction pairs (Ccl2:Ccr2; Ccl4:Ccr5, Ccl7:Ccr2) suggested that following Ccl5:Ccr5 interaction, Ccl2 and Ccl7 arising from SMC_cAdar1_KD2, Fibro_Sca1_hi, and macrophage cell populations appeared to interact with Ccr2 within macrophage and T lymphocyte populations (Fig S9G-L). These data suggested that Ccl4 from resident and infiltrating macrophage clusters further signaled through Ccr5 on these same cell clusters (Fig S9J&K). Activation of CCR5 has been previously identified as a key regulator of antiapoptotic signaling in macrophage cells in response to viral infection[47]. Our observation provides a unique insight that following the loss of SMC *Adar1*, Ccl5:Ccr5 acts as a dominant regulator of macrophage infiltration and macrophage cell survival.

Homozygous deletion of MDA5 (Ifih1) prevents transcriptomic effect of SMC-

Adar1 KO seen with single cell RNA sequencing

Our data indicates that loss of SMC *Adar1* elicits severe vascular effect and mortality due to dsRNA activation of MDA5 (Fig 4). Further, our in vitro HCASMC data suggests that loss of *ADAR1* with concurrent MDA5 KD produces only a minor effect on cellular transcriptome (Fig 2). However, it is unclear if this is true in vivo and there remains question as to if loss of RNA editing by ADAR1 may mediate significant transcriptomic effect despite concurrent MDA5 deletion. To investigate this, we performed additional

scRNAseq captures in 9 *SMC-Adar1^{-/-} + Mda5^{-/-}* mice at this same timepoint of 14-days

post tamoxifen as previously performed, for a total of 6 scRNAseq captures. Data was

integrated together with control and *SMC-Adar1^{-/-}* captures (16 total sequencing

libraries). Remarkably, UMAP visualization of single cell transcriptomic data from *SMC-*

Adar1^{-/-} + Mda5^{-/-} mice show nearly identical pattern compared to control (S10A).

Infiltrating macrophage and T cell populations and Sca1-hi fibroblast populations are no

longer present. When evaluating expression of key ISGs such as *Isg15* and *Ccl5*,

homozygous *Mda5* (*Ifih1*) KO prevents the entirety of ISG induction across all cell types

(S10B-C) and when analysis is focused on tdTomato clusters along (S10D-E). We

interrogated expression of specific SMC marker genes such as *Myh11* and *Cnn1* in

tdTomato lineage traced cells, where we observe that downregulation of contractile

markers seen with *SMC-Adar1* KO is prevented with *Mda5* KO (S10F-G), indicating this

effect is due to activation of MDA5. *Findmarker* differential gene expression analysis

and heatmap visualization of marker genes from the three genotypes show near

identical patterns between control and *SMC-Adar1^{-/-} + Mda5^{-/-}* mice (S10H). These data

demonstrate most effect of *SMC-Adar1* KO is secondary to MDA5 activation. A small

gene set where expression is observed to be downregulated in both *SMC-Adar1^{-/-}* and

SMC-Adar1^{-/-} + Mda5^{-/-} mice compared to control (S10H), suggesting that a small

fraction of genes may be downregulated with loss of SMC Adar1 RNA editing

independent of MDA5. A second set of genes appear to be downregulated with *SMC-*

Adar1^{-/-} however are increased with *SMC-Adar1^{-/-} + Mda5^{-/-}* (S10H), suggesting a role

for MDA5 in regulating the expression of these genes independent of *Adar1*.

SMC specific haploinsufficiency of Adar1 activates the ISG response in

atherosclerosis and increases vascular chondromyoctye formation

Although conditional homozygous deletion of *Adar1* in SMCs caused a profound and

rapid onset phenotype *in vivo* in a manner dependent on Mda5, we aimed to understand

if haploinsufficiency of *Adar1* in SMCs, where mice appear to have no discernable

phenotype, may lead to MDA5 activation, SMC phenotypic modulation, and progression

of atherosclerosis. To do this, we used the *SMC-Adar1^{-/+}* (*Adar1^{fl/WT}*, *Myh11^{CreERT2}*,

ROSAtdTomato, *ApoE^{-/-}*) mice with genotype controls (*Adar1^{WT/WT}*, *Myh11^{CreERT2}*,

ROSAtdTomato, *ApoE⁻ᐟ⁻*), dosed with tamoxifen at 7.5 weeks of age, and placed on high

fat diet for 16 weeks (Fig 6A). In this model, mice developed pronounced

atherosclerosis as we have shown previously[31,32,43,44]. *SMC-Adar1⁻ᐟ⁺* and control mice

had no differences in weight or lipid analysis (cholesterol, LDL, HDL, triglycerides) at

this 16 week timepoint (Fig S11). To understand the cellular resolution of the vessel

wall, at 16 weeks, we performed scRNAseq of the aortic root and ascending aorta. With

a total of 12 scRNAseq captures analyzing over 70,000 cells comparing *SMC-Adar1⁻ᐟ⁺*

and control genotypes, we observed expected distinct cell clusters of SMCs, fibroblasts,

endothelial cells, macrophage cells, and T cells (Fig 6B). Lineage traced SMCs

demonstrated distinct SMC clusters as we have previously observed and have been

consistent with phenotypic modulation[32,43,44], with a mature SMC cluster, a fibromyocyte

(FMC) cluster, and a calcification promoting chondromyocyte (CMC) cluster of cells (Fig

6B). Importantly, *SMC-Adar1⁻ᐟ⁺* mice showed a large and distinct new SMC population

residing between the mature SMC population and the FMC population (Fig 6B).

Characterization of this population by *findmarker* analysis demonstrated this cluster to

express numerous markers of ISG activation (i.e. *Ifit3*, *Ifit1*, *Isg15*) consistent with MDA5

activation. Analysis of *Isg15* as a marker of ISG activation revealed notable expression

within this SMC cluster, termed 'SMC_ISG' (Fig 6B&C).

To focus on the lineage traced SMC population, we performed an analysis on the

'SMC', 'SMC_ISG', 'FMC', and 'CMC' clusters. With subsequent dimensionality

reduction and repeat clustering, UMAP visualization highlighted the marked effect of

SMC *Adar1* haploinsufficiency in SMC phenotypic states in atherosclerosis (Fig 6D&E).

Top 10 marker genes and GO term enrichment for 'SMC', 'SMC_ISG', 'FMC', and

'CMC' clusters reveals the SMC_ISG cluster to be marked by ISGs including *Ifit3, Isg15,*

Ifit1, and *Irf7* and GO terms highlighting "response to virus", "response to interferon-

beta", and "defense response to virus" (Fig S12A&B). Evaluation of SMC marker *Myh11*

demonstrated downregulation of mature contractile markers with phenotypic modulation

(Fig S13A) with associated upregulation of FMC and CMC markers (i.e. *Tnfrsf11b*,

Col2a1, and *Acan*) (Fig S13B-D). Separation by genotype further highlighted that with

SMC-Adar1⁻ᐟ⁺ mice, there was a shift away from mature SMCs, marked increase in the

SMC_ISG cluster, and an increase in CMC (Fig 6E&F). A featureplot of *Isg15*

expression in SMC subsets highlighted a possible relationship between the 'SMC_ISG'

and 'CMC' clusters (Fig 6G). Comparison of the expression of CMC markers *Col2a1*

and *Acan* between genotypes demonstrated an increase in CMC cells with *SMC-Adar1⁻*

⁄⁺ mice consistent with this finding (Fig S13E&F). This finding suggests that ISG

activation from SMC *Adar1* haploinsufficiency in atherosclerosis may be contributing to

CMC formation and vascular calcification.

Lineage trajectory analysis of vascular chondrogenesis reveals a distinct ISG

dependent trajectory

To investigate the potential for this SMC lineage relationship, we performed

pseudotime inference and cellular trajectory analysis with Slingshot[48]. By increasing

clustering resolution we generated 14 high resolution clusters of all tdTomato lineage

traced SMCs and performed Slingshot inference to understand the cellular trajectory

between mature SMC to CMC clusters. In this analysis, we observed only 2 distinct

trajectories between SMC and CMC (Fig 6H). Importantly, one trajectory excludes all

SMC_ISG clusters, while a second is dependent on SMC_ISG clusters (Fig 6H). This

suggests two distinct cellular trajectories to CMC formation, with one trajectory

dependent on ISG activation, while a second is ISG independent. This observation of an

ISG dependent trajectory to CMC formation with increased CMC formation with SMC

Adar1 haploinsufficiency in atherosclerosis implicates a disease context specific

activation of MDA5 with inadequate RNA editing and disease progression.

To better understand the top genes driving this ISG dependent trajectory, we

performed findmarker analysis on all high-resolution clusters and evaluated the top

markers for each cluster in sequence of the ISG dependent pathway (Fig S14A). This

analysis highlights that as SMC lose expression of contractile markers such as *Myh11*

and *Acta2*, they begin to express numerous ISGs. Initially, there is high expression of

Irf7, *Isg15*, *Ifit1*, which is then followed by increasing expression of *Ly6a*, *C3*, *Cxcl12*,

and other FMC markers such as *Dcn* and *Tnfrsf11b*, before SMCs have marked

expression of key CMC genes such as *Col2a1*, *Chad*, *Fmod*, and *Spp1* (Fig S14A). To

further understand the pathways driving this ISG dependent trajectory, we selected the

gene markers of the high-resolution clusters that are distinct to the ISG trajectory and

performed gene ontology pathway analysis (Fig S14B). Here, we identified top viral

immune response pathways including *'Coronavirus Disease'*, *'Influenza A'*, and

'Epstein-Barr Virus infection', driven by genes such as *Jun, Ccl2*, *Il6*, *Nfkb1*, and this

analysis also reveals key JAK/STAT signaling molecules such as *Stat3* and *Jak1* (Fig

S14B). In addition to immune pathways, *'Focal Adhesion'* and *'Regulation of Actin*

Cytoskeleton' are highlighted, driven by genes such as *Rock1*, *Itgb1*, *Egfr,* PDGFs, and

Map2k1, suggesting a distinct interrelationship between activation of JAK/STAT and key

immune signaling pathways influencing focal adhesion pathways (Fig S14B).

To determine the location of MDA5 activation within the plaque and vessel wall,

we performed RNA scope on aortic root sections for *Isg15*. In control mice, we observed

scattered *Isg15* expression within the media, the base of the plaque and extending up to

the region beneath the cap, as well as in the adventitia (Fig 6I). However, in *SMC*

Adar1$^{-/+}$ mice we observed a marked increase in *Isg15* signal, with intense *Isg15*

expression within the media, throughout the plaque, and within the adventitia (Fig 6J).

Signal of *Isg15* RNAscope positive area in the vessel wall was quantified (Fig S15A)

and location of *Isg15* signal was compared to signal of chondromyocyte marker *Col2a1*,

where we reveal that cells that express *Isg15* reside in spatially distinct regions

compared to *Col2a1* (Fig S15B-C). These data suggest ISG activation in

atherosclerosis in control mice, however this effect is markedly increased with

haploinsufficiency of *Adar1* within the vascular SMC.

SMC specific haploinsufficiency in Adar1 increases plaque size and calcification

in atherosclerosis

We performed histological analysis of the aortic root for *SMC-Adar1$^{-/+}$* and control

genotype mice after 16 weeks of high fat diet (n=13 each group). Using fluorescent

imaging of tdTomato and autofluorescence (elastin), we quantified plaque area and

plaque area normalized to total area of the internal elastic lamina area (IEL) (Fig 7A-D).

We observed that *SMC-Adar1$^{-/+}$* mice had a significant increase in plaque area, an

effect that remained following normalization to overall area of the IEL (Fig 7A-D).

Notably, quantification of the area of the media (area between IEL and external elastic

lamina, EEL), we observed *SMC-Adar1$^{-/+}$* mice to have a significant reduction, an effect

that remained following normalization to the total area of the EEL (Fig 7E&F). When

plaque area was normalized to media area, the observed effect with *SMC-Adar1$^{-/+}$* mice

was further enhanced (Fig 7G). In addition, we observed total area of the EEL to be

increased (Fig 7H), suggesting an increased outward remodeling of the plaque. Given

our observation from single cell data that suggested increased chondromyocyte

formation, we evaluated the effect of SMC *Adar1* haploinsufficiency on calcification by

Ferangi Blue Chromogen staining (Fig 7I-L). We observed *SMC-Adar1$^{-/+}$* mice to have a

significant increase in total Ferangi Blue staining by area (Fig 7K), an effect that

remained with normalization to plaque area (Fig 7L). These findings demonstrate that

SMC *Adar1* haploinsufficiency accelerate the formation atherosclerotic plaque area and

calcification.

SMC specific haploinsufficiency of Adar1 increased SMC lineage traced cell

content in plaque without a change in acellular area

Our data suggests that impaired RNA editing in vascular SMC leads to increased

plaque area and decreased media thickness, implicating increased SMC migration out

of the media and into the plaque. To investigate this, we quantified tdTomato positive

area within plaque as a marker of SMC lineage traced cells integrating into the plaque.

Here, we observed that in *SMC Adar1$^{-/+}$* mice, the percent tdTomato area within plaque

is increased compared to control mice (p=0.0024)(Fig S16A-B). To then understand if

this increased SMC migration into the plaque results in a similar increase in tdTomato

area within the cap of the plaque, we measured the tdTomato area of the top 30μm

segment of the cap (Fig S16C). Here, we did not observe a significant difference,

although there was a greater mean tdTomato area in the SMC Adar1$^{-/+}$ mice

(p=0.0749).

To evaluate if there is increased acellular area within the plaque, we performed

Masson's Trichrome stains on lesions and quantified acellularity by tracing the plaque

and subtracting all area that has stain to then be able to quantify all acellular area. Here,

when normalizing acellular area by plaque area, we observed no difference between

genotypes (Fig S16D-E).

SMC specific haploinsufficiency in Adar1 has minimal effect on macrophage content in atherosclerosis

With a subsetted analysis on all non-tdTomato lineage traced cell populations, we observed that *SMC-Adar1*$^{-/+}$ mice have a small increase in total macrophage cell populations in this 16-week atherosclerosis model (39.7% vs 35.2% of all non-tdTomato cell population) (χ^2 *p=3.66E-26*) with minimal change in cell numbers across all other cell types (Fig S17A&B). As we have shown that CCL signaling and Ccl5:Ccr5 appeared to be the dominant driver of macrophage recruitment in *SMC-Adar1*$^{-/-}$ mice, we investigated if these signaling molecules were similarly upregulated in this 16-week *SMC-Adar1*$^{-/+}$ atherosclerosis model. When performing receptor-ligand interaction analysis using CellChat[46], we observed the 'SMC_ISG' cluster to have absent CCL signaling and did not appear to act as a sender, receiver, mediator, or influencer (Fig S17C&D). We then performed Cd68 immunohistochemical staining on aortic root sections to evaluate for Cd68 positive area within the plaque between control and *SMC-Adar1*$^{-/+}$ mice (Fig S17E-I). We observed that the total Cd68 positive area within the plaque was significantly greater in the *SMC-Adar1*$^{-/+}$ mice (Fig S17G), however this effect only trended towards significance when normalized to plaque area (Fig S17H) or area of the total vessel at the internal elastic lamina (Fig S17I). The lack of a major effect on macrophage content at this timepoint for this 16-week atherosclerosis *Adar1* haploinsufficiency model is in contrast to our 2-week SMC-*Adar1* KO model (Fig 3) and likely reflects a 'dose-dependent' response of SMC *Adar1* gene deletion, MDA5 activation, and *Ccl5* expression.

Haploinsufficiency of Mda5 prevents ISG activation and chondromyocyte formation in atherosclerosis with SMC-Adar1$^{-/+}$

Homozygous as well as heterozygous deletion of *Mda5* (*Ifih1*) prevents mortality seen with *SMC-Adar1* KO (Fig 4) and homozygous deletion of *Mda5* (*Ifih1*) prevents the entirety of ISG activation in our single cell RNA sequencing analysis (Fig S10), raising the question as to if deletion of *Mda5* can similarly prevent ISG induction and prevent CMC formation observed with atherosclerosis with *SMC-Adar1* haploinsufficiency. To investigate this, we further bread *SMC-Adar1*$^{-/+}$ mice onto *Mda5* (*Ifih1*) heterozygous

background (*Adar1*^{fl/WT}, *Ifih1*^{-/+}, *Myh11*^{CreERT2}, *ROSAtdTomato*, *ApoE*^{-/-}) mice and

performed additional 16-week HFD atherosclerosis studies following tamoxifen dosing

at 7.5 weeks following our established protocol (Fig 6). We chose to use the *Mda5*

(*Ifih1*) heterozygous background over homozygous deletion given haploinsufficiency is

more akin to what may be achieved from future therapeutic inhibition. We performed

additional single cell RNA sequencing on the aortic root and ascending aorta with 9

mice for a total of 6 additional scRNAseq captures. Following data integration with a

total of more than 96K cells with >50,000 reads per cell (18 total scRNAseq libraries),

we evaluated how *Mda5* haploinsufficiency affects cellular phenotype and lineage

trajectory during atherosclerosis compared to control and *SMC-Adar1*^{-/+} genotypes. With

UMAP visualization, we demonstrate near identical overlap between the control and

SMC-Adar1^{-/+} + *Mda5*^{-/+} genotypes (Fig 8A), suggesting Mda5 haploinsufficiency to

largely prevent global transcriptomic effect of SMC *Adar1* haploinsufficiency. With

evaluation of key ISGs such as *Isg15*, we observe that *Mda5* haploinsufficiency

prevents the entirety of ISG induction (Fig 8B). With SMC subset analysis and

visualization of SMC clusters ('SMC', 'SMC_ISG', 'FMC', and 'CMC') we observe a clear

effect where *SMC-Adar1*^{-/+} + *Mda5*^{-/+} genotype eliminates the 'SMC_ISG' cluster seen

with *SMC-Adar1*^{-/+} mice (Fig 8C), with further reduction in CMC formation (Fig 8C).

Upon evaluation of key marks of SMC, SMC_ISG, FMC, and CMC populations, we

observe that *Mda5* haploinsufficiency decreases the expression of CMC markers such

as *Col2a1* and *Acan* (Fig 8D-E) with near elimination of expression of 'SMC_ISG'

marker *Isg15* (Fig 8F). In quantifying the proportion of cells that belong to each SMC

cluster in the control, *SMC-Adar1*^{-/+}, and *SMC-Adar1*^{-/+} + *Mda5*^{-/+} genotypes, we

observe that the control and *SMC-Adar1*^{-/+} genotypes have 17% and 24% CMCs

whereas the *SMC-Adar1*^{-/+} + *Mda5*^{-/+} genotype significantly reduces this proportion to

12% (Fig 8G). This significant reduction in CMC population in the *SMC-Adar1*^{-/+} + *Mda5*⁻

^{/+} genotype is significantly lower than not only the *SMC-Adar1*^{-/+} genotype, but also the

control genotype (Fig 8G). This may suggest that even in control genotype, Mda5 is

being activated and contributing to CMC formation. Differential findmarker gene

expression analysis in the tdTomato population and heatmap visualization shows how

the majority of marker genes increased in *SMC-Adar1*^{-/+} genotype compared to control

are further decreased in the *SMC-Adar1$^{-/+}$ + Mda5$^{-/+}$* genotype, suggesting the majority

of gene expression effect is dependent on Mda5 (Fig 8H). However, similar to what we

had observed in the SMC Adar1 KO + Mda5 KO study (Fig S10), a portion of genes

appear to be regulated by Adar1 independent of Mda5 and similarly a portion of genes

appear to be regulated by Mda5 independent of Adar1 (Fig 8H). Upon evaluation of top

ISGs (ie. *Isg15*, *Ifit3*, *Irf7*, *Stat1*), we observe a consistent pattern of increased

expression in *SMC-Adar1$^{-/+}$* genotype compared to control which is then blocked in

entirety with the *SMC-Adar1$^{-/+}$ + Mda5$^{-/+}$* genotype (Fig 8I). However, importantly, when

evaluating key markers of SMC phenotypic transition and FMC/CMC formation (ie.

Spp1, *Fn1*, *Tnfrsf11b*, *Dcn*) we see a baseline moderate level of expression in the

control group, a significant increase in the *SMC-Adar1$^{-/+}$* genotype, and then a decrease

in expression to a level lower than the control in the *SMC-Adar1$^{-/+}$ + Mda5$^{-/+}$* genotype

(Fig 8J). This is strong evidence to suggest that Mda5 is playing a significant role in

regulating SMC phenotypic transition in atherosclerosis even with genetically intact

Adar1 RNA editing, consistent with our noted finding of increasing ISG activation with

SMC phenotypic transition in both human and mouse in atherosclerosis (Fig 1) and

human genetics data.

DISCUSSION

Both academic and industry leaders are leveraging the power of human genetics to

drive discovery in the goal to define causal disease biology[49]. The rapid pace of

identifying novel genomic loci that meet genome wide level significance for complex

disease has resulted in notable challenges of target prioritization within a locus[50], as

well as prioritization for therapeutic drug development[51]. However, the association of

edQTL and CAD risk, where impaired RNA editing increases CAD risk while increasing

ISG activation, implicates a common disease pathway that is largely dependent on one

gene product — MDA5. Strikingly, 20% of CAD risk loci examined colocalized with

edQTL loci[12]. With a new understanding of the genetic determinants of RNA editing and

dsRNA sensing as an underlying etiology of coronary artery disease and other

inflammatory disorders[12], there is a significant need to define how this mechanism

influences disease risk.

Here, we provide a crucial link between the human genetics that implicate RNA

editing and dsRNA sensing by MDA5 in atherosclerosis and the biological

underpinnings of this disease association. We reveal a unique aspect of biology, where

the vascular SMC has a distinct requirement for ADAR1 mediated RNA editing to evade

dsRNA sensing by MDA5. Analysis of human atherosclerotic plaque scRNAseq data

indicate a pattern where SMCs express immunogenic RNA at high levels and that ISG

activation — suggestive of MDA5 activation with inadequate RNA editing — occurs with

SMC phenotypic modulation in both human and mouse data. We further show that

deletion of SMC specific *Adar1* in adult mice results in loss of vascular integrity and

significant mortality due to activation of MDA5 (Figs 3 and 4). Further, scRNAseq

analysis of aortic tissue reveals distinct ISG activation and a complex transcriptional

response and inflammatory cell recruitment within the vessel wall (Fig 5), an effect

blocked in its near entirety with deletion of MDA5 (Fig S10). This highlights the absolute

requirement for *Adar1* within the vascular SMC, but its role appears to be primarily to

prevent dsRNA activation of MDA5. Importantly, in 16-week atherosclerosis studies in

Adar1 haploinsufficient mice, scRNAseq reveals ISG activation in SMCs in a cell type

and context specific mechanism driving phenotypic modulation, vascular

chondromyocyte formation, and to a lesser extent macrophage recruitment (Fig 6), and

that this coincides with increased plaque area, decreased thickness of the media, and

increased calcification (Fig 7). This effect of ISG activation and chondromyocyte

formation in SMC *Adar1* haploinsufficiency is prevented with only haploinsufficiency of

Mda5, suggesting the importance of Mda5 in mediating this effect (Fig 8). This indicates

that SMC *Adar1* haploinsufficiency and Mda5 activation facilitates thinning of the media

while increasing plaque area, increased SMC migration into the plaque, and vascular

calcification.

These findings highlight an important new concept, that the need for RNA editing

to evade MDA5 activation not only occurs in a cell type specific manner, but a cellular

context specific manner. Two prior studies have investigated the role of SMC *Adar1* in

haploinsufficiency in SMC phenotypic modulation, focused on carotid injury[41] and an

angiotensin II model of abdominal aortic aneurysm (AAA)[40]. In these reports, the

authors reported that loss of SMC Adar1 prevented phenotypic modulation, however no

investigation into potential MDA5 activation was performed. Our data reported here

appears in contrast to this finding, however our findings implicate that the effect of loss

of ADAR1 on phenotypic transition is dependent on MDA5 activation and occurs in a

'dose dependent' mechanism where increasing MDA5 activation has a greater effect.

The SMC *Adar1[-/+]* model is unique in that in haploinsufficiency there is no immediate

activation of MDA5, however in specific cellular contexts immunogenic RNA formation

appears to 'max out' the RNA editing capacity of the single *Adar1* allele leading to

MDA5 activation. This effect is overt in a 16-week high fat diet model of atherosclerosis,

however this was perhaps not evident in either of the prior short term carotid injury or

AAA models. However, a recent discovery that MDA5 (*IFIH1*) is a key locus in a large

GWAS of AAA raises an intriguing hypothesis that ADAR1 and MDA5 are causal

mechanisms in AAA formation[52].

Our data indicate that SMC phenotypic modulation is a cellular context that

requires heightened RNA editing and that deficient RNA editing with SMC *Adar1*

heterozygous mice leads to MDA5 activation that accelerates SMC phenotypic changes

and disease state. Although our data suggests that expression of *ADAR1* increases with

SMC phenotypic modulation in vitro and in atherosclerosis (Figs S1 and S4), this likely

reflects inadequate compensatory response leading to an overall accumulation of

dsRNA and MDA5 activation in specific cell states, influencing SMC phenotype. These

data reveal distinct trajectories to SMC formation of the vascular chondromyocyte

(CMC), where ISG activation through MDA5 facilitates the formation of CMCs and

vascular calcification. This effect of MDA5 activation on SMC phenotypic state is

consistent with prior work investigating the role of ADAR1 and MDA5 on cellular

proliferation and cell death[13]. ADAR1, through its role of editing dsRNA and preventing

MDA5 activation, is crucial in regulating proliferation and deterring endogenous dsRNA

from triggering transcriptional shutdown and apoptotic cell death through activation of

PKR (*EIF2AK2*)[13]. Intriguingly, in this report, with initial KD of *ADAR1*, cell proliferation

was found to be increased, however with mounting MDA5 activation there was

ultimately PKR activation and transcriptional shut down and cell death[13]. This effect,

where initial loss of ADAR1 stimulates proliferation and cell state transitions, is perhaps

consistent with modifying a cell state similar to SMC to FMC transition in atherosclerosis

in our SMC $Adar1^{-/+}$ model. However, with overwhelming MDA5 activation as seen in

our SMC $Adar1^{-/-}$ model, there is transcriptional shut down, cell death, and major

vascular effect.

Importantly, although classical inflammatory mechanisms have been implicated

in the pathogenesis of atherosclerosis and tremendous interest has been placed in the

potential to inhibit these pathways to treat disease[53], only a small minority of GWAS loci

in CAD implicate inflammatory pathways[2-5]. This finding raises the question as to the

overall weight of effect for disease causality for such inflammatory mediators (i.e. IL6,

TNFα, and IL1β). The relatively minimal benefit of IL1β antagonism in CAD[54] may

suggest an overall limited potential to treat disease through inhibition of these

inflammatory mediators. However, dsRNA sensing by MDA5 and ISG activation as an

overarching mechanism provides a new avenue of understanding as to how

inflammatory mechanisms are driving disease biology. Our data demonstrates that ISG

activation occurs in SMCs during atherogenesis, however the exact function of these

ISGs remains unclear. The function of ISGs has been previously reviewed[55], however,

ultimately IFN and ISGs serve key roles to regulate cellular viral immune response,

likely through driving transcriptional shutdown in a PKR dependent mechanism of cells

infected by a viral pathogen. Importantly, Type I interferon induces JAK/STAT

intracellular signaling pathways[55], a finding we observed in our ISG dependent cellular

trajectory in SMC phenotypic transition (Fig 6H & Fig S14). This finding suggests that

JAK/STAT is an important regulator of this trajectory. Specific ISGs have distinct roles in

fighting viral infection, for example on inhibiting viral entry (IFITM1/2/3), mRNA

synthesis (IFI16), protein synthesis (PKR), and mRNA degradation (ISG20)[55]. The

mechanism by which specific ISGs may modify atherosclerosis within the vessel wall

remains unclear. However, an important observation from human genetics is that

edQTL loci, which ultimately implicates dsRNA formation, MDA5 activation, and ISG

induction, to be a greater driver of disease heritability than all eQTLs in CAD[12]. This

data presented here further implicates a specific mechanism of disease while

highlighting a unique opportunity to leverage this biology for therapeutic development by

prioritizing the inhibition of dsRNA sensing and downstream activation of MDA5.

MATERIALS AND METHODS

Data availability

All raw and processed sequencing data are deposited in the National Center for Biotechnology Information Gene Expression Omnibus. Bulk and single cell RNA sequencing data is available through accession IDs GSE254862, GSE255661, and GSE280641.

Mouse Models

To generate SMC-specific deletion of Adar we used the *Adar$^{flox/flox}$* allele (exon 7-9 floxed; MGI allele: Adar$^{tm1.1Phs}$; MGI:3828307) as we have reported previously[56], crossed onto the validated SMC-specific mouse Cre allele (Myh11^{CreERT2}) combined with the floxed tandem dimer tomato (tdT) fluorescent reporter protein gene knocked into the ROSA26 locus (B6.Cg-Gt(ROSA)26Sortm14(CAGtdTomato)Hze/J) and ApoE$^{-/-}$ alleles (Adar$^{flox/flox}$, Myh11^{CreERT2}, ROSAtdT/+, ApoE$^{-/-}$). To induce homozygous or heterozygous deletion of *Adar1* in SMCs (*SMC-Adar1$^{-/-}$*: *Adar1*$^{fl/fl}$, *Myh11^{CreERT2}*, *ROSAtdTomato*, *ApoE$^{-/-}$* or *SMC-Adar1$^{-/+}$*: *Adar1*$^{fl/WT}$, *Myh11^{CreERT2}*, *ROSAtdTomato*, *ApoE$^{-/-}$*), mice receive 2 doses of tamoxifen (200mg/kg, P.O.) at 7.5-8 weeks of age (dose 1 - day 0; and dose 2 - day 2) and started on a high fat diet as we have done previously.[43,44] To evaluate the effect of MDA5 KO on *Adar1* KO background, we crossed the *Ifih1$^{-/-}$* mouse (B6.Cg-*Ifih1*$^{tm1.1Cln}$/J) that we have used previously[56], onto the above background in a heterozygous manner to generate SMC *Adar1$^{-/-}$* x MDA5$^{-/+}$ mice (Adar$^{flox/flox}$, *Ifih1$^{-/+}$*, Myh11^{CreERT2}, ROSAtdT/+, ApoE$^{-/-}$), SMC *Adar1$^{-/-}$* x MDA5$^{-/-}$ mice (Adar$^{flox/flox}$, *Ifih1$^{-/-}$*, Myh11^{CreERT2}, ROSAtdT/+, ApoE$^{-/-}$), or SMC *Adar1$^{-/+}$* x MDA5$^{-/+}$ mice (Adar$^{flox/WT}$, *Ifih1$^{-/+}$*, Myh11^{CreERT2}, ROSAtdT/+, ApoE$^{-/-}$).

Mouse aortic root/ascending aorta cell dissociation

Immediately after sacrifice, mice were perfused with phosphate buffered saline (PBS). The aortic root and ascending aorta were excised, up to the level of the brachiocephalic artery. Tissue was washed three times in PBS, placed into an enzymatic dissociation cocktail (2 U ml−1 Liberase (5401127001; Sigma–Aldrich) and 2 U ml−1 elastase (LS002279; Worthington) in Hank's Balanced Salt Solution (HBSS)) and incubated at 37

°C for 45min, and subsequently minced. The cell suspension was further dissociated

with 1ml pipette 20X and single cell suspension was confirmed under microscope. Cells

were then pelleted by centrifugation at 500g for 5 min. The enzyme solution was then

discarded, and cells were resuspended in fresh HBSS and passed through a 35um filter

into a FACS tube. To increase biological replication, a single mouse was used to obtain

single-cell suspension and three mice were used in combination for each scRNA

capture. 3 separate pairs of isolation were performed for control, *SMC-Adar1$^{-/+}$*, *SMC-*

Adar1$^{-/-}$, *SMC-Adar1$^{-/-}$ + Mda5$^{-/-}$,* or *SMC-Adar1$^{-/+}$ + Mda5$^{-/+}$* mice. Cells were FACS

sorted based on tdTomato expression. tdT+ cells (considered to be of SMC lineage)

and tdT− cells were then captured on separate but parallel runs of the same scRNA-

Seq workflow (gating strategy and threshold identical to those published in previous

work by Wirka et al[32]), and datasets were later combined for all subsequent analyses.

Mouse aortic root histology

IHC was performed according to standard protocol. Primary antibodies for CD68 rabbit

polyclonal antibody (1:400 dilution; ab125212; Abcam) with secondary rabbit-on-rodent

HRP polymer (RMR622; BioCare Medical) for CD68 evaluation. Hematoxylin and eosin

(H&E) and Trichrome staining was performed using standard protocols using and H&E

and Trichrome Stain Kits (Abcam, ab245880 and ab150686 respectively). The

processed sections were visualized using a Leica DM5500 microscope objective

magnifications, and images were obtained using Leica Application Suite X software. For

quantification of atherosclerosis plaque area, sections obtained at equal distance

measured from the superior margin of the coronary sinus were used for comparison to

ensure equal level sections were obtained. Images with FITC autofluorescence and

tdTomato with red channel representing SMC lineage traced cells were obtained. Areas

of interest were quantified using ImageJ (National Institutes of Health) software, and

compared using a two-sided t-test. Lesion size was defined by the area encompassing

the intimal edge of the lesion to the border of SMC-marker positive intima-media

junction. Calcification was evaluated using Ferangi Blue Chromogen Kit 2 (Biocare

Medical, FB813) following manufacturers protocol. For RNAscope, slides were

processed and hybridized according to the manufacturer's protocol, with reagents from

ACD Bio. All area quantification was performed in a genotype blinded fashion with

image J using length information embedded in exported files. All biological replicates for

each staining were performed simultaneously on position-matched aortic root sections

to limit intra-experimental variance.

Single-cell capture and library preparation and sequencing

All single-cell capture and library preparation was performed in the Quertermous lab,

sequencing was performed by Medgenome Inc (Foster City, CA). Cells were loaded into

a 10x Genomics microfluidics chip and encapsulated with barcoded oligo-dT-containing

gel beads using the 10x Genomics Chromium controller according to the manufacturer's

instructions. Single-cell libraries were then constructed according to the manufacturer's

instructions (Illumina). Libraries from individual samples were multiplexed into one lane

before sequencing on an Illumina platform with targeted depth of 50,000 reads per cell

for RNA or 400 million paired end reads per sequencing library.

Analysis of scRNA-Seq data

Fastq files from each experimental time point and mouse genotype were aligned to the

reference genome (mm10) individually using CellRanger Software (10x Genomics). The

dataset was then analyzed using the R package Seurat 30 and datasets were merged

in Seurat using standard Merge function. The dataset was trimmed of cells expressing

fewer than 2000 genes, and genes expressed in fewer than 50 cells. The number of

genes, number of unique molecular identifiers and percentage of mitochondrial genes

were examined to identify outliers. As an unusually high number of genes can result

from a 'doublet' event, in which two different cell types are captured together with the

same barcoded bead, cells with >9000 genes were discarded. Cells containing >7.5%

mitochondrial genes were presumed to be of poor quality and were also discarded. The

gene expression values then underwent library-size normalization and normalized using

established LogNormalize function in Seurat. No additional batch correction was

performed. Principal component analysis was used for dimensionality reduction,

followed by clustering in principal component analysis space using a graph-based

clustering approach via Louvain algorithm. UMAP was then used for two-dimensional

visualization of the resulting clusters. Pseudotime trajectory analysis was performed
using Slingshot with available software. Analysis, visualization and quantification of
gene expression and generation of gene modulescores were performed using Seurat's
built-in function such as "FeaturePlot", "VlnPlot", "AddModuleScore", and "FindMarker."
For immunogenic RNA score, we generated an AddModuleScore 'score' that
summates the expression of all 'immunogenic' RNA from a putative list that was
generated by combining our data from experimental *in vitro* models[26] along with human
genetic edQTL analysis[12].

HCASMC culture and in vitro genomic studies
Cells were cultured in smooth muscle growth medium (Lonza; catalog number: CC-
3182) supplemented with human epidermal growth factor, insulin, human basic
fibroblast growth factor and 5% FBS, according to the manufacturer's instructions. All
HCASMC lines were used at passages 4–8. HCASMCs immortalized with hTert
transgene were acquired from Dr. Clint Miller (UVA) were used for all calcification
assays with HCASMCs between passages 31 and 35. siRNA knockdowns were
performed using Lipofectamine RNAiMax (Life Technologies) using manufacturer's
recommended protocol at 50pg siRNA / 100,000cells.

Phenotypic transition, calcification, and TGFβ assays: For phenotypic transition
assays, HCASMC are cultured in 5% BSA and then subjected to serum starvation for 2
days. Following 2 days of serum starvation (0% BSA) HCASMCs develop a mature
SMC phenotype, reintroduction of serum (5%) stimulates a more proliferative phenotype
and RNA/protein is isolated 48hrs following reintroduction of serum. For calcification
assays, HCASMCs are cultured in 1% BSA with 10mM beta-glycerophosphate and
8mM CaCl2 for 7 to 10 days. For TGFβ stimulation assays, we cultured HCASMC to
confluency, treated with siRNA for 48 hours, and transitioned cells to a serum free
condition for 48 hours. We then stimulated cells with TGFβ (10ng/mL) or control for 72
hours before we isolated RNA and performed quantitative PCR.

RNA isolation and Bulk RNA sequencing

RNA was isolated from disrupted HCASMC cells using standard RNeasy mini kit

protocol (Qiagen). Isolated RNA underwent quality assessment, cDNA conversion, and

library construction by Novogene (San Jose, CA). cDNA libraries were sequenced using

Illumina Novoseq 6000 achieving 20-30 million paired-end reads per sample. RNAseq

data were processed using a workflow starting from fastq files and code is available on

github (https://github.com/zhaoshuoxp/Pipelines-Wrappers/blob/master/RNAseq.sh). In

brief, the quality control was performed using FastQC, mapping to the human genome

hg19/GRCh37 was performed using STAR second pass mapping to increase the

percentage of mapped reads, and counting was done with featureCounts using

GENCODE gtf annotation. Next, differential analysis was performed using DESeq2.

970
971 **RNA editing analysis**
972
RNA editing analysis in bulk RNA seq data was performed following a pipeline as we

have done previously[12]. Briefly, we compiled a list of reference editing sites for

quantification. By incorporating known sites in the RADAR database[57], tissue-specific

sites identified in GTEx V6p17, and recently published hyper-editing sites[58], we finalized

a list of 2,802,572 human editing sites. To quantify the editing levels, we computed the

ratio of G reads divided by the sum of A and G reads at each site. Because the bulk

RNA-seq data from HCASMC cells was not strand specific, the reads cannot be

automatically assigned to sense and antisense strands, so that the quantification of

editing level may be affected by expression of either of the two overlapping genes.

Therefore, we developed a method to measure the RNA editing level of a given site

using the reads derived from the corresponding transcript (sense for A-to-G edits and

antisense for T-to-C edits), with less influence by the other overlapping transcript. For

reads that are edited, they would be derived from the sense transcript if the edits are A-

to-G, and from the antisense transcript if the edits are T-to-C. For the rest of the reads

that are unedited, we estimated the numbers derived from the sense versus antisense

transcripts in proportion to their expression levels. Overall RNA editing frequency was

reported as percentage of total RNA editing sites across the genome.

List of Supplementary Materials

- Supplemental Figure 1. *SMC ADAR1 and MDA5 expression increase in atherosclerosis in mice*

- Supplemental Figure 2. *Loss of ADAR1 in human coronary artery SMCs (HCASMCs) regulates viral sensing transcriptomic response in vitro*

- Supplemental Figure 3. Global RNA editing is decreased in HCASMC following phenotypic modulation in vitro

- Supplemental Figure 4. *Serum and TGFβ stimulation upregulate ADAR1 expression with differential effect on ISG induction in HCASMC.*

- Supplemental Figure 5. *Loss of ADAR1 regulates transcriptomic and inflammatory response to calcification in vitro*

- Supplemental Figure 6. *Quantification of histological findings for control and SMC-Adar1 KO mice*

- Supplemental Figure 7. *Aortas from SMC Adar1 KO with Mda5 KO are indistinguishable from control*

- Supplemental Figure 8. *SMC Adar1 KD induces antiviral defense transcriptional pathways*

- Supplemental Figure 9. *Loss of SMC Adar1 coordinates distinct response throughout vessel wall*

- Supplemental Figure 10. *Homozygous deletion of MDA5 (Ifih1) prevents transcriptomic effect of SMC-Adar1 KO*

- Supplemental Figure 11. *SMC Adar1 haploinsufficiency has no effect on weight or cholesterol in atherosclerosis high fat diet model*

- Supplemental Figure 12. *SMC specific haploinsufficiency of Adar1 in SMC subset analysis shows ISG activation with phenotypic modulation in atherosclerosis*

- Supplemental Figure 13. *SMC specific haploinsufficiency of Adar1 increases vascular chondromyoctye formation*

- Supplemental Figure 14. ISG dependent trajectory analysis from SMC to CMC implicates distinct gene ontologies

1022   •   Supplemental Figure 15. *Isg15 RNAscope reveals increased Isg15 signal in*

1023   *plaque of SMC Adar1$^{-/+}$ mice*

1024   •   Supplemental Figure 16. *SMC specific haploinsufficiency in Adar1 increased*

1025   *SMC lineage traced cell content in plaque without change in acellular area*

1026   •   Supplemental Figure 17. *SMC specific haploinsufficiency of Adar1 has minimal*

1027   *effect on macrophage infiltration in atherosclerosis*

1028

References

Diseases, G. B. D. & Injuries, C. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* **396**, 1204-1222 (2020). https://doi.org/10.1016/S0140-6736(20)30925-9

Nikpay, M. *et al.* A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet* **47**, 1121-1130 (2015). https://doi.org/10.1038/ng.3396

Schunkert, H. *et al.* Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet* **43**, 333-338 (2011). https://doi.org/10.1038/ng.784

van der Harst, P. & Verweij, N. Identification of 64 Novel Genetic Loci Provides an Expanded View on the Genetic Architecture of Coronary Artery Disease. *Circ Res* **122**, 433-443 (2018). https://doi.org/10.1161/CIRCRESAHA.117.312086

Tcheandjieu, C. *et al.* Large-scale genome-wide association study of coronary artery disease in genetically diverse populations. *Nat Med* **28**, 1679-1692 (2022). https://doi.org/10.1038/s41591-022-01891-3

Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and disease. *Nat Rev Genet* **16**, 197-212 (2015). https://doi.org/10.1038/nrg3891

Consortium, G. T. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318-1330 (2020). https://doi.org/10.1126/science.aaz1776

Eisenberg, E. & Levanon, E. Y. A-to-I RNA editing - immune protector and transcriptome diversifier. *Nat Rev Genet* **19**, 473-490 (2018). https://doi.org/10.1038/s41576-018-0006-1

Jonkhout, N. *et al.* The RNA modification landscape in human disease. *RNA* **23**, 1754-1769 (2017). https://doi.org/10.1261/rna.063503.117

Walkley, C. R. & Li, J. B. Rewriting the transcriptome: adenosine-to-inosine RNA editing by ADARs. *Genome Biol* **18**, 205 (2017). https://doi.org/10.1186/s13059-017-1347-3

1060   11   Gatsiou, A. & Stellos, K. RNA modifications in cardiovascular health and disease.
1061   *Nat Rev Cardiol* **20**, 325-346 (2023). https://doi.org/10.1038/s41569-022-00804-8

1062   12   Li, Q. *et al.* RNA editing underlies genetic risk of common inflammatory diseases.
1063   *Nature* **608**, 569-577 (2022). https://doi.org/10.1038/s41586-022-05052-x

1064   13   Chung, H. *et al.* Human ADAR1 Prevents Endogenous RNA from Triggering
1065   Translational Shutdown. *Cell* **172**, 811-824 e814 (2018).
1066   https://doi.org/10.1016/j.cell.2017.12.038

1067   14   Liddicoat, B. J. *et al.* RNA editing by ADAR1 prevents MDA5 sensing of
1068   endogenous dsRNA as nonself. *Science* **349**, 1115-1120 (2015).
1069   https://doi.org/10.1126/science.aac7049

1070   15   Goubau, D., Deddouche, S. & Reis e Sousa, C. Cytosolic sensing of viruses.
1071   *Immunity* **38**, 855-869 (2013). https://doi.org/10.1016/j.immuni.2013.05.007

1072   16   Chalk, A. M., Taylor, S., Heraud-Farlow, J. E. & Walkley, C. R. The majority of A-
1073   to-I RNA editing is not required for mammalian homeostasis. *Genome Biol* **20**,
1074   268 (2019). https://doi.org/10.1186/s13059-019-1873-2

1075   17   Nishikura, K. Functions and regulation of RNA editing by ADAR deaminases.
1076   *Annual review of biochemistry* **79**, 321-349 (2010).
1077   https://doi.org/10.1146/annurev-biochem-060208-105251

1078   18   Jain, M. *et al.* RNA editing of Filamin A pre-mRNA regulates vascular contraction
1079   and diastolic blood pressure. *EMBO J* **37** (2018).
1080   https://doi.org/10.15252/embj.201694813

1081   19   Pestal, K. *et al.* Isoforms of RNA-Editing Enzyme ADAR1 Independently Control
1082   Nucleic Acid Sensor MDA5-Driven Autoimmunity and Multi-organ Development.
1083   *Immunity* **43**, 933-944 (2015). https://doi.org/10.1016/j.immuni.2015.11.001

1084   20   Mannion, N. M. *et al.* The RNA-editing enzyme ADAR1 controls innate immune
1085   responses to RNA. *Cell reports* **9**, 1482-1494 (2014).
1086   https://doi.org/10.1016/j.celrep.2014.10.041

1087   21   Feigenbaum, A. *et al.* Singleton-Merten syndrome: an autosomal dominant
1088   disorder with variable expression. *Am J Med Genet A* **161A**, 360-370 (2013).
1089   https://doi.org/10.1002/ajmg.a.35732

22   Rutsch, F., Buers, I. & Nitschke, Y. Hereditary Disorders of Cardiovascular
Calcification. *Arterioscler Thromb Vasc Biol* **41**, 35-47 (2021).
https://doi.org/10.1161/ATVBAHA.120.315577

23   Stellos, K. *et al.* Adenosine-to-inosine RNA editing controls cathepsin S
expression in atherosclerosis by enabling HuR-mediated post-transcriptional
regulation. *Nat Med* **22**, 1140-1150 (2016). https://doi.org/10.1038/nm.4172

24   Vlachogiannis, N. I. *et al.* Adenosine-to-inosine Alu RNA editing controls the
stability of the pro-inflammatory long noncoding RNA NEAT1 in atherosclerotic
cardiovascular disease. *J Mol Cell Cardiol* **160**, 111-120 (2021).
https://doi.org/10.1016/j.yjmcc.2021.07.005

25   Emdin, C. A. *et al.* Analysis of predicted loss-of-function variants in UK Biobank
identifies variants protective for disease. *Nat Commun* **9**, 1613 (2018).
https://doi.org/10.1038/s41467-018-03911-8

26   Sun, T. *et al.* A Small Subset of Cytosolic dsRNAs Must Be Edited by ADAR1 to
Evade MDA5-Mediated Autoimmunity. *bioRxiv*, 2022.2008.2029.505707 (2022).
https://doi.org/10.1101/2022.08.29.505707

27   Alsaigh, T., Evans, D., Frankel, D. & Torkamani, A. Decoding the transcriptome of
calcified atherosclerotic plaque at single-cell resolution. *Commun Biol* **5**, 1084
(2022). https://doi.org/10.1038/s42003-022-04056-7

28   Alexander, M. R. & Owens, G. K. Epigenetic control of smooth muscle cell
differentiation and phenotypic switching in vascular development and disease.
Annu Rev Physiol **74**, 13-40 (2012). https://doi.org/10.1146/annurev-physiol-
012110-142315

29   Basatemur, G. L., Jorgensen, H. F., Clarke, M. C. H., Bennett, M. R. & Mallat, Z.
Vascular smooth muscle cells in atherosclerosis. *Nat Rev Cardiol* **16**, 727-744
(2019). https://doi.org/10.1038/s41569-019-0227-9

30   Cheng, P. *et al.* ZEB2 Shapes the Epigenetic Landscape of Atherosclerosis.
Circulation (2022). https://doi.org/10.1161/CIRCULATIONAHA.121.057789

31   Kim, J. B. *et al.* Environment-Sensing Aryl Hydrocarbon Receptor Inhibits the
Chondrogenic Fate of Modulated Smooth Muscle Cells in Atherosclerotic

Lesions. *Circulation* **142**, 575-590 (2020).

https://doi.org/10.1161/CIRCULATIONAHA.120.045981

32   Wirka, R. C. *et al.* Atheroprotective roles of smooth muscle cell phenotypic

modulation and the TCF21 disease gene as revealed by single-cell analysis. *Nat*

Med **25**, 1280-1289 (2019). https://doi.org/10.1038/s41591-019-0512-5

33   Cheng, P. *et al.* Smad3 regulates smooth muscle cell fate and mediates adverse

remodeling and calcification of the atherosclerotic plaque. *Nature Cardiovascular*

Research **1**, 322-333 (2022). https://doi.org/10.1038/s44161-022-00042-8

34   Kim, H. J. *et al.* Molecular mechanisms of coronary artery disease risk at the

PDGFD locus. *Nat Commun* **14**, 847 (2023). https://doi.org/10.1038/s41467-023-

36518-9

35   Sharma, D. *et al.* Comprehensive Integration of Multiple Single-Cell

Transcriptomic Data Sets Defines Distinct Cell Populations and Their Phenotypic

Changes in Murine Atherosclerosis. *Arterioscler Thromb Vasc Biol* **44**, 391-408

(2024). https://doi.org/10.1161/ATVBAHA.123.320030

36   Ramaswami, G. *et al.* Accurate identification of human Alu and non-Alu RNA

editing sites. *Nat Methods* **9**, 579-581 (2012). https://doi.org/10.1038/nmeth.1982

37   Shankman, L. S. *et al.* KLF4-dependent phenotypic modulation of smooth muscle

cells has a key role in atherosclerotic plaque pathogenesis. *Nat Med* **21**, 628-637

(2015). https://doi.org/10.1038/nm.3866

38   Maurano, M. *et al.* Protein kinase R and the integrated stress response drive

immunopathology caused by mutations in the RNA deaminase ADAR1. *Immunity*

54, 1948-1960 e1945 (2021). https://doi.org/10.1016/j.immuni.2021.07.001

39   Wong, D. *et al.* FHL5 Controls Vascular Disease-Associated Gene Programs in

Smooth Muscle Cells. *Circ Res* **132**, 1144-1161 (2023).

https://doi.org/10.1161/CIRCRESAHA.122.321692

40   Cai, D. *et al.* A Novel Mechanism Underlying Inflammatory Smooth Muscle

Phenotype in Abdominal Aortic Aneurysm. *Circ Res* **129**, e202-e214 (2021).

https://doi.org/10.1161/CIRCRESAHA.121.319374

41   Fei, J., Cui, X. B., Wang, J. N., Dong, K. & Chen, S. Y. ADAR1-Mediated RNA

Editing, A Novel Mechanism Controlling Phenotypic Modulation of Vascular

Smooth Muscle Cells. *Circ Res* **119**, 463-469 (2016).
https://doi.org/10.1161/CIRCRESAHA.116.309003

42   Cai, D. & Chen, S. Y. ADAR1 Is Essential for Smooth Muscle Homeostasis and
Vascular Integrity. *Cells* **13** (2024). https://doi.org/10.3390/cells13151257

43   Cheng, P. *et al.* Smad3 regulates smooth muscle cell fate and mediates adverse
remodeling and calcification of the atherosclerotic plaque. *Nat Cardiovasc Res* **1**,
322-333 (2022). https://doi.org/10.1038/s44161-022-00042-8

44   Cheng, P. *et al.* ZEB2 Shapes the Epigenetic Landscape of Atherosclerosis.
Circulation **145**, 469-485 (2022).
https://doi.org/10.1161/CIRCULATIONAHA.121.057789

45   Lendahl, U., Muhl, L. & Betsholtz, C. Identification, discrimination and
heterogeneity of fibroblasts. *Nat Commun* **13**, 3409 (2022).
https://doi.org/10.1038/s41467-022-30633-9

46   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat.
Nat Commun **12**, 1088 (2021). https://doi.org/10.1038/s41467-021-21246-9

47   Tyner, J. W. *et al.* CCL5-CCR5 interaction provides antiapoptotic signals for
macrophage survival during viral infection. *Nat Med* **11**, 1180-1187 (2005).
https://doi.org/10.1038/nm1303

48   Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genomics* **19**, 477 (2018). https://doi.org/10.1186/s12864-
018-4772-0

49   Plenge, R. M., Scolnick, E. M. & Altshuler, D. Validating therapeutic targets
through human genetics. *Nat Rev Drug Discov* **12**, 581-594 (2013).
https://doi.org/10.1038/nrd4051

50   Mountjoy, E. *et al.* An open approach to systematically prioritize causal variants
and genes at all published human GWAS trait-associated loci. *Nat Genet* **53**,
1527-1533 (2021). https://doi.org/10.1038/s41588-021-00945-5

51   Estrada, K. *et al.* Identifying therapeutic drug targets using bidirectional effect
genes. *Nat Commun* **12**, 2224 (2021). https://doi.org/10.1038/s41467-021-
21843-8

52   Roychowdhury, T. *et al.* Genome-wide association meta-analysis identifies risk
loci for abdominal aortic aneurysm and highlights PCSK9 as a therapeutic target.
Nat Genet **55**, 1831-1842 (2023). https://doi.org/10.1038/s41588-023-01510-y
53   Soehnlein, O. & Libby, P. Targeting inflammation in atherosclerosis - from
experimental insights to the clinic. *Nat Rev Drug Discov* **20**, 589-610 (2021).
https://doi.org/10.1038/s41573-021-00198-1
54   Ridker, P. M. *et al.* Antiinflammatory Therapy with Canakinumab for
Atherosclerotic Disease. *N Engl J Med* **377**, 1119-1131 (2017).
https://doi.org/10.1056/NEJMoa1707914
55   Schoggins, J. W. Interferon-Stimulated Genes: What Do They All Do? *Annu Rev*
Virol **6**, 567-584 (2019). https://doi.org/10.1146/annurev-virology-092818-015756
56   Hu, S. B. *et al.* ADAR1p150 prevents MDA5 and PKR activation via distinct
mechanisms to avert fatal autoinflammation. *Mol Cell* **83**, 3869-3884 e3867
(2023). https://doi.org/10.1016/j.molcel.2023.09.018
57   Ramaswami, G. & Li, J. B. RADAR: a rigorously annotated database of A-to-I
RNA editing. *Nucleic Acids Res* **42**, D109-113 (2014).
https://doi.org/10.1093/nar/gkt996
58   Porath, H. T., Knisbacher, B. A., Eisenberg, E. & Levanon, E. Y. Massive A-to-I
RNA editing is common across the Metazoa and correlates with dsRNA
abundance. *Genome Biol* **18**, 185 (2017). https://doi.org/10.1186/s13059-017-
1315-y

1202

1203
1204   **Acknowledgements**
1205
1206   **Funding:**
1207

National Institutes of Health grant F32HL160067 (CW)
National Institutes of Health grant L30HL159413 (CW)
National Institutes of Health grant K08HL167699 (CW)
National Institutes of Health grant K08HL153798 (PC)
National Institutes of Health grant R01HL134817 (TQ)
National Institutes of Health grant R01HL139478 (TQ)
National Institutes of Health grant R01HL156846 (TQ)
National Institutes of Health grant R01HL151535 (TQ)

National Institutes of Health grant R01HL145708 (TQ)
National Institutes of Health grant UM1 HG011972 (TQ)
National Institutes of Health grant R35GM144100 (JBL)
National Institutes of Health grant R01MH115080 (JBL)
National Institutes of Health grant R01GM102484 (JBL)
American Heart Association grant 20CDA35310303 (PC)
American Heart Association grant 23CDA1042900 (CW)
American Heart Association grant 23POST1018991 (WG).

1224
1225 **Author Contributions**
1226

Conceptualization: C.W., Q.L., P.C., J.B.L., and T.Q.

Methodology: C.W., Q.L., J.M.

Investigation: C.W., Q.L., J.M., H.G., D.G., W.G., M.W., Q.Z., M.R., D.L., B.P., A.B.,

R.K., T.N.

Visualization: C.W.

Funding acquisition: C.W., J.B.L., T.Q.

Project administration: C.W., J.B.L., T.Q.

Supervision: J.B.L., T.Q.

Writing — original draft: C.W., J.B.L., T.Q.

Writing — review and editing: C.W., Q.L., J.B.L., T.Q.

1237
1238
Competing interests:

C.W. is a consultant for AiRNA Bio and Avidity Biosciences. T.Q. serves on the scientific

advisory board for Amgen. J.B.L. is a co-founder of AIRNA Bio and a consultant for

Risen Pharma. J.B.L. and Q.L. are named inventors of a provisional patent filed by

Stanford University (serial no. 63/473,678), describing a method related to RNA editing.

The other authors declare no competing interests.

1245

1246 **Data and materials availability:**

1247 All raw and processed sequencing data are deposited in the National Center for

1248 Biotechnology Information Gene Expression Omnibus. Bulk and single cell RNA

1249 sequencing data is available through accession IDs GSE254862, GSE255661, and

1250 GSE280641.

Figure 1. *Smooth muscle cells in atherosclerosis express immunogenic RNA and show evidence of ISG induction in phenotypic modulation.* (A) Schematic of

ADAR1 RNA editing of dsRNA. (B) UMAP of scRNAseq data from human carotid

atherosclerotic plaque (Alsaigh et al., 2022). (C) Module score of immunogenic RNA

expression. (D) Clustering of SMC subset with (E) violin plot showing SMC marker

MYH11 with (F) fibromyocyte/chondromyocyte marker *TNFRSF11B*. (G-I) Violin plots of

ISGs *ISG15* (G), *IF135* (H), and *IFI16* (I). (J) GO-term enrichment analysis for human

athero SMC clustering. (K-N) In mouse comprehensive integrated dataset of lineage

traced SMCs (from Sharma et al., 2024), featureplot of *Myh11* (K) and *Col2a1* (L) at 0

and 16 weeks of high fat diet. (L-M) Featureplots of *Isg15* (L) and *Ifit3* (M) in SMC

phenotypic modulation.

1263

1264

Figure 2. *ADAR1 regulates RNA editing in human coronary SMCs and RNA editing requirement is dependent on cell context.* (A) Schematic of SMC phenotypic modulation in vitro assay. (B) Linear regression analysis of ADAR1 expression and global RNA editing frequency in primary human coronary artery SMCs (HCASMCs) following siRNA KD of *ADAR*. (C) Grouped bar chart of RNA editing frequency with cellular treatment of scramble, *ADAR*, and *ADAR + IFIH1* (MDA5) siRNA. (D) Volcano

plot of bulk RNAseq DE gene analysis comparing scramble vs *ADAR* siRNA in serum

starved conditions, (E) volcano plot comparing scramble vs *ADAR + IFIH1* siRNA in

serum starved conditions, (F) volcano plot comparing scramble vs *ADAR* siRNA in

serum fed conditions, and (G) volcano plot comparing scramble vs *ADAR + IFIH1*

siRNA in serum fed conditions. (H) Principal component analysis (PCA) of bulk RNAseq

data of principal components 1 and 2 for each HCASMC treatment. Grouped bar chart

of normalized RNAseq reads across HCASMC treatments for (I) *ACTA2*, (J) *CNN1*, (K)

ISG15, (L) *KLF4*, (M) *EGR1*, and (N) *ATF3*. *P*-values represent simple linear regression

(B) or (C, I-N) two-way ANOVA with multiple comparisons post hoc analysis. N = 3 RNA

seq libraries per condition.

1281

1282

Figure 3. *SMC specific ADAR1 is required to maintain vascular integrity and survival.* (A) Schematic of SMC-*Adar1⁻/⁻* model and accumulation of dsRNA and MDA5 activation. (B) Survival curves for 56 mice treated with tamoxifen at 8 weeks of age, comparing *SMC-Adar1⁻/⁻*, *SMC-Adar1⁻/⁺* (*Adar1*$^{fl/WT}$, *Myh11*^{CreERT2}, *ROSAtdTomato*, *ApoE⁻/⁻*), and control (*Adar1*$^{WT/WT}$, *Myh11*^{CreERT2}, *ROSAtdTomato*, *ApoE⁻/⁻*) genotypes. (C) Gross image of aortic root and ascending aorta of *SMC-Adar1*^{KO} mouse at 2 weeks

following tamoxifen treatment. (D-E) Representative images of hematoxylin and eosin

(H&E) staining from *SMC-Adar1*^{-/-} (D) and control genotype mice (E). (F-G) Fluorescent

imaging with autofluorescence indicating elastin (green) and Myh11 derived cells

labeled with tdTomato (red) in two mice with adventitial ballooning in *SMC-Adar1*^{-/-} (F-

G). (H) Bright field imaging of aorta cross section from *SMC-Adar1*^{-/-} mouse

corresponding to fluorescent imaging in panel G. (I) Cd68 immunostaining of aorta

cross section from *SMC-Adar1*^{-/-} mouse showing macrophage infiltration. (J) Masson

trichrome stain of aorta cross section from *SMC-Adar1*^{-/-} mouse showing intra-medial

collagen deposition. *P*-value represents comparison of survival curves, Log-rank

(Mantel-Cox) test (B).

1299

1300

Figure 4. *Haploinsufficiency in MDA5 is adequate to prevent mortality in SMC-Adar1$^{-/-}$ mice.* (A) Survival curves for *SMC-Adar1$^{-/-}$* (*Adar1*$^{fl/fl}$, *Myh11*^{CreERT2}, *ROSAtdTomato*, *ApoE*$^{-/-}$), *SMC-Adar1$^{-/-}$ + MDA5$^{-/+}$* (*Adar1*$^{fl/fl}$, *Ifih1*$^{-/+}$, *Myh11*^{CreERT2}, *ROSAtdTomato*, *ApoE*$^{-/-}$), and *SMC-Adar1$^{-/-}$ + MDA5$^{-/-}$* (*Adar1*$^{fl/fl}$, *Ifih1*$^{-/-}$, *Myh11*^{CreERT2}, *ROSAtdTomato*, *ApoE*$^{-/-}$) genotypes following tamoxifen treatment at 8 weeks of age. (B) Weight curve at 0, 7, and 14 days following tamoxifen treatment between genotypes of surviving mice. (C) Individual weight measures at 0, 7, and 14 days following tamoxifen treatment between genotypes of surviving mice. (D) Schematic diagram of *SMC-Adar1$^{-/-}$* and MDA5 haploinsufficiency or global KO protecting against mortality. *P-*

1310   values represent comparison of survival curves Log-rank (Mantel-Cox) test (A-B). (B)

1311   Data point represents means ($\pm$SEM).

1312

1313

1314

Figure 5. *Single cell RNA sequencing reveals SMC Adar1 to control ISG*

activation, phenotypic response, and macrophage infiltration. (A) scRNAseq

UMAP from aortic root and ascending aorta grouped by genotype of *SMC-Adar1$^{-/-}$*

(*Adar1$^{fl/fl}$*, *Myh11^{CreERT2}*, *ROSAtdTomato*, *ApoE$^{-/-}$*) and control (*Adar1$^{WT/WT}$*, *Myh11^{CreERT2}*,

ROSAtdTomato, *ApoE$^{-/-}$*) genotypes at 2 weeks following tamoxifen treatment. (B)

UMAP with clustering and cell type identification of combined *SMC-Adar1$^{-/-}$* and control

genotype data. (C) Featureplot of tdTomato showing SMC lineage tracing. (D)

Featureplot of *Isg15*. (E) RNAscope of *Isg15* in aortic tissue of *SMC-Adar1$^{-/-}$* mice

showing highest *Isg15* expression at the lumen aspect of media, with additional

adventitial expression. (F) Featureplot of *Ifnb1*. (G) Heatmap of network centrality

scores for CCL signaling network across cell clusters. (H) Contributions of each

receptor-ligand pair for CCL signaling. (I) Chord diagram of receptor-ligand interaction

between clusters for CCL5:CCR5. (J-K) Featureplot of (J) *Ccl5* and (K) *Ccr5*.

1328

1329

1330

1331

Figure 6. *SMC specific haploinsufficiency of Adar1 activates the ISG response in*

atherosclerosis increasing vascular chondromyoctye formation. (A) UMAP of

scRNAseq data from atherosclerotic aortic root and ascending aorta at 16 weeks high

fat diet split by *SMC-Adar1$^{-/+}$* (*Adar1$^{fl/WT}$*, *Myh11^{CreERT2}*, *ROSAtdTomato*, *ApoE$^{-/-}$*) and

control (*Adar1$^{WT/WT}$*, *Myh11^{CreERT2}*, *ROSAtdTomato*, *ApoE$^{-/-}$*) genotypes. (B) UMAP of

grouped *SMC-Adar1$^{-/+}$* and control genotypes. (C) Featureplot of *Isg15* split by

genotype. (D) UMAP of SMC subset analysis grouped by genotype. (E) UMAP of SMC

subset cluster analysis split by genotype showing SMC, SMC_ISG (interferon

stimulated gene), fibromyocyte (FMC), and chondromyocyte (CMC) cell clusters. (F)

Stacked bar chart showing cell populations of SMC subset for SMC, SMC_ISG, FMC,

and CMC clusters. (G) Featureplot of *Isg15* in SMC subset split by genotype. (H)

Lineage trajectory analysis using slingshot within SMC subset showing two distinct

lineages (SMC_ISG dependent and independent) from SMC to CMC. (I&J) RNA scope

for *Isg15* in aortic root sections in control (I) and *SMC-Adar1$^{-/+}$* (J) mice.

1346

Figure 7. *SMC specific haploinsufficiency in Adar1 increases plaque size and*

calcification in atherosclerosis. Representative fluorescent images of aortic root

plaque histology in *SMC-Adar1$^{-/+}$* and control genotypes (A&B) with quantification of

plaque area (C), plaque area normalized to internal elastic lamina area (IEL) (D), media

area (E), media area normalized to external elastic lamina area (EEL) (F), plaque area

normalized to media area (G), and EEL area (H). Representative images of Ferangi

Blue calcification staining in *SMC-Adar1$^{-/+}$* and control genotypes (I&J), quantification of

Ferangi Blue area (K), and Ferangi Blue area normalized to plaque area (L). (M)

Schematic diagram showing mechanism of dsRNA formation, MDA5 activation, and

impaired RNA editing by ADAR1 to accelerate SMC phenotypic modulation,

calcification, and atherosclerosis. Data represent mean +/- SEM, N = 13 per group. *P*-

values represent unpaired two-tailed T-test.

1361

1362

1363

1364
Figure 8. *Haploinsufficiency of Mda5 prevents ISG activation and*
chondromyocyte formation in atherosclerosis with SMC-Adar1^{-/+}. (A) UMAP of
integrated scRNAseq data set between control (*Adar1^{WT/WT}, Myh11^{CreERT2},*

ROSAtdTomato, ApoE$^{-/-}$), SMC-Adar1 het (Adar1^{flWTl}, Myh11^{CreERT2}, ROSAtdTomato,

ApoE$^{-/-}$), and SMC-Adar1 het + Mda5 het (Adar1^{flWTl}, Ifih1$^{-/+}$, Myh11^{CreERT2},

ROSAtdTomato, ApoE$^{-/-}$) genotypes following 16 weeks high fat diet. (B) Featureplot of

Isg15 split by genotype. (C) UMAP tdTomato+ subset analysis with SMC, SMC_ISG,

FMC, and CMC clusters split by genotype. Featureplots split by genotype for (D)

Col2a1, (E) *Acan*, and (F) *Isg15*. (G) Stacked bar charts showing proportion of cells

within 'SMC', 'SMC_ISG', 'FMC', and 'CMC' clusters split by genotype. (H) Heatmap

visualization for normalized expression of marker genes for control, *SMC-Adar1 het*,

and *SMC-Adar1 het + Mda5 het* genotypes in tdTomato+ subset analysis. Dotplot gene

expression splity by genotype in SMC subset for (I) ISG genes and (J) FMC/CMC

marker genes.

1379

1380