<u>Title</u>

Development of a DSM test battery to determine depression-like marmoset

<u>Authors</u>

Hajime Yamanaka[1,2,*], Hidetoshi Ishibashi[1,3], Katsuki Nakamura[1,4]

<u>Affiliations</u>

[1]Department of Animal Models for Human Disease, National Institute of

Neuroscience, National Center for Neurology and Psychiatry, Kodaira, Tokyo

187-8502, Japan

[2]Neural Circuit Unit, Okinawa   Institute of Science and   Technology Graduate

University, Onna-son, Okinawa 904-0495, Japan

[3]Pre-clinical Research Center, Tokyo Medical University, Shinjuku-ku, Tokyo,

160-8402, Japan

[4]Section of Cognitive Neuroscience, Center for the Evolutionary Origins of

Human Behavior, Kyoto University, Inuyama, Aichi, 484-8506, Japan

*Corresponding Author: Hajime Yamanaka, Ph.D.

Neural Circuit Unit, Okinawa Institute of Science and Technology Graduate

University, Onna-son, Okinawa

Email: hajime.yamanaka@oist.jp

The number of words in the text of the paper (Introduction, Results, Discussion);

3203 words.

The number of display items; 7 figures and 1 table

Abstract

Depression is a serious and high-incidence mental disorder that can lead to suicide. Despite much depression research using humans and rodents to date, it has been difficult to completely overcome depression and elucidate its etiology. Marmoset monkeys have been utilized to study neuropsychiatric disorders in recent years. Since the clinical diagnostic criteria for psychiatric disorders are mainly based on multiple behavioral changes, test battery systems, a multifaceted and comprehensive strategy, are applied to depression research. Here, we devised a test battery with six tests to investigate depression-like symptoms corresponding to clinical criteria and set criteria for a test indicator-based judgment. We did a usability confirmation experiment using a drug and judged three of nine marmosets as depression-like. This rate is comparable with the incidence rate of human patients; i.e., 10-30% of patients who received reserpine treatment suffer from depression. Thus, we propose the test battery we constructed in this study will contribute to the study of depression.

Introduction

Major depressive disorder (depression) is a disease with numerous problems

and in need of further research. Its lifetime prevalence is as high as 8-12%, and

it affects about 280 million people worldwide[1-3]. The social and economic losses

caused by depression are enormous, since depression can lead to suicide,

social withdrawal, and absence from work. The World Health Organization has

warned that depression will be one of the top risk diseases of global burden in

the future, and the Lancet-World Psychiatric Association Committee has issued

a declaration calling for efforts to reduce the global burden of depression[4,5]. The

Diagnostic and Statistical Manual of Mental Disorders (DSM) diagnostic criteria

for depression lists nine symptoms, implying that this disease exhibits a

complex pathological condition in which multiple symptoms overlap at the same

time[6]. Depression is a disease that is elusive and offers few clues, because not

only has no clear etiology been identified, no clear abnormal findings in clinical

tests, such as imaging or biomarkers, or postmortem brain examinations have

been identified. Antidepressants, which are typical treatment, take two weeks to

work, and one-third of patients are refractory to treatment[7,8]. Currently, patients

continue to suffer from complex and long-term depressive symptoms, thereby

prompting a vigorous pursuit of etiological elucidation within the scientific

community. However, it is extremely difficult to find an etiology that goes beyond

the realm of hypothesis. Given that depression has not been completely

controlled or eradicated to date, expectations for future progress in depression

research are high.

Depression studies typically use rodent models or clinical specimens;

however, monkey models are beginning to be realized. Numerous rodent

models based on drug and genetic manipulation techniques are reported[9,10].

Although the development of antidepressants using rodents has contributed

significantly to depression research, no drug has been developed that can

completely control depression. The genetic analysis of big data of human

specimens has sought genetic predispositions and include large-scale genome-

wide association studies (GWAS) that have reported associations with many

genome sequences[11]. The reported etiological factors are being studied for

treatment and diagnosis. On the other hand, monkey studies with depression

are gradually beginning to progress, especially in model. Marmosets share

many characteristics with humans and have been used extensively to study

many human diseases but rarely depression[12-17]. One of the reasons is the lack

of a well-developed research platform. Such a platform would provide its own

unique insights while bridging rodent and human studies.

   We devised the "DSM Test Battery for Depressed Monkeys" in this study in

accordance with the policies and methods of the DSM, the international

diagnostic standard for mental disorders. The DSM provides specific symptoms

and diagnostic criteria and operationally makes a diagnosis according to them

(symptom-based diagnosis). On the other hand, our test battery provides

specific test indicators and criteria, and judgments of depression-like

marmosets are made operationally according to them (test indicator-based

judgment). Test indicators cover as many DSM depressive symptoms as

possible, and model judgment is conducted on an individual basis. The

Peephole Test (PhT) for visual exploratory behavior (VEB) and the Sucrose

Preference Test (SPT) for anhedonic behavior were devised to examine "loss of

interest and pleasure" in the DSM. The Eat Style (ES) test for food sorting

behavior was devised to examine "fatigue or loss of energy" in the DSM. We

added three tests corresponding to DSM depressive symptoms for a total of six

tests developed. Some of these tests were developed based on the behavioral

characteristics of monkeys and previous studies of depression. VEB dysfunction

and anhedonic behavior closely mimic the loss of interest and pleasure, a core

symptom of major depression episodes according to the DSM. VEB is

spontaneous behavior exploring an external environment using vision and was

first examined in rhesus monkey[18]. The same group later showed that the

frequency of VEB increases when another monkey or a moving toy train is

present and that the duration of VEB is prolonged when moving pictures are

shown, indicating that exploratory behavior intensity depends on the type of

stimulus[19,20]. Thus, it is argued that VEB is closely related to curiosity, interest,

and motivation[21]. However, the behavioral characteristics of other species

including marmosets, remain unclear. Anhedonic behavior describes the

decreased ability to feel or experience pleasure[22]. The SPT is used to assess

anhedonic behavior in models of depression, including those stress-induced,

drug-induced, and genetically-induced[23-26]. Generally, the models display lower

sucrose intake/preference, but this decline is recovered by chronic treatment

with antidepressants. However, there are no SPT studies of marmosets. Other

tests for motor activity (Activity) and food intake (FI) are already established,

and there are standard methods routinely used for body weight (BW)[27,28]. In

addition, the ES test is a test developed independently to examine food sorting

behavior.

In the present study, a three-step work process was carried out to develop a

new DSM-based research platform to determine depressive-like marmosets.

First, several tests suitable for marmosets were designed and developed, and

preliminary experiments were conducted to obtain basic data for each test. Next,

the tests were combined to construct a test battery, and the criteria, order of

implementation, and period were set. Finally, a usability confirmation

experiment was conducted using drugs. Here, a drug that causes depression-

like symptoms was administered to the marmosets, and changes were

examined during the drug administration (ON) and non-administration (OFF)

periods to confirm the drug's efficacy. Depression-like marmosets were

determined according to a simple and specific procedure (algorithm) based on

patterns of change. The drug used was reserpine, which causes depression-like

symptoms in laboratory animals and humans and has abundant human data

[29,30]. Using our new platform, we clarified how many of the nine drug-treated

marmosets were depression-like marmosets and compared the incidence with

previous human findings. This research platform, the "DSM Test Battery for

1   Depressed Monkeys," has the potential to establish new criteria for determining

2   depression-like models and unify etiological testing and comparisons.

3
4

Results

Experiment-1: Duration and frequency of VEB by means of PhT

In experiment-1, a test was developed to measure the duration and frequency

of peeping in response to curiosity while the test subject was in its home cage

(Fig. 1). We examined how the VEB of marmoset differed in response to

different stimuli **(Fig. 2)**. The duration and frequency of VEB were investigated

among five testing stimuli: a model insect, a model car, and three conspecifics

(Conspecific 1, 2 and 3). A one-way repeated measures ANOVA showed that

there were significant differences among the type of stimulus in both the

duration ($F_{(4, 16)}$ = 26.69, $p < 0.001$) and frequency ($F_{(4, 16)}$ = 19.63, $p < 0.001$).

By scrutinizing the data, the stimuli were divided into two main groups: object

and conspecific. A multiple comparison Scheffe's test showed significant

differences between the object and conspecific groups (duration, $p < 0.001$;

frequency, $p < 0.001$), indicating that the marmosets showed longer and more

frequent VEB to the conspecifics than to the objects.

Experiment-2: Sucrose concentration-response by means of SPT

The SPT is easily performed using a strong spring-loaded fixture, disposable

bottle, and limiting the supply of water from other sources **(Fig. 3a)**. The animal

intakes water from two bottles, one with sucrose solution and the other with tap

water, over 24 days (4 days per concentration) **(Fig. 3b and c)**. A one-way

repeated measures ANOVA showed significant differences in sucrose solution

intake ($F_{(1.80, 10.79)} = 49.79$, $p < 0.001$), sucrose preference percentage ($F_{(1.56, 9.35)} = 34.18$, $p < 0.001$), and tap water intake ($F_{(5, 30)} = 11.49$, $p < 0.001$)

depending on the sucrose concentration. Since a sucrose preference of 0% is

assumed to be the anhedonic state, we compared sucrose solution intake and

sucrose preference percentage with the 0% condition. A post hoc analysis

revealed that sucrose solution intake was higher for sucrose concentrations of

0.375, 0.75, 1.5, and 3.0% compared with 0% (Shaffer method, $p = 0.001$-

0.022), and the sucrose preference percentage was higher for 1.5 and 3.0%

(Shaffer method; 1.5%, $p = 0.017$; 3.0%, $p < 0.001$). These results revealed that

a sucrose concentration of more than 1.5% is needed to assess anhedonia in

marmoset monkeys. Hence, we chose a sucrose concentration of 1.8% for the

following experiments.

Experiment-3: Preliminary experiment for ES test and FI

During the ES test, marmosets took pellets (food) from a feedbox and ate the

food or dropped some into a mesh tray under their cages **(Fig. 4a and b)**. We

successfully developed a measurement procedure consisting of five steps,

which enabled us to simultaneously examine two indicators: the food drop index

and intake **(Supplementary Fig. 1a)**. In this preliminary experiment of over 3

days, when 40.1 g of food was fed, the amount of FI was 12.5 g (SD 3.3 g), and

the total amount of leftovers was 27.6 g (SD 3.5 g) **(Fig. 4c and**

Supplementary Fig. 1b). The amount of dropped leftover was 17.0 g (SD 7.5

g) resulting in a food drop index of 65% (SD 32.9%, range 19-100%) **(Fig. 4d**

and Supplementary Fig. 1b). Also, none of the 12 marmosets had a dropped

leftover rate (food drop index) of 0%, indicating that they have the habit of

dropping food if they are given enough food **(Fig. 4e)**.

Experiment-4: Preliminary experiment for motor activity using an infrared

sensor.

Each marmoset's hourly activity is shown in **Fig. 5.** Activity began in the

morning, remained high during the day, and disappeared when the lights went

out in the evening. The circadian rhythm of activity levels associated with

changes in light and dark was consistent with previous reports about

marmoset[27,31].

Experiment-5: Usability confirmation experiment by a test battery

consisting of six tests

To confirm how many marmosets were depression-like upon reserpine

treatment, we identified individual marmosets showing depression-like

symptoms using a test battery and specific judgment criteria. This battery

consists of six tests: PhT, SPT, ES test, FI, BW, and Activity **(Fig. 6** and

Supplementary Table 1). Three to six marmosets qualified as responders for

each test (for details, see **Table 1** and **Supplementary Tables 2-8**). Three

marmosets, Monkeys B, C and I, met the criteria that depression-like

marmosets are responders in four or more tests, giving us an incidence of

depression-like marmosets by reserpine treatment of 33%. This incidence rate

resembles the reported 19% and 26% human clinical incidence[32,33]. Notably,

Monkey C qualified as a responder for all six tests (**Fig. 7**), and Monkeys B and

I qualified as responders for four tests (**Supplementary Fig. 2**). Plasma cortisol

1   levels showed many marmosets qualified as responders, consistent with a

2   previous rat study (**Supplementary Table 9**)[34].

Discussion

Here we report the "DSM Test Battery for Depressed Monkeys", a new research

platform for the study of depression using marmoset. We designed   and

developed several tests for marmosets and collected their basic data. We

clarified in these preliminary experiments that VEB is efficiently induced when a

conspecific is presented as the stimulus in the PhT and that sucrose water at a

concentration of 1.5% or higher should be used to identify anhedonic behavior

in the SPT. Next, we combined tests and set the criteria, order of

implementation, and period for this platform. Finally, in a usability confirmation

experiment using reserpine, the incidence rate of depression-like marmosets

detected by this battery test was 33%, which is comparable with the human

incidence[32,33].

   This battery covers four of the nine DSM diagnostic criteria (**Supplementary**

Table 1)**.** Among the remaining five, it is possible to add a new test for (4)

abnormality of sleep and (8) diminished ability to the battery. The former can be

measured using sleep polysomnography, and the latter can be examined using

a cognitive function test[35,36]. Because the outstanding three criteria are based

on the patient's chief complaint and information from their families, they are

difficult to introduce to the battery. Importantly, this battery is close to a clinical

diagnosis in that the symptoms are evaluated for each and every individual,

facilitating translational research.

   Using the battery as a screening strategy for an exploratory study will enable

the isolation and identification of individuals, etiology hypotheses, and the

selection of antidepressants with certain characteristics. For example, since the

depressive symptoms caused by reserpine do not appear in all individuals even

at the same dose, it is generally assumed that there is a hidden regulator

system determining the vulnerability or responsiveness in the background of the

depression. In order to identify this system, it is necessary to efficiently obtain

and use a depression model with high phenomenological similarity. Based on

our finding, we therefore concluded that Monkey C should be bred over several

generations. In this regard, the battery will play a critical role in selective

breeding. Among primates closely related to humans, the marmoset is

advantageous for selective breeding because of its earlier sexual maturation

and higher fecundity. In addition, this strategy may selectively breed individuals

who are not only highly responsive to reserpine but also vulnerable to

depression.

As for the test battery system`s use to study etiology hypotheses, it has been reported that synaptic transmission and neuronal activity can be switched on and off by drug administration in DREADD (Designer Receptors Exclusively Activated by Designer Drug) and Tet-on systems using a virus vector. This capability allows for an analysis of V-shaped patterns using doxycycline or clozapine-N-oxide[37,38].

Finally, regarding use of our battery in preclinical antidepressant screening, new antidepressant candidates have been screened mainly using rodent models. Moreover, the battery may be able to pick up other antidepressants that do not show positive effects in rodents. For example, the teratogenicity of thalidomide and the appearance of major signs of MPTP are observed in primates but not in rodents[39]. By setting a sufficiently long washout period between treatments, different drugs can be evaluated by the same assessment tool using an identical individual, resulting in more strict and effective comparisons of drugs.

The PhT revealed that VEB in marmosets has similar characteristics as rhesus monkeys. Thus, this test will be a useful tool not only for evaluating symptoms of depression but also for researching interpersonal relationships,

fear, and the development of antidepressants and anxiolytics. The presentation

of a conspecific individual has been reported to increase the frequency of VEB

2.7-fold compared with the absence of the individual[19]. In the present study, we

found the mean frequency was 16 counts under the presence of a conspecific in

the stimulus phase and 5 counts in the empty chamber condition in the baseline

phase for a 3.2-fold change in experiment-1. The similarity of this fold change to

that observed in rhesus monkeys supports the idea that the PhT is a useful tool

for measuring VEB in marmosets. Also, the PhT is useful for objectively

evaluating attachment, interest, and curiosity by focusing on VEB as well as

research on social aversion and withdrawal, anxiety, and fear. For example, an

ethologically based test using a taxidermized predator (e.g., cat, snake, or

hawk) that is placed on a large arena was reported to study anxiolytic action in

marmoset[40]. Similarly, it is possible to measure the duration and frequency of

VEB only by replacing the stimulus in the chamber with a taxidermized predator

in the PhT apparatus. Hence, in a smaller space, it is possible to examine the

effects of psychotropic drugs such as anxiolytic drugs on behavior. On the other

hand, we found no responder to the conspecific-induced frequency difference of

VEB in experiment-5. Furthermore, the frequency in control block-1 (2.1 ± 5.0

counts, n = 9) was low compared with that in experiment-1 (11.5 ± 6.0 counts, n

= 5), and a large number of marmosets showed values near zero or negative. In

contrast, the duration difference had some responders, remained high and

stable, and was significantly unaffected by the time course from experiment-1 to

experiment-5. Thus, the frequency of VEB might be a labile indicator with an

elusive property. In addition, the PhT in the next generation of the battery should

use only conspecifics as the presenting stimulus.

We also reported, for the first time, the sucrose response of the SPT in

marmoset, although it has been reported in rhesus monkeys and rodents.

Although the response curve in the marmoset showed a concentration-

dependent increase similar to that of the rhesus monkey, there was a difference

in slope, with the marmoset showing a weaker relationship. Additionally, the

curve has been studied in a maternal-deprivation depression model, with the

consumption of sucrose solution reducing at 1.5% and 3.0% concentrations,

consistent with our findings[41]. The sucrose concentration used in the present

study was 1.8%, which is within these two concentrations. On the other hand,

the sucrose concentration used in the SPT for rodents is typically 0.5–2%. If the

concentration is too low, individuals will not show a preference for sweetness.

Conversely, if it is too high, the sensitivity will be poor and it will be difficult to

detect anhedonic behavior. Thus, 1–2% sucrose concentration might be optimal

for investigating the anhedonic behavior of marmoset in the SPT.

   The food drop index in the ES test is an essential test in depression research

but is also underdeveloped. The food drop index is defined as the percentage of

pellets thrown on the floor after being picked up once, nibbled, and checked for

favorite flavor and softness and describes a series of actions referred to as food

sorting behavior. In the present study, numerous pellets were observed under

the animals' home cages, and some of the pellets showed evidence of nibbling.

This observation suggests that the marmosets engaged in sorting the pellets by

grasping and evaluating their seasoning and texture. A decline in dropped foods

is associated with symptoms such as diarrhea and disturbed fur. Therefore,

monitoring the amount of dropped food serves as a crucial indicator in rearing

practices for the early detection of poor health. In addition, since this behavior is

a simple routine task and does not require a high level of concentration or

thinking, it can be used to evaluate fatigue and loss of energy (symptom

number "6") in the DSM criteria **(Supplementary Table 1)**. The food drop index

was high (65%), and experiment-3 revealed that healthy marmosets always

dropped food selectively. Maintaining stable high levels in this food drop index is

essential for disease modeling research that focuses on assessing reductions in

the food drop index related to DSM symptom 6. Initially, it was hypothesized

that the food drop index serves as the same test indicator as food intake.

However, it was determined that the two indicators are different, as clear

differences were observed between responders **(Table 1 and Supplementary**

Table 5-6). For example, Monkey F qualifies as a responder according to the

food drop index, but not according to food intake. Additionally, Monkey F

displays a clear V-shaped pattern in the food drop index, whereas its food

intake continues to increase across blocks. By studying the food drop index with

food intake, which can be measured simultaneously, we expect that the

characteristics of the food index can be more clearly elucidated.

In conclusion, the present test battery is a valuable tool for assessing

depression-like symptoms in common marmoset models, testing the etiology of

depression and developing novel antidepressants. The test battery we propose

has the potential to become a powerful translational tool to fill the gap between

experimental animal models and clinical depression.

Methods

Animals

Experiments were performed on 14 marmoset monkeys (*C. jacchus*, aged 2.2 ±

0.5 years, 12 male and 2 female). Three additional male marmoset monkeys

(conspecific, aged 2.4 ± 0.8 years) were used as stimuli in the PhT. All monkeys

were housed indoors at room temperature (25–28 °C with 40–60% humidity).

Light was provided from 07:00 to 19:00. Food pellets (CMS-1M, CLEA Japan

Inc, Tokyo, Japan) softened with warm water and supplemented with honey,

vitamins, and BIOFERMIN (intestinal regulators) were fed daily. Water was

provided *ad libitum*, except in the SPT. The animals were maintained and

handled in accordance with the "Guide for the Care and Use of Laboratory

Animals" by the National Research Council in the United States and the

guidelines of the National Institute of Neuroscience, National Centre of

Neurology and Psychiatry in Japan. All experimental procedures were approved

by the Animal Experimental Committee of the National Institute of Neuroscience.

Peephole Test (PhT)

An acrylic plastic apparatus (150 x 350 x 150 mm) was placed into a stainless-

steel transport cage (200 x 365 x 190 mm) that was attached to the subject's

home cage **(Fig. 1)**. The apparatus consisted of a testing chamber and a

presenting chamber with a black wall in between. The wall had a 20 mm

diameter circular window that serves as a peephole. The subject animal could

enter the testing chamber and view a stimulus in the presenting chamber

through the peephole. A transparent acrylic plate was set in the peephole to

avoid direct physical contact between the subject individual and the stimulus

individual. The subject, upon entering the test chamber, needed to duck under

the camera box and step on the step box to peep inside the presenting chamber.

The absence of extra space enabled detection of the subject's head by an

optoelectronic switch. The most important feature of the PhT device is its design,

which requires the subject's head to be positioned in a narrow space to peep

through the hole. An optoelectronic switch (E3Z-T66, OMRON Co., Kyoto,

Japan) and a small pinhole camera (KPC-S500P1, KT & C Co., Seoul, Korea)

on the back of the hole detected the response of the subject. The optoelectronic

switch consisted of an infrared emitter and receiver to detect if there was an

obstruction in between. The camera monitored from behind the subject to

simultaneously detect the subject's presence. One test trial consisted of two

distinct phases (stimulus and baseline phases), and only one stimulus a day

was presented. The stimulus was present in the stimulus phase, and the

presenting chamber was empty in the baseline phase. Two 300-s phases were

performed in a trial; the baseline phase followed by a principal stimulus phase.

The duration and frequency of VEB were recorded with an acquisition program

written in LabView (National Instruments, Austin, TX) in another room. The

duration and frequency of the baseline phase were subtracted from those of the

stimulus phase throughout the analysis. Prior to the experiment, each marmoset

was acclimated by exposure to the apparatus 13 times, with 20 min acclimation

trials once per day. Five animals were subjected to experiment-1, and all

measurements were made between 13:00 and 17:00. The PhT was conducted

at intervals of less than or equal to 7 days, and two repeated measurements

were performed for each of the five stimuli. Inanimate stimuli (object) consisted

of a model car and a model insect, and three marmosets (Conspecific 1, 2 and

3) different from the subject marmosets were prepared for the animate stimuli

(Fig. 2). The data for two trials using each stimulus was averaged and used for

further analysis.

In experiment-5, two PhT trials were performed for each block; a stimulus

animal and an object were presented one by one as the stimulus in a block. A

stimulus animal contained two marmosets (Conspecific 1 and 3), which were

different from the subject marmosets, and only a model insect was used as the

stimulus object. The presentation order was at random, and daily tests were

performed between 15:00 and 17:00 **(Fig. 6)**.

Sucrose Preference Test (SPT)

Two Falcon tubes with water nozzles each containing either tap water or

sucrose solution were attached to the front side of the home cage **(Fig. 3)**. The

automatic water supply was stopped during the test period. All fluid intake was

normalized by body weight. The preference rate was calculated according to the

following formula:

Sucrose preference percentage (%) = [(Sucrose solution intake - Tap water

intake)/Total Intake) x 100].

In experiment-2, referring to a previous study with rhesus monkey, six

concentrations (0, 0.375, 0.5625, 0.75, 1.5, and 3.0%) of sucrose solutions

were set and presented at random[41]. Seven animals were used in this

experiment. Each concentration was presented for 4 days, and the left and right

presentation positions were counterbalanced and changed daily. Each daily test

trial was performed between 14:30 on the first day and 14:00 on the second day

(23.5 hours a day). Weighing, washing the bottle, and replacement of the fluids

were performed between 14:00 and 14:30. The data over the 4 days for each

concentration was summed and used for the analysis.

In experiment-5, two test trials were performed for each block, and the

position of the 1.8% sucrose solution was changed daily. Daily tests were

performed between 14:30 and 17:30 **(Fig. 6)**. Two test results per block were

averaged and used for further analysis.

Eat Style (ES) Test and Food Intake (FI)

Food was presented for 19.5 h from 12:30 to 8:00 the following day, and the

leftover food was collected separately from two places, a feedbox and a mesh

tray **(Fig. 4a and b)**. The feedbox leftovers were defined as leftovers collected

from the feedbox, and the dropped leftovers were defined as leftovers dropped

on the mesh tray suspended below the floor of the home cage. The sum of

these two was referred to as total leftovers. Leftover food was collected and

then dried at 65 °C for one day before being weighed **(Supplementary Fig. 1a)**.

The FI was the value obtained by subtracting the total leftover food from the

given 40 g of feed. The food drop index (%) was calculated according to the

following formula:

Food drop index (%) = (Dropped leftovers (g)/Total leftovers (g)) x 100].

The food used was ordinary pellets given in daily life, and no other food or

reward was given during the experiment. Basic daily data for three consecutive

days was obtained in experiment-3, the preliminary experiment, in which 12

animals were subjected. The same procedure was used to obtain daily data in

experiment-5, in which 9 animals were used.

Motor Activity

We used home cages equipped with an infrared motion sensor (Panasonic

Electric Works Co., Osaka, Japan) on the ceiling to continually record the

motion of the animals in the animal rearing room **(Fig. 5)**. The system utilized a

passive thermographic infrared sensor to monitor the heat emitted from the

animals. The 3D localization of the heat source was monitored, and changes in

this localization were recorded as movement. Animal data was monitored and

recorded on a computer using a data acquisition program written in LabView.

Data for one week were obtained from 12 animals every 24 hours in

preliminary experiment-4. In experiment-5, activity data were acquired between

12:30 and 14:30. The 60-min activity data between 40 and 100 minutes after

the administration time of reserpine was utilized **(Fig. 6)**. The motor activity in

this time zone was converted to a one-minute average to obtain average motor

activity. Two trials for two days were averaged as block data of motor activity

and used for the analysis.

Drug Administration

Reserpine depletes monoamines in the brain by functionally inhibiting vesicular

monoamine transporter 2[29]. Reserpine (Sigma, St. Louis, MO) was dissolved in

glacial acetic acid and then diluted to 0.10 mg/mL with sterile water. Marmosets

were injected intramuscularly with 0.015 and 0.030 mg/kg reserpine or vehicle

once daily at 12:00-12:30 on the day of an experiment **(Fig. 6)**. All reserpine

and vehicle solutions contained 3.75% glacial acetic acid.

Responder and Depression-like Marmoset

Prior to experiment-5, a test battery of six principal tests was constructed by

determining the experimental schedule (timing, sequence, frequency, etc.) and

the criteria for responders and depressive-like marmosets **(Fig. 6)**. Individuals

displaying depression-like symptoms were determined using the judgment

criteria based on DSM in experiment-5 **(Supplementary Table 1)**. Ten tests

were incorporated into the experimental schedule, ensuring that all were

performed at least once in one block, and four blocks of continuous

measurement for 8 days before and after administration of the drug were

performed. Reserpine was administered in block-2 and block-3 with a dose-

escalation method, and the dosing schedule was set to be sandwiched the

reserpine blocks between block-1 and block-4, in which the vehicle was

administered. Marmosets were evaluated individually between the four blocks.

Responders to reserpine were qualified by whether the test data displayed a V-

shaped pattern over the four blocks. Individuals whose measurement data

satisfied the following conditions were qualified to be responders: Block-1 >

Block-3, Block-4 > Block-3, and Block-2 > Block-3. Depression-like marmosets

were defined as individuals who were responders in more than half (≥ four

tests) of the six tests. This condition corresponds to the DSM criteria for

depression, which requires more than half (≥ five) of the nine symptoms be

present.

Statistical Analysis

Experiment-1 and -2 data were analysed using a one-way repeated-measures

analysis of variance (ANOVA) followed by Scheffe's test[42,43] (when comparing

two groups, object vs. conspecific groups) or Shaffer's[44] (pairwise comparison)

post hoc test. The data used were confirmed to be normally distributed by

passing the Shapiro-Wilk normality test. Homoscedasticity assumptions were

investigated before the use of the ANOVA tests. $p < 0.05$ was regarded as

significant. All analyses were performed using SPSS software (Version 11.5,

SPSS Inc., Chicago, IL).

<u>Data availability statement</u>

The datasets generated during and/or analyzed during the current study are

available from the corresponding author upon reasonable request.

<u>Figure Legends</u>

Figure 1

Experimental setup and apparatus of the peephole test (PhT) designed for

common marmoset. (a) Left: the entire test apparatus during the experiment.

Right: a subject marmoset and transport box attached to the subject's home

cage. (b) Ground plan of the apparatus. (c) Section plan on the center line

drawn in (b). The inset photograph taken from the presenting chamber shows

the peeping marmoset. The red arrow and spot indicate the infrared LED beam

that detects the presence of the head.

Figure 2

Characteristics of visual exploratory behavior (VEB) depending on the type of

stimulus by means of the PhT. (a) Duration differences of VEB are shown for

five different stimuli (two objects and three conspecifics). (b) Frequency

differences of VEB are shown for five different stimuli. Data are expressed as

the mean and SD. *$p < 0.05$, Scheffe's contrast test (object vs. conspecific).

Figure 3

Setup of the sucrose preference test (SPT) and concentration-response curves.

(a) A marmoset drinking from a bottle in its home cage. (b) Consumption of tap

water and sucrose solutions against sucrose concentrations for four days. (c)

Percentage of sucrose solution intake against concentration. Data are

expressed as the mean and SD. *$p < 0.05$ compared to 0% sucrose

(Shaffer's test).

Figure 4

The eat style (ES) test and food intake (FI) in experiment-3. (a) A mesh tray

placed under the floor of the home cage retrieves dropped food pellets. A

feedbox is located on the front door of the cage, and marmosets have free

access to the given food. (b) Two marmosets reaching out from the home cage

and taking food from the feedbox. (c) The average FI and total leftover from a

preliminary experiment. (d) The average dropped and feedbox leftovers from a

preliminary experiment. Data are expressed as the mean and SD (n = 12). (e)

The distribution of all 36 data points (12 animals, three-day trials each) for the

food drop index.

Figure 5

Measurement system and circadian rhythm of motor activity. (a) An infrared

sensor is set above the ceiling of each home cage, and the acquired signal data

is sent to a control computer in a separate room. (b) 24-hour observation of

motor activity using the infrared sensor (n = 12). Black squares and bold lines

indicate hourly activity acquired over the course of a week. The yellow area

indicates the light period, and the grey areas indicate the dark period. Data are

expressed as the mean and SD.

Figure 6

Experimental schedule for the depression-like marmoset in the test battery

system in experiment-5. One block consists of 2 days, and 10 tests are

conducted in a block. Continuous measurements for 8 days (four blocks; 2 days

each) are performed, and reserpine-treated blocks (Block-2 and -3) are

sandwiched between the two vehicle-treated blocks (Block-1 and -4). Most of

the six tests are performed daily, and vital measurements and blood sampling

for cortisol are performed once per block as additional reference data.

Figure 7

Depression-like symptoms of Monkey C in the six tests of the battery. Drug-

induced V-shaped change patterns were observed in the (a) duration difference

of VEB in the PhT, (b) sucrose preference percentage in SPT, (c) food drop

index in the ES test, (d) food intake, (e) body weight, and (f) motor activity.

Table

Table 1   Responder list for each test of the battery in experiment-5.

Test name	Monkey name								
	A	**B**	**C**	D	E	F	G	H	**I**
Peephole test (PhT)	–	**R**	**R**	–	–	R	–	–	**R**
Sucrose preference test (SPT)	–	–	**R**	–	R	–	–	R	**R**
Eat style (ES) test	–	**R**	**R**	–	R	R	–	–	**R**
Food intake (FI)	R	**R**	**R**	–	R	–	–	–	**R**
Body weight (BW)	–	–	**R**	R	–	–	–	R	–
Activity	R	**R**	**R**	R	–	–	R	R	–
Number of tests qualified as responders	2	**4**	**6**	2	3	2	1	3	**4**

"R" indicates responder marmosets who displayed a V-shaped pattern,

"-" indicates non-responders who did not display a V-shaped pattern.

Bold letters indicate depression-like marmosets. If in more than half of the tests (at least four tests) the subject qualified as a responder, then the individual was judged to be a depressed-like marmoset.

References

1 Andrade, L. *et al.* The epidemiology of major depressive episodes: results

from the International Consortium of Psychiatric Epidemiology (ICPE)

Surveys. *Int J Methods Psychiatr Res* **12**, 3-21 (2003).

2 Kessler, R. C. *et al.* The epidemiology of major depressive disorder: results

from the National Comorbidity Survey Replication (NCS-R). *JAMA* **289**,

3095-3105 (2003).

3 "WHO Depression." Accessed: Mar. 20, 2024. [Online]. Available:

https://www.who.int/news-room/fact-sheets/detail/depression

4 Mathers, C. D. & Loncar, D. Projections of Global Mortality and Burden of

Disease from 2002 to 2030. *PLOS Medicine* **3**, e442 (2006).

5 Herrman, H. *et al.* Time for united action on depression: a Lancet-World

Psychiatric Association Commission. *Lancet* **399**, 957-1022 (2022).

6 American Psychiatric Association (APA). *Diagnostic and statistical manual

of mental disorders* (DSM-5). (American Psychiatric Publishing, 2013).

7 Trivedi, M. H. *et al.* Evaluation of outcomes with citalopram for depression

using measurement-based care in STAR*D: implications for clinical

practice. *Am J Psychiatry* **163**, 28-40 (2006).

Heerlein, K. *et al.* Real-world evidence from a European cohort study of patients with treatment resistant depression: Treatment patterns and clinical outcomes. *J Affect Disord* **290**, 334-344 (2021).

Willner, P. The validity of animal models of depression. *Psychopharmacology* **83**, 1-16 (1984).

Becker, M., Pinhasov, A. & Ornoy, A. Animal models of depression: what can they teach us about the human disease? *Diagnostics* **11**, 123 (2021).

"GWAS Catalog, The NHGRI-EBI Catalog of human genome-wide association studies, Major Depressive Disorder." Accessed: April 20 2024. [Online]. Available: https://www.ebi.ac.uk/gwas/efotraits/ MONDO_0002009

Mansfield, K. Marmoset models commonly used in biomedical research. *Comp Med* **53**, 383-392 (2003).

Iwanami, A. *et al.* Establishment of graded spinal cord injury model in a nonhuman primate: the common marmoset. *J Neurosci Res* **80**, 172-181 (2005).

Roeling, T. A. *et al.* Effects of unilateral 6-hydroxydopamine lesions on neuropeptide immunoreactivity in the basal ganglia of the common

marmoset, Callithrix jacchus, a quantitative immunohistochemical analysis. *J Chem Neuroanat* **9**, 155-164 (1995).

Gnanalingham, K. K., Smith, L. A., Hunter, A. J., Jenner, P. & Marsden, C. D. Alterations in striatal and extrastriatal D-1 and D-2 dopamine receptors in the MPTP-treated common marmoset: an autoradiographic study. *Synapse* **14**, 184-194 (1993).

Genain, C. P. & Hauser, S. L. Experimental allergic encephalomyelitis in the New World monkey Callithrix jacchus. *Immunol Rev* **183**, 159-172 (2001).

t Hart, B. A. *et al.* A new primate model for multiple sclerosis in the common marmoset. *Immunol Today* **21**, 290-297 (2000).

Butler, R. A. Discrimination learning by rhesus monkeys to visual-exploration motivation. *J Comp Physiol Psychol* **46**, 95-98 (1953).

Butler, R. A. Incentive conditions which influence visual exploration. *J Exp Psychol* **48**, 19-23 (1954).

Butler, R. A. & Woolpy, J. H. Visual attention in the rhesus monkey. *J Comp Physiol Psychol* **56**, 324-328 (1963).

Butler, R. A. Acquired drives and the curiosity-investigative motives. In

Principles of Comparative Psychology (eds Waters, R. H. *et al.*) 144-176

(McGraw-Hill Company, 1960).

22   Katz, R. J. Animal model of depression: pharmacological sensitivity of a

hedonic deficit. *Pharmacol Biochem Behav* **16**, 965-968 (1982).

23   Alexander, B. *et al.* Reversal of depressed behaviors in mice by p11 gene

therapy in the nucleus accumbens. *Sci Transl Med* **2**, 54ra76 (2010).

24   Nautiyal, K. M. *et al.* A lack of serotonin 1B autoreceptors results in

decreased anxiety and depression-related behaviors.

Neuropsychopharmacology **41**, 2941-2950 (2016).

25   Walker, A. K. *et al.* NMDA receptor blockade by ketamine abrogates

lipopolysaccharide-induced depressive-like behavior in C57BL/6J mice.

Neuropsychopharmacology **38**, 1609-1616 (2013).

26   Willner, P., Towell, A., Sampson, D., Sophokleous, S. & Muscat, R.

Reduction of sucrose preference by chronic unpredictable mild stress, and

its restoration by a tricyclic antidepressant. *Psychopharmacology (Berl)* **93**,

358-364 (1987).

Tomioka, I. *et al.* Transgenic monkey model of the polyglutamine diseases recapitulating progressive neurological symptoms. *eNeuro* **4**, e0250-16.2017; 10.1523/ENEURO.0250-16.2017 (2017).

Ross, C. N., Power, M. L., Artavia, J. M. & Tardif, S. D. Relation of food intake behaviors and obesity development in young common marmoset monkeys. *Obesity* **21**, 1891-1899 (2013)

Kandel, E.R. Disorders of mood: depression, mania, and anxiety disorders. In *Principles of Neural Science* (eds Kandel, E.R. *et al.*) 1209-1226 (McGraw-Hill Company, 2000)

Goodwin, F. K., Ebert, M. H., Bunney, W.E. Mental effects of reserpine in man: A review. In *Psychiatric Complications of Medical Drugs* (ed by Shader, R.I.) 73–101 (Raven Press, 1972).

Mann, T. M., Williams, K. E., Pearce, P. C. & Scott, E. A. A novel method for activity monitoring in small non-human primates. *Lab Anim* **39**, 169-177 (2005).

Lemieux, G., Davignon, A. & Genest, J. Depressive states during Rauwolfia therapy for arterial hypertension; a report of 30 cases. *Can Med Assoc J* **74**, 522-526 (1956).

Quetsch, R. M., Achor, R. W., Litin, E. M. & Faucett, R. L. Depressive reactions in hypertensive patients; a comparison of those treated with Rauwolfia and those receiving no specific antihypertensive treatment. *Circulation* **19**, 366-375 (1959).

McKinney, W. T., Jr., Prange, A. J., Jr., Majchowicz, E. & Schlesinger, K. Plasma corticosterone changes following alterations in brain norepinephrine and serotonin. *Dis Nerv Syst* **32**, 308-313 (1971).

Takemoto, A., Izumi, A., Miwa, M. & Nakamura, K. Development of a compact and general-purpose experimental apparatus with a touch-sensitive screen for use in evaluating cognitive functions in common marmosets. *J Neurosci Methods* **199**, 82-86 (2011).

Oikonomidis, L. *et al.* A dimensional approach to modeling symptoms of neuropsychiatric disorders in the marmoset monkey. *Dev Neurobiol* **77**, 328-353 (2017).

Kinoshita, M. *et al.* Genetic dissection of the circuit for hand dexterity in primates. *Nature* **487**, 235-238 (2012).

Nagai, Y. *et al.* PET imaging-guided chemogenetic silencing reveals a critical role of primate rostromedial caudate in reward evaluation. *Nat Commun* **7**, 13605; 10.1038/ncomms13605 (2016).

Schardein, J. L., Schwetz, B. A. & Kenel, M. F. Species sensitivities and prediction of teratogenic potential. *Environmental Health Perspectives* **61**, 55-67 (1985).

Barros, M., Boere, V., Huston, J. P. & Tomaz, C. Measuring fear and anxiety in the marmoset (Callithrix penicillata) with a novel predator confrontation model: effects of diazepam. *Behav Brain Res* **108**, 205-211 (2000).

Paul, I. A., English, J. A. & Halaris, A. Sucrose and quinine intake by maternally-deprived and control rhesus monkeys. *Behav Brain Res* **112**, 127-134 (2000).

Escartin, C. *et al.* Activation of astrocytes by CNTF induces metabolic plasticity and increases resistance to metabolic insults. *J Neurosci* **27**, 7094-7104 (2007).

Sun, Y. J., Kim, Y. J., Ibrahim, L. A., Tao, H. W. & Zhang, L. I. Synaptic mechanisms underlying functional dichotomy between intrinsic-bursting

and regular-spiking neurons in auditory cortical layer 5. *J Neurosci* **33**,

5326-5339 (2013).

44   Mitsuyasu, H. *et al.* Flexion contracture persists if the contracture is more

than 15 degrees at 3 months after total knee arthroplasty. *J Arthroplasty* **26**,

639-643 (2011).

Acknowledgements

We greatly thank all members in the Animal Models for Human Disease, National Institute of Neuroscience, National Center for Neurology and Psychiatry, for their technical assistance. This study was funded by research grant 20B-10 for Nervous and Mental Disorders from the Ministry of Health, Labor and Welfare of Japan, "Development of nonhuman primate models for mental and nervous diseases," and partly supported by a research grant from the Astellas Foundation for Research on Metabolic Disorders.

Author contributions

All authors conceived the experiments. H.Y. and H.I. performed experiments. K.N. and H.I. acquired financial support. H.Y. analyzed the data. All authors wrote the manuscript, discussed the data and commented on the manuscript.

Competing interests

The authors declare no competing interests.

Figure 1

a

b

c

Figure 2

a

b

Figure 3

a

b

c

Figure 4

a

Feedbox   Mesh tray

b

Mesh tray   Feedbox

c

d

e

Figure 5

a

b

Figure 6

Figure 7

a

b

c

d

e

f