Acute rapamycin treatment reveals novel mechanisms of behavioral, physiological, and functional dysfunction in a maternal inflammation mouse model of autism and sensory over-responsivity

JE Le Belle[a,b,m,] M Condro[a], C Cepeda[a,i], KD Oikonomou[a], K Tessema[a], L Dudley[a], J Schoenfield[b], R Kawaguchi[a], D Geschwind[a,e,f,k], AJ Silva[g,h,i,j,k], Z Zhang[l], K Shokat[l], NG Harris[b,k,m], & HI Kornblum[a,c,d,i,k,m]

[a] Semel Institute for Neuroscience & Human Behavior, Department of Psychiatry, David Geffen School of Medicine, UCLA, Los Angeles, CA
[b] The UCLA Brain Injury Research Center, Department of Neurosurgery, David Geffen School of Medicine, UCLA, Los Angeles, CA
[c] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095, USA
[d] Jonsson Comprehensive Cancer Center, David Geffen School of Medicine at UCLA, Los Angeles, CA
[e] Program in Neurogenetics, Department of Neurology, David Geffen School of Medicine, UCLA, Los Angeles, CA
[f] Department of Human Genetics, UCLA, Los Angeles, CA
[g] Department of Neurobiology, UCLA, Los Angeles, CA
[h] Department of Psychology, UCLA, Los Angeles, CA
[i] Department of Psychiatry and Biobehavioral Sciences, David Geffen School of Medicine, UCLA, Los Angeles, CA
[j] Integrative Center for Learning and Memory, UCLA, Los Angeles, CA
[k] Brain Research Institute, UCLA, Los Angeles, CA
[l] Department of Cellular and Molecular Pharmacology, Howard Hughes Medical Institute, University of California, San Francisco, CA
[m] UCLA Intellectual and Developmental Disabilities Research Center, Semel Institute, Department of Psychiatry, David Geffen School of Medicine, UCLA, Los Angeles, CA

Abstract

Maternal inflammatory response (MIR) during early gestation in mice induces a cascade of physiological and behavioral changes that have been associated with autism spectrum disorder (ASD). In a prior study and the current one, we find that mild MIR results in chronic systemic and neuro-inflammation, mTOR pathway activation, mild brain overgrowth followed by regionally specific volumetric changes, sensory processing dysregulation, and social and repetitive behavior abnormalities. Prior studies of rapamycin treatment in autism models have focused on chronic treatments that might be expected to alter or prevent physical brain changes. Here, we have focused on the acute effects of rapamycin to uncover novel mechanisms of dysfunction and related to mTOR pathway signaling. We find that within 2 hours, rapamycin treatment could rapidly rescue neuronal hyper-excitability, seizure susceptibility, functional network connectivity and brain community structure, and repetitive behaviors and sensory over-responsivity in adult offspring with persistent brain overgrowth. These CNS-mediated effects are also associated with alteration of the expression of several ASD-,

ion channel-, and epilepsy-associated genes, in the same time frame. Our findings suggest that mTOR dysregulation in MIR offspring is a key contributor to various levels of brain dysfunction, including neuronal excitability, altered gene expression in multiple cell types, sensory functional network connectivity, and modulation of information flow. However, we demonstrate that the adult MIR brain is also amenable to rapid normalization of these functional changes which results in the rescue of both core and comorbid ASD behaviors in adult animals without requiring long-term physical alterations to the brain. Thus, restoring excitatory/inhibitory imbalance and sensory functional network modularity may be important targets for therapeutically addressing both primary sensory and social behavior phenotypes, and compensatory repetitive behavior phenotypes.

Introduction

Neurodevelopmental disorders result from the disruption of brain development *in utero* or in early life[1]. Genetic, environmental, epigenetic, and immunological factors have all been demonstrated to contribute to their complex pathogenesis[1–5]. For example, there are numerous different anatomical abnormalities in brain structure and cellular changes that have been observed in the brains of autistic individuals and in animal models of autism that indicate a multifactorial etiology[3,4,6–19]. Despite this complex etiology and the spectrum of phenotypes in Autism Spectrum Disorders (ASD), a common pathogenic mechanism has been identified for both genetic and non-genetic factors contributing to several neurodevelopmental disorders: dysregulated mTOR pathway signaling in the brain[6,7,20–31].

Abnormal PI3K/Akt/mTOR signaling pathway activity in response to divergent underlying causes has been shown to lead to aberrant brain structure, synapse formation, synaptic pruning, excitatory/inhibitory balance, neuro-inflammatory signaling, and brain size[7,7,30,32–55]. Hyperactive mTOR signaling is known to contribute to seizures, sensory sensitivity, intellectual disability, macrocephalic brain growth, and a number of ASD-associated

behavioral abnormalities[4,5,26,56–59]. This is thought to be related to the role of this important pathway in controlling cell growth, survival, energy balance, proliferation, autophagy, and ion channel expression. The mTOR pathway can become hyper-activated by several known mutations in ASD-associated genes such as PTEN, TSC1, TSC2, FMR1, Mecp2, DEPDC5, CACNA1C, UBE3A, and CNTNAP2[28,60–66]. ASD behaviors, macrocephaly, and epilepsy have also been recapitulated in a variety of PTEN and TSC murine knockout models in different cellular populations, developmental timeframes, and quantity of gene deletion. Furthermore, the Simons Foundation Autism Research Initiative (SFARI) has also identified a number of high-confidence autism risk genes in humans with several belonging to mTOR signaling gene sets[63,67–70].

However, genetic mutations are not the only cause of increased mTOR activation and brain overgrowth. Hyper-activated mTOR signaling can both induce and be induced by neuro-inflammation of varying etiology[57,71–82]. There is considerable evidence of an elevated inflammatory state in ASD from gene expression studies, pro-inflammatory cytokines found in the blood and CSF and from increased activated microglia found throughout post-mortem autistic brains in both children and adults[19,83–97]. There is also evidence of brain overgrowth in ASD that is not linked to any specific genetic variation. For example, Courchesne et al., and others have demonstrated that there is a limited period of increased , albeit not macrocephalic, brain growth early in development that has no known genetic or non-genetic etiology to date[98–100]. Maternal immune system activation is a good candidate mechanism for this phenomenon because it is also known to contribute to brain overgrowth and ASD in humans and in rodent models[19,85,101–117]. We have previously shown that exposure to a low-level Maternal Inflammatory Response (MIR) induced by lipopolysaccharide early in gestation produces offspring with mild brain overgrowth, hyperactive mTOR signaling in the neurogenic SVZ niche, and abnormal ASD-related behaviors in mice[118].

In addition to the core features of ASD (atypical social, communication, and stereotyped behaviors), there are many common comorbid phenotypes including sensory over-responsivity (SOR), epilepsy, disordered sleep, gastrointestinal disturbances, anxiety, depression, and irritability[119] (Hazen et al., 2014). Rather than approaching treatment as a "cure" for ASD, a common approach for current pharmacological therapy is focused on treating these non-core symptoms with existing approved drugs. The rationale for treating each aberrant phenotype separately rather than tackling core symptomology is the complex and not fully understood etiology of ASD, the spectrum of ASD phenotypes and severity, and a lack of a core molecular target that underlies multiple aberrant behaviors. However, mTOR signaling is potentially one such convergent target that may impact both core and non-core phenotypes. Rapamycin, an inhibitor of mTOR, is known to be immunosuppressive, anti-proliferative, and an inducer of autophagy. It has been used in pre-clinical models of PTEN and TSC gene mutations to rescue the epileptic phenotype and several ASD-related behaviors in young mice by successfully preventing the development of aberrant brain physiological changes[9,120–124]. For example, developmental inhibition of mTOR over several weeks prevents brain overgrowth, loss of inhibitory neurons, seizures, and abnormal synaptic connectivity[9,79,125–130]. Rapamycin has, perhaps surprisingly, also been shown to rescue several abnormal phenotypes in adult mice despite the persistence of physiological brain abnormalities associated with PTEN and TSC gene mutations and mTOR activation, including macrocephalic brain overgrowth[9,19,124,125,131,132]. To date it has been used clinically to treat patients with ASD and PTEN or TSC gene mutations for weeks to months with mixed results, improving some cognitive functions, reducing seizure frequency, and improving social behaviors in some studies but not in others[133–145]. Chronic Rapamycin treatment has also been shown to rescue some phenotypes in idiopathic ASD mouse models with elevated mTOR signaling such as in utero valproic acid exposure, ASD-seizure, and BTBR models[122,125,146–149].

While there has been encouraging progress at rescuing mTOR-mediated brain pathology and behavioral abnormalities in pre-clinical models with chronic rapamycin treatments, there could still be risks to normal brain

development with juvenile treatment in humans without specific mTOR pathway-related gene mutations and adverse side effects from the use of such a potent growth inhibitor and immune suppressor. Therefore, we wanted to further understand the mechanisms that may underlie the effects of adult mTOR inhibition that isn't aimed at preventing or reversing structural brain abnormalities. Furthermore, our study focuses on the effects of mTOR inhibition within an acute timeframe, which has revealed new mechanisms and functional changes that are different to those reported with a chronic inhibition treatment approach. Specifically, we have assessed the effects of acute (2 hour) rapamycin treatment of mTOR activation in adult offspring following mild MIR on behavior, sensory processing, brain functional connectivity and modularity, electrophysiological properties, and effects on specific cellular compartments and expression of ASD-, epilepsy-, and ion channel-associated genes. Thus, with these findings we have identified several potentially less disruptive targets for pharmacological or neuromodulatory intervention in several ASD physiological and behavioral phenotypes in the CNS.

Materials and Methods

Animals

Time-mated, pregnant CD-1 wild-type mice were obtained from Charles River Laboratories (CA, USA). Maternal Inflammatory Response (MIR) was induced with low-dose lipopolysaccharide (LPS, E. coli serotype 0111:B4; 0.008 mg/kg, Sigma) administered in a single I.P. injection on E9 of gestation as described in Le Belle, et al., 2014. Offspring were housed 4 per cage on a 12-hour light/dark cycle with ad libitum access to food and water. Behavioral experiments were performed during the light phase. All experimental procedures were conducted in accordance with the University of California, Los Angeles, (UCLA) Chancellor's Animal Research Committee and the Public Health Service Policy on Humane Care and Use of Laboratory Animals.

Blood Cytokine Analysis

Blood was collected from pregnant dams after LPS injection and from adult MIR and control offspring. Blood plasma was separated and frozen before analysis by the UCLA Pathology Immune Assessment Core using the Luminex's xMAP® Mouse 32-plex Cytokine/Chemokine Panel Immunoassay according to standard assay methods.

Regional Brain Microdissection and Western Blot

The sensory-motor cortex, amygdala, and striatum were microdissected from adult offspring and flash frozen for standard western blot analysis. Tissue homogenates (cytosol) were lysed in RIPA buffer (25 mM Tris-HCl, 150 mM NaCl, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS, pH 7.6 at 4℃) containing a cocktail of protease inhibitors (Calbiochem). Equal amounts of protein were separated by SDS-PAGE (4–12% Bis-Tris gels, Invitrogen) and transferred to PVDF membranes (Invitrogen), blocked in TBST plus 5% non-fat milk and then incubated with the following primary antibodies: phospho-S6 (S235/236; 1:2000, Cell Signaling, #4858), S6 (1:2000, cell signaling, #2217), and b-actin (1:2000, Cell Signaling, #4970) overnight at 4℃. The membranes were then washed with TBST and incubated for 1 h at room temperature. The washed membranes were then treated with an enhanced chemiluminescence detection reagent (Thermo Scientific). All Blots were developed using ChemiDoc XRS+ Molecular Imager (BioRad) and analyzed using Quantity One software (BioRad). Band densities were normalized to the total amount of protein loaded per lane using Sypro Ruby (BioRad).

Behavioral Testing

The three-chamber social interaction test was performed on adult mice according to the methods of Silverman, et al.[150]. Measurement of repetitive behaviors in an open field chamber was performed according to the methods of Peñagarikano et al.[151]. Pre-pulse inhibition and habituation using a tactile air-puff prepulse and audible startle was tested according to the methods of Orefice, et al.[152]. The Von Frey fiber tactile sensitivity test was performed according to Bonin, et al.[153]. The tactile avoidance light/dark box is a novel method developed

for this study to examine tactile sensitivity of the paws and avoidance behaviors. This test uses the classic light/dark chamber box which is traditionally a test for anxiety in mice where normal mice will explore both light and dark chambers but mice prefer the dark chamber and mice with heighted anxiety will spend much more time in the "safe" dark chamber compared to normal controls[154]. We adapted this test to use either a smooth plastic floor or a rough textured floor in one chamber using rough, anti-slip stair adhesive strips from a local hardware store. We quantified the time mice spent in the left or right chamber when both were lit and the floor was smooth plastic (trial 1), when both were lit but the floor in one chamber was the rough texture (trial 2), when one chamber was dark but both floors were smooth (trial 3, the classic light/dark box test), and finally when the dark (preferred) chamber had the rough textured floor and the lit chamber had a smooth floor (trial 4). The time spent in each chamber during a 10-minute test was quantified and compared between groups. Spending less time in the normally preferred dark chamber when the floor was a rough texture but not when the floor was smooth demonstrates an aversive tactile sensitivity. See Figure 4A for a visual summary of this sensory test. Mice that were examined with multiple behaviors were given the tests on different days from the least to most stressful: open field, 3-chamber social, tactile light/dark box, Von Frey, and finally pre-pulse inhibition of startle. The relative severity of behavioral abnormalities is categorized as follows: mild (15-48% difference to control average), moderate (46-80% difference to control average), and severe (81-300% difference to control average).

Single Cell and Bulk Genetic Sequencing and Analysis

Mice were euthanized according to approved protocols and sensory-motor cortex tissue was microdissected from MIR and control offspring and flash frozen before being prepared for sequencing analysis. For bulk RNA sequencing, 12 MIR and 12 control mice were given rapamycin or vehicle 2 hours before euthanizing and flash freezing the brain tissue for sequencing. RNA is isolated using a Qiagen RNeasy Mini kit (Cat. 74106) followed by library preparation using a KAPA Stranded mRNA-Seq Kit (Cat. KR0960). The sequencing libraries were pooled

and sequenced to generate minimum of 220 M 50 paired-end reads on NovaSeq6000 (Illumina). Reads were aligned to mouse mm10 genome reference using STAR aligner (v2.4.0). Read counts for refSeq genes were generated by HT-seq 0.6.1. Various QC were performed after alignment to examine the level of mismatch rate, mapping rate to the whole genome, repeats, chromosomes, key transcriptomic regions (exons, introns, UTRs, genes), insert sizes, AT/GC dropout, transcript coverage and GC bias. Outliers were removed based on QC results. Differential expression analysis was conducted with R-project and the Bioconductor package EdgeR. Differentially expressed genes (DEGs) between groups were defined as those meeting all of the following criteria: false discovery rate (FDR) < 0.05, LogFC > 0.5, and p < 0.05 threshold (see supplemental data for complete gene list). We also tested for overlap between these DEGs and high-confidence autism-associated genes in ASD databases[68,96].

For single-nucleus RNA sequencing, 6 MIR and 6 control mice were analyzed. Single nuclei were isolated from frozen brain tissue using iodixanol-based density gradient centrifugation and submitted to UCLA Technology Center for Genomics and Bioinformatics for library preparation (via Chromium Single Cell 3' v3 kit from 10x Genomics) and sequencing (via NovaSeq 6000 S2 platform from Illumina). Cell Ranger (10x Genomics) was used for alignment, filtering, barcode identification, and generation of gene-cell matrices for each sample. Ambient RNA was removed using CellBender[155]. Pre-processing (minimum cells 3, minimum features 200, doublet removal via scDblFinder[156], maximum mitochondrial content 5%), integration (3000 features), and clustering (resolution 0.8) were performed using the R package Seurat[157–159]. Canonical lineage markers were used to identify cell types (microglia, astrocytes, oligodendrocytes, pericytes, endothelial cells, upper layer and deep layer excitatory neurons, and interneurons). Differential abundance of cell types in MIR vs CTRL was assessed using the R package propeller[160]. Differential expression testing was performed using the pseudobulk approach (edgeR with likelihood ratio test) via the R package Libra[161]. Gene set enrichment analysis (GSEA)[162] was performed to assess differential enrichment of the "Hallmark mTORC1 Signaling" gene set[163,164] using the R

package fgsea[165]. The above comparisons underwent correction for multiple comparisons using the native functionality of the respective packages. Significance was defined as FDR or adjusted-p < 0.05.

Brain Slice Electrophysiology

Adult (2-6 mo) control and MIR mice were deeply anesthetized with isoflurane and perfused transcardially with ice-cold, high-sucrose slicing solution containing (in mM): 26 $NaHCO_3$, 1.25 NaH_2PO_4, 208 sucrose, 10 glucose, 2.5 KCl, 1.3 $MgCl_2$, and 8 $MgSO_4$. Mice were swiftly decapitated, brains extracted and immediately placed in oxygenated high-sucrose slicing solution. Coronal slices containing both sensorimotor cortex and dorsolateral striatum were cut at 300 μm and transferred to an incubating chamber containing artificial cerebrospinal fluid (ACSF) (in mM): 130 NaCl, 5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 2 $MgCl_2$, 2 $CaCl_2$, and 10 glucose) oxygenated with 95% O_2–5% CO_2 (pH 7.2–7.4, osmolality 290-310 mOsm/L, 32–34°C). Slices were allowed to recover for 60 min and electrophysiological recordings were obtained at room temperature. Cortical pyramidal neurons (CPNs) in layers 2/3 or striatal medium-sized spiny neurons (MSNs) were visually identified using infrared illumination with differential interference contrast optics (IR-DIC). Whole-cell patch clamp recordings in voltage- or current-clamp modes were obtained using a MultiClamp 700B Amplifier (Molecular Devices) and the pCLAMP 10.5 acquisition software (Molecular Devices). For CPN recordings (in voltage and current clamp), the internal pipette solution contained (in mM): 112.5 K-gluconate, 4 NaCl, 17.5 KCl, 0.5 $CaCl_2$, 1 $MgCl_2$, 5 ATP (dipotassium salt), 1 NaGTP, and 10 HEPES (pH 7.2, 270–280 mOsm/L). For striatal MSN recordings (in voltage clamp only), the internal pipette solution contained (in mM): 130 Cs-methanesulfonate, 10 CsCl, 4 NaCl, 1 $MgCl_2$, 5 MgATP, 5 EGTA, 10 HEPES, 5 GTP, 10 phosphocreatine, and 0.1 leupeptin (pH 7.2, 270 mOsm). Electrode impedance was typically 4-5 MΩ in the bath. Cells with access resistance exceeding 25 MΩ were discarded. Cells were first voltage clamped at -70 mV to assess passive membrane properties (membrane capacitance, input resistance and decay time constant) and measured by applying a 10 mV depolarizing step voltage command and using the membrane test function integrated in the pClamp10 software. Active membrane properties of CPNs

(resting membrane potential, intrinsic excitability, and rheobase) were determined in current clamp mode and by injecting current pulses of increasing intensity. Rheobase (minimal current intensity required to evoke an action potential), consisted of a single depolarizing current pulse (5 ms duration) of increasing intensity (Δ=15 pA). Spontaneous synaptic currents were recorded at -70 mV holding potential in the presence (MSNs) or absence (CPNs) of Bicuculline (BIC, 10-20 µM), a competitive antagonist of GABAA receptors, to better isolate glutamatergic events. Spontaneous synaptic currents were analyzed using the MiniAnalysis software (version 6.0, Synaptosoft, Fort Lee, NJ). The effects of rapamycin *ex vivo* (1-2 µM) on synaptic activity were examined by incubating slices from control or MIR mice for 2 hr before and during electrophysiological recordings. In vivo rapamycin treatment was also given to MIR mice (5 mg/kg, I.P.) or vehicle (DMSO) 2 hours before decapitation and slice electrophysiology.

Magnetic Resonance Imaging

Adult MIR and Control offspring were imaged using a 7 Tesla (T) Biospec small animal MRI system using Paravision 5.2 software (Bruker) at the UCLA Brain Mapping Center. Mice were briefly anesthetized with 2% isoflurane vaporized in oxygen flowing at 1 L/min, then placed on an MRI-compatible cradle and a single-channel surface coil (Bruker) secured over the head. For resting-state scans, a 3 mm-thick agar gel cap (Sigma, 3% in distilled water) was placed between the head and the surface coil, in order to reduce signal distortion in the blood-oxygen-level-dependent (BOLD) signal[166]. To minimize time-dependent effects of the anesthetic on the BOLD signal, these initial steps were performed within a 10–15-minute time window[167]. Isoflurane was gradually discontinued and sedation was initiated with a single subcutaneous (s.c.) injection of dexmedetomidine (Dexdomitor®, Zoetis; 0.15 mg/kg) followed by continuous s.c. infusion at 0.3 mg/kg/hr throughout the duration of the imaging as described in a previous publication[166]. Respiration and body temperature of the mouse were continuously monitored remotely, and maintained in a physiological range (37 ± 1 °C; Small Animal Instruments Inc.) by a homeothermically-controlled forced warm air over the body (SA11 Instr, Inc., USA). At the end of the

imaging session, sedation was reversed by atipamezole (Antisedan®, Pfizer) at 1.5 mg/kg (i.p.). Data were

acquired using the S116 Bruker gradients (400 mT/m) in combination with a single-channel surface coil

(described above) and a 72 mm birdcage transmit coil. An initial series of scans was performed to confirm

proper head position, then localized FASTMAP shimming was performed to improve field homogeneity. T2-

weighted structural scans were acquired with a Rapid-Relaxation-with-Enhancement (RARE) sequence (RARE

factor=8, Echo time (TE)=56 ms, repetition time (TR)=6,018 ms, 4 averages, field-of-view (FOV)=1.8× 1.2cm, slice

thickness=0.5 mm, 18 slices, FA=90 deg, bandwidth (BW)=50 kHz, matrix=60×40). Then, 45-minute long

functional (BOLD) data were acquired using the same image geometry as the structural scans, with a one-shot,

interleaved, gradient-echo echo planar imaging sequence with the following parameters: TE=16 ms, TR=2,000

ms, FA=70 degrees, BW 400 kHz and a data matrix of 60×40). 10 dummy scans were used to allow the T1 signal

to reach steady-state prior to signal acquisition. Resting-state functional data were acquired using above-

described parameters, with the following modifications: field-of-view (FOV)=3.0×3.0 cm, matrix=128×64;

functional (BOLD) scans acquired using a two-shot gradient-echo echo-planar imaging (GE-EPI) sequence, TE=19

ms, TR=1,000 ms (2,000 ms per volume), FA=70 degrees, BW 400 kHz and a data matrix of 128×64, Fourier

transformed to 128×128, voxel resolution 0.23×0.23×0.5 mm). For each mouse, baseline data were collected

and on the following week rapamycin (5mg/kg) or vehicle control (DMSO) was injected i.p. 2 hours before they

were imaged in an identical manner.

MRI Analyses

Data were converted to nifti format and entered into a python-scripted, preprocessing pipeline to do the

following steps: brain extraction[168], movement correction[169] and co-registration to a study-created brain

template[170], slice timing correction, Gaussian smoothing to 0.8mm, bandpass filtering 0.01-0.2Hz using the FSL

Toolbox[171]. Data were masked for inclusion of brain voxels common to all mice to prevent bias. A mouse brain

atlas (Ekam Solutions, USA) was registered to the study brain template, masked for common brain regions, and

then inverse warped back to subject space. Functional adjacency matrices were calculated from the pre-processed, time-series rsfMRI data using 136 brain regions common to all brains to obtain Pearson correlation coefficients that were then Fisher z-transformed. Statistical contrasts were generated between MIR and Control mice and within group between baseline and RAPA intervention using network-based statistics[172] implemented in GraphVar[173]. Subnetwork edge connectivity was evaluated at P<0.01, 2-tailed, using non-parametric permutation testing over 1000 iterations. Seed analysis was conducted by temporal correlation of whole brain data to mean signal extraction from regions of interest at the pre-processed subject-level data, and then at the group-level within the general linear model framework implemented by using FSL Feat, with statistical correct at 2.1z, P<0.01 cluster-based correction[174]. Distribution of brain networks into a community architecture was evaluated by calculation of modularity using 1000 iterations of the Louvain algorithm[175]. Modules derived from this process were then further classified at the nodal level by computing the classification diversity and classification consistency as described previously[176] as implemented under GraphVar.

Tensor-based deformation analysis: Structural RARE data were converted to nifti image format, brain extracted from extracranial tissue using FSL BET[177]. All data were used to construct a study-specific, mean deformation template (MDT) images using the ants Multivariate Template Construction script[178] from Advanced Normalization Tools (ANTs, v.2.3.3)[179] that consisted of bias field correction and a three-stage coregistration procedure of rigid, affine and non-linear deformable registrations. The raw data were then separately coregistered to the MDT template using a two-stage affine and symmetric diffeomorphic registration under ANTs. The resulting deformation field was then used to derive the Jacobian determinant that provides an indication of the local tissue expansion or atrophy compared to the MDT. All data were masked for a common space to ensure a contribution from all mice to each voxel, following which statistical group differences in voxel-based Jacobian values were computed using non-linear permutation testing implemented under FSL Randomise[180] under the general linear model framework using cluster-based thresholding set at z=3.1 (P<0.001)

and variance smoothing of 0.1mm. Data were then corrected for multivoxel comparison using false discovery rate correction at q=0.01[181,182].

Rapamycin Treatments

Mice were treated with 5 mg/kg dose of rapamycin (Sigma-Aldrich) I.P. or vehicle control (DMSO)[124]. All behavioral analyses, electrophysiology, MRI, and tissue collection for sequencing was performed at 2-hours post-injection except for the 3-chamber social interaction test which was performed after 3 daily injections of rapamycin at the same concentration. See figure 3A for experimental paradigm with acute rapamycin treatments. Mice treated with chronic rapamycin for 5 weeks also received a daily 5 mg/kg dose of rapamycin (Sigma-Aldrich) I.P. or vehicle control (DMSO; Figure 3F).

RapaBlock Treatments

RapaBlock is a brain impermeable small molecule [183,184] that binds FKBP12, and acts to prevent access to the necessary factor for mTOR inhibition in the peripheral nervous system. Therefore, mice co-administered both rapamycin and RapaBlock will still have mTOR inhibition in the central but not in the peripheral nervous system. RapaBlock is dissolved in DMSO with and made into solution with Tween80, PEG300, and DPBS. Mice were injected simultaneously with 5 mg/kg dose of rapamycin or vehicle control (DMSO) and with 40 mg/kg of RapaBlock or vehicle control, given via intraperitoneal injection in opposite flanks. RapaBlock was a generous gift from the Shokat lab (UC San Francisco).

S6K1 inhibitor Treatment

The S6K1 inhibitor PF-4708671 was dissolved in DMSO. Both vehicle and PF-4708671 were formulated in 10% DMSO, 10% tween 80 and 80% water. PF-4708671 (75mg/kg) or vehicle was given via intraperitoneal injection of PF-4708671 as described in Huang, et al.[185].

Plexxikon Microglia Depletion

Mice were fed a research diet containing the CSF1R-inhibitor, Plexxikon 5622 (Plexxikon-Daiichi, San Francisco) to deplete brain microglia. PLX5622 was provided by Plexxikon Inc. (Berkeley, CA) and was formulated in standard AIN-76A rodent chow at a concentration of 1200 mg/kg (Research Diets, New Brunswick, NJ) and provided ad libitum. Control mice were given standard AIN-76A rodent chow. Mice were fed the Plexxikon or control diet from Research Diets for 3 weeks prior to and during behavioral testing of young and older adult mice. Immunohistochemistry was used to confirm >98% microglia depletion from the diet at the end of the experiments.

Immunohistochemistry

Brains were flash frozen and cut on a cryostat at 30uM. Brain sections were post-fixed on slides with 4% paraformaldehyde before being permeabilized with 3% TritonX-100 and immunostained with IBA1 (AbCam), NeuN (ThermoFischer), and Egr1 (Cell Signaling Technology) primary antibodies followed by the appropriate Alexa fluorescent secondary antibodies (ThermoFischer), and Hoescht counterstain. Positive cells were quantified on a Zeiss AxioImager Z1 fluorescent microscope with Apotome 2.0, using the unbiased optical fractionator approach (StereoInvestigator; MicroBrightField).

Seizure Susceptibility

Mice were tested for relative seizure susceptibility using a modified protocol from Shimada and Yamagata[186] in which an escalating dose of the seizure-inducing drug, Pentylenetetrazole (PTZ), a $GABA_A$ receptor antagonist, is used to establish the relative seizure susceptibility of different groups of animals. An intraperitoneal injection of PTZ into an animal induces an acute, severe seizure at a high dose, but sequential injections of a subconvulsive dose are used for the development of chemical kindling, an epilepsy model. With this method,

vulnerability to PTZ-mediated seizures can be estimated. PTZ was dissolved sy 2 mg/mL in sterile 0.9% (w/v) NaCl and prepared on the day of use. Mice were injected i.p. with a starting dose of 10 mg/kg and then injected again every 2 days, increasing the dose by 10mg/kg. Mice were observed for the following seizure-related escalating behaviors for 30 minutes following the injection: (0) no seizure behaviors, (1) immobilization, (2) head nodding and partial myoclonus (sudden, brief involuntary twitching or jerking of a muscle or group of muscles), (3) continuous whole body myoclonus, (4) rearing or tonic seizure (falling on the side followed by forelimb tonic contraction and hindlimb tonic extension), (5) tonic-clonic seizure with rushing and jumping. Both control and MIR mice with and without acute rapamycin treatment were given escalating doses of PTZ until all mice from one group were displaying seizure-like activity (1-5) and the number and severity of seizure symptoms were then compared between groups to determine relative induced seizure susceptibility.

Statistical Analyses

   All data are expressed as mean ± SEM unless otherwise indicated; t tests were performed using Microsoft Excel to determine the statistical significance of treatment sets. For multiple comparisons, one- or two-way ANOVA was performed as appropriate, and Bonferroni or Holm-Sidak post hoc t tests were done to determine significance. The Spearman rank order test was used for correlation analysis. Alpha values were 0.05 except when adjusted by the post hoc tests.

Results

Chronic pathophysiology in MIR offspring

In our mouse model of mild maternal inflammation, we have previously shown that juvenile and young adult mice have mild brain overgrowth and increased PI3K/AKT/mTOR pathway activation in the neurogenic SVZ niche[118]. In our current study we have observed that the pathophysiologic changes caused by exposure to MIR continue to persist and evolve throughout adulthood in the offspring, including chronic systemic (Fig 1B) and brain (Fig 1D) neuro-inflammation, chronic mTOR pathway activation (Fig 1E), abnormal social and repetitive behaviors (Fig 1F-G), enhanced sensory sensitivity (Fig 1H), and changes in regional volumetric brain sizes (Fig2). In order to determine what might be driving this long-term dysfunction in our neuro-inflammatory MIR model we examined blood cytokines in the young and old adult MIR offspring compared to age-matched control offspring and found evidence of an ongoing, chronically elevated inflammatory state with several pro-inflammatory cytokines remaining elevated in blood at all ages (Fig 1B). Large-fold increases in several pro-inflammatory cytokines were observed in blood samples from MIR offspring compared to controls in both young and old adults in G-CSF, IFN-g, IL-6, MCP-1, MIG, KC, M-CSF, GM-CSF and there were also smaller but significant increases in IL-9, IL-1beta, TNF-alpha, VEGF, IP-10, and LIF (Fig 1B; $p<0.01$). The largest increases in pro-inflammatory cytokines in the young adult MIR mice were in IFN-g, MCP-1, and IL-6 and in old adult mice were M-CSF, MCP-1, and G-CSF (Fig 1B; $p<0.001$). Acute 2-hour treatment with the mTOR inhibitor rapamycin had negligible effects on blood cytokines in adult MIR and control offspring (Supplemental Fig S1A). Therefore, the effects of acute rapamycin in our experiments are not directly related to reducing the on-going systemic inflammation in MIR offspring. Immunohistochemistry staining and quantification of microglia (IBA1+) further showed that this chronic, systemic inflammation is also affecting the brain with chronically elevated brain microglia populations from before birth to old adulthood in MIR offspring compared to controls (Fig 1D). Furthermore, western blot analysis of mTOR pathway activation in tissue microdissections from sensory-motor

cortex, striatum, and amygdala, regions associated with abnormal ASD behaviors, using phospho-S6 antibody show elevated mTOR pathway activity in the combined brain regions in MIR compared to control adult offspring (Fig 1E). Thus, both systemic and brain-specific inflammation and mTOR pathway signaling is elevated throughout the lifespan in this mild maternal inflammation model.

One of our hypotheses about the mechanisms behind the effects of MIR on behavior is that it may be driven by the increased inflammation seen in the blood and brain throughout the lifespan of these mice. Increased microglia in the brain in response to this chronic state could be one mechanism driving abnormal mTOR pathway activation in the brain and the abnormal behaviors observed in MIR offspring. Therefore, we used the CSFR1 inhibitor, Plexxikon5622, to deplete brain microglia in order to test this hypothesis. We found that ablation of microglia in young adult mice partially rescues repetitive, social, and sensory abnormalities (Fig 1F-H), suggesting that microglia do play a role in the pathophysiology of MIR exposure. However, microglia ablation does not significantly rescue these behaviors in old adult MIR mice even though rapamycin treatment does rescue them at this age (Fig 1F-H), suggesting that while microglia may play a role in the development and evolution of behavioral abnormalities, they are unlikely to be the primary mediator of ongoing behavioral alterations.

Exposure to mild maternal inflammation is the only pre-clinical model to date that produces a pattern of brain growth changes that have been reported to occur in 50-70% of humans with ASD, namely mild brain overgrowth at young ages followed by a decrease in growth at older ages[98]. This can be seen in the MIR mice in both gross brain wet weight measurements from birth to just over a year in age (Fig.2A) and in regional volumetric analysis of individual brain structures using MRI structural imaging (Fig. 2B-C). Adult MIR mice show significant increases in the volumetric size of the lateral ventricles, the hippocampus, caudate nucleus, and the amygdala but decreases in the thalamus and in sensory, motor, and association cortical areas (Fig. 2B-D). These regional

changes are consistent with human volumetric studies which have identified similar changes in these structures in different adult autistic populations[187–193].

Acute Rapamycin Treatment Effects on ASD-related Behaviors

Because of the persistently elevated mTOR pathway activation in MIR offspring, we examined the effects of the mTORC1 inhibitor, rapamycin, on animal physiology and behavior. While most studies of the effects of rapamycin on mouse behavior have used repeated or chronic administration of the drug, we have focused on its acute effects in order to more fully understand the mechanisms of its effects, which don't necessarily involve any significant structural changes to the brain or physical synaptic connectivity. Therefore, we examined the effect of an acute 2-hour treatment with rapamycin on multiple ASD-associated behaviors. See Figure 3A for the experimental treatment and behavioral testing paradigm. First we used phospho-S6 western blot from tissue microdissections in the cortex, striatum, and amygdala to demonstrate that a 2-hour treatment time frame allows for the rapamycin to cross the blood brain barrier and significantly inhibit the mTOR pathway in the brain (Fig 3B). We then tested the rapamycin and vehicle-treated MIR and control offspring for social approach (Fig 3C) and repetitive behaviors (grooming and circling in an open field) in both young and old adult mice (Fig. 3D). We have previously demonstrated that MIR offspring have significantly reduced social approach and increased repetitive behaviors[118]. Here we found high levels of repetitive behaviors in MIR offspring that become more severe in old MIR offspring compared to young offspring (Fig 3D) and that acute rapamycin treatment significantly reduces this behavior and social approach deficits in the 3-chamber test at both young and old ages (Fig 3C-D). A single dose of rapamycin has a temporary rescue effect on repetitive behaviors with significant behavioral abnormalities returning after 72 hours in MIR offspring (Fig 3E). However, repeated daily treatment with rapamycin leads to an apparent tolerance effect over 5 weeks leading to a loss of effectiveness on repetitive behaviors (Fig 3F).

In addition to social and repetitive behaviors we also determined that MIR offspring display behaviors consistent with sensory over-responsivity (SOR), a common ASD phenotype[194–197] that has also been demonstrated in genetic mouse models of Fragile X and Rett Syndrome[152,198]. We showed that compared to control offspring, MIR offspring have a significantly increased aversion to rough tactile textures on the paws in a novel light/dark avoidance test (Fig 4A-B), increased sensitivity to tactile touch/pressure in the Von Frey fiber sensitivity test (Fig 4C), and abnormal sensory gating and lack of habituation in the pre-pulse inhibition startle test using tactile (air puff) pre-pulse and auditory and tactile startle (Fig 4D-F). These tests demonstrate significant sensory processing abnormalities in MIR offspring. These abnormalities were all at least partially rescued by acute rapamycin treatment in both young and old adult mice (Fig 4B-F).

Central Nervous System Actions of Rapamycin

There is good evidence that sensory abnormalities in ASD can be mediated specifically by peripheral neuron abnormalities[152,198]. Therefore, given the strong rescue effects that we observed on sensory over-responsivity in addition to other ASD-related behaviors, in the MIR mice we wanted to assess whether the drug was acting on the peripheral or central nervous system. To do this we tested the effects of RapaBlock, a small molecule that does not cross the blood brain barrier and prevents the effects of rapamycin on the peripheral nervous system[183,184] on behavior in MIR and control animals. We found that there was no significant change in rapamycin rescue of repetitive behaviors, social approach, or sensory aversion with simultaneous RapaBlock treatment (Fig 5A-C), suggesting that the main effect of rapamycin is on the central nervous system in MIR mice.

mTOR Pathway Specificity of Effects

To confirm that rapamycin is having its rescue effects on MIR mice via its normal signaling to downstream mTOR targets like S6 Kinase and to rule out significant off-target effects, we administered the S6K inhibitor, PF4708671, according to the methods of Huang et al.[185] and evaluated the effect on repetitive behaviors. We

found that it does partially rescue repetitive behaviors after an acute 6-hour treatment (Fig 5D; p<0.05).

However, a comparison of the level of rescue of repetitive behaviors between rapamycin-treated, S6 kinase

inhibition, and microglia ablation in young adult MIR offspring indicates that rapamycin is significantly more

effective than the other treatments (Fig 5E).

Neuronal Hyper-excitability & Seizure Susceptibility

In order to understand potential physiological mechanisms underlying the behavioral changes in MIR mice and

the effects of rapamycin we examined the electrophysiological properties of neurons in brain slice preparations.

We found that cortical pyramidal neurons (CPNs) in layer 2/3 of the somatosensory cortex of adult MIR offspring

displayed a depolarization shift, resulting in a resting membrane potential that is closer to the action potential

threshold and decreased Rheobase (Table 1) and significantly increased frequency, amplitude, and area in

spontaneous excitatory postsynaptic potentials (EPSPs, P=0.038, Fig 6A) and voltage-evoked responses

(Supplemental Fig S1B-C), which indicate a marked increase in neuronal hyper-excitability compared to controls.

Under current-clamp conditions (depolarizing current injection of pulses 500-1000 ms pulses) the cortical

neurons from MIR brains produced 4 different categories of action potential activity (see Otsuka and Kawaguchi,

2008): regular spiking-slowly adapting (RS-SA), regular spiking-fast adapting (RS-FA), regular spiking-doublet (RS-

Doublet--when increasing current intensity evokes two action potentials at the beginning of the pulse), and

intrinsic bursting (IB). In layers 2/3, CPNs generally belong to the RS-SA category which was the case for the

neurons from control mice which had 75% RS-SA, 15% RS-FA, 10% RS-Doublet, and no intrinsic bursting (Fig 6B).

CPNs from MIR brains had a dramatically altered spiking profile with mainly RS-Doublets (45%), only 40% RS-SA,

3% RS-FA, and 12% intrinsic bursting patterns which are not normally observed in upper layer neurons and was

not seen in CPNs from control mice, indicating significantly altered firing patterns (Fig 6C, P=0.0014). The effects

of rapamycin on the hyper-excitable phenotype of MIR neurons was assessed via *in vitro* treatment where the

rapamycin is added to the slice media (2uM) and via *in vivo* treatment (5 mg/kg) where the mice were injected

with rapamycin I.P., 2 hours prior to euthanasia and tissue preparation. In vitro rapamycin significantly rescued the resting membrane properties of the CNPs from MIR offspring, normalizing the resting membrane potential and increasing Rheobase (Table 1). The increased frequency, amplitude, and area of spontaneous EPSCs in cortical neurons was significantly rescued following the *in vitro* rapamycin treatment (Fig 6C; P<0.5) and the frequency of EPSCs was similarly significantly reduced following *in vivo* rapamycin treatment (Fig 6D, P<0.5). Spontaneous EPSCs in striatal medium spiny neurons (MSNs) showed a trend of increasing frequency that didn't reach significance (Supplemental Fig S2A) and was reduced by acute *in vitro* rapamycin treatment (Supplemental Fig S2B).

Because the striatum receives its main excitatory input from the cerebral cortex and, to a lesser extent, from the thalamus we wanted to examine if the cortical hyper-excitability observed in MIR mice was affecting the striatal MSNs. To do this we used Bicuculline (10uM) to block GABA$_A$ receptors *ex vivo* in the slice media. We observed that the CPNs display paroxysmal (epileptic-like) discharges that propagate to the striatum and are manifested by large synaptic currents and bursts of spontaneous synaptic currents in MSNs. This indicates unmasked cortical hyper-excitability because control brains rarely displayed this type of MSN activity (Supplemental Fig S2C). Acute treatment with rapamycin *ex vivo* also reduced Bicuculline-induced hyper-excitability in MSNs in MIR brains (Supplemental Fig S2D).

Given this neuronal hyper-excitability observed in MIR offspring we also tested the effect of a dose escalation paradigm using the seizure-inducing drug, pentylenetetrazol (PTZ) to determine if MIR mice have brains that are more susceptible to seizure induction. The number of animals per group at each progressive seizure score following PTZ treatment was used to indicate relative seizure susceptibility between MIR and Control offspring. We found that MIR mice progressed up the seizure score ranking at lower doses of PTZ than control offspring with all MIR mice (8/8) experiencing some level of seizure activity at 40 mg/kg compared to only 2/8 control

mice (Table 2). Rapamycin treatment reduced this seizure susceptibility in MIR mice, lowering the seizure score

of mice but not eliminating the seizure-inducing effect of PTZ on the mice (Table 2). Thus adult MIR mice display

abnormal neuronal hyper-excitability in cortical and striatal neurons that is normalized by acute rapamycin

treatment, suggesting that one mechanism by which rapamycin is acting acutely on the brain is via the rapid

normalization of inhibitory/excitatory balance in the brain.

Circuit-level functional dysregulation of MIR brains is rescued by acute rapamycin treatment

Another potential mechanism by which acute rapamycin treatment could rapidly affect brain function and

behavior is through effects at the level of functional connectivity and network organization. We assessed young

adult MIR and control offspring before and 2 hours after rapamycin treatment using resting state functional

Magnetic Resonance Imaging (rsfMRI). Functional connectivity (r values transformed to Fisher z scores) across

the brain were calculated as an indication of the connectivity strength between cortical and subcortical regions

before and after acute Rapamycin treatment (Fig 7A). This comparison demonstrates that at baseline prior to

rapamycin, MIR mice show a general pattern of increased FC brain wide compared to controls, especially within

cortical and subcortical regions (white rectangles Fig. 7A) and within hemispheres between cortical-subcortical

regions (Fig. 7A, yellow rectangles). The effect of acute rapamycin treatment dramatically increases FC within

subcortical regions but decreases FC within cortical regions and between cortical and subcortical regions within

each hemisphere. The effects of rapamycin on control brains is small by comparison (Fig. 7A). Thus, the changes

on brain FC in MIR offspring from acute rapamycin treatment are large but do not produce a return to control FC

conditions.

Network based statistics (NBS) methodology (P<0.01 NBS, 2-tailed, Zalesky et al., 2010) was used to

quantitatively examine these FC differences at a detailed, subnetwork level (Fig. 7B,C). This showed that the

altered FC in MIR mice at baseline was primarily due to increases between cortical primary and higher-order

(secondary) sensory processing areas (primary somatosensory and somatosensory for hind limbs, forelimbs, whiskers, trunk, and neck/shoulder, visual, auditory, taste, and olfactory) and subcortical structures (thalamus and basal ganglia). However, there were notable decreases in FC between sensory cortical areas and amygdala (Fig. 7B). Despite the strong repetitive behavior phenotype in the MIR mice, there were surprisingly more limited motor-related FC differences identified in MIR offspring compared to controls. The primary motor cortex only showed increased FC to retrosplenial cortex (sensory integration), subicular complex (memory), and the striatum (motor). The largest proportion of the FC increases in MIR mice relative to controls was seen between subcortical structures, primarily basal ganglia connections with thalamus and sensory cortex (Fig. 7B). FC decreases between MIR and control mice were sparser, and were again primarily sensory cortex changes, but to the thalamus and amygdala (Fig. 7B).

The largest effects of acute rapamycin on FC in MIR mice was a pattern of within subcortical increases in FC (Fig. 7C) and the opposite effect on the cortex with significant FC decreases (Fig. 7B,C). Acute rapamycin treatment had only a limited effect on control mice (Supplemental Fig.S3A). The significant decreases in FC in MIR mice from rapamycin were between somatosensory cortical areas (primary, forelimb, hind limb, trunk, barrel) and basal ganglia and thalamic structures (Fig 7C). The increases in FC in MIR mice from rapamycin were between left and right hemisphere primary somatosensory cortex, from insular cortex (sensory salience), piriform cortex (olfaction), and endo-piriform nucleus (taste) to thalamic structures and between amygdala (limbic emotion) and thalamic and basal ganglia structures (Fig. 7C).

A functional modularity analysis was used to identify the community network architecture (groups of brain regions that are more strongly connected to each other than to other regions) for each experimental group where the number of modules identified indicate the functional specialization or segregation within that group's brain network. MIR brains were found to have four main modules: cortical, (red), two subcortical/cortical

(yellow, green) and a thalamic and basal ganglia module (blue) and a modularity index of 0.693 (Fig 7D(ii)). Control mice were found to only have three modules: anterior cortical (green), posterior cortical (yellow), and a mainly subcortical module (red) and a significantly lower modularity index of 0.511 (Fig 7D(i), p<0.01). The brain regions (nodes) that make up these modules were further quantitatively evaluated using classification consistency and diversity (CC, CD) parameters which respectively reflect: the degree to which each node was classified to the same versus different modules across all mice within a group (7D(i,ii,iii,iv)). With the exception of the cortex in control mice, MIR mice were characterized by all modules having high CD values relative to the controls, indicating increased inter-module connectivity in MIR mice that likely facilitates greater integration between the modules than in controls. (7D(ii)).

Acute rapamycin treatment had a dramatic effect on this community structure in the brain by normalizing the number of modules in MIR mice to 3 main modules that became broadly, spatially similar to controls (Fig 7D(iii)). This was marked by a whole brain reduction in modularity from 0.693 at baseline to 0.552 following acute rapamycin intervention (P<.05), which was approaching control baseline modularity of 0.511, indicating reduced segregation in the network. Within modules, rapamycin shifted the cortical nodes in the MIR mice to a significantly higher CD value that was very similar to the cortex in control mice at baseline (Fig 7D(ii,iii)). The two subcortical network nodes in MIR mice treated with rapamycin were shifted in the opposite direction to lower CD values and closer to controls values. In control mice, while rapamycin had no significant effect on modularity (0.511 to 0.533 in baseline and rapamycin treated conditions respectively), it had the opposite effect on nodes within modules compared to in MIR mice. It significantly reduced the cortical module CD values compared to baseline, but significantly increased subcortical CD values, indicating much greater flexibility (Fig 7D(i,iv).

A functional connectivity seed analysis from the posterior thalamus (multi-sensory processing, attention, motor control) to whole brain was performed (P<.01, cluster corrected) on the data from MIR and Control mice

before and after acute rapamycin treatment. MIR mice at baseline were found to have significantly increased

cortico-thalamic FC for primary motor cortex, piriform cortex (olfactory), and primary somatosensory cortex (S1,

S1DZ, hind limb, forelimb, barrel, shoulder/neck) compared to controls (Fig 7E). There were also increases in FC

from posterior thalamus to external globus pallidus (multi-sensory processing and motor control), dorsal

striatum (multi-sensory processing and motor control), and contralateral anterior thalamus (somatosensory and

proprioception sensory processing) compared to controls (Fig 7E). However, following acute rapamycin

treatment these MIR FC increases were reversed for cortico-thalamic FC between primary motor cortex and

primary somatosensory cortex (hind limb, forelimb), and from posterior thalamus to left striatum and to

contralateral anterior thalamus (Fig 7E). Finally, rapamycin further increased cortico-thalamic FC in primary

somatosensory cortex (S1, and barrel cortex) and piriform cortex and between posterior thalamus and dorsal

striatum and external globus pallidus compared to MIR mice at baseline (Fig 7E). Rapamycin treatment

bilaterally increased FC for several cortico-thalamic connections that weren't differentially connected in MIR

mice pre-treatment (insular cortex (salience network), and entorhinal cortex (memory, sensory salience, spatial

navigation) and between posterior thalamus and amygdala, bilateral dorsal striatum and external globus pallidus

(Fig 7E). Rapamycin treatment also bilaterally decreased posterior thalamus FC to the anterior cingulate cortex

(salience network), retrosplenial cortex (visual/auditory sensory processing and associations), primary auditory

cortex, primary and secondary visual cortex, temporal association cortex (memory, visual/auditory processing),

ventral striatum (sensory integration with emotion/reward processing), and hippocampus (memory) which also

weren't differentially connected to the posterior thalamus in MIR mice prior to rapamycin treatment (Fig 7E).

Therefore, rapamycin treatment's effect on brain functional connectivity was largely found to be a reduction to

thalamic-mediated sensory processing regions. Voxelwise rsfMRI mapping of the thalamic seed FC data showed

a pattern of increased FC strength and hyper-connectivity in MIR mice compared to controls. Global histogram

analysis of this data using the difference between MIR and control mice (Fig. 7E) confirmed a large shift in FC

strength for both cortical and subcortical structures in MIR mice and a divergent effect of rapamycin treatment,

both decreasing cortical hyper connectivity with the posterior thalamus and increasing connectivity for subcortical structures (Fig 7F).

In summary, our analysis of resting state functional connectivity and circuitry demonstrates that MIR mice have increased FC in widespread areas of the brain and this is largely normalized towards controls following an acute 2-hour treatment with Rapamycin, at the expense of dramatically increased subcortical FC (Fig. 7A). Specific brain nodes that have increased functional synchrony in MIR mice compared to controls include retrosplenial cortex, auditory and somatosensory cortex (sensory) and cingulate cortex and amygdala (limbic), and insular cortex (sensory salience) areas (Fig. 7B) which is congruent with the sensory, anxiety, and social behavioral abnormalities seen in the MIR mice that are rescued by the acute rapamycin treatment[118] (Fig. 3C-D). Significant decreases in FC in MIR mice compared to controls were seen between forelimb (motor), medial amygdala, striatum, globus pallidus, and periaqueductal gray areas which have been implicated in repetitive behaviors, and is also rescued by rapamycin (Fig. 7C). There were very few significant changes in FC in control mice following rapamycin treatment (Supplemental Fig S3A) which aligns with our observation that there were no significant behavioral effects of rapamycin on control mice (Fig 3C-E, 4B-C).

We validated several of these imaging findings with immediate early gene (IE) immunostaining for Egr1+ neurons in the brains of mice euthanized 30 minutes after behaving freely in an open field environment where MIR mice displayed characteristic repetitive behaviors and control mice performed normal exploratory behaviors. Using Egr1 expression as a correlate of recent neuronal activity we found that the motor cortex and striatum of MIR mice had increased egr1 positive neurons, which also confirms our electrophysiology findings of neuron hyper-excitability in those brain regions (Supplemental Fig S3B). We also found increased egr1+ neurons in MIR brains compared to controls in the retrosplenial cortex and decreased egr1+ neurons in the medial amygdala (Supplemental Fig S3B). Thus, another mechanism by which rapamycin is acutely affecting brain

function is by the rapid alterations in functional activation and connectivity and normalization of network flexibility that is affected by mTOR inhibition.

Single Nucleus and Bulk mRNA Sequencing of Genetic Alterations in MIR Mice and After Rapamycin Treatment

In order to identify further potential mechanisms underlying MIR-induced behavioral changes and their mitigation with rapamycin, we performed single nucleus RNA sequencing (snRNA seq) of adult MIR and control mice and bulk RNA sequencing of MIR and control animals. SnRNA sequencing of tissue from the somatosensory cortex of adult MIR and control offspring did not indicate significant changes in the numbers of cells found in the different cell populations in this cortical brain region. Volcano plots visualize the most significantly differentially expressed neuronal (Fig 8C) genes in MIR mice relative to controls. Although there weren't significant changes in the cellular makeup of the cortex, the number of differentially expressed genes (DEGs) within each cell type analyzed between MIR and Control offspring shows that there are many significant cell intrinsic changes with the greatest number of DEGs found in oligodendrocyte, microglia, upper layer and deep layer excitatory neuron populations, and other excitatory neurons (Fig 8D, p<0.05). Overall, there are more significant DEGs in the excitatory neurons compared to inhibitory neurons in MIR mice (Fig 8D). mTOR pathway-related genes are significantly enriched in MIR mice in oligodendrocytes and their precursor cells, microglia, PV interneurons, excitatory upper layer neurons, other excitatory (neurogranin) neurons, endothelial cells, and astrocytes (Fig 8F; p<0.05). A number of autism-associated, epilepsy/seizure-associated, and neuro-inflammatory-associated genes identified by cell type were significantly differentially expressed in MIR compared to control offspring using pseudobulk analysis of the snRNA seq data (LogFC>1, p_val_adj <0.05, FDR<0.1, Table 3). Two of these genes, Kirrel and GRIP1, that were significantly down-regulated in MIR mice are also known to have similarly reduced function in people and other mouse models of sensory over-responsivity and repetitive behaviors (Gandhi and Lee, 2020; Hisaoka et al., 2018; Mejias et al., 2011, 2019). Gene set enrichment analysis (GSEA) of snRNA seq data by pathway for all cell types indicate significant increases in genes for the energy-dependent regulation of

mTOR and activity of kainate receptors, among others (Supplemental Fig S4A). GSEA of pathways by cell type shows (1) enriched gene expression involved in ion channel transport and homeostasis and decreases in TGF-beta and potassium channel signaling in all neurons in MIR mice (Supplemental Fig S4B), (2) enriched gene expression involved in NF-kappa beta, Notch, and Toll-like receptors 5 and 8 and decreased expression of cytosolic calcium level and AP-2 transcription factors in excitatory MIR neurons (Supplemental Fig S4C), and (3) enriched gene expression involved in GABA receptor activation and interleukin-1 signaling but decreased expression of pathways for voltage-gated potassium channels, NCAM-regulated neurite outgrowth, and gap junction regulation in inhibitory neurons from MIR mice (Supplemental Fig S4D). Thus, the lack of significant cellular population changes in MIR mice indicates that the neuronal hyperexcitability found in these mice is more likely due to alterations in cell intrinsic properties because there is no loss or over-abundance of a particular inhibitory or excitatory neuronal population. Therefore, the mechanism of acute rapamycin treatment in rescuing the abnormal phenotypes in MIR mice is likely to be targeting intrinsic neurotransmission or ion channel phenotypes in the neural cell types with activated mTOR signaling which includes PV interneurons and upper layer excitatory neurons which can be rapidly altered (Fig 8E).

Bulk RNA sequencing was performed on somatosensory cortical tissue from MIR and Control offspring with and without acute rapamycin treatment. Several DEGs that can rapidly affect neuronal function within the short time period of the acute rapamycin treatment were significantly altered in MIR mice including GABA and glutamate-related genes, transcription factors, and ion channel genes (LogFC>1, p<0.05, FDR<0.1, Table 4). Bulk sequencing data was also used to examine ASD-associated risk factor, high confidence genes using the gene sets from (Gandal et al., 2018, 2019; Basu et al., 2009). We found a small number of these ASD-associated genes were significantly up- or down-regulated in MIR mice, including TEK, PLAUR, ID1, Homer1, ASPM, TEKT4, CNTNAP3, Slc35D3, Grin2c, CHD7, CDKN1A (p21), and NeuroD1 (Table 5). Rapamycin rescued the over-expression of several ASD-associated genes, including TEK, Plaur, CDKN1A (p21), and ID1, and the under-

expression of TEKT4, CHD7, and CNTNAP3 in MIR mice (Table 6). Rapamycin treatment also altered expression of several genes that were differentially expressed in MIR mice relative to controls that aren't known to be specifically associated with autism, including decreasing Nestin, GEM, TIPARP, and APOLD1 expression and increasing HIF3A and PLIN4 expression (Table 6). GSEA analysis of pathways in these bulk sequencing experiments show significantly enriched expression of genes involved in glutamate receptor signaling, oxytocin signaling, PI3K pathway signaling, apoptosis, and angiogenesis pathways in MIR mice (Supplemental Fig S4E). Gene ontology for biological processes show that MIR brains are enriched for G-protein coupled receptor signaling and negative regulation of hydrolase activity among others but decreased expression of regulation of cytokines, amine metabolic processes, and autonomic nervous system development in MIR mice (Supplemental Fig S4F). Following rapamycin treatment, GPCR signaling, innate immune system, signal transduction, post-translational, and protein modification signaling are all increased (Supplemental Fig S4G). GSEA for pathways indicate significantly elevated gene sets involved in regulation of neurotransmitter levels, lipid regulation, and response to growth factors but decreases in those involved in epithelial and skeletal systems following acute rapamycin treatment (Supplemental Fig S4H).

   Thus, we found a number of differentially expressed genes in MIR offspring which control cellular mechanisms like excitatory & inhibitory neurotransmitter production, transcription factors, and ion channel functions which could contribute to the pathophysiological neuron excitability and aberrant functional connectivity that we have observed in MIR mice and which can be rapidly altered by an acute rapamycin treatment. There were also a number of autism-associated genes that were differentially regulated in MIR offspring that are consistent with the behavioral phenotypes of these mice. Although changes at the level of gene expression within a 2-hour time frame of our rapamycin treatment might not be expected to be large, we did see the rescue of several of the autism-associated genes that are differentially expressed in MIR mice, identifying a number of promising new therapeutic targets.

Discussion

Exposure to mild maternal inflammatory response (MIR) in early gestation in CD-1 mice recapitulates multiple ASD associated physiological and behavioral phenotypes. This includes mild brain overgrowth in juvenile ages followed by regionally variable volumetric brain changes in later adulthood like cortical thinning and larger hippocampal, basal ganglia, and ventricular structures. Similar growth trajectories have been reported in clinical ASD and neuro-inflammation studies[187–193,199]. MIR mice also demonstrate other common ASD pathology like chronically elevated blood cytokines and brain microglia which have also been reported in human ASD populations[19,83–97,200–205]. Furthermore, this model produces offspring with multiple ASD-related behavioral phenotypes like social, communication (vocalization), anxiety, and repetitive behaviors[118]. We have now further determined that MIR mice demonstrate significant sensory over-responsivity (SOR) and sensory processing dysfunction as well. SOR is an extremely impairing clinical condition marked by avoidance and/or sensitivity to sensations such as scratchy clothing, loud noises, or visually stimulating environments. SOR is present at high rates across the autism spectrum and is associated with other core features of ASD, as well as difficulties with behavioral and emotion regulation[206–209]. Despite its significance, very little is known about the causes or underlying biological mechanisms of SOR and there are essentially no evidence-based treatments for SOR. Several genetic murine models of mutations in autism risk factor genes like Mecp2 (Rett Syndrome) and FMR1 (Fragile X)[152,198] are known to display SOR behaviors but ours is the first report of SOR in an acquired model of ASD with a neuro-inflammatory origin.

Although it may be surprising that a single exposure to low-level maternal inflammation could cause such lasting effects on the brain and behavior, we and others have found that early *in utero* exposure sets up chronic

dysregulation in systemic immune function[210–215]. Our analysis of both young and older adult MIR offspring showed a persistent elevation in pro-inflammatory cytokines in the blood and increased brain-wide increases in microglia, suggesting a chronic inflammatory state across the life-span compared to age-matched controls. Both clinical and pre-clinical studies have previously reported some success in the clinical treatment of ASD in children with an anti-inflammatory drug, pioglitazone, which improved behavioral phenotypes along with reducing neuro-inflammation[216–224]. We also found a concomitant chronic elevation of mTOR signaling in the neurogenic adult subventricular niche[118] and now in several more regions in adult (cortex, amygdala, and striatum). Induction of systemic and brain inflammation in adult wildtype/control mice does not produce any of the ASD-associated abnormal behaviors that are produced in MIR mice where inflammation was induced *in utero* and persists chronically[225]. Therefore, an elevated inflammatory state in the adult brain does not, on its own, explain most of the MIR phenotypes, suggesting that there are likely important vulnerable periods in brain development where inflammatory exposure is especially deleterious and/or the effects of chronic inflammation and increased mTOR pathway activity have stronger effects. It is well known that chronic neuro-inflammation can contribute to significant brain pathophysiology[214]. Maternal inflammation is also known to be a significant risk factor for both ASD and brain overgrowth[19,85,101–117]. Therefore, one of our initial hypotheses was that the chronically increased brain microglia underlies some or all of the abnormal brain pathology and behaviors in the MIR model as microglia have been demonstrated to influence several related behaviors and mTOR signaling[90,226–232]. However, our experiments which ablated microglia in the adult brain suggest that the role of microglia in the etiology of MIR-induced abnormalities is complex. Firstly, microglia ablation with a CSF1 inhibitor can rescue some abnormal behaviors in young adult mice but not in older adult mice suggesting that they likely play a role in setting up physical and functional brain changes but are not the primary cause of behavioral dysfunction since acute treatment with the mTOR inhibitor, rapamycin, can significantly reduce abnormal MIR phenotypes in both young and old MIR offspring. Thus the effects of MIR exposure on behavior are complex and evolving over the lifespan as we can see in the physical changes to the MIR brain such as the

early overgrowth followed by later regional decrease in cortical brain volume. Chronic post-natal neuro-inflammation and mTOR activation in the brain has also been associated with epilepsy and a disturbed excitatory/inhibitory balance[199,233–235]. In keeping with a putative neuro-inflammatory etiology, our MIR mice also display significant cortical hyper-excitability and susceptibility to seizure induction.

Given the long-term elevation in mTOR signaling that we have seen throughout the brain in MIR offspring and the similarity of some ASD-associated phenotypes in genetic models of autism (TSC and PTEN mutants) that also have increased mTOR pathway activation, we examined the role of mTOR signaling in MIR pathology. It is known that the mTOR inhibitor, rapamycin, can rescue several abnormal ASD-associated behaviors in murine genetic models of PTEN and TSC mutations even in adult mice with existing/ongoing brain physiological abnormalities like brain overgrowth[9,19,124,125,131,132]. Therefore, we hypothesized a similar rescue effect was possible in our MIR model. However, we observed an unanticipated and potentially useful rapid effect of rapamycin on several abnormal behaviors in MIR offspring. We confirmed via microdissection of several brain regions and western blot analysis that the rapamycin treatment was able to access the CNS and significantly reduce mTOR signaling within the brain during this acute time course. Additionally, it has been reported that sensory abnormalities in Mecp2 and FMR1 mutant mice can be reversed through entirely peripheral neuron interventions[152,198]. Therefore, to test this in our maternal inflammation model we used a newly developed FKBP12 antagonist, RapaBlock, which prevents rapamycin effects in the peripheral nervous system to demonstrate a CNS-mediated effect on MIR mice. These insights allowed us to focus our search for the causes of MIR abnormalities and rapamycin effects in the brain on rapidly changeable physiological and functional mechanisms which would not be apparent from more chronic rapamycin treatment paradigms that have been shown to induce permanent physical transformations such as restoring normal dendritic and synaptic complexity[9,22,120–124]. In this way, studying the rapid effects of rapamycin may reveal novel biological and functional mechanisms underlying multiple ASD-associated abnormal phenotypes that can be overlooked with chronic treatment paradigms.

While synaptic strengthening/weakening through altered levels of neurotransmitter release or post-synaptic receptor expression can happen on short (minutes to hours) time scales[236–238], widespread physical synaptic remodeling is an unlikely mechanism for our acute rapamycin effects. This led us to examine changes in neuronal gene expression, excitation, and functional activation and organization at the circuit level as possible fast-acting mechanisms that could underlie the brain and behavioral dysfunction rescued by acute rapamycin treatment in the MIR model. Our single cell sequencing results did not find that MIR mice have significant neuronal or glial cell population size changes. However, there was significantly upregulated gene expression related to mTOR pathway activation in multiple cell types including microglia, astrocytes, oligodendrocytes and their progenitors, excitatory upper layer neurons, and inhibitory parvalbumin neurons in MIR mice. This suggests that cell-intrinsic changes are responsible for mTOR dysregulation in MIR offspring rather than alterations in the balance of specific cell population numbers. Additionally, we found that a number of autism, epilepsy, and neuro-inflammation associated genes alterations were differentially expressed in upper and deep layer excitatory neurons from MIR offspring that have been associated with repetitive behaviors, sensory processing dysfunction, synaptic plasticity, and epilepsy. Bulk sequencing with and without rapamycin treatment suggests several gene expression alterations occur in several rapidly changeable cellular properties such as transcription factors, ion channels, and neurotransmitter synthesis-related genes in MIR mice. In addition, several high confidence ASD risk genes[68,96,239] were altered in MIR offspring and a number of these were reversed by acute rapamycin treatment. Several of these ASD genes also have associations with alterations in brain size (ASPM, KIF14), epilepsy (PLAUR, ASPM, MAST3, HOMER1, ACE, ID1, CHD7, CDKN1A (p21)), intellectual disability (CHD7, TEKT4), sensory processing (PLAUR), synaptic plasticity (HOMER1, CNTNAP3, IQGAP3) and neuroinflammation (TEK/Tie2, CNTNAP3, ACE, CDKN1A[240–268]. Although the single cell sequencing results show that there were no significant changes in the relative balance of excitatory and inhibitory neurons in MIR mice, we did observe a significant hyper-excitability in cortical neurons in our electrophysiology experiments, which could be related to

the cell-intrinsic gene expression changes that we saw in excitatory and inhibitory neurotransmitter- and ion channel-related genes in MIR neurons. Rapid reduction of this neuronal hyper-excitability and seizure susceptibility with rapamycin treatment within the same acute time-frame as behavioral rescue suggests that restoring excitatory-inhibitory regulation in the brain may underlie the normalization of sensory dysfunction and ASD-related behaviors in the MIR model. In support of this, others have shown that an acute 2-hour pre-treatment with the rapalogue Everolimus also reduces seizure susceptibility and suppressed neuro-inflammatory responses in a kainic acid rat epilepsy model[269].

Functional connectivity (FC) is another level of brain regulation which can be rapidly altered by acute rapamycin treatment. The majority of the FC changes in MIR offspring compared to controls in our resting-state fMRI data were increases in FC and only a few brain regions showed decreases. We found that the largest proportion of FC differences between MIR and control offspring involved largely sensory rather than motor circuits despite the strong repetitive motor phenotype in MIR mice. While there was an increase in M1 motor cortex and basal ganglia FC in MIR brains the majority of the hyper-connectivity in cortico-striatal cortico-limbic functional connections were from sensory cortical areas. Additionally, the basal ganglia regions with significantly increased FC connections to motor cortex in MIR were the dorsolateral striatum and external GP which are known to have sensory processing roles in addition to motor functions. These basal ganglia regions receive projections from the left thalamus and sensorimotor cortex, and integrate sensorimotor information to learn and perform motor habits. The most significant FC changes in MIR mice relative to controls were found in the thalamus which prompted us to do a seed analysis from this region. Thalamic sensory FC was altered in multiple primary sensory cortical areas as well as sensory salience network structures (insula and anterior cingulate) in MIR brains. We interpret this dominant sensory phenotype in FC alterations to be in keeping with the behavioral SOR findings in the MIR model. However, it does not entirely fit with the strong repetitive behavior phenotype even though there is considerable comorbidity for sensory dysregulation with repetitive behaviors[270–274]. We

propose that recent clinical findings may offer an explanation for the lack of a stronger functional motor phenotype in our mice. A recent human study showed that reductions in repetitive behaviors in individuals are associated with an increase in other autism phenotypes like anxiety and social difficulties and vice versa, suggesting that some repetitive behaviors may be a self-regulation mechanism rather than a primary phenotype in autism[275]. Given the close correlations that have been found between repetitive behaviors and sensory sensitivity and the beneficial effects of repetitive behaviors[276], our finding that sensory dysregulation is the more dominant brain phenotype in rsfMRI of MIR mice with SOR, supports the idea of an emergence of repetitive behaviors as coping mechanism rather than being a cause of dysfunction. The acute rapamycin treatment in MIR mice reduced FC towards control levels in the cortex but curiously caused even greater FC differences in subcortical structures. This shows that the rescue effects of the treatment are not entirely via a restoration back to control levels and patterns of functional connectivity across the brain. This is likely related to the fact that brain structural abnormalities remain, such as the increases in hippocampus, caudate, and amygdala size and decreased thalamus and cortical volumes and rapamycin is acting in the context of these physical changes. Longer treatments with rapamycin like those in Pagani et al.[22] in TSC mutant mice show that rescuing the physical synaptic abnormalities produces a greater return to control FC patterns. These circuit-level insights into mechanisms of dysregulation and the pattern of changes that are associated with behavioral recovery have identified potentially novel neuromodulatory targets for non-pharmaceutical based treatments like TMS and focused ultrasound.

Analysis of network community architecture is a data-driven approach to evaluate brain wide synchrony and network properties rather than using a strictly anatomical analysis of functional connectivity that relies on predefined brain regions. These circuit-level properties can be rapidly modulated and significantly affect brain function. Graphical analysis of this functional circuitry allows the unbiased detection of communities of brain regions with strong functional connectivity, which may represent specialized functional units or systems in the

brain. Graph theory can also address cognitive flexibility and information integration by examining how hub nodes (highly connected nodes within a module that maintain network integrity and facilitate information flow between brain regions) are structured, the strength of this connectivity, and the size (specialization) of modules. For example, classification diversity (H-index), is a measure of module diversity that indicates how evenly distributed a node's connections are across different modules relative to the number of connections within its own module. This measure is helpful in identifying nodes that play a crucial role in integrating information across different functional modules in the brain (like multisensory integration), which is an essential aspect of cognitive flexibility. This network analysis of resting-state fMRI of MIR mice revealed that the offspring have a larger number of modules compared to control offspring, suggesting that their brain network is more segregated into distinct functional sub-networks within the subcortical regions, representing an increased specialization of function. However, the nodes within these subcortical modules also have a significantly shifted higher classification diversity values than in controls, indicating nodes more frequently transition membership between modules within the MIR group, reflecting potentially increased information transfer between modules, and greater functional integration within the brain. However, the opposite is true within the cortex of MIR mice, with a lower classification index indicating reduced information transfer to other network modules of the brain. The combination of increased module number and both higher and lower intra-module nodal diversity suggests a complex reorganization of brain networks which may reflect both deficits and compensatory adaptive changes in functional connectivity. We are unaware of other resting state fMRI studies in preclinical models like ours that have examined this type of global functional reorganization at the network level. However, there have been clinical studies that have undertaken this type of network analysis that have similar findings of increased numbers of functional modules, increased connections between modules, and increased module diversity in individuals with ASD with resting-state fMRI[277–282]. Our MIR offspring share this aberrant network architecture with humans, demonstrating another cross-species phenotype that is present in our inflammatory, acquired model of ASD. It has been suggested that the functional modular reorganization in the ASD brain may be a

compensatory mechanism for restoring functional homeostasis at the circuit level in response to physical

connectivity changes, and that therapeutic normalization of this altered modular organization would therefore

be likely to worsen ASD dysfunction[278]. However, our acute rapamycin treatment data suggest otherwise. The

rapid rescue of this abnormal network modularity in MIR mice following rapamycin treatment was one of the

largest effects of the pharmacological treatment on the MIR model of brain dysfunction. Both the number of

modules and the distribution of the nodal connectivity within these modules was returned to control levels after

acute rapamycin, suggesting that at a macro circuit level, the treatment restores functional information flow and

network stability within the same time frame that we observe behavioral and electrophysiological rescue from

this treatment. Therefore, our findings strongly suggest that modular organization of functional brain networks

are related to behavioral traits in ASD and that this may be a central functional mechanism underlying

inflammation-mediated behavioral dysfunction in several core and comorbid autism phenotypes. There is

evidence in the literature that supports the theory that these functional brain alterations that we have observed

in the MIR model are induced by maternal inflammation and sustained by the chronic inflammation in the

offspring. For example, neuro-inflammation has been increasingly recognized as a factor contributing to atypical

functional connectivity patterns and network abnormalities[283–289]. Neuro-inflammation can affect synaptic

function by modulating the release of neurotransmitters, altering receptor expression, and ion channel function

which impacts long-term potentiation and long-term depression[72,86,290–295]. These changes can disrupt the

formation and maintenance of functional connections leading to functional network abnormalities. For example,

enhanced long-term depression in humans and animal ASD models have been shown to underlie hyper-reactive

sensory processing[296]. Neuro-inflammation during critical periods of brain development can also interfere with

processes like axonal growth and synaptogenesis[87,219,297–301] which can contribute to the increased modularity

and module diversity seen in ASD and in our MIR model as the brain adapts to these physical changes with

functional compensation towards the goal of homeostasis. Disruption to normal glutamate and GABAergic

neurotransmission and receptor expression can also cause imbalances in neuronal excitation thresholds that

further impacts functional network[199,235,302–307] which is consistent with the phenotypes that we have observed in MIR offspring.

   In summary, our experiments demonstrate a plausible etiology for MIR-induced brain and behavioral dysfunction that is revealed through the acute effects of rapamycin: (1) exposure to MIR *in utero* causes neuroinflammation and chronic post-natal neuroinflammation in the offspring that, (2) elevates mTOR signaling chronically, (3) increases neurogenesis and synaptogenesis at a critical period of early brain development leading to, (4) brain overgrowth, abnormal physical connectivity, and (5) a compensatory functional reorganization of the brain connectome. Our data also suggest that exposure to maternal inflammation could account for the non-macrocephalic brain overgrowth in ASD that has been identified in early childhood[98–100]and that even mild alterations in brain growth can produce aberrant functional connectivity and cellular dysfunction that leads to brain-wide functional network changes and behavioral abnormalities. Although we have shown that aberrant brain and behavioral phenotypes can be understood as dysfunction at the level of large-scale networks and their information integration, we have also established in this MIR model and with acute rapamycin treatment that this global activity is dependent on processes happening at the level of physical systems like the immune system, of cells and their excitability, and of sub-cellular processes like pathway activation and gene expression regulation. Therefore, although longer interventions in suppressing excessive mTOR signaling have been shown to also impact the physical features of maladaptive plasticity[22], we have identified a number of potential therapeutic targets at various levels of dysfunction. We demonstrate that the MIR brain is amenable to rapid normalization of functional organization and neuronal excitability with short rapamycin treatments which results in rescue of core and comorbid ASD behaviors in adult animals without requiring long-term physical alterations to the brain. Thus, restoring excitatory/inhibitory imbalance and sensory functional network modularity may be key mTOR-related targets for therapeutically addressing both primary sensory and social behavior phenotypes and compensatory repetitive behavior phenotypes.

Acknowledgements:

Dr. Miriam and Sheldon G. Adelson Medical Research Foundation
UCLA Intellectual and Developmental Disabilities Research Center NIH grant HD103557
Center for Autism Research and Treatment (CART) Pilot Grant Award, which is supported by NIH/NICHD grant P50-HD-055784
UCLA Brain Injury Research Center
Simons Foundation Cross-Species Studies of ASD Grant
UCLA Clinical and Translational Science Institute (CTSI) Accelerator Core Voucher Program supported by grant number UL1TR001881
Autism Speaks Basic and Clinical grant
Autism Speaks Environmental Sciences grant
Gene sequencing was done by the UCLA Technology Center for Genomics and Bioinformatics, Dr. Xinmin Li Director
Z.Z. is a Damon Runyon Fellow supported by the Damon Runyon Cancer Research Foundation (DRG-2281-17)
Electrophysiology experiments were supported by the Cell, Circuits and Systems Analysis Core (NIH P50HD103557)

1. Tanaka, M., Spekker, E., Szabó, Á., Polyák, H. & Vécsei, L. Modelling the neurodevelopmental pathogenesis in neuropsychiatric disorders. Bioactive kynurenines and their analogues as neuroprotective agents-in celebration of 80th birthday of Professor Peter Riederer. *J Neural Transm (Vienna)* **129**, 627–642 (2022).

2. Li, Y.-J., Zhang, X. & Li, Y.-M. Antineuroinflammatory therapy: potential treatment for autism spectrum disorder by inhibiting glial activation and restoring synaptic function. *CNS Spectr* **25**, 493–501 (2020).

3. Chen, J., Alberts, I. & Li, X. Dysregulation of the IGF-I/PI3K/AKT/mTOR signaling pathway in autism spectrum disorders. *Int J Dev Neurosci* **35**, 35–41 (2014).

4. Thomas, S. D., Jha, N. K., Ojha, S. & Sadek, B. mTOR Signaling Disruption and Its Association with the Development of Autism Spectrum Disorder. *Molecules* **28**, 1889 (2023).

5. Arenella, M. *et al.* Immunogenetics of autism spectrum disorder: A systematic literature review. *Brain Behav Immun* **114**, 488–499 (2023).

6. Lugo, J. N. *et al.* Deletion of PTEN produces autism-like behavioral deficits and alterations in synaptic proteins. *Front Mol Neurosci* **7**, 27 (2014).

7. Kwon, C.-H. *et al.* Pten Regulates Neuronal Arborization and Social Interaction in Mice. *Neuron* **50**, 377–388 (2006).

8. Rademacher, S. & Eickholt, B. J. PTEN in Autism and Neurodevelopmental Disorders. *Cold Spring Harb Perspect Med* **9**, a036780 (2019).

9. Zhou, J. *et al.* Pharmacological inhibition of mTORC1 suppresses anatomical, cellular, and behavioral abnormalities in neural-specific Pten knock-out mice. *J Neurosci* **29**, 1773–1783 (2009).

10. Ljungberg, M. C., Sunnen, C. N., Lugo, J. N., Anderson, A. E. & D'Arcangelo, G. Rapamycin suppresses seizures and neuronal hypertrophy in a mouse model of cortical dysplasia. *Dis Model Mech* **2**, 389–398 (2009).

11. Nguyen, L. H. *et al.* mTOR inhibition suppresses established epilepsy in a mouse model of cortical dysplasia. *Epilepsia* **56**, 636–646 (2015).

12. Sunnen, C. N. *et al.* Inhibition of the mammalian target of rapamycin blocks epilepsy progression in NS-Pten conditional knockout mice. *Epilepsia* **52**, 2065–2075 (2011).

13. Amaral, D. G., Schumann, C. M. & Nordahl, C. W. Neuroanatomy of autism. *Trends Neurosci* **31**, 137–145 (2008).

14. Lee, J. K. *et al.* Longitudinal Evaluation of Cerebral Growth Across Childhood in Boys and Girls With Autism Spectrum Disorder. *Biol Psychiatry* **90**, 286–294 (2021).

15. Libero, L. E. *et al.* Persistence of megalencephaly in a subgroup of young boys with autism spectrum disorder. *Autism Res* **9**, 1169–1182 (2016).

16. van Rooij, D. *et al.* Cortical and Subcortical Brain Morphometry Differences Between Patients With Autism Spectrum Disorder and Healthy Individuals Across the Lifespan: Results From the ENIGMA ASD Working Group. *Am J Psychiatry* **175**, 359–369 (2018).

17. Allen, M. *et al.* Astrocytes derived from ASD individuals alter behavior and destabilize neuronal activity through aberrant Ca2+ signaling. *Mol Psychiatry* **27**, 2470–2484 (2022).

18. Vakilzadeh, G. & Martinez-Cerdeño, V. Pathology and Astrocytes in Autism. *Neuropsychiatr Dis Treat* **19**, 841–850 (2023).

19. López-Aranda, M. F. *et al.* Postnatal immune activation causes social deficits in a mouse model of tuberous sclerosis: Role of microglia and clinical implications. *Sci Adv* **7**, eabf2073 (2021).

20. Specchio, N. *et al.* Autism and Epilepsy in Patients With Tuberous Sclerosis Complex. *Front Neurol* **11**, 639 (2020).

21. Cummings, K., Watkins, A., Jones, C., Dias, R. & Welham, A. Behavioural and psychological features of PTEN mutations: a systematic review of the literature and meta-analysis of the prevalence of autism spectrum disorder characteristics. *J Neurodev Disord* **14**, 1 (2022).

22. Pagani, M. *et al.* mTOR-related synaptic pathology causes autism spectrum disorder-associated functional hyperconnectivity. *Nat Commun* **12**, 6084 (2021).

23. Zahedi Abghari, F., Moradi, Y. & Akouchekian, M. PTEN gene mutations in patients with macrocephaly and classic autism: A systematic review. *Med J Islam Repub Iran* **33**, 10 (2019).

24. Kim, W.-Y. Brain size is controlled by the mammalian target of rapamycin (mTOR) in mice. *Commun Integr Biol* **8**, e994377 (2015).

25. LiCausi, F. & Hartman, N. W. Role of mTOR Complexes in Neurogenesis. *Int J Mol Sci* **19**, 1544 (2018).

26. Winden, K. D., Ebrahimi-Fakhari, D. & Sahin, M. Abnormal mTOR Activation in Autism. *Annu Rev Neurosci* **41**, 1–23 (2018).

27. Zoghbi, H. Y. & Bear, M. F. Synaptic dysfunction in neurodevelopmental disorders associated with autism and intellectual disabilities. *Cold Spring Harb Perspect Biol* **4**, a009886 (2012).

28. Onore, C., Yang, H., Van de Water, J. & Ashwood, P. Dynamic Akt/mTOR Signaling in Children with Autism Spectrum Disorder. *Front Pediatr* **5**, 43 (2017).

29. Costa-Mattioli, M. & Monteggia, L. M. mTOR complexes in neurodevelopmental and neuropsychiatric disorders. *Nat Neurosci* **16**, 1537–1543 (2013).

30. Lipton, J. O. & Sahin, M. The neurology of mTOR. *Neuron* **84**, 275–291 (2014).

31. Sato, A. & Ikeda, K. Genetic and Environmental Contributions to Autism Spectrum Disorder Through Mechanistic Target of Rapamycin. *Biol Psychiatry Glob Open Sci* **2**, 95–105 (2022).

32. Groszer, M. *et al.* PTEN negatively regulates neural stem cell self-renewal by modulating G0-G1 cell cycle entry. *Proc Natl Acad Sci U S A* **103**, 111–116 (2006).

33. Gregorian, C. *et al.* Pten Deletion in Adult Neural Stem/Progenitor Cells Enhances Constitutive Neurogenesis. *J Neurosci* **29**, 1874–1886 (2009).

34. Hui, K. K. & Tanaka, M. Autophagy links MTOR and GABA signaling in the brain. *Autophagy* **15**, 1848–1849 (2019).

35. Antoine, M. W., Langberg, T., Schnepel, P. & Feldman, D. E. Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models. *Neuron* **101**, 648-661.e4 (2019).

36. Chaudry, S. & Vasudevan, N. mTOR-Dependent Spine Dynamics in Autism. *Front Mol Neurosci* **15**, 877609 (2022).

37. Kassai, H. *et al.* Selective activation of mTORC1 signaling recapitulates microcephaly, tuberous sclerosis, and neurodegenerative diseases. *Cell Rep* **7**, 1626–1639 (2014).

38. Sadowski, K., Kotulska-Jóźwiak, K. & Jóźwiak, S. Role of mTOR inhibitors in epilepsy treatment. *Pharmacological Reports* **67**, 636–646 (2015).

39. Marotta, R. *et al.* The Neurochemistry of Autism. *Brain Sci* **10**, 163 (2020).

40. Luo, C. *et al.* Perfect match: mTOR inhibitors and tuberous sclerosis complex. *Orphanet J Rare Dis* **17**, 106 (2022).

41. Laplante, M. & Sabatini, D. M. mTOR signaling in growth control and disease. *Cell* **149**, 274–293 (2012).

42. Angliker, N., Burri, M., Zaichuk, M., Fritschy, J.-M. & Rüegg, M. A. mTORC1 and mTORC2 have largely distinct functions in Purkinje cells. *Eur J Neurosci* **42**, 2595–2612 (2015).

43. McCabe, M. P. *et al.* Genetic inactivation of mTORC1 or mTORC2 in neurons reveals distinct functions in glutamatergic synaptic transmission. *Elife* **9**, e51440 (2020).

44. Backman, S. A. *et al.* Deletion of Pten in mouse brain causes seizures, ataxia and defects in soma size resembling Lhermitte-Duclos disease. *Nat Genet* **29**, 396–403 (2001).

45. Williams, K. A. & Swedo, S. E. Post-infectious autoimmune disorders: Sydenham's chorea, PANDAS and beyond. *Brain Res* **1617**, 144–154 (2015).

46. Weston, M. C., Chen, H. & Swann, J. W. Loss of mTOR repressors Tsc1 or Pten has divergent effects on excitatory and inhibitory synaptic transmission in single hippocampal neuron cultures. *Front Mol Neurosci* **7**, 1 (2014).

47. Kwon, C. H. *et al.* Pten regulates neuronal soma size: a mouse model of Lhermitte-Duclos disease. *Nat Genet* **29**, 404–411 (2001).

48. Barrows, C. M., McCabe, M. P., Chen, H., Swann, J. W. & Weston, M. C. PTEN Loss Increases the Connectivity of Fast Synaptic Motifs and Functional Connectivity in a Developing Hippocampal Network. *J Neurosci* **37**, 8595–8611 (2017).

49. Trifonova, E. A., Mustafin, Z. S., Lashin, S. A. & Kochetov, A. V. Abnormal mTOR Activity in Pediatric Autoimmune Neuropsychiatric and MIA-Associated Autism Spectrum Disorders. *Int J Mol Sci* **23**, 967 (2022).

50. Zhao, J.-P. & Yoshii, A. Hyperexcitability of the local cortical circuit in mouse models of tuberous sclerosis complex. *Mol Brain* **12**, 6 (2019).

51. Malik, R. *et al.* Tsc1 represses parvalbumin expression and fast-spiking properties in somatostatin lineage cortical interneurons. *Nat Commun* **10**, 4994 (2019).

52. Artinian, J. *et al.* Regulation of Hippocampal Memory by mTORC1 in Somatostatin Interneurons. *J Neurosci* **39**, 8439–8456 (2019).

53. Fu, C. *et al.* GABAergic Interneuron Development and Function Is Modulated by the Tsc1 Gene. *Cereb Cortex* **22**, 2111–2119 (2012).

54. Haji, N. *et al.* Tsc1 haploinsufficiency in Nkx2.1 cells upregulates hippocampal interneuron mTORC1 activity, impairs pyramidal cell synaptic inhibition, and alters contextual fear discrimination and spatial working memory in mice. *Mol Autism* **11**, 29 (2020).

55. Weston, M. C., Chen, H. & Swann, J. W. Multiple Roles for Mammalian Target of Rapamycin Signaling in Both Glutamatergic and GABAergic Synaptic Transmission. *J Neurosci* **32**, 11441–11452 (2012).

56. Marchese, M. *et al.* Autism-epilepsy phenotype with macrocephaly suggests PTEN, but not GLIALCAM, genetic screening. *BMC Med Genet* **15**, 26 (2014).

57. Jansen, L. A. *et al.* PI3K/AKT pathway mutations cause a spectrum of brain malformations from megalencephaly to focal cortical dysplasia. *Brain* **138**, 1613–1628 (2015).

58. Koboldt, D. C. *et al.* PTEN somatic mutations contribute to spectrum of cerebral overgrowth. *Brain* **144**, 2971–2978 (2021).

59. Rosina, E. *et al.* Disruption of mTOR and MAPK pathways correlates with severity in idiopathic autism. *Transl Psychiatry* **9**, 1–10 (2019).

60. Arenella, M., Mota, N. R., Teunissen, M. W. A., Brunner, H. G. & Bralten, J. Autism spectrum disorder and brain volume link through a set of mTOR-related genes. *J Child Psychol Psychiatry* **64**, 1007–1014 (2023).

61. Aldinger, K. A., Plummer, J. T., Qiu, S. & Levitt, P. SnapShot: genetics of autism. *Neuron* **72**, 418-418.e1 (2011).

62. Xing, X. *et al.* Hyperactive Akt-mTOR pathway as a therapeutic target for pain hypersensitivity in Cntnap2-deficient mice. *Neuropharmacology* **165**, 107816 (2020).

63. Trifonova, E. A., Klimenko, A. I., Mustafin, Z. S., Lashin, S. A. & Kochetov, A. V. The mTOR Signaling Pathway Activity and Vitamin D Availability Control the Expression of Most Autism Predisposition Genes. *Int J Mol Sci* **20**, 6332 (2019).

64. Bi, X., Sun, J., Ji, A. X. & Baudry, M. Potential therapeutic approaches for Angelman syndrome. *Expert Opin Ther Targets* **20**, 601–613 (2016).

65. Kabitzke, P. A. *et al.* Comprehensive analysis of two Shank3 and the Cacna1c mouse models of autism spectrum disorder. *Genes Brain Behav* **17**, 4–22 (2018).

66. Roy, B., Amemasor, E., Hussain, S. & Castro, K. UBE3A: The Role in Autism Spectrum Disorders (ASDs) and a Potential Candidate for Biomarker Studies and Designing Therapeutic Strategies. *Diseases* **12**, 7 (2023).

67. Canales, C. P. *et al.* Sequential perturbations to mouse corticogenesis following in utero maternal immune activation. *Elife* **10**, e60100 (2021).

68. Basu, S. N., Kollu, R. & Banerjee-Basu, S. AutDB: a gene reference resource for autism research. *Nucleic Acids Res* **37**, D832-836 (2009).

69. Leblond, C. S. *et al.* Operative list of genes associated with autism and neurodevelopmental disorders based on database review. *Mol Cell Neurosci* **113**, 103623 (2021).

70. Wen, Y., Alshikho, M. J. & Herbert, M. R. Pathway Network Analyses for Autism Reveal Multisystem Involvement, Major Overlaps with Other Diseases and Convergence upon MAPK and Calcium Signaling. *PLoS One* **11**, e0153329 (2016).

71. Hu, Y. *et al.* mTOR-mediated metabolic reprogramming shapes distinct microglia functions in response to lipopolysaccharide and ATP. *Glia* **68**, 1031–1045 (2020).

72. Hodges, S. L. & Lugo, J. N. Therapeutic role of targeting mTOR signaling and neuroinflammation in epilepsy. *Epilepsy Res* **161**, 106282 (2020).

73. Haidinger, M. *et al.* A versatile role of mammalian target of rapamycin in human dendritic cell function and differentiation. *J Immunol* **185**, 3919–3931 (2010).

74. Lehman, J. A., Calvo, V. & Gomez-Cambronero, J. Mechanism of ribosomal p70S6 kinase activation by granulocyte macrophage colony-stimulating factor in neutrophils: cooperation of a MEK-related, THR421/SER424 kinase and a rapamycin-sensitive, m-TOR-related THR389 kinase. *J Biol Chem* **278**, 28130–28138 (2003).

75. Weichhart, T., Hengstschläger, M. & Linke, M. Regulation of innate immune cell function by mTOR. *Nat Rev Immunol* **15**, 599–614 (2015).

76. Xiao, Z., Peng, J., Wu, L., Arafat, A. & Yin, F. The effect of IL-1β on synaptophysin expression and electrophysiology of hippocampal neurons through the PI3K/Akt/mTOR signaling pathway in a rat model of mesial temporal lobe epilepsy. *Neurol Res* **39**, 640–648 (2017).

77. Argandona Lopez, C. & Brown, A. M. Microglial- neuronal crosstalk in chronic viral infection through mTOR, SPP1/OPN and inflammasome pathway signaling. *Front Immunol* **15**, 1368465 (2024).

78. Lashgari, N.-A. *et al.* TLR/mTOR inflammatory signaling pathway: novel insight for the treatment of schizophrenia. *Can J Physiol Pharmacol* **102**, 150–160 (2024).

79.	Sharma, A. & Mehan, S. Targeting PI3K-AKT/mTOR signaling in the prevention of autism. *Neurochem Int* **147**, 105067 (2021).

80.	Yang, H. *et al.* Modulation of TSC-mTOR signaling on immune cells in immunity and autoimmunity. *J Cell Physiol* **229**, 17–26 (2014).

81.	Byles, V. *et al.* The TSC-mTOR pathway regulates macrophage polarization. *Nat Commun* **4**, 2834 (2013).

82.	Vergadi, E., Ieronymaki, E., Lyroni, K., Vaporidi, K. & Tsatsanis, C. Akt Signaling Pathway in Macrophage Activation and M1/M2 Polarization. *J Immunol* **198**, 1006–1014 (2017).

83.	Krakowiak, P. *et al.* Neonatal Cytokine Profiles Associated With Autism Spectrum Disorder. *Biol Psychiatry* **81**, 442–451 (2017).

84.	Estes, M. L. & McAllister, A. K. Immune mediators in the brain and peripheral tissues in autism spectrum disorder. *Nat Rev Neurosci* **16**, 469–486 (2015).

85.	Patterson, P. H. Immune involvement in schizophrenia and autism: etiology, pathology and animal models. *Behav Brain Res* **204**, 313–321 (2009).

86.	Xiong, Y., Chen, J. & Li, Y. Microglia and astrocytes underlie neuroinflammation and synaptic susceptibility in autism spectrum disorder. *Front Neurosci* **17**, 1125428 (2023).

87.	Matta, S. M., Hill-Yardin, E. L. & Crack, P. J. The influence of neuroinflammation in Autism Spectrum Disorder. *Brain Behav Immun* **79**, 75–90 (2019).

88.	Vargas, D. L., Nascimbene, C., Krishnan, C., Zimmerman, A. W. & Pardo, C. A. Neuroglial activation and neuroinflammation in the brain of patients with autism. *Ann Neurol* **57**, 67–81 (2005).

89.	Gesundheit, B. *et al.* Immunological and autoimmune considerations of Autism Spectrum Disorders. *J Autoimmun* **44**, 1–7 (2013).

90.	Theoharides, T. C., Asadi, S. & Patel, A. B. Focal brain inflammation and autism. *J Neuroinflammation* **10**, 46 (2013).

91. Voineagu, I. *et al.* Transcriptomic analysis of autistic brain reveals convergent molecular pathology. *Nature* **474**, 380–384 (2011).

92. Morgan, J. T. *et al.* Microglial activation and increased microglial density observed in the dorsolateral prefrontal cortex in autism. *Biol Psychiatry* **68**, 368–376 (2010).

93. Laurence, J. A. & Fatemi, S. H. Glial fibrillary acidic protein is elevated in superior frontal, parietal and cerebellar cortices of autistic subjects. *Cerebellum* **4**, 206–210 (2005).

94. Edmonson, C., Ziats, M. N. & Rennert, O. M. Altered glial marker expression in autistic post-mortem prefrontal cortex and cerebellum. *Mol Autism* **5**, 3 (2014).

95. Masi, A., Glozier, N., Dale, R. & Guastella, A. J. The Immune System, Cytokines, and Biomarkers in Autism Spectrum Disorder. *Neurosci Bull* **33**, 194–204 (2017).

96. Gandal, M. J. *et al.* Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science* **362**, eaat8127 (2018).

97. Li, X. *et al.* Elevated immune response in the brain of autistic patients. *J Neuroimmunol* **207**, 111–116 (2009).

98. Courchesne, E. Brain development in autism: early overgrowth followed by premature arrest of growth. *Ment Retard Dev Disabil Res Rev* **10**, 106–111 (2004).

99. Courchesne, E., Campbell, K. & Solso, S. Brain growth across the life span in autism: age-specific changes in anatomical pathology. *Brain Res* **1380**, 138–145 (2011).

100. Redcay, E. & Courchesne, E. When is the brain enlarged in autism? A meta-analysis of all brain size reports. *Biol Psychiatry* **58**, 1–9 (2005).

101. Fatemi, S. H. *et al.* Prenatal viral infection leads to pyramidal cell atrophy and macrocephaly in adulthood: implications for genesis of autism and schizophrenia. *Cell Mol Neurobiol* **22**, 25–33 (2002).

102. Smith, S. E. P., Li, J., Garbett, K., Mirnics, K. & Patterson, P. H. Maternal Immune Activation Alters Fetal Brain Development through Interleukin-6. *J Neurosci* **27**, 10695–10702 (2007).

103. Shi, L., Fatemi, S. H., Sidwell, R. W. & Patterson, P. H. Maternal influenza infection causes marked behavioral and pharmacological changes in the offspring. *J Neurosci* **23**, 297–302 (2003).

104. Malkova, N. V., Yu, C. Z., Hsiao, E. Y., Moore, M. J. & Patterson, P. H. Maternal immune activation yields offspring displaying mouse versions of the three core symptoms of autism. *Brain Behav Immun* **26**, 607–616 (2012).

105. Choi, G. B. *et al.* The maternal interleukin-17a pathway in mice promotes autism-like phenotypes in offspring. *Science* **351**, 933–939 (2016).

106. Atladóttir, H. O. *et al.* Association of family history of autoimmune diseases and autism spectrum disorders. *Pediatrics* **124**, 687–694 (2009).

107. Atladóttir, H. O. *et al.* Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord* **40**, 1423–1430 (2010).

108. Brown, A. S. *et al.* Elevated maternal C-reactive protein and autism in a national birth cohort. *Mol Psychiatry* **19**, 259–264 (2014).

109. Ashwood, P., Wills, S. & Van de Water, J. The immune response in autism: a new frontier for autism research. *J Leukoc Biol* **80**, 1–15 (2006).

110. Lee, B. K. *et al.* Maternal hospitalization with infection during pregnancy and risk of autism spectrum disorders. *Brain Behav Immun* **44**, 100–105 (2015).

111. Modabbernia, A., Velthorst, E. & Reichenberg, A. Environmental risk factors for autism: an evidence-based review of systematic reviews and meta-analyses. *Mol Autism* **8**, 13 (2017).

112. Goines, P. E. & Ashwood, P. Cytokine dysregulation in autism spectrum disorders (ASD): possible role of the environment. *Neurotoxicol Teratol* **36**, 67–81 (2013).

113. Jiang, H.-Y. *et al.* Maternal infection during pregnancy and risk of autism spectrum disorders: A systematic review and meta-analysis. *Brain Behav Immun* **58**, 165–172 (2016).

114. Kolevzon, A., Gross, R. & Reichenberg, A. Prenatal and perinatal risk factors for autism: a review and integration of findings. *Arch Pediatr Adolesc Med* **161**, 326–333 (2007).

115. Kancherla, V. & Dennis, L. K. A Meta-analysis of Prenatal, Perinatal, and Neonatal Risks for Autism. *American Journal of Epidemiology* **163**, S20 (2006).

116. Gardener, H., Spiegelman, D. & Buka, S. L. Prenatal Risk Factors for Autism: A Comprehensive Meta-analysis. *Br J Psychiatry* **195**, 7–14 (2009).

117. Gardener, H., Spiegelman, D. & Buka, S. L. Perinatal and neonatal risk factors for autism: a comprehensive meta-analysis. *Pediatrics* **128**, 344–355 (2011).

118. Le Belle, J. E. *et al.* Maternal inflammation contributes to brain overgrowth and autism-associated behaviors through altered redox signaling in stem and progenitor cells. *Stem Cell Reports* **3**, 725–734 (2014).

119. Hazen, E. P., Stornelli, J. L., O'Rourke, J. A., Koesterer, K. & McDougle, C. J. Sensory symptoms in autism spectrum disorders. *Harv Rev Psychiatry* **22**, 112–124 (2014).

120. Narvaiz, D. A. *et al.* Rapamycin improves social and stereotypic behavior abnormalities induced by pre-mitotic neuronal subset specific Pten deletion. *Genes Brain Behav* **22**, e12854 (2023).

121. Zhang, J., Zhang, J.-X. & Zhang, Q.-L. PI3K/AKT/mTOR-mediated autophagy in the development of autism spectrum disorder. *Brain Res Bull* **125**, 152–158 (2016).

122. Lieberman, O. J. *et al.* mTOR Suppresses Macroautophagy During Striatal Postnatal Development and Is Hyperactive in Mouse Models of Autism Spectrum Disorders. *Front Cell Neurosci* **14**, 70 (2020).

123. Ehninger, D. & Silva, A. J. Rapamycin for treating Tuberous sclerosis and Autism spectrum disorders. *Trends Mol Med* **17**, 78–87 (2011).

124. Ehninger, D. *et al.* Reversal of learning deficits in a Tsc2+/- mouse model of tuberous sclerosis. *Nat Med* **14**, 843–848 (2008).

125. Kotajima-Murakami, H. *et al.* Effects of rapamycin on social interaction deficits and gene expression in mice exposed to valproic acid in utero. *Mol Brain* **12**, 3 (2019).

126. Meikle, L. *et al.* Response of a Neuronal Model of Tuberous Sclerosis to Mammalian Target of Rapamycin (mTOR) Inhibitors: Effects on mTORC1 and Akt Signaling Lead to Improved Survival and Function. *J Neurosci* **28**, 5422–5432 (2008).

127. Getz, S. A., DeSpenza, T., Li, M. & Luikart, B. W. Rapamycin prevents, but does not reverse, aberrant migration in Pten knockout neurons. *Neurobiol Dis* **93**, 12–20 (2016).

128. Amegandjin, C. A. *et al.* Sensitive period for rescuing parvalbumin interneurons connectivity and social behavior deficits caused by TSC1 loss. *Nat Commun* **12**, 3653 (2021).

129. Tsai, P. T. *et al.* Autistic-like behaviour and cerebellar dysfunction in Purkinje cell Tsc1 mutant mice. *Nature* **488**, 647–651 (2012).

130. Zeng, L.-H., Xu, L., Gutmann, D. H. & Wong, M. Rapamycin prevents epilepsy in a mouse model of tuberous sclerosis complex. *Ann Neurol* **63**, 444–453 (2008).

131. Tsai, P. T. *et al.* Sensitive Periods for Cerebellar-Mediated Autistic-like Behaviors. *Cell Rep* **25**, 357-367.e4 (2018).

132. Tang, G. *et al.* Loss of mTOR-dependent macroautophagy causes autistic-like synaptic pruning deficits. *Neuron* **83**, 1131–1143 (2014).

133. Krueger, D. A. *et al.* Everolimus for subependymal giant-cell astrocytomas in tuberous sclerosis. *N Engl J Med* **363**, 1801–1811 (2010).

134. Krueger, D. A. *et al.* Everolimus treatment of refractory epilepsy in tuberous sclerosis complex. *Ann Neurol* **74**, 679–687 (2013).

135. Cardamone, M. *et al.* Mammalian target of rapamycin inhibitors for intractable epilepsy and subependymal giant cell astrocytomas in tuberous sclerosis complex. *J Pediatr* **164**, 1195–1200 (2014).

136. Kotulska, K. *et al.* Long-term effect of everolimus on epilepsy and growth in children under 3 years of age treated for subependymal giant cell astrocytoma associated with tuberous sclerosis complex. *Eur J Paediatr Neurol* **17**, 479–485 (2013).

137. MacKeigan, J. P. & Krueger, D. A. Differentiating the mTOR inhibitors everolimus and sirolimus in the treatment of tuberous sclerosis complex. *Neuro Oncol* **17**, 1550–1559 (2015).

138. Srivastava, S. *et al.* A randomized controlled trial of everolimus for neurocognitive symptoms in PTEN hamartoma tumor syndrome. *Hum Mol Genet* **31**, 3393–3404 (2022).

139. Overwater, I. E. *et al.* A randomized controlled trial with everolimus for IQ and autism in tuberous sclerosis complex. *Neurology* **93**, e200–e209 (2019).

140. Davies, D. M. *et al.* Sirolimus therapy for angiomyolipoma in tuberous sclerosis and sporadic lymphangioleiomyomatosis: a phase 2 trial. *Clin Cancer Res* **17**, 4071–4081 (2011).

141. Mizuguchi, M. *et al.* Everolimus for epilepsy and autism spectrum disorder in tuberous sclerosis complex: EXIST-3 substudy in Japan. *Brain Dev* **41**, 1–10 (2019).

142. Hwang, S.-K. *et al.* Everolimus improves neuropsychiatric symptoms in a patient with tuberous sclerosis carrying a novel TSC2 mutation. *Mol Brain* **9**, 56 (2016).

143. Kilincaslan, A. *et al.* Beneficial Effects of Everolimus on Autism and Attention-Deficit/Hyperactivity Disorder Symptoms in a Group of Patients with Tuberous Sclerosis Complex. *J Child Adolesc Psychopharmacol* **27**, 383–388 (2017).

144. Ess, K. C. & Franz, D. N. Everolimus for cognition/autism in children with tuberous sclerosis complex: Definitive outcomes deferred. *Neurology* **93**, 51–52 (2019).

145. Saffari, A. *et al.* Safety and efficacy of mTOR inhibitor treatment in patients with tuberous sclerosis complex under 2 years of age - a multicenter retrospective study. *Orphanet J Rare Dis* **14**, 96 (2019).

146. Hodges, S. L. *et al.* Rapamycin, but not minocycline, significantly alters ultrasonic vocalization behavior in C57BL/6J pups in a flurothyl seizure model. *Behav Brain Res* **410**, 113317 (2021).

147. Qin, L., Dai, X. & Yin, Y. Valproic acid exposure sequentially activates Wnt and mTOR pathways in rats. *Mol Cell Neurosci* **75**, 27–35 (2016).

148. McMahon, J. J. *et al.* Seizure-dependent mTOR activation in 5-HT neurons promotes autism-like behaviors in mice. *Neurobiol Dis* **73**, 296–306 (2015).

149. Burket, J. A., Benson, A. D., Tang, A. H. & Deutsch, S. I. Rapamycin improves sociability in the BTBR T+Itpr3tf/J mouse model of autism spectrum disorders. *Brain Res Bull* **100**, 70–75 (2014).

150. Silverman, J. L., Yang, M., Lord, C. & Crawley, J. N. Behavioural phenotyping assays for mouse models of autism. *Nat Rev Neurosci* **11**, 490–502 (2010).

151. Peñagarikano, O. *et al.* Absence of CNTNAP2 leads to epilepsy, neuronal migration abnormalities, and core autism-related deficits. *Cell* **147**, 235–246 (2011).

152. Orefice, L. L. *et al.* Peripheral Mechanosensory Neuron Dysfunction Underlies Tactile and Behavioral Deficits in Mouse Models of ASDs. *Cell* **166**, 299–313 (2016).

153. Bonin, R. P., Bories, C. & De Koninck, Y. A simplified up-down method (SUDO) for measuring mechanical nociception in rodents using von Frey filaments. *Mol Pain* **10**, 26 (2014).

154. Crawley, J. & Goodwin, F. K. Preliminary report of a simple animal behavior model for the anxiolytic effects of benzodiazepines. *Pharmacol Biochem Behav* **13**, 167–170 (1980).

155. Fleming, S. J. *et al.* Unsupervised removal of systematic background noise from droplet-based single-cell experiments using CellBender. *Nat Methods* **20**, 1323–1335 (2023).

156. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet identification in single-cell sequencing data using scDblFinder. *F1000Res* **10**, 979 (2021).

157. Heinemann, J. Cluster Analysis of Untargeted Metabolomic Experiments. *Methods Mol Biol* **1859**, 275–285 (2019).

158. Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902.e21 (2019).

159. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

160. Phipson, B. *et al.* propeller: testing for differences in cell type proportions in single cell data. *Bioinformatics* **38**, 4720–4726 (2022).

161. Squair, J. W. *et al.* Confronting false discoveries in single-cell differential expression. *Nat Commun* **12**, 5692 (2021).

162. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545–15550 (2005).

163. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417–425 (2015).

164. Castanza, A. S. *et al.* Extending support for mouse data in the Molecular Signatures Database (MSigDB). *Nat Methods* **20**, 1619–1620 (2023).

165. Korotkevich, G. *et al.* Fast gene set enrichment analysis. 060012 Preprint at https://doi.org/10.1101/060012 (2021).

166. Adamczak, J. M., Farr, T. D., Seehafer, J. U., Kalthoff, D. & Hoehn, M. High field BOLD response to forepaw stimulation in the mouse. *Neuroimage* **51**, 704–712 (2010).

167. Magnuson, M. E., Thompson, G. J., Pan, W.-J. & Keilholz, S. D. Time-dependent effects of isoflurane and dexmedetomidine on functional connectivity, spectral characteristics, and spatial distribution of spontaneous BOLD fluctuations. *NMR Biomed* **27**, 291–303 (2014).

168. Oguz, I., Zhang, H., Rumple, A. & Sonka, M. RATS: Rapid Automatic Tissue Segmentation in rodent brain MRI. *J Neurosci Methods* **221**, 175–182 (2014).

169. Jenkinson, M. & Smith, S. A global optimisation method for robust affine registration of brain images. *Med Image Anal* **5**, 143–156 (2001).

170. Avants, B. B. *et al.* The Insight ToolKit image registration framework. *Front Neuroinform* **8**, 44 (2014).

171. Smith, S. M. *et al.* Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* **23 Suppl 1**, S208-219 (2004).

172. Zalesky, A., Fornito, A. & Bullmore, E. T. Network-based statistic: identifying differences in brain networks. *Neuroimage* **53**, 1197–1207 (2010).

173. Kruschwitz, J. D., List, D., Waller, L., Rubinov, M. & Walter, H. GraphVar: a user-friendly toolbox for comprehensive graph analyses of functional brain connectivity. *J Neurosci Methods* **245**, 107–115 (2015).

174. Worsley, K. J. Statistical analysis of activation images. in *Functional Magnetic Resonance Imaging: An Introduction to Methods* (eds. Jezzard, P., Matthews, P. M. & Smith, S. M.) 0 (Oxford University Press, 2001). doi:10.1093/acprof:oso/9780192630711.003.0014.

175. Blondel, V. D., Guillaume, J.-L., Lambiotte, R. & Lefebvre, E. Fast unfolding of communities in large networks. *J. Stat. Mech.* **2008**, P10008 (2008).

176. Fornito, A., Zalesky, A. & Breakspear, M. Graph analysis of the human connectome: promise, progress, and pitfalls. *Neuroimage* **80**, 426–444 (2013).

177. M, J. BET2 : MR-Based Estimation of Brain, Skull and Scalp Surfaces. *Eleventh Annual Meeting of the Organization for Human Brain Mapping, 2005* **17**, 167 (2005).

178. Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage* **54**, 2033–2044 (2011).

179. Avants, B. B., Epstein, C. L., Grossman, M. & Gee, J. C. Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Med Image Anal* **12**, 26–41 (2008).

180. Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M. & Nichols, T. E. Permutation inference for the general linear model. *Neuroimage* **92**, 381–397 (2014).

181. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300 (1995).

182. Genovese, C. R., Lazar, N. A. & Nichols, T. Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage* **15**, 870–878 (2002).

183. Zhang, Z. *et al.* Brain-restricted mTOR inhibition with binary pharmacology. *Nature* **609**, 822–828 (2022).

184. Ehinger, Y. *et al.* Brain-specific inhibition of mTORC1 eliminates side effects resulting from mTORC1 blockade in the periphery and reduces alcohol intake in mice. *Nat Commun* **12**, 4407 (2021).

185. Huang, W.-C., Chen, Y. & Page, D. T. Hyperconnectivity of prefrontal cortex to amygdala projections in a mouse model of macrocephaly/autism syndrome. *Nat Commun* **7**, 13421 (2016).

186. Shimada, T. & Yamagata, K. Pentylenetetrazole-Induced Kindling Mouse Model. *J Vis Exp* 56573 (2018) doi:10.3791/56573.

187. Braden, B. B. & Riecken, C. Thinning Faster? Age-Related Cortical Thickness Differences in Adults with Autism Spectrum Disorder. *Res Autism Spectr Disord* **64**, 31–38 (2019).

188. Tamura, R., Kitamura, H., Endo, T., Hasegawa, N. & Someya, T. Reduced thalamic volume observed across different subgroups of autism spectrum disorders. *Psychiatry Res* **184**, 186–188 (2010).

189. Qiu, T. *et al.* Two years changes in the development of caudate nucleus are involved in restricted repetitive behaviors in 2-5-year-old children with autism spectrum disorder. *Dev Cogn Neurosci* **19**, 137–143 (2016).

190. Banker, S. M., Gu, X., Schiller, D. & Foss-Feig, J. H. Hippocampal contributions to social and cognitive deficits in autism spectrum disorder. *Trends Neurosci* **44**, 793–807 (2021).

191. Schumann, C. M. *et al.* The amygdala is enlarged in children but not adolescents with autism; the hippocampus is enlarged at all ages. *J Neurosci* **24**, 6392–6401 (2004).

192. Schumann, C. M. & Amaral, D. G. Stereological analysis of amygdala neuron number in autism. *J Neurosci* **26**, 7674–7679 (2006).

193. Turner, A. H., Greenspan, K. S. & van Erp, T. G. M. Pallidum and lateral ventricle volume enlargement in autism spectrum disorder. *Psychiatry Res Neuroimaging* **252**, 40–45 (2016).

194. Green, S. A. & Ben-Sasson, A. Anxiety Disorders and Sensory Over-Responsivity in Children with Autism Spectrum Disorders: Is There a Causal Relationship? *J Autism Dev Disord* **40**, 1495–1504 (2010).

195. Simmons, D. H. *et al.* Sensory Over-responsivity and Aberrant Plasticity in Cerebellar Cortex in a Mouse Model of Syndromic Autism. *Biol Psychiatry Glob Open Sci* **2**, 450–459 (2022).

196. Tavassoli, T., Miller, L. J., Schoen, S. A., Nielsen, D. M. & Baron-Cohen, S. Sensory over-responsivity in adults with autism spectrum conditions. *Autism* **18**, 428–432 (2014).

197. Tavassoli, T. *et al.* Sensory over-responsivity: parent report, direct assessment measures, and neural architecture. *Molecular Autism* **10**, 4 (2019).

198. Orefice, L. L. *et al.* Targeting peripheral somatosensory neurons to improve tactile-related phenotypes in ASD models. *Cell* **178**, 867-886.e24 (2019).

199. Lewis, M. L., Kesler, M., Candy, S. A., Rho, J. M. & Pittman, Q. J. Comorbid epilepsy in autism spectrum disorder: Implications of postnatal inflammation for brain excitability. *Epilepsia* **59**, 1316–1326 (2018).

200. Masi, A. *et al.* Cytokine aberrations in autism spectrum disorder: a systematic review and meta-analysis. *Mol Psychiatry* **20**, 440–446 (2015).

201. Williams, J. A. *et al.* Inflammation and Brain Structure in Schizophrenia and Other Neuropsychiatric Disorders: A Mendelian Randomization Study. *JAMA Psychiatry* **79**, 498–507 (2022).

202. Than, U. T. T. *et al.* Inflammatory mediators drive neuroinflammation in autism spectrum disorder and cerebral palsy. *Sci Rep* **13**, 22587 (2023).

203. Kern, J. K., Geier, D. A., Sykes, L. K. & Geier, M. R. Relevance of Neuroinflammation and Encephalitis in Autism. *Front Cell Neurosci* **9**, 519 (2015).

204. Onore, C., Careaga, M. & Ashwood, P. The role of immune dysfunction in the pathophysiology of autism. *Brain Behav Immun* **26**, 383–392 (2012).

205. Pardo, C. A., Vargas, D. L. & Zimmerman, A. W. Immunity, neuroglia and neuroinflammation in autism. *Int Rev Psychiatry* **17**, 485–495 (2005).

206. Ben-Sasson, A., Cermak, S. A., Orsmond, G. I., Carter, A. S. & Fogg, L. Can we differentiate sensory over-responsivity from anxiety symptoms in toddlers? perspectives of occupational therapists and psychologists. *Infant Ment Health J* **28**, 536–558 (2007).

207. Ben-Sasson, A. *et al.* Sensory clusters of toddlers with autism spectrum disorders: differences in affective symptoms. *J Child Psychol Psychiatry* **49**, 817–825 (2008).

208. Glod, M., Riby, D. M., Honey, E. & Rodgers, J. Psychological Correlates of Sensory Processing Patterns in Individuals with Autism Spectrum Disorder: A Systematic Review. *Rev J Autism Dev Disord* **2**, 199–221 (2015).

209. Ausderau, K. K. *et al.* Sensory subtypes and associated outcomes in children with autism spectrum disorders. *Autism Res* **9**, 1316–1327 (2016).

210. Siniscalco, D., Schultz, S., Brigida, A. L. & Antonucci, N. Inflammation and Neuro-Immune Dysregulations in Autism Spectrum Disorders. *Pharmaceuticals (Basel)* **11**, 56 (2018).

211. Tartaglione, A. M. *et al.* Maternal immune activation induces autism-like changes in behavior, neuroinflammatory profile and gut microbiota in mouse offspring of both sexes. *Transl Psychiatry* **12**, 384 (2022).

212. Kim, E. *et al.* Maternal gut bacteria drive intestinal inflammation in offspring with neurodevelopmental disorders by altering the chromatin landscape of CD4+ T cells. *Immunity* **55**, 145-158.e7 (2022).

213. Shimizu, Y., Sakata-Haga, H., Saikawa, Y. & Hatta, T. Influence of Immune System Abnormalities Caused by Maternal Immune Activation in the Postnatal Period. *Cells* **12**, 741 (2023).

214. Lucchina, L. & Depino, A. M. Altered peripheral and central inflammatory responses in a mouse model of autism. *Autism Res* **7**, 273–289 (2014).

215. Hsiao, E. Y., McBride, S. W., Chow, J., Mazmanian, S. K. & Patterson, P. H. Modeling an autism risk factor in mice leads to permanent immune dysregulation. *Proc Natl Acad Sci U S A* **109**, 12776–12781 (2012).

216. Kirsten, T. B., Casarin, R. C., Bernardi, M. M. & Felicio, L. F. Pioglitazone abolishes autistic-like behaviors via the IL-6 pathway. *PLoS One* **13**, e0197060 (2018).

217. Kirsten, T. B., Casarin, R. C., Bernardi, M. M. & Felicio, L. F. Pioglitazone abolishes cognition impairments as well as BDNF and neurotensin disturbances in a rat model of autism. *Biol Open* **8**, bio041327 (2019).

218. Sandhu, A. *et al.* Ameliorating effect of pioglitazone on prenatal valproic acid-induced behavioral and neurobiological abnormalities in autism spectrum disorder in rats. *Pharmacol Biochem Behav* **237**, 173721 (2024).

219. Buehler, M. R. A proposed mechanism for autism: an aberrant neuroimmune response manifested as a psychiatric disorder. *Med Hypotheses* **76**, 863–870 (2011).

220. Emanuele, E., Lossano, C., Politi, P. & Barale, F. Pioglitazone as a therapeutic agent in autistic spectrum disorder. *Medical Hypotheses* **69**, 699 (2007).

221. Boris, M. *et al.* Effect of pioglitazone treatment on behavioral symptoms in autistic children. *J Neuroinflammation* **4**, 3 (2007).

222. Mirza, R. & Sharma, B. Beneficial effects of pioglitazone, a selective peroxisome proliferator-activated receptor-γ agonist in prenatal valproic acid-induced behavioral and biochemical autistic like features in Wistar rats. *Int J Dev Neurosci* **76**, 6–16 (2019).

223. Mirza, R. & Sharma, B. A selective peroxisome proliferator-activated receptor-γ agonist benefited propionic acid induced autism-like behavioral phenotypes in rats by attenuation of neuroinflammation and oxidative stress. *Chem Biol Interact* **311**, 108758 (2019).

224. Capano, L. *et al.* A pilot dose finding study of pioglitazone in autistic children. *Mol Autism* **9**, 59 (2018).

225. Reed, M. D. *et al.* IL-17a promotes sociability in mouse models of neurodevelopmental disorders. *Nature* **577**, 249–253 (2020).

226. Luo, Y. *et al.* Minocycline improves autism-related behaviors by modulating microglia polarization in a mouse model of autism. *Int Immunopharmacol* **122**, 110594 (2023).

227. Kim, H.-J. *et al.* Deficient autophagy in microglia impairs synaptic pruning and causes social behavioral defects. *Mol Psychiatry* **22**, 1576–1584 (2017).

228. Xu, X., Hanganu-Opatz, I. L. & Bieler, M. Cross-Talk of Low-Level Sensory and High-Level Cognitive Processing: Development, Mechanisms, and Relevance for Cross-Modal Abilities of the Brain. *Frontiers in Neurorobotics* **14**, (2020).

229. Patel, A. B., Tsilioni, I., Leeman, S. E. & Theoharides, T. C. Neurotensin stimulates sortilin and mTOR in human microglia inhibitable by methoxyluteolin, a potential therapeutic target for autism. *Proc Natl Acad Sci U S A* **113**, E7049–E7058 (2016).

230. Dong, J. *et al.* The effect of intrauterine inflammation on mTOR signaling in mouse fetal brain. *Dev Neurobiol* **80**, 149–159 (2020).

231. Zhao, X. *et al.* Noninflammatory Changes of Microglia Are Sufficient to Cause Epilepsy. *Cell Rep* **22**, 2080–2093 (2018).

232. Zimmer, T. S. *et al.* Tuberous Sclerosis Complex as Disease Model for Investigating mTOR-Related Gliopathy During Epileptogenesis. *Front Neurol* **11**, 1028 (2020).

233. Huang, A. S. *et al.* Characterizing effects of age, sex and psychosis symptoms on thalamocortical functional connectivity in youth. *Neuroimage* **243**, 118562 (2021).

234. Villasana-Salazar, B. & Vezzani, A. Neuroinflammation microenvironment sharpens seizure circuit. *Neurobiol Dis* **178**, 106027 (2023).

235. Jiang, P. *et al.* Icariin alleviates autistic-like behavior, hippocampal inflammation and vGlut1 expression in adult BTBR mice. *Behav Brain Res* **445**, 114384 (2023).

236. O'Donnell, C. Nonlinear slow-timescale mechanisms in synaptic plasticity. *Current Opinion in Neurobiology* **82**, 102778 (2023).

237. Gonzalez, K. C., Negrean, A., Liao, Z., Polleux, F. & Losonczy, A. Synaptic Basis of Behavioral Timescale Plasticity. 2023.10.04.560848 Preprint at https://doi.org/10.1101/2023.10.04.560848 (2023).

238. Duarte, R., Seeholzer, A., Zilles, K. & Morrison, A. Synaptic patterning and the timescales of cortical dynamics. *Current Opinion in Neurobiology* **43**, 156–165 (2017).

239. Gandal, M. J. *et al.* Shared molecular neuropathology across major psychiatric disorders parallels polygenic overlap. *Science* **359**, 693–697 (2018).

240. Okur, Z. *et al.* Control of neuronal excitation-inhibition balance by BMP-SMAD1 signalling. *Nature* **629**, 402–409 (2024).

241. Peddada, S., Yasui, D. H. & LaSalle, J. M. Inhibitors of differentiation (ID1, ID2, ID3 and ID4) genes are neuronal targets of MeCP2 that are elevated in Rett syndrome. *Hum Mol Genet* **15**, 2003–2014 (2006).

242. van Ravenswaaij-Arts, C. M., Hefner, M., Blake, K. & Martin, D. M. CHD7 Disorder. in *GeneReviews®* (eds. Adam, M. P. et al.) (University of Washington, Seattle, Seattle (WA), 1993).

243. Chen, H., Xu, G., Du, H., Yi, M. & Li, C. Integrative analysis of gene expression associated with epilepsy in human epilepsy and animal models. *Mol Med Rep* **13**, 4920–4926 (2016).

244. Lee, T.-S. *et al.* Gene Expression in Temporal Lobe Epilepsy is Consistent with Increased Release of Glutamate by Astrocytes. *Mol Med* **13**, 1–13 (2007).

245. Ryan, R. *et al.* Functional characterization of tektin-1 in motile cilia and evidence for TEKT1 as a new candidate gene for motile ciliopathies. *Hum Mol Genet* **27**, 266–282 (2018).

246. Khalil, R. *et al.* PSMD12 haploinsufficiency in a neurodevelopmental disorder with autistic features. *Am J Med Genet B Neuropsychiatr Genet* **177**, 736–745 (2018).

247. Montgomery, S. H. & Mundy, N. I. Evolution of ASPM is associated with both increases and decreases in brain size in primates. *Evolution* **66**, 927–932 (2012).

248. Bishop, D. V. M. Genes, cognition, and communication: insights from neurodevelopmental disorders. *Ann N Y Acad Sci* **1156**, 1–18 (2009).

249. Shen, J. *et al.* ASPM mutations identified in patients with primary microcephaly and seizures. *J Med Genet* **42**, 725–729 (2005).

250. Eagleson, K. L. *et al.* Genetic disruption of the autism spectrum disorder risk gene PLAUR induces GABAA receptor subunit changes. *Neuroscience* **168**, 797–810 (2010).

251. Eagleson, K. L., Campbell, D. B., Thompson, B. L., Bergman, M. Y. & Levitt, P. The autism risk genes MET and PLAUR differentially impact cortical development. *Autism Res* **4**, 68–83 (2011).

252. Spinelli, E. *et al.* Pathogenic MAST3 Variants in the STK Domain Are Associated with Epilepsy. *Ann Neurol* **90**, 274–284 (2021).

253. Xu, Q. *et al.* MAST3 modulates the inflammatory response and proliferation of fibroblast-like synoviocytes in rheumatoid arthritis. *Int Immunopharmacol* **77**, 105900 (2019).

254. Wang, Y., Li, Y., Wang, G., Lu, J. & Li, Z. Overexpression of Homer1b/c induces valproic acid resistance in epilepsy. *CNS Neurosci Ther* **29**, 331–343 (2023).

255. Phadke, L. *et al.* A primary rodent triculture model to investigate the role of glia-neuron crosstalk in regulation of neuronal activity. *Front Aging Neurosci* **14**, 1056067 (2022).

256. Du, Y., Brennan, F. H., Popovich, P. G. & Zhou, M. Microglia maintain the normal structure and function of the hippocampal astrocyte network. *Glia* **70**, 1359–1379 (2022).

257. Shan, W. *et al.* Neuronal PAS domain protein 4 (Npas4) controls neuronal homeostasis in pentylenetetrazole-induced epilepsy through the induction of Homer1a. *J Neurochem* **145**, 19–33 (2018).

258. Gomez Perdiguero, E. *et al.* Tissue-resident macrophages originate from yolk-sac-derived erythro-myeloid progenitors. *Nature* **518**, 547–551 (2015).

259. Wang, S. *et al.* IQGAP3, a novel effector of Rac1 and Cdc42, regulates neurite outgrowth. *J Cell Sci* **120**, 567–577 (2007).

260. Reilly, M. L. *et al.* Loss-of-function mutations in KIF14 cause severe microcephaly and kidney development defects in humans and zebrafish. *Hum Mol Genet* **28**, 778–795 (2019).

261. Zheng, E., Cai, Z., Li, W., Ni, C. & Fang, Q. Achaete-scute complex-like 2 regulated inflammatory mechanism through Toll-like receptor 4 activating in stomach adenocarcinoma. *World J Surg Oncol* **20**, 266 (2022).

262. Tong, D.-L. *et al.* The critical role of ASD-related gene CNTNAP3 in regulating synaptic development and social behavior in mice. *Neurobiol Dis* **130**, 104486 (2019).

263. Firouzabadi, N. *et al.* Genetic Variants of Angiotensin-Converting Enzyme Are Linked to Autism: A Case-Control Study. *PLoS One* **11**, e0153667 (2016).

264. Pereira, M. G. A. G. *et al.* Inhibition of the renin-angiotensin system prevents seizures in a rat model of epilepsy. *Clin Sci (Lond)* **119**, 477–482 (2010).

265. Yu, S., Wang, G., Yao, B., Xiao, L. & Tuo, H. Arc and Homer1 are involved in comorbid epilepsy and depression: A microarray data analysis. *Epilepsy Behav* **132**, 108738 (2022).

266. Tărlungeanu, D. C. *et al.* Impaired Amino Acid Transport at the Blood Brain Barrier Is a Cause of Autism Spectrum Disorder. *Cell* **167**, 1481-1494.e18 (2016).

267. Fudge, N. J. & Mearow, K. M. Extracellular matrix-associated gene expression in adult sensory neuron populations cultured on a laminin substrate. *BMC Neurosci* **14**, 15 (2013).

268. Chénier, S. *et al.* CHD2 haploinsufficiency is associated with developmental delay, intellectual disability, epilepsy and neurobehavioural problems. *J Neurodev Disord* **6**, 9 (2014).

269. Cepeda, C. *et al.* Cellular antiseizure mechanisms of everolimus in pediatric tuberous sclerosis complex, cortical dysplasia, and non-mTOR-mediated etiologies. *Epilepsia Open* **3**, 180–190 (2018).

270. Boyd, B. A. *et al.* Sensory features and repetitive behaviors in children with autism and developmental delays. *Autism Res* **3**, 78–87 (2010).

271. Joosten, A. V. & Bundy, A. C. Sensory processing and stereotypical and repetitive behaviour in children with autism and intellectual disability. *Aust Occup Ther J* **57**, 366–372 (2010).

272. Riby, D. M., Janes, E. & Rodgers, J. Brief report: exploring the relationship between sensory processing and repetitive behaviours in Williams Syndrome. *J Autism Dev Disord* **43**, 478–482 (2013).

273. Fetta, A. *et al.* Relationship between Sensory Alterations and Repetitive Behaviours in Children with Autism Spectrum Disorders: A Parents' Questionnaire Based Study. *Brain Sciences* **11**, 484 (2021).

274. Williams, K. L., Campi, E. & Baranek, G. T. Associations among sensory hyperresponsiveness, restricted and repetitive behaviors, and anxiety in autism: An integrated systematic review. *Research in Autism Spectrum Disorders* **83**, 101763 (2021).

275. Waizbard-Bartov, E. *et al.* Changes in the severity of autism symptom domains are related to mental health challenges during middle childhood. *Autism* **28**, 1216–1230 (2024).

276. Nwaordu, G. & Charlton, R. A. Repetitive Behaviours in Autistic and Non-Autistic Adults: Associations with Sensory Sensitivity and Impact on Self-Efficacy. *J Autism Dev Disord* (2023) doi:10.1007/s10803-023-06133-0.

277. Rudie, J. D. *et al.* Reduced functional integration and segregation of distributed neural systems underlying social and emotional information processing in autism spectrum disorders. *Cereb Cortex* **22**, 1025–1037 (2012).

278. Sigar, P., Uddin, L. Q. & Roy, D. Altered global modular organization of intrinsic functional connectivity in autism arises from atypical node-level processing. *Autism Res* **16**, 66–83 (2023).

279. You, X. *et al.* Atypical modulation of distant functional connectivity by cognitive state in children with Autism Spectrum Disorders. *Front Hum Neurosci* **7**, 482 (2013).

280. Harlalka, V., Bapi, R. S., Vinod, P. K. & Roy, D. Age, Disease, and Their Interaction Effects on Intrinsic Connectivity of Children and Adolescents in Autism Spectrum Disorder Using Functional Connectomics. *Brain Connect* **8**, 407–419 (2018).

281. Henry, T. R., Dichter, G. S. & Gates, K. Age and Gender Effects on Intrinsic Connectivity in Autism Using Functional Integration and Segregation. *Biol Psychiatry Cogn Neurosci Neuroimaging* **3**, 414–422 (2018).

282. Keown, C. L. *et al.* Network organization is globally atypical in autism: A graph theory study of intrinsic functional connectivity. *Biol Psychiatry Cogn Neurosci Neuroimaging* **2**, 66–75 (2017).

283. Passamonti, L. *et al.* Neuroinflammation and Functional Connectivity in Alzheimer's Disease: Interactive Influences on Cognitive Performance. *J Neurosci* **39**, 7218–7226 (2019).

284. Colasanti, A. *et al.* Hippocampal Neuroinflammation, Functional Connectivity, and Depressive Symptoms in Multiple Sclerosis. *Biol Psychiatry* **80**, 62–72 (2016).

285. Marsland, A. L. *et al.* Systemic inflammation and resting state connectivity of the default mode network. *Brain Behav Immun* **62**, 162–170 (2017).

286. Alvarez, G. M., Hackman, D. A., Miller, A. B. & Muscatell, K. A. Systemic inflammation is associated with differential neural reactivity and connectivity to affective images. *Social Cognitive and Affective Neuroscience* **15**, 1024–1033 (2020).

287. Wang, H. *et al.* Functional brain rewiring and altered cortical stability in ulcerative colitis. *Mol Psychiatry* **27**, 1792–1804 (2022).

288. Wei, Y. *et al.* Applying dimensional psychopathology: transdiagnostic prediction of executive cognition using brain connectivity and inflammatory biomarkers. *Psychological Medicine* **53**, 3557–3567 (2023).

289. Liu, Y. *et al.* Altered functional connectivity and topology structures in default mode network induced by inflammatory exposure in aged rat: A resting-state functional magnetic resonance imaging study. *Frontiers in Aging Neuroscience* **14**, (2022).

290. Harutyunyan, A. A., Harutyunyan, H. A. & Yenkoyan, K. B. Novel Probable Glance at Inflammatory Scenario Development in Autistic Pathology. *Front Psychiatry* **12**, 788779 (2021).

291. Pape, K., Tamouza, R., Leboyer, M. & Zipp, F. Immunoneuropsychiatry - novel perspectives on brain disorders. *Nat Rev Neurol* **15**, 317–328 (2019).

292. Miller, A. H., Haroon, E., Raison, C. L. & Felger, J. C. Cytokine Targets in the Brain: Impact on Neurotransmitters and Neurocircuits. *Depress Anxiety* **30**, 297–306 (2013).

293. Breitinger, U. & Breitinger, H.-G. Excitatory and inhibitory neuronal signaling in inflammatory and diabetic neuropathic pain. *Mol Med* **29**, 53 (2023).

294. Eren-Koçak, E. & Dalkara, T. Ion Channel Dysfunction and Neuroinflammation in Migraine and Depression. *Front. Pharmacol.* **12**, (2021).

295. Min, S. S. *et al.* Chronic brain inflammation impairs two forms of long-term potentiation in the rat hippocampal CA1 area. *Neurosci Lett* **456**, 20–24 (2009).

296. Wilson, J. F., Lodhia, V., Courtney, D. P., Kirk, I. J. & Hamm, J. P. Evidence of hyper-plasticity in adults with Autism Spectrum Disorder. *Research in Autism Spectrum Disorders* **43–44**, 40–52 (2017).

297. Rodriguez, J. I. & Kern, J. K. Evidence of microglial activation in autism and its possible role in brain underconnectivity. *Neuron Glia Biol* **7**, 205–213 (2011).

298. Skaper, S. D., Facci, L., Zusso, M. & Giusti, P. An Inflammation-Centric View of Neurological Disease: Beyond the Neuron. *Front Cell Neurosci* **12**, 72 (2018).

299. Deverman, B. E. & Patterson, P. H. Cytokines and CNS development. *Neuron* **64**, 61–78 (2009).

300. Di Marco, B., Bonaccorso, C. M., Aloisi, E., D'Antoni, S. & Catania, M. V. Neuro-Inflammatory Mechanisms in Developmental Disorders Associated with Intellectual Disability and Autism Spectrum Disorder: A Neuro-Immune Perspective. *CNS Neurol Disord Drug Targets* **15**, 448–463 (2016).

301. Wei, H. *et al.* IL-6 is increased in the cerebellum of autistic brain and alters neural cell adhesion, migration and synaptic formation. *Journal of Neuroinflammation* **8**, 52 (2011).

302. Thion, M. S. *et al.* Biphasic Impact of Prenatal Inflammation and Macrophage Depletion on the Wiring of Neocortical Inhibitory Circuits. *Cell Rep* **28**, 1119-1126.e4 (2019).

303. Thion, M. S. & Garel, S. On place and time: microglia in embryonic and perinatal brain development. *Curr Opin Neurobiol* **47**, 121–130 (2017).

304. Win-Shwe, T.-T. *et al.* Social behavior, neuroimmune markers and glutamic acid decarboxylase levels in a rat model of valproic acid-induced autism. *J Toxicol Sci* **43**, 631–643 (2018).

305. Mittli, D. Inflammatory processes in the prefrontal cortex induced by systemic immune challenge: Focusing on neurons. *Brain Behav Immun Health* **34**, 100703 (2023).

306. Filipello, F. *et al.* The Microglial Innate Immune Receptor TREM2 Is Required for Synapse Elimination and Normal Brain Connectivity. *Immunity* **48**, 979-991.e8 (2018).

307. Hunt, R. F., Boychuk, J. A. & Smith, B. N. Neural circuit mechanisms of post-traumatic epilepsy. *Front Cell Neurosci* **7**, 89 (2013).

A. Maternal Inflammatory Response Experimental Paradigm

Embryonic Day 9 (E9) — Birth (P0) — Young Adult (P60) — Old Adult (P200-400)

0.008 mg/kg LPS or Vehicle I.P. injection in pregnant dams

Brain microglia assessed

Experimental assessments of adult MIR offspring and the response to acute rapamycin treatment

Behavior, western blot, single cell & bulk sequencing, electrophysiology, MRI morphometry, rs-fMRI, microglia ablation, mTOR inhibition +/- RapaBlock, S6K inhibition, immunohistochemistry, blood cytokines, seizure susceptibility

B. Blood Cytokine Levels in Young and Old Adult Offspring

Fold Change MIR:CNTL

Young Adult / Old Adult

G-CSF, IFN-g, IL-6, IP-10, MCP-1, MIG, VEGF, KC, M-CSF, GM-CSF, IL-9, IL-1b, TNF-a, LIF

C. 12 hours after LPS Injection in Pregnant Dams

Fold Change LPS:VEH

G-CSF, IFN-g, IL-6, IP-10, MCP-1, MIG, VEGF, KC, M-CSF, GM-CSF, IL-9, IL-1b, TNF-a, LIF

D. Cortical IBA1+ Microglia: chronic increase in MIR offspring

% Control Offspring

E18, P0, P200, P400

E. pS6/S6 Western Blot: Flash-frozen Brain

Phospho-S6/S6 normalized protein levels

CNTL / MIR

F. Repetitive Behaviors Before and After Plex5622

% Total Time

CNTL, MIR, CNTL, MIR (+Plex) — Old Adult; CNTL, MIR, CNTL, MIR (+Plex) — Young Adult

G. 3 Chamber Ratio of Social Approach Before and After Plex5622

Ratio of interaction time (mouse : empty cup)

CNTL, MIR, CNTL, MIR (+Plex) — Old Adult; CNTL, MIR, CNTL, MIR (+Plex) — Young Adult

H. Von Frey Fiber Sensitivity Before and After Plex5622

Mechanical Threshold (g)

Young, +Plex, Old, +Plex — Control Offspring; Young, +Plex, Old, +Plex — MIR Offspring

Figure 1.   MIR offspring have a chronically increased inflammatory state and mTOR pathway activation and abnormal repetitive and social behaviors and sensory sensitivity in adult mice following a single, low-dose LPS injection early in gestation that is reduced by microglia ablation in young adult but not old adult offspring (**A**) Experimental timeline summary; (**B**) Inflammatory blood cytokines are significantly increased in both young and old adult MIR offspring relative to age matched controls (all p<0.001 MIR greater than control values) and are expressed as fold-change in MIR relative to controls (G-CSF=granulocyte colony stimulating factor, IFN-g=interferon gamma, IL-6=interleukin 6, IP-10=interferon gamma induced protein, MCP-1=monocyte chemoattractant, MIG= monokine induced by gamma protein interferon, VEGF=vascular endothelial growth factor, KC=keratinocyte chemoattractant, M-CSF=macrophage colony stimulating factor, GM-CSF=granulocyte macrophage colony stimulating factor, IL-9=interleukin 9, IL-1b=interleukin 1 beta, TNF-a=tumor necrosis factor alpha, LIF=leukemia inhibitory factor; all cytokines; (**C**) Blood cytokine changes in LPS injected pregnant dams (0.008 mg/kg) graphed as fold-change relative to vehicle-injected control pregnant dams at 12 hours post-injection, all cytokines (p<0.01); (**D**) Stereoinvestigator quantification of IBA+ immunohistochemistry staining of microglia in the sensory-motor cortex of adult MIR offspring, graphed as % control, all *p<0.05 (MIR v control values); (**E**) Phospho-S6 protein measured by western blot on microdissected, flash-frozen brain tissue from cortex, amygdala, and striatum (combined),*p=0.022;); (**F**) Repetitive behaviors that are significantly increased in young and old MIR mice compared to controls (*p<0.01) are significantly reduced after 3 week Plexxikon 5622 treatment to ablate brain microglia in the young (**p<0.05) but not the old adult MIR offspring; (**G**) Social approach (ratio of time spent interacting with mouse relative to empty cup) in the 3 chamber social preference test is significantly reduced in young and old adult MIR mice (*p<0.01) but is significantly increased after 3 week Plexxikon 5622 treatment in young (**p<0.05) but not old MIR offspring; (**H**) Sensitivity to touch/pressure on their hind paws in the Von Frey fiber sensitivity test is significantly lower in young and old adult MIR mice (*p<0.05) but is only increased toward control levels in young MIR mice after 3 week Plexxikon 5622 treatment (**p<0.05); all data mean ± SEM

A.

Brain Overgrowth: Birth to Old Adulthood

% Control brain wet weight normalized to body weight

P0 (*), P20 (**), P100 (***), P200 (****)

B.

Regional Volumetric Changes in Old Adult MIR Offspring Relative to Controls

MIR>Con   MIR<Con

C.

Change from Control (% of mean)

Cortex   Amygdala   Caudate   Hippocampus   Thalamus

Control MIR   Control MIR   Control MIR   Control MIR   Control MIR

Figure 2. **MIR offspring demonstrate growth patterns consistent with reports in humans with ASD of early brain overgrowth followed by undergrowth with cortical thinning and enlargement in other subcortical structures in adulthood.** (**A**) Brain wet weights normalized to body weight of MIR offspring relative to control offspring from birth to post-natal day 400 (P400) show early overgrowth (*,**,***,****$p<0.05$), (**B**) Diffusion weighted MRI regional volumetric brain analysis of adult offspring (P400) showing mean percent differences between MIR and Control mice (blue = decreased red = increased brain volume) and variable size changes based on brain region with significant ($p<0.001$) decreased cortical thickness (CTX), thalamus (Thal) and amygdala (Amg) volumes but significant ($p<0.05$) increases in lateral ventricle (LV), caudate (Cau), and hippocampus (Hipp) size, (**C**) Quantification of these regional deformation changes that were significant in MIR mice relative to controls in cortex, hippocampus, caudate, and amygdala; all data mean ± SEM

A. **Acute Rapamycin Treatment and Testing**

Adult Offspring — 5 mg/kg Rapa I.P.

2 hours — Measure: Repetitive Behaviors (grooming/circling), SOR Behaviors, Western Blots, Electrophysiology, fMRI, mRNA sequencing

Day 2 — 5 mg/kg Rapa I.P.

Day 3 — 5 mg/kg Rapa I.P.

Day 4 — 5 mg/kg Rapa I.P.

Measure: Social Approach

B. **Acute (2-hour) Rapamycin Treatment Decreases mTOR Pathway Activation In Vivo**

C. **3-Chamber Social Approach**

D. **Open-Field Repetitive Behaviors Increase with Age and Are Rescued by Acute Rapamycin**

E. **Effect Longevity of Single Rapamycin Dose on Repetitive Behaviors**

F. **Chronic Effect of Daily Rapamycin Treatment on Repetitive Behaviors Shows Habituation**

Figure 3. Acute (2-hour) rapamycin treatment rescues elevated brain mTOR signaling and repetitive and social behavior abnormalities in young and old adult MIR offspring. (**A**) Acute rapamycin treatment and testing paradigm; (**B**) Western blots showing that MIR mice have elevated mTOR pathway activation in adult offspring and that rapamycin treatment (Rapa) significantly reduces this elevated phospho-S6 protein levels in the primary somatosensory cortex (CTX), amygdala (AMG), and striatum (STR) at 2 hours post-injection (*,**,***$p<0.01$), demonstrating significant mTOR signaling pathway inhibition in this acute timeframe; (**C**) MIR mice have significantly reduced social approach (ratio of time spend with empty cup: stranger mouse, *,**,***$p<0.05$) and acute rapamycin treatment rescues the social approach deficits in the 3-chamber social preference test in adult offspring; (**D**) Young adult MIR offspring have significantly increased repetitive behaviors (grooming and circling) in an open-field test (*$p<0.01$) and old adult MIR offspring have an even greater increase in these repetitive behaviors (**$p<0.01$). Acute rapamycin treatment significantly reduces these repetitive behaviors in both young and old adult offspring (***,****$p<0.01$); (**E**) The longevity of the significant effect of a single dose of rapamycin (5mg/kg) on repetitive behavior abnormalities in MIR mice is approximately 72 hours with a significant effect still seen at 2 and 24 hours post-injection (*,**$p<0.01$); (**F**) Repeat daily doses of rapamycin (5mg/kg) show the development of a tolerance and gradual loss of efficacy on repetitive behavior abnormalities chronically at 5 weeks

Figure 4. **MIR mice display significant sensory over-responsivity (SOR), a common phenotype in ASD, which is rescued by acute rapamycin treatment.** (**A**) A novel tactile avoidance test was used to assess MIR and control offspring for aversion to rough tactile textures on the hind-limbs and fore-limbs using a light/dark box with varying light and floor textures. This demonstrates that MIR mice have normal exploratory behavior (trial 1), that they spend significantly less time in a chamber with rough flooring (Trial 2; *p<0.05), that they spend more time in a dark rather than light chamber when the flooring is the same, as do control mice (Trial 3; *p<0.05), but they avoid the normally preferred dark chamber when the flooring is a rough, aversive texture, unlike control mice (Trial 4; ***p<0.05); (**B**) Following acute rapamycin treatment young and old adult MIR mice no longer show a significant avoidance of the dark but rough textured floor chamber (Trial 4; *,**p<0.05) is significantly reduced by acute rapamycin treatment in young and old adult offspring; (**C**) MIR mice show increased sensitivity to touch/pressure on their hind paws in the Von Frey fiber sensitivity test that is rescued by acute rapamycin treatment in adult offspring; (**D**) MIR mice show a lack of habituation to a tactile startle (air puff blown on the back of the mouse at the beginning, middle, and end of the PPI test) that is restored by acute rapamycin treatment; (**E**) MIR mice show abnormal sensory gating in an air-puff pre-pulse inhibition (PPI) test of startle that is rescued by acute rapamycin treatment; (**F**) MIR mice also show a lack of habituation to an audible startle at the beginning, middle, and end of the PPI test that is rescued by acute rapamycin treatment.

A
Repetitive Behaviors: CNS Rapamycin Effects

B
3 Chamber Ratio of Social Approach: CNS Rapamycin Effects

C
Tactile Aversion Box (Trial 4): CNS Rapamycin Effects

D
Repetitive Behaviors after S6K Inhibitor Treatment

E
Relative Rescue Effect of Treatments on Repetitive Behaviors

Figure 5. **RapaBlock co-treatment with Rapamycin confirms the central nervous system action of rapamycin rescue of MIR offspring and S6K inhibition confirms that rescue of repetitive behaviors occurs via the mTOR signaling pathway.** (A) Repetitive behaviors, (B) Social approach, and (C) Tactile avoidance are significantly abnormal in MIR offspring compared to controls (*$p<0.01$) which are significantly improved following treatment with rapamycin (**$p<0.05$), and these improvements with rapamycin treatment are not significantly affected by simultaneous administration of RapaBlock, indicating that the peripheral nervous system effects of rapamycin are small relative to the contribution of central nervous system actions in MIR offspring; (D) Repetitive behaviors are significantly decreased 6 hours after treatment with the phospho-S6 kinase inhibitor PF4708671 confirming that downstream targets in the mTOR pathway may also produce some improvement in behavior (*$p<0.05$); (E) The relative rescue effects of rapamycin, Plexxikon 5622, and PF4708671 are compared demonstrating that the greatest effect on repetitive behaviors is from rapamycin.

A Spontaneous EPSCs in Cortical Pyramidal Neurons

B Firing Patterns of Cortical Pyramidal Neurons Following Injection of Depolarizing Current Pulses

C Spontaneous EPSCs in Cortical Pyramidal Neurons Are Reduced Following *Ex Vivo* Rapamycin Treatment

D Spontaneous EPSCs in Cortical Pyramidal Neurons Are Reduced Following *In Vivo* Rapamycin Treatment

Figure 6. **Cortical neurons in MIR offspring display abnormal electrophysiology indicating hyper-excitability which is rescued by acute rapamycin treatment.** (**A**) Traces and quantification of the frequency (*p<0.05), amplitude(**p<0.05), and area (***p<0.01) of spontaneous excitatory postsynaptic currents in cortical pyramidal neurons are all significantly increased in adult MIR mice compared to control offspring; (**B**) Firing Patterns of Cortical Pyramidal Neurons Following Injection of Depolarizing Current Pulses (500-1000 ms) show that layer 2/3 neurons from MIR mice have decreases in regular spiking slow and fast adaption neuron firing, increased doublet firing, and ectopic burst firing that is not present in the neurons from control mice; (**C**) Traces and quantification of the frequency (*p<0.01), amplitude (**p<0.05), and area (***p<0.05) of spontaneous excitatory postsynaptic currents in cortical pyramidal neurons from adult MIR mice demonstrate a significant rescue of neuronal hyper-excitability when treated with rapamycin *ex vivo* (2uM); (**D**) Traces and quantification of the frequency (*p<0.05), amplitude, and area of spontaneous excitatory postsynaptic currents in cortical pyramidal neurons from adult MIR mice treated acutely with rapamycin *in vivo* (5mg/kg, 2 hours prior to brain slice preparation) significantly rescues spiking frequency; The following abbreviations were used: Cm: membrane capacitance, pF: picofarads, Rm: membrane resistance, $M\Omega$: ohms, Tau: time constant, ms: milliseconds, RMP: resting membrane potential, mV: millivolts, pA: picoamperes. All data mean ± SEM.

A. Connectivity Strength Between Regions

B. Functional Connectivity: MIR v CNTL

D. Network Modularity

E. Thalamic Seed FC

F. Posterior Thalamus FC Strength

Figure 7. **Disordered functional connectivity seen in MIR mice are significantly rescued by acute rapamycin treatment.** (**A**) Fisher z scores (transformed r values, 0.2-0.6) indicating functional connectivity strength between cortical and subcortical brain regions in MIR and control mice before and after acute rapamycin treatment; (**B**) Regional increases in functional connectivity in resting state fMRI from MIR and control mice before and after acute rapamycin treatment (P<0.01 NBS, 2-tailed); (C) Regional decreases in functional connectivity in resting state fMRI from MIR and control mice before and after acute rapamycin treatment (P<0.01 NBS, 2-tailed); (D) Differential network modularity organization across the brain is increased in MIR mice compared to control offspring and is rescued by acute rapamycin treatment; (E) Differential functional connectivity by brain region from a left posterior thalamus seed in MIR mice compared to control offspring before and after acute rapamycin treatment; (F) Voxel-wise quantification of posterior thalamus functional connectivity is significantly increased in MIR mice compared to control offspring and is differentially affected by acute rapamycin treatment with cortical regions showing decreased thalamic FC and subcortical regions showing increased thalamic FC following treatment. Abbreviations used as follows: **acc**=anterior cingulate cortex; **rsc**=retrosplenial cortex; **sc**=subicular complex; **v2c**=secondary visual cortex; **s1s**=primary somatosensory cortex; **s2s**=secondary somatosensory cortex; **s1b**=S1 barrel cortex; **s1a**=primary auditory cortex; **dthal**=dorsal thalamus; **vthal**=ventral thalamus; **pthal**=posterior thalamus; **pvthal**=paraventricular thalamus; **mdthal**=medial dorsal thalamus; **rthal**=reticular thalamus; **lpthal**=lateral posterior thalamus; **tac**=temporal association cortex; **stm**=stria terminalis; **mgen**=medial geniculate; **lgen**=lateral geniculate; **str**=striatum; **cmthal**=central median thalamus; **clau**=claustrum; **gp**=globus pallidus; **sn**=substantia nigra; **amg** =amygdala; **lamg**=lateral amygdala; **hypo**=hypothalamus; **vpal**=ventral pallidum; **entc**=entorhinal cortex; **insc**=insular cortex; **s1f** = primary face somatosensory cortex; **s1fl**=primary forelimb somatosensory cortex; **s1hl**=primary hindlimb somatosensory cortex; **s1t**=primary trunk somatosensory cortex; **m1**=primary motor cortex; **m2**=secondary motor cortex; **ot**=olfactory tubercles; **aamg**=anterior amygdala; **camg**=central amygdala; **bamg**=basal amygdala; **mamg** =medial amygdala; **pfc**=piriform cortex; **epn**=endopiriform nucleus; **s2c**=secondary somatosensory cortex; **supc**=superior colliculus; **fx**=hippocampal fornix; fim=hippocampal fimbria; **cgm**=cingulum; **hab**=habenular; **dg**=dentate gyrus; **ca3**=hippocampus CA3; **paqu**=periaqueductal gray; **rtf**=reticular formation; **lsep**= lateral septum

Figure 8. **Cell sequencing from sensorimotor cortex microdissections from MIR and control adult offspring.** Clustering of single-cell RNA seq data from sensorimotor cortex microdissections as volcano plots of differentially expressed genes in MIR offspring compared to Controls in (**A**) upper layer and (**B**) deep layer excitatory neurons and in (**C**) parvalbumin (PV) inhibitory interneurons; (**D**) The number of genes enriched in MIR mice by cell type; (**E**) Relative enrichment (fold change) of mTOR pathway regulated genes significantly enriched in MIR mice by cell type indicates that MIR mice have an increase in these genes in microglia, neurogranin neurons, PV+ interneurons, oligodendrocytes, oligodendrocyte precursor cells (OPC), endothelial cells, upper layer excitatory neurons, and astrocytes compared to control offspring.

	Cm(pF)	RM(MΩ)	Tau(ms)	RMP(mV)	Rheobase(pA)
Control	208.8 ± 21	109.2 ± 16	4.0 ± 0.4	-78.1 ± 2.4	725.6 ± 98
MIR	207.8 ± 82	129.0 ± 16	3.6 ± 0.3	**-70.6 ± 1.3***	**499.2 ± 52****
MIR+Rapa	214.5 ± 8	106.9 ± 8	3.7 ± 0.16	73.3 ± 1.4	590.6 ± 61

Table 1. Passive Membrane Properties of Cortical Pyramidal Neurons. Passive membrane properties (membrane capacitance, input resistance, time constant, and resting membrane potential (RMP)), and Rheobase) of cortical pyramidal neurons show MIR mice have significantly higher resting membrane potential (*$p<0.05$, closer to action potential threshold) and decreased Rheobase (**$p<0.05$, lower minimum current to depolarize) indicating neuronal hyper-excitability in the cortex and increased membrane capacitance in the striatum compared to controls

Maximum Seizure Score	PTZ dosage (mg/kg)									
	10		20		30		40		40 + Rapamycin	
	CNTL	MIR	CNTL	MIR	CNTL	MIR	CNTL	MIR	CNTL	MIR
0	8/8	8/8	8/8	7/8	7/8	4/8	6/8	0/8	7/8	0/8
1	0/8	0/8	0/8	1/8	1/8	3/8	2/8	1/8	1/8	3/8
2	0/8	0/8	0/8	0/8	0/8	1/8	0/8	2/8	0/8	4/8
3	0/8	0/8	0/8	0/8	0/8	0/8	0/8	3/8	0/8	1/8
4	0/8	0/8	0/8	0/8	0/8	0/8	0/8	2/8	0/8	0/8
5	0/8	0/8	0/8	0/8	0/8	0/8	0/8	0/8	0/8	0/8
6	0/8	0/8	0/8	0/8	0/8	0/8	0/8	0/8	0/8	0/8

Table 2. Seizure Susceptibility by PTZ Dosage. Seizure scores during pentylenetetrazol (PTZ) dose escalation indicates a greater severity and larger number of mice per group displaying a seizure behavior in MIR mice compared to Control offspring that is decreased by acute rapamycin treatment.

Cell Type	ASD-associated Genes		Epilepsy/Seizure-associated Genes		Neuroinflammation-associated Genes	
	Up MIR	Down MIR	Up MIR	Down MIR	Up MIR	Down MIR
Excitatory Deep Neurons	DBI HMGA2, CCND3, MAFB	NPAS4, MYOM2, EGR4, KIRREL*	EGR4, DBI, MAFB	NPAS4		
Excitatory Upper Neurons	FKBP5, CCND3, CD7	NPAS4, MYOM2, EGR4, COL11A1	FKBP5, EGR4	NPAS4, ACTN2		
Other Excitatory Neurons	TBR1	ACTN2, MCM6, MRC2, DISP1, NOSTRIN, DISP1, KCTD18, KCNMB2, TTC28, NKAIN3, GPC6, CDK6, GRIP1*, EGFR, FILIP1, ST8SIA2	TBR1, DISP1	MCM6, DISP1, KCNMB2, NKAIN3, GRIP1*, ST8SIA2, EGFR, FILIP1		
Interneurons	WIPF1, SLC15A2, RPL4	Exd1, SLC17A8	WIPF1			SLC17A8
Microglia	FKBP5, E2F1, MCM6, C4B, PIK3C2G, FAM111A, DIAPH3, CCl12, HSPD1, TUBB5	SEMA4B, SLC44A2	FKBP5, FAM111A, DIAPH3	SEMA4B	XDH, CCL12, C4B, TUBB5	
Astrocytes	IRS2					
Oligodendrocytes	HIF3A, FKBP5, C4B, CDH6, FGF1, DUSP1, BCAT1, KCNJ3	LYPD6, SLPI	DUSP1,KCNMA1		PHYHD1, C4B, MAT2a	SLPI
OPCs	HIF3A, FKBP5, NPAS1, HSP90AA1		FKBP5, ARF6			

Table 3. Single nucleus sequencing analysis of significantly differentially expressed genes in MIR mice compared to control offspring are shown for Autism-associated genes, Epilepsy/Seizure-associated genes, and neuro-inflammation-associated genes; All data use (LogFC >1, P_val_adj <0.05, FDR<0.1) for abundance and significance criteria. *Sensory over-responsivity and repetitive behavior associated genes

Differentially Expressed Genes MIR vs CNTL Offspring													
Transcription Factors		GABAergic Genes		Glutamatergic Genes		Ion Channels		Potassium Channels		Sodium Channels		Calcium Channels	
Up MIR	Down MIR	Up MIR	Down MIR	Up MIR	Down MIR	Up MIR	Down MIR	Up MIR	Down MIR	Up MIR	Down MIR	Up MIR	Down MIR
Runx2 Nkx1-2 E2f8 Trpc6 Tcf19	Nkx2-1 Nkx2-2 Myt1 Helt Eya2 Asic4 Tfap20 Tfap2a		Gabre Gabrq		Grik1 Grik2 Grin2c Grid2 Grm1 Grm4	Slc22a3 Slc22a4 Asic4 Clcnka Slco2b1 Trpc6	Trpc6 Clic6 Trpc7 Trpv1 Trpm3 Slc35d3	Kcnh3 Kcnv1	Kcne1l Kcnj5 Kcnab1 Kcnk10 Kcnj13		Slc4a5 Slc10a4 Slc8a3 Scn5a Scnn1g	Cacnb3	Cacng5 Cacna2d2

Table 4. Bulk sequencing analysis of significantly differentially expressed genes in MIR mice compared to control offspring are shown for transcription factors, GABAergic genes, Glutamatergic genes, ion channel genes, potassium ion genes, sodium ion genes, and calcium ion genes which may affect neuronal excitability; All data use (LogFC >1, P<0.05, FDR<0.1) for abundance and significance criteria.

ASD Associated Genes					
MIR vs CNTL Offspring		CNTL Rapa vs CNTL Veh		MIR Rapa vs MIR Veh	
UP in MIR	**Down in MIR**	**UP in CR**	**Down in CR**	**Up in MR**	**Down in MR**
TEK (Tie2)	ACE	CDH22	TEK (Tie2)	PCDHA7	TEK (Tie2)
PLAUR (CD87)	CNTNAP3	PLAUR (CD87)	NTNG1	TEKT4	PLAUR (CD87)
ID1	MFRP	PPP1R1B	MFRP		ID1
HOMER1	PCDH11X	ADA	CSMD3		SIK1
ASPM	CHD7	MEGF11	GPR37		CDKN1A (p21)
CDKN1A (p21)	TEKT4		PCDH15		
IQGAP3	NEUROD1		DCC		
KIF14	GRIN2C		KHDRBS2		
MAST3	SLC35D3				

Table 5. (F) Bulk sequencing analysis of autism-associated genes that are significantly enriched in MIR mice relative to control offspring; All data use (LogFC >1, P<0.05, FDR<0.1) for abundance and significance criteria.

MIR v Control DEGs Reversed by Rapamycin	
Decreased by Rapa	**Increased by Rapa**
TEK (Tie 2)	HIF3A
PLAUR (uPAR, CD87)	PLIN4
NES	TEKT4
GEM	CHD7
ID1	CNTNAP3
TIPARP	
APOLD1	
CDKN1A (p21)	

Table 6. (H) Bulk sequencing analysis of differentially expressed genes that were enriched in MIR offspring relative to control offspring that are rescued (reversed) by acute rapamycin treatment. All data use (LogFC >1, P<0.05, FDR<0.1) for abundance and significance criteria.

A.

Negligible Effect of 2 hour Rapamycin Treatment on Blood Cytokines

B. EPSCs from Cortical Pyramidal Neurons from MIR Mice under Voltage Clamp and Current Clamp Conditions

C. Current Evoked Action Potential Bursts in Cortical Pyramidal Neurons

Supplemental Figure 1. Acute Rapamycin Doesn't Decrease Inflammatory Cytokines in MIR Mice and Cortical Somatosensory Pyramidal Neurons Display Multiple Signs of Hyper-Excitability in MIR Brains

(A) Blood cytokines are unchanged by 2-hour Rapamycin treatment in adult MIR mice, see Figure 1 legend for cytokine abbreviations and full names; (**B**) Excitatory post-synaptic current activity in cortical pyramidal neurons under voltage clamp and current clamp conditions. The arrow indicates a putative NMDA spike, which also suggests CPN hyper-excitability in MIR mice; (**C**) Action potential bursting from cortical pyramidal neurons under current clamp conditions.

A.

Spontaneous EPSCs in Striatal MS Neurons

B.

Spontaneous EPSCs in Striatal Medium Spiny neurons Following *Ex Vivo* Rapamycin Treatment

C.

Striatal Medium Spiny Neuron EPSCs in Bicuculline

D. Striatal Medium Spiny Neuron EPSCs During GABAA Receptor Inhibition by Bicuculline With and Without Rapamycin in vitro

Supplemental Figure 2. Electrophysiology of Medium Spiny Neurons in the Striatum Show Trends Towards Hyper-excitability that is Reduced by Acute Rapamycin Treatment. (A) Traces and quantification of the frequency, amplitude, and area of spontaneous excitatory postsynaptic currents in striatal medium spiny neurons from adult MIR and control offspring show a trend for increases in MIR mice compared to control but do not reach significance; (B) Traces and quantification of the frequency, amplitude, and area of spontaneous excitatory postsynaptic currents in striatal medium spiny neurons from adult MIR mice treated acutely with rapamycin ex vivo (2uM) compared to controls shows a trend for decreased spiking frequency that does not reach significance; (C) Traces showing treatment with Bicuculline, a competitive antagonist of GABA A receptors, increases excitatory post-synaptic currents in medium spiny neurons in the striatum, (D) Traces showing acute *ex vivo* rapamycin treatment given simultaneously with Bicuculline prevents the increase in medium spiny neurons EPSCs.

A rs-fMRI FC (Control v Control+Rapa)

B

EGR1+/NeuN+ Active Neurons in Brain Regions During Open Field Behavior

Supplemental Figure 3. Functional Connectivity Changes in Resting State fMRI in Control Offspring and Immunohistochemistry Validation of Increased Neuron Activity Indicated by fMRI Results. (A) The changes in FC in control mice after acute rapamycin treatment were small compared to MIR mice (Fig. 6C) and were mostly decreases (blue lines) between cortical and subcortical regions with few increases (red lines) between retrosplenial cortex and basal ganglia structures; (B) Stereological quantification of immediate early gene protein EGR1 in neurons (NeuN+) were quantified using stereological counting indicating increased neuronal activity in MIR mice in the motor cortex, somatosensory cortex, retrosplenial cortex, and the striatum and decreased activity in the amygdala in agreement with functional MRI findings.

A. SnRNA-Seq GSEA: Pathway All Cells

B. SnRNA-Seq GSEA: Pathway All

C. SnRNA-Seq GSEA: Pathway Excitatory Neurons

D. SnRNA-Seq GSEA: Pathway Inhibitory Neurons

E. Bulk Seq GSEA: Pathway MIR vs CNTL Offspring

F. Bulk Seq GSEA: Geneontology biological processes MIR vs CNTL Offspring

G. Bulk Seq GSEA: Pathway MIR + Rapa vs MIR Offspring

H. Bulk Seq GSEA: Geneontology biological processes MIR + Rapa vs MIR Offspring

Supplemental Figure 4. Gene Set Enrichment Analysis of snRNA seq and Bulk sequencing of Genes that Are Significantly Differentially Expressed in MIR Offspring.
(A) Gene set enrichment analysis (GSEA) of pseudo-bulk expression from single-nucleus RNA sequencing in cellular pathways for all cell types in MIR mice relative to control offspring; (B) GSEA of pathways enriched in MIR mice in all neurons; (C) GSEA of pathways enriched in MIR mice in excitatory neurons; (D) GSEA of pathways enriched in MIR mice in inhibitory neurons; (E) Gene set enrichment analysis (GSEA) of pathways enriched in MIR mice compared to control offspring from bulk sequencing; (F) Geneontology of biological processes enriched in MIR mice compared to control offspring; (G) GSEA of pathways differentially expressed in MIR mice treated with rapamycin compared to vehicle treated MIR mice ; (H) Geneontology of biological processes differentially expressed in MIR mice treated with rapamycin compared to vehicle-treated MIR mice; expression and significance criteria used for all are LogFC>1, P≤0.05, FDR ≤0.1

	Rank	logFC_mir_vs_ctrl	pVal_mir_vs_ctrl	FDR_pVal_mir_vs_ctrl
Microglia_1	6	0.024198538	0.730088556	0.988144314
Microglia_2	15	0.001630415	0.981174139	0.988144314
Endothelial	13	0.007223573	0.898156918	0.988144314
Pericyte	14	-0.036032657	0.575069239	0.988144314
Unclassified	12	0.066055026	0.406867978	0.988144314
Oligodendrocyte	3	-0.045686542	0.658849051	0.988144314
Astrocyte_1	4	-0.053901143	0.656765516	0.988144314
Astrocyte_2	17	0.01232224	0.848949092	0.988144314
OPC_1	8	-0.011208874	0.86612819	0.988144314
OPC_2	16	-0.000997818	0.988144314	0.988144314
Interneuron-SST	7	-0.012737802	0.870655188	0.988144314
Interneuron-PV	5	0.057783538	0.479172194	0.988144314
Interneuron-VIP	10	0.125438347	0.120224361	0.841010574
Interneuron-SV2C	11	0.069811105	0.365002986	0.988144314
Neurogranin	9	0.362809282	0.148413631	0.841010574
Excitatory-deep	2	-0.497502762	0.000931602	0.01583724
Excitatory-upper	1	0.114508117	0.457762268	0.988144314

Supplemental Table 1. Single nucleus sequencing comparison of cell populations between MIR and Control offspring sensory-motor cortex indicates that there are no significant differences between treatment groups for any cell type.