Endogenous FGFs drive ERK-dependent cell fate patterning in 2D human gastruloids

Kyoung Jo[1]*, Zong-Yuan Liu[1]*, Gauri Patel[4], Zhiyuan Yu[2], LiAng Yao[1], Seth Teague[4], Craig Johnson[1], Jason Spence[1,3-5], Idse Heemskerk[1-4,6^]

1. Department of Cell and Developmental Biology, University of Michigan Medical School, Ann Arbor, Michigan
2. Department of Computational Medicine and Bioinformatics, University of Michigan Medical School, Ann Arbor, Michigan
3. Center for Cell Plasticity and Organ Design, University of Michigan Medical School, Ann Arbor, Michigan
4. Department of Biomedical Engineering, University of Michigan, Ann Arbor, Michigan
5. Department of Internal Medicine, Gastroenterology, University of Michigan Medical School, Ann Arbor, Michigan
6. Department of Physics, University of Michigan, Ann Arbor, Michigan

* equal contributions
^ for correspondence: iheemske@umich.edu

Abstract

The role of FGF is the least understood of the morphogens driving mammalian gastrulation. Here we investigated the function of FGF in a stem cell model for human gastrulation known as a 2D gastruloid. We found a ring of FGF-dependent ERK activity that closely follows the emergence of primitive streak (PS)-like cells but expands further inward. We showed that this ERK activity pattern is required for PS-like differentiation and that loss of PS-like cells upon FGF receptor inhibition can be rescued by directly activating ERK. We further demonstrated that the ERK-ring depends on localized activation of basolaterally positioned FGF receptors (FGFR) by endogenous FGF gradients. We confirmed and extended previous studies in analyzing expression of FGF pathway components, showing FGFR1 is the main receptor, FGF2 is highly expressed across several cell types, and FGF4/17 are the main FGF ligands expressed in the PS-like cells, similar to the human and monkey embryo but different from the mouse. We found that knockdown of FGF4 greatly reduced PS-like differentiation while FGF17 knockdown primarily affected subsequent mesoderm differentiation. FGF8 expression was spatially displaced from PS-markers and FGF4 expression and peaked earlier, while knockdown led to an expansion in PS-like cells, suggesting FGF8 may counteract FGF4 to limit PS-like differentiation. Thus, we have identified a previously unknown role for FGF-dependent ERK signaling in 2D gastruloids and possibly the human embryo, driven by a mechanism where FGF4 and FGF17 signal through basally localized FGFR1 to induce PS-like cells and their derivatives, potentially restricted by FGF8.

Introduction

During gastrulation the three germ layers are established, and the body plan is laid out. In amniotes this involves the formation of the primitive streak on the posterior side of the embryo from which cells ingress beneath the epiblast and migrate to form the mesodermal and endodermal layers. An important role in gastrulation is played by evolutionarily conserved morphogens from the BMP, Wnt, Nodal, and FGF families[1]. The function of the first three signaling pathways is comparatively well understood: in both mouse and human a transcriptional hierarchy between BMP4, Wnt3, and Nodal plays a central role in inducing dynamic signaling gradients along the anterior posterior axis that control cell fate patterning[1–3]. For BMP and Nodal, basal receptor localization in the epiblast is crucial to form signaling activity gradients[3–7]. Wnt and Nodal signaling are directly required for mesoderm and endoderm differentiation, while the balance between BMP and Nodal signaling is important for patterning the germ layers along the body axes[8].

In contrast, the role of FGF in mammalian and especially human gastrulation has been more elusive. Secreted FGFs signal through FGF receptor tyrosine kinases (FGFRs) in a heparan sulfate proteoglycan (HSPG)-dependent manner to activate several signaling pathways including MAPK/ERK, PI3K, and PLCγ[9]. There is evidence for a conserved requirement for FGF/ERK signaling in mesoderm formation across vertebrate model organisms[10–13]. However, it is unknown if there are FGF- and phosphorylated ERK gradients across the mammalian primitive streak[14,15], whether there are functionally relevant ERK signaling dynamics like in other contexts, or whether receptor localization is important for establishing signaling gradients[16–19]. It is also unclear to what extent the requirement in mesoderm induction is direct, or indirect by modulating other signals[14,15,20]. In addition, several FGFs are co-expressed and the roles of specific FGFs are uncertain. Moreover, there may be significant interspecies differences in this respect. Notably, FGF8 is the only FGF which has been shown to be required for gastrulation in the mouse[21], and its requirement in gastrulation appears to be conserved across vertebrate model organisms[22–24], yet FGF8 is not highly expressed in primate embryos[25,26].

Mutants for FGF8, FGFR1, HSPG synthesis, and ERK2, as well as pharmacological inhibition of MEK and FGFR give rise to similar phenotypes in mouse gastrulation, suggesting that FGFR1 and FGF8 are the only FGF ligand-receptor pair required for gastrulation and that these act through the MAPK/ERK pathway[12,20,21,27–31]. FGF4 expression during gastrulation is also lost in FGF8 mutants, leaving the possibility of a role for FGF4, but this

remains unknown as FGF4 is also required in the blastocyst and mutants therefore do not reach gastrulation[32]. In
all mutants and for pharmacological inhibition of FGF/ERK signaling, expression of the primitive streak marker
brachyury (TBXT, also BRA, or T in mouse) is lost or severely diminished in the posterior streak and mesoderm
fails to migrate away, piling up in this region. In addition, TBX6 expression is lost completely, and no lateral or
paraxial mesoderm are formed. In contrast, TBXT expression in the anterior streak and extraembryonic mesoderm
is not dependent on FGF/ERK signaling; extraembryonic mesoderm development is normal, and the axial
mesoderm is expanded[20,21,27,28].
In vitro models offer an opportunity to disentangle the complexity of FGF signaling in human development. Here,
we explore the role of FGF/ERK signaling in micropatterned human pluripotent stem cells treated with BMP4, also
known as 2D human gastruloids. These give rise to concentric rings of different cell types associated with
gastrulation after 2 days of differentiation. Pluripotent cells in the center are surrounded by a primitive streak-like
ring, which itself is surrounded by primordial germ cells and amnion-like cells on the colony edge[33–35]. Due to its
simplicity and reproducibility, this system has proven powerful in deciphering the mechanisms underlying aspects
of human gastrulation. It was used to show that BMP-Wnt-Nodal hierarchy that controls primitive streak formation
during mouse gastrulation is preserved in human while providing a range of new insights into how these signals
function[2,3]. For example, the crucial role for BMP and Nodal receptor localization in patterning was discovered in
2D gastruloids and later confirmed in the mouse embryo. This system also revealed that BMP, Wnt, and Nodal
signaling patterns are highly dynamic[3–7]. It therefore also provides a promising approach to determine how FGF
may function in human gastrulation.
In the 2D human gastruloid model, we show that a ring of phosphorylated ERK forms in an FGF-dependent
manner and expands largely in lockstep with the formation of primitive streak-like cells but extends further inward.
We demonstrate that this pattern of ERK activity is due to localized FGF receptor (FGFR) activation that requires
basal FGFR localization. We then show that exogenous FGF2 in the media is required for pattern formation to occur
but can be removed well before the phosphorylated (active) ERK (pERK) and PS-like rings appear, suggesting the
pERK ring instead depends on endogenous FGF gradients. Using single cell RNA-sequencing, we determine the
expression of FGF pathway components including FGFR1 as the main receptor and FGF2, FGF4, and FGF17 as
the most highly expressed ligands. Our analysis corroborates previous studies describing expression of FGF2 and
FGF17 in human embryos and gastruloids but for the first time recognizes a possible role for FGF4 and includes a
broad analysis of genes involved in FGF signaling modulation[25,36]. We find that the FGF4 and FGF17 expression
are restricted to the PS-like ring and that knockdown of FGF4 significantly reduces PS-like differentiation, while
FGF17 knockdown primarily affects downstream mesoderm differentiation. Given its importance in the mouse, we
also analyze FGF8 and find its expression is spatially and temporally displaced from FGF4 and PS-markers, while
FGF8 knockdown leads to an increase in PS-like differentiation, suggesting it may counteract FGF4. In summary,
we have greatly advanced our understanding of how FGFs function in a model for human gastrulation.
95
96   **Results**
97
FGF/ERK signaling is required for primitive streak-like differentiation.
We previously observed that pharmacological inhibition of MEK (the kinase upstream of ERK) or FGF receptors
leads to a loss of differentiation to primitive streak-like and primordial germ cell-like cells, suggesting a crucial role
for FGF/ERK signaling[34]. Moreover, a ring of active ERK (pERK) has been observed in 2D gastruloids but its
connection to FGF signaling or differentiation remains unclear[33]. Therefore, we wanted to investigate the role of
FGF/ERK signaling further.
To determine the dynamics of pERK and relate it to PS-like differentiation, we first performed time series
immunofluorescence co-staining for pERK and the PS marker TBXT after BMP4 treatment in 700um diameter
colonies. This revealed the pERK ring first emerges around 24h, around the same time as TBXT expression, and
gradually gets brighter and wider (Fig.1a). By 42h the outer edge of the ERK ring coincided with TBXT expression,
while the inner edge was less sharp and extended further towards the center than TBXT, reminiscent of the domains
of WNT and NODAL signaling that extend further in than the PS-like region[3,6] (Fig.1a). Cell fate patterning in 2D
gastruloids has a fixed length scale from the edge, with the central part absent in smaller colonies due to the fixed
length scale of the BMP signaling gradient from the edge. Consistently, we found that ERK signaling also has a
fixed pattern from the edge regardless of colony size, so that in smaller colonies the ERK ring becomes a domain
of high ERK signaling extending through the colony center (Supp. Fig.1a). MEK or FGFR inhibition for 30 minutes
after 41.5h eliminated ERK activity, suggesting ERK activity depends entirely on FGF and that continuous FGF
signaling is necessary to maintain the ERK signaling pattern (Fig.1bc). Inhibition for the full duration of differentiation
eliminated TBXT expression as well as other PS markers (Fig.1bc, Supp. Fig.1b). However, we were able to rescue
TBXT expression by directly and uniformly activating ERK (Fig. 1d-f, Supp. Fig.1cd) in cells stably expressing the
doxycycline-inducible membrane-targeted catalytic domain of SOS (dox-SOS[cat])[37]. Despite uniform induced ERK

activity, TBXT expression remained excluded from the center, showing that ERK activation alone is not sufficient
for TBXT induction. However, TBXT levels were much higher in the rescue (Supp. Fig. 1cd), possibly reflecting the
much higher level of ERK activity induced by dox-SOS^{cat}. MEK and FGFR inhibition also led to a reduction in cell
number which staining for BrdU and cleaved Cas3 suggested was primarily due to reduced proliferation (Supp. Fig.
1ef). In contrast to TBXT expression, reduced cell number caused by FGFR inhibition was only partially rescued by
dox-SOS^{cat}, indicating additional downstream pathways regulate cell number (Fig. 1e). Altogether, these data show
the formation of a PS-like ring is dependent on FGF signaling through ERK.
This raised several questions. First, is FGF/ERK signaling required directly for PS-like differentiation, or
does it act indirectly? These possibilities are not mutually exclusive. For example, FGF/ERK could be required
directly but also act indirectly by controlling Wnt or Nodal expression, as both Wnt and Nodal signaling are required
for PS-like differentiation. Second, which are the FGFs activating the ERK pathway? The pathway could be
activated by exogenous FGF2 present in the pluripotency medium mTeSR1 or by endogenously expressed ligands.
Third, how is ERK activity restricted to a ring?
To determine if there is a direct requirement of FGF/ERK signaling and whether this depends on exogenous
FGF, we performed differentiation to PS-like cells in standard culture by exogenous activation of Wnt and Nodal
signaling, bypassing the need to form endogenous Wnt and Nodal gradient as in 2D gastruloids[38]. We used a
minimal base medium (E6) without exogenous FGF or other growth factors. To rule out any additional requirement
for endogenous Nodal and Wnt which may be FGF/ERK-dependent, we used Nodal knockout cells[3] treated with
the Wnt secretion inhibitor IWP2, thereby eliminating any endogenous activity of these pathways. We also inhibited
BMP receptors using LDN193189 to rule out any indirect effect through endogenous BMP. Under these conditions,
with exogenous Wnt and Nodal activation for 24h in the absence of exogenous FGF, cells efficiently differentiated
to TBXT+ PS-like cells (Fig.1g, Supp. Fig.1g). These PS-like cells had high active ERK levels just as in the 2D
gastruloids. Inhibition of either FGFR or MEK abrogated ERK activity and TBXT expression while FGFR inhibition
also severely reduced proliferation. In both cases significant expression of the pluripotency and ectoderm marker
SOX2 was maintained (Supp. Fig. 1g). Given that FGF/ERK is known to be involved in pluripotency maintenance,
this suggests inhibition may not be complete and that much lower ERK activity is required for pluripotency
maintenance than PS-like differentiation, consistent with lower ERK activity in the pluripotent center of gastruloids.
MEK and FGFR inhibition reduce proliferation and thereby cell density, raising the possibility that their lack
of differentiation is an indirect effect that depends on cell density: a critical density may be required for differentiation
due to a community effect. To rule this out we performed the experiment across a range of densities, which
confirmed that PS differentiation in the control was efficient even at the lowest densities while FGFR and MEK
inhibition effectively blocked differentiation when compared to controls with similar final cell densities (Supp.Fig.1h).
We also tested whether proliferation and differentiation require similar levels of ERK signaling by determining the
MEKi dose response of gastruloids. This revealed that lower doses of MEKi effectively block differentiation without
strongly affecting cell number, further supporting that the effects of ERK signaling on proliferation and differentiation
are independent (Supp.Fig.1i). Altogether, our results show that endogenous FGF expression is both necessary
and sufficient for PS-like differentiation by exogenous Wnt and Nodal stimulation and demonstrate a direct
requirement for ERK signaling in PS-like differentiation.
(Next page)
Figure 1. a) Immunofluorescence stainings for pERK and TBXT and their radial intensity profiles in 2D gastruloids at different
times. Error bars in graphs are standard deviations over N=4 colonies. **b,c)** Effect FGF receptor inhibition (FGFRi, b) or MEK
inhibition (MEKi, c) on pERK and TBXT in colonies fixed at 42h after BMP treatment. Inhibition for either 30 minutes at 41.5h
(rounded to 42 in figure) or throughout differentiation. **d)** BMP only control and FGFRi inhibition in cells expressing doxycycline-
inducible SOS^{cat} with and without addition of doxycycline. **e-f)** Cell numbers (e) and radial profiles of TBXT positive cells (f) for
conditions in (d). Error bars represent standard deviation. Statistical significance was assessed using one-way ANOVA followed
by Tukey's HSD test for pairwise comparisons. **g)** PS marker TBXT, pERK and pluripotency marker SOX2 in Nodal knockout
cells with inhibitors of Wnt secretion (WntSeci) and BMP receptors (BMPRi) with or without MEKi or FGFRi. All scale bars 50um.

a

| | 16h | 24h | 28h | 32h | 36h | 40h | 44h | 48h |

pERK

TBXT

overlay + DAPI

intensity (a.u.) — edge distance (μm)

Phospho-ERK depends on FGF and is required for formation of a primitive streak-like ring in 2D gastruloids

FGFR inhibition

b ctrl / 30 min @ 42h / continuous — TBXT — pERK — overlay + DAPI

MEK inhibition

c ctrl / 30 min @ 42h / continuous — TBXT — pERK — overlay + DAPI

Rescue of FGFR inhibition by direct ERK activation with dox-SOS^{cat}

d ctrl / FGFRi - dox / FGFRi + dox — TBXT — pERK — overlay + DAPI

e ×10^3 cell number — cells per colony — BMP4, +FGFRi -dox, +dox — *** ** n.s.

f TBXT — positive fraction — edge distance (μm) — BMP4, +FGFRi, +FGFRi +dox

There is a *direct* requirement for FGF/ERK signaling in primitive streak-like differentiation

g

| | DAPI | TBXT | pERK | SOX2 | overlay + DAPI |

NodalKO + BMPRi + WNTSeci + CHIR + Activin

+ MEKi

+ FGFRi

Figure 2. a) Schematic of the FGF/ERK signaling pathway. **b)** Distribution of pERK levels in TBXT+ cells versus ISL1+ and other cells based on data and thresholds in Supp. Fig. 2b. **c)** Expression of key markers in PHATE[39] projection of scRNA-seq data for 42h colony. **d)** Cell type annotation. PS-LC: primitive streak-like cells, NasMe: nascent mesoderm, AE-LC: amniotic ectoderm-like cells: PGC-LC: primordial germ cell-like cells, endo: definitive endoderm. **e)** Expression of canonical FGF ligands (log transformed and smoothened). **f)** Comparison of FGF ligand expression in 2D human gastruloids to human, monkey, and mouse embryo, normalized within each type of sample (see methods). **g-i)** Expression of several genes involved in FGF signaling with strong differential expression between PS-like cells or nascent mesoderm and pluripotent cells, split by gene category.

Single cell RNA-sequence reveals differentially expressed FGF pathway components.

The above results suggest endogenous FGF expression may contribute to elevated ERK signaling in the PS-like ring. However, FGF/ERK signaling may be modulated at many levels (Fig. 2a). A spatial pattern of ERK activation could reflect a concentration gradient of FGF. Alternatively, differential expression of receptors and HSPG-modifying enzymes could lead to spatially patterned receptor activation and downstream ERK activity even in the absence of an FGF gradient. Even if receptor activation is uniform, downstream ERK activity could be spatially patterned by downstream negative regulators of ERK activity such as Sprouty-family proteins (SPRY) and dual-specificity phosphatases (DUSPs)[40].

To understand which genes modulating FGF/ERK signaling could be responsible for the elevated ERK signaling in TBXT positive cells and required for TBXT expression (Fig.1, Fig. 2b), we used single cell RNA-sequencing data from one previously published sample[34] and one newly collected replicate (Supp. Fig. 2a) to determine expression of FGF pathway components in human 2D gastruloids, extending earlier work describing FGF expression[25,35,36,41]. TBXT-positive PS-like cells give rise to nascent mesoderm expressing TBX6 as well as definitive endoderm expressing SOX17 and FOXA2, both of which may still express TBXT at 42h[42]. To understand how *TBXT* expression is initiated in an FGF-dependent manner, we focused on differential expression in the PS-LCs relative to pluripotent cells, reasoning that FGF modulation specific to nascent mesoderm or endoderm is a consequence rather than a cause of PS-LC differentiation. We identified different cell types based on marker genes with the PS-like cells defined as *TBXT+MIXL1+POU5F1+TBX6-SOX17-FOXA2-* (Fig. 2cd, Supp. Table 4). We then analyzed expression of different categories of FGF/ERK regulators.

We found four FGF ligands with significant expression: *FGF2, 4, 8,* and *17* (Fig. 2e, Supp. Fig. 2b). We then compared the expression of these ligands in PS-like cells to reference data of CS7 human, CS8 monkey, and E6.5-E7.5 mouse gastrulation (Fig. 2f, Supp. Fig. 2cd). We also included *FGF3* and *FGF5*, which are expressed during mouse gastrulation[32,43,44]. High expression of *FGF4* and *FGF17* and low expression of FGF8 was consistent between human, monkey, and 2D gastruloid. *FGF2* expression was also high in the PS-like populations in 2D gastruloids and the CS8 monkey embryo, but much lower in the human embryo data. Furthermore, the monkey and human embryos expressed *FGF3*, which was minimally expressed in gastruloids. A similar comparison in nascent mesoderm showed much lower *FGF2/4* and much higher *FGF17* compared to PS across all primate samples (Supp. Fig. 2e). Expression in 2D gastruloids was consistent with published RNA-seq data for hPSC-derived PS-like cells, which expressed high *FGF2* and *FGF4*, some *FGF8* and *FGF17*, and no *FGF3* (Supp. Fig. 2f). Comparison with

mouse confirmed significant differences from primates, with high expression of *FGF5, 8*, much lower expression of
FGF4,17, and no expression of *FGF2* in the mouse[21,43,44].
Of the FGF receptors, only *FGFR1* and *FGFR4* were strongly expressed, with *FGFR1* expression about 5-
fold higher (Fig. 2g, Supp. Fig. 2g). Both receptors showed graded expression that decreased from pluripotent to
PS-like cells and much lower expression in amnion-like cells (Fig. 2g). Therefore, elevated ERK activity in PS-like
cells cannot be accounted for by overall receptor expression. Differential expression of genes regulating heparin-
sulfate proteoglycans which modulate FGF receptor binding was mild except for glypicans 3 and 4 which were
downregulated in PS-like cells relative to pluripotent cells (Fig. 2h, Supp. Fig. 2g). In contrast, several intracellular
negative regulators of ERK signaling were strongly differentially expressed between cell clusters, including *SPRY1*
and *PEBP1* which were significantly higher in pluripotent cells (Fig. 2i, Supp. Fig. 2g). Surprisingly, however,
expression of *SPRY4* was not elevated in PS-like cells where ERK signaling is high, suggesting that transcription
of *SPRY4* may not be a good readout of ERK activity in human hPSCs like it is in the mouse[45]. On the other hand,
expression of the ERK negative feedback regulator *IL17RD* (*SEF*) does seem to approximately reflect expected
ERK activity in each population. Components of the core ERK pathway including ERK itself were highly expressed
across all clusters (Supp. Fig. 2g).
Thus, while scRNA-seq identified specific genes involved, it is consistent with several non-exclusive
mechanisms for generating an ERK pattern. For example, uniform FGF2 could form an ERK ring by equally
activating receptors across the colony but with ERK response reduced by elevated expression of negative
regulators in the pluripotent center. Alternatively, based on gene expression we cannot rule out HSPGs increasing
receptor binding specifically in the ring. Finally, localized expression of FGF4 or FGF17 in the PS-like ring could
lead to a concentration gradient responsible for increased receptor activation.
240
The phosphorylated ERK pattern is due to differential activation of FGF receptors.
To distinguish between the different possibilities, we asked whether the spatial pattern of pERK signaling is
controlled primarily below, at, or above the level of the receptors. To address this, we first visualized phosphorylated
FGFR1 (pFGFR1) and found its spatial pattern strongly resembles that of pERK (Fig.3a, Supp. Fig.3a). As expected,
pFGFR1 was lost upon brief FGFR inhibition (Fig.3a, Supp. Fig.3a). In contrast, after MEK inhibition, high pFGFR1
remained while pERK was greatly reduced (Supp. Fig.3bc). This suggests the spatial pattern of ERK reflects a
spatial pattern of receptor activity.
The spatial pattern in receptor activity could either be due to differential receptor expression or differential
receptor activation. There was no significant upregulation of FGF receptors in PS-like cells according to our single
cell RNA-sequencing data (Fig 2e). In fact, the most highly expressed receptor, FGFR1, was elevated in pluripotent
cells. This suggests the ERK activity pattern is not due to a receptor expression pattern with two caveats. First,
mRNA expression may not reflect protein expression. Second, multiple isoforms exist for FGF receptors 1-3 (but
not 4) that have different affinity for FGF ligands, leaving open the possibility that PS-like cells could express a
different FGFR1 isoform than pluripotent cells, despite overall FGFR1 expression being approximately uniform.
To determine if the spatial pattern in pFGFR1 could be explained by a spatial pattern of FGFR1 protein, we
performed immunofluorescence staining of FGFR1. Consistent with the scRNA-seq data, and in sharp contrast to
the pFGFR1 stain, FGFR1 protein expression was graded from center to edge with its highest levels in the
pluripotent center and therefore could not explain the signaling pattern (Fig.3b, Supp. Fig.3d). To determine if
different isoforms of FGFR1 are differentially expressed in PS-like vs. pluripotent cells, we designed qPCR primers
to detect the IIIb and IIIc isoforms of *FGFR1* (Supp. Fig. 3e-g). These are the most common isoforms, also known
as the epithelial and mesenchymal isoforms, and therefore appeared the mostly likely candidates for differential
expression. We then analyzed expression of these isoforms in 42h micropatterns versus pluripotent cells and PS-
like cells differentiated directly as in Fig.1d. However, the change in expression between samples was similar for
each isoform, with all isoforms decreasing in 2D gastruloids and directly differentiated PS-like cells relative to
pluripotent cells (Fig. 3c). To corroborate this and also determine absolute FGFR isoform expression in pluripotent
and PS-like cells, we analyzed our single cell RNA-sequencing data and previously published bulk RNA-sequencing
data for isoform expression in these two cell types. This revealed that for both pluripotent and PS-like cells FGFR
IIIc is the dominant isoform with on average almost 900-fold higher expression than IIIb, while IIIa was not expressed
at all (Fig. 3d). Altogether, these data suggest that spatial patterning in ERK is due to differential activation of FGFR
receptors, either due to FGF concentration gradients or possibly spatially patterned HSPGs. Immunostaining for
HS showed no clear correlation with TBXT or the expected ERK signaling pattern which is similar but extends
further inward (Supp. Fig.4). This does not rule out a role for HSPGs in spatial modulation, since FGF signaling
depends on specific sulfation patterns rather than overall HS levels[46,47]. Nevertheless, it led us to focus on FGF
gradients as the most likely cause for the ERK signaling pattern.
275

Figure 3. a) phospho-FGFR1 stain with and without 1h FGFRi treatment at 41h, radial intensity profile on the right. **b)** FGFR1 with and without 1h FGFRi treatment at 41h, radial intensity profile on the right. Error bars in (b,c) represent standard deviation over N=4 colonies. **c)** qPCR data for relative expression of FGF receptor 1 isoforms IIIb, IIIc and total FGFR1 in pluripotent cells (pluri), gastruloids at 42h (MP), and directly differentiated PS-like cells (dPS). Error bars represent standard deviation of technical triplicates. **d)** Absolute FGFR1 isoform expression from RNA-seq for anterior/mid primitive streak and pluripotent cells from Loh 2016[38] (Loh_APS/MPS/HESC) as well as the pluripotent and PS-LC clusters from Fig.2. All scale bars 50 um.

FGFR1-IIIc is the dominant isoform across the colony

FGF receptors are localized basolaterally.

To maintain an endogenous FGF concentration gradient on the apical side of the colony which faces the medium, would require ligands to remain tethered to the surface. However, FGF has been reported to diffuse freely in the intercellular space[48]. Moreover, there is FGF2 present in the medium that would be expected to uniformly activate FGF receptors. The situation is similar for BMP4, which is present uniformly in the medium for 2D gastruloids but does not cause uniform signaling activity[4]. In the case of BMP, that is because receptors are localized basolaterally[4] and tight junctions seal the basolateral side from the medium. Consequently, response to exogenous ligands in the medium is restricted to the colony edge, while endogenous ligands are confined to the basal side and prevented from diffusing away into the medium everywhere but the edge. This was found to be critical for patterning in both 2D gastruloids and the mouse embryo[4,5]. To explain the pERK pattern, we therefore hypothesized FGF receptors were similarly restricted to the basolateral side.

To test our hypothesis, we first starved pluripotent colonies of FGF by maintenance in FGF-free E6 medium for 12h and then treated with FGF2. Consistent with our hypothesis, phospho-FGFR and phosphor-ERK were both restricted to the colony edge (Fig. 4a). To determine if this could be explained by diffusion of exogenous FGF from the medium, we repeated the experiment with micropattern pluripotent cells and added fluorescent dextran with a similar molecular weight as FGF. We observed fluorescent dextran diffusing into the intercellular space from the edge inward over the same distance that ERK signaling was observed (Fig. 4b-d). We then acquired high resolution z-stacks of FGFR1 co-stained with the tight junction protein TJP1 (ZO-1) which revealed the majority of FGFR1 was below the tight junctions (Fig. 4e). To directly test that cells are more responsive to basal FGF than apical FGF, we then performed a transwell assay, where pluripotent cells were grown on filters to enable separate apical and basal stimulation. We found that only basal stimulation with FGF led to a significant increase in pERK (Fig. 4f). Finally, to determine if the center of the micropatterned colony is similarly able to respond to basal FGF stimulation and pERK levels there can be attributed to reduced FGF ligand concentration, we performed a scratch assay. Strong ERK response to FGF2 in the media was seen in the colony center around a scratch, but this response was blocked by FGFR inhibitor (Fig.4gh). If HSPGs rather than FGF ligand were limiting ERK response in the center, we would not have expected strong response around a scratch. The initial ERK levels around the scratch greatly exceeded those anywhere in the control. However, we found this was due to adaptive ERK response to FGF. After six hours ERK levels around the scratch were similar to those in the ERK ring around the PS-like cells (Fig. 4ij). We therefore concluded that the pERK pattern is most likely due to a basolateral FGF ligand gradient which has its maximum in the PS-like ring.

Response to exogenous FGF2 treatment is restricted to the colony edge

FGF receptors are localized basolaterally

ERK response to FGF2 is adaptive

Figure 4. a) pERK and pFGFR1 response of pluripotent colonies in standard culture grown in FGF-free (E6) medium for 24 hrs and then treated with FGF2 for 30 minutes. **b,c)** Fluorescent dextran (b) and pERK (c) in micropatterned pluripotent colony 24h after media change containing both FGF2 and dextran. **d)** Radial intensity profile of dextran and pERK for conditions in b,c), averaged over 2 colonies. **e)** Cross-section of 2D gastruloid shows FGF receptor 1 predominantly below the tight junction marked by ZO-1. White box marks are magnified in inset e'. **f)** pERK and pFGFR1 response in transwell experiment with cell treated with FGF either apically or basally. **g)** ERK signaling in scratched colonies with or without FGFRi. **h)** Quantification of pERK intensity as a function of distance from the scratch. Error bars represent standard deviation over colonies. **i)** ERK signaling stains at different times after scratching. **j)** Quantification of pERK intensity as a function of distance from the scratch at different times (left) and peak pERK level over time (right). Error bars represent standard deviation over N=4 colonies. Scale bars 50 micron.

347
Endogenous FGF gradients underly the ERK activity pattern.
348
To identify which FGF ligands are required for PS-like differentiation, we first determined the requirement for
exogenous FGF2 in the media. Differentiation of 2D gastruloids in FGF-free medium was severely reduced but
restored by addition of FGF2 (Fig. 5a, Supp. Fig. 5a). However, if FGF2 was removed 24h after BMP treatment,
differentiation was normal. The fact that removal 2h after BMP treatment significantly reduced differentiation
suggests this was not a failure to wash out the FGF2. Furthermore, MEK inhibition at 24h severely reduces TBXT
expression[34] while FGF2 is known to have a half-life on the order of hours[49] and no media changes are performed
during standard differentiation. Altogether this suggests that exogenous FGF2 is required at the time of BMP
treatment but that endogenous FGFs are sufficient for the ERK activity ring and PS-like differentiation that start a
day later.
We reasoned that FGFs whose expression correlates with the presence of PS-like cells are more likely to
be responsible for the observed ERK ring and PS-like differentiation. Our analysis in figure 2 had implicated FGF4
and FGF17 as most strongly expressed in the PS-like cells and nascent mesoderm, respectively (Fig. 2c). However,
it is unclear whether expression of these ligands is a cause or consequence of PS-like differentiation, or both, in the
case of a feedback loop. To get more information about the relationship between FGF ligands and PS versus
mesoderm differentiation, we performed pseudotime analysis of our scRNA-seq data (Fig. 5b, Supp. Fig 5b). Along
a pseudotime trajectory from pluripotent cells to nascent mesoderm, *FGF2* expression was nearly constant. *FGF8*
was upregulated at lower levels prior to *TBXT* and declined as *TBXT* was upregulated. *FGF4* upregulation
approximately coincided with *TBXT*, but *FGF4* began decreasing as *TBX6* expression increased. In contrast, *FGF17*
expression started after *FGF4* but continued to increase after *FGF4* started decreasing. Direct comparison of *FGF*
and *TBXT* expression also showed strong positive correlation between *TBXT* and both *FGF4* and *FGF17* but
opposing relations with *TBX6* (Supp. Fig. 5c). Together this suggested to us initial TBXT expression may depend
more on FGF8 or FGF4, while FGF17 could play a later role in maintaining or expanding the PS-like population or
facilitating subsequent mesoderm differentiation.
Pseudotime may not accurately predict true dynamics. Therefore, we verified expression at two different
times using RNA FISH (fluorescent in situ hybridization). We found that *FGF4* expression was similar at 32h and
40h (Fig.5cf, Supp. Fig. 5d). This is consistent with its decrease in pseudotime as cell differentiate to nascent
mesoderm, since in real time differentiation is asynchronous and new PS-like cells form while older ones
differentiate to nascent mesoderm, which would yield a flatter temporal profile. In contrast, peak *FGF8* levels
decreased while *FGF17* strongly increased over time, both matching their pseudotime trends (Fig.5d-f, Supp. Fig.
5ef). As expected, the spatial profiles of both *FGF4* and *FGF17* resembled TBXT, although the peaks were
displaced slightly inward, while *FGF8* expression was highest in the inner part of the ERK ring where TBXT
expression was low (Fig. 5c-f). Co-staining showed *FGF4* and *FGF8* expressed in concentric rings with little overlap
at 42h (Supp.Fig.5g). Scatterplots further confirmed that high expression of both *FGF4* and *FGF17* but not *FGF8*
was restricted to TBXT positive cells (Supp. Fig. 5h).
To test the functions of these FGFs we decreased their expression. Knockdown (KD) of *FGF4* with small
interfering RNAs (siRNA) led to a strong decrease in TBXT and pERK (Fig. 5g, Supp.Fig.6a). We repeated this in
a different cell line and also created stable cell lines expressing short hairpin RNA (shRNA) to knock down *FGF4* in
both cell lines with similar effect (Supp.Fig. 6b-d). Strikingly, *FGF8* KD had the opposite effect and led to an increase
in TBXT expression without a strong effect on pERK visible at 42h (Fig.5h, Supp.Fig.6e). Finally, *FGF17* KD caused
only a small decrease in TBXT or pERK, again consistent between different cell lines (Fig.5i, Supp.Fig.6f-i). We
further interrogated the endoderm and PGC marker SOX17 since both cell types require ERK signaling to
differentiate and transiently express TBXT, as well as the mesoderm marker TBX6. TBX6 expression at 42h was
near completely lost upon either *FGF4* KD or *FGF17* KD (Fig.5j, Supp.Fig.6j). This suggests that while FGF17 is
not required for PS induction (Fig.5i), it is needed for subsequent mesoderm differentiation. In contrast, there was
only a small reduction of SOX17 (Fig.5j). Since we previously found with our protocol most SOX17 positive cells at
42h represent PGC-LCs, while most endoderm differentiates later[34], this suggests FGF4/17 are not required for
PGC-LC induction but does not rule out a requirement for these FGFs in endoderm differentiation. A double KD of
FGF4 and *FGF17* had a similar phenotype as *FGF4* KD alone, suggesting FGF17 may be downstream of FGF4
(Supp. Fig. 6k). Combined, these data support distinct functions for FGF4, FGF8, and FGF17 in our human
gastrulation model, where FGF4 and FGF8 may play opposing roles in PS induction while FGF17 is important
during continued differentiation to mesoderm.
400

Exogenous FGF2 is only required at the start of differentiation

FGF4 and FGF17 expression correlate with TBXT, while FGF8 peaks earlier and is displaced inward

Reduction in FGF4 or FGF17 leads to reduction in PS-like cells and derivatives, while FGF8 reduction increase PS

Figure 5. a) Pattern at 42h with exogenous FGF2 in the culture medium for different amounts of time. **b)** Pseudotime analysis of FGFs and TBXT, TBX6 expression. **c-e)** FGF4 (c), FGF8 (d), and FGF17 (e) FISH co-stained for TBXT and pERK at 30h or 32h and 40h or 42h after BMP4 treatment. **f)** Radial intensity corresponding to (c-e). **g-i)** FGF4 (g), FGF8 (h), and FGF17 (i) FISH co-stained for TBXT and pERK in control siRNA versus FGF4 siRNA (g), FGF8 siRNA (h) and FGF17 siRNA (i) with corresponding radial intensity profiles. **j)** TBX6 and SOX17 expression in control shRNA versus FGF4 and FGF17 shRNA. Error bars in all panels except (b) represent standard deviation over N=4 colonies. Error bars in (b) represent the 95% confidence interval of the generalized additive model (see methods).

Discussion

We have shown that in a stem cell model for early human gastrulation, FGF-dependent ERK signaling is directly required for differentiation to PS-like cells and their derivatives, and that loss of differentiation upon FGF inhibition can be rescued by directly activating ERK. We also showed that FGF receptors are polarized basolaterally, preventing the colony center from responding to exogenous FGF in the medium. Furthermore, FGF receptor activation is increased in the PS-like ring, matching the pattern of active ERK despite overall receptor expression being lower in the PS-like ring than in the pluripotent center. This suggests elevated ERK activity is due to increased binding of FGFs to the receptors in this region, possibly due to increased FGF concentration. We found that both FGF4 and FGF17 are highly expressed in PS-like cells, while FGF8 is expressed earlier and displaced inward (Fig.6). Knockdown of FGF4 leads to a strong reduction in ERK activity and PS-like cells, suggesting it may be responsible for increased ERK signaling in the PS-like region. In contrast, FGF8 knockdown led to an expansion of ERK activity and PS-like differentiation. Finally, FGF17 knockdown leds to a loss of mesoderm differentiation.

Figure 6. Graphical summary. Exogenous FGF2 acts on the colony edge and is prevented from reaching the basal FGFR1 receptors elsewhere by tight junctions. Primitive streak cells express FGF4 and FGF17 and these FGFs are associated with high ERK activity. FGF8 is expressed further inside the colony.

Many questions remain. Although we showed that FGF gradients are likely the dominant cause for the spatial pattern of ERK signaling, our scRNA-seq data suggest other levels of regulation may contribute, which remain to be explored. Furthermore, although we showed FGF/ERK signaling is directly required for PS-like differentiation, it may also act indirectly by affecting the expression of Wnt and Nodal. We will explore this and the more general question of how FGF integrates with the hierarchy of BMP, Wnt, and Nodal in a separate manuscript. Related is the indirect effect FGF may have on patterning by its control of cell growth and thereby cell density. Cell density affects the response to exogenous BMP and downstream expression of Wnt, Nodal, and various inhibitors[4,50,51], as well as FGF itself, possibly causing a "community effect"[52–54]. It likely has other effects as well. However, the rescue of TBXT expression but not growth by ERK activation when FGF receptors are blocked suggests the FGF-dependent cell growth is not required to generate a primitive streak-like ring. The ability of exogenous Wnt and Nodal to induce PS-LCs in very sparse culture while MEK and FGFR inhibition block differentiation at the same density similarly argues against a requirement for FGF-dependent cell growth in PS-LC differentiation (Supp.Fig.1h).

Another important question is to what extent different FGFs act redundantly or perform different functions. Our data suggest a picture where different FGFs are required consecutively, starting with exogenous FGF2, then endogenous FGF4, to support initial differentiation of PS-like cells, and finally FGF17 to maintain and expand the PS-like ring and support mesoderm and possibly endoderm differentiation. Whether these FGFs could substitute for each other if they were expressed at similar times and levels is unclear. It also remains to be determined whether the nearly constant high expression of FGF2 we observed is required for patterning. Most puzzling is the role of FGF8, whose knockdown led to an increase in PS-like differentiation. Given that FGF8 expression is displaced inward, one possibility is that it induces inhibitors of other PS-inducing signals there in order to limit PS differentiation, e.g., the Nodal inhibitor Lefty, which is expressed in a similar domain[55]. The rescue of PS-like differentiation by uniform activation of ERK to a supraphysiological level suggests that the role of the different FGFs is simply to sufficiently activate ERK signaling. However, future work will have to investigate the dox-SOS[cat] rescue phenotype in much greater detail to see how differentiation and other cell behaviors such as migration are affected.

It is interesting to ask if there is a simple explanation for the seemingly large differences in FGF expression between primates and other species. FGF8 is the only FGF shown to be required for mouse gastrulation and is upstream of FGF4 in both chick and mouse, but it is barely expressed in human and monkey primitive streak and appears to play a different role in our stem cell model. If FGFs are interchangeable in simply activating ERK for primitive streak induction, variation in FGF expression is not surprising. However, FGF4 and FGF8 are the main FGFs in mouse, chick and frog gastrulation[56–58], suggesting a high degree of conservation. It is possible FGF8 has simply been substituted by FGF17, which is in the same subfamily[9], but this does not explain why the role of FGF8 in gastrulation otherwise appears strongly conserved, or why FGF8 is upstream of FGF4 in the mouse while FGF4

is expressed before FGF17 in 2D gastruloids. Similarly, it is possible that FGF2 in primates substitutes for FGF5 in
the mouse as the FGF that is highly expressed in the epiblast. Although these are not in the same subfamily and
generally have different receptor affinities, they both have highest affinity for FGFR1[9]. That leaves FGF4 as the only
FGF that may be conserved across vertebrate gastrulation, and it is tempting to speculate on why. In Xenopus,
FGF4 and TBXT (Xbra) function in a positive feedback loop[22,59]. The strongly correlated initial transcriptional
dynamics of TBXT and FGF4 in 2D gastruloids are consistent with a similar feedback loop. However, an FGF8
feedback loop has also been proposed and recent work suggests TBXT is not required for FGF4 or FGF17
expression[54,60,61]. Future work will therefore have to investigate further if FGF4 plays a unique role.
A final related question is the origin of the discrepancies we and others[36] found between the primate (i.e.,
non-mouse) samples in Fig. 2 and Supp. Fig. 2: FGF3 appears only significantly expressed in the human embryo,
but not in the monkey embryo and human stem cell model. This could also reflect genetic variation: if FGFs function
redundantly, their relative expression levels could be relatively unconstrained between individuals. On the other
hand, in the nascent mesoderm, FGF2 is only high in human gastruloids. This may reflect developmental stage: the
in vivo data is developmentally more advanced and nascent mesoderm may lose FGF2 expression later. It is also
possible the maintenance conditions of hPSCs lead to in vitro artefacts in FGF expression.
In conclusion, the complexity of FGF signaling, with its many ligands, modulators, and functions, has
prevented a clear understanding of its role in mammalian gastrulation in general and in human gastrulation
specifically. Here we showed that 2D human gastruloids provide a powerful approach to make headway and
although many open questions remain, we have taken significant steps towards this goal.
493
494

Acknowledgements

We thank Lila Solnica-Krezel, Blerta Stringa, Aryeh Warmflash, and Ben Allen for discussions. We thank Jared Toettcher for the dox-SOScat plasmid. Library prep and next-generation sequencing was carried out in the Advanced Genomics Core at the University of Michigan. This work was supported by the National Institute of General Medical Sciences (NIGMS R35GM138346), and the Branco Weiss Fellowship – Society in Science. KJ was partially supported by the Michigan Pioneer Postdoctoral Fellowship and the NIH F32 Ruth L. Kirschstein Postdoctoral National Research Service Award (5F32HD108980-02).

Data availability

Raw single cell RNA sequencing data generated in this study was deposited in GEO (GSE271604), analyzed data will be added upon publication. Raw image data are available upon request.

Code availability

All code for data analysis and model simulations is available on https://github.com/idse/FGF

Competing Interests Statement

The authors declare no competing interests.

Methods

Cell lines

The cell lines used were the embryonic stem cell line ESI017 (XX), and the induced pluripotent stem cell line PGP1 (XY). The pluripotency of these cells was confirmed by immunostaining of pluripotency markers OCT3/4, SOX2, NANOG. All cells were routinely tested for mycoplasma contamination, and negative results were recorded. ESI017 cells stably expressing doxycycline inducible SOS[cat] were made using a published Piggybac plasmid gifted by Jared Toettcher[37].

Cell culture and differentiation

Human pluripotent stem cells were cultured in the commercially defined pluripotency maintaining media mTeSR1 (STEMCELL Technologies #85850) on Cultrex (R&D Systems)-coated tissue culture plates. For FGF2 starvation we used Essential 6 Medium (Thermofisher, A1516401), or mTeSR1 without bFGF, TGF-beta, LiCl, GABA and pipecolic acid (STEMCELL Technologies #05896). For routine cell maintenance, cell passaging was performed every 3 days. For whole-colony passaging, L7[62] was used, while single-cell passaging was performed with Accutase or TrypLE (Gibco). Single-cell suspensions were used to seed for all the differentiation experiments to control initial cell number.

To directly differentiate to primitive streak-like cells, cells were resuspended in a single-cell suspension and seeded in cultrex coated wells in mTeSR with ROCK inhibitor (RI) Y-27632 (MeChemExpress, cat# HY-10583). Cells were then maintained in mTeSR with RI for 24h before adding CHIR (3-5uM) and Activin (100ng/ml) treatment for 24-30h. Experiments were performed in 18-well Ibidi slides (cat# 81818).

To differentiate cells for 2D gastruloids by micropatterning, we followed the protocol in[42]. Briefly, cells were resuspended in a single-cell suspension and seeded in laminin-coated micropatterned wells in mTeSR with ROCK inhibitor (RI) (MeChemExpress, cat# HY-10583). Micropatterned wells were washed with PBS-/- 30 minutes after the initial seeding to wash off the non-attached cells. Cells were then maintained for 24 hours in mTeSR before adding the BMP4 treatment by a full medium change. For time series micropatterned differentiation, full media changes with designated treatment(s) were performed every 24 hours. Experiments were performed in micropatterned 18-well Ibidi slides (cat# 81818) prepared as previously described[63]. Cell signaling reagents and doses used are listed in Supplementary Table 1.

Immunofluorescence staining

Antibodies can be found in Supplementary Tables 2 and 3. For stains including pERK, we used methanol fixation, for all other stains we used PFA fixation. PFA fixation: Samples from the 18-well Ibidi slides were rinsed with PBS, fixed for 20 min in 4% paraformaldehyde, rinsed twice with PBS, and blocked for 30 min at room temperature with 3% donkey serum and 0.1% Triton X-100 in 1× PBS. After blocking, cells were incubated with primary antibodies

at 4°C overnight, followed by three washes in PBST (PBS with 0.1% Tween 20). They were then incubated with
secondary antibodies and DAPI for 30 min at room temperature and washed twice in PBST at room temperature.
Methanol Fixation: Samples from the 18-well Ibidi slides were rinsed with PBS, fixed for 20 mins in cold methanol,
then rinsed with PBS for 5 min two times, after which 10% phosphate buffered formalin was added for 20 min at
room temperature. Cells were then rinsed with TBS twice followed by 5 mins of incubation in cold methanol and
rehydrate in TBS for 20 mins. Blocking was done for 30 mins at room temperature with 5% donkey serum, 1% BSA,
0.2% Triton-X-100 in 1x TBS after which staining proceeded the same way as with PFA fixation.
559

Microscopy
561   Fixed sample imaging was performed with an Andor Dragonfly/Leica DMI8 spinning disk confocal microscope with
562   a 40x, 1.1NA water and 20x 0.8NA air objectives using Andor Fusion software version 2.3.0.31 and a
563   Nikon/Yokogawa spinning dish confocal microscope with a 40x silicon oil objective using NIS Elements AR software
564   version 5.41.02. Live-cell imaging was performed with an Andor Dragonfly/Leica DMI8 spinning dish confocal
565   microscope under the controlled temperature (37°C), CO_2 concentration (5%), and humidity (>60%).
566

General image analysis
We segmented nuclei in individual z-slices based on nuclear fluorescence stained with DAPI in fixed cells using a
pipeline we previously described, which integrates two machine learning approaches: Ilastik pixel classification[64]
and Cellpose[65] (v1). Fluorescence intensities were then calculated per nucleus as mean intensities in the 3D
nuclear again with our established custom image-processing pipeline. To calculate the radial fluorescence
distributions in micropatterned colonies, colonies were subdivided into radial bins with equal numbers of cells, mean
or median and variance were then calculated within these bins. To calculate positive fractions, markers were first
thresholded based on their intensity distribution, after which the fraction of cells positive for each marker was
calculated within different bins. Standard deviations were then calculated over multiple replicate colonies. For
visualization purposes only, background subtraction using grayscale opening on a 100 micron scale and 5 pixel
wide median filter was applied slice by slice to z-stacks of micropatterned colonies before making maximal intensity
projections.
579

(Single-cell) RNA sequencing and analysis
Cells were collected using accutase and resuspended in ice-cold PBS. Single-cell RNA-sequencing was performed
by the University of Michigan Advanced Genomics Core. Cells were barcoded using the 10X Genomics Chromium
system (part numbers 1000268, 1000120, 1000215). For quality control, cDNA was quantified by Qubit High
Sensitivity DNA assay and Agilent TapeStation. Sequencing was performed on the Illumina NovaSeq 6000 with
NovaSeq S4 flowcell and Control Software version 1.7.0. Reads were aligned using cellranger-4.0.0 with the
GRCh38 reference. Further processing was mostly done in Python, primarily using the Scanpy[66] and SCVI[67]
packages. Analysis was performed on log transformed data. Library size normalization was performed on the
intersection of the genes present in our data and the reference CS7 human embryo data to facilitate comparison of
expression levels. We integrated the two scRNA-seq datasets for 42h 2D human gastruloids using SCVI based on
the top 2000 highly variable genes, which were selected using the default function in Scanpy using the seurat_v3
flavor. Cell cycle genes were regressed out during integration. Dimensional reduction for visualization was
performed using PHATE[39] on the SCVI latent space. Gene expression was smoothened with MAGIC[68] for
visualization on PHATE plots in Fig. 2e,g-i. To annotate different cell types, we used MAGIC to impute that data
and get smooth distributions of gene expression. We then thresholded marker genes to assign cell fate (Table 4),
which produced better results than the more common unsupervised clustering using Leiden, and in particular more
consistent results between datasets from different species. For interspecies comparison of FGF ligands we
normalized the to the maximal FGF expression within each species, where for the mouse data, the three different
time points were first integrated using SCVI. Thus we compare between species the relative expression of FGF
ligands within each species, e.g., FGF8 expression is much higher than FGF4 at any time in the mouse, but much
lower than FGF4 in human.
For pseudotime analysis (Fig. 5), we applied the trajectory inference tool Slingshot[69] in R to compute the
pseudotime trajectory in the SCVI latent representation. Slingshot determines the membership of a cell to a lineage
at each branching point by its projection distance to the fitted principal curve of that lineage. We projected the
mesoderm developmental trajectory onto the PHATE map for visualization. We further used TradeSeq[70], which fits
a generalized additive model (GAM), to infer the pseudotime gene expression trends along the mesoderm lineage
trajectory from Slingshot. For visualizing the correlation between the expression of FGF ligands and marker genes
(Supp. Fig. 5), we used our SCVI model to denoise the expression and mutual nearest neighbor to further harmonize
the difference between samples.

To quantify isoforms from RNA sequencing salmon v1.10.0 was used[71]. First, gencode.v45.transcripts.fa
was downloaded from gencodegenes.org. The reference was customized by removing lines for FGFR1 and
replacing them with the transcript sequence for FGFRIIIa, FGFRIIIb, FGFRIIIc (sequences included as
supplementary file).   Three samples from GEO dataset GSM2257301 were downloaded through SRA
(SRX1725562 (Pluripotent), SRX1725577 (APS), SRX1725643 (MPS))[38]. Adapters and the first 10 bases were
trimmed from the FASTQ files using trim_galore. Finally, the Salmon "quant" command with the --validateMappings
option was used to quantify transcripts from trimmed FASTQ files. For single cell datasets,
https://timoast.github.io/sinto/ was used to subset the 10x aligned file by cluster. Reads for clusters of interest were
converted back to raw FASTQ files and run through the same alignment-based mode of salmon but as single end
reads.
619
Quantitative real-time PCR
RNA were extracted using RNAqueous™-Micro Total RNA Isolation Kit (CAT# AM1931, Thermo Fisher Scientific)
and then cDNA synthesis was prepared from it using SuperScript™ VILO™ cDNA Synthesis Kit (CAT# 11754250,
Thermo Fisher Scientific) following the manufacturer's protocol with adjustment for optimization. Measurements
were performed with SYBR green and the primers in the Table 5. GAPDH was used for normalization in all
experiments.
626
Fluorescent dextran assays
Fluorescent dextran essays were performed by adding 10 ug/ml fluorescein isothiocyanate-dextran (10KD, Sigma
Aldrich) to the media. To calculate fluorescence intensity as a function of distance from the colony edge, Otsu
thresholding followed by several morphological operations was used to create a mask for the colony in each z-slice,
thereby excluding bright dextran fluorescence outside the colony. Background subtraction on each z-slice of the
image was performed to further remove background signal from dextran outside the colony while maintaining signal
in the intercellular space. The radial intensity profile was then calculated from the maximal intensity projection of
the masked, background subtracted image.
635
Scratch assays
Human embryonic or induced pluripotent stem cells were differentiated on micropattern wells for 41h, then each
micropattern colony was manually scratched with a serological needle. After 60mins, they were fixed with PFA or
methanol followed by immunostaining with appropriate antibodies. To calculate the intensity profile as a function of
distance from a scratch, we created a mask for the scratch based on the difference between an overall mask for
the colony and the convex hull, calculated the distance transform of the scratch mask, then binned pixels by distance
from the scratch and calculated their mean ERK intensity.
643
Transmembrane assays
Human embryonic or induced pluripotent stem cells were seeded into Corning 6.5 mm transwell with 0.4 μm pore
polyester membrane insert (cat# 3470) with E6 supplemented with TGF1b, FGF2, and Y-27632 for 16h, then media
was changed to E6 with TGF1b. After 24h, the media was changed to E6 supplemented with FGF2 from the top or
the bottom side of the transwell. 30 mins later, cells were fixed followed by immunofluorescence staining. The
transwell cartoon in Fig 4 was generated with Biorender.
650
651 **Fluorescence In Situ Hybridization**
652 The fluorescence in situ hybridization protocol (FISH) was performed according to the manufacturer's instructions
653 (ACDbio, RNAscope multiplex fluorescent manual). A list of probes and reagents can be found in Supplementary
654 Table 6.
655
Gene targeting by shRNA
We used the constructs shRNA F4: pPB[shRNA]-EGFP-U6>hFGF4[shRNA#1] and shRNA F17: pPB[3shRNA]-
U6>hFGF17[shRNA#1]-U6>hFGF17[shRNA#2]-U6>{hFGF17_shRNA3}-hPGK>Bsd   both   purchased   from
VectorBuilder in a PiggyBac Transposon Vectors with puromycin resistance which was stably integrated into ESI17
hESCs and PGP1 hiPSCs. Transfections were performed using Lipofectamine™ Stem Transfection Reagent
(CAT# STEM00001, Thermo Fisher Scientific), following the manufacturer's protocol.
662
663 **Gene targeting by siRNA**
664 We purchased predesigned siRNA from Millipore Sigma; NM_002007 (FGF4), NM_006119 (FGF8), NM_008004
665 (FGF17). The siRNA transfections were performed using Lipofectamine™ RNAiMAX Transfection Reagent (CAT#
666 13778075, Thermo Fisher Scientific), following the manufacturer's protocol with adjustment for optimization.

667
Statistics and reproducibility
669 All experiments were performed twice or more. All the attempts in replicating experiments yielded consistent results.
670 Unless specifically noted, all the quantifications were performed with N=4 colonies within the same experimental
671 condition.
672
Ethics statement
674 We comply with all the ethical regulations in our research. We were granted approval by the Human Pluripotent
675 Stem Cell Research Oversight (HPSCRO) Committee at the University of Michigan to work with human embryonic
676 stem cells and human induced pluripotent stem cells.
677
678 **Supplementary Table 1: Cell signaling reagents.**

Reagent	Nickname	Vendor, cat #	Dose	Function
rhBMP4	BMP	R&D Systems, 314BP/CF	50ng/ml	Activate BMP signaling
rhFGF2	FGF2	Gibco, PHG6015	100ng/ml	Activate FGF receptors
rhTGFβ1	TGFβ	Peprotech, 100-21C- 2UG	40ng/ml	Activate TGFb receptors
rhActivinA	Activin	R&D Systems, 338AC050	100ng/ml	Activate Activin/Nodal pathway
LDN193189	BMPRi	MedChemExpress, HY-12071	250 nM	Inhibit BMP receptors
IWP2	WNTSeci	ApexBio, A3512-5	5 μM	Block Wnt secretion
PD0325901	MEKi	ESIBIO, ST10009	5uM	Inhibit MEK function
PD173074	FGFRi	Medchemexpress,HY10321	1uM	Inhibit FGF receptors
SB 431542	SB	Fisher, 50-101-2514	10uM	Inhibit Nodal receptors
CHIR 99021	CHIR	Tocris, 4423	See figures	Canonical Wnt agonist

679
680 **Supplementary Table 2: Primary antibodies used for immunofluorescence.**

Antigen	Species	Dilution	Cat #	Vendor
ISL1	Mouse	1:200	39.4D5	DSHB
SOX2	Rabbit	1:200	3579S	Cell Signaling Tech
TBXT (BRA)	Goat	1:300	AF2085	R&D Systems
EOMES	Mouse	1:200	MAB6166	R&D Systems
MIXL1	Rabbit	1:150	A17223	ABclonal
pERK	Rabbit	1:200	9101S	Cell signaling Tech
pFGFR1	Rabbit	1:100	52928	Cell signaling Tech

FGFR1	Rabbit	1:100	9740T	Cell signaling Tech	
Heparan Sulfate	Mouse	1:100	370255-S	AMS Bio	

682 **Supplementary Table 3: Secondary antibodies.**

Antigen	Species	Dilution	Cat #	Vendor
Alexa Fluor 488 anti-mouse	Donkey IgG	1:500	A21202	ThermoFisher Scientific
Alexa Fluor 555 anti-rabbit	Donkey IgG	1:500	A31572	ThermoFisher Scientific
Alexa Fluor 647 anti-goat	Donkey IgG	1:500	A21447	ThermoFisher Scientific
Alexa Fluor 647 anti-rabbit	Donkey IgG	1:500	A31573	ThermoFisher Scientific
Alexa Fluor 555 anti-goat	Donkey IgG	1:500	A21432	ThermoFisher Scientific

684 **Supplementary Table 4: Marker genes for cell fate annotation**

Cell type	Markers
Pluripotent	SOX2+POU5F1+TBXT-MIXL1-SOX17-
Amnion-like cell (AM-LC)	ISL1+POU5F1+TBXT-MIXL1-SOX17-
Primitive streak-like cell (PS-LC)	TBXT+MIXL1+POU5F1+TBX6-SOX17-FOXA2-
Nascent mesoderm (NasMe)	TBXT+MIXL1+TBX6+SOX17-
Endoderm	SOX17+EOMES+TFAP2C-
Ectoderm	SOX2+POU5F1-TBXT-MIXL1-SOX17-
Primordial germ cell-like cells (PGC-LC)	SOX17+TFAP2C+

686 **Supplementary Table 5: qPCR primers**

Gene	Forward	Reverse
FGFR1	AATGAGTACGGCAGCATCAAC	ACCTCGATGTGCTTTAGCCAC
FGFR1 IIIb	GCCCAGACAACCTGCCTTAT (Binds to FGFR1-IIIa)	CACACATACTCCCCGCTCTG (Binds to FGFR1-IIIb)
FGFR1 IIIc	GCCCAGACAACCTGCCTTAT (Binds to FGFR1-IIIa)	AGAGTGATGGGATCCGA (Binds to FGFR1-IIIc)
GAPDH	ACAACTTTGGTATCGTGGAAGG	GCCATCACGCCACAGTTTC

689 **Supplementary Table 6: Probes for Fluorescent In Situ Hybridization**

Probe	TSA dilution	Cat #	Vendor
Hs-FGF4	1:2000	RNAScope #512291	ACDBio

| Hs-FGF8-C2 | 1:2000 | RNAScope 415791-C2 | ACDBio |
| Hs-FGF17-C1 | 1:2000 | RNAScope #1148351-C1 | ACDBio |

690
691
References

1.　Arnold, S. J. & Robertson, E. J. Making a commitment: cell lineage allocation and axis patterning
in the early mouse embryo. *Nat. Rev. Mol. Cell Biol.* **10**, 91–103 (2009).
2.　Martyn, I., Kanno, T. Y., Ruzo, A., Siggia, E. D. & Brivanlou, A. H. Self-organization of a human
organizer by combined Wnt and Nodal signalling. *Nature* (2018) doi:10.1038/s41586-018-0150-y.
3.　Chhabra, S., Liu, L., Goh, R., Kong, X. & Warmflash, A. Dissecting the dynamics of signaling
events in the BMP, WNT, and NODAL cascade during self-organized fate patterning in human
gastruloids. *PLoS Biol.* **17**, e3000498 (2019).
4.　Etoc, F. *et al.* A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid Self-
Organization. *Dev. Cell* **39**, 302–315 (2016).
5.　Zhang, Z., Zwick, S., Loew, E., Grimley, J. S. & Ramanathan, S. Embryo geometry drives
formation of robust signaling gradients through receptor localization. (2018) doi:10.1101/491290.
6.　Heemskerk, I. *et al.* Rapid changes in morphogen concentration control self-organized
patterning in human embryonic stem cells. *Elife* **8**, (2019).
7.　Massey, J. *et al.* WNT ligands stimulate transient signaling in human pluripotent cells and
synergize with TGF-β ligands to stimulate sustained signaling during differentiation. *bioRxiv* 406306
(2018) doi:10.1101/406306.
8.　Yang, R. *et al.* Amnion signals are essential for mesoderm formation in primates. *Nat Commun*
12, 5126 (2021).
9.　Ornitz, D. M. & Itoh, N. The Fibroblast Growth Factor signaling pathway. *Wiley Interdiscip Rev*
Dev Biol **4**, 215–266 (2015).
10.　LaBonne, C. & Whitman, M. Mesoderm induction by activin requires FGF-mediated intracellular
signals. *Development* **120**, 463–472 (1994).
11.　Krens, S. F. G., Corredor-Adámez, M., He, S., Snaar-Jagalska, B. E. & Spaink, H. P. ERK1 and ERK2
MAPK are key regulators of distinct gene sets in zebrafish embryogenesis. *BMC Genomics* **9**, 196
(2008).
12.　Yao, Y. *et al.* Extracellular signal-regulated kinase 2 is necessary for mesoderm differentiation.
Proc Natl Acad Sci U S A **100**, 12759–12764 (2003).
13.　Mitrani, E., Gruenbaum, Y., Shohat, H. & Ziv, T. Fibroblast growth factor during mesoderm
induction in the early chick embryo. *Development* **109**, 387–393 (1990).
14.　Corson, L. B., Yamanaka, Y., Lai, K.-M. V. & Rossant, J. Spatial and temporal patterns of ERK
signaling during mouse embryogenesis. *Development* **130**, 4527–4537 (2003).
15.　Morgani, S. M. & Hadjantonakis, A.-K. Signaling regulation during gastrulation: Insights from
mouse embryos and *in vitro* systems. in *Current Topics in Developmental Biology* (eds Small, S. &
Briscoe, J.) vol. 137 391–431 (Academic Press, 2020).
16.　De Simone, A. *et al.* Control of osteoblast regeneration by a train of Erk activity waves. *Nature*
590, 129–133 (2021).
17.　Marshall, C. J. Specificity of receptor tyrosine kinase signaling: transient versus sustained
extracellular signal-regulated kinase activation. *Cell* **80**, 179–185 (1995).
18.　Ender, P. *et al.* Spatiotemporal control of ERK pulse frequency coordinates fate decisions during
mammary acinar morphogenesis. *Dev Cell* **57**, 2153-2167.e6 (2022).

19.   Aoki, K. *et al.* Propagating Wave of ERK Activation Orients Collective Cell Migration. *Dev Cell* **43**,
305-317.e5 (2017).
20.   Ciruna, B. & Rossant, J. FGF Signaling Regulates Mesoderm Cell Fate Specification and
Morphogenetic Movement at the Primitive Streak. *Dev. Cell* **1**, 37–49 (2001).
21.   Sun, X., Meyers, E. N., Lewandoski, M. & Martin, G. R. Targeted disruption of Fgf8 causes failure
of cell migration in the gastrulating mouse embryo. *Genes Dev.* **13**, 1834–1846 (1999).
22.   Fletcher, R. B. & Harland, R. M. The role of FGF signaling in the establishment and maintenance
of mesodermal gene expression in Xenopus. *Dev Dyn* **237**, 1243–1254 (2008).
23.   Yang, X., Dormann, D., Münsterberg, A. E. & Weijer, C. J. Cell Movement Patterns during
Gastrulation in the Chick Are Controlled by Positive and Negative Chemotaxis Mediated by FGF4 and
FGF8. *Developmental Cell* **3**, 425–437 (2002).
24.   Fürthauer, M., Reifers, F., Brand, M., Thisse, B. & Thisse, C. sprouty4 acts in vivo as a feedback-
induced antagonist of FGF signaling in zebrafish. *Development* **128**, 2175–2186 (2001).
25.   Tyser, R. C. V. *et al.* Single-cell transcriptomic characterization of a gastrulating human embryo.
Nature **600**, 285–289 (2021).
26.   Zhai, J. *et al.* Primate gastrulation and early organogenesis at single-cell resolution. *Nature* **612**,
732–738 (2022).
27.   Deng, C. X. *et al.* Murine FGFR-1 is required for early postimplantation growth and axial
organization. *Genes Dev* **8**, 3045–3057 (1994).
28.   Yamaguchi, T. P., Harpal, K., Henkemeyer, M. & Rossant, J. fgfr-1 is required for embryonic
growth and mesodermal patterning during mouse gastrulation. *Genes Dev* **8**, 3032–3044 (1994).
29.   Ciruna, B. G., Schwartz, L., Harpal, K., Yamaguchi, T. P. & Rossant, J. Chimeric analysis of
fibroblast growth factor receptor-1 (Fgfr1) function: a role for FGFR1 in morphogenetic movement
through the primitive streak. *Development* **124**, 2829–2841 (1997).
30.   García-García, M. J. & Anderson, K. V. Essential role of glycosaminoglycans in Fgf signaling
during mouse gastrulation. *Cell* **114**, 727–737 (2003).
31.   Oki, S., Kitajima, K. & Meno, C. Dissecting the role of Fgf signaling during gastrulation and left-
right axis formation in mouse embryos using chemical inhibitors. *Dev Dyn* **239**, 1768–1778 (2010).
32.   Feldman, B., Poueymirou, W., Papaioannou, V. E., DeChiara, T. M. & Goldfarb, M. Requirement
of FGF-4 for postimplantation mouse development. *Science* **267**, 246–249 (1995).
33.   Warmflash, A., Sorre, B., Etoc, F., Siggia, E. D. & Brivanlou, A. H. A method to recapitulate early
embryonic spatial patterning in human embryonic stem cells. *Nat. Methods* **11**, 847–854 (2014).
34.   Jo, K. *et al.* Efficient differentiation of human primordial germ cells through geometric control
reveals a key role for Nodal signaling. *Elife* **11**, e72811 (2022).
35.   Minn, K. T. *et al.* High-resolution transcriptional and morphogenetic profiling of cells from
micropatterned human ESC gastruloid cultures. *Elife* **9**, (2020).
36.   Minn, K. T., Dietmann, S., Waye, S. E., Morris, S. A. & Solnica-Krezel, L. Gene expression
dynamics underlying cell fate emergence in 2D micropatterned human embryonic stem cell gastruloids.
Stem Cell Reports **16**, 1210–1227 (2021).
37.   Underhill, E. J. & Toettcher, J. E. Control of gastruloid patterning and morphogenesis by the Erk
and Akt signaling pathways. *Development* **150**, dev201663 (2023).
38.   Loh, K. M. *et al.* Mapping the Pairwise Choices Leading from Pluripotency to Human Bone,
Heart, and Other Mesoderm Cell Types. *Cell* **166**, 451–467 (2016).
39.   Moon, K. R. *et al.* Visualizing structure and transitions in high-dimensional biological data. *Nat.
Biotechnol.* **37**, 1482–1492 (2019).

40.	Neben, C. L., Lo, M., Jura, N. & Klein, O. D. Feedback regulation of RTK signaling in development.
Dev Biol **447**, 71–89 (2019).
41.	Tyser, R. C. V. & Srinivas, S. Recent advances in understanding cell types during human
gastrulation. *Seminars in Cell & Developmental Biology* **131**, 35–43 (2022).
42.	Chen, B. *et al.* Extended culture of 2D gastruloids to model human mesoderm development.
bioRxiv 2024.03.21.585753 (2024) doi:10.1101/2024.03.21.585753.
43.	Khoa, L. T. P. *et al.* Visualization of the Epiblast and Visceral Endodermal Cells Using Fgf5-P2A-
Venus BAC Transgenic Mice and Epiblast Stem Cells. *PLOS ONE* **11**, e0159246 (2016).
44.	Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early organogenesis.
Nature **566**, 490–495 (2019).
45.	Morgani, S. M. *et al.* A Sprouty4 reporter to monitor FGF/ERK signaling activity in ESCs and
mice. *Developmental Biology* **441**, 104–126 (2018).
46.	Xu, D. & Esko, J. D. Demystifying heparan sulfate-protein interactions. *Annu Rev Biochem* **83**,
129–157 (2014).
47.	Habuchi, H. & Kimata, K. Functions of Heparan Sulfate Proteoglycans in Morphogenesis. in
Experimental Glycoscience 303–308 (Springer, Tokyo, 2008). doi:10.1007/978-4-431-77922-3_72.
48.	Yu, S. R. *et al.* Fgf8 morphogen gradient forms by a source-sink mechanism with freely diffusing
molecules. *Nature* **461**, 533–536 (2009).
49.	Benington, L., Rajan, G., Locher, C. & Lim, L. Y. Fibroblast Growth Factor 2-A Review of
Stabilisation Approaches for Clinical Applications. *Pharmaceutics* **12**, 508 (2020).
50.	Teague, S. *et al.* Time-integrated BMP signaling determines fate in a stem cell model for early
human development. *Nat Commun* **15**, 1471 (2024).
51.	Ortiz-Salazar, M. A., Camacho-Aguilar, E. & Warmflash, A. Endogenous Nodal switches Wnt
interpretation from posteriorization to germ layer differentiation in geometrically constrained human
pluripotent cells. *bioRxiv* 2024.03.13.584912 (2024) doi:10.1101/2024.03.13.584912.
52.	Gurdon, J. B. A community effect in animal development. *Nature* **336**, 772–774 (1988).
53.	Nemashkalo, A., Ruzo, A., Heemskerk, I. & Warmflash, A. Morphogen and community effects
determine cell fates in response to BMP4 signaling in human embryonic stem cells. *Development* **144**,
3042–3053 (2017).
54.	Gattiglio, M., Protzek, M. & Schröter, C. Population-level antagonism between FGF and BMP
signaling steers mesoderm differentiation in embryonic stem cells. *Biology Open* **12**, bio059941 (2023).
55.	Liu, L. *et al.* Nodal is a short-range morphogen with activity that spreads through a relay
mechanism in human gastruloids. *Nat Commun* **13**, 497 (2022).
56.	Isaacs, H. V., Pownall, M. E. & Slack, J. M. eFGF is expressed in the dorsal midline of Xenopus
laevis. *Int J Dev Biol* **39**, 575–579 (1995).
57.	Christen, B. & Slack, J. M. FGF-8 is associated with anteroposterior patterning and limb
regeneration in Xenopus. *Dev Biol* **192**, 455–466 (1997).
58.	Hardy, K. M., Yatskievych, T. A., Konieczka, J., Bobbs, A. S. & Antin, P. B. FGF signalling through
RAS/MAPK and PI3K pathways regulates cell movement and gene expression in the chicken primitive
streak without affecting E-cadherin expression. *BMC Dev. Biol.* **11**, 20–17 (2011).
59.	Isaacs, H. V., Pownall, M. E. & Slack, J. M. eFGF regulates Xbra expression during Xenopus
gastrulation. *The EMBO Journal* **13**, 4469–4481 (1994).
60.	Bulger, E. A., Muncie-Vasic, I., Libby, A. R. G., McDevitt, T. C. & Bruneau, B. G. TBXT dose
sensitivity and the decoupling of nascent mesoderm specification from EMT progression in 2D human
gastruloids. *Development* **151**, dev202516 (2024).

61.   Evans, A. L. *et al.* Genomic Targets of Brachyury (T) in Differentiating Mouse Embryonic Stem
Cells. *PLOS ONE* **7**, e33346 (2012).

62.   Nie, Y., Walsh, P., Clarke, D. L., Rowley, J. A. & Fellner, T. Scalable passaging of adherent human
pluripotent stem cells. *PLoS ONE* **9**, e88012 (2014).

63.   Freeburne, E. *et al.* Spatial Single Cell Analysis of Proteins in 2D Human Gastruloids Using
Iterative Immunofluorescence. *Curr Protoc* **3**, e915 (2023).

64.   Sommer, C., Straehle, C., Kothe, U. & Hamprecht, F. A. Ilastik: Interactive learning and
segmentation toolkit. in *2011 8th IEEE International Symposium on Biomedical Imaging (ISBI 2011)*
230–233 (IEEE, 2011). doi:10.1109/ISBI.2011.5872394.

65.   Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nat. Methods* **18**, 100–106 (2021).

66.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol* **19**, 15 (2018).

67.   Gayoso, A. *et al.* A Python library for probabilistic analysis of single-cell omics data. *Nat
Biotechnol* **40**, 163–166 (2022).

68.   van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell*
174, 716-729.e27 (2018).

69.   Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics.
BMC Genomics **19**, 477 (2018).

70.   Van den Berge, K. *et al.* Trajectory-based differential expression analysis for single-cell
sequencing data. *Nat Commun* **11**, 1201 (2020).

71.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-
aware quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

72.   Eswarakumar, V. P., Lax, I. & Schlessinger, J. Cellular signaling by fibroblast growth factor
receptors. *Cytokine Growth Factor Rev.* **16**, 139–149 (2005).

Supplementary Figures

The length scale of the ERK signaling pattern is independent of colony size

a

b

All primitive streak markers are lost upon FGF/ERK inhibition

Rescue of FGFRi with dox-SOS^cat

FGF/ERK affects cell number primarily through proliferation

c **d** **e** **f**

There is a *direct* requirement for FGF/ERK signaling in primitive streak-like differentiation

g **h**

i

Supplementary Figure 1. a) Stains and quantification for pERK and TBXT in smaller colonies, diameter from left to right:
500um, 350um, 250um. **b)** Staining for three primitive streak markers with or with FGFRi or MEKi. **c)** DAPI stain corresponding
to Fig. 1d. **d)** Radial intensity profile for TBXT corresponding to Fig.1d. **e**) BrdU and cleaved Cas3 stainings after MEK or
FGFR inhibition. **f)** Quantification of conditions in e. **g)** Quantification of conditions in Fig.1g for five images each, N indicates
total number of cells. **h**) Low density differentiation with and without MEK and FGFR inhibition. **i**) pERK and TBXT stains and
quantification for different doses of MEK inhibitor shows low doses effectively block differentiation with minimal impact on cell
number.

Supplementary Figure 2. a) PHATE projection of scRNA-seq data colored for sample. **b,c)** Expression of canonical FGF ligands
across different cell types in 2D gastruloids (b) and human CS7 embryo (c). Expression across clusters is normalized by row,
while total expression is shown in a separate column to the directly to left. The leftmost column indicates differential expression
between PS-like and pluripotent cells. The color scale of (differential) expression is cut off for better contrast. **d)** Threshold-
based annotation (left) for human, monkey, and mouse embryos used in interspecies comparison (Fig. 2f), compared to original
annotation (right). **e)** Interspecies comparison as in Fig. 2f but for nascent mesoderm and PS-LC + nascent mesoderm. **f)** Bulk
RNA-seq data by Loh et al[38] for expression of different FGFs in directed differentiation to mid primitive streak (MPS) or anterior
primitive streak (APS) relative the human embryonic stem cells (hESCs). **g)** Expression of genes involved in FGF signaling.

The phosphorylated ERK pattern reflects receptor activation

FGFR1-IIIc is the dominant isoform across the colony

Supplementary Figure 3. a) DAPI image corresponding to Fig. 3a. **b-c)** pFGFR1 (b) or pERK (c) and TBXT stains with and without continuous MEKi treatment. **d)** DAPI image corresponding to Fig. 3b. **e)** Design of qPCR primers to detect FGFR1 isoforms, diagram adapted from Eswarakumar et al[72]. **f)** Primer sequence and properties. **g)** Amplicon sizes match prediction. Scale bars 50um.

886 **Supplementary Figure 4. a,b)** Heparan sulfate (HS) stain (a) and radial intensity profile (b). Scale bar 50um. Error bands
887 represent standard deviation over N=4 colonies.

888
889
890 **Supplementary Figure 5. a)** DAPI image corresponding to Fig. 5a. **b)** Pseudotime dynamics from Fig.5b with each gene
891 normalized individually. **c)** Correlations between FGFs and TBXT, TBX6 colored for clusters, with cluster colors matching Fig.
892 2d. **d-f)** DAPI images and overlays including DAPI corresponding to Fig.5c-e, respectively. **g)** Combined FISH for FGF4 and
893 FGF8. **h)** Scatterplots of FGF4, FGF8, and FGF17 mRNA versus TBXT colored for cell density.
894
895

896
897
898 **Supplementary Figure 6. a)** Additional channels for the colony shown in Fig.5g. **b)** FGF4 shRNA in ESI17 cells. **c-d)** FGF4
899 siRNA (c) and shRNA (d) in PGP1 cells. **e-f)** additional channels for Fig.5h-i. **g**) FGF17 shRNA in ESI17 cells. **h-i)** FGF17 siRNA
900 (h) and shRNA (i) in PGP1 cells. **j)** additional channels for the TBX6 ctrl and FGF17 shRNA in Fig.5j. **k)** FGF4 + FG17 shRNA
901 double knockdown (stains from multiple colonies).