Endogenous FGFs drive ERK-dependent cell fate patterning in 2D human gastruloids

Kyoung Jo[1]*, Zong-Yuan Liu[1]*, Gauri Patel[4], Zhiyuan Yu[2], LiAng Yao[1], Seth Teague[4], Craig Johnson[1], Jason Spence[1,3-5], Idse Heemskerk[1-4,6^]

1. Department of Cell and Developmental Biology, University of Michigan Medical School, Ann Arbor, Michigan
2. Department of Computational Medicine and Bioinformatics, University of Michigan Medical School, Ann Arbor, Michigan
3. Center for Cell Plasticity and Organ Design, University of Michigan Medical School, Ann Arbor, Michigan
4. Department of Biomedical Engineering, University of Michigan, Ann Arbor, Michigan
5. Department of Internal Medicine, Gastroenterology, University of Michigan Medical School, Ann Arbor, Michigan
6. Department of Physics, University of Michigan, Ann Arbor, Michigan

* equal contributions
^ for correspondence: iheemske@umich.edu

Abstract

The role of FGF is the least understood of the morphogens driving mammalian gastrulation. Here we investigated the function of FGF in a stem cell model for human gastrulation known as a 2D gastruloid. We found a ring of FGF-dependent ERK activity that closely follows the emergence of primitive streak (PS)-like cells but expands further inward. We showed that this ERK activity pattern is required for PS-like differentiation and that loss of PS-like cells upon FGF receptor inhibition can be rescued by directly activating ERK. We further demonstrated that the ERK-ring depends on localized activation of basally localized FGF receptors (FGFR) by endogenous FGF gradients. We confirm and extend previous studies in analyzing expression of FGF pathway components, showing the main receptor to be FGFR1 and the key ligands FGF2/4/17, similar to the human and monkey embryo but different from the mouse. In situ hybridization and scRNA-seq revealed that *FGF4* and *FGF17* expression colocalize with the PS marker TBXT but only *FGF17* is maintained in nascent mesoderm and endoderm. FGF4 and FGF17 reduction both reduced ERK activity and differentiation to PS-like cells and their derivatives, indicating overlapping function. Thus, we have identified a previously unknown role for FGF-dependent ERK signaling in 2D gastruloids and possibly the human embryo, driven by a mechanism where FGF4 and FGF17 signal through basally localized FGFR1 to induce PS-like cells.

Introduction

During gastrulation the three germ layers are established, and the body plan is laid out. In amniotes this involves the formation of the primitive streak on the posterior side of the embryo from which cells ingress beneath the epiblast and migrate to form the mesodermal and endodermal layers. An important role in gastrulation is played by evolutionarily conserved morphogens from the BMP, Wnt, Nodal, and FGF families[1]. The function of the first three signaling pathways is comparatively well understood: in both mouse and human a transcriptional hierarchy between BMP4, Wnt3, and Nodal plays a central role in inducing dynamic signaling gradients along the anterior posterior axis that control cell fate patterning[1–3]. For BMP and Nodal, basal receptor localization in the epiblast is crucial to form signaling activity gradients[3–7]. Wnt and Nodal signaling are directly required for mesoderm and endoderm differentiation, while the balance between BMP and Nodal signaling is important for patterning the germ layers along the body axes[8].

In contrast, the role of FGF in mammalian and especially human gastrulation has been more elusive. Secreted FGFs signal through FGF receptor tyrosine kinases (FGFRs) in a heparan sulfate proteoglycan (HSPG)-dependent manner to activate several signaling pathways including MAPK/ERK, PI3K, and PLCγ[9]. There is evidence for a conserved requirement for FGF/ERK signaling in mesoderm formation across vertebrate model organisms[10–13]. However, it is unknown if there are FGF- and phosphorylated ERK gradients across the mammalian primitive streak[14,15], whether there are functionally relevant ERK signaling dynamics like in other contexts, or whether receptor localization is important for establishing signaling gradients[16–19]. It is also unclear to what extent the requirement in mesoderm induction is direct, or indirect by modulating other signals[14,15,20]. In addition, several FGFs are co-expressed and the roles of specific FGFs are uncertain. Moreover, there may be significant interspecies differences in this respect. Notably, FGF8 is the only FGF which has been shown to be required for gastrulation in the mouse[21], and its requirement in gastrulation appears to be conserved across vertebrate model organisms[22–24], yet FGF8 is not expressed in primate embryos[25,26].

Mutants for FGF8, FGFR1, HSPG synthesis, and ERK2, as well as pharmacological inhibition of MEK and FGFR give rise to similar phenotypes in mouse gastrulation, suggesting that FGFR1 and FGF8 are the only FGF ligand-receptor pair required for gastrulation and that these act through the MAPK/ERK pathway[12,20,21,27–31]. FGF4 expression during gastrulation is also lost in FGF8 mutants, leaving the possibility of a role for FGF4, but this remains unknown as FGF4 is also required in the blastocyst and mutants therefore do not reach gastrulation[32]. In all mutants and for pharmacological inhibition of FGF/ERK signaling, expression of the primitive streak marker

brachyury (TBXT, also BRA, or T in mouse) is lost or severely diminished in the posterior streak and mesoderm fails to migrate away, piling up in this region. In addition, TBX6 expression is lost completely, and no lateral or paraxial mesoderm are formed. In contrast, TBXT expression in the anterior streak and extraembryonic mesoderm is not dependent on FGF/ERK signaling; extraembryonic mesoderm development is normal, and the axial mesoderm is expanded.

In vitro models offer an opportunity to disentangle the complexity of FGF signaling in human development. Here, we explore the role of FGF/ERK signaling in micropatterned human pluripotent stem cells treated with BMP4, also known as 2D human gastruloids. These give rise to concentric rings of different cell types associated with gastrulation after 2 days of differentiation. Pluripotent cells in the center are surrounded by a primitive streak-like ring, which itself is surrounded by primordial germ cells and amnion-like cells on the colony edge[33–35]. Due to its simplicity and reproducibility, this system has proven powerful in deciphering the mechanisms underlying aspects of human gastrulation. It was used to show that BMP-Wnt-Nodal hierarchy that controls primitive streak formation during mouse gastrulation is preserved in human while providing a range of new insights into how these signals function[2,3]. For example, the crucial role for BMP and Nodal receptor localization in patterning was discovered in 2D gastruloids and later confirmed in the mouse embryo. This system also revealed that BMP, Wnt, and Nodal signaling patterns are highly dynamic[3–7]. It therefore also provides a promising approach to determine how FGF may function in human gastrulation.

In the 2D human gastruloid model, we show that a ring of phosphorylated ERK forms in an FGF-dependent manner and expands largely in lockstep with the formation of primitive streak-like cells but extends further inward. We demonstrate that this pattern of ERK activity is due to localized FGF receptor (FGFR) activation that requires basal FGFR localization. We then show that exogenous FGF2 in the media is required for pattern formation to occur but can be removed well before the phosphorylated (active) ERK (pERK) and PS-like rings appear, suggesting the pERK ring instead depends on endogenous FGF gradients. Using single cell RNA-sequencing, we determine the expression of FGF pathway components including FGFR1 as the main receptor and FGF2, FGF4, and FGF17 as the main ligands. Our analysis corroborates previous studies describing expression of FGF2 and FGF17 in human embryos and gastruloids but for the first time recognizes a possible role for FGF4 and includes a broad analysis of genes involved in FGF signaling modulation[25,36]. We find that the FGF4 and FGF17 expression are restricted to the PS-like ring and that reduction of FGF4 significantly reduces PS-like differentiation, while FGF17 reduction leads to a smaller reduction in PS-like cells and their derivatives. In summary, we have greatly advanced our understanding of how FGFs function in a model for human gastrulation.

Results

FGF/ERK signaling is required for primitive streak-like differentiation.

We previously observed that pharmacological inhibition of MEK (the kinase upstream of ERK) or FGF receptors leads to a loss of differentiation to primitive streak-like and primordial germ cell-like cells, suggesting a crucial role for FGF/ERK signaling[34]. Moreover, a ring of active ERK (pERK) has been observed in 2D gastruloids but its connection to FGF signaling or differentiation remains unclear[33]. Therefore, we wanted to investigate the role of FGF/ERK signaling further.

To determine the dynamics of pERK and relate it to PS-like differentiation, we first performed time series immunofluorescence co-staining for pERK and the PS marker TBXT after BMP4 treatment. This revealed the pERK ring first emerges around 24h, around the same time as TBXT expression, and gradually gets brighter and wider (Fig.1a-c). By 42h the outer edge of the ERK ring coincided with TBXT expression, while the inner edge was less sharp and extended further towards the center than TBXT, reminiscent of the domains of WNT and NODAL signaling that extend further in than the PS-like region[3,6] (Fig.1a, Supp. Fig.1ab). MEK or FGFR inhibition for 30 minutes after 41.5h eliminated ERK activity, suggesting ERK activity depends entirely on FGF and that continuous FGF signaling is necessary to maintain the ERK signaling pattern (Fig.1bc). Inhibition for the full duration of differentiation eliminated TBXT expression as well as other PS markers (Fig.1bc, Supp. Fig.1c). However, we were able to rescue TBXT expression by directly and uniformly activating ERK (Fig. 1d-f, Supp. Fig.1de) in cells stably expressing the doxycycline-inducible membrane-targeted catalytic domain of SOS (dox-SOS^{cat})[37]. Although TBXT expression was rescued by ERK activation, reduced proliferation caused by FGFR inhibition was not, indicating proliferation may be regulated by additional downstream pathways (Fig. 1d). Furthermore, despite uniform induced ERK activity, TBXT expression remained excluded from the center, showing that ERK activation alone is not sufficient for TBXT induction. However, TBXT levels were much higher in the rescue (Supp. Fig. 1E), possibly reflecting the much higher level of ERK activity induced by dox-SOS^{cat}. Altogether, these data show the formation of a PS-like ring is dependent on FGF signaling through ERK.

This raised several questions. First, is FGF/ERK signaling required directly for PS-like differentiation, or
does it act indirectly, for example, by controlling the expression of Wnt or Nodal which are both required for PS-like
differentiation? These possibilities are not mutually exclusive. Second, which are the FGFs activating the ERK
pathway? The pathway could be activated by exogenous FGF2 present in the pluripotency medium mTeSR1 or by
endogenously expressed ligands. Third, how is ERK activity restricted to a ring?
To determine if there is a direct requirement of FGF/ERK signaling and whether this depends on exogenous
FGF, we performed differentiation to PS-like cells in standard culture by exogenous activation of Wnt and Nodal
signaling, bypassing the need to form endogenous Wnt and Nodal gradient as in 2D gastruloids[38]. We used a
minimal base medium (E6) that did not include exogenous FGF or other growth factors. To rule out any additional
requirement for endogenous Nodal and Wnt which may be FGF/ERK-dependent, we used Nodal knockout cells[3]
treated with the Wnt secretion inhibitor IWP2, thereby eliminating any endogenous activity of these pathways. We
also inhibited BMP receptors using LDN193189 to rule out any indirect effect through endogenous BMP.
Under these conditions, with exogenous Wnt and Nodal activation for 24h in the absence of exogenous
FGF, cells efficiently differentiated to TBXT+ PS-like cells (Fig.1g, Supp. Fig.1f). These PS-like cells had high active
ERK levels just as in the 2D gastruloids. Inhibition of either FGFR or MEK abrogated ERK activity and TBXT
expression while FGFR inhibition also severely reduced proliferation. In both cases significant expression of the
pluripotency and ectoderm marker SOX2 was maintained, although SOX2 levels were much lower with FGFR
inhibition (Supp. Fig. 1f). Given that FGF/ERK is known to be involved in pluripotency maintenance, this suggests
inhibition may not be complete and that much lower ERK activity is required for pluripotency maintenance than PS-
like differentiation, consistent with lower ERK activity in the pluripotent center of gastruloids. The greater loss of
SOX2 with inhibition of FGFR than MEK could reflect more efficient inhibition of ERK signaling but could also
indicate that other pathways downstream of FGFR, such as PI3K, play a role in maintaining SOX2 expression.
Altogether, these results show that endogenous FGF expression is both necessary and sufficient for PS-like
differentiation by exogenous Wnt and Nodal stimulation and demonstrate a direct requirement for ERK signaling in
PS-like differentiation.
(Next page)
Figure 1. a) Immunofluorescence stainings for pERK and TBXT and their radial intensity profiles in 2D gastruloids at different
times. Error bars in graphs are standard deviations over N=4 colonies. **b,c)** Effect FGF receptor inhibition (FGFRi, b) or MEK
inhibition (MEKi, c) on pERK and TBXT in colonies fixed at 42h after BMP treatment. Inhibition for either 30 minutes at 41.5h
(rounded to 42 in figure) or throughout differentiation. **d)** BMP only control and FGFRi inhibition in cells expressing doxycycline-
inducible SOS^cat with and without addition of doxycycline. **e-f)** Cell numbers (e) and radial profiles of TBXT positive cells for
conditions in (d). **g)** PS marker TBXT, pERK and pluripotency marker SOX2 in Nodal knockout cells with inhibitors of Wnt
secretion (WntSeci) and BMP receptors (BMPRi) with or without MEKi or FGFRi. All scale bars 50um.

A ring of phosporylated ERK first appears around 24h and expands inward over time

a

	16h	24h	28h	32h	36h	40h	44h	48h

pERK

TBXT

overlay + DAPI

intensity (a.u.) — edge distance (μm)

Phospho-ERK depends on FGF and is required for formation of a primitive streak-like ring in 2D gastruloids

FGFR inhibition

b ctrl / 30 min @ 42h / continuous — TBXT — pERK — overlay + DAPI

MEK inhibition

c ctrl / 30 min @ 42h / continuous — TBXT — pERK — overlay + DAPI

Rescue of FGFR inhibition by direct ERK activation with dox-SOS^{cat}

d ctrl / FGFRi - dox / FGFRi + dox — TBXT — pERK — overlay + DAPI

e x10³ cell number — cells per colony — BMP4, +FGFRi -dox +dox

f TBXT — positive fraction — edge distance (μm) — BMP4, +FGFRi, +FGFRi +dox

There is a *direct* requirement for FGF/ERK signaling in primitive streak-like differentiation

g

	DAPI	TBXT	pERK	SOX2	overlay
NodalKO + BMPRi + WNTSeci + CHIR + Activin					
+ MEKi					
+ FGFRi					

Figure 2. a) Schematic of the FGF/ERK signaling pathway. **b)** Distribution of pERK levels in TBXT+ cells versus ISL1+ and other cells based on data and thresholds in Supp. Fig. 2b. **c)** Expression of key markers in PHATE[39] projection of scRNA-seq data for 42h colony. **d)** Cell type annotation. PS-LC: primitive streak-like cells, NasMe: nascent mesoderm, AE-LC: amniotic ectoderm-like cells: PGC-LC: primordial germ cell-like cells, endo: definitive endoderm. **e)** Expression of canonical FGF ligands (log transformed and smoothened). **f)** Comparison of FGF ligand expression in 2D human gastruloids to human, monkey, and mouse embryo, normalized within each type of sample. **g-i)** Expression of several genes involved in FGF signaling with strong differential expression between PS-like cells or nascent mesoderm and pluripotent cells, split by gene category.

Single cell RNA-sequence reveals differentially expressed FGF pathway components.

The above results suggest endogenous FGF expression may contribute to elevated ERK signaling in the PS-like ring. However, FGF/ERK signaling may be modulated at many levels (Fig. 2a). A spatial pattern of ERK activation could reflect a concentration gradient of FGF. Alternatively, differential expression of receptors and HSPG-modifying enzymes could lead to spatially patterned receptor activation and downstream ERK activity even in the absence of an FGF gradient. Even if receptor activation is uniform, downstream ERK activity could be spatially patterned by downstream negative regulators of ERK activity such as Sprouty-family proteins (SPRY) and dual-specificity phosphatases (DUSPs)[40].

To understand which genes modulating FGF/ERK signaling could be responsible for the elevated ERK signaling in TBXT positive cells and required for TBXT expression (Fig.1, Fig. 2b), we extended previous work using single cell RNA-sequencing to determine expression of FGF pathway components in human gastrulation[25,35,36]. TBXT-positive PS-like cells give rise to nascent mesoderm expressing TBX6 as well as definitive endoderm expressing SOX17 and FOXA2, both of which may still express TBXT at 42h[41]. To understand how TBXT expression is initiated in an FGF-dependent manner, we focused on differential expression in the PS-LC relative to pluripotent cells, reasoning that FGF modulation specific to nascent mesoderm or endoderm is a consequence rather than a cause of PS-LC differentiation. We identified different cell types based on marker genes with the PS-like cells defined as TBXT+MIXL1+POU5F1+TBX6-SOX17-FOXA2- (Fig. 2cd, Supp. Table 4). For our analysis, we combined one previously published dataset and one new replicate sample at 42h (Supp. Fig. 2a). We then analyzed expression of different categories of FGF/ERK regulators.

We found four FGF ligands with significant expression: FGF2, 4, 8, and 17 (Fig. 2e, Supp. Fig. 2b). We then compared the expression of these ligands in PS-like cells to reference data of CS7 human, CS8 monkey, and E6.5-E7.5 mouse gastrulation (Fig. 2f, Supp. Fig. 2cd). We also included FGF3 and FGF5, which are expressed during mouse gastrulation[32,42,43]. High expression of FGF4 and FGF17 and low expression of FGF8 was consistent between human, monkey, and 2D gastruloid. FGF2 expression was also high in the PS-like populations in 2D gastruloids and the CS8 monkey embryo, but much lower in for human embryo data. Furthermore, the monkey and human embryos expressed FGF3, which was minimally expressed in gastruloids. A similar comparison in nascent mesoderm showed much lower FGF2/4 and much higher FGF17 compared to PS across all primate samples (Supp. Fig. 2e). Expression in 2D gastruloids was consistent with published RNA-seq data for hPSC-derived PS-like cells,

which expressed high FGF2 and FGF4, some FGF8 and FGF17, and no FGF3 (Supp. Fig. 2f). Comparison with
mouse confirmed significant differences from primates, with high expression of FGF5, 8, much lower expression of
FGF4,17, and no expression of FGF2 in the mouse[21,42,43].
Of the FGF receptors, only FGFR1 and FGFR4 were strongly expressed, with FGFR1 expression about 5-
fold higher (Fig. 2g, Supp. Fig. 2g). Both receptors showed graded expression that decreased from pluripotent to
PS-like cells and much lower expression in amnion-like cells (Fig. 2g). Therefore, elevated ERK activity in PS-like
cells cannot be accounted for by overall receptor expression. Differential expression of genes regulating heparin-
sulfate proteoglycans which modulate FGF receptor binding was mild except for glypicans 3 and 4 which were
downregulated in PS-like cells relative to pluripotent cells (Fig. 2h, Supp. Fig. 2g). In contrast, several intracellular
negative regulators of ERK signaling were strongly differentially expressed between cell clusters, including SPRY1
and PEBP1 which were significantly higher in pluripotent cells (Fig. 2i, Supp. Fig. 2g). Surprisingly, however,
expression of SPRY4 was not elevated in PS-like cells where ERK signaling is high, suggesting that transcription
of SPRY4 may not be a good readout of ERK activity in human hPSCs like it is in the mouse[44]. On the other hand,
expression of the ERK negative feedback regulator IL17RD (SEF) does seem to approximately reflect expected
ERK activity in each population. Components of the core ERK pathway including ERK itself were highly expressed
across all clusters (Supp. Fig. 2g).
Thus, while scRNA-seq identified specific genes involved, it is consistent with several non-exclusive
mechanisms for generating an ERK pattern. For example, uniform FGF2 could form an ERK ring by equally
activating receptors across the colony but with ERK response reduced by elevated expression of negative
regulators in the pluripotent center. Alternatively, based on gene expression we cannot rule out HSPGs increasing
receptor binding specifically in the ring. Finally, localized expression of FGF4 or FGF17 in the PS-like ring could
lead to a concentration gradient responsible for increased receptor activation.
240
241 **The phosphorylated ERK pattern is due to differential activation of FGF receptors.**
242
To distinguish between the different possibilities, we asked whether the spatial pattern of pERK signaling is
controlled primarily below, at, or above the level of the receptors. To address this, we first visualized phosphorylated
FGFR1 (pFGFR1) and found its spatial pattern to strongly resembles that of pERK (Fig.3a, Supp. Fig.3a). As
expected, pFGFR1 was lost upon brief FGFR inhibition (Fig.3a, Supp. Fig.3a). In contrast, after MEK inhibition, high
pFGFR1 remained while pERK was greatly reduced (Supp. Fig.3bc). This suggests the spatial pattern of ERK
reflects a spatial pattern of receptor activity.
The spatial pattern in receptor activity could either be due to differential receptor expression or differential
receptor activation. There was no significant upregulation of FGF receptors in PS-like cells according to our single
cell RNA-sequencing data (Fig 2e). In fact, the most highly expressed receptor, FGFR1, was elevated in pluripotent
cells. This suggests the ERK activity pattern is not due to a receptor expression pattern with two caveats. First,
mRNA expression may not reflect protein expression. Second, multiple isoforms exist for FGF receptors 1-3 (but
not 4) that have different affinity for FGF ligands, leaving open the possibility that PS-like cells could express a
different FGFR1 isoform than pluripotent cells, despite overall FGFR1 expression being approximately uniform.
To determine if the spatial pattern in pFGFR1 could be explained by a spatial pattern of FGFR1 protein, we
performed immunofluorescence staining of FGFR1. Consistent with the scRNA-seq data, and in sharp contrast to
the pFGFR1 stain, FGFR1 protein expression was graded from center to edge with its highest levels in the
pluripotent center, and therefore could not explain the signaling pattern (Fig.3b, Supp. Fig.3d). To determine if
different isoforms of FGFR1 are differentially expressed in PS-like vs. pluripotent cells, we designed qPCR primers
to detect the IIIb and IIIc isoforms of FGFR1 (Supp. Fig. 3e-g). These are the most common isoforms, also known
as the epithelial and mesenchymal isoforms, and therefore appeared the mostly likely candidates for differential
expression. We then analyzed expression of these isoforms in 42h micropatterns versus pluripotent cells and PS-
like cells differentiated directly as in Fig.1d. However, the change in expression between samples was similar for
each isoform, with all isoforms decreasing in 2D gastruloids and directly differentiated PS-like cells relative to
pluripotent cells (Fig. 3c). To corroborate this and also determine absolute FGFR isoform expression in pluripotent
and PS-like cells, we analyzed our single cell RNA-sequencing data and previously published bulk RNA-sequencing
data for isoform expression in these two cell types. This revealed that for both pluripotent and PS-like cells FGFR
IIIc is the dominant isoform with on average almost 900-fold higher expression than IIIb, while IIIa was not expressed
at all (Fig. 3d). Altogether, these data suggest that spatial patterning in ERK is due to differential activation of FGFR
receptors, either due to FGF concentration gradients or possibly spatially patterned HSPGs.
272
273

The phosphorylated ERK pattern reflects receptor activation

Figure 3. a) phospho-FGFR1 stain with and without 1h FGFRi treatment at 41h, radial intensity profile on the right. **b)** FGFR1 with and without 1h FGFRi treatment at 41h, radial intensity profile on the right. **c)** qPCR data for relative expression of FGF receptor 1 isoforms IIIb, IIIc and total FGFR1 in pluripotent cells, gastruloids at 42h and directly differentiated PS-like cells. **d)** RNA-seq analysis of absolute FGFR1 isoform expression. All scale bars 50 um.

FGFR1-IIIc is the dominant isoform across the colony

FGF receptors are localized basolaterally.

To maintain an endogenous FGF concentration gradient on the apical side of the colony which faces the medium, would require ligands to remain tethered to the surface. However, FGF has been reported to diffuse freely in the intercellular space[45]. Moreover, there is FGF2 present in the medium that would be expected to uniformly activate FGF receptors. The situation is similar for BMP4, which is present uniformly in the medium for 2D gastruloids but does not cause uniform signaling activity[4]. In the case of BMP, that is because receptors are localized basolaterally[4] and tight junctions seal the basolateral side from the medium. Consequently, response to exogenous ligands in the medium is restricted to the colony edge, while endogenous ligands are confined to the basal side and prevented from diffusing away into the medium everywhere but the edge. This was found to be critical for patterning in both 2D gastruloids and the mouse embryo[4,5]. To explain the pERK pattern, we therefore hypothesized FGF receptors were similarly restricted to the basolateral side.

To test our hypothesis, we first starved pluripotent colonies of FGF by maintenance in FGF-free E6 medium for 12h and then treated with FGF2. Consistent with our hypothesis, phospho-FGFR and phosphor-ERK were both restricted to the colony edge (Fig. 4a). To determine if this could be explained by diffusion of exogenous FGF from the medium, we repeated the experiment with micropattern pluripotent cells and added fluorescent dextran with a similar molecular weight as FGF. We observed fluorescent dextran diffusing into the intercellular space from the edge inward over the same distance that ERK signaling was observed (Fig. 4b-d). We then acquired high resolution z-stacks of FGFR1 co-stained with the tight junction protein TJP1 (ZO-1) which revealed the majority of FGFR1 was below the tight junctions (Fig. 4e). To directly test that cells are more responsive to basal FGF than apical FGF, we then performed a transwell assay, where pluripotent cells were grown on filters to enable separate apical and basal stimulation. We found that only basal stimulation with FGF led to a significant increase in pERK (Fig. 4f). Finally, to determine if the center of the micropatterned colony is similarly able to respond to basal FGF stimulation and pERK levels there can be attributed to reduced FGF ligand concentration, we performed a scratch assay. Strong ERK response to FGF2 in the media was seen in the colony center around a scratch, but this response was blocked by FGFR inhibitor (Fig.4gh). If HSPGs rather than FGF ligand were limiting ERK response in the center, we would not have expected strong response around a scratch. We therefore concluded that the pERK pattern is most likely due to a basolateral FGF ligand gradient which has its maximum in the PS-like ring.

Response to exogenous FGF2 treatment is restricted to the colony edge

FGF receptors are localized basolaterally

Figure 4. a) pERK and pFGFR1 response of pluripotent colonies in standard culture grown in FGF-free (E6) medium for 24 hrs and then treated with FGF2 for 30 minutes. **b)** Fluorescent dextran in micropatterned pluripotent colony with quantification. **c)** pERK response in micropatterned pluripotent colony after 42h in E6 with TGFβ and FGF2. **d)** Radial intensity profile of dextran and pERK for conditions in b,c), averaged over 2 colonies. **e)** Cross-section of 2D gastruloid shows FGF receptor 1 predominantly below the tight junction marked by ZO-1. White box marks are magnified in inset e'. **f)** pERK and pFGFR1 response in transwell experiment with cell treated with FGF either apically or basally. **g)** ERK signaling in scratched colonies with or without FGFRi. **h)** Quantification of pERK intensity as a function of distance from the scratch. Error bars represent standard deviation over colonies. Scale bars 50 micron.

Endogenous FGF4 and FGF17 gradients underly the ERK activity pattern.

To identify which FGF ligands are required for PS-like differentiation, we first determined the requirement for exogenous FGF2 in the media. Differentiation of 2D gastruloids in FGF-free medium was severely reduced but restored by addition of FGF2 (Fig. 5a, Supp. Fig. 5a). However, if FGF2 was removed 24h after BMP treatment, differentiation was normal. The fact that removal 2h after BMP treatment significantly reduced differentiation suggests this was not a failure to wash out the FGF2. Furthermore, MEK inhibition at 24h severely reduces TBXT

expression[34] while FGF2 is known to have a half-life on the order of hours[46] and no media changes are performed
during standard differentiation. Altogether this suggests that exogenous FGF2 is required at the time of BMP
treatment but that endogenous FGFs are sufficient for the ERK activity ring and PS-like differentiation that start a
day later.
We reasoned that FGFs whose expression correlates with the presence of PS-like cells are more likely to
be responsible for the observed ERK ring and PS-like differentiation. Our analysis in figure 2 had implicated FGF4
and FGF17 as most strongly expressed in the PS-like cells and nascent mesoderm, respectively (Fig. 2c). However,
it is unclear whether expression of these ligands is a consequence of PS-like differentiation, its cause, or both, in
the case of a feedback loop. To get more information about the relationship between FGF ligands and PS vs.
mesoderm differentiation, we performed pseudotime analysis of our scRNA-seq data (Fig. 5b, Supp. Fig 5b). Along
a pseudotime trajectory from pluripotent cells to nascent mesoderm, FGF2 expression was nearly constant. FGF8
was upregulated prior to TBXT and declined as TBXT was upregulated although its levels always remained low.
FGF4 upregulation approximately coincided with TBXT, but FGF4 began decreasing as TBX6 expression
increased. In contrast, FGF17 expression started after FGF4 but continued to increase after FGF4 started
decreasing. Direct comparison of FGF and TBX expression also showed strong positive correlation between TBXT
and both FGF4 and FGF17 but opposing relations with TBX6 (Supp. Fig. 5c). Together this suggested to us that
FGF4 and FGF17 may have overlapping functions, but that initial TBXT expression may depend more on FGF4
while FGF17 could play a later role in maintaining or expanding the PS-like population or facilitating subsequent
mesoderm differentiation.
Pseudotime may not accurately predict true dynamics. Therefore, we verified expression at different two
different times using RNA FISH (fluorescent in situ hybridization). We found that FGF4 expression was similar at
32h and 40h (Fig.5c-e, Supp. Fig. 5d). This is consistent with its decrease in pseudotime as cell differentiate to
nascent mesoderm, since in real time differentiation is asynchronous and new PS-like cells form while older ones
differentiate to nascent mesoderm, which would yield a flatter temporal profile. In contrast, FGF17 strongly
increased from 32h to 40h, again consistent with the pseudotime analysis (Fig.5f-h, Supp. Fig. 5e). As expected,
the spatial profiles of both FGF4 and FGF17 resembled TBXT, although the peaks were displaced slightly inward
(Fig.5d,g). Scatterplots further showed that expression of both FGF4 and FGF17 was restricted to TBXT positive
cells with high pERK levels (Fig. 5e,h). Conversely, however, high pERK was not restricted to FGF expressing cells
since the pERK ring extended significantly further inward than FGF expression. One possible explanation for this
is inward diffusion of FGF ligands.
To functionally test the requirement for these FGFs we decreased their expression. We created a cell line
stably expressing short hairpin RNA to knock down FGF4 which led to a decrease in TBXT and pERK (Fig. 5i). We
also decreased expression by transfection with small interfering RNAs (siRNA) with a similar outcome for FGF4
(Supp. Fig. 5f). Finally, in the absence of a homozygous knockout, we characterized a heterozygous knockout for
FGF17, which also led to decrease in TBXT (Fig. 5j, Supp. Fig. 5g-h, Supp. Fig. 6). The strength of this effect varied
from experiment to experiment, partially related to cell density variation (Supp. Fig. 5hj). We further interrogated
endoderm and primordial germ cell markers since both cell types require ERK signaling to differentiate and
transiently express TBXT (Fig. 5k, Supp. Fig. 5ij). Consistent with our previous work, we observed very little
expression of FOXA2 at 42h. However, SOX17 positive cells both positive for TFAP2C (primordial germ cell-like
cells) and negative for TFAP2C (prospective endoderm) were reduced. Combined, these data support partially
redundant functions for FGF4 and FGF17 in PS-like differentiation with FGF17 more important continued
differentiation to mesoderm and endoderm.

393
394 **Discussion**
395
We have shown that in a stem cell model for early human gastrulation, FGF-dependent ERK signaling is directly
required for differentiation to PS-like cells and their derivatives, and that loss of differentiation upon FGF inhibition
can be rescued by directly activating ERK. We also showed that FGF receptors are polarized basolaterally,
preventing the colony center from responding to exogenous FGF in the medium. Furthermore, FGF receptor
activation is increased in the PS-like ring, matching the pattern of active ERK despite overall receptor expression
being lower in the PS-like ring than in the pluripotent center. This suggests elevated ERK activity is due increased
binding of FGFs to the receptors in this region, possibly due to increased FGF concentration. We found that both
FGF4 and FGF17 have strongly increased expression in the PS-like cells, and decreased expression of either leads
to a reduction in PS-like cells or their derivatives, suggesting they might be responsible for increased ERK signaling
in the PS-like region.
Many questions remain. Although we showed that FGF gradients are likely the dominant cause for the
spatial pattern of ERK signaling, our scRNA-seq data suggest other levels of regulation may contribute, which
remains to be explored further. Furthermore, although we showed FGF/ERK signaling is directly required in PS-like

differentiation, it may also act indirectly by affecting the expression of Wnt and Nodal. We will explore this and the
more general question of how FGF integrates with the hierarchy of BMP, Wnt, and Nodal in a separate manuscript.
Related is the indirect effect FGF may have on patterning by its control of cell growth and thereby cell density. Cell
density affects the response to exogenous BMP and downstream expression of Wnt, Nodal, and various
inhibitors[4,47,48]. It likely has other effects as well. However, the rescue of TBXT expression but not growth by ERK
activation when FGF receptors are blocked suggests the FGF-dependent cell growth is not required to generate a
primitive streak-like ring.
Another important question is to what extent different FGFs act redundantly or perform different functions.
Our data suggest a picture where different FGFs are required consecutively, starting with exogenous FGF2, then
endogenous FGF4, possibly preceded by FGF8, to support initial differentiation of PS-like cells, and finally FGF17
to maintain and expand the PS-like ring and support mesoderm and endoderm differentiation. Whether these FGFs
could substitute for each other if they were expressed at similar times and levels is unclear. It also remains to be
determined whether the nearly constant high expression of FGF2 we observed is required for patterning. The fact
that PS-like differentiation was largely rescued by uniformly activating ERK to a supraphysiological level suggests
that the role for the different FGFs is simply to sufficiently activate ERK signaling. However, future work will have to
investigate the dox-SOS^{cat} rescue phenotype in much greater detail to see how differentiation and other cell
behaviors such as migration are affected.
It is interesting to ask if there is a simple explanation for the seemingly large differences in FGF expression
between primates and other species. FGF8 is the only FGF shown to be required for mouse gastrulation and is
upstream of FGF4 in both chick and mouse, but it is barely expressed in human and monkey. If FGFs are
interchangeable in simply activating ERK for primitive streak induction, variation in FGF expression is not surprising.
However, FGF4 and FGF8 are the main FGFs in mouse, chick and frog gastrulation[49–51], suggesting a high degree
of conservation. It is possible FGF8 has simply been substituted by FGF17, which is in the same subfamily[9], but
this does not explain why the role of FGF8 in gastrulation otherwise appears strongly conserved, or why FGF8 is
upstream of FGF4 in the mouse while FGF4 is expressed before FGF17 in 2D gastruloids. Similarly, it is possible
that FGF2 in primates substitutes for FGF5 in the mouse as the FGF that is highly expressed in the epiblast.
Although these are not in the same subfamily and generally have different receptor affinities, they both have highest
affinity for FGFR1[9]. That leaves FGF4 as the only FGF that may be conserved across vertebrate gastrulation, and
it is tempting to speculate on why. In Xenopus, FGF4 and TBXT (Xbra) function in a positive feedback loop. The
strongly correlated initial transcriptional dynamics of TBXT and FGF4 in 2D gastruloids are consistent with a similar
feedback loop. However, recent work suggests TBXT is not required for FGF4 or FGF17 expression[52]. One
possibility is that FGF expression is controlled by a different primitive streak transcription factor such as
EOMES/TBR2. Future work will have to investigate this further.
A final related question is the origin of the discrepancies we and others[36] found between the primate (i.e.,
non-mouse) samples in Fig. 2 and Supp. Fig. 2: FGF3 appear only significantly expressed in the human embryo,
but not in the monkey embryo and human stem cell model. This could also reflect genetic variation: if FGFs function
redundantly, their relative expression levels could be relatively unconstrained between individuals. On the other
hand, in the nascent mesoderm, FGF2 is only high in human gastruloids. This may reflect developmental stage: the
in vivo data is developmentally more advanced and nascent mesoderm may lose FGF2 expression later. It is also
possible the maintenance conditions of hPSCs lead to in vitro artefacts in FGF expression.
In conclusion, the complexity of FGF signaling, with its many ligands, modulators, and functions, has
prevented a clear understanding of its role in mammalian gastrulation in general and in human gastrulation
specifically. Here we showed that 2D human gastruloids provide a powerful approach to make headway and
although many open questions remain, we have taken significant steps towards this goal.
453
454

Exogenous FGF2 is only required at the start of differentiation

FGF4 and FGF17 expression correlate with TBXT

Reduction in FGF4 or FGF17 leads to reduction in PS-like cells and derivatives

Figure 5. a) Pattern at 42h with exogenous FGF2 in the culture medium for different amounts of time. **b)** Pseudotime analysis of FGFs and TBXT, TBX6 expression. **c,f)** FGF4 (c), and FGF17 (f) FISH co-stained for TBXT and pERK at 32h and 40h after BMP4 treatment. **d,g)** Radial intensity profiles for conditions in (c,f) respectively. **e,h)** Scatterplots of FGF4 (e), and FGF17 (h) mRNA versus pERK colored for TBXT. **i)** FGF4 FISH co-stained for TBXT and pERK in ctrl (scramble) shRNA versus FGF4 shRNA cells and corresponding radial intensity profiles.**j)** pERK and TBXT expression in WT versus FGF17+/- cells. **k)** SOX17 and TFAP2C expression in WT versus FGF17+/- cells.

Acknowledgements

We thank Lila Solnica-Krezel, Blerta Stringa, Aryeh Warmflash, and Ben Allen for discussions. We thank Jared Toettcher for the dox-SOScat plasmid. Library prep and next-generation sequencing was carried out in the Advanced Genomics Core at the University of Michigan. This work was supported by the National Institute of General Medical Sciences (NIGMS R35GM138346), and the Branco Weiss Fellowship – Society in Science. KJ was partially supported by the Michigan Pioneer Postdoctoral Fellowship and the NIH F32 Ruth L. Kirschstein Postdoctoral National Research Service Award (5F32HD108980-02).

Data availability

Raw single cell RNA sequencing data generated in this study was deposited in GEO (GSE271604), analyzed data will be added upon publication. Raw image data are available upon request.

Code availability

All code for data analysis and model simulations is available on https://github.com/idse/FGF

Competing Interests Statement

The authors declare no competing interests.

Methods

Cell lines
The cell lines used were the embryonic stem cell line ESI017 (XX), and the induced pluripotent stem cell line PGP1 (XY). The pluripotency of these cells was confirmed by immunostaining of pluripotency markers OCT3/4, SOX2, NANOG. All cells were routinely tested for mycoplasma contamination, and negative results were recorded. ESI017 cells stably expressing doxycycline inducible SOS^{cat} were made using a published Piggybac plasmid gifted by Jared Toettcher[37].

Cell culture and differentiation
Human pluripotent stem cells were cultured in the commercially defined pluripotency maintaining media mTeSR1 (STEMCELL Technologies #85850) on Cultrex (R&D Systems)-coated tissue culture plates. For FGF2 starvation we used Essential 6 Medium (Thermofisher, A1516401), or mTeSR1 without bFGF, TGF-beta, LiCl, GABA and pipecolic acid (STEMCELL Technologies #05896). For routine cell maintenance, cell passaging was performed every 3 days. For whole-colony passaging, L7[53] was used, while single-cell passaging was performed with Accutase or TrypLE (Gibco). Single-cell suspensions were used to seed for all the differentiation experiments to control initial cell number.

To directly differentiate to primitive streak-like cells, cells were resuspended in a single-cell suspension and seeded in cultrex coated wells in mTeSR with ROCK inhibitor (RI) Y-27632 (MeChemExpress, cat# HY-10583). Cells were then maintained in mTeSR with RI for 24h before adding CHIR (3-5uM) and Activin (100ng/ml) treatment for 24-30h. Experiments were performed in 18-well Ibidi slides (cat# 81818).

To differentiate cells for 2D gastruloids by micropatterning, we followed the protocol in[41]. Briefly, cells were resuspended in a single-cell suspension and seeded in laminin-coated micropatterned wells in mTeSR with ROCK inhibitor (RI) (MeChemExpress, cat# HY-10583). Micropatterned wells were washed with PBS-/- 30 minutes after the initial seeding to wash off the non-attached cells. Cells were then maintained for 24 hours in mTeSR before adding the BMP4 treatment by a full medium change. For time series micropatterned differentiation, full media changes with designated treatment(s) were performed every 24 hours. Experiments were performed in micropatterned 18-well Ibidi slides (cat# 81818) prepared as previously described[54]. Cell signaling reagents and doses used are listed in Supplementary Table 1.

Immunofluorescence staining
Antibodies can be found in Supplementary Tables 2 and 3. For stains including pERK, we used methanol fixation, for all other stains we used PFA fixation. PFA fixation: Samples from the 18-well Ibidi slides were rinsed with PBS, fixed for 20 min in 4% paraformaldehyde, rinsed twice with PBS, and blocked for 30 min at room temperature with 3% donkey serum and 0.1% Triton X-100 in 1× PBS. After blocking, cells were incubated with primary antibodies

at 4°C overnight, followed by three washes in PBST (PBS with 0.1% Tween 20). They were then incubated with
secondary antibodies and DAPI for 30 min at room temperature and washed twice in PBST at room temperature.
Methanol Fixation: Samples from the 18-well Ibidi slides were rinsed with PBS, fixed for 20 mins in cold methanol,
then rinsed with PBS for 5 min two times, after which 10% phosphate buffered formalin was added for 20 min at
room temperature. Cells were then rinsed with TBS twice followed by 5 mins of incubation in cold methanol and
rehydrate in TBS for 20 mins. Blocking was done for 30 mins at room temperature with 5% donkey serum, 1% BSA,
0.2% Triton-X-100 in 1x TBS after which staining proceeded the same way as with PFA fixation.
527
Microscopy
Fixed sample imaging was performed with an Andor Dragonfly/Leica DMI8 spinning disk confocal microscope with
a 40x, 1.1NA water and 20x 0.8NA air objectives using Andor Fusion software version 2.3.0.31 and a
Nikon/Yokogawa spinning dish confocal microscope with a 40x silicon oil objective using NIS Elements AR software
version 5.41.02. Live-cell imaging was performed with an Andor Dragonfly/Leica DMI8 spinning dish confocal
microscope under the controlled temperature (37°C), CO_2 concentration (5%), and humidity (>60%).
534
General image analysis
We segmented nuclei in individual z-slices based on nuclear fluorescence stained with DAPI in fixed cells using a
pipeline we previously described, which integrates two machine learning approaches: Ilastik pixel classification[55]
and Cellpose[56] (v1). Fluorescence intensities were then calculated per nucleus as mean intensities in the 3D
nuclear again with our established custom image-processing pipeline. To calculate the radial fluorescence
distributions in micropatterned colonies, colonies were subdivided into radial bins with equal numbers of cells, mean
or median and variance were then calculated within these bins. To calculate positive fractions, markers were first
thresholded based on their intensity distribution, after which the fraction of cells positive for each marker was
calculated within different bins. Standard deviations were then calculated over multiple replicate colonies. For
visualization purposes only, background subtraction using grayscale opening on a 100 micron scale and 5 pixel
wide median filter was applied slice by slice to z-stacks of micropatterned colonies before making maximal intensity
projections.
547
(Single-cell) RNA sequencing and analysis
Cells were collected using accutase and resuspended in ice-cold PBS. Single-cell RNA-sequencing was performed
by the University of Michigan Advanced Genomics Core. Cells were barcoded using the 10X Genomics Chromium
system (part numbers 1000268, 1000120, 1000215). For quality control, cDNA was quantified by Qubit High
Sensitivity DNA assay and Agilent TapeStation. Sequencing was performed on the Illumina NovaSeq 6000 with
NovaSeq S4 flowcell and Control Software version 1.7.0. Reads were aligned using cellranger-4.0.0 with the
GRCh38 reference. Further processing was mostly done in Python, primarily using the Scanpy[57] and SCVI[58]
packages. We integrated the two scRNA-seq datasets for 42h 2D human gastruloids using SCVI based on the top
2000 highly variable genes, which were selected using the default function in Scanpy using the seurat_v3 flavor.
Cell cycle genes were regressed out during integration. Dimensional reduction for visualization was performed using
PHATE[39] on the SCVI latent space. Gene expression was smoothened with MAGIC[59] for visualization on PHATE
plots in Fig. 3c, Supp. Fig.3b. To annotate different cell types, we used MAGIC to impute that data and get smooth
distributions of gene expression. We then thresholded marker genes to assign cell fate (Table 4), which produced
better results than the more common unsupervised clustering using Leiden, as well as more consistent results
between different datasets.
For pseudotime analysis (Fig. 5), we applied the trajectory inference tool Slingshot[60] in R to compute the
pseudotime trajectory in the SCVI latent representation. Slingshot determines the membership of a cell to a lineage
at each branching point by its projection distance to the fitted principal curve of that lineage. We projected the
mesoderm developmental trajectory onto the PHATE map for visualization. We further used TradeSeq[61], which fits
a generalized additive model (GAM), to infer the pseudotime gene expression trends along the mesoderm lineage
trajectory from Slingshot. For visualizing the correlation between the expression of FGF ligands and marker genes
(Supp. Fig. 5), we used our SCVI model to denoise the expression and mutual nearest neighbor to further harmonize
the difference between samples.
To quantify isoforms from RNA sequencing salmon v1.10.0 was used[62]. First, gencode.v45.transcripts.fa
was downloaded from gencodegenes.org. The reference was customized by removing lines for FGFR1 and
replacing them with the transcript sequence for FGFRIIIa, FGFRIIIb, FGFRIIIc (sequences included as
supplementary file). Three samples from GEO dataset GSM2257301 were downloaded through SRA
(SRX1725562 (Pluripotent), SRX1725577 (APS), SRX1725643 (MPS))[38]. Adapters and the first 10 bases were
trimmed from the FASTQ files using trim_galore. Finally, the Salmon "quant" command with the --validateMappings
option was used to quantify transcripts from trimmed FASTQ files. For single cell datasets,

https://timoast.github.io/sinto/ was used to subset the 10x aligned file by cluster. Reads for clusters of interest were
converted back to raw FASTQ files and run through the same alignment-based mode of salmon but as single end
reads.
581
Quantitative real-time PCR
583 RNA were extracted using RNAqueous™-Micro Total RNA Isolation Kit (CAT# AM1931, Thermo Fisher Scientific)
584 and then cDNA synthesis was prepared from it using SuperScript™ VILO™ cDNA Synthesis Kit (CAT# 11754250,
585 Thermo Fisher Scientific) following the manufacturer's protocol with adjustment for optimization. Measurements
586 were performed with SYBR green and the primers in the Table 5. GAPDH was used for normalization in all
587 experiments.
588
Fluorescent dextran assays
Fluorescent dextran essays were performed by adding 10 ug/ml fluorescein isothiocyanate-dextran (10KD, Sigma
Aldrich) to the media. To calculate fluorescence intensity as a function of distance from the colony edge, Otsu
thresholding followed by several morphological operations was used to create a mask for the colony in each z-slice,
thereby excluding bright dextran fluorescence outside the colony. Background subtraction on each z-slice of the
image was performed to further remove background signal from dextran outside the colony while maintaining signal
in the intercellular space. The radial intensity profile was then calculated from the maximal intensity projection of
the masked, background subtracted image.
597
Scratch assays
Human embryonic or induced pluripotent stem cells were differentiated on micropattern wells for 41h, then each
micropattern colony was manually scratched with a serological needle. After 60mins, they were fixed with PFA or
methanol followed by immunostaining with appropriate antibodies. To calculate the intensity profile as a function of
distance from a scratch, we created a mask for the scratch based on the difference between an overall mask for
the colony and the convex hull, calculated the distance transform of the scratch mask, then binned pixels by distance
from the scratch and calculated their mean ERK intensity.
605
Transmembrane assays
607 Human embryonic or induced pluripotent stem cells were seeded into Corning 6.5 mm transwell with 0.4 µm pore
608 polyester membrane insert (cat# 3470) with E6 supplemented with TGF1b, FGF2, and Y-27632 for 16h, then media
609 was changed to E6 with TGF1b. After 24h, the media was changed to E6 supplemented with FGF2 from the top or
610 the bottom side of the transwell. 30 mins later, cells were fixed followed by immunofluorescence staining. The
611 transwell cartoon in Fig 4 was generated with Biorender.
612
Fluorescence In Situ Hybridization
614 The fluorescence in situ hybridization protocol (FISH) was performed according to the manufacturer's instructions
615 (ACDbio, RNAscope multiplex fluorescent manual). A list of probes and reagents can be found in Supplementary
616 Table 6.
617
Gene targeting by shRNA
619 We used the construct hFGF4[shRNA#1] from VectorBuilder in a PiggyBac Transposon Vectors with puromycin
620 resistance which was stably integrated into ESI17 hESCs. Transfections were performed using Lipofectamine™
621 Stem Transfection Reagent (CAT# STEM00001, Thermo Fisher Scientific), following the manufacturer's protocol.
622
Gene targeting by siRNA
624 We purchased predesigned siRNA from Millipore Sigma; NM_002007 (FGF4), NM_008004 (FGF17). The siRNA
625 transfections were performed using Lipofectamine™ RNAiMAX Transfection Reagent (CAT# 13778075, Thermo
626 Fisher Scientific), following the manufacturer's protocol with adjustment for optimization.
627
Statistics and reproducibility
629 All experiments were performed twice or more. All the attempts in replicating experiments yielded consistent results.
630 Unless specifically noted, all the quantifications were performed with N=4 colonies within the same experimental
631 condition.
632
Ethics statement

634 We comply with all the ethical regulations in our research. We were granted approval by the Human Pluripotent
635 Stem Cell Research Oversight (HPSCRO) Committee at the University of Michigan to work with human embryonic
636 stem cells and human induced pluripotent stem cells.
637
638 **Supplementary Table 1: Cell signaling reagents.**

Reagent	Nickname	Vendor, cat #	Dose	Function
rhBMP4	BMP	R&D Systems, 314BP/CF	50ng/ml	Activate BMP signaling
LDN193189	BMPRi	MedChemExpress, HY-12071	250 nM	Inhibit BMP receptors
IWP2	WNTSeci	ApexBio, A3512-5	5 µM	Block Wnt secretion
rActivinA	Activin	R&D Systems, 338AC050	100ng/ml	Activate Activin/Nodal pathway
PD0325901	MEKi	ESIBIO, ST10009	5uM	Inhibit MEK function
PD173074	FGFRi	Medchemexpress,HY10321	1uM	Inhibit FGF receptors
SB 431542	SB	Fisher, 50-101-2514	10uM	Inhibit Nodal receptors
CHIR 99021	CHIR	Tocris, 4423	See figures	Canonical Wnt agonist

639
640 **Supplementary Table 2: Primary antibodies used for immunofluorescence.**

Antigen	Species	Dilution	Cat #	Vendor
ISL1	Mouse	1:200	39.4D5	DSHB
SOX2	Rabbit	1:200	3579S	Cell Signaling Tech
TBXT (BRA)	Goat	1:300	AF2085	R&D Systems
EOMES	Mouse	1:200	MAB6166	R&D Systems
MIXL1	Rabbit	1:150	A17223	ABclonal
pERK	Rabbit	1:200	9101S	Cell signaling Tech
pFGFR1	Rabbit	1:100	52928	Cell signaling Tech
FGFR1	Rabbit	1:100	9740T	Cell signaling Tech

641
642 **Supplementary Table 3: Secondary antibodies.**

Antigen	Species	Dilution	Cat #	Vendor
Alexa Fluor 488 anti-mouse	Donkey IgG	1:500	A21202	ThermoFisher Scientific
Alexa Fluor 555 anti-rabbit	Donkey IgG	1:500	A31572	ThermoFisher Scientific

Alexa Fluor 647 anti-goat	Donkey IgG	1:500	A21447	ThermoFisher Scientific
Alexa Fluor 647 anti-rabbit	Donkey IgG	1:500	A31573	ThermoFisher Scientific
Alexa Fluor 555 anti-goat	Donkey IgG	1:500	A21432	ThermoFisher Scientific

Supplementary Table 4: Marker genes for cell fate annotation

Cell type	Markers
Pluripotent	SOX2+POU5F1+TBXT-MIXL1-SOX17-
Amnion-like cell (AM-LC)	ISL1+POU5F1+TBXT-MIXL1-SOX17-
Primitive streak-like cell (PS-LC)	TBXT+MIXL1+POU5F1+TBX6-SOX17-FOXA2-
Nascent mesoderm (NasMe)	TBXT+MIXL1+TBX6+SOX17-
Endoderm	SOX17+EOMES+TFAP2C-
Ectoderm	SOX2+POU5F1-TBXT-MIXL1-SOX17-
Primordial germ cell-like cells (PGC-LC)	SOX17+TFAP2C+

Supplementary Table 5: qPCR primers

Gene	Forward	Reverse
FGFR1	AATGAGTACGGCAGCATCAAC	ACCTCGATGTGCTTTAGCCAC
FGFR1 IIIb	GCCCAGACAACCTGCCTTAT (Binds to FGFR1-IIIa)	CACACATACTCCCCGCTCTG (Binds to FGFR1-IIIb)
FGFR1 IIIc	GCCCAGACAACCTGCCTTAT (Binds to FGFR1-IIIa)	AGAGTGATGGGATCCGA (Binds to FGFR1-IIIc)
GAPDH	ACAACTTTGGTATCGTGGAAGG	GCCATCACGCCACAGTTTC

Supplementary Table 6: Probes for Fluorescent In Situ Hybridization

Probe	TSA dilution	Cat #	Vendor
Hs-FGF4	1:2000	RNAScope #512291	ACDBio
Hs-FGF17-C1	1:2000	RNAScope #1148351-C1	ACDBio

References

1.   Arnold, S. J. & Robertson, E. J. Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. *Nat. Rev. Mol. Cell Biol.* **10**, 91–103 (2009).
2.   Martyn, I., Kanno, T. Y., Ruzo, A., Siggia, E. D. & Brivanlou, A. H. Self-organization of a human organizer by combined Wnt and Nodal signalling. *Nature* (2018) doi:10.1038/s41586-018-0150-y.
3.   Chhabra, S., Liu, L., Goh, R., Kong, X. & Warmflash, A. Dissecting the dynamics of signaling events in the BMP, WNT, and NODAL cascade during self-organized fate patterning in human gastruloids. *PLoS Biol.* **17**, e3000498 (2019).

4.   Etoc, F. *et al.* A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid Self-
Organization. *Dev. Cell* **39**, 302–315 (2016).
5.   Zhang, Z., Zwick, S., Loew, E., Grimley, J. S. & Ramanathan, S. Embryo geometry drives formation of
robust signaling gradients through receptor localization. (2018) doi:10.1101/491290.
6.   Heemskerk, I. *et al.* Rapid changes in morphogen concentration control self-organized patterning in
human embryonic stem cells. *Elife* **8**, (2019).
7.   Massey, J. *et al.* WNT ligands stimulate transient signaling in human pluripotent cells and synergize with
TGF-β ligands to stimulate sustained signaling during differentiation. *bioRxiv* 406306 (2018) doi:10.1101/406306.
8.   Yang, R. *et al.* Amnion signals are essential for mesoderm formation in primates. *Nat Commun* **12**, 5126
(2021).
9.   Ornitz, D. M. & Itoh, N. The Fibroblast Growth Factor signaling pathway. *Wiley Interdiscip Rev Dev Biol* **4**,
215–266 (2015).
10.   LaBonne, C. & Whitman, M. Mesoderm induction by activin requires FGF-mediated intracellular signals.
Development **120**, 463–472 (1994).
11.   Krens, S. F. G., Corredor-Adámez, M., He, S., Snaar-Jagalska, B. E. & Spaink, H. P. ERK1 and ERK2
MAPK are key regulators of distinct gene sets in zebrafish embryogenesis. *BMC Genomics* **9**, 196 (2008).
12.   Yao, Y. *et al.* Extracellular signal-regulated kinase 2 is necessary for mesoderm differentiation. *Proc Natl*
Acad Sci U S A **100**, 12759–12764 (2003).
13.   Mitrani, E., Gruenbaum, Y., Shohat, H. & Ziv, T. Fibroblast growth factor during mesoderm induction in
the early chick embryo. *Development* **109**, 387–393 (1990).
14.   Corson, L. B., Yamanaka, Y., Lai, K.-M. V. & Rossant, J. Spatial and temporal patterns of ERK signaling
during mouse embryogenesis. *Development* **130**, 4527–4537 (2003).
15.   Morgani, S. M. & Hadjantonakis, A.-K. Signaling regulation during gastrulation: Insights from mouse
embryos and *in vitro* systems. in *Current Topics in Developmental Biology* (eds. Small, S. & Briscoe, J.) vol. 137
391–431 (Academic Press, 2020).
16.   De Simone, A. *et al.* Control of osteoblast regeneration by a train of Erk activity waves. *Nature* **590**, 129–
133 (2021).
17.   Marshall, C. J. Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular
signal-regulated kinase activation. *Cell* **80**, 179–185 (1995).
18.   Ender, P. *et al.* Spatiotemporal control of ERK pulse frequency coordinates fate decisions during
mammary acinar morphogenesis. *Dev Cell* **57**, 2153-2167.e6 (2022).
19.   Aoki, K. *et al.* Propagating Wave of ERK Activation Orients Collective Cell Migration. *Dev Cell* **43**, 305-
317.e5 (2017).
20.   Ciruna, B. & Rossant, J. FGF Signaling Regulates Mesoderm Cell Fate Specification and Morphogenetic
Movement at the Primitive Streak. *Dev. Cell* **1**, 37–49 (2001).
21.   Sun, X., Meyers, E. N., Lewandoski, M. & Martin, G. R. Targeted disruption of Fgf8 causes failure of cell
migration in the gastrulating mouse embryo. *Genes Dev.* **13**, 1834–1846 (1999).
22.   Fletcher, R. B. & Harland, R. M. The role of FGF signaling in the establishment and maintenance of
mesodermal gene expression in Xenopus. *Dev Dyn* **237**, 1243–1254 (2008).
23.   Yang, X., Dormann, D., Münsterberg, A. E. & Weijer, C. J. Cell Movement Patterns during Gastrulation in
the Chick Are Controlled by Positive and Negative Chemotaxis Mediated by FGF4 and FGF8. *Developmental Cell*
3, 425–437 (2002).
24.   Fürthauer, M., Reifers, F., Brand, M., Thisse, B. & Thisse, C. sprouty4 acts in vivo as a feedback-induced
antagonist of FGF signaling in zebrafish. *Development* **128**, 2175–2186 (2001).
25.   Tyser, R. C. V. *et al.* Single-cell transcriptomic characterization of a gastrulating human embryo. *Nature*
600, 285–289 (2021).
26.   Zhai, J. *et al.* Primate gastrulation and early organogenesis at single-cell resolution. *Nature* **612**, 732–738
(2022).
27.   Deng, C. X. *et al.* Murine FGFR-1 is required for early postimplantation growth and axial organization.
Genes Dev **8**, 3045–3057 (1994).
28.   Yamaguchi, T. P., Harpal, K., Henkemeyer, M. & Rossant, J. fgfr-1 is required for embryonic growth and
mesodermal patterning during mouse gastrulation. *Genes Dev* **8**, 3032–3044 (1994).
29.   Ciruna, B. G., Schwartz, L., Harpal, K., Yamaguchi, T. P. & Rossant, J. Chimeric analysis of fibroblast
growth factor receptor-1 (Fgfr1) function: a role for FGFR1 in morphogenetic movement through the primitive
streak. *Development* **124**, 2829–2841 (1997).
30.   García-García, M. J. & Anderson, K. V. Essential role of glycosaminoglycans in Fgf signaling during
mouse gastrulation. *Cell* **114**, 727–737 (2003).
31.   Oki, S., Kitajima, K. & Meno, C. Dissecting the role of Fgf signaling during gastrulation and left-right axis

formation in mouse embryos using chemical inhibitors. *Dev Dyn* **239**, 1768–1778 (2010).
32. Feldman, B., Poueymirou, W., Papaioannou, V. E., DeChiara, T. M. & Goldfarb, M. Requirement of FGF-4
for postimplantation mouse development. *Science* **267**, 246–249 (1995).
33. Warmflash, A., Sorre, B., Etoc, F., Siggia, E. D. & Brivanlou, A. H. A method to recapitulate early
embryonic spatial patterning in human embryonic stem cells. *Nat. Methods* **11**, 847–854 (2014).
34. Jo, K. *et al.* Efficient differentiation of human primordial germ cells through geometric control reveals a key
role for Nodal signaling. *Elife* **11**, e72811 (2022).
35. Minn, K. T. *et al.* High-resolution transcriptional and morphogenetic profiling of cells from micropatterned
human ESC gastruloid cultures. *Elife* **9**, (2020).
36. Minn, K. T., Dietmann, S., Waye, S. E., Morris, S. A. & Solnica-Krezel, L. Gene expression dynamics
underlying cell fate emergence in 2D micropatterned human embryonic stem cell gastruloids. *Stem Cell Reports*
16, 1210–1227 (2021).
37. Underhill, E. J. & Toettcher, J. E. Control of gastruloid patterning and morphogenesis by the Erk and Akt
signaling pathways. *Development* **150**, dev201663 (2023).
38. Loh, K. M. *et al.* Mapping the Pairwise Choices Leading from Pluripotency to Human Bone, Heart, and
Other Mesoderm Cell Types. *Cell* **166**, 451–467 (2016).
39. Moon, K. R. *et al.* Visualizing structure and transitions in high-dimensional biological data. *Nat.*
Biotechnol. **37**, 1482–1492 (2019).
40. Neben, C. L., Lo, M., Jura, N. & Klein, O. D. Feedback regulation of RTK signaling in development. *Dev*
Biol **447**, 71–89 (2019).
41. Chen, B. *et al.* Extended culture of 2D gastruloids to model human mesoderm development. *bioRxiv*
2024.03.21.585753 (2024) doi:10.1101/2024.03.21.585753.
42. Khoa, L. T. P. *et al.* Visualization of the Epiblast and Visceral Endodermal Cells Using Fgf5-P2A-Venus
BAC Transgenic Mice and Epiblast Stem Cells. *PLOS ONE* **11**, e0159246 (2016).
43. Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early organogenesis. *Nature*
566, 490–495 (2019).
44. Morgani, S. M. *et al.* A Sprouty4 reporter to monitor FGF/ERK signaling activity in ESCs and mice.
Developmental Biology **441**, 104–126 (2018).
45. Yu, S. R. *et al.* Fgf8 morphogen gradient forms by a source-sink mechanism with freely diffusing
molecules. *Nature* **461**, 533–536 (2009).
46. Benington, L., Rajan, G., Locher, C. & Lim, L. Y. Fibroblast Growth Factor 2-A Review of Stabilisation
Approaches for Clinical Applications. *Pharmaceutics* **12**, 508 (2020).
47. Teague, S. *et al.* Time-integrated BMP signaling determines fate in a stem cell model for early human
development. *Nat Commun* **15**, 1471 (2024).
48. Ortiz-Salazar, M. A., Camacho-Aguilar, E. & Warmflash, A. Endogenous Nodal switches Wnt
interpretation from posteriorization to germ layer differentiation in geometrically constrained human pluripotent
cells. *bioRxiv* 2024.03.13.584912 (2024) doi:10.1101/2024.03.13.584912.
49. Isaacs, H. V., Pownall, M. E. & Slack, J. M. eFGF is expressed in the dorsal midline of Xenopus laevis. *Int*
J Dev Biol **39**, 575–579 (1995).
50. Christen, B. & Slack, J. M. FGF-8 is associated with anteroposterior patterning and limb regeneration in
Xenopus. *Dev Biol* **192**, 455–466 (1997).
51. Hardy, K. M., Yatskievych, T. A., Konieczka, J., Bobbs, A. S. & Antin, P. B. FGF signalling through
RAS/MAPK and PI3K pathways regulates cell movement and gene expression in the chicken primitive streak
without affecting E-cadherin expression. *BMC Dev. Biol.* **11**, 20–17 (2011).
52. Bulger, E. A., Muncie-Vasic, I., Libby, A. R. G., McDevitt, T. C. & Bruneau, B. G. TBXT dose sensitivity
and the decoupling of nascent mesoderm specification from EMT progression in 2D human gastruloids.
Development **151**, dev202516 (2024).
53. Nie, Y., Walsh, P., Clarke, D. L., Rowley, J. A. & Fellner, T. Scalable passaging of adherent human
pluripotent stem cells. *PLoS ONE* **9**, e88012 (2014).
54. Freeburne, E. *et al.* Spatial Single Cell Analysis of Proteins in 2D Human Gastruloids Using Iterative
Immunofluorescence. *Curr Protoc* **3**, e915 (2023).
55. Sommer, C., Straehle, C., Kothe, U. & Hamprecht, F. A. Ilastik: Interactive learning and segmentation
toolkit. in *2011 8th IEEE International Symposium on Biomedical Imaging (ISBI 2011)* 230–233 (IEEE, 2011).
doi:10.1109/ISBI.2011.5872394.
56. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular
segmentation. *Nat. Methods* **18**, 100–106 (2021).
57. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis.
Genome Biol **19**, 15 (2018).

58.   Gayoso, A. *et al.* A Python library for probabilistic analysis of single-cell omics data. *Nat Biotechnol* **40**,
163–166 (2022).
59.   van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell* **174**,
716-729.e27 (2018).
60.   Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC*
Genomics **19**, 477 (2018).
61.   Van den Berge, K. *et al.* Trajectory-based differential expression analysis for single-cell sequencing data.
Nat Commun **11**, 1201 (2020).
62.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware
quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).
63.   Eswarakumar, V. P., Lax, I. & Schlessinger, J. Cellular signaling by fibroblast growth factor receptors.
Cytokine Growth Factor Rev. **16**, 139–149 (2005).

Supplementary Figures

A ring of phosporylated ERK first appears around 24h and expands inward over time

All primitive streak markers are lost upon FGF/ERK inhibition

Rescue of FGFR inhibition by direct ERK activation with dox-SOScat

There is a *direct* requirement for FGF/ERK signaling in primitive streak-like differentiation

Supplementary Figure 1. a) DAPI images and ISL1 stains corresponding to Fig. 1a. **b)** Scatter plots of TBXT and ISL1 expression. **c)** Staining for three primitive streak markers with or with FGFRi or MEKi. **d)** DAPI stain corresponding to Fig. 1d. **e)** Radial intensity profile for TBXT corresponding to Fig.1d. **f)** Quantification of conditions in Fig.1g for five images each, N indicates total number of cells.

Supplementary Figure 2. a) PHATE projection of scRNA-seq data colored for sample. **b,c)** Expression of canonical FGF ligands across different cell types in 2D gastruloids (b) and human CS7 embryo (c). Expression across clusters is normalized by row, while total expression is shown in a separate column to the directly to left. The leftmost column indicates differential expression between PS-like and pluripotent cells. The color scale of (differential) expression is cut off for better contrast. **d)** Threshold-based annotation (left) for human, monkey, and mouse embryos used in interspecies comparison (Fig. 2f), compared to original annotation (right). **e)** Interspecies comparison as in Fig. 2f but for nascent mesoderm and PS-LC + nascent mesoderm. **f)** Bulk RNA-seq data by Loh et al[38] for expression of different FGFs in directed differentiation to mid primitive streak (MPS) or anterior primitive streak (APS) relative the human embryonic stem cells (hESCs). **g)** Expression of genes involved in FGF signaling.

The phosphorylated ERK pattern reflects receptor activation

FGFR1-IIIc is the dominant isoform across the colony

812
813
814 **Supplementary Figure 3. a)** DAPI image corresponding to Fig. 3a. **b-c)** pFGFR1 (b) or pERK (c) and TBXT stains with and
815 without continuous MEKi treatment. **d)** DAPI image corresponding to Fig. 3b. **e)** Design of qPCR primers to detect FGFR1
816 isoforms, diagram adapted from Eswarakumar et al[63]. **f)** Primer sequence and properties. **g)** Amplicon sizes match prediction.
817 Scale bars 50um.
818

Exogenous FGF2 is only required at the start of differentiation

FGF4 expression correlates most strongly with TBXT (and therefore pERK)

Reduction in FGF4 or FGF17 leads to reduction in PS-like cells and derivatives

Supplementary Figure 5. a) DAPI image corresponding to Fig. 5a. **b)** Pseudotime dynamics from Fig.5b with each gene normalized individually. **c)** Correlations between FGFs and TBXT, TBX6 colored for clusters, with cluster colors matching Fig. 2d. **d,e)** DAPI images and overlays including DAPI corresponding to Fig.5c,f respectively. **f)** Representative stainings and quantification for FGF4 siRNA versus controls. **g)** DAPI stain, individual channels and overlay with DAPI for images corresponding to Fig. 5k. **h)** Scatterplot of fraction of TBXT-positive cells in repeated experiments with FGF17+/- cells colored for genotype (left) or experiment (right). **i)** Individual channels including DAPI and FOXA2 for images corresponding to Fig.5l. **j)** Radial profiles of cells positive for different markers in FGF17+/- versus WT.

Reduction in FGF4 or FGF17 leads to reduction in PS-like cells and derivatives

828
829
830   **Supplementary Figure 6. a)** Genotyping for FGF17+/- cells.
831