1 **Strength of enemy release from parasitoids is context-dependent in the invasive African Fig**
2 **Fly, *Zaprionus indianus***
3

4 Camille R. Walsh-Antzak[1] and Priscilla A. Erickson[1*]
5
6   1. University of Richmond, Richmond, VA, 23173
7   * Corresponding author; perickso@richmond.edu

8
Abstract

Understanding the mechanisms underlying the success of biological invasions is essential to
employ effective prediction and management strategies. Escape from natural enemies in invaded
regions (enemy release hypothesis, ERH) and increased competitive ability are hallmarks of
invasive species; however, these two processes are rarely studied within the same context. Here,
we examined the effect of enemy release on the competition outcomes of a successful invasive
insect pest in North America, the African fig fly (*Zaprionus indianus*). Parasitoid wasps such as
Leptopilina heterotoma that parasitize drosophilid larvae may seek out established species with
known host suitability over a novel species, so we hypothesized *Z. indianus* may have low
susceptibility to parasitoids, giving them a competitive advantage over co-occurring
drosophilids. We tested this hypothesis by comparing the adult emergence rates from *Z. indianus*
larvae reared alone or in competition with *Drosophila hydei* or *D. simulans* larvae in the
presence and absence of parasitoid wasps under low and high larval densities. At low larval
densities, *Z. indianus* emerged at equal rates to *D. hydei* but outcompeted *D. simulans,* and these
outcomes were not affected by parasitoids. However, at high densities, the addition of parasitoids
shifted competition outcomes in favor of *Z. indianus*, suggesting enemy release provides a
competitive advantage under some circumstances. These results indicate that the strength of
enemy release in *Z. indianus* is widely dependent on contextual factors such as density and
competitor species. This study emphasizes how a community approach to testing the ERH is
vital as the overall interpretation of the presence and strength of enemy release differed between
intraspecific and interspecific experiments. Further investigation of how these results apply to
field environments could offer insight into how *Z. indianus* alters ecosystems and how
productive biological control may limit the spread of *Z. indianus*.

32
33
Short Abstract

Invasive species may succeed in new environments in part because they are less susceptible to
diseases and parasites that have co-evolved with local hosts, giving invaders a competitive
advantage. We tested this hypothesis by competing an invasive fruit fly against established
species in the presence of parasitoid wasps that lay their eggs in fruit fly larvae. We found that
the invasive species generally outcompeted other species in the presence of parasitoids, but the

extent of its advantage depended on the species it was competing against and the number of
larvae present.
42
Introduction
The spread of invasive species poses significant ecological, environmental, and economic
threats. Invasive species have largely contributed to biodiversity loss and the extinction of native
species (Bradshaw et al. 2016) and are destructive in novel environments, leading to agricultural
damage and economic costs (Diagne et al. 2021). Furthermore, invasive species raise human
health concerns as they can alter disease-vector dynamics and facilitate the spread of pathogens
(Roy et al. 2022). Despite the well-known risks of biological invasions, managing invasive
species remains a challenge. A comprehensive understanding of the mechanisms underlying
invasive species' success is vital for effective conservation, management, and mitigation
practices.
One of the most widely proposed hypotheses explaining the success of invasive species is
the enemy release hypothesis (Keane & Crawley 2002, Enders et al. 2018). This hypothesis
proposes that invasive species in novel environments are no longer impacted by their natural
enemies (e.g., predators, parasites, or pathogens from the invader's native range). As a result the
invader population grows as local species, but not the invader, are restrained by natural enemies.
Due to the loss of natural enemies, invasive species may also succeed in invaded communities by
relocating resources from defence mechanisms to growth and development (evolution of
increased competitive ability hypothesis, Blossey & Notzold 1995). Competition with native
species also plays a crucial role in determining the success or failure of invasive species (Clarke
& McGeoch 2023). The inherent competitive ability of invasive species is often superior to
native species, facilitating their dominance in invaded communities (Gioria & Osborne 2014,
Fridley & Sax 2014). Although components of enemy release and competition overlap, there are
few studies examining both as factors for invasion success. Studying these two processes within
the same context may provide a more cohesive understanding of the spread and success of
invasive species.
The African fig fly, *Zaprionus indianus*, is a member of the family Drosophilidae and is
native to tropical regions of central Africa. *Z. indianus* is a polyphagous consumer of decaying
fruit and has over 80 documented host species, though it can also be a pest on ripening figs
(Pfeiffer et al. 2019) and potentially raspberries (Stazzonelli et al. 2023). In the 1990s, *Z.*
indianus was first detected in South America in Brazil, where it quickly spread across the
continent and damaged fig crops (Vilela 1999). Following a northward invasion pattern, *Z.*
indianus was detected in Florida in 2005 (Van der Linde et al. 2006) and has since been detected
across eastern North America (Pfeiffer et al. 2019, Rakes et al. 2023). Sampling data from apple
and peach orchards reveal repeated and rapid population growth relative to other drosophilids,
suggesting *Z. indianus* outcompetes co-occurring drosophilid species (Rakes et al. 2023).
However, the factors that facilitate the success of *Z. indianus* remain unknown. Evaluating the
role of enemy release in competition outcomes could shed light on the mechanisms underlying

the repeated success of *Z. indianu*s in novel environments. *Zaprionus indianus* is an optimal
study system as its relatively recent arrival to North America and lack of permanent
establishment in temperate regions (Pfeiffer et al. 2019) suggests it has not co-evolved with
North American enemies, providing an opportunity to assess whether reduced enemy pressure in
a novel environment mediates their population success.
Parasitoid wasps are natural enemies of drosophilids and exert strong selective pressures
on fly populations, with up to 90% of drosophilids parasitized by parasitoids in certain
environments (Fleury et al. 2004). Additionally, host-parasitoid interactions exhibit
coevolutionary arms races, and each have highly specialized attack and defense mechanisms
(Godfray & Shimada 1999). Invasive species lack this coevolutionary relationship with
parasitoids present in the introduced range which raises questions about how they interact with
these novel parasitoids (Kraaijeveld et al. 1998). *Leptopilina heterotoma* is a generalist parasitoid
wasp found in North America that parasitizes larvae of numerous drosophilid species due to its
low selection threshold (Quicray et al. 2023, de Bruijn et al. 2022). The life cycle of *L.
heterotoma* begins with an adult parasitoid ovipositing into fly larvae. If the parasitization is
successful (e.g., the parasitoid overcomes the flies' immune defense), then the parasitoid
develops inside the host and slowly consumes it, ultimately resulting in host death during fly
pupation. An adult parasitoid typically emerges from the fly pupae ~21-23 days after oviposition
(Quicray et al. 2023). Although *L. heterotoma* is a generalist, not all hosts are suitable for
parasitoid development (Quicray et al. 2023). While *L. heterotoma* detects *Z. indianus* as
potential hosts (e.g., ovipositing in larvae), the majority of *Z. indianus* successfully emerge as
adults, suggesting that the immune system of *Z. indianus* is resistant to *L. heterotoma*, perhaps
due to the presence of giant hemocytes (Kacsoh et al. 2014). Additionally, no *L. heterotoma*
adults emerge from *Z. indianus*, indicating *Z. indianus* is an unsuitable host. Therefore, invasive
populations of *Z. indianus* might benefit relative to other drosophilids by escaping parasitism
from *L. heterotoma*.
Resource competition is another major driving force behind the success and spread of
invasive species as they often have greater capacities to exploit resources more efficiently than
local species (Mooney & Cleland 2001, Gioria & Osborne 2014). Ephemeral food substrates,
such as rotting fruits and their associated microbiota, result in both intra- and interspecific
competition for drosophilids (Rohlfs & Hoffmeister 2004). *Z. indianus* likely experiences high
rates of resource competition with other drosophilid species because it occupies similar niches
(e.g., opportunistic oviposition in rotting host fruits) (Bragard et al. 2022). Competition
outcomes in drosophilids are widely context-dependent and may fluctuate based on abiotic and
biotic factors such as air temperature or fruit substrate (Comeault & Matute 2021, de Paiva
Mendonça 2023). Host-parasitoid interactions can also influence competition outcomes. For
example, *D. melanogaster* outperforms its closest relative, *D. simulans,* in the absence of
parasitoids, but these competition outcomes are inverted when parasitoids are present
(Boulêtreau et al. 1991). However, it is unknown how parasitoids may alter competition
outcomes between recent invaders and local populations. If *L. heterotoma* preferentially

parasitizes established species in the presence of *Z. indianus*, then *Z. indianus* may capitalize on
reduced competition stressors.
In this study, we evaluated how parasitoids affect the competition outcomes of *Z.*
indianus under controlled laboratory conditions. As our main goal was to evaluate how enemy
release influences the dominance of *Z. indianus* in invaded environments, we used a community-
comparison approach as described by Brian & Catford (2023). We measured adult emergence
rates in the absence and presence of *L. heterotoma* to compare competition outcomes both
between and within species. We conducted assays at low density and high density treatments to
account for varying fly densities observed in the field across different locations and time periods,
which may provide insight about host-parasitoid interactions in spatially and temporally
heterogeneous environments (Rakes et al. 2023, Gleason et al. 2019). In the interspecific
competition assays, we offered parasitoids the opportunity to parasitize the larvae of *Z. indianus*
and another drosophilid species. We used *D. simulans* and *D. hydei* as co-occurring, established
species as they are both cosmopolitan in distribution (Werner et al. 2018). Although not native to
North America (Zhao & Begun 2017, Lachaise et al. 1988), *D. simulans* and *D. hydei* have both
been widespread in the eastern US since at least 1921 (Sturtevant 1921), allowing substantially
more time for coevolution with North American parasitoids relative to *Z. indianus*. Genomic data
suggest that despite a population bottleneck, introduced North American *Z. indianus* populations
retain substantial genetic variation relative to congeners (Comeault et al. 2020, 2021). Therefore,
the strength of enemy release may vary due to genetic variation. We compared the parasitized
emergence rates of a low-latitude population from Florida with a high-latitude population from
Connecticut, as flies from Florida and temperate regions are genetically differentiated (Erickson
et al. 2024). Lastly, we evaluated emergence rates with differing parasitoid exposure times to test
whether *L. heterotoma* exhibits host-switching (Murdoch 1969), in which a parasitoid initially
oviposits in a favorable host species, but then switches to a different host after the original host
becomes saturated.
Given the lack of coevolutionary history and previously described host incompatibility of
L. heterotoma to *Z. indianus*, we hypothesized that *Z. indianus* would be less susceptible to
parasitization than established drosophilids, leading to higher emergence rates for *Z. indianus*.
Given the rapid population growth of *Z. indianus* observed in orchards (Rakes et al. 2023), we
predicted that *Z. indianus* performs equal to or better than established species in the absence of
parasitoids, and this advantage is further intensified by the presence of *L. heterotoma*, alleviating
competition stressors for *Z. indianus*.

Materials & Methods

Insect rearing

Adult *Z. indianus* were obtained from isofemale lines derived from females collected in North
America in 2022 and 2023 (Rakes et al. 2023). *D. simulans* were collected from orchards in
Florida in 2022, and *D. hydei* were collected from orchards in Virginia in 2023. All flies were
maintained on a cornmeal-molasses artificial diet at room temperature. All parents of the

experimental offspring were raised in controlled lab conditions for at least two generations. *L.*
heterotoma (Lh14) stocks were obtained from Todd Schlenke at the University of Arizona. The
Lh14 strain was originally collected in 2002 from Winters, California (Kacsoh & Schlenke
2012). Parasitoids were maintained on laboratory strains of *D. melanogaster* (w1118 and
Canton-S). To elongate parasitoid lifespan, 500 μL of 50% honey-water solution was provided
on vial plugs.

Intraspecific competition assays

To test how parasitoid presence affects intraspecific competition outcomes of *Z. indianus,* we
prepared vials with only *Z. indianus* larvae in the absence and presence of parasitoid wasps. We
allowed adult flies to oviposit on 3% agar plates containing 10% grape juice concentrate for 24
hours. A paste of baker's yeast and water was added to each plate to encourage oviposition. We
collected two-day-old larvae from the plates and inoculated vials filled with 10 mL of cornmeal-
molasses substrate. We compared emergence rates at low densities (50 total larvae) and high
densities (150 total larvae). For parasitized treatments, we immediately added six adult *L.*
heterotoma (three female, three male) to the inoculated vials for 24 hours. Parasitoids were not
offered hosts prior to experiments. In the control treatments, we followed the same procedure but
did not add parasitoids. We counted emerging adult flies every 1-2 days until no further adults
emerged. Intraspecific assays had two replicates per treatment. We conducted two independent
intraspecific assays (one alongside each interspecific assay, see below) and combined the data
for a total of four vials per density and parasitoid treatment.

Interspecific competition assay

To test how parasitoid presence affects interspecific competition outcomes, we compared the
larva-to-adult survival rates of *Z. indianus* and co-occurring drosophilid species (*D. hydei or D.*
simulans) reared together in the absence and presence of parasitoid wasps. Low and high total
densities were the same as the intraspecific assays (50 and 150 larvae); however, interspecific
assays contained equal numbers of each species (25 larvae of each species for the low density
experiment and 75 of each species for high density). We collected larvae as described above
from the two species separately and combined them into a single vial with or without parasitoids.
We added six parasitoids (three female, three male) for 24 hours. We counted adults of both
species every 1-2 days as they emerged. Each interspecific treatment had five replicates, totaling
20 vials per species comparison. All experiments were incubated at room temperature.

Comparing parasitoid susceptibility across Z. indianus *populations*

To test for potential genetic differences in parasitoid susceptibility, we compared emergence
rates of *Z. indianus* from five Florida (FL) isofemale lines, and four Connecticut (CT) isofemale
lines collected in 2022 (Rakes et al. 2023). These isofemale lines were reared in the lab for
approximately 16 months (~20 generations) prior to the experiments described here. We added a
total of 25 *Z. indianus* larvae to each vial. We prepared three replicates of parasitized vials for

each line, resulting in a total of 27 vials. Two parasitoids (one female, one male) were added to
vials for 24 hours.

Assessing potential parasitoid host-switching

As *D. simulans* is a known suitable host of *L. heterotoma* (Quicray et al. 2023), we predicted that
L. heterotoma might initially oviposit in *D. simulans* before parasitizing *Z. indianus*. To assess
parasitoid foraging behavior, we compared the emergence rates of *Z. indianus* and *D. simulans*
following 1 hr, 8 hr, and 24 hr of parasitoid exposure. All vials contained a low density of 20
larvae (10 of each species) to increase the likelihood of host saturation. Two parasitoids (one
female, one male) were added to parasitized treatments. Non-parasitized treatments experienced
0 hr of parasitoid exposure. Each parasitoid duration treatment had three replicates (9 total vials),
and the non-parasitized controls had 12 replicates.

Data Analysis

All analyses were performed using the R statistical software (v. 4.4.0; R Core Team 2024). We
used the proportion of adult flies relative to starting larvae to calculate emergence rates of each
treatment. We used a generalized linear model (GLM) with a binomial distribution to compare *Z.*
indianus emergence rates for the intraspecific assays. Emerged adult flies were modeled as a
function of *density* (as a factor), *parasitoid presence*, and a *density*parasitoid* interaction. We
analyzed the interspecific data in separate models for each density and competing species
comparison. We used GLMs with a quasibinomial distribution to compare species emergence
rates between treatments. The quasibinomial distribution corrected for overdispersion in the
interspecific data (variance/mean ratios ranged from 5-10). Emerged adult flies were modeled as
a function of *species*, *parasitoid presence*, and a *species*parasitoid* interaction. We used a
generalized linear mixed model (GLMM) implemented in *lme4* (Bates et al. 2015) with a
binomial distribution to compare *Z. indianus* emergence rates from Florida (FL) and Connecticut
(CT) populations. The model included *location* as a main effect and *isofemale line* as a random
effect. We used a GLM with a binomial distribution to assess differences in *Z. indianus* and *D.*
simulans emergence rates at various parasitoid exposure times as a function of the *species*,
exposure time, and a *species*exposure time* interaction.

We calculated estimated marginal means (least squared means) for each model and
performed pairwise linear contrasts between groups using the *emmeans* package (Lenth et al.
2020) in R. We focused on comparing groups that differed by only a single variable (e.g. same
species but different parasitoid treatment, or same treatment but different species, Appendix
Table 1). We generated Tukey-adjusted linear contrast P values; statistical significance was
determined by $P_{adjusted} < 0.05$. All plots were generated with *ggplot* (Wickham 2016) and data
were managed with *data.table* (Dowle & Srinivasan 2012).

Results

Intraspecific competition assay

High density (GLM; Z = -8.326, P < 0.001) and presence of *L. heterotoma* (Z = -8.600, P <
0.001) significantly decreased *Z. indianus* emergence rates (Fig. 1A, Appendix Table 1). There
was a significant interaction between parasitoid presence and density, with parasitoids causing a
greater reduction in emerging adults at low density relative to high density (Z = 4.729, P <
0.001).

Interspecific competition assay

At low density, parasitoid presence significantly decreased the emergence rates of co-reared *Z.*
indianus and *D. hydei* (linear contrast of GLM; Z = 4.234, P < 0.001 & Z = 2.837, P = 0.024,
respectively; Fig. 1B top panel, Appendix Table 1). However, there were no significant
differences between *Z. indianus* and *D. hydei* emergence rates in either the absence or presence
of parasitoid wasps (Z = -0.654, P = 0.914 & Z = -1.748, P = 0.299, respectively). At high
density, the presence of parasitoids significantly decreased the emergence rates of *D. hydei* (Z =
7.402, P < 0.001; Fig. 1B bottom panel), but did not affect *Z. indianus* emergence rates (Z =
0.061, P = 0.999). *Z. indianus* had significantly lower emergence rates than *D. hydei* in the
absence of parasitoids (Z = 5.445, P < 0.001), but had significantly higher emergence rates in the
presence of parasitoids (Z = -4.383, P < 0.001).

At low density, parasitoid presence significantly decreased the emergence rates of co-
reared *Z. indianus* and *D. simulans* (Z = 6.360, P < 0.001 & Z = 4.924, P < 0.001, respectively;
Fig. 1C top panel, Appendix Table 1). However, *Z. indianus* emergence rates were significantly
higher than those of *D. simulans* in both the absence and presence of parasitoids (Z = -2.737, P =
0.032 & Z = -3.898, P < 0.001). At high density, parasitoid presence also significantly decreased
the emergence rates of both *Z. indianus* and *D. simulans* (Z = 6.288, P < 0.001 & Z = 4.586, P <
0.001, respectively; Fig. 1C bottom panel). There were no significant differences between *Z.*
indianus and *D. simulans* emergence rates in the absence of parasitoids (Z = -0.787, P = 0.861).
However, *Z. indianus* had significantly higher emergence rates than *D. simulans* in the presence
of parasitoids (Z = -3.098, P = 0.011).

Comparing parasitoid susceptibility across Z. indianus *populations*

There were no significant differences in adult emergence rates between parasitized Connecticut
and Florida populations (Fig. 2; Appendix Table 1; GLMM; P = 0.561).

Host switching assay

Exposure to parasitoids for one hour significantly decreased the emergence rates of *D. simulans*
(Z = -3.819, P < 0.001), but not *Z. indianus* (Z = -1.560, P = 0.119), when compared to non-
parasitized controls of each species (Fig. 3; Appendix Table 1). However, exposure to parasitoids

for 8 hours and 24 hours significantly decreased the emergence rates of both *D. simulans* ($P_{8h} <$
0.001, $P_{24h} < 0.001$) and *Z. indianus* ($P_{8h} < 0.001$, $P_{24h} = 0.004$).

Discussion

We explored the effect of enemy release on influencing competition outcomes by comparing the
adult emergence rates of a recent invasive insect pest, *Z. indianu*s, and co-occurring
cosmopolitan drosophilid species in the absence and presence of *L. heterotoma* parasitoid wasps.
Here, we demonstrate that consideration of competition interactions and environmental context is
essential when assessing the relative strength of enemy release. Our results suggest that enemy
release in the presence of competing species may contribute to the rapid population growth of *Z.*
indianus that is commonly observed in the field, and drosophilid density affects the relative
strength of enemy release in benefiting *Z. indianus* populations. We found that under most
circumstances, *Z. indianus* is an equal or better competitor than *D. hydei* and *D. simulans*, but
competition outcomes are widely dependent on factors such as parasitoid presence, density, and
competitor species. Our findings support the enemy release hypothesis because the emergence
rates of *Z. indianus* were equal to or higher than *D. hydei* and *D. simulans* in the presence of
parasitoids (Fig. 1B-C), indicating that parasitoids were often more detrimental to the established
species (a known host) than to *Z. indianus* (a novel host), especially at high densities. Further,
our results demonstrate the importance of taking a community approach when studying enemy
release because *Z. indianus* was susceptible to parasitization by *L. heterotoma*, but the relative
impact of this susceptibility depended on other contextual factors.

We observed that the relative strength of enemy release differed between densities,
suggesting the benefit of enemy release to *Z. indianus* is density-dependent. At low densities, the
addition of parasitoids did not alter competition outcomes for *Z. indianus* with *D. hydei* or *D.*
simulans. Therefore, at low densities, the benefit of enemy release is small for *Z. indianus* as
parasitoids have approximately equal effects on both competing species and do not alter
competition dynamics. However, at high densities, we observed that parasitoid presence shifted
the competition outcomes in favor of *Z. indianus* in both interspecific assays. Since competition
outcomes switched to benefit *Z. indianus* at high densities, but not low densities, we conclude
that enemy release was stronger at high densities compared to low densities.

A possible explanation for differing competition outcomes based on density is that *L.*
heterotoma switched from a known host (D. *simulans* or *D. hydei*) to a novel host (*Z. indianus)*.
As host switching is likely initiated by optimal foraging behavior (Křivan 1996), an increase in
the abundance of suitable hosts (e.g., high density) might decrease the number of parasitized
suboptimal hosts. We tested the possibility of host switching by examining the effects of
parasitoid exposure time on the emergence rates of *Z. indianus* and *D. simulans* (Fig. 3). Our
data suggest parasitoids initially oviposited in *D. simulans,* but switched to *Z. indianus* once

suitable hosts were saturated. The difference in parasitization rates might result from variations

in behavioral avoidance strategies between *Z. indianus* and *D. simulans* larvae, or *L. heterotoma*

may prefer to oviposit in *D. simulans*. Although we did not directly assess parasitoid host

preference, parasitoids are highly sensitive to host-selection and can accept or reject hosts based

on their suitability and quality (Wertheim 2022). *D. simulans* primarily uses lamellocytes for its

anti-parasitoid immune response, rather than multinucleated giant hemocytes like *Z. indianus*

(Salazar-Jaramillo et al. 2014, Cinege et al. 2020). Virulence factors of *L. heterotoma* have

evolved mechanisms to combat the lamellocyte immune response (McGonigle et al. 2017),

suggesting that species using lamellocytes may be more favorable hosts than species using

multinucleated giant hemocytes. Therefore, it is possible that the density-dependent results

observed in the interspecific assays reflect host switching behavior of *L. heterotoma*. Although

host-switching is commonly observed in generalist parasitoids (Murdoch 1969, Jones et al.

2015), specialist parasitoid species may be more reluctant to oviposit in suboptimal hosts due to

increased host selectivity. Therefore, evaluation of how different parasitoid species interact with

Z. indianus alongside other drosophilids would be beneficial for a broader understanding of

enemy release since parasitoid life-history strategies may produce different competition

outcomes.

Additionally, the strength of enemy release differed based on the competing species. At

high densities, the strength of enemy release was greatest when *Z. indianus* was competing with

D. hydei as the presence of parasitoids shifted competition outcomes from *Z. indianus* having

significantly lower emergence rates to significantly higher emergence rates than *D. hydei* (Fig.

1B). Although parasitoid presence also shifted competition outcomes in favor of *Z. indianus*

when competing with *D. simulans*, the magnitude of the shift was smaller as *Z. indianus* already

performed equal to *D. simulans* in the absence of parasitoids. Although the presence of

parasitoids decreased *D. simulans* and *D. hydei* survivorship to zero, parasitized larvae likely still

competed with *Z. indianus* larvae as host death does not occur until after the fly has pupated

(Quicray et al. 2023). We did not directly test for the presence of competition, and the density-

dependent effects on survival we observed may be due to other factors such as accumulation of

waste. However, a similar study directly tested for interspecific competition between *Z. indianus*

and *D. suzukii*, suggesting competition is present between other species (de Paiva Mendonça

2023). Further evaluation of how competing species' traits may alter the strength of enemy

release would be beneficial for identifying the specific mechanisms that influence enemy release

strength in community environments. These nuances emphasize how using a community

approach for testing the enemy release hypothesis is crucial as interpretation of the strength of

enemy release may differ based on factors such as density and competing species. Our results

extend the work of Brian & Catford (2023) as we found that enemy diversity and the relative

enemy impact can independently contribute to the strength of enemy release.

Interestingly, we observed that parasitoids significantly decreased the emergence rates of

Z. indianus in intraspecific assays (Fig. 1), and *Z. indianus* emergence rates were never higher

than 50% in the presence of parasitoids. Our *Z. indianu*s results differ from a study by Kacsoh et
al. (2014), which reported the majority (> 50%) of *Z. indianus* escaped parasitism by *L.*
heterotoma, but they are similar to a study by Cinege et al. (2023) which reported fewer than
30% of *Z. indianus* emerged in the presence of *L. heterotoma*. It is unlikely that genetic variation
between *Z. indianus* lines is responsible for these differences, as we found no reproducible
variation in emergence rates across isofemale lines (Fig. 2). Differences in laboratory conditions
or insect rearing may have produced different parasitization rates across studies. Within our
experiments, there were more vials with 0% emergence in the geographic variation experiment
(Fig. 2) compared to the low density treatment of the intraspecific experiments (Fig. 1A). It is
possible that a lower density of flies in the geographic variation experiment (N=25) led to a
faster saturation rate of larvae (Mitsui & Kimura, 2000). Alternatively, an uncontrolled
experimental factor, such as parasitoid health or age, may have affected parasitization efficiency
between experiments.

All experiments in this study were conducted under controlled laboratory conditions,
which limits applicability to field environments. We used larva-to-adult emergence rates as an
indicator for population success and did not directly measure parasitization, but our use of
control treatments allows us to infer that parasitization occurred. Additionally, we did not test
other fitness components such as growth, fecundity, and reproductive output, which may also be
used to assess enemy impact. Assessment of multigenerational effects using multiple fitness
factors may more comprehensively determine the impact of parasitoid presence on invader
population performance. In *D. melanogaster,* adult females that successfully defend themselves
from parasitism have several reduced fitness traits such as smaller body size and lower fecundity
(Fellowes et al. 1999). Additionally, investment in parasitoid defense mechanisms (e.g.,
increased hemocyte counts) is associated with tradeoffs in competitive ability in *D. melanogaster*
(Kraaijeveld et al. 2001). These tradeoffs might particularly affect *Z. indianus*, because of their
higher resistance to parasitoids due to the role of multinucleated giant hemocytes in their
immune response (Kacsoh et al. 2014, Cinege et al. 2023). Therefore, increased defense
mechanisms may lead to diminished fitness outcomes for *Z. indianus*, which could contribute to
drops in wild population sizes seen later in the season (Rakes et al. 2023). In addition to
physiological defense mechanisms, behavioral responses are also vital for reducing parasitism in
hosts (Hart 1990). Assessing whether the presence of parasitoids or competing species can
influence *Z. indianus'* oviposition or larval foraging behavior may provide insights into potential
behavioral resistance strategies of *Z. indianus* in field environments. Further evaluation of how
parasitism affects *Z. indianus* population dynamics over time would be beneficial to understand
the strength and frequency of enemy release in the field. Additionally, using a more natural
experimental design (e.g., natural fruit substrates, multiple co-occurring drosophilid species) to
mirror environmental heterogeneity could provide more information on how these results apply
to the field. Specifically, the use of artificial food substrates limited our evaluation of how larval
predator avoidance (e.g., escape mechanisms) may alter competition outcomes in parasitized
treatments. Other abiotic environmental conditions are also known to influence *L. heterotoma*

host selection (Roitberg et al. 1992). Unfavorable conditions such as changes in barometric
pressure and humidity may diminish foraging strategies, leading to increased parasitization of
suboptimal hosts in the field. Overall, complex and variable field conditions could potentially
produce different ecological outcomes than those observed in this study.

Conclusion

The enemy release hypothesis has raised controversy due to its mixed support within and
across taxa (Colautti et al. 2004, Liu & Stiling 2006, Brian & Catford 2023). Our findings further
emphasizes how varying contexts (e.g., density or competitor species) may alter species
interactions and consequently affect the interpretation of enemy release. Here, we demonstrate
that enemy release may contribute to the success of *Z. indianus*, and larval density influences the
relative strength of enemy release. We confirm that *L. heterotoma* effectively decreases the
emergence rates of *Z. indianus*, which may have major implications for management of *Z.*
indianus in the field. However, these benefits may be diminished when more suitable hosts are
present, especially at high larval densities. Due to the increasing threat of *Z. indianus* as an
agricultural pest, productive biological control may control the growth and spread of *Z. indianus*
populations. Verification of these results in a natural environment would be highly beneficial for
understanding how this invasive species interacts with native ecosystems, and for determining if
biological control could be an effective strategy for managing *Z. indianus* populations.

Figure Legends

Figure 1: Enemy release from *L. heterotoma* depends on competing species and larval
density in *Z. indianus*. A) Intraspecific competition assay results comparing emergence rates of
unparasitized and parasitized *Z. indianus* larvae. Gray labels indicate density as the total number
of larvae present. B) Interspecific competition assay results comparing *Z. indianus* and *D. hydei*
and C) *Z. indianus* and *D. simulans* adult emergence rates. The low density (top) is 50 total
larvae (25 of each species in B-C) and the high density (bottom) is 150 total larvae (75 of each
species in B-C). Asterisks indicate statistical significance by linear contrast of a generalized
linear model ($P_{adjusted} < 0.05$). Solid lines with asterisks indicate significant effects of parasitoids
within species. Dotted lines with asterisks indicate significant differences between species raised
in direct competition. N=5 replicate vials per treatment.

Figure 2: No geographic variation in *Z. indianus* susceptibility to *L. heterotoma*.
Adult emergence rates from parasitized Connecticut (CT) and Florida (FL) *Z. indianus*
isofemale lines. Boxplots indicate combined emergence rates of all lines and replicates
from each state. Points (with horizontal jitter added for clarity) indicate individual

replicate vials; colors indicate unique isofemale lines from CT (N =4 lines) or FL (N=5
lines) with three replicates of 25 larvae per line.

Figure 3: *L. heterotoma* parasitizes *D. simulans* more rapidly than it parasitizes *Z.*
***indianus*.** Interspecific assay comparing the emergence rates of *Z. indianus* (red bars) and
D. simulans (yellow bars) at different parasitoid exposure times. Hashed lines represent
control vials with no parasitoids added (N=12 total). Parasitized treatments had three
replicates for each exposure time. A density of 20 larvae (10 of each species) was used
for all vials. Error bars represent standard deviation. Asterisks indicate significant
differences between control and parasitized treatments by linear contrast ($P_{adjusted} < 0.05$).

Data Accessibility statement
All raw data used for these analyses and code to replicate the figures and statistics are available
at https://doi.org/10.5061/dryad.6hdr7sr8n. Data and code are also available at
https://github.com/camille-wa/Zaprionus-parasitoid-experiments.

Competing interests statement

The authors declare no competing interests.

Author contributions

CRW and PAE conceptualized the study and acquired funding. CRW developed methodology,
conducted all experiments, collected and analyzed all data, created figures, and wrote the
manuscript with supervision from PAE. CRW and PAE revised the manuscript.

Acknowledgements
446
We are grateful to Dr. Todd Schlenke for providing parasitoid wasp lines and rearing advice. We
thank Adam Lenhart and Dr. Alan Bergland for *D. simulans* lines. We thank Logan Rakes and
Weston Gray for assistance rearing *Z. indianus* lines. Dr. John Orrock and members of the
Erickson lab provided helpful feedback on this project and manuscript. This work was funded by
award # 1R15GM146208-01 to PAE from the National Institutes of Health, startup funds to PAE
from the University of Richmond, a University of Richmond Summer Fellowship to CRW, and a
Robert F. Smart Student Research Award to CRW.

Appendix

Appendix Table 1: Generalized linear model results for each experiment. Z-values are

reported for all experiments except for the interspecific competition assays (Fig. 1B-C), where T-

values are given in italics due to use of quasibinomial errors. Significant terms were determined

by $P < 0.05$ and are indicated in bold.

Experiment	Term	Estimate	SE	Z/T	P-value
Z. indianus intraspecific (Fig. 1A)	Intercept	1.325	0.174	7.632	**< 0.001**
	Parasitoid	-1.966	0.229	-8.600	**< 0.001**
	Density	-1.600	0.192	-8.326	**< 0.001**
	Parasitoid:Density	1.230	0.260	4.729	**< 0.001**
Z. indianus vs *D. hydei* low density (Fig. 1B-top)	Intercept	0.717	0.364	*1.971*	0.066
	Parasitoid	-5.538	1.952	*-2.837*	**0.012**
	Species	0.350	0.535	*0.654*	0.522
	Parasitoid:Species	3.084	2.036	*1.515*	0.149
Z. indianus vs *D. hydei* high density (Fig. 1B-bottom)	Intercept	-0.005	0.153	*-0.035*	0.973
	Parasitoid	-3.404	0.460	*-7.402*	**< 0.001**
	Species	-1.316	0.242	*-5.445*	**< 0.001**
	Parasitoid:Species	3.388	0.531	*6.380*	**< 0.001**
Z. indianus vs *D. simulans* low density (Fig. 1C-top)	Intercept	0.209	0.180	*1.159*	0.263
	Parasitoid	-5.029	1.021	*-4.924*	**< 0.001**
	Species	0.736	0.269	*2.737*	**0.015**
	Parasitoid:Species	3.256	1.059	*3.076*	**0.007**
Z. indianus vs *D. simulans* high density (Fig. 1C-bottom)	Intercept	-0.317	0.156	*-2.030*	0.059
	Parasitoid	-4.911	1.071	*-4.586*	**0.003**
	Species	0.173	0.220	*0.787*	0.443
	Parasitoid:Species	3.184	1.106	*2.879*	**0.011**
Z. indianus Florida vs Connecticut (Fig. 2)	Intercept	-2.753	0.400	-6.890	**< 0.001**
	State	-0.307	0.528	-0.581	0.561
Z. indianus vs. *D. simulans* exposure time (Fig. 3)	Intercept	0.547	0.219	2.499	**0.013**
	Species	1.063	0.358	2.973	**0.003**
	1 hour	-1.933	0.506	-3.819	**< 0.001**
	8 hours	-3.914	1.040	-3.762	**< 0.001**
	24 hours	-3.914	1.040	-3.762	**< 0.001**
	Species:1 hour	1.171	0.704	1.664	0.096
	Species:8 hours	1.899	1.141	1.665	0.096
	Species:24 hours	2.573	1.139	2.258	**0.024**

References

Allori Stazzonelli E, Funes CF, Corral González MN, Gibilisco SM, Kirschbaum DS. Population fluctuation and infestation levels of Zaprionus indianus Gupta (Diptera: Drosophilidae) in berry crops of northwestern Argentina. Acta Hortic. 2023;(1381):141-148. doi:10.17660/ActaHortic.2023.1381.19

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software, 67(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Boulétreau, M., P. Fouillet & R. Allemand. 1991. "Parasitoids Affect Competitive Interactions between the Sibling Species, Drosophila Melanogaster and D. Simulans" 74:171–77.

Bradshaw, Corey J. A., Boris Leroy, Céline Bellard, David Roiz, Céline Albert, Alice Fournier, Morgane Barbet-Massin, Jean-Michel Salles, Frédéric Simard, and Franck Courchamp. 2016. "Massive yet Grossly Underestimated Global Costs of Invasive Insects." *Nature Communications* 7 (1): 12986. https://doi.org/10.1038/ncomms12986.

Brian, Joshua I., and Jane A. Catford. 2023. "A Mechanistic Framework of Enemy Release." *Ecology Letters* 26 (12): 2147–66. https://doi.org/10.1111/ele.14329.

Clarke, David A., and Melodie A. McGeoch. 2023. "Invasive Alien Insects Represent a Clear but Variable Threat to Biodiversity." *Current Research in Insect Science* 4:100065. https://doi.org/10.1016/j.cris.2023.100065.

Cinege, Gyöngyi, Lilla B. Magyar, Henrietta Kovács, Viktória Varga, László Bodai, Nóra Zsindely, Gábor Nagy, Zoltán Hegedűs, Dan Hultmark, and István Andó. 2023. "Distinctive Features of *Zaprionus indianus* Hemocyte Differentiation and Function Revealed by Transcriptomic Analysis." *Frontiers in Immunology* 14 (December):1322381. https://doi.org/10.3389/fimmu.2023.1322381.

Colautti, Robert I., Anthony Ricciardi, Igor A. Grigorovich, and Hugh J. MacIsaac. 2004. "Is Invasion Success Explained by the Enemy Release Hypothesis?" *Ecology Letters* 7 (8): 721–33. https://doi.org/10.1111/j.1461-0248.2004.00616.x.

Comeault, Aaron A., Andreas F. Kautt, and Daniel R. Matute. 2021. "Genomic Signatures of Admixture and Selection Are Shared among Populations of *Zaprionus indianus* across the Western Hemisphere." *Molecular Ecology* 30 (23): 6193–6210. https://doi.org/10.1111/mec.16066.

Comeault, Aaron A, Jeremy Wang, Silas Tittes, Kristin Isbell, Spencer Ingley, Allen H Hurlbert, and Daniel R Matute. 2020. "Genetic Diversity and Thermal Performance in Invasive and Native Populations of African Fig Flies." Edited by John Parsch. *Molecular Biology and Evolution* 37 (7): 1893–1906. https://doi.org/10.1093/molbev/msaa050.

De Bruijn, Jessica A. C., Louise E. M. Vet, Hans M. Smid, and Jetske G. De Boer. 2022. "Effects of Oviposition in a Non-host Species on Foraging Behaviour of the Parasitoid *Cotesia Glomerata*." *Ecological Entomology* 47 (4): 668–78. https://doi.org/10.1111/een.13151.

De Paiva Mendonça, Larine, Khalid Haddi, and Wesley Augusto Conde Godoy. 2023. "Effects of
Co-Occurrence and Intra- and Interspecific Interactions between *Drosophila suzukii* and
Zaprionus Indianus." Edited by Kyung-Jin Min. *PLOS ONE* 18 (3): e0281806.
https://doi.org/10.1371/journal.pone.0281806.

Diagne, Christophe, Boris Leroy, Anne-Charlotte Vaissière, Rodolphe E. Gozlan, David Roiz, Ivan
Jarić, Jean-Michel Salles, Corey J. A. Bradshaw, and Franck Courchamp. 2021. "High and
Rising Economic Costs of Biological Invasions Worldwide." *Nature* 592 (7855): 571–76.
https://doi.org/10.1038/s41586-021-03405-6.

Dowle M, Srinivasan A. data.table: extension of 'data.frame'. R package, version 1.14.8. 2021.
https://CRAN.R-project.org/package=data.table.

EFSA Panel on Plant Health (PLH), Claude Bragard, Paula Baptista, Elisavet Chatzivassiliou,
Francesco Di Serio, Paolo Gonthier, Josep Anton Jaques Miret, et al. 2022. "Pest Categorisation
of *Zaprionus indianus*." *EFSA Journal* 20 (3). https://doi.org/10.2903/j.efsa.2022.7144.

Enders, Martin, Marc-Thorsten Hütt, and Jonathan M. Jeschke. 2018. "Drawing a Map of Invasion
Biology Based on a Network of Hypotheses." *Ecosphere* 9 (3): e02146.
https://doi.org/10.1002/ecs2.2146.

Fellowes, Kraaijeveld, and Godfray. 1999. "The Relative Fitness of *Drosophila Melanogaster*
(Diptera, Drosophilidae) That Have Successfully Defended Themselves against the Parasitoid
Asobara tabida (Hymenoptera, Braconidae)." *Journal of Evolutionary Biology* 12 (1): 123–28.
https://doi.org/10.1046/j.1420-9101.1999.00018.x.

Fleury, Frederic, Nicolas Ris, R. Allemand, P. Fouillet, Yves Carton, and M. Bouletreau. 2004.
"Ecological and Genetic Interactions in Drosophila–Parasitoids Communities: A Case Study
with D. Melanogaster, D. Simulans and Their Common Leptopilina Parasitoids in South-
Eastern France." *Genetica* 120 (April):181–94. https://doi.org/10.1007/978-94-007-0965-2_15.

Gardos, G., and J. O. Cole. 1976. "Maintenance Antipsychotic Therapy: Is the Cure Worse than the
Disease?" *The American Journal of Psychiatry* 133 (1): 32–36.
https://doi.org/10.1176/ajp.133.1.32.

Gioria, Margherita, and Bruce A. Osborne. 2014. "Resource Competition in Plant Invasions:
Emerging Patterns and Research Needs." *Frontiers in Plant Science* 5 (September).
https://doi.org/10.3389/fpls.2014.00501.

Gleason JM, Roy PR, Everman ER, Gleason TC, Morgan TJ. Phenology of Drosophila species
across a temperate growing season and implications for behavior. Desneux N, ed. PLoS ONE.
2019;14(5):e0216601. doi:10.1371/journal.pone.0216601

Godfray, H. C. J., and Masakazu Shimada. 1999. "Parasitoids as Model Organisms for Ecologists."
Population Ecology 41 (1): 3–10. https://doi.org/10.1007/PL00011980.

Hart, Benjamin L. 1990. "Behavioral Adaptations to Pathogens and Parasites: Five Strategies."
Neuroscience & Biobehavioral Reviews 14 (3): 273–94. https://doi.org/10.1016/S0149-
7634(05)80038-7.

Jones, Thomas S., Adam R. Bilton, Lorraine Mak, and Steven M. Sait. 2015. "Host Switching in a Generalist Parasitoid: Contrasting Transient and Transgenerational Costs Associated with Novel and Original Host Species." *Ecology and Evolution* 5 (2): 459–65. https://doi.org/10.1002/ece3.1333.

Kacsoh, Balint Z., Julianna Bozler, and Todd A. Schlenke. 2014. "A Role for Nematocytes in the Cellular Immune Response of the Drosophilid *Zaprionus indianus*." *Parasitology* 141 (5): 697–715. https://doi.org/10.1017/S0031182013001431.

Kacsoh, Balint Z., and Todd A. Schlenke. 2012. "High Hemocyte Load Is Associated with Increased Resistance against Parasitoids in Drosophila Suzukii, a Relative of D. Melanogaster." Edited by Robert DeSalle. *PLoS ONE* 7 (4): e34721. https://doi.org/10.1371/journal.pone.0034721.

Keane, Ryan M., and Michael J. Crawley. 2002. "Exotic Plant Invasions and the Enemy Release Hypothesis." *Trends in Ecology & Evolution* 17 (4): 164–70. https://doi.org/10.1016/S0169-5347(02)02499-0.

Kraaijeveld, A. R., J. J. M. Van Alphen, and H. C. J. Godfray. 1998. "The Coevolution of Host Resistance and Parasitoid Virulence." *Parasitology* 116 (S1): S29–45. https://doi.org/10.1017/S0031182000084924.

Kraaijeveld, A. R., E. C. Limentani, and H. C. J. Godfray. 2001. "Basis of the Trade–off between Parasitoid Resistance and Larval Competitive Ability in *Drosophila Melanogaster*." *Proceedings of the Royal Society of London. Series B: Biological Sciences* 268 (1464): 259–61. https://doi.org/10.1098/rspb.2000.1354.

Křivan, Vlastimil. 1996. "Optimal Foraging and Predator–Prey Dynamics." Theoretical Population Biology 49 (3): 265–90. https://doi.org/10.1006/tpbi.1996.0014.

Lachaise D, Cariou ML, David JR, Lemeunier F, Tsacas L, Ashburner M. Historical Biogeography of the Drosophila melanogaster Species Subgroup. In: Hecht MK, Wallace B, Prance GT, eds. Evolutionary Biology. Springer US; 1988:159-225. doi:10.1007/978-1-4613-0931-4_4

Lenth R, Singmann H, Love J, Buerkner P, Herve M (2020) Package 'emmeans' R topics documented. Am Stat 34:216–22

Liu, Hong, and Peter Stiling. 2006. "Testing the Enemy Release Hypothesis: A Review and Meta-Analysis." *Biological Invasions* 8 (7): 1535–45. https://doi.org/10.1007/s10530-005-5845-y.

McGonigle, John E., Alexandre B. Leitão, Sarah Ommeslag, Sophie Smith, Jonathan P. Day, and Francis M. Jiggins. 2017. "Parallel and Costly Changes to Cellular Immunity Underlie the Evolution of Parasitoid Resistance in Three Drosophila Species." Edited by Elizabeth Ann McGraw. *PLOS Pathogens* 13 (10): e1006683. https://doi.org/10.1371/journal.ppat.1006683.

Mitsui H, Kimura MT. Coexistence of drosophilid flies: Aggregation, patch size diversity and parasitism. Ecological Research. 2000;15(1):93-100. doi:10.1046/j.1440-1703.2000.00328.x

Mooney HA, Cleland EE. The evolutionary impact of invasive species. Proc Natl Acad Sci USA. 2001;98(10):5446-5451. doi:10.1073/pnas.091093398

Murdoch, William W. 1969. "Switching in General Predators: Experiments on Predator Specificity
and Stability of Prey Populations." *Ecological Monographs* 39 (4): 335–54.
https://doi.org/10.2307/1942352.
Pfeiffer, Douglas G, Meredith E Shrader, James C E Wahls, Brittany N Willbrand, Ian Sandum, Kim
Van Der Linde, Curtis A Laub, Ryan S Mays, and Eric R Day. 2019. "African Fig Fly (Diptera:
Drosophilidae): Biology, Expansion of Geographic Range, and Its Potential Status as a Soft
Fruit Pest." *Journal of Integrated Pest Management* 10 (1): 20.
https://doi.org/10.1093/jipm/pmz018.
Quicray, Maude, Léonore Wilhelm, Thomas Enriquez, Shulin He, Mathilde Scheifler, and Bertanne
Visser. 2023. "The *Drosophila*-parasitizing Wasp *Leptopilina heterotoma* : A Comprehensive
Model System in Ecology and Evolution." *Ecology and Evolution* 13 (1): e9625.
https://doi.org/10.1002/ece3.9625.
R Core Team. R: a language and environment for statistical computing, version 4.4.0. Vienna
(Austria): R Foundation for Statistical Computing; 2024. https://www.R-project.org/.
Rakes, Logan M, Megan Delamont, Christine Cole, Jillian A Yates, Lynsey J Blevins, Fatima
Naureen Hassan, Alan O Bergland, and Priscilla A Erickson. 2023. "A Small Survey of
Introduced *Zaprionus indianus* (Diptera: Drosophilidae) in Orchards of the Eastern United
States." Edited by Brian Aukema. *Journal of Insect Science* 23 (5): 21.
https://doi.org/10.1093/jisesa/iead092.
Rohlfs, Marko, and Thomas S. Hoffmeister. 2004. "Spatial Aggregation across Ephemeral Resource
Patches in Insect Communities: An Adaptive Response to Natural Enemies?" *Oecologia* 140
(4): 654–61. https://doi.org/10.1007/s00442-004-1629-9.
Roitberg, Bernard D., Marc Mangel, Robert G. Lalonde, Carol A. Roitberg, Jacques J. M. Van
Alphen, and Louise Vet. 1992. "Seasonal Dynamic Shifts in Patch Exploitation by Parasitic
Wasps." *Behavioral Ecology* 3 (2): 156–65. https://doi.org/10.1093/beheco/3.2.156.
Roy, Helen E., Elena Tricarico, Richard Hassall, Charlotte A. Johns, Katy A. Roy, Riccardo Scalera,
Kevin G. Smith, and Bethan V. Purse. 2023. "The Role of Invasive Alien Species in the
Emergence and Spread of Zoonoses." *Biological Invasions* 25 (4): 1249–64.
https://doi.org/10.1007/s10530-022-02978-1.
Salazar-Jaramillo, Laura, Angeliki Paspati, Louis Van De Zande, Cornelis Joseph Vermeulen, Tanja
Schwander, and Bregje Wertheim. 2014. "Evolution of a Cellular Immune Response in
Drosophila: A Phenotypic and Genomic Comparative Analysis." *Genome Biology and*
Evolution 6 (2): 273–89. https://doi.org/10.1093/gbe/evu012.
Sturtevant AH. The North American Species of Drosophila. The Carnegie institution of Washington;
1921. doi:10.5962/bhl.title.33564
Van Der Linde, Kim, Gary J. Steck, Ken Hibbard, Jeffrey S. Birdsley, Linette M. Alonso, and David
Houle. 2006. "First Records of *Zaprionus indianus* (Diptera: Drosophilidae), a Pest Species on

Commercial Fruits from Panama and the United States of America" *Florida Entomologist* 89 (3): 402–4. https://doi.org/10.1653/0015-4040(2006)89[402:FROZID]2.0.CO;2.

Vilela, Carlos Ribeiro. 1999. "Is *Zaprionus indianus* Gupta, 1970 (Diptera, Drosophilidae) Currently Colonizing the Neotropical Region?" *Drosophila Information Service* 82:37–39. https://repositorio.usp.br/item/001064912.

Werner, Thomas, Tessa Steenwinkel, and John Jaenike. 2018. *The Encyclopedia of North American Drosophilids Volume 1: Drosophilids of the Midwest and Northeast*. Michigan Technological University.

Wertheim, Bregje. 2022. "Adaptations and Counter-Adaptations in Drosophila Host–Parasitoid Interactions: Advances in the Molecular Mechanisms." *Current Opinion in Insect Science* 51 (June):100896. https://doi.org/10.1016/j.cois.2022.100896.

Wickham H. ggplot2: elegant graphics for data analysis. New York (NY): Springer-Verlag; 2016.

Zhao L, Begun DJ. Genomics of parallel adaptation at two timescales in Drosophila. Matzkin L, ed. PLoS Genet. 2017;13(10):e1007016. doi:10.1371/journal.pgen.1007016