Title: Serum amyloid A3 deficiency modulates aortic immune composition and

attenuates murine atherosclerosis progression

Authors: Tzu-Yin Chou[1,2], Yu-Zhen Ye[1,2], Chia-Jung Lien[1], Jui-Chun Wang[1], Chen-

Hsuan Yang[1], Yen-Ting Chen[1], Pei-An Chao[1], Jian-Da Lin[1]*

Affiliations:

[1]Department of Biochemical Science and Technology, National Taiwan University,

Taipei 10617, Taiwan

[2]These authors contributed equally.

* Correspondence to:

jdlin@ntu.edu.tw

Abstract

Atherosclerosis is a growing concern in developed nations, necessitating the identification of therapeutic targets for advancing personalized medicine. Serum amyloid A3 (*Saa3*) has been linked to accelerated plaque progression by affecting cholesterol metabolism and modulation of inflammation. We hypothesize that knocking out *Saa3* (*Saa3*$^{-/-}$) could mitigate plaque development by regulating aortic immune cell compositions during atherosclerosis progression. Using a murine model, we induced atherosclerosis via a gain-of-function mutant PCSK9-encoding adeno-associated viral vector (AAVmPCSK9) in female wild-type (WT) and *Saa3*$^{-/-}$ mice. *Saa3*$^{-/-}$ mice developed smaller plaques than WT mice, and single-cell RNA sequencing revealed significant differences in aortic immune cell populations, particularly among aortic macrophages. Aortic macrophages in atherosclerotic *Saa3*$^{-/-}$ mice represent an anti-inflammatory and tissue-repairing phenotype and the Trem2^{hi} macrophages, characterized by high *Gpnmb*, *Lpl*, and *Spp1* expressions, predominated over the typical foamy macrophages in *Saa3*$^{-/-}$ compared to WT mice. Notably, SAA3 regulates cholesterol metabolism and inflammatory responses in foamy macrophages. Our study highlights *Saa3* as a key modulator of aortic immune cells that impact atherosclerosis progression.

Introduction

Atherosclerosis, a major cardiovascular concern in developed countries [1], is a chronic arterial disease characterized by plaque formation within arterial walls. This process involves complex interplays among various cellular components and inflammatory mediators that drive inflammation and lipid accumulation. Atherosclerosis initiation involves low-density lipoprotein (LDL) accumulation in the coronary artery intima, which upon oxidation to ox-LDL by risk factors, activates nuclear factor kappa B (NF-κB) [2, 3]. This leads to the expression of adhesion molecules like vascular cell adhesion molecule-1 (VCAM-1) and intercellular adhesion molecule-1 (ICAM-1) on the endothelium, aiding monocyte recruitment into the vascular wall [4]. These recruited monocytes then convert into macrophages under the influence of macrophage colony-stimulating factor (M-CSF), and release inflammatory cytokines such as interleukin (IL)-1β, IL-12, and tumor necrosis factor (TNF)-α, escalating the inflammatory response [5]. As the disease progresses, these macrophages become foam cells by uptaking ox-LDL, which is crucial for plaque formation and intimal thickening.

Macrophages are the predominant type of immune cell found in aortic plaques but display a wide range of phenotypes and functions. At the onset of atherosclerosis,

macrophages ingest ox-LDL via receptors like CD36 and SR-A, forming cholesterol-

rich foam cells. Continued lipid intake without effective cholesterol removal leads to

foam cell death, creating a necrotic core in advanced plaques [6, 7]. Single-cell RNA

sequencing (scRNA-seq) of monocyte-derived macrophages has revealed that

macrophage heterogeneity is beyond M1/M2 definitions and associated with different

states of atherosclerosis, where progression is linked to more complex states than

regression [8]. This technique has revealed various macrophage subsets within

atherosclerotic plaques, each characterized by unique markers and roles contributing

to aortic plaque progression or regression. Notably, three main groups are identified:

tissue-resident-like macrophages, marked by *Folr2* and *Lyve1*, which promote tissue

stability; inflammatory macrophages, with markers such as *Tnf* and *Il1b*, which

activate immune responses; and anti-inflammatory TREM2[hi] macrophages, associated

with cholesterol metabolism and identified by markers such as *Trem2* [9, 10]. This

diversity in macrophage functions underscores the complexity of atherosclerosis and

offers multiple targets for therapy.

The Serum Amyloid A (SAA) protein family, including SAA3, represents a group of

acute-phase proteins central to immune responses and inflammatory processes. Unlike

humans, who lack a functional SAA3 protein due to a premature stop codon and

resulting frameshift mutation that renders the SAA3 gene a pseudogene [11], mice

express a functional SAA3 and mouse SAA3 shares 70% amino acid homology with

human SAA1 [12]. Its tissue distribution and expression patterns, particularly in

adipose tissue, align more closely with human SAA1 than with mouse SAA1 or

SAA2, suggesting a functional similarity between mouse SAA3 and human SAA1

[12]. Furthermore, sex-dimorphisms have been reported in the context of

atherosclerosis, SAA3-associated inflammatory responses, and microbiota-mediated

regulation of immune and metabolic processes. Smit et al. (2024) found that in aged

atherosclerotic *Ldlr*[-/-] mice, plaque volume was similar between males and females,

but female mice presented plaques with higher collagen and cholesterol crystal

content, along with a smaller necrotic core and fewer macrophages compared to males

[13]. *Saa3* has been shown to have sexually dimorphic effects on atherosclerosis, with

deficiency being atheroprotective in male mice but being proatherogenic in female

mice [14]. This response is suggested to be influenced by sex steroids, as 17β-

estradiol promotes a pro-inflammatory profile in male *Saa3*[-/-] macrophages [14].

However, another study indicates a significant upregulation of SAA3, but not SAA1

or SAA2, in the adipose tissue of conventional mice, compared to germ-free mice

[15]. The presence of intestinal microbiota significantly upregulated SAA3 expression

in adipose tissue and colonic tissue, with evidence indicating that its colonic

regulation is partially mediated through the Toll-like receptor (TLR)/MyD88/NF-κB

signaling axis [12]. These data suggest that *Saa3*-associated sex-specific differences

in atherosclerosis progression and inflammatory responses may result from its

interaction with confounding factors, such as the microbiota, which modulate SAA3

levels and sex dimorphism in atherosclerosis.

Previous studies suggest that mouse *Saa3* modulates local injury and inflammatory

responses, as its mRNA and protein expression in macrophages increases with the

dose of lipopolysaccharide (LPS) [12]. It also indicates that increased SAA expression

correlates with the aortic plaque progression in *ApoE$^{-/-}$* mice, accompanied by

elevated levels of IL-6 and TNF-α. This upregulation potentially facilitates the

infiltration of M1 macrophages into atherosclerotic regions, thereby promoting

atherosclerosis progression [16]. Mouse *Saa3* is actively expressed, predominantly in

macrophages and adipose tissues [17], and the activation of the *Saa3* promoter is

regulated through C/EBPβ signaling, which in turn activates macrophages,

underscoring the complex interaction within adipose tissue [18]. Pulmonary exposure

to nanoparticles induces pronounced acute phase responses, with mouse *Saa3*

identified as the most differentially regulated gene, demonstrating prolonged

upregulation in lung tissue and correlating with neutrophil infiltration and systemic

SAA3 protein levels [19]. Notably, SAA3 has been reported to promote

atherosclerosis, and overexpressed SAA3 increases atherosclerotic plaque size [20].

Overall, these studies support the role of SAA3 in regulating macrophage function

during atherosclerosis, though the specific type of aortic macrophages that regulate

aortic plaque progression and the underlying mechanisms remain elusive.

To assess how *Saa3* deficiency impacts aortic immune cell composition and

functionalities in contributing to atherosclerosis progression, we applied scRNA-seq

using the BD Rhapsody platform on immune cells from aortic arches of female *Saa3*

knockout (*Saa3$^{-/-}$*) mice and littermate wild-type (WT) control mice during

atherosclerosis. Our investigation seeks to delineate the impact of *Saa3* deficiency on

the phenotypic and functional landscape of aortic macrophages and other immune cell

populations. By comparing the immune cell profiles between *Saa3$^{-/-}$* and WT mice,

we aim to uncover the molecular mechanisms by which SAA3 influences macrophage

or other immune cell types-mediated aortic plaque progression.

Results

***Saa3* deletion reduced aortic plaque progression but increased peripheral lipid**

levels

To determine the phenotype of atherogenesis after *Saa3* deficiency, we utilized a

C57BL/6 wild-type (WT) mouse-based atherosclerosis model independent of *Ldlr$^{-/-}$*

mice breeding [8, 21, 22]. Plaque progression was initiated by injecting the mice with

an adeno-associated viral vector expressing a gain-of-function mutant of protein

convertase subtilisin/kexin type 9 (AAVmPCSK9), resulting in LDL receptor

deficiency and hypercholesterolemia. Both female C57BL/6 WT and C57BL/6 *Saa3*[-/-]

mice were treated with AAVmPCSK9 and fed the Western diet (WD) for 18 weeks to

induce hypercholesterolemia and aortic plaque formation. We longitudinally collected

plasma from mice at 0, 2, 4, 8, 12, and 16 weeks on WD, and the immune profiling

was performed on sorted CD45+ cells from aortic arches by single-cell RNA-seq at

18 weeks on WD (Fig. 1a). We first determined total cholesterol levels in the plasma

of WT and *Saa3*[-/-] mice and a baseline comparison at weeks 0 revealed no significant

differences, with levels below 100 mg/dL (Fig. 1b). Over time, *Saa3*[-/-] mice

developed higher cholesterol levels than WT, with a slight increase at weeks 2

through 12, and a significant rise by weeks 16 (Fig. 1b).

Research demonstrates that serum amyloid A (SAA) attenuates the anti-

inflammatory and antioxidative functions of high-density lipoprotein (HDL) by

facilitating its binding to adipocyte and macrophage-derived glycoproteins [23]. We

conducted high-density lipoprotein (HDL)/ low-density lipoprotein LDL analyses that

revealed an increase in peripheral HDL levels during the initial phase of

atherosclerosis progression (Fig. 1c). Not only did HDL levels demonstrate a rising

trend with statistically significant increases in the *Saa3*[-/-] group compared to WT at

weeks 2 and 4, but the absence of *Saa3* also correlated with a potential increase in

LDL levels in the bloodstream (Fig. 1c).

Overexpression or suppression of SAA3 can increase or decrease aortic lesion area

in *ApoE*[-/-] mice [20]. We also determined aortic plaque sizes between WT and *Saa3*[-/-]

mice by histologically evaluating aortic root sections stained with Oil Red O (Fig.

1d). It demonstrated a significant decrease in plaque area in *Saa3*[-/-] compared to WT

mice, with mean values of 31.50% and 45.48%, respectively (Fig. 1e). These

observations demonstrate that *Saa3* deficiency increases peripheral lipid levels but

attenuates aortic plaque formation, suggesting the complex roles of *Saa3* in

modulating lipid metabolism and atherogenesis.

159 *Saa3* deficiency alters intercellular immune communications in aortic arches

160 during atherosclerosis progression

All immune cells express CD45, a major transmembrane glycoprotein expressed on

all hematopoietic cells [24] and some cardiac fibroblasts [25]. To elucidate the

immunological impact of *Saa3* deficiency within the aortic arch and its putative role

in the pathogenesis of atherosclerosis, we performed a single-cell RNA sequencing on

FACS-purified CD45+ cells (Supplementary Fig. 1). After data processing, possible

contaminated cell removal, anchor-based dataset integration [26], and cell

clusters visualized by Uniform Manifold Approximation and Projection (UMAP), it

revealed the immune cell compositional variances in aortic arches between female

WT and *Saa3*$^{-/-}$mice during atherosclerosis progression (Fig. 2). Utilizing the cell

annotation tool by *SingleR* [27], we classified the cell phenotypes, enabling initial

taxonomical segregation within the distinct clusters (Fig. 2a). Our analysis identified

9 cell subpopulations, encompassing B cells, dendritic cells (DCs), innate lymphoid

cells (ILCs), macrophages, monocytes, neutrophils, natural killer T (NKT) cells, T

cells, and γδ T cells (Tgd) (Fig. 2a). Despite similar UMAP topologies between WT

and *Saa3*$^{-/-}$ aortic cell populations (Fig. 2a), *Saa3*$^{-/-}$ mice showed increased T cells, B

cells, monocytes, γδ T cell, and neutrophils proportions but decreased macrophages,

innate lymphoid cells (ILC), and NKT cells compared to WT mice (Fig. 2b).

Heatmaps of differential gene expression profiles supported these *SingleR*-annotated

cell type differences, highlighting the top 5 marker genes per subset (Fig. 2c).

Cellchat has been used to infer, analyze, and visualize cell-cell communication

networks in scRNA-seq datasets [27]. To dissect the variances in signaling dynamics

in aortic cell types between WT and *Saa3*$^{-/-}$ mice, we employed *CellChat* to evaluate

the intercellular communication networks and assess the number and intensity of

cellular interactions within the respective cohorts (Figs. 2d-2e). The resultant metrics

demonstrated 1,822 and 1,787 interactions across all cell types in aortic arches

between WT and *Saa3*$^{-/-}$ mice, with interaction strength for *Saa3*$^{-/-}$ cell types

quantified at 83.957 compared to 64.762 for WT (Fig. 2d). To rigorously dissect the

discrete intercellular communication patterns responsible for the observed differential

signaling, we interrogate the heatmaps that delineate both the frequency and

magnitude of interaction strengths across aortic cell types (Fig. 2e). Increased

magnitude of intercellular interaction strengths was observed in the aortic arches of

Saa3$^{-/-}$ mice compared to their WT mice (Fig. 2d). The increased interaction

magnitudes were primarily directed from most immune cells toward γδ T cells (Tgd),

while reduced interactions were directed from macrophages toward other immune

cells in *Saa3*$^{-/-}$ mice compared to WT mice (Fig. 2e). These patterns suggest that the

absence of the *Saa3* gene exerts a profound regulatory influence on the intercellular

communication network among immune cells, potentially revealing critical insights

into the regulatory mechanisms of immune function within the aortic niche.

***Saa3* depletion increases biological processes in aortic macrophages that are**

highly associated with lipid metabolism and inflammatory recruitment during

atherosclerosis

CellChat analysis revealed significant differences in cell-cell interaction signals

between aortic macrophages and other aortic immune cell populations within the

$Saa3^{-/-}$ and WT group (Figs. 2d-2e). To delve deeper, we assessed the transcriptomic

variances in macrophages from $Saa3^{-/-}$ versus WT mice. Our volcano plot delineated

distinct gene expression profiles, with genes such as *Fabp5* and *Gpnmb* are

upregulated in $Saa3^{-/-}$ mice, exceeding the established $\log_2$ fold change threshold of

>1 and p-values < 0.01 (Fig. 2f). *Fabp5*, which plays a central role in the intracellular

transport of lipids, is integral to the uptake, transport, and metabolism of fatty acids

[28]. *Gpnmb* has been shown to protect against obesity-related metabolic disorders by

mitigating macrophage inflammatory capacity [29]. Conversely, genes such as

Cxcl13, *Lyz1*, *Tmem176a*, *Clec4a1*, *C3*, and *Cfp* are downregulated in $Saa3^{-/-}$

macrophages (Fig. 2f). Studies have demonstrated that a lack of *Cxcl13* correlates

with a diminished presence of atherosclerotic conditions, likely due to lower

recruiting inflammatory granulocytes within the lesions [30] [31]. *C3* expression in

macrophages has been linked to exacerbating atherosclerosis by limiting macrophage

efferocytosis, contributing to plaque development and inflammation [32]. *Cfp* has

been linked to regulating complement activation and orchestrating iron homeostasis

[33]. Gene set enrichment analysis, conducted using the *ClusterProfiler* package [34],

revealed distinct gene ontology (GO) pathways in aortic macrophages (Fig. 2g). WT

aortic macrophages were enriched for pathways associated with enhanced immune

cell recruitment and chemotaxis, whereas $Saa3^{-/-}$ aortic macrophages exhibited

enrichment for pathways related to lipid localization and transport (Fig. 2g,

Supplementary Table 1). Overall, these data suggest the phenotypes of aortic

macrophages were impacted by reduced recruitment of inflammatory macrophages,

which helped mitigate inflammation potentially influencing lesion formation and

stability.

Altered macrophage compositions and enhanced Trem2^{hi} macrophage presence

linked to diminished aortic plaque progression in *Saa3$^{-/-}$* mice

Macrophage (Mφ) polarization, along with several distinct clusters, has been

identified as increasingly prevalent during the atherosclerosis progression than

regression [8, 35]. To dissect the functional heterogeneity of aortic macrophage

populations in WT and *Saa3$^{-/-}$* mice during atherosclerosis, we classified macrophages

into subpopulations and compared the top differentially expressed genes from each

cluster against our previously published dataset, where we profiled single-cell sorted

Cx3cr1-derived aortic macrophages from lineage-tracing mice during atherosclerosis

progression and regression (hereinafter referred to as "*Lin regression study*")[8].

There are five distinct macrophage clusters in the aortic arches of WT and *Saa3$^{-/-}$*

mice during atherosclerosis (Fig. 3a). CD74^{hi}MHCII^{hi} Mφ is the most abundant

macrophage subset in both genotypes, with CD74^{hi}MHCII^{hi}, Folr2^{hi}, and

Retnla^{hi}Ear2^{hi} Mφ being more prevalent in WT mice compared to *Saa3$^{-/-}$* mice (Fig.

3b). Conversely, Trem2^{hi} and Apbb2^{hi} Mφ are found in higher percentages within the

aortic arches of *Saa3$^{-/-}$* mice than WT mice (Fig. 3b).

The Cd74^{hi}MHCII^{hi} Mφ, a subset identified through the *Lin regression study* [8], is

characterized by high expression of genes such as *Tmem176b*, *Lilra5*, *H2-Eb1*, and

H2-Aa (Supplementary Fig. 2a and Table 2). Folr2^{hi} Mφ exhibit unique transcriptional

profiles with *F13a1*, *Pf4*, *Folr2*, and *Lyve1* (Supplementary Fig. 2a and Table 2).

These markers are indicative of a tissue-resident-like macrophage phenotype [36-39]

and align with the molecular signature of the Folr2[hi] macrophage cluster previously

delineated in the *Lin regression study* [8]. Retnla[hi]Ear2[hi] Mφ is distinguished by their

elevated expression of genes such as *Retnla* and *Ear2*, which suggest an early-stage

transition of monocytes to macrophages within tissues [40, 41]. Additionally, these

macrophages show enrichment in genes like *Gm2a* and *Fn1*, confirming their

specialized state as identified in the *Lin regression study* [8]. Trem2[hi] Mφ is defined

by a gene expression signature that mirrors the population described by *Cochain et al.*

[42] and *Lin regression* [8] studies and is exemplified by heightened expression of

Ctsd, Lgals3, Spp1, and *Lpl* (Supplementary Fig. 2a and Table 2). Apbb2[hi] Mφ is

characterized by the predominant expression of *Cd9, Cd72*, and *Hexb* genes. While

sharing traits with the "Trem2[hi] Mφ," they are uniquely distinguished by their

expression of *Apbb2* (Supplementary Fig. 2a and Table 2).

On the other hand, we identified differentially expressed genes (DEGs) across five

macrophage populations and conducted gene ontology (GO) enrichment analysis on

biological pathways to explore potential functions (Supplementary Fig. 2b and Table

3). Compared to other macrophage clusters, Cd74[hi]MHCII[hi] Mφ in the aortic arches

showed significant enrichment in pathways related to leukocyte-mediated immunity,

positive regulation of cell activation, activation of the immune response, and

immunoglobulin-mediated immune responses (Supplementary Fig. 2b and Table 3).

Folr2[hi] Mφ is distinguished by activation of immune response, ameboidal-type cell

migration, cell chemotaxis, regulation of angiogenesis, and regulation of viral

processes (Supplementary Fig. 2b and Table 3). Retnla[hi]Ear2[hi] Mφ were noted for

their leukocyte proliferation, mononuclear cell proliferation, positive regulation of

leukocyte cell-cell adhesion, positive regulation of T cell activation, and response to

type II interferon (Supplementary Fig. 2b and Table 3). Trem2[hi] Mφ is characterized

by the generation of precursor metabolites and energy, aerobic respiration, oxidative

phosphorylation, purine ribonucleoside triphosphate biosynthetic process, and ATP

biosynthetic process (Supplementary Fig. 2b and Table 3). These functions highlight

the metabolic specialization of Trem2^{hi} Mφ in energy production and biosynthesis.

Apbb2^{hi} Mφ demonstrated upregulation in myeloid cell differentiation, leukocyte

migration, actin filament organization, regulation of protein-containing complex

assembly, and mononuclear cell migration (Supplementary Fig. 2b and Table 3).

In *Lin regression study* [8], Cd74^{hi}MHCII^{hi} Mφ and Retnla^{hi}Ear2^{hi} Mφ are abundant

in aortic arches of progression group and we also observed these two subsets are

higher in aortic arches of WT mice compared to *Saa3$^{-/-}$* mice (Fig. 3b). *Folr2*

expression is increased in human atherosclerotic plaques compared to normal arterial

tissue[43] and we also observed aortic Folr2^{hi} Mφ are higher in WT mice than *Saa3$^{-/-}$*

mice (Fig. 3b). Trem2^{hi} Mφ exhibits comparable prevalence between the progression

and regression cohorts in the *Lin regression study* [8], whereas both Trem2^{hi} and

Apbb2^{hi} Mφ are more abundant in the aortic arches of *Saa3$^{-/-}$* mice compared to WT

mice (Fig. 3b). Trem2^{hi} Mφ is previously identified as lipid-laden foam cells within

atherosclerotic lesions, underscoring their capacity for lipid processing and

phagocytosis [9, 10, 35]. They are associated with anti-inflammatory responses,

which may help prevent the establishment of a proinflammatory environment within

the plaque [9, 35, 44]. Similar to previous findings, Trem2^{hi} Mφ expresses high *Spp1,*

a gene associated with fibrotic responses indicative of tissue repair mechanisms [45,

46] (Fig. 3c, Supplementary Fig. 2a). Additionally, this subset expresses *Gpnmb*, a

gene implicated in anti-inflammatory processes [29, 47], along with *Fabp5*, which is

involved in the uptake and transport of fatty acids [28] (Fig. 3c). Notably, enzymes

such as *Lpl*, central to lipid metabolism[48] , and *Mmp12*, a contributor to

extracellular matrix remodeling [49], further characterize this population (Fig. 3c).

Hence, these data indicate that *Saa3*$^{-/-}$ mice exhibit higher compositions of aortic

macrophages with tissue-repairing and anti-inflammatory properties, leading to

reduced aortic plaque progression.

***Saa3* modulates aortic macrophage functionalities during atherosclerosis**

Investigating the transition of macrophage phenotypes from M1-like to M2-like as a

therapeutic strategy recognizes the complexity of aortic macrophage phenotypes in

atherosclerosis, which extend beyond conventional M1 and M2 polarization states [8,

35]. We use typical markers for Chemokine, Interferon (IFN), Inflammatory, and

Type II responses to visualize the aortic macrophage subsets from WT and *Saa3*$^{-/-}$

mice (Fig. 3d). Among the comparisons between WT and *Saa3*$^{-/-}$ aortic Mφ,

pronounced differences were noted in the increased expression levels of chemokines

including *Cxcl1*, *Ccl2*, *Ccl3*, and *Ccl4* in *Saa3*$^{-/-}$ than WT Folr2^{hi} Mφ (Fig. 3d).

Chemokines are signaling molecules that attract various types of immune cells to sites

of infection or injury. Increased expression of chemokines by tissue-resident

macrophages enhances the recruitment of neutrophils and lymphocytes, which is

crucial for initiating and sustaining an effective immune response [50]. Additionally,

there was a higher expression of type II response markers including *Cd163, Mrc1*, and

Msr1 [51-54] in *Saa3*$^{-/-}$ than WT Folr2^{hi} Mφ (Fig. 3d). Moreover, Il10ra showed

higher expression in *Saa3*$^{-/-}$ than WT Cd74^{hi}MHCII^{hi} Mφ, and Il10ra is required for

protection against atherosclerosis in low-density lipoprotein receptor knockout mice

[55]. Furthermore, we observed higher expression levels of interferon regulatory and

downstream markers such as *Irf5, Irf7, Isg15*, and *Stat1* in WT than *Saa3*$^{-/-}$

Retnla^{hi}Ear2^{hi} Mφ (Figure 3d), which may indicate an enhanced interferon response in

these cells [56, 57]. These expression patterns indicate a significant alteration in gene

expression profiles and macrophage composition between the two genotypes,

suggesting that deletion of the *Saa3* may modulate macrophage recruitment, type 2

immune responses, and inflammatory/anti-inflammatory phenotypes.

Saa3 modulates aortic macrophage communications that impact inflammatory

processes

To determine the role of *Saa3* in aortic macrophage communication during

atherosclerosis, we also performed a comparative analysis of cell interactions using

CellChat [27] in both WT and *Saa3$^{-/-}$* macrophages. Our findings indicate a

substantial decline in intercellular communication in *Saa3$^{-/-}$* macrophages, with

interaction events decreasing from 482 in WT to 290 in *Saa3$^{-/-}$* cells (Fig. 3e).

Correspondingly, we observed a significant drop in the strength of these interactions,

with the average interaction strength in WT macrophages being 17.997 compared to a

lower average of 14.348 in *Saa3$^{-/-}$* macrophages, mirroring the decrease in the number

of interaction events (Fig. 3e).

The heatmap depicts the overall signaling patterns observed in various macrophage

subpopulations from WT and *Saa3$^{-/-}$* mice (Fig. 3f), and the bar graph shows the

relative information flow of these signaling pathways between WT and *Saa3$^{-/-}$*

macrophages (Fig. 3g). In the context of inflammatory-mediated atherosclerosis

progression, WT macrophages present higher signaling activity in THBS, ANGPTL,

LAMININ, PVR, CD226, OSM, NOTCH, CADAM, and JAM pathways compared to

Saa3$^{-/-}$ macrophages (Figs. 3f-3g). In THBS signaling, *Thbs1* is a critical regulator of

macrophage interleukin-1β (IL-1β) production and activation via CD47, highlighting

the complex interplay between thrombospondins and immune modulation [58]. In

ANGPTL signaling, ANGPTL2 promotes vascular inflammation and accelerates

atherosclerosis progression, with increased cardiac ANGPTL2 production linked to

heart failure and pathological cardiac remodeling by influencing inflammatory

responses and affecting both cardiomyocytes and macrophages[59, 60]. In LAMININ

and JAM signaling, LAMININ and junctional adhesion molecules (JAM) are

involved in monocyte-to-macrophage differentiation and macrophage migration,

playing roles in immune cell dynamics and atherosclerotic disease modulation [61,

62]. PVR significantly contributes to atherosclerosis by facilitating leukocyte

migration across the endothelium, thereby promoting inflammation [63]. When loss of

CD226, a costimulatory molecule, has been observed to protect against

atherosclerosis by preventing foam cell formation and reducing plaque buildup [64].

Oncostatin M (OSM) signaling promotes atherogenesis and plaque vulnerability by

promoting the differentiation of vascular smooth muscle cells into osteoblastic

phenotypes via M1 macrophages, potentially leading to plaque instability [35].

NOTCH signaling plays a significant role in atherosclerosis development, with *Notch-*

1 deletion inducing anti-inflammatory M2 effects through the PI3K-Akt pathway,

contrasting with its increased presence in vulnerable human plaques [65].

In contrast to their WT counterparts, *Saa3^-/-* macrophages exhibit elevated

expression of signaling molecules such as COLLAGEN, GRN, SN, TGFb, and

CLEC. Collagen plays a crucial role in modulating macrophage functions and

improving plaque stability, with an increased presence in M2-polarizing macrophages

within the plaque [66, 67]. GRN exerts anti-atherogenic effects by enhancing nitric

oxide production in endothelial cells through the Akt/eNOS signaling pathway,

forming a complex with apolipoprotein A-I to stabilize atherosclerotic plaques. In

addition, in atherosclerotic mice, its deficiency leads to increased expression of

adhesion molecules and inflammatory cytokines [68-70]. TGF-β signaling contributes

to the reduction and stabilization of atherosclerotic plaques by altering macrophage

functions, and its overexpression is linked to smaller lesions, increased collagen

deposition, smooth muscle cell infiltration, and reduced macrophage presence,

ultimately fostering a more stable lesion phenotype [71]. CLEC signaling, specifically

through Clec4a2, a C-type lectin receptor, protects against atherosclerosis by

promoting homeostatic functions within the vessel wall, enhancing cholesterol

metabolism, and limiting immune responses to toll-like receptors [72]. In summary,

this analysis confirms that aortic macrophages in WT mice are more associated with

inflammatory and migration processes, whereas aortic macrophages in $Saa3^{-/-}$ mice

are more related to vascular protection and aortic plaque stabilization.

Differential effects of *Saa3* deficiency on macrophage subpopulations and ligand-

receptor interactions

We further investigated the differential effects of *Saa3* deficiency on five

macrophage subpopulations. Our observations revealed that, aside from Trem2^{hi} Mφ,

the gene expression profiles of CD74^{hi}MHCII^{hi} Mφ, Folr2^{hi} Mφ, Retnla^{hi}Ear2^{hi} Mφ,

and Apbb2^{hi} Mφ in $Saa3^{-/-}$ mice exhibited substantial downregulation of numerous

genes when compared to their WT counterparts (Figs. 4a-4d). We also performed GO

term enrichment pathway analysis on the downregulated DEGs in these four $Saa3^{-/-}$

macrophage subpopulations compared to WT (Figs. 4e-4h). The analysis revealed

common downregulated genes in $Saa3^{-/-}$ mice, including *Lyz1*, *Lyz2*, other lysozyme-

related genes, *Cxcl13*, and *Ccl8*, which are associated with pathways involved in cell

killing and the destruction of other organisms, suggesting that the absence of *Saa3*

impairs macrophage functions in combating pathogens as previous studies [73, 74]

(Figs. 4e-4h). Specifically, Folr2^{hi} Mφ in *Saa3*$^{-/-}$ mice showed significant

downregulation of *Cxcl13* (Fig. 4b). Previous study indicates patients with carotid

atherosclerosis exhibit elevated plasma levels of CXCL13, especially in symptomatic

cases, and higher CXCL13 expression within atherosclerotic carotid plaques

compared to non-atherosclerotic vessels [75]. CCL8 is a potent chemoattractant for

monocytes and macrophages, directing them to areas of inflammation and infection

[76, 77]. Retnla^{hi}Ear2^{hi} Mφ in WT mice show higher enrichment in ERK1 and ERK2

cascade, positive regulation of angiogenesis, and lipid storage (Figs. 4c, 4g). ERK1

and ERK2 signaling pathways regulate macrophage function in atherosclerosis.

Deletion of *Erk1* reduces macrophage migration and plaque lipid content in LDL

receptor-deficient mice on a high-fat diet [78, 79]. Apbb2^{hi} Mφ in WT mice enrich in

defense response to virus and interferon (Figs. 4d, 4h). This may indicate that Apbb2^{hi}

Mφ in WT mice has more inflammatory response features than in *Saa3*$^{-/-}$ mice,

contributing to chronic inflammation, plaque instability, and progression [80]. To

elucidate the specific ligand-receptor interactions contributing to the observed

differences in pathway patterns between *Saa3*$^{-/-}$ and WT groups, we conducted a

comparative analysis of ligand-receptor signaling within five subpopulations of

macrophages (Supplementary Figs 3a-3e). The analysis encompassed an assessment

of the differential expression of ligands and receptors and the computation of

interaction probabilities, thereby enabling the dissection of signaling discrepancies

between the two genotypic conditions. Since aortic Trem2^{hi} Mφs are more abundant

in *Saa3*$^{-/-}$ mice, we focus on ligand-receptor interactions within Trem2^{hi} Mφ under

both WT and *Saa3*$^{-/-}$ conditions (Supplementary Fig. 3d). Prominently, pairs such as

Spp1-Cd44 and Spp1-integrins (Itgav+Itgb1, Itga5+Itgb1, Itga4+Itgb1), exhibit a

heightened probability of interaction in the signal transduction pathways of Trem2^{hi}

Mφ within the *Saa3$^{-/-}$* mice (Supplementary Fig. 3d). Spp1 is established to bind

integrins Itgav, Itgb1, Itgb5, and Itga4 (Col4a2-Sdc44, Col4a2-Cd44, Col4a2-

(Itga9+Itgb1), Col4a1-Sdc44, Col4a1-Cd44, Col4a1-(Itga9+Itgb1)), influencing

cellular adhesion and migration crucially [81, 82] (Supplementary Fig. 3d).

Furthermore, interactions involving Col4a2 and Col4a1 with their partners are

exclusively observed in the *Saa3$^{-/-}$* condition (Supplementary Fig. 3d). As major

constituents of the basement membrane, type IV collagen may have implications for

the stability of atherosclerotic plaques [83]. In contrast, the CCL-CCR signaling axis,

encompassing Ccl9-Ccr1, Ccl8-Ccr1, Ccl7-Ccr1, Ccl6-Ccr2, Ccl6-Ccr1, Ccl3-Ccr5,

and Ccl3-Ccr1, is exclusively active in the WT condition within Trem2^{hi} Mφ

transitioning to Retnla^{hi}Ear2^{hi} Mφ (Supplementary Fig. 3d). These chemokine-

mediated communications facilitate the recruitment of macrophages or monocytes to

sites of inflammation, contributing to the formation and progression of atherosclerotic

lesions and plaque vulnerability [84, 85]. Parallel observations were noted for the

ligand-receptor pairs Cd74^{hi}MHCII^{hi} Mφ, Folr2^{hi} Mφ, Retnla^{hi}Ear2^{hi} Mφ, and Apbb2^{hi}

Mφ to Retnla^{hi}Ear2^{hi} Mφ (Supplementary Figs. 3a-3e). It is posited that these altered

signaling patterns may underlie the observed reductions in plaque burden and

macrophage recruitments in *Saa3$^{-/-}$* mice.

Trem2^{hi} Mφ potentially mitigates aortic plaque development by enhancing lipid

uptake, promoting an anti-inflammatory environment, and increasing tissue

remodeling.

In atherosclerosis, *Saa3* deletion precipitates substantial changes in macrophage gene

expression profiles, within the Trem2^{hi} Mφ, as evidenced by differential expression

between WT and *Saa3*[-/-] groups (Fig. 5a). The upregulation of genes in *Saa3*[-/-] than

WT Trem2[hi] Mφ such as *Fabp4, Fabp5,* and *Lpl*, which are central to lipid

metabolism [86, 87], as well as *Ctsk*, has been implicated in promoting disturbed

flow-induced endothelial inflammation [88]. Other genes including *Gpnmb,*

Atp6v0d2, Mrc1, Mmp12, and *Spp1* also exhibit altered expression (Fig. 5a). Studies

suggest that *Gpnmb* may be anti-inflammatory by promoting inflammation resolution,

as it can attenuate pro-inflammatory responses and promote M2 polarization in

macrophages [47, 89, 90]. The suppression of *Gpnmb* has been associated with

enhanced atherosclerotic plaque formation, whereas its overexpression correlates with

decreased plaque development [91]. Augmented *Atp6v0d2* expression activates

autophagy as a compensatory response to hemodynamic pressure, which indicates its

role in maintaining plaque stability and potentially curtailing the progression of

atherosclerosis [92, 93]. *Mrc1* appears to modulate macrophage polarization toward

an anti-inflammatory phenotype [94], while *Mmp12* plays a pivotal role in attenuating

inflammation and fostering resolution via targeted proteolysis of chemokines [95, 96].

Spp1+ foamy macrophages have been linked to signaling pathways that might abate

plaque progression and stabilize susceptible lesions [97]. These genes correspond to

GO enrichment pathways such as lipid localization, fatty acid transport, and the

positive regulation of macrophage-derived foam cell differentiation (Fig. 5b). These

results suggest that *Saa3*[-/-] Trem2[hi] Mφ may possess enhanced lipid uptake and

metabolism capabilities, which could contribute to the significant reduction in plaque

size. Additionally, genes like *Ctsk*, *Gpnmb*, *Igf1*, and *Spp1* are associated with tissue

remodeling pathways, indicating a potential shift of *Saa3*[-/-] Trem2[hi] Mφ towards an

anti-inflammatory phenotype (Fig. 5b). Differential gene expression analysis utilizing

violin plots demonstrated that genes associated with tissue repair and lipid

metabolism—specifically *Gpnmb*, *Fabp5*, *Lpl*, *Lgals3*, *Cd36*, *Spp1*, and *Pparg*—are

more prominently expressed in *Saa3*$^{-/-}$ Trem2^{hi} Mφ compared to their WT

counterparts (Fig. 5c). CD36 is a scavenger receptor widely expressed on the surface

of various cell types, including macrophages. CD36 binds to oxLDL particles,

facilitating their uptake into the macrophage, particularly in the context of

atherosclerosis and inflammation [98]. Concomitantly, there is a downregulation of

genes in *Saa3*$^{-/-}$ than WT Trem2^{hi} Mφ such as *Thbs1*, *Kdm7a*, *Vegfa*, *Slc7a11*, *Fn1*,

and *Chil3*. These genes correspond to GO enrichment pathways such as positive

regulation of chemotaxis, positive regulation of vasculature development, and

angiogenesis (Fig. 5b). Angiogenesis is associated with inflammatory cells and can

contribute to plaque instability [99, 100]. This complex gene expression network

underscores the multifaceted influence of *Saa3* in atherosclerosis and pinpoints

interconnected genes geared toward modulating macrophage function to combat

disease progression.

Analysis of aortic Trem2^{hi} macrophages from WT and *Saa3*$^{-/-}$ mice also revealed a

consistent expression pattern for a cohort of genes associated with lipid metabolism

and cholesterol processing (Figs 5b-5c). The violin plots compare the expression

levels of genes central to cholesterol efflux pathways, including *Abca1, Abcg1,*

Nr1h2, Scarb, Nr1h3, and *Apoe* (Fig. 5c, Supplementary Fig. 4). There were no

statistically significant differences observed between the two genotypes for most

genes (Fig. 5c).

Analysis through *CellChat* has additionally revealed alterations in the growth

differentiation factor (GDF) signaling pathway in all subsets of the aortic

macrophages (Figs. 5d-5e). In the *Saa3*$^{-/-}$ Trem2^{hi} macrophage phenotype, there is a

notable shift towards the sender role in the GDF signaling network (Fig. 5d). In

contrast, Folr2^{hi} macrophages predominantly assume the signaling initiative in the

WT condition (Figs. 5d-5e). This shift in signaling behavior suggests that the absence

of *Saa3* leads to a reprogramming of the Trem2^{hi} macrophage population, endowing

them with a more prominent role in initiating signaling within the GDF pathway.

GDF-15 has been reported to reduce cardiac damage and rupture after myocardial

infarction by mitigating inflammation and apoptosis via the PI3K/Akt pathway. It also

protects against atherosclerosis by modulating apoptosis, inhibiting monocyte

recruitment and macrophage activation, and enhancing plaque stability [101-103].

Other studies suggest GDF-15 may reduce macrophage accumulation in *Ldlr*$^{-/-}$ or

Apoe$^{-/-}$ mice [104-106]. This shift in the gene expression landscape highlights the

critical role of *Saa3* in regulating macrophage function and may provide insight into

the molecular mechanisms underpinning the immune responses in aortic plaque

progression.

SAA3 modulates cholesterol efflux, synthesis, and inflammatory responses in

macrophages under steady-state condition or during foamy macrophage

differentiation

SAA3 has been shown to exacerbate atherosclerosis in *ApoE*$^{-/-}$ mice, as evidenced by

increased lesion area upon overexpression and reduced disease severity with

suppression [20]. Another study also observed reduced aortic plaque lesions in

atherosclerotic *Saa3*$^{-/-}$*Ldlr*$^{-/-}$ mice and indicated that these effects are sex-dependent

[14]. To validate our scRNA-seq observations on aortic macrophages and investigate

the role of *Saa3* in foamy macrophages, we performed qRT-PCR to assess the

expression of inflammatory genes (*Il1b* and *Tnf*), cholesterol efflux genes (*Abca1* and

Abcg1), and cholesterol synthesis genes (*Cyp51* and *Hmgcr*) in bone marrow-derived

macrophages (BMDMs) from WT and *Saa3-/-* mice (Fig. 6a). These BMDMs were

treated overnight with either media alone or 20 µg/ml soluble cholesterol to

differentiate cells into foamy macrophages (Fig. 6a). We also conducted scRNA-seq

analysis on WT BMDMs treated with SAA3 in either media alone or 20 µg/ml

soluble cholesterol (Fig. 6a). Overall, gender-specific effects were generally minimal

under these experimental conditions, while genes involved in cholesterol efflux and

synthesis were significantly lower in female WT foamy macrophages treated with

SAA3 compared to males (Figs. 6b-c). Consistent with the previous study [14], *Il1b*

expression was significantly upregulated in BMDMs from female *Saa3-/-* mice

compared to WT under media-alone conditions, with no such effect observed in male

mice, while *Tnf* expression remained unchanged between genotypes (Fig. 6b).

Following soluble cholesterol treatment, both WT and *Saa3-/-* BMDMs present

minimal differences, with upregulated expression of the cholesterol efflux gene

(*Abca1*) and downregulated expression of the cholesterol synthesis gene (*Cyp51*),

indicative of the foamy macrophage phenotype (Fig. 6c). SAA3 treatment

significantly reduced cholesterol efflux gene expression (*Abca1* and *Abcg1*) while

increasing cholesterol synthesis gene expression (*Hmgcr*) in WT BMDMs undergoing

foamy macrophage differentiation (Fig. 6c). We next performed scRNA-seq analysis

on SAA3-treated WT BMDMs to confirm that genes associated with tissue repair and

lipid metabolism (*Gpnmb*, *Lpl*, *Lgals3*, *Spp1*, *Cd36*, *Pparg*) were significantly

downregulated compared to untreated controls (Fig. 6d). This aligns with the *in vivo*

phenotypes, where Trem2hi aortic macrophages from *Saa3-/-* mice exhibited

significantly higher expression of these genes compared to Trem2hi macrophages

from WT mice (Figs. 5c, 6d). We further observed the genes associated with acute

inflammatory response, interferon responses, and cell chemotaxis are increased in

SAA3-treated WT BMDMs (Figs. 6e-6f). Notably, exogenous SAA3 significantly

increases *Il1b* expression in both foamy and non-foamy macrophages (Fig. 6b).

These data indicate that SAA3 modulates inflammatory and lipid metabolism

pathways in macrophages, particularly during foamy macrophage differentiation, by

reducing cholesterol efflux, increasing cholesterol synthesis, and enhancing

inflammatory and interferon responses, suggesting that SAA3 potentially exacerbates

atherosclerosis progression through these mechanisms.

Discussion

Gut microbiota plays significant roles in inducing and maintaining sex dimorphism,

particularly in relation to immune responses, metabolism, and hormone regulation. In

NOD mice, the sex difference in type 1 diabetes susceptibility disappears under germ-

free conditions, while gut microbiota transfer from males to females elevates

testosterone levels and protects against disease [107]. This protection, mediated by

microbiota-regulated sex hormones, is lost upon androgen receptor blockade[107].

Metabolic syndrome is also mediated by the gut microbiota in a sex-dependent

manner, as demonstrated by the transfer of this sex-specific effect through fecal

transplantation between male and female mice, with antibiotic treatment eliminating

these differences and highlighting the causative role of microbiota in sexual

dimorphism [108]. Furthermore, the microbiota regulates bone marrow mesenchymal

stem cells (BMMSCs) by inhibiting osteogenesis and promoting adipogenesis, as

shown by increased osteogenic differentiation in germ-free (GF) mice, while

colonization with specific-pathogen-free (SPF) microbiota restored normal BMMSC

proliferation, differentiation, and immunomodulatory functions [109]. The microbiota

regulates hematopoietic stem cell (HSC) self-renewal and differentiation under stress

by modulating local bone marrow iron availability through red blood cell recycling by

macrophages [110]. Notably, the gut microbiota modulates the expression of SAA3 in

colonic and adipose tissue via Toll-like receptor signaling [15]. These findings

suggest that the gut microbiota may influence confounding factors, such as iron

availability and SAA3 levels, within the tissue microenvironment, thereby affecting

sex dimorphism and immune cell differentiation. The SAA3-induced sex dimorphism

observed in previous studies may also be influenced by microbiota-driven effects on

bone marrow cell differentiation during early life.

Previous study indicates that SAA3 promotes atherosclerosis in male mice while

providing protective effects in female mice [14]. Jylhava et. al. suggest a potential

sex-dimorphism in the relationship between serum amyloid A (SAA) and early

atherosclerosis, as SAA levels correlated with carotid intima-media thickness (IMT)

in men but not in women, indicating differing underlying mechanisms of

atherosclerotic risk between the sexes [111]. The absence of SAA3 in female mice led

to a significant decrease in adipose tissue macrophage content and inflammation,

while similar effects were not observed in male mice [112]. Another study highlights

a significant sex dimorphism in the association between SAA levels and adipocyte

size, with pronounced correlations observed in lean and overweight women, while no

such associations were found in men [113]. Our results indicate there is still less

aortic plaque progression in female atherosclerotic *Saa3$^{-/-}$* than in WT mice (Fig. 1).

The gender differences were minimal but SAA3 significantly reduced cholesterol

efflux genes and increased synthesis genes, particularly in female WT foamy

macrophages (Fig. 6). SAA3-treated WT BMDMs showed downregulation of tissue

repair and lipid metabolism genes and upregulation of acute inflammatory and

interferon-response genes, aligning with *in vivo* phenotypes of Trem2^{hi} macrophages

in *Saa3^{-/-}* mice represented the higher expression of lipid metabolism genes (Fig. 6).

These results suggest that SAA3 modulates inflammatory and lipid pathways in

foamy macrophages, which may be exacerbating atherosclerosis progression *in vivo*.

The single-cell RNA sequencing (scRNA-seq) analysis of CD45+ cells from aortic

arches revealed significant alterations in the aortic immune cell landscape of *Saa3^{-/-}*

mice. Increased proportions of T cells, B cells, monocytes, and γδ T cells were noted,

while macrophages and innate lymphoid cells (ILCs) were reduced compared to WT

mice (Fig. 2). This shift in immune cell composition suggests that *Saa3* deficiency

alters the inflammatory responses within the aortic arches, potentially contributing to

the observed reduction in plaque progression. *CellChat* analysis further highlighted

the impact of *Saa3* deficiency on intercellular communication within the aortic

microenvironment (Figs. 2, 3). *Saa3^{-/-}* mice exhibited enhanced intercellular

interaction strength, particularly among γδ T cells and all other immune cell types,

suggesting a more reactive immune milieu. Conversely, the reduced macrophage-to-

all other immune cell signalings were observed, indicating a dampened inflammatory

response (Fig. 2). These findings suggest that *Saa3* deficiency promotes shifts towards

reparative and less inflammatory phenotypes in aortic immune cells, potentially

influencing atherosclerosis progression.

Compared to our previously published dataset, where we profiled single-cell sorted

Cx3cr1-derived aortic macrophages during atherosclerosis progression and regression

[8], we identified 11 aortic macrophage clusters, while we only identified five distinct

clusters with notable differences in gene expression between WT and *Saa3^{-/-}* mice

(Fig. 3a). This may be due to the difference in the platform used to capture single cells

between 10X Genomics and BD Rhapsody. Nonetheless, we still observed that

Trem2^{hi} macrophages were more abundant in *Saa3^{-/-}* mice (Fig. 3b) and revealed gene

expression profiles indicative of enhanced lipid metabolism and tissue repair

functions (Fig. 5). These macrophages may be crucial in mitigating plaque

progression by promoting an anti-inflammatory environment and enhancing

extracellular matrix production. Our findings also highlight the differential ligand-

receptor interactions in WT and $Saa3^{-/-}$ mice macrophages (Supplementary Fig. 3).

$Saa3$ deficiency led to altered signaling dynamics, particularly in Trem2^{hi}

macrophages, which showed increased interactions involving $Spp1$ and integrins, as

well as type IV collagen, suggesting enhanced adhesion and migration capabilities. In

contrast, chemokine-mediated communications were more prominent in WT

macrophages, facilitating the recruitment of inflammatory cells to atherosclerotic

lesions (Supplementary Fig. 3d). The observed changes in gene expression and

intercellular communication underscore the complex role of $Saa3$ in modulating

Trem2^{hi} macrophage functions during atherosclerosis. $Saa3$ deficiency appears to

shift Trem2^{hi} macrophages towards a phenotype that promotes tissue repair and

plaque stability (Fig. 5), which may explain the reduced plaque progression despite

elevated cholesterol levels.

In conclusion, our study reveals that $Saa3$ deficiency has a dual impact on

atherosclerosis, reducing aortic plaque progression in female mice while increasing

peripheral lipid levels. The enhanced reparative and anti-inflammatory properties of

macrophages in $Saa3^{-/-}$ mice likely contribute to the observed protective effects

against atherosclerosis. $Saa3$ is exclusively expressed in macrophages within the

immune cell population and SAA3 modulates inflammatory and lipid pathways in

foamy macrophages. These findings provide valuable insights into the regulatory

mechanisms of immune function within the aortic niche and highlight the potential of

targeting $Saa3$ for therapeutic strategies in atherosclerosis. Further research is

warranted to elucidate the precise molecular pathways involved and to explore the

therapeutic potential of modulating *Saa3* expression in cardiovascular diseases.

The observed alterations in immune cell composition and intercellular communication

in *Saa3*-deficient mice highlight a shift toward reparative and anti-inflammatory

phenotypes, potentially stabilizing plaques despite elevated cholesterol levels.

Author contributions

TYC, YZY, and JDL designed experiments and interpreted the data. TYC and YZY

wrote the manuscript draft, followed by JDL editing. TYC, YZY, CJL, JCW, CHY,

YTC, and PAC performed experiments. TYC and YZY coordinated and performed

the single-cell RNA-seq experiments and then analyzed single-cell RNA-seq data.

Acknowledgments

We thank the Animal Resource Center and the Consortium of Integrative Biomedical

Science Key Technology at National Taiwan University for their technical support.

We also acknowledged the AAV Core Facility of Academia Sinica in generating

recombinant AAV (Grant AS-CFII112-204) and National Center for High-

performance Computing (NCHC) of National Applied Research Laboratories

(NARLabs) in Taiwan for providing computational and storage resources. Funding:

This work was financially supported by the National Science and Technology

Council, Taiwan (110-2636-B-002-021, 111-2636-B-002-024, 112-2636-B-002-015,

113-2636-B-002-009), National Taiwan University (113L894501, 113-UN0066),

Ministry of Education (Yushan Young Fellow Program, Featured Areas Research

Center Program of "Center for Advanced Computing and Imaging in Biomedicine"

(NTU-113L900701))

Material and Methods

Animals

All animal procedures were approved by the National Taiwan University IACUC Committee. The Saa3-knockout mice ($Saa3^{-/-}$; C57BL/6N-Saa3tm1.1(KOMP)Vlcg/MbpMmucd) were developed by the Mutant Mouse Resource & Research Centers (MMRRC). The ES cell clone 13807A-A1 was injected into morulae or blastocysts to generate chimeric mice. These chimeras were then bred with C57BL/6N mice, resulting in the production of heterozygous tm1 (Deletion) mice. To achieve recombination at the LoxP sites, heterozygous tm1 mice were crossed with a ubiquitous Cre deleter mouse line, generating tm1.1 (CRE-mediated Deletion) allele mice. Notably, the archived germplasm does not carry the Cre transgene. Cryo-recovery of the strain will be conducted using C57BL/6N females by the MMRRC. Detailed information about the Cre-mediated deletion process can be found at the following reference: Velocigene KOMP Repository, clone 13807, http://velocigene.com/komp/detail/13807. The cryopreserved sperm were acquired and recovered by the National Laboratory Animal Center in Taiwan. We generated heterozygous $Saa3^{+/-}$ mice and bred them to obtain 12-week-old Wild-type C57BL/6J mice wild-type (WT) and $Saa3^{-/-}$ littermates used for the experiments. Murine atherosclerosis was induced by injecting intraperitoneally once with AAVmPCSK9 (ssAAV8/hAAT-mPCSK9D377Y, AAV core, Institute of Biomedical Sciences, Academia Sinica, using a plasmid acquired from Addgene (Plasmid #58376) and kindly provided by Dr. Mi-Hua Tao) at 5×10^{11} viral particles/mouse and fed on Western diet (Research Diets, Cat# D12079Bi) for 18 weeks. All mice were housed in a specific pathogen-free (SPF) environment within individually ventilated cages

700   (IVC) at a ventilation rate of 70-100 air changes per hour (ACH) under positive

701   pressure. The facility maintained a constant temperature of $22 \pm 2°C$ and a relative

702   humidity of $55 \pm 10\%$. A 12-hour light-dark cycle was regulated by timers (7:00 AM

703   to 7:00 PM light, 7:00 PM to 7:00 AM dark).

Cholesterol level measurements

Blood samples were collected from mice and total cholesterol levels in plasma were

assessed in WT and *Saa3*[-/-] mice subjected to AAVmPCSK9D377Y and fed on

Western Diet for 0, 4, 8, 12, and 16 weeks (n=5 mice per group). Cholesterol levels

were measured using a colorimetric assay (FUJIFILM/Wako, Cat# 639-50981), with

absorbance read at 600 nm following a 5-minute incubation at 37 °C. HDL and

LDL/VLDL cholesterol levels in plasma were quantified using the ab65390

Cholesterol Assay Kit (Abcam, Version 13a, Updated 17 April 2023), following the

manufacturer's protocol with adaptations for our setup. Plasma (20-30 μL) was mixed

with an equal volume of 2X Precipitation Buffer to separate the HDL and LDL/VLDL

cholesterol fractions. After adding reaction mixes, samples were incubated at 37 °C

for 60 minutes away from light. Absorbance was measured at 570 nm using a

microplate reader, and cholesterol concentrations were determined by correlating

absorbance readings with the standard curve. Statistically significant differences

between the two groups at respective time points were determined using the unpaired

Mann-Whitney U-test ($*p < 0.05$).

Histology

At harvesting, the aortic arch was meticulously excised from the experimental mice

for the subsequent single-cell isolation, and then aortic roots were embedded in OCT

(Catalog # NC0696746). Serial sections were performed at 6 μm on embedded aortic

roots and stained with Oil Red O solution (Sigma) for 10 minutes at 37°C. Intimal

lesions and stained areas were imaged by an inverted fluorescence microscope

(OLYMPUS CKX53) and quantified by ImageJ [17].

Aortic cells flow cytometry sorting and single-cell RNA-seq library preparation

The protocol for isolating aortic cells from mice is as previously described[8]. Briefly,

the procedure began with aorta perfusion using phosphate-buffered saline (PBS).

After perfusion, the aortas were stored in FACS buffer (1X HBSS, 1% BSA, 1mM

EDTA, 20mM HEPES, and 1mM Sodium Pyruvate). The enzymatic digestion

processes involved immersing the aorta in a digestion buffer containing 4U/ml

liberase (Roche, Cat. 54001127001), 100 ug/ml hyaluronidase (Sigma, Cat. H3506),

and 100 ug/ml DNase I (Sigma, Cat. DN25), followed by fine mincing. The mixture

was incubated at 37°C to facilitate effective digestion, with the process repeated 5-6

times to ensure complete tissue breakdown. After digestion, the mixture was filtered

through a 35μm strainer to obtain a single-cell suspension. The collected cells were

then stained with CD45 (Alexa Fluor 488 anti-mouse CD45, Clone: 30-F11,

Biolegend, Cat. 103122) and Live/Dead (L/D) Blue markers (Thermo Fisher, Cat.

L34962), and the CD45-positive, L/D Blue-negative cells were sorted as

Supplementary Figure 1 to prepare single-cell libraries.

Single-cell samples were labeled using the BD™ Mouse Immune Single-Cell

Multiplexing Kit (Cat. 633793) for each cartridge run on the BD Rhapsody Express

Single Cell Analysis System for single-cell capture, followed by cDNA synthesis with

the BD Rhapsody cDNA Kit (Cat. 633773). The mRNA Whole Transcriptome

Analysis (WTA) and Sample Tag libraries were prepared using the BD Rhapsody™

WTA Amplification Kit (Cat. 633801), according to the manufacturer's guidelines.

The quality and quantity of the libraries were assessed using the Agilent 4200

TapeStation System with Agilent High Sensitivity D1000 ScreenTape (Agilent

Technologies, Santa Clara, CA) and a Qubit Fluorometer with the Qubit dsDNA HS

Assay (Invitrogen, MA, USA). Finally, the libraries were sequenced using the

NovaSeq PE150 sequencer (Illumina, San Diego, CA).

Aortic single-cell sequencing data reads alignment and analysis

The fastq files generated from sequencing were processed using Seven Bridges

Genomics. The data were aligned to the mouse_GRCm38-PhiX-gencodevM19-

20181206 and demultiplexed using hashtags to separate samples. Subsequent steps

included filtering low-quality cells and identifying unique molecule identifiers

(UMIs) to produce filtered matrices, which were then utilized as inputs for further

analysis. The filtered matrices were analyzed using the Seurat package (v.4.3.0)[103] in

R (v.4.3.1). Initially, to exclude low-quality cells, we filtered out cells with

nFeature_RNA < 200 and percent.mt > 15, tailoring filter parameters for each sample

based on violin plots of RNA feature numbers and the percentage of mitochondria in

each cell. We used the FindVariableFeatures() function in each sample to select the

top 2000 genes with the highest variance using the variance variance-stabilizing

transformation (VST) method for downstream analysis. Next, we applied a linear

transformation to scale the data, ensuring equal weight to each gene expression level

in downstream analysis. A linear dimensional reduction (PCA) was performed, and

the Elbow plot was used to decide which PCs to include in the following steps. We

cluster cells using the Louvain algorithm (resolution = 0.5). The non-linear

dimensional reduction method was employed for visualizing cell clustering, such as

UMAP.

To analyze two or more single-cell datasets together, we followed the Seurat

integration analysis pipeline. We used the FindIntegrationAnchors() function to

identify anchors, followed by the IntegrateData() function to use these anchors to

integrate the datasets. To automatically annotate cell types, SingleR (v.2.8.0) was

utilized with the ImmGen database, which specifically focuses on immunological cell

types. To ensure the analysis targeted the immune population, contaminating cell

types such as fibroblasts and stromal cells were subsequently excluded. To identify

differentially expressed features, we employed the FindAllarkers() or FindMarkers()

functions to identify biomarkers for each cell cluster (min.pct = 0.25, logfc.threshold

= 0.25). We used visualization tools such as VlnPlot, feature plot, dotplot, and

heatmap to display intergroup gene expression differences. Additionally, to

comprehend the potential biological pathway connections of gene sets expressed by

each cell population and their differences between two genotypes, we conducted a

Gene Ontology (GO) analysis using the ClusterProfiler package (v4.8.3)[104]. This was

done after identifying differentially expressed genes (DEGs) in each cell population

compared to the cell population of another genotype. The results were visualized

using a Dotplot. Moreover, the analysis of intercellular communications was

conducted using the CellChat package (v1.6.1)[105]. This approach allowed us to

understand the roles played by each cell group in signal transduction pathways and

compare the differences in cellular signaling among cell groups under different

genotypes.

Bone marrow-derived macrophages (BMDMs) culture

Bone marrow cells harvested from C57BL/6J mice were cultured in RPMI 1640

medium supplemented with 10% heat-inactivated fetal bovine serum (FBS), 1%

penicillin/streptomycin/amphotericin B (P/S/A), and 20 ng/mL macrophage colony-

stimulating factor (M-CSF). On day 3, the medium was replaced with 10 mL fresh

medium. On day 5, an additional 2 mL of medium was added. After seven days of

culture, adherent macrophages were detached using Accutase and treated with 100

ng/mL SAA3 protein. The cell suspension was adjusted to a density of 5×10^6

cells/mL, seeded into a 48-well plate, and cultured for 24 hours.

Real-time RT-PCR

Total RNA was extracted from BMDMs using TRIzol™ Reagent, and cDNA was

synthesized using SuperScript™ IV Reverse Transcriptase. Real-time RT-PCR was

performed with TaqMan™ Fast Advanced Master Mix on a CFX Connect Real-Time

PCR Detection System. The assays utilized the following TaqMan probes: *Gapdh*

(Mm99999915_g1), *Saa3* (Mm00441203_m1), *Il-1β* (Mm00434228_m1), and *Tnf*

(Mm00443258_m1). Additionally, real-time RT-PCR was also conducted using

iTaq™ Universal SYBR® Green Supermix with primers targeting *Abca1*, *Abcg1*,

Hmgcr, and *Cyp51*. The fold change in each target mRNA was determined by

normalizing its expression level to *Gapdh* within the same sample. Relative gene

expression was calculated using the comparative method ($2^{-\Delta\Delta CT}$).

BMDM single-cell RNA library preparation and sequencing

BMDMs were fixed and permeabilized according to the Evercode™ Fixation v2

protocol (Parse Biosciences). The split-pool barcoding process, following the

Evercode™ WT v2 protocol, consisted of three rounds. In Round 1, cells were

distributed into wells, and sample-specific barcodes were applied to the fixed cells

using an in-cell reverse transcription (RT) reaction. Cells from all wells were then

pooled together. In Round 2, cells were redistributed across a plate, and an in-cell

ligation was performed to append the second barcode. The cells from each well were

again pooled together. In Round 3, cells were split across a plate, and a third barcode

was appended with another in-cell ligation. After pooling cells from each well, the

pooled cells were divided across multiple sublibraries. The cells were then lysed, and

the library was fragmented, end-repaired, and went through A-tailing followed by the

ligation of the sequencing adaptor. The library was then labeled with a fourth

sublibrary-specific barcode. The quality of dsDNA libraries was analyzed using the

High Sensitivity D5000 ScreenTape Kit (Agilent) and concentrations were assessed

with the Qubit dsDNA HS Kit (Thermo Fisher Scientific). Libraries were sequenced

on a NovaSeq X Plus (Illumina) with 10% PhiX.

BMDM single-cell RNA data processing

The sequencing data was demultiplexed and mapped to database (mm10, GRCm38)

with the commercial pipeline (ParseBiosciences-Pipeline.v1.2.1). The sequence reads

from different samples was then distributed based on different Round 1 barcode with

R(v4.4.2). To exclude low-quality cells, we filtered cells with nFeature_RNA > 200,

nFeature_RNA < 2000, and percent.mt < 45, adjusting the filter parameters for each

sample based on violin plots of RNA feature counts and mitochondrial percentages

per cell. The FindVariableFeatures() function was used to select the top 2000 genes

with the highest variance, applying the variance-stabilizing transformation (VST)

method for downstream analysis. The data were then scaled using a linear

transformation to ensure equal weighting of gene expression levels in subsequent

analyses. Principal component analysis (PCA) was performed for linear dimensional

reduction, and the Elbow plot was used to determine the number of principal

components (PCs) to retain. Cell clustering was conducted using the Louvain

algorithm with a resolution of 0.5. For visualization, we applied the non-linear

dimensional reduction method UMAP. For integrating two or more single-cell

datasets, we followed the Seurat integration analysis pipeline. The IntegrateLayers()

function with the CCAIntegration method was used for dataset integration, followed

by the JoinLayers() function to merge the layers. Cell types were annotated

automatically using SingleR (v2.8.0) with the ImmGen database, which specializes in

immunological cell types. Contaminating cell types, such as fibroblasts, were

excluded to ensure the analysis focused on the immune population. To identify

potential biological pathway connections and differences following SAA3 protein

treatment, we performed Gene Ontology (GO) analysis using the ClusterProfiler

package (v4.10.1) on differentially expressed genes (DEGs) from BMDMs under

different conditions. Results were visualized using DotPlot representations.

Data and code availability

Raw sequence data from single-cell RNA sequencing experiments are deposited in the

NCBI Sequence Read Archive under BioProject accession number PRJNA1129034

and gene expression omnibus (GEO) accession number GSE270922. All processing

was performed in R and analysis scripts can be found on Github at

(https://github.com/JDLinLab/Saa3KO-Arch-scRNAseq-pepline-)

Figure legend

Fig. 1 Effects of *Saa3* deletion on peripheral lipid profiles and aortic plaque

progression during atherosclerosis.

(a) Schematic of an experimental model for the time points of plasma collections and

single-cell RNA sequencing (scRNA-seq) on aortic immune cells. C57BL/6 wild-type

(WT) and *Saa3*$^{-/-}$ female mice were injected with AAVmPCSK9 and fed with the

Western diet for 18 weeks. Then aortic arches were digested to single-cell

suspensions and FACS-sorted aortic CD45+ cells were subjected to scRNA-seq on

the BD Rhapsody platform from WT and *Saa3*$^{-/-}$ mice.

(b) Plasma total cholesterol levels in WT and *Saa3*$^{-/-}$ mice subjected with

ssAAV8/hAAT-mPCSK9 and fed with the Western diet for 0, 4, 8, 12, and 16 weeks.

(c) Measurement of high-density lipoprotein (HDL) cholesterol levels and low-

density lipoprotein (LDL) cholesterol levels in plasma from WT and *Saa3*$^{-/-}$ mice over

a period of 0, 2, 4, 8, 12, and 16 weeks.

(d) Representative Oil Red O staining of aortic root sections in WT and *Saa3*$^{-/-}$ mice

at 18 weeks post Western diet. Scale bar: 100 μm.

(e) As in (d), quantification of the lesion area of aortic root sections stained with Oil

Red O.

All data are presented as mean ± SEM; Statistically significant differences between

the two groups at respective time points were denoted by asterisks (*$p < 0.05$),

determined using an unpaired Mann-Whitney U-test. Each symbol represents an

individual mouse; WT mice (n=5), *Saa3*$^{-/-}$ mice (n=5).

Fig. 2 *Saa3* deficiency alters immune intercellular communication in aortic

arches during atherosclerosis progression.

(a) UMAP visualization compares the distribution of cell types in aortic arches

between WT and *Saa3*$^{-/-}$ mice. Unbiased cell type annotations against the IMMGEN

database are colored and labeled accordingly on the plot.

(b) Cell type composition by percentages of WT and *Saa3*$^{-/-}$ mice (blue, WT; red,

Saa3$^{-/-}$) in total sorted CD45+ cells.

(c) Heatmap of top 5 marker genes of each cell type in aortic arches from WT and

Saa3$^{-/-}$ mice. The selection of these top marker genes was based on the results of a

Wilcoxon test for differential expression.

(d) Bar plots show the number (left) and strength (right) of inferred interactions of

intercellular communication networks in aortic arches from WT and *Saa3*$^{-/-}$ mice.

(e) Heatmap visualization of intercellular communication networks in aortic arches

from WT and *Saa3*$^{-/-}$ mice. The red and blue colors represent increased or

decreased signaling in the aortic arches of *Saa3*$^{-/-}$ mice compared to the WT mice. The

top-colored bar plot represents the sum of the columns of values (incoming signaling),

and the right-colored bar plot represents the sum of a row of values (outgoing

signaling).

(f) Volcano plot illustrating differential gene expression in total macrophages within

aortic arches of *Saa3*$^{-/-}$ versus WT mice. The x-axis displays the log$_2$ fold change,

while the y-axis shows the -Log$_{10}$ *p*-value. Grey dots denote genes without significant

differences, green dots indicate genes with significant fold changes, and red dots mark

genes significant in both fold change and *p*-value. The dashed vertical lines represent

the fold change boundaries, and the dashed horizontal line demonstrates the p-value

significance threshold.

(g) Top 10 biological processes enriched in total macrophages within the aortic arches

of WT mice compared to *Saa3*[-/-] mice (upper panel) or *Saa3*[-/-] mice compared to wild-

type mice (lower panel). Enrichment analysis utilized GO pathways to identify

significant biological processes. The gene ratio is plotted along the x-axis, and the

enriched biological processes are listed on the y-axis. A color gradient has been

applied to indicate the -$\log_{10}$ adjusted *p*-value (p.adjust) of each term, with the size of

each point reflecting the number of genes associated with the respective pathway.

Fig. 3 Delineation of macrophage sub-cluster in aortic arches between WT and

***Saa3*[-/-] mice during atherosclerosis progression.**

(a) UMAP plots show the sub-clusters of macrophages (Mφ) in aortic arches between

WT (left) and *Saa3*[-/-] (right) mice. Macrophages are delineated based on their gene

expression profiles: CD74[hi]MHCII[hi] Mφ in navy blue, Folr2[hi] Mφ in red,

Retnla[hi]Ear2[hi] Mφ in green, Trem2[hi] Mφ in sky blue, and Apbb2[hi] Mφ in pink.

(b) The bar graph displays the percentage composition of various macrophage

subpopulations within total macrophages from WT and *Saa3*[-/-] mice. Different colors

represent distinct macrophage subpopulations as indicated in **(a)**.

(c) Dot plot representing the expression profiles of five sub-macrophage populations.

The x-axis lists genes with differential expression, while the y-axis indicates the sub-

macrophage clusters.

(d) Dot plots show the expression profiles of macrophage markers involved in

pathways of chemokine, IFN (interferon), inflammatory, and type II responses across

sub-macrophage clusters. For **(c)** and **(d)**, the size of each dot corresponds to the

percentage of the indicated cell cluster expressing a particular gene marker, and the

color intensity signifies the mean expression level within that cell cluster.

(e) Bar plots show the number (left) and strength (right) of inferred intercellular

communication networks in total aortic macrophages from WT and *Saa3*$^{-/-}$ mice.

(f) Heatmap visualization of overall signaling patterns in aortic macrophages of WT

and *Saa3*$^{-/-}$ mice. The top-colored bar plot represents the sum of each column of

values (signaling the strength of overall pathways) across sub-clusters. The gradient

blue colored bar represents the relative strength of the indicated pathway.

(g) The bar graph displays the relative information flow of various signaling

molecules in aortic macrophages of WT and *Saa3*$^{-/-}$ mice. Blue bars indicate higher

relative information flow in WT macrophages, while red bars indicate higher

information flow in *Saa3*$^{-/-}$ macrophages.

Fig. 4 *Saa3* deficient impacts gene expression profiles of aortic macrophages

(a)-(d) Volcano plot illustrating differential gene expression in four sub-macrophage

populations (CD74^{hi} MHCII^{hi} , Folr2^{hi}, Retnla^{hi} Ear2^{hi}, and Apbb2^{hi} macrophage(Mφ))

within aortic arches of WT versus *Saa3*$^{-/-}$ mice. The x-axis displays the log$_2$ fold

change, while the y-axis shows the -Log$_{10}$ p-value. Grey dots denote genes without

significant differences, green dots indicate genes with significant fold changes, and

red dots mark genes significant in both fold change and *p*-value. The dashed vertical

lines represent the fold change boundaries, and the dashed horizontal line

demonstrates the *p*-value significance threshold.

(e)-(h) The bubble plot presents the GO enrichment analysis for CD74^{hi} MHCII^{hi} ,

Folr2^{hi}, Retnla^{hi} Ear2^{hi}, and Apbb2^{hi} aortic macrophage (Mφ) from WT and *Saa3*$^{-/-}$

mice. The x-axis represents the GeneRatio. The y-axis lists the enriched GO terms.

Bubble size corresponds to the count of DEGs associated with each GO term. The

color gradient from blue to red represents the -$\log_{10}$ (adjusted *p*-value) of the

enrichment.

Fig. 5 Aortic Trem2^{hi} macrophages in *Saa3$^{-/-}$* mice express more lipid metabolism

and tissue repair-associated genes.

(a) Volcano plot illustrating differential gene expression in aortic Trem2^{hi}

macrophage (Mφ) population of *Saa3$^{-/-}$* versus WT mice.

(b) The bubble plot presents the GO enrichment analysis for Trem2^{hi} Mφ from WT

versus *Saa3$^{-/-}$* (upper panel) or *Saa3$^{-/-}$* versus WT (lower panel) mice.

(c) Violin plots depicting the expression levels of tissue repair-associated genes in

Trem2^{hi} Mφ from WT and *Saa3$^{-/-}$* mice aortas. Each plot indicates the gene expression

level for *Gpnmb*, *Fabp5*, *Lpl*, *Lgals3*, *Spp1*, *Cd36*, and *Pparg*, with blue representing

WT and red for *Saa3$^{-/-}$* group. The central white bar denotes the median expression

level, while the thick bar within the violins indicates the interquartile range. The

width of the violins reflects the density of data points at different expression levels.

(d) The circular diagram illustrates the GDF signaling pathway networks in aortic

macrophage subpopulations from WT (left) and *Saa3$^{-/-}$* (right) mice. Each segment of

the circle represents a different macrophage subpopulation. The connections between

segments indicate interactions and signaling pathways shared among these

subpopulations.

(e) Heatmap representation of the inferred GDF signaling pathway network across

five sub-populations of aortic macrophages. Each column represents a specific

macrophage sub-population, while each row corresponds to the role of the population

in the signaling pathway as a sender, receiver, mediator, or influencer.

Fig. 6 SAA3 modulates cholesterol metabolism, inflammatory gene expression,

and immune response pathways in macrophages under steady-state condition or

during foamy macrophage differentiation.

(a) Schematic overview of the experimental design. Bone marrow cells were isolated

from WT and $Saa3^{-/-}$ mice. Then cultured and differentiated into bone marrow-

derived macrophages (BMDMs), followed by treatment with soluble cholesterol to

induce foamy macrophages or recombinant SAA3 protein. Downstream analyses

included real-time PCR and single-cell RNA sequencing (scRNA-seq) using Parse

Evercode™.

(b-c) As treatment conditions in **(a)**, Real-time PCR analysis of genes involved in

inflammatory cytokines (*Il1b*, *Tnf*) and cholesterol metabolism (*Abca1*, *Abcg1*,

Cyp51, *Hmgcr*) and under various treatment conditions. Data are represented as

relative gene expression normalized to the control group, with significant differences

marked by asterisks (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$). The symbols represent the

following comparisons: φ indicates SAA3-treated male WT BMDMs versus male WT

BMDMs cultured in media alone, § indicates SAA3-treated female WT BMDMs

versus female WT BMDMs cultured in media alone, # indicates SAA3-treated male

WT BMDMs in the presence of soluble cholesterol versus male WT BMDMs in

soluble cholesterol media alone, and † indicates SAA3-treated female WT BMDMs in

the presence of soluble cholesterol versus female WT BMDMs in soluble cholesterol

media alone. The number of symbols denotes the level of statistical significance,

where one symbol indicates $p < 0.05$, two symbols indicate $p < 0.01$, and three

symbols indicate $p < 0.001$.

(d-f) scRNA-seq analysis of WT BMDMs treated with media alone or SAA3 protein.

(d-e) Dot plots show average expression (color intensity) and the percentage of cells

expressing key genes (dot size) related to tissue repair, lipid metabolism, and

inflammatory pathways.

(f) Gene Ontology (GO) enrichment analysis of pathways upregulated in SAA3-

treated BMDMs compared to controls. The size of the dots represents the gene ratio,

and color intensity represents statistical significance ($-\log_{10}$ (adjusted p-value)).

Statistical significance was determined using ANOVA or t-tests where appropriate.

1028

1029 **Supplementary Figure Legends**

1030

Supplementary Fig. 1 Sorting strategy of aortic CD45+ cells.

Flow cytometry analysis was performed for cell sorting to isolate CD45+ immune

cells from aortic arches stained for CD45 Alexa 488 and Live/Dead Blue. The top left

plot shows forward scatter (FSC-A) versus side scatter (SSC-A), differentiating cells

based on size and granularity, with a population gated as P1. The top right plot

presents FSC-A versus FSC-H, with the discrimination gate P2 to exclude cell

aggregates. The bottom left plot displays SSC-A against the Live/Dead Blue, with live

cells gated as P7. The bottom right plot demonstrates SSC-A against the fluorescence

intensity of CD45 Alexa-488, with the CD45+ population gated as P8. The CD45+

cells were sorted from P8 gated population.

1041

1042

1043 **Supplementary Fig. 2 Heatmap of top 10 gene markers and GO enrichment**

1044 **analysis of five aortic macrophage sub-cluster.**

(a) Heatmap of top 10 marker genes of each sub-macrophage population in aortic arches from WT and *Saa3*-/- mice. The selection of these top marker genes was based on the results of a Wilcoxon test for differential expression.

(b) Gene Ontology (GO) enrichment analysis of biological pathways in 5 sub-populations of aortic macrophage (Mφ), using differentially expressed genes (DEGs) identified when comparing one of specific Mφ sub-population against the rest of all other Mφ sub-populations. It can represent the functional characteristics of each Mφ sub-population.

Supplementary Fig. 3 Comparison of ligand-receptor pair differences across five sub-macrophage populations between WT and *Saa3*-/- mice.

The bubble plot illustrating the protein-protein interaction probabilities between various ligand-receptor pairs across **(a)** CD74hi MHCIIhi macrophage (Mφ), **(b)** Folr2hi Mφ, **(c)** Retnlahi Ear2hi Mφ, **(d)** Trem2hi Mφ, and **(e)** Apbb2hi Mφ between WT and *Saa3*-/- mice. The x-axis categorizes the conditions under which the interactions were assessed. The y-axis lists the ligand-receptor pairs. Each bubble's color corresponds to the communication probability, with a gradient from minimum (blue) to maximum (red). The communication probability can be seen as a numerical value representing how strong or likely this interaction is between specific cell types. The size of each bubble represents the *p*-value of the interaction.

Supplementary Fig. 4 Comparison of differences in the expression level of cholesterol efflux and lipid metabolism genes from Trem2hi macrophages between WT and *Saa3*-/- mice.

Differential gene expression in aortic Trem2hi macrophages (Mφ) from WT and *Saa3*-/- mice. Violin plots illustrate the distribution of expression levels for genes involved in cholesterol efflux and lipid metabolism in Trem2hi macrophages. Blue represents WT macrophages, while red indicates *Saa3*-/- macrophages.

References

References

1. Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, et
al. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart
Association. Circulation. 2019;139(10):e56-e528. doi:
10.1161/CIR.0000000000000659. PubMed PMID: 30700139.

2. Malekmohammad K, Bezsonov EE, Rafieian-Kopaei M. Role of Lipid
Accumulation and Inflammation in Atherosclerosis: Focus on Molecular and Cellular
Mechanisms. Front Cardiovasc Med. 2021;8:707529. Epub 20210906. doi:
10.3389/fcvm.2021.707529. PubMed PMID: 34552965; PubMed Central PMCID:
PMCPMC8450356.

3. Libby P, Buring JE, Badimon L, Hansson GK, Deanfield J, Bittencourt MS, et al.
Atherosclerosis. Nat Rev Dis Primers. 2019;5(1):56. Epub 20190816. doi:
10.1038/s41572-019-0106-z. PubMed PMID: 31420554.

4. Steinl DC, Kaufmann BA. Ultrasound imaging for risk assessment in
atherosclerosis. Int J Mol Sci. 2015;16(5):9749-69. Epub 20150429. doi:
10.3390/ijms16059749. PubMed PMID: 25938969; PubMed Central PMCID:
PMCPMC4463615.

5. Sekalska B. [Aortic expression of monocyte chemotactic protein-1 (MCP-1) gene
in rabbits with experimental atherosclerosis]. Ann Acad Med Stetin. 2003;49:79-90.
PubMed PMID: 15552841.

6. Zhang Y, Dong D, Xu X, He H, Zhu Y, Lei T, et al. Oxidized high-density
lipoprotein promotes CD36 palmitoylation and increases lipid uptake in macrophages.
J Biol Chem. 2022;298(6):102000. Epub 20220429. doi: 10.1016/j.jbc.2022.102000.
PubMed PMID: 35500650; PubMed Central PMCID: PMCPMC9144050.

7. Javadifar A, Rastgoo S, Banach M, Jamialahmadi T, Johnston TP, Sahebkar A.
Foam Cells as Therapeutic Targets in Atherosclerosis with a Focus on the Regulatory
Roles of Non-Coding RNAs. Int J Mol Sci. 2021;22(5). Epub 20210303. doi:
10.3390/ijms22052529. PubMed PMID: 33802600; PubMed Central PMCID:
PMCPMC7961492.

8. Lin JD, Nishi H, Poles J, Niu X, McCauley C, Rahman K, et al. Single-cell
analysis of fate-mapped macrophages reveals heterogeneity, including stem-like
properties, during atherosclerosis progression and regression. JCI Insight. 2019;4(4).
Epub 20190221. doi: 10.1172/jci.insight.124574. PubMed PMID: 30830865; PubMed
Central PMCID: PMCPMC6478411.

9.   Zhang K, Wang Y, Chen S, Mao J, Jin Y, Ye H, et al. TREM2(hi) resident
macrophages protect the septic heart by maintaining cardiomyocyte homeostasis. Nat
Metab. 2023;5(1):129-46. Epub 20230112. doi: 10.1038/s42255-022-00715-5.
PubMed PMID: 36635449; PubMed Central PMCID: PMCPMC9886554.

10.   Jung SH, Hwang BH, Shin S, Park EH, Park SH, Kim CW, et al. Spatiotemporal
dynamics of macrophage heterogeneity and a potential function of Trem2(hi)
macrophages in infarcted hearts. Nat Commun. 2022;13(1):4580. Epub 20220806. doi:
10.1038/s41467-022-32284-2. PubMed PMID: 35933399; PubMed Central PMCID:
PMCPMC9357004.

11.   den Hartigh LJ, May KS, Zhang XS, Chait A, Blaser MJ. Serum amyloid A and
metabolic disease: evidence for a critical role in chronic inflammatory conditions. Front
Cardiovasc Med. 2023;10:1197432. Epub 20230615. doi: 10.3389/fcvm.2023.1197432.
PubMed PMID: 37396595; PubMed Central PMCID: PMCPMC10311072.

12.   Reigstad CS, Lunden GO, Felin J, Backhed F. Regulation of serum amyloid A3
(SAA3) in mouse colonic epithelium and adipose tissue by the intestinal microbiota.
PLoS One. 2009;4(6):e5842. Epub 20090609. doi: 10.1371/journal.pone.0005842.
PubMed PMID: 19513118; PubMed Central PMCID: PMCPMC2688757.

13.   Smit V, de Mol J, Kleijn M, Depuydt MAC, de Winther MPJ, Bot I, et al. Sexual
dimorphism in atherosclerotic plaques of aged Ldlr(-/-) mice. Immun Ageing.
2024;21(1):27. Epub 20240502. doi: 10.1186/s12979-024-00434-3. PubMed PMID:
38698438; PubMed Central PMCID: PMCPMC11064395.

14.   Chait A, Wang S, Goodspeed L, Gomes D, Turk KE, Wietecha T, et al. Sexually
Dimorphic Relationships Among Saa3 (Serum Amyloid A3), Inflammation, and
Cholesterol Metabolism Modulate Atherosclerosis in Mice. Arterioscler Thromb Vasc
Biol. 2021;41(6):e299-e313. Epub 20210325. doi: 10.1161/ATVBAHA.121.316066.
PubMed PMID: 33761762; PubMed Central PMCID: PMCPMC8159856.

15.   Reigstad CS, Backhed F. Microbial regulation of SAA3 expression in mouse colon
and adipose tissue. Gut Microbes. 2010;1(1):55-7. doi: 10.4161/gmic.1.1.10514.
PubMed PMID: 21327118; PubMed Central PMCID: PMCPMC3035134.

16.   Dong Z, Wu T, Qin W, An C, Wang Z, Zhang M, et al. Serum amyloid A directly
accelerates the progression of atherosclerosis in apolipoprotein E-deficient mice. Mol
Med. 2011;17(11-12):1357-64. Epub 20110921. doi: 10.2119/molmed.2011.00186.
PubMed PMID: 21953420; PubMed Central PMCID: PMCPMC3321823.

17.   Meek RL, Eriksen N, Benditt EP. Murine serum amyloid A3 is a high density
apolipoprotein and is secreted by macrophages. Proc Natl Acad Sci U S A.

1992;89(17):7949-52. doi: 10.1073/pnas.89.17.7949. PubMed PMID: 1518819;
PubMed Central PMCID: PMCPMC49832.

18. Sanada Y, Yamamoto T, Satake R, Yamashita A, Kanai S, Kato N, et al. Serum
Amyloid A3 Gene Expression in Adipocytes is an Indicator of the Interaction with
Macrophages. Sci Rep. 2016;6:38697. Epub 20161208. doi: 10.1038/srep38697.
PubMed PMID: 27929048; PubMed Central PMCID: PMCPMC5144138.

19. Hadrup N, Zhernovkov V, Jacobsen NR, Voss C, Strunz M, Ansari M, et al. Acute
Phase Response as a Biological Mechanism-of-Action of (Nano)particle-Induced
Cardiovascular Disease. Small. 2020;16(21):e1907476. Epub 20200329. doi:
10.1002/smll.201907476. PubMed PMID: 32227434.

20. Thompson JC, Wilson PG, Shridas P, Ji A, de Beer M, de Beer FC, et al. Serum
amyloid A3 is pro-atherogenic. Atherosclerosis. 2018;268:32-5. Epub 20171117. doi:
10.1016/j.atherosclerosis.2017.11.011. PubMed PMID: 29175652; PubMed Central
PMCID: PMCPMC5839639.

21. Keeter WC, Carter NM, Nadler JL, Galkina EV. The AAV-PCSK9 murine model
of atherosclerosis and metabolic dysfunction. Eur Heart J Open. 2022;2(3):oeac028.
Epub 20220420. doi: 10.1093/ehjopen/oeac028. PubMed PMID: 35919346; PubMed
Central PMCID: PMCPMC9242032.

22. Peled M, Nishi H, Weinstock A, Barrett TJ, Zhou F, Quezada A, et al. A wild-type
mouse-based model for the regression of inflammation in atherosclerosis. PLoS One.
2017;12(3):e0173975. Epub 20170314. doi: 10.1371/journal.pone.0173975. PubMed
PMID: 28291840; PubMed Central PMCID: PMCPMC5349694.

23. Han CY, Kang I, Omer M, Wang S, Wietecha T, Wight TN, et al. Serum amyloid
A-containing HDL binds adipocyte-derived versican and macrophage-derived biglycan,
reducing its antiinflammatory properties. JCI Insight. 2020;5(20). Epub 20200924. doi:
10.1172/jci.insight.142635. PubMed PMID: 32970631; PubMed Central PMCID:
PMCPMC7605543.

24. Holmes N. CD45: all is not yet crystal clear. Immunology. 2006;117(2):145-55.
doi: 10.1111/j.1365-2567.2005.02265.x. PubMed PMID: 16423050; PubMed Central
PMCID: PMCPMC1782222.

25. Haudek SB, Xia Y, Huebener P, Lee JM, Carlson S, Crawford JR, et al. Bone
marrow-derived fibroblast precursors mediate ischemic cardiomyopathy in mice. Proc
Natl Acad Sci U S A. 2006;103(48):18284-9. Epub 20061117. doi:
10.1073/pnas.0608799103. PubMed PMID: 17114286; PubMed Central PMCID:
PMCPMC1643845.

26. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019;177(7):1888-902 e21. Epub
20190606. doi: 10.1016/j.cell.2019.05.031. PubMed PMID: 31178118; PubMed
Central PMCID: PMCPMC6687398.

27. Aran D, Looney AP, Liu L, Wu E, Fong V, Hsu A, et al. Reference-based analysis
of lung single-cell sequencing reveals a transitional profibrotic macrophage. Nat
Immunol. 2019;20(2):163-72. Epub 20190114. doi: 10.1038/s41590-018-0276-y.
PubMed PMID: 30643263; PubMed Central PMCID: PMCPMC6340744.

28. Xu B, Chen L, Zhan Y, Marquez KNS, Zhuo L, Qi S, et al. The Biological
Functions and Regulatory Mechanisms of Fatty Acid Binding Protein 5 in Various
Diseases. Front Cell Dev Biol. 2022;10:857919. Epub 20220404. doi:
10.3389/fcell.2022.857919. PubMed PMID: 35445019; PubMed Central PMCID:
PMCPMC9013884.

29. Nickl B, Qadri F, Bader M. Anti-inflammatory role of Gpnmb in adipose tissue of
mice. Sci Rep. 2021;11(1):19614. Epub 20211004. doi: 10.1038/s41598-021-99090-6.
PubMed PMID: 34608215; PubMed Central PMCID: PMCPMC8490452.

30. Ding C, Xiao T, Deng Y, Yang H, Xu B, Li J, et al. The Teleost CXCL13-CXCR5
Axis Induces Inflammatory Cytokine Expression through the Akt-NF-kappaB, p38-AP-
1, and p38-NF-kappaB Pathways. J Immunol. 2024;212(2):317-34. doi:
10.4049/jimmunol.2300237. PubMed PMID: 38054894.

31. Han JH, Suh CH, Jung JY, Nam JY, Kwon JE, Yim H, et al. Association of
CXCL10 and CXCL13 levels with disease activity and cutaneous manifestation in
active adult-onset Still's disease. Arthritis Res Ther. 2015;17(1):260. Epub 20150919.
doi: 10.1186/s13075-015-0773-4. PubMed PMID: 26385705; PubMed Central PMCID:
PMCPMC4575437.

32. Kiss MG, Papac-Milicevic N, Porsch F, Tsiantoulas D, Hendrikx T, Takaoka M, et
al. Cell-autonomous regulation of complement C3 by factor H limits macrophage
efferocytosis and exacerbates atherosclerosis. Immunity. 2023;56(8):1809-24 e10.
Epub 20230726. doi: 10.1016/j.immuni.2023.06.026. PubMed PMID: 37499656;
PubMed Central PMCID: PMCPMC10529786.

33. Suzuki M, Becker L, Pritchard DK, Gharib SA, Wijsman EM, Bammler TK, et al.
Cholesterol accumulation regulates expression of macrophage proteins implicated in
proteolysis and complement activation. Arterioscler Thromb Vasc Biol.
2012;32(12):2910-8. Epub 20121004. doi: 10.1161/ATVBAHA.112.300383. PubMed
PMID: 23042816; PubMed Central PMCID: PMCPMC3501207.

34. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal

enrichment tool for interpreting omics data. Innovation (Camb). 2021;2(3):100141.

Epub 20210701. doi: 10.1016/j.xinn.2021.100141. PubMed PMID: 34557778;

PubMed Central PMCID: PMCPMC8454663.

35. Hou P, Fang J, Liu Z, Shi Y, Agostini M, Bernassola F, et al. Macrophage

polarization and metabolism in atherosclerosis. Cell Death Dis. 2023;14(10):691. Epub

20231020. doi: 10.1038/s41419-023-06206-z. PubMed PMID: 37863894; PubMed

Central PMCID: PMCPMC10589261.

36. Cochain C, Vafadarnejad E, Arampatzi P, Pelisek J, Winkels H, Ley K, et al.

Single-Cell RNA-Seq Reveals the Transcriptional Landscape and Heterogeneity of

Aortic Macrophages in Murine Atherosclerosis. Circ Res. 2018;122(12):1661-74. Epub

20180315. doi: 10.1161/CIRCRESAHA.117.312509. PubMed PMID: 29545365.

37. Nalio Ramos R, Missolo-Koussou Y, Gerber-Ferder Y, Bromley CP, Bugatti M,

Nunez NG, et al. Tissue-resident FOLR2(+) macrophages associate with CD8(+) T cell

infiltration in human breast cancer. Cell. 2022;185(7):1189-207 e25. Epub 20220323.

doi: 10.1016/j.cell.2022.02.021. PubMed PMID: 35325594.

38. Ensan S, Li A, Besla R, Degousee N, Cosme J, Roufaiel M, et al. Self-renewing

resident arterial macrophages arise from embryonic CX3CR1(+) precursors and

circulating monocytes immediately after birth. Nat Immunol. 2016;17(2):159-68. Epub

20151207. doi: 10.1038/ni.3343. PubMed PMID: 26642357.

39. Beckers CML, Simpson KR, Griffin KJ, Brown JM, Cheah LT, Smith KA, et al.

Cre/lox Studies Identify Resident Macrophages as the Major Source of Circulating

Coagulation Factor XIII-A. Arterioscler Thromb Vasc Biol. 2017;37(8):1494-502.

Epub 20170608. doi: 10.1161/ATVBAHA.117.309271. PubMed PMID: 28596376;

PubMed Central PMCID: PMCPMC5526434.

40. Sanin DE, Ge Y, Marinkovic E, Kabat AM, Castoldi A, Caputa G, et al. A common

framework of monocyte-derived macrophage activation. Sci Immunol.

2022;7(70):eabl7482. Epub 20220415. doi: 10.1126/sciimmunol.abl7482. PubMed

PMID: 35427180.

41. Loke P, Lin JD. Redefining inflammatory macrophage phenotypes across stages

and tissues by single-cell transcriptomics. Sci Immunol. 2022;7(70):eabo4652. Epub

20220415. doi: 10.1126/sciimmunol.abo4652. PubMed PMID: 35427177.

42. Cochain C, Vafadarnejad E, Arampatzi P, Jaroslav P, Winkels H, Ley K, et al.

Single-Cell RNA-Seq Reveals the Transcriptional Landscape and Heterogeneity of

Aortic Macrophages in Murine Atherosclerosis. Circ Res. 2018;122:1661-74. Epub

2018/03/17. doi: CIRCRESAHA.117.312509 [pii]

10.1161/CIRCRESAHA.117.312509. PubMed PMID: 29545365.

43.   Muller A, Beck K, Rancic Z, Muller C, Fischer CR, Betzel T, et al. Imaging
atherosclerotic plaque inflammation via folate receptor targeting using a novel 18F-
folate radiotracer. Mol Imaging. 2014;13:1-11. PubMed PMID: 24622812.

44.   Kim K, Shim D, Lee JS, Zaitsev K, Williams JW, Kim KW, et al. Transcriptome
Analysis Reveals Nonfoamy Rather Than Foamy Plaque Macrophages Are
Proinflammatory in Atherosclerotic Murine Models. Circ Res. 2018;123(10):1127-42.
doi: 10.1161/CIRCRESAHA.118.312804. PubMed PMID: 30359200; PubMed Central
PMCID: PMCPMC6945121.

45.   Qi J, Sun H, Zhang Y, Wang Z, Xun Z, Li Z, et al. Single-cell and spatial analysis
reveal interaction of FAP(+) fibroblasts and SPP1(+) macrophages in colorectal cancer.
Nat Commun. 2022;13(1):1742. Epub 20220401. doi: 10.1038/s41467-022-29366-6.
PubMed PMID: 35365629; PubMed Central PMCID: PMCPMC8976074.

46.   Hoeft K, Schaefer GJL, Kim H, Schumacher D, Bleckwehl T, Long Q, et al.
Platelet-instructed SPP1(+) macrophages drive myofibroblast activation in fibrosis in a
CXCL4-dependent manner. Cell Rep. 2023;42(2):112131. Epub 20230218. doi:
10.1016/j.celrep.2023.112131. PubMed PMID: 36807143; PubMed Central PMCID:
PMCPMC9992450.

47.   Saade M, Araujo de Souza G, Scavone C, Kinoshita PF. The Role of GPNMB in
Inflammation.   Front   Immunol.   2021;12:674739.   Epub   20210512.   doi:
10.3389/fimmu.2021.674739. PubMed PMID: 34054862; PubMed Central PMCID:
PMCPMC8149902.

48.   Kumari A, Kristensen KK, Ploug M, Winther AL. The Importance of Lipoprotein
Lipase Regulation in Atherosclerosis. Biomedicines. 2021;9(7). Epub 20210706. doi:
10.3390/biomedicines9070782. PubMed PMID: 34356847; PubMed Central PMCID:
PMCPMC8301479.

49.   Wang X, Khalil RA. Matrix Metalloproteinases, Vascular Remodeling, and
Vascular   Disease.   Adv   Pharmacol.   2018;81:241-330.   Epub   20170919.   doi:
10.1016/bs.apha.2017.08.002. PubMed PMID: 29310800; PubMed Central PMCID:
PMCPMC5765875.

50.   Zou Y, Hu M, Lee J, Nambiar SM, Garcia V, Bao Q, et al. Nrf2 is essential for
timely M phase entry of replicating hepatocytes during liver regeneration. Am J Physiol
Gastrointest   Liver   Physiol.   2015;308(4):G262-8.   Epub   20141218.   doi:

10.1152/ajpgi.00332.2014. PubMed PMID: 25524062; PubMed Central PMCID:
PMCPMC4329475.

51.   Buechler C, Ritter M, Orso E, Langmann T, Klucken J, Schmitz G. Regulation of
scavenger receptor CD163 expression in human monocytes and macrophages by pro-
and antiinflammatory stimuli. J Leukoc Biol. 2000;67(1):97-103. PubMed PMID:
10648003.

52.   Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time
for reassessment. F1000Prime Rep. 2014;6:13. Epub 20140303. doi: 10.12703/P6-13.
PubMed PMID: 24669294; PubMed Central PMCID: PMCPMC3944738.

53.   Gudgeon J, Marin-Rubio JL, Trost M. The role of macrophage scavenger receptor
1 (MSR1) in inflammatory disorders and cancer. Front Immunol. 2022;13:1012002.
Epub 20221017. doi: 10.3389/fimmu.2022.1012002. PubMed PMID: 36325338;
PubMed Central PMCID: PMCPMC9618966.

54.   Zhao SJ, Kong FQ, Jie J, Li Q, Liu H, Xu AD, et al. Macrophage MSR1 promotes
BMSC osteogenic differentiation and M2-like polarization by activating
PI3K/AKT/GSK3beta/beta-catenin pathway. Theranostics. 2020;10(1):17-35. Epub
20200101. doi: 10.7150/thno.36930. PubMed PMID: 31903103; PubMed Central
PMCID: PMCPMC6929615.

55. Kurilenko N, Fatkhullina AR, Mazitova A, Koltsova EK. Act Locally, Act
Globally-Microbiota, Barriers, and Cytokines in Atherosclerosis. Cells. 2021;10(2).
Epub 20210207. doi: 10.3390/cells10020348. PubMed PMID: 33562334; PubMed
Central PMCID: PMCPMC7915371.

56. Chen HJ, Tas SW, de Winther MPJ. Type-I interferons in atherosclerosis. J Exp
Med. 2020;217(1). doi: 10.1084/jem.20190459. PubMed PMID: 31821440; PubMed
Central PMCID: PMCPMC7037237.

57. Edsfeldt A, Swart M, Singh P, Dib L, Sun J, Cole JE, et al. Interferon regulatory
factor-5-dependent CD11c+ macrophages contribute to the formation of rupture-prone
atherosclerotic plaques. Eur Heart J. 2022;43(19):1864-77. Epub 20220209. doi:
10.1093/eurheartj/ehab920. PubMed PMID: 35567557; PubMed Central PMCID:
PMCPMC9113304.

58. Stein EV, Miller TW, Ivins-O'Keefe K, Kaur S, Roberts DD. Secreted
Thrombospondin-1 Regulates Macrophage Interleukin-1beta Production and
Activation through CD47. Sci Rep. 2016;6:19684. Epub 20160127. doi:
10.1038/srep19684. PubMed PMID: 26813769; PubMed Central PMCID:
PMCPMC4728557.

59. Tian Z, Miyata K, Kadomatsu T, Horiguchi H, Fukushima H, Tohyama S, et al.
ANGPTL2 activity in cardiac pathologies accelerates heart failure by perturbing
cardiac function and energy metabolism. Nat Commun. 2016;7:13016. Epub 20160928.
doi: 10.1038/ncomms13016. PubMed PMID: 27677409; PubMed Central PMCID:
PMCPMC5052800.

60. Horio E, Kadomatsu T, Miyata K, Arai Y, Hosokawa K, Doi Y, et al. Role of
endothelial cell-derived angptl2 in vascular inflammation leading to endothelial
dysfunction and atherosclerosis progression. Arterioscler Thromb Vasc Biol.
2014;34(4):790-800. Epub 20140213. doi: 10.1161/ATVBAHA.113.303116. PubMed
PMID: 24526691.

61. Li L, Song J, Chuquisana O, Hannocks MJ, Loismann S, Vogl T, et al. Endothelial
Basement Membrane Laminins as an Environmental Cue in Monocyte Differentiation
to Macrophages. Front Immunol. 2020;11:584229. Epub 20201030. doi:
10.3389/fimmu.2020.584229. PubMed PMID: 33193400; PubMed Central PMCID:
PMCPMC7662115.

62. Wang J, Liu H. The Roles of Junctional Adhesion Molecules (JAMs) in Cell
Migration. Front Cell Dev Biol. 2022;10:843671. Epub 20220309. doi:
10.3389/fcell.2022.843671. PubMed PMID: 35356274; PubMed Central PMCID:
PMCPMC8959349.

63. Mishra BH, Mishra PP, Raitoharju E, Marttila S, Mononen N, Sievanen H, et al.
Modular genome-wide gene expression architecture shared by early traits of
osteoporosis and atherosclerosis in the Young Finns Study. Sci Rep. 2021;11(1):7111.
Epub 20210329. doi: 10.1038/s41598-021-86536-0. PubMed PMID: 33782480;
PubMed Central PMCID: PMCPMC8007808.

64. Zhang Y, Xu X, Ma J, Liu Y, Duan C, Liu Y, et al. Loss of CD226 protects
apolipoprotein E-deficient mice from diet-induced atherosclerosis. Biochim Biophys
Acta Mol Basis Dis. 2022;1868(9):166452. Epub 20220523. doi:
10.1016/j.bbadis.2022.166452. PubMed PMID: 35618182.

65. Zhang M, Yue X, Zhao X, Lu Y, Liu H, Zhang Z, et al. Macrophage-specific
deletion of Notch-1 induced M2 anti-inflammatory effect in atherosclerosis via
activation of the PI3K-oxidative stress axis. Aging (Albany NY). 2023;15(24):15196-
212. Epub 20231226. doi: 10.18632/aging.205342. PubMed PMID: 38149979;
PubMed Central PMCID: PMCPMC10781475.

66. Barrett TJ. Macrophages in Atherosclerosis Regression. Arterioscler Thromb Vasc
Biol. 2020;40(1):20-33. Epub 20191114. doi: 10.1161/ATVBAHA.119.312802.

PubMed PMID: 31722535; PubMed Central PMCID: PMCPMC6946104.

67. Scott TE, Lewis CV, Zhu M, Wang C, Samuel CS, Drummond GR, et al. IL-4 and
IL-13 induce equivalent expression of traditional M2 markers and modulation of
reactive oxygen species in human macrophages. Sci Rep. 2023;13(1):19589. Epub
20231110. doi: 10.1038/s41598-023-46237-2. PubMed PMID: 37949903; PubMed
Central PMCID: PMCPMC10638413.

68. Huang G, An L, Fan M, Zhang M, Chen B, Zhu M, et al. Potential role of full-
length and nonfull-length progranulin in affecting aortic valve calcification. J Mol Cell
Cardiol. 2020;141:93-104. Epub 20200402. doi: 10.1016/j.yjmcc.2020.03.012.
PubMed PMID: 32247641.

69. Saeedi-Boroujeni A, Purrahman D, Shojaeian A, Poniatowski LA, Rafiee F,
Mahmoudian-Sani MR. Progranulin (PGRN) as a regulator of inflammation and a
critical factor in the immunopathogenesis of cardiovascular diseases. J Inflamm (Lond).
2023;20(1):1. Epub 20230119. doi: 10.1186/s12950-023-00327-0. PubMed PMID:
36658641; PubMed Central PMCID: PMCPMC9851114.

70. Nguyen AD, Nguyen TA, Singh RK, Eberle D, Zhang J, Abate JP, et al.
Progranulin in the hematopoietic compartment protects mice from atherosclerosis.
Atherosclerosis. 2018;277:145-54. Epub 20180830. doi:
10.1016/j.atherosclerosis.2018.08.042. PubMed PMID: 30212683; PubMed Central
PMCID: PMCPMC6432779.

71. Reifenberg K, Cheng F, Orning C, Crain J, Kupper I, Wiese E, et al.
Overexpression of TGF-ss1 in macrophages reduces and stabilizes atherosclerotic
plaques in ApoE-deficient mice. PLoS One. 2012;7(7):e40990. Epub 20120719. doi:
10.1371/journal.pone.0040990. PubMed PMID: 22829904; PubMed Central PMCID:
PMCPMC3400574.

72. Park I, Goddard ME, Cole JE, Zanin N, Lyytikainen LP, Lehtimaki T, et al. C-type
lectin receptor CLEC4A2 promotes tissue adaptation of macrophages and protects
against atherosclerosis. Nat Commun. 2022;13(1):215. Epub 20220111. doi:
10.1038/s41467-021-27862-9. PubMed PMID: 35017526; PubMed Central PMCID:
PMCPMC8752790.

73. Chauvin C, Radulovic K, Boulard O, Delacre M, Waldschmitt N, Regnier P, et al.
Loss of NOD2 in macrophages improves colitis and tumorigenesis in a lysozyme-
dependent manner. Front Immunol. 2023;14:1252979. Epub 20231009. doi:
10.3389/fimmu.2023.1252979. PubMed PMID: 37876927; PubMed Central PMCID:
PMCPMC10590911.

74. Alam MS, Getz M, Safeukui I, Yi S, Tamez P, Shin J, et al. Genomic expression
analyses reveal lysosomal, innate immunity proteins, as disease correlates in murine
models of a lysosomal storage disorder. PLoS One. 2012;7(10):e48273. Epub
20121019. doi: 10.1371/journal.pone.0048273. PubMed PMID: 23094108; PubMed
Central PMCID: PMCPMC3477142.

75. Smedbakken LM, Halvorsen B, Daissormont I, Ranheim T, Michelsen AE,
Skjelland M, et al. Increased levels of the homeostatic chemokine CXCL13 in human
atherosclerosis - Potential role in plaque stabilization. Atherosclerosis.
2012;224(1):266-73. Epub 20120720. doi: 10.1016/j.atherosclerosis.2012.06.071.
PubMed PMID: 22840692.

76. Dangi A, Husain I, Jordan CZ, Yu S, Natesh N, Shen X, et al. Blocking CCL8-
CCR8-Mediated Early Allograft Inflammation Improves Kidney Transplant Function.
J Am Soc Nephrol. 2022;33(10):1876-90. Epub 20220816. doi:
10.1681/ASN.2022020139. PubMed PMID: 35973731; PubMed Central PMCID:
PMCPMC9528333.

77. Luo H, Li L, Han S, Liu T. The role of monocyte/macrophage chemokines in
pathogenesis of osteoarthritis: A review. Int J Immunogenet. 2024;51(3):130-42. Epub
20240310. doi: 10.1111/iji.12664. PubMed PMID: 38462560.

78. Ricard N, Zhang J, Zhuang ZW, Simons M. Isoform-Specific Roles of ERK1 and
ERK2 in Arteriogenesis. Cells. 2019;9(1). Epub 20191221. doi: 10.3390/cells9010038.
PubMed PMID: 31877781; PubMed Central PMCID: PMCPMC7017123.

79. Al-Ahmadi W, Webberley TS, Joseph A, Harris F, Chan YH, Alotibi R, et al. Pro-
atherogenic actions of signal transducer and activator of transcription 1 serine 727
phosphorylation in LDL receptor deficient mice via modulation of plaque inflammation.
FASEB J. 2021;35(10):e21892. doi: 10.1096/fj.202100571RR. PubMed PMID:
34569651; PubMed Central PMCID: PMCPMC9549671.

80. Jin M, Fang J, Wang JJ, Shao X, Xu SW, Liu PQ, et al. Regulation of toll-like
receptor (TLR) signaling pathways in atherosclerosis: from mechanisms to targeted
therapeutics. Acta Pharmacol Sin. 2023;44(12):2358-75. Epub 20230807. doi:
10.1038/s41401-023-01123-5. PubMed PMID: 37550526; PubMed Central PMCID:
PMCPMC10692204.

81. Frank JW, Seo H, Burghardt RC, Bayless KJ, Johnson GA. ITGAV (alpha v
integrins) bind SPP1 (osteopontin) to support trophoblast cell adhesion. Reproduction.
2017;153(5):695-706. Epub 20170301. doi: 10.1530/REP-17-0043. PubMed PMID:
28250242.

82. Erikson DW, Burghardt RC, Bayless KJ, Johnson GA. Secreted phosphoprotein 1 (SPP1, osteopontin) binds to integrin alpha v beta 6 on porcine trophectoderm cells and integrin alpha v beta 3 on uterine luminal epithelial cells, and promotes trophectoderm cell adhesion and migration. Biol Reprod. 2009;81(5):814-25. Epub 20090701. doi: 10.1095/biolreprod.109.078600. PubMed PMID: 19571258.

83. Vallejo J, Duner P, To F, Engelbertsen D, Goncalves I, Nilsson J, et al. Activation of immune responses against the basement membrane component collagen type IV does not affect the development of atherosclerosis in ApoE-deficient mice. Sci Rep. 2019;9(1):5964. Epub 20190412. doi: 10.1038/s41598-019-42375-8. PubMed PMID: 30979943; PubMed Central PMCID: PMCPMC6461614.

84. Li M, Wang M, Wen Y, Zhang H, Zhao GN, Gao Q. Signaling pathways in macrophages: molecular mechanisms and therapeutic targets. MedComm (2020). 2023;4(5):e349. Epub 20230911. doi: 10.1002/mco2.349. PubMed PMID: 37706196; PubMed Central PMCID: PMCPMC10495745.

85. Ruytinx P, Proost P, Van Damme J, Struyf S. Chemokine-Induced Macrophage Polarization in Inflammatory Conditions. Front Immunol. 2018;9:1930. Epub 20180907. doi: 10.3389/fimmu.2018.01930. PubMed PMID: 30245686; PubMed Central PMCID: PMCPMC6137099.

86. Furuhashi M, Saitoh S, Shimamoto K, Miura T. Fatty Acid-Binding Protein 4 (FABP4): Pathophysiological Insights and Potent Clinical Biomarker of Metabolic and Cardiovascular Diseases. Clin Med Insights Cardiol. 2014;8(Suppl 3):23-33. Epub 20150202. doi: 10.4137/CMC.S17067. PubMed PMID: 25674026; PubMed Central PMCID: PMCPMC4315049.

87. Wieczorek E, Cwiklinska A, Kuchta A, Kortas-Stempak B, Gliwinska A, Jankowski M. The Differential Effects of HDL Subpopulations on Lipoprotein Lipase (LPL)-Mediated VLDL Catabolism. Biomedicines. 2021;9(12). Epub 20211205. doi: 10.3390/biomedicines9121839. PubMed PMID: 34944655; PubMed Central PMCID: PMCPMC8698418.

88. Fang F, Feng T, Li J, Zhang H, Wang Q, Chen Y, et al. Cathepsin K contributed to disturbed flow-induced atherosclerosis is dependent on integrin-actin cytoskeleton-NF-kappaB pathway. Genes Dis. 2023;10(2):583-95. Epub 20220425. doi: 10.1016/j.gendis.2022.03.020. PubMed PMID: 37223522; PubMed Central PMCID: PMCPMC10201601.

89. Zhang H, Zhang S, Dang X, Lin L, Ren L, Song R. GPNMB plays an active role in the M1/M2 balance. Tissue Cell. 2022;74:101683. Epub 20211117. doi:

10.1016/j.tice.2021.101683. PubMed PMID: 34800878.

90. Wang J, Zhang X, Long M, Yuan M, Yin J, Luo W, et al. Macrophage-derived
GPNMB trapped by fibrotic extracellular matrix promotes pulmonary fibrosis.
Commun Biol. 2023;6(1):136. Epub 20230202. doi: 10.1038/s42003-022-04333-5.
PubMed PMID: 36732560; PubMed Central PMCID: PMCPMC9893197.

91. Nickl B, Qadri F, Bader M. Role of Gpnmb in atherosclerosis of female mice.
Biochem Biophys Res Commun. 2022;621:20-4. Epub 20220702. doi:
10.1016/j.bbrc.2022.06.082. PubMed PMID: 35809343.

92. Chen P, Chen Y, Wu W, Chen L, Yang X, Zhang S. Identification and validation of
four hub genes involved in the plaque deterioration of atherosclerosis. Aging (Albany
NY). 2019;11(16):6469-89. Epub 20190826. doi: 10.18632/aging.102200. PubMed
PMID: 31449494; PubMed Central PMCID: PMCPMC6738408.

93. Huang K, Chen S, Yu LJ, Wu ZM, Chen QJ, Wang XQ, et al. Serum secreted
phosphoprotein 1 level is associated with plaque vulnerability in patients with coronary
artery disease. Front Immunol. 2024;15:1285813. Epub 20240215. doi:
10.3389/fimmu.2024.1285813. PubMed PMID: 38426091; PubMed Central PMCID:
PMCPMC10902157.

94. Orecchioni M, Ghosheh Y, Pramod AB, Ley K. Macrophage Polarization:
Different Gene Signatures in M1(LPS+) vs. Classically and M2(LPS-) vs. Alternatively
Activated Macrophages. Front Immunol. 2019;10:1084. Epub 20190524. doi:
10.3389/fimmu.2019.01084. PubMed PMID: 31178859; PubMed Central PMCID:
PMCPMC6543837.

95. Collison J. MMP12 makes the cut. Nat Rev Rheumatol. 2018;14(9):501. doi:
10.1038/s41584-018-0056-y. PubMed PMID: 30022107.

96. Dean RA, Cox JH, Bellac CL, Doucet A, Starr AE, Overall CM. Macrophage-
specific metalloelastase (MMP-12) truncates and inactivates ELR+ CXC chemokines
and generates CCL2, -7, -8, and -13 antagonists: potential role of the macrophage in
terminating polymorphonuclear leukocyte influx. Blood. 2008;112(8):3455-64. Epub
20080725. doi: 10.1182/blood-2007-12-129080. PubMed PMID: 18660381.

97. Willemsen L, de Winther MP. Macrophage subsets in atherosclerosis as defined
by single-cell technologies. J Pathol. 2020;250(5):705-14. Epub 20200311. doi:
10.1002/path.5392. PubMed PMID: 32003464; PubMed Central PMCID:
PMCPMC7217201.

98. Chen Y, Zhang J, Cui W, Silverstein RL. CD36, a signaling receptor and fatty acid

transporter that regulates immune cell metabolism and fate. J Exp Med. 2022;219(6). Epub 20220419. doi: 10.1084/jem.20211314. PubMed PMID: 35438721; PubMed Central PMCID: PMCPMC9022290.

99. Herrmann J, Lerman LO, Mukhopadhyay D, Napoli C, Lerman A. Angiogenesis in atherogenesis. Arterioscler Thromb Vasc Biol. 2006;26(9):1948-57. Epub 20060622. doi: 10.1161/01.ATV.0000233387.90257.9b. PubMed PMID: 16794218.

100. Khurana R, Simons M, Martin JF, Zachary IC. Role of angiogenesis in cardiovascular disease: a critical appraisal. Circulation. 2005;112(12):1813-24. doi: 10.1161/CIRCULATIONAHA.105.535294. PubMed PMID: 16172288.

101. Asrih M, Wei S, Nguyen TT, Yi HS, Ryu D, Gariani K. Overview of growth differentiation factor 15 in metabolic syndrome. J Cell Mol Med. 2023;27(9):1157-67. Epub 20230329. doi: 10.1111/jcmm.17725. PubMed PMID: 36992609; PubMed Central PMCID: PMCPMC10148061.

102. Adela R, Banerjee SK. GDF-15 as a Target and Biomarker for Diabetes and Cardiovascular Diseases: A Translational Prospective. J Diabetes Res. 2015;2015:490842. Epub 20150727. doi: 10.1155/2015/490842. PubMed PMID: 26273671; PubMed Central PMCID: PMCPMC4530250.

103. Kempf T, Eden M, Strelau J, Naguib M, Willenbockel C, Tongers J, et al. The transforming growth factor-beta superfamily member growth-differentiation factor-15 protects the heart from ischemia/reperfusion injury. Circ Res. 2006;98(3):351-60. Epub 20060105. doi: 10.1161/01.RES.0000202805.73038.48. PubMed PMID: 16397141.

104. Preusch MR, Baeuerle M, Albrecht C, Blessing E, Bischof M, Katus HA, et al. GDF-15 protects from macrophage accumulation in a mousemodel of advanced atherosclerosis. Eur J Med Res. 2013;18(1):19. Epub 20130624. doi: 10.1186/2047-783X-18-19. PubMed PMID: 23800095; PubMed Central PMCID: PMCPMC3701574.

105. Kempf T, Zarbock A, Widera C, Butz S, Stadtmann A, Rossaint J, et al. GDF-15 is an inhibitor of leukocyte integrin activation required for survival after myocardial infarction in mice. Nat Med. 2011;17(5):581-8. Epub 20110424. doi: 10.1038/nm.2354. PubMed PMID: 21516086.

106. Johnen H, Kuffner T, Brown DA, Wu BJ, Stocker R, Breit SN. Increased expression of the TGF-b superfamily cytokine MIC-1/GDF15 protects ApoE(-/-) mice from the development of atherosclerosis. Cardiovasc Pathol. 2012;21(6):499-505. Epub 20120302. doi: 10.1016/j.carpath.2012.02.003. PubMed PMID: 22386250.

107. Markle JG, Frank DN, Mortin-Toth S, Robertson CE, Feazel LM, Rolle-

Kampczyk U, et al. Sex differences in the gut microbiome drive hormone-dependent
regulation of autoimmunity. Science. 2013;339(6123):1084-8. Epub 20130117. doi:
10.1126/science.1233521. PubMed PMID: 23328391.

108. Kaliannan K, Robertson RC, Murphy K, Stanton C, Kang C, Wang B, et al.
Estrogen-mediated gut microbiome alterations influence sexual dimorphism in
metabolic syndrome in mice. Microbiome. 2018;6(1):205. Epub 20181113. doi:
10.1186/s40168-018-0587-0. PubMed PMID: 30424806; PubMed Central PMCID:
PMCPMC6234624.

109. Xiao E, He L, Wu Q, Li J, He Y, Zhao L, et al. Microbiota regulates bone marrow
mesenchymal stem cell lineage differentiation and immunomodulation. Stem Cell Res
Ther. 2017;8(1):213. Epub 20170929. doi: 10.1186/s13287-017-0670-7. PubMed
PMID: 28962644; PubMed Central PMCID: PMCPMC5622543.

110. Zhang D, Gao X, Li H, Borger DK, Wei Q, Yang E, et al. The microbiota regulates
hematopoietic stem cell fate decisions by controlling iron availability in bone marrow.
Cell Stem Cell. 2022;29(2):232-47 e7. Epub 20220121. doi:
10.1016/j.stem.2021.12.009. PubMed PMID: 35065706; PubMed Central PMCID:
PMCPMC8818037.

111. Jylhava J, Haarala A, Eklund C, Pertovaara M, Kahonen M, Hutri-Kahonen N, et
al. Serum amyloid A is independently associated with metabolic risk factors but not
with early atherosclerosis: the Cardiovascular Risk in Young Finns Study. J Intern Med.
2009;266(3):286-95. doi: 10.1111/j.1365-2796.2009.02120.x. PubMed PMID:
19702793.

112. den Hartigh LJ, Wang S, Goodspeed L, Ding Y, Averill M, Subramanian S, et al.
Deletion of serum amyloid A3 improves high fat high sucrose diet-induced adipose
tissue inflammation and hyperlipidemia in female mice. PLoS One. 2014;9(9):e108564.
Epub 20140924. doi: 10.1371/journal.pone.0108564. PubMed PMID: 25251243;
PubMed Central PMCID: PMCPMC4177399.

113. Sjoholm K, Lundgren M, Olsson M, Eriksson JW. Association of serum amyloid
A levels with adipocyte size and serum levels of adipokines: differences between men
and women. Cytokine. 2009;48(3):260-6. Epub 20090915. doi:
10.1016/j.cyto.2009.08.005. PubMed PMID: 19758820.

1557

1558

1559

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Figure 1

Supplementary Figure 2

a

b

Supplementary Figure 3

a

Supplementary Figure 3

Supplementary Figure 3

C

RetnlahiEar2hi Mφ

Supplementary Figure 3

d

Supplementary Figure 3

Supplementary Figure 4