The RRM domain-containing protein Rbp3 interacts with ribosomes and the 3′ ends of mRNAs encoding photosynthesis proteins

Luisa Hemm[1], Elisabeth Lichtenberg[2], Stefan Tholen[3], Viktoria Reimann[1], Kenta

Kakazu[4], Sotaro Machida[4], Moontaha Mahbub[5], Oliver Schilling[3], Annegret Wilde[2]

Satoru Watanabe[4], Conrad W. Mullineaux[5] and Wolfgang R. Hess[1,*]

[1]Genetics and Experimental Bioinformatics, Faculty of Biology, Institute of Biology III,

University of Freiburg, Freiburg, Germany;

[2]Molecular Genetics of Prokaryotes, Faculty of Biology, Institute of Biology III,

University of Freiburg, Freiburg, Germany;

[3]Institute for Surgical Pathology, Medical Center – University of Freiburg, Faculty of

Medicine, Freiburg, Germany;

[4]Department of Bioscience, Tokyo University of Agriculture, 1-1-1 Sakuragaoka,

Setagaya-ku, Tokyo, Japan

[5]School of Biological and Behavioural Sciences, Queen Mary University of London,

London, United Kingdom

*corresponding author: Wolfgang R. Hess

Email: wolfgang.hess@biologie.uni-freiburg.de

Author Contributions: WRH designed the project and secured funding. ST and OS

carried out MS-based proteomic analyses. SW, SM and KK provided the Δ*rbp2/3*

mutant and the BLI data. VR performed the microarray analyses, MM and CWM

provided the spectroscopic analyses. EL and AW performed the RNA-FISH and

microscopical analyses and contributed to the data interpretation. All other

experiments and analyses were performed by LH. LH and WRH wrote the manuscript

with input from all authors.

Competing Interest Statement: The authors declare that they have no competing

interests.

Classification: Biological Sciences, Microbiology

Keywords cyanobacteria, gene expression, mRNA transport, photosynthesis, RNA binding proteins

This PDF file includes:

Main Text

Figures 1 to 6

Abstract

RNA recognition motif (RRM) domain proteins are crucial RNA-binding proteins (RBPs) across all domains of life. In cyanobacteria, single RRM domain proteins are involved in mRNA targeting to the thylakoid membrane and acclimation to certain stress conditions, but many details of their physiological functions and molecular targets have remained unknown. The model cyanobacterium *Synechocystis* sp. PCC 6803 has a family of three genes encoding the RRM domain-containing proteins Rbp1, Rbp2 and Rbp3. Here, we verified the RNA-binding activity of Rbp3 *in vivo* and show that cells of a Δ*rbp3* deletion strain had a lower PSI:PSII ratio and decreased pigment content and were significantly smaller than wild-type cells. To identify the set of interacting molecules, co-immunoprecipitation experiments were performed with a strain expressing a C-terminally FLAG-tagged Rbp3. Mass spectrometry of the elution fraction suggested physical proximity between Rbp3, ribosomes, and a very small number of other proteins. The most highly enriched transcript in the co-eluting RNA fraction was the *psaAB* mRNA. This was corroborated by fluorescent *in situ* hybridization (FISH) analyses showing decreased *psaA* mRNA signals in Δ*rbp3*, and colocalization with Rbp3-GFP in the wild type. Other mRNAs co-enriched with Rbp3 encode thylakoid, plasma membrane and carboxysome proteins. Binding assays using Bio-layer Interferometry validated the Rbp3-*psaAB* mRNA interaction, indicating a preference for folded RNA segments near or overlapping the respective stop codons.

Significance statement

The mechanisms by which proteins are produced at specific sites and inserted into the intricate membrane systems of photosynthetic cyanobacteria are only partially understood. While RRM domain proteins are well-studied RNA-binding proteins in eukaryotes, their functions in bacteria remain underexplored. This study reveals that the RRM domain protein Rbp3 in the cyanobacterium *Synechocystis* sp. PCC 6803 binds mRNAs encoding photosynthetic proteins, plasma membrane proteins and carboxysome proteins and localizes near ribosomes. The bound RNA segments are typically near the ends of coding regions, or in 5′ untranslated regions. These findings suggest that Rbp3 is involved in targeting mRNAs to various intracellular locations by interacting with structural elements within these mRNA molecules.

Introduction

RNA-binding proteins in bacteria

RNA-binding proteins (RBPs) play crucial roles in the physiological functions, processing and stability of RNA molecules. Although prokaryotes lack the extensive processing of pre-mRNAs that is typical of eukaryotes, they have a variety of RBPs that act in different steps of RNA-related processes. Among the best-studied RBPs in bacteria are the Rho factors, which terminate transcription (1), Hfq and ProQ, which mediate sRNA/mRNA interactions and can alter RNA stability (2, 3), and cold shock protein A, an important transcription antitermination factor (4). Little is known about putative homologs of these well-known RBPs in cyanobacteria. A structural homolog of Hfq exists in most cyanobacteria including the model *Synechocystis* sp. PCC 6803 (hereafter *Synechocystis* 6803), but the residues interacting with RNA in other bacteria are not conserved (5), and RNA-binding activity was not detected (6). However, systematic studies revealed several candidate RBPs in cyanobacteria (7–9). Some of these RBPs have recently been characterized, such as YlxR/RnpM interacting with RNase P (10), or PatR, a member of the larger OB-fold protein family binding several sRNAs and impacting the pattern of cell differentiation pattern in multicellular *Nostoc* sp. PCC 7120 (8), but only a few cyanobacterial RBPs have been analyzed in more detail thus far.

RRM domain-containing proteins in cyanobacteria

The RNA recognition motif (RRM; also called ribonucleoprotein domain (RNP) or RNA-binding domain (RBD)) (11, 12) is the most common RNA-binding domain. It consists of a four-stranded antiparallel beta-sheet, stacked on two alpha helices (13). The domain was first described in proteins of heterogeneous nuclear ribonucleoprotein particles, where it is involved in the splicing of heterogeneous nuclear RNA (11). In addition to splicing, eukaryotic RRM proteins are now known to be involved in a variety of RNA-related processes. Eukaryotic RRM proteins are associated with mRNA stability (14), poly(A)-binding (15), mRNA transport (16), translation (17) and degradation (18). In prokaryotes, most proteins harboring an RRM domain are found in species in the phylum cyanobacteria (19). These bacterial RRM domain proteins typically have a single RRM domain, whereas eukaryotic proteins usually have multiple RRM domains.

RRM-domain proteins in cyanobacteria were discovered in *Synechococcus* sp.
PCC 6301 (20), and homologs were soon found in many other cyanobacteria (21–24).
Genes encoding these RBPs are strikingly conserved, with at least three members
throughout the entire cyanobacterial phylum **(*SI Appendix*, Fig. S1)**, suggesting a
function very relevant to the physiology of cyanobacteria. Distinct types of these RBPs
have been recognized (25, 26), proteins with a C-terminal glycine-rich domain,
frequently becoming expressed under cold stress (22, 27, 28), short proteins that
consist of only the RRM domain and the longer Type II proteins with a relatively long
C-terminal region of unknown function.

The presence of RRM domains in these proteins has prompted multiple
attempts to verify their RNA binding experimentally. Sugita and Sugiura presented
evidence of Rbp1 and Rbp2 from *Synechococcus elongatus* PCC 6301 to bind RNA
in vitro (20). Utilizing a gel-filtration assay, RNA-binding activity was shown for the
RBPs of *Anabaena variabilis* (27). Mutsuda et al. presented evidence of Rbp1 and
Rbp2 binding to RNA *in vivo* based on native gel electrophoresis of nuclease-treated
and -untreated cell extracts in *Synechococcus elongatus* PCC 7942 (29). Using an
RNA-binding assay with homopolymeric RNA molecules, RNA-binding activity was
demonstrated for RbpA1 of *Nostoc* sp. PCC 7120 *in vitro* (25). Mahbub et al. performed
RT-PCR on RNA co-immunoprecipitated with Rbp2 and Rbp3 from *Synechocystis*
6803 (30), and Zhang et al. applied RNA-co-immunoprecipitation and high-throughput
sequencing (RIP-Seq) to recombinant Rbp1 and Rbp2 from the same strain (26).
Finally, evidence of *Synechococcus elongatus* PCC 7942 Rbp2 binding to RNA was
obtained by incubating recombinant Rbp2 and poly(rU), followed by gel filtration (31).
Based on these results gained for several different RBPs in various cyanobacteria, it
is now timely to determine the population of interacting transcripts *in vivo*.

Three RRM domain-containing proteins are present in *Synechocystis* 6803,
Rbp1, Rbp2, and Rbp3 (encoded by the genes *sll0517, ssr1480* and *slr0193*,
respectively; **SI Appendix, Fig. S1**). Rbp1 is important under cold stress (32), although
Rbp2 and Rbp3 are also somewhat cold-inducible (33). Rbp3 is involved in the
acclimation to salt stress and the cold-induced lipid peroxidation (34). Indeed, the
levels of some mRNAs encoding proteins involved in fatty acid biosynthesis and salt
stress differed between a Δ*rbp3* mutant and the wild type, although the molecular basis
of these differences remained unclear (34). Rbp3 consists of the RRM domain in the
N-terminal 85 amino acids and an uncharacterized C-terminal section of 66 residues.

It could be shown, that the salt tolerance in Δ*rbp3* mutants was restored when the
Rbp3 C-terminus was fused to the N-terminus of Rbp1 suggesting that the salt-related
aspects were not connected to the RRM domain (26). Conversely, the glycine-rich C-
terminus of Rbp1 was dispensable under cold stress conditions (26).

Using fluorescent *in situ* hybridization (FISH) to probe the subcellular location of
mRNAs encoding core subunits of the photosystems, we observed that Rbp2 and
Rbp3 are involved in the targeting of *psaA* and *psbA* mRNAs towards the thylakoid
membrane (30). Here, we analyzed the function of Rbp3 at the molecular and
physiological level. We demonstrate actual RNA binding *in vivo* and show that this
protein is relevant for the accumulation of a subset of mRNAs and determined the
interacting proteins and transcripts. The data suggest that Rbp3 binds to the 3′ regions
of mRNAs encoding photosynthesis core subunits and interacts closely with
ribosomes. Our results further suggest specialized functions for the different members
of this protein family, but point also at some functional complementarity. Finally, we
present a working model for the function of Rbp3.

159
160

Results

Rbp3 mutant strains show differences in growth and pigment content compared
to wild type

To phenotype the Δ*rbp3* deletion mutant, spot assays were performed in triplicate with
stepwise 10-fold dilutions (**Fig. 1A, B**). Growth of the *Synechocystis* 6803 wild type on
agar medium was compared to that of Δ*rbp3* and a complementation strain in which a
plasmid pVZ322-localized *rbp3* copy was introduced into Δ*rbp3* and expressed from
its native promoter. Growth of the Δ*rbp3* strain appeared slightly impaired compared
to the other two strains. However, closer inspection revealed similar numbers of
colonies in the three strains, while the colony sizes in Δ*rbp3* were significantly
decreased compared to the wild type and complementation strain. After 8 days of
growth under standard conditions, wild-type colonies had a median colony size of
~0.6 mm, while that of the Δ*rbp3* mutant was ~0.36 mm. This effect was rescued in the
complementation strain Δ*rbp3*+Rbp3 under low light (**Fig. 1A**), and partially rescued
under high light conditions (**Fig. 1B**). We conclude that the size, but not the number of

colonies, was different in the absence of Rbp3 indicating that lack of Rbp3 slows
growth, but does not induce cell death.

Whole-cell absorption spectra normalized to OD_{750} and 77K fluorescence emission
spectra with chlorophyll excitation at 435 nm normalized to the PSI peak, or
phycocyanin excitation at 600 nm indicated a lower cellular content of chlorophyll *a*
and phycocyanin in Δ*rbp3* compared to the GT-I wild type (**Fig. 1C–E**). Moreover,
these spectra indicated that the Δ*rbp3* cells had a lower PSI/PSII ratio. Introduction of
an untagged Rbp3 restored spectra almost identical with the wild type (**Fig. 1C–E**).
From these experiments, we conclude that the loss of Rbp3 affects the growth of cells
as well as their photosynthetic properties, as indicated by the lower pigment content.
This phenotype could be rescued by introducing a gene encoding Rbp3.

To allow the detection of Rbp3 by Western blotting and confocal microscopy, we
complemented Δ*rbp3* mutants also with gene variants expressing a C-terminally fused
3xFLAG or sfGFP tag. To exclude that these tags altered the Rbp3 function, we verified
complementation of the Δ*rbp3* mutant phenotype by expressing Rbp3-3xFLAG or
Rbp3-GFP using absorption and 77 K fluorescence emission spectroscopy (**Fig. 2D,**
E and *SI Appendix,* **Fig. S2).** Notably, the Δ*rbp3* cells were significantly smaller than
the wild-type cells, a phenotype that was also complemented by the ectopic expression
of Rbp3-sfGFP (*SI Appendix,* **Fig. S3).**

Verification of *in-vivo* RNA binding of Rbp3

Previous analyses suggested that Rbp3 is involved in the targeting of *psaA* and *psbA2*
mRNAs towards the thylakoid membrane (30), but its actual RNA binding has not been
directly demonstrated. To test the *in vivo* RNA binding of Rbp3, cultures expressing
triple FLAG-tagged Rbp3 (strain Rbp3-3xFLAG) were used in an assay using the
enzyme polynucleotide kinase (PNK) (35). The principle is outlined in **Fig. 2A.** After
size-fractionating the complex and transfer to a nitrocellulose membrane, the labeled
RNA was visualized by autoradiography and the protein by incubating the membrane
with anti-FLAG antisera (**Fig. 2B**). For Rbp3, several bands were present in the protein
panel, both for extracts from the crosslinked cultures and for non-crosslinked cultures
used as negative controls. The most prominent band at ~20 kDa corresponded to the
molecular mass of the Rbp3-3xFLAG monomer (19.6 kDa). Below this, three other
bands were observed, which were likely N-terminally truncated Rbp3-3xFLAG

fragments. Two bands >20 kDa, one at ~35 kDa and one at ~70 kDa, were common to both Rbp3 protein lanes, that is, with and without UV treatment. For the crosslinked sample, two bands were specific: a somewhat diffuse band at ~30 kDa and another at ~55 kDa. In the autoradiogram, a single prominent band was observed at ~30 kDa (**Fig. 2B**). Its position and somewhat diffuse appearance indicated that it belonged to the Rbp3-RNA ribonucleoprotein complex, which was also visible in the protein lane at the same molecular mass.

The sfGFP-3xFLAG control exhibited a prominent band just above the 35 kDa marker and several faint bands between ~70 kDa and ~100 kDa in the crosslinked protein sample, absent in the non-crosslinked sample, likely resulting from non-specific UV effects. A weaker band below the primary one indicated degraded sfGFP, implying UV treatment reduced protein stability. In the autoradiogram, only a very faint band was visible, possibly crosslinked sfGFP mRNA or RNA that had been near sfGFP during the crosslinking step. In comparison, the cross-linked Rbp3-3xFLAG sample showed a distinct band in the autoradiogram, matching a band in the western blot. This suggested that Rbp3 bound unknown RNA molecules *in vivo*, which could be isotope-labeled after protein purification and were protected from RNase digestion. The ~10 kDa mass difference between RNA-free Rbp3 (~20 kDa) and the RNA-Rbp3 complex indicated protected RNA fragments of approximately 60 nt. The diffused appearance of this band suggested a heterogeneous RNA population in length, composition, or both. The fact that Rbp3 was detected both with and without bound RNA suggested the existence of two pools of this protein.

The majority of Rbp3 RNA interactors are transcripts encoding membrane-associated proteins

The results of the PNK assay suggested that Rbp3 was interacting *in vivo* with an RNA population heterogenous in length, composition, or both (**Fig. 2B**). To identify these transcripts, CLIP-Seq analyses were performed. Cultures expressing Rbp3-3xFLAG or sfGFP-3xFLAG were crosslinked using 1.25 J cm^{-2} UV-C. After enrichment of the samples with anti-FLAG magnetic beads (***SI Appendix,* Fig. S4**), the protein was proteinase K digested and the RNA purified. On average, we obtained 1130 ± 269 ng of RNA from 200 mL of UV-treated Rbp3-3xFLAG cultures and 583 ± 107 ng from sfGFP-3xFLAG cultures. For control, we prepared total RNA from 25 mL aliquots of

the same cultures after UV treatment. All samples were prepared in biological

triplicates. After reverse transcription and sequencing, the RNA sequencing data were

mapped to the genome of *Synechocystis* 6803, followed by a DESeq2 (36) analysis of

transcripts enriched in the Rbp3-3xFLAG pulldown compared to the sfGFP-3xFLAG

pulldown. A total of 229 transcripts were significantly enriched (log$_2$FC > 1, p ≤ 0.05),

including many mRNAs encoding membrane-associated proteins (**Fig. 2C**). Of these

enriched transcripts, 41 were mRNAs encoding proteins associated with the thylakoid

membrane, 21 encoded proteins associated with the plasma membrane, and seven

proteins involved in carboxysome formation and carbon fixation. Additionally, 16

transcripts encode cytoplasmic proteins, for all other no location was annotated by a

GO term. The transcripts with the highest enrichment were *psaA* and *psaB*, organized

in a dicistronic operon and encoding the core proteins of photosystem (PS) I. Other

highly enriched transcripts were *epsB* (*sll0923*) encoding a cell membrane protein, the

tyrosine kinase Wzc related to the production of exopolysaccharides (37), *atpH*

(ssl2615) encoding the small, membrane-intrinsic ATP synthase subunit c, *apcE*

encoding the core-membrane linker polypeptide ApcE (or L$_{CM}$), which anchors the giant

phycobilisome light-harvesting structure to the membrane (38), *cpcL/cpcG2* encoding

an alternative linker protein that facilitates energy transfer from a small phycobilisome

type directly to PSI (39), and *psbA* encoding the PSII core protein D1. Additionally, the

transcript for the cysteine-rich small protein IsiE, which is encoded in the *isiA-isiB*

intergenic region (40) was highly enriched. For the complete list of 229 transcripts

enriched in the co-IP with Rbp3 after UV cross-linking, see ***SI Appendix,*** **Table S1**.

In order to test whether deletion of *rbp3* had an effect on the *psaA* mRNA content per

cell we used *in situ* FISH analysis. The results showed a significantly reduced *psaA*

mRNA content per cell in Δ*rbp3* compared to the wild type and the complementation

control Δ*rbp3*+Rbp3-sfGFP (**Fig. 2F, G**).

Based on these analyses, we conclude that Rbp3 binds several different mRNAs, the

most abundant of which are the *psaA* and *psaB* mRNAs. The interaction of Rbp3 with

these mRNAs is relevant for their accumulation level per cell and this should influence

the number of PSI units.

Specific transcript regions are enriched in the Rbp3 interactome

The enrichment of certain mRNAs in the analysis shown in **Fig. 2C** gives no information
whether certain regions in these transcripts were more highly enriched than others.
However, coverage plots of selected mRNAs indicated a highly uneven distribution of
reads retained from crosslinking to Rbp3. For the dicistronic *psaAB* transcript, a peak
was observed close to the 3′ region and overlapping the stop codon. Enrichment in
regions close to the mRNA 3′ end was also observed for *psbA2* (*SI Appendix,* **Fig.**
S5A, B). In contrast, we observed for *rne* encoding RNase E, a clear peak in the middle
of its 583 nt long 5′UTR, beginning exactly at the previously defined U-rich site (41) (***SI***
Appendix, **Fig. S5C**). A peak within the 5'UTR was also observed for *desC* (***SI***
Appendix, **Fig. S5D, E**).
Since no sequence similarity was detected between the peaks, we calculated the
secondary structures of selected peak regions and the surrounding segments using
RNAfold (42). For the transcript segments enriched for *psaAB* and *psbA2*, stem-loop
structures were predicted that included the respective stop codon and short stretches
of   unpaired   nucleotides.   These   structures   are   located   upstream   of   the
Rho-independent terminators (**Fig. 3B,** *SI Appendix,* **Fig. S5B**). For the element at
the end of the *psaB* mRNA, we tested these interactions by Bio-layer Interferometry
(BLI) using recombinant Rbp3 and biotin-labeled 50 nt ribooligonucleotides
representing the most-enriched region of the *psaB* mRNA. For comparison, we tested
the 50 nt sequences immediately before or after the peak segment (**Fig. 3C**). The
binding height towards the sequence of the coverage peak containing the STOP codon
and the predicted Rbp3 binding site was higher than the binding height towards
neighboring sequences (**Fig. 3D)**. To gain further insight in the likely interacting RNA
motifs, we used RRMScorer, a recently developed algorithm that predicts the most
probable interacting nucleotide segments (13). RRMScorer identified a 5′-UUNGN-3′
motif (*psaAB*: UUAGG and UUCGG; *psbA2*: UUGGU; *desC*: UUCGA) in the three
shown enriched transcript segments with confidence scores between -0.248 and -
0.374, placing them clearly in the likely-binder region (13). These motifs include or are
directly adjacent to unpaired nucleotide(s). Previously characterized RRM-domain
containing RBPs bind single-stranded as well as structured RNA segments (13).
Therefore, these results are in line with previous reports on other RRM-containing
RBPs.
The two high-scoring motifs predicted by RRMScorer for the *psaB* mRNA are located
immediately before the stop codon (motif 2) and shortly after (motif 1; **Fig. 3B**), i.e. in

a functionally sensitive region. To verify their relevance for the interaction with Rbp3,
we mutated motif 2 (UUCGG) to CCUAA and performed another analysis by BLI. The
result clearly showed a much-reduced interaction with Rbp3 (**Fig. 3E**). Therefore, motif
2 is indeed relevant for the interaction, either directly, or via an effect on the secondary
structure, or both.

Different from these results, we observed enrichment along longer transcript stretches
for mRNAs encoding carboxysome shell proteins or RuBisCO subunits (***SI Appendix,***
Fig. S5F–H).

Rbp3 co-immunoprecipitates with the ribosome and LabA/NYN-domain proteins

The appearance of differently-sized signals for Rbp3 binding RNA and not binding RNA
in our PNK assay (**Fig. 2B**) and the previous observation of involvement in thylakoid
membrane targeting (30) posed the question if one or the other Rbp3 fraction would
interact with other cellular structures. To identify possibly interacting proteins, co-
immunoprecipitation (co-IP) analysis of Rbp3-3xFLAG was performed followed by
mass spectrometry-based identification of the enriched proteins. A parallel analysis
using sfGFP-3xFLAG as bait protein was used as control. Both analyses were done in
triplicates and in total 686 proteins were detected. From these, 64 were significantly
enriched ($\log_2$FC > 1, p = 0.05) for co-immunoprecipitating with Rbp3 (**Fig. 4, *SI***
Appendix, **Table S2**). Forty-one of these 62 proteins belong to the ribosome and were
enriched with a $\log_2$FC > 2. Four ribosomal proteins could not or only be detected in
one or two replicates in the sfGFP control (**Fig. 4B**). Two proteins, Slr1870 and
Slr1859, were detected only in the Rbp3 pulldown. Slr1870 is an uncharacterized
protein that harbors a LabA/NYN domain in the N-terminal and central two thirds, and
an OST-HTH domain in the C-terminal third. We noticed that another highly enriched
protein, Slr0650 ($\log_2$FC of +3.97), has a very similar structure. The enriched protein
Slr1859 was characterized as an anti-sigma factor antagonist (43). Its gene *slr1859*
belongs to a cluster that previously was characterized as involved in the regulation of
carbon metabolism, particularly glucose metabolism (44). Besides the enrichment of
ribosomal proteins, proteins involved in RNA degradation like RNase J (Rnj) or the
polyribonucleotide nucleotidyltransferase (Pnp) were enriched. Another enriched
protein was the flavodiiron protein Flv2, which is involved in alternative electron
transport and photoprotection (45). The uncharacterized protein Sll0518 that harbors

two oligonucleotide/oligosaccharide-binding (OB-fold) domains and is encoded directly
downstream of Rbp1, had a log$_2$FC of +3.69. In addition, seven more such potentially
relevant other proteins were enriched, Slr0006, a TsaC fold protein involved in the N6-
threonylcarbamoyladenosine modification of tRNA (46), the PilL-C protein Slr0322
(47), the thylakoid-associated ssDNA-binding protein Slr1034, as well as Slr0635,
Slr1223 and Slr1854.

The ribosome-associated protein Rps1A was significantly enriched, whereas Rps1B
showed no significant enrichment due to its high abundance in the second replicate of
the sfGFP pulldown. Both Rps1A and Rps1B were suggested to be involved in the
Shine–Dalgarno-independent initiation of translation (48, 49).

To independently verify the association of ribosomes and Rbp3, RNA-FISH analysis of
16S rRNA was performed in cells expressing Rbp3-sfGFP. The Pearson's correlation
coefficient in the overlay between the sfGFP and the TAMRA (16S rRNA) signal was
close to 1 (median: 0.935), strongly supporting the close interaction between
ribosomes and Rbp3 (**Fig. 4C**). FISH labelling of the *psaA* mRNA in the same strain
and microscopic analysis revealed a significant overlap with Rbp3-sfGFP fluorescence
(**Fig. 4D**). The Pearson's correlation coefficient in the overlay between the sfGFP
signal and the *psaA* mRNA was positive (median: 0.352) supporting their interaction.
That the correlation was not higher is consistent with the detection of two Rbp3 forms
in the PNK assay, one binding RNA and the other non-binding (**Fig. 2B**), as well as the
fact that our UV cross-linking experiment revealed interaction also with other
transcripts than the *psaAB* mRNA (**Fig. 2C**).

Transcriptomic analyses of the Δ*rbp2*Δ*rbp3* double mutant

To study the effect of Rbp3 deletion and to prevent complementation by Rbp2 we
analyzed the transcriptome of the Δ*rbp2*Δ*rbp3* double mutant that was used previously
in the investigation of mRNA thylakoid membrane targeting (30). The Rbps in
cyanobacteria, especially Rbp1, are considered to be important factors in cold stress
acclimation (19, 32). However, Rbp2 and Rbp3 are also somewhat cold-inducible (33,
34). Therefore, we analyzed the transcript levels 1 h, 2 h and 4 h after transferring the
cultures from 30°C to 20°C. In addition, since the lack of Rbp3 affected photosynthetic
properties (**Fig. 1C–E**), we performed an analysis 0.5 h, 1 h and 2 h following a shift to

high light (250 µmol photons m^{-2} s^{-1}). Subsequently, we performed microarray

analyses of selected samples to identify the full set of transcriptomic differences.

Transcriptomic differences in the comparison of Δ*rbp2*Δ*rbp3* versus wild type

under non-stress conditions and after shift to 20°C

Effects on *psaAB* mRNA accumulation were already observed 1 h after cold stress

induction and 0.5 h after shift to high light (**Fig. 5A, B**). Therefore, we focused on the

comparison of transcriptomic data from these time points to the respective non-stress

control. Under non-stress conditions (T0), the levels of 8 transcripts were significantly

increased, and of 37 transcripts decreased in the Δ*rbp2*Δ*rbp3* to wild type comparison

(**Fig. 5C, *SI Appendix,* Table S3**). The *psaA* and *psaB* mRNA levels ($\log_2$FC of -0.3)

showed only a small decrease under these steady-state conditions (***SI Appendix,***

Table S3). The most decreased transcript levels ($\log_2$FC < 2.0), besides *rbp2* and *rbp3*,

were observed for *sll0788*, which encodes a DUF305 membrane protein and the

plasmid pSYSX-located genes *slr6040* and *slr6039* (*hik31a* operon*),* along with a

slightly decreased level for the neighboring *slr6041* (**Fig. 5C**). The two *hik31* operons

are involved in the control of heterotrophic growth, of acclimation to dark and low O_2

conditions, and in the responses to several stress conditions (50). Moreover, the *hik31*

operon is involved in copper resistance of *Synechocystis* 6803 (51). The eight

upregulated transcripts included the complete transcriptional unit *pilA1*/*pilA2*/*sll1696*

(***SI Appendix,* Table S3**). In addition, *ndhC* encoding the NDH-1 complex core subunit

NdhC was upregulated (which was also enriched in the Rbp3 CLIP-Seq analysis, **Fig.**

5C, *SI Appendix,* Table S1). Interestingly, the *rbp1* transcript was upregulated when

rbp2 and *rbp3* were lacking, indicating a complementary function of the only remaining

RRM-domain-containing protein. Furthermore, *slr0552* and *slr0601,* encoding proteins

of unknown function and *slr0460* encoding an IS701-like transposase were

upregulated.

After 1 h of cold stress, 61 transcripts showed a significantly increased level and 64 a

significantly decreased level in the double mutant compared to wild type (**Fig. 5D**).

Notably, *psaA* ($\log_2$FC = -1.64) and *psaB* ($\log_2$FC = -2.17), the most enriched mRNAs

in our CLIP-Seq analysis (**Fig. 2C**), showed the most decreased transcript levels. The

levels of two other transcripts encoding PSI proteins were decreased in Δ*rbp2*Δ*rbp3*

as well, *psaJ* (*sml0008*) with a $\log_2$FC of -1.43 and *psaF* (*sll0819*) with a $\log_2$FC of -

1.25. Other highly downregulated genes ($\log_2$FC < -2) were again the *hik31a* operon,

two mRNAs encoding proteins of unknown function (*sll0788* and *sll8018*), as well as

sll1621 encoding peroxiredoxin. Several downregulated transcripts were identified as

Rbp3 interaction partners in CLIP-Seq, such as *atpA*, *atpH, atpG* and *atpD* encoding

ATP synthase subunits, mRNAs encoding components of the phycobilisome (*cpcD*),

the cytochrome b_6f complex (*petA* and *petD*), and proteins involved in carbon fixation

(*rbcS* and *ccmM*) (**SI Appendix, Table S3**).

Five transcripts showed highly increased levels ($\log_2$FC > 1.5, *p* = 0.05) in Δ*rbp2*Δ*rbp3*

compared to wild type (**Fig. 5D**). Two of these transcripts (*pilA2* and *sll1696*) already

showed higher levels under normal conditions. The other three transcripts included the

sRNA *csiR1* and *slr1214* encoding the PatA-type response regulator NixC (47), also

called LsiR (52). Gene *slr1214* is located downstream of the highly abundant *csiR1*

sRNA and does not have an own promoter. The end of the *csiR1* transcript was

significantly enriched in the Rbp3-3xFLAG CLIP-Seq pulldown, indicating a regulation

at the RNA level (**SI Appendix, Fig. S5I**). The full transcriptome dataset is visualized

in **Dataset 1 Cold**.

Transcriptomic differences in the comparison of Δ*rbp2*Δ*rbp3* versus wild type

after shift to high light

The T0 samples from the cold shift and the light shift experiments were similar, but not

identical due to different cell densities in the start cultures (OD_{750} of ~ 1.0 in the

temperature experiment and ~0.6 in the light experiment). In cells grown under

standard light conditions (T0), the steady-state levels of 58 transcripts were

significantly increased and of 58 transcripts, besides *rbp2* and *rbp3,* significantly

decreased ($\log_2$FC ≥ $\lvert 1 \rvert$, *p* = 0.05) in the double mutant compared to wild type (**Fig.**

5E, SI Appendix, Table S4). Notably, the *hik31a* operon, *sll0788, sll8018*, and *sbtA*

and *sbtB*, encoding the bicarbonate transporter SbtA and its regulator SbtB (53), were

strongly decreased.

Among the 58 transcripts with increased levels were three non-coding RNAs: *ncl0430*,

a small asRNA transcribed from the complementary strand in the *psaA-psaB* intergenic

spacer, *ncl0680,* constituting the 5′UTR of the *kaiC2B2* operon and *ncr0860*

constituting the 5′UTR of the *hik34* gene (*slr1285).* Moreover, four transcripts encoding

proteins of unknown function (*sll1696, slr0959, slr1915* and *slr1674*). Among the most

increased transcripts were *slr0853* encoding the peptide-alanine-alpha-N-acetyltransferase RimI, *rbp1* and *ssr2227*, which encodes a putative transposase.

After 30 minutes of high light exposure (250 µmol photons m^{-2} s^{-1}), the levels of 22 transcripts were significantly increased ($log_2FC > 1$) and the levels of 40 transcripts decreased ($log_2FC < -1$) (**Fig. 5F**). Again, the *hik31a* operon, as well as the transcripts of *sll8018*, *sll1621* and *sll0788* showed the strongest decreases. Additionally, transcript levels of the cell division gene *sepF* (54) were notably lower in Δ*rbp2*Δ*rbp3*. The transcript levels of some photosynthetic proteins showed a strong decrease ($log_2FC < -1.5$), namely *cpcD*, *psaB* and *cpcC1*. These three transcripts were also significantly enriched in the Rbp3 CLIP-Seq dataset, as were other transcripts showing a decreased level upon light stress (***SI Appendix,* Table S1, S4**).

Of the 22 transcripts with increased levels, only four exhibited a $log_2FC > 1.5$ and only *rbp1* had a $log_2FC > 2$, underscoring the importance of the only remaining RRM-domain protein in Δ*rbp2*Δ*rbp3*. Among the transcripts with highly increased levels were furthermore the transcript for the phycobilisome degrading protein NblD (55) and the non-coding RNA *ncr0380*. The full transcriptome dataset is visualized in **Dataset 2 HL**. In summary, in cells deficient in both the *rbp3* and *rbp2* genes, the transcript levels of *psaA* and *psaB* were significantly lower under cold stress and high light stress compared to wild type (**Fig. 5**). The most pronounced reduction in *psaAB* transcript levels occurred 1 h after shifting from 30°C to 20°C (**Fig. 5A**). This decrease was detected by microarray analysis (**Fig. 5C–F,** ***SI Appendix,* Tables S3, S4**) and northern blot analysis (**Fig. 5A, B**). Transcript levels of several stress response proteins were also altered following *rbp2* and *rbp3* deletion.

Discussion

All sequenced cyanobacterial genomes harbor a gene family encoding RRM domain-containing RBPs, typically consisting of three members, but sometimes more, such as eight in the N_2-fixing, multicellular *A. variabilis* (27). RNA binding was previously verified for these eight Rbps using an *in vitro* assay (27). More recently, *in vitro* RIP-Seq analyses of *Synechocystis* 6803 recombinant Rbp1 and Rbp2 after total RNA incubation identified transcripts mainly related to iron uptake (26).

Here, we show that Rbp3, one of the three different RRM-type RBPs in *Synechocystis* 6803, binds RNA *in vivo* (**Fig. 2B**) and used CLIP-Seq to determine the interacting

RNA population (**Fig. 2C**). The results are consistent with our previous report that Rbp3 is involved in targeting mRNAs encoding the core subunits of both photosystems to the thylakoid membrane, and demonstrate that Rbp3 physically interacts with them.

The lack of Rbp3 has consequences. Whole cell absorption spectra showed that the Δ*rbp3* mutant had a lower chlorophyll-to-phycocyanin ratio and a lower chlorophyll content per cell as suggested by the lower A_{680}:OD_{750} ratio (**Fig. 1C, 2D** and **SI Appendix, Fig. S2A**). Furthermore, 77 K fluorescence emission spectra normalized to the PSI peak showed that the Δ*rbp3* mutant has a lower PSI:PSII ratio compared to the wild type (**Fig. 1D, 2E** and **SI Appendix, Fig. S2B**). Most of the chlorophyll in the cell is associated with PSI and therefore, this combination of effects is indicative of a reduced cellular PSI content (56, 57). This result is furthermore consistent with our finding that the *psaAB* mRNAs were the most-enriched transcripts in our CLIP-Seq analysis (**Fig. 2C**). Our CLIP-Seq results also identified several other mRNAs interacting with Rbp3. Most prominent were mRNAs encoding linker proteins of the phycobilisome, such as *apcE, cpcC1* and *cpcL*, as well as a protein of the ATP synthase complex that locates to the same membrane system. It is not unlikely that thylakoid membrane interactions of these proteins follow similar principles. This is in line with recent observations in an *Nostoc* sp. PCC 7120 deletion mutant lacking RbpG, the ortholog of *Synechocystis* 6803 Rbp3, which showed perturbed thylakoid membrane organization and function (58).

However, some mRNAs enriched by CLIP-Seq, such as *epsB*, encode proteins that are more likely to interact with the plasma membrane and even non-membrane proteins, most notably carboxysome shell proteins and RuBisCO (*rbcLXS* operon, see **SI Appendix, Fig. S5F**). These findings suggest a broader role for Rbp3, while supporting its previously proposed function, along with Rbp2, in localizing specific photosynthetic mRNAs to the thylakoid surface (30).

As interacting sites within the bound transcripts, we found mainly predicted folded secondary structure elements in which the RRMscorer algorithm identified U-rich pentanucleotide motifs (UUNGN) with at least one unpaired nucleotide or U-G base pair. Eukaryotic RRM proteins are highly flexible, typically binding single-stranded RNA, and occasionally single-stranded DNA (59, 60), although structured RNA elements are also documented (13). It is interesting to note that sequences within mRNA 3′UTRs have been recognized to contribute to the correct targeting of the encoded proteins to intracellular membranes (61, 62). The term memRBPs has been

proposed for the "membrane-associated RNA-binding proteins" involved (63), and
well-characterized memRBPs, like the fungal Rrm4, contain RRM domains (64). The
analogy between Rbp3, which interacts with the 3′UTRs of mRNAs encoding key
proteins of the photosynthetic machinery, and eukaryotic RRM proteins, which are
involved in transport by targeting stem-loop elements in mRNA 3′UTRs, is striking.
Thus, Rbp3 can be considered as a bacterial memRBP.

To perform their transport functions, memRBPs must interact with other cellular
structures; for instance, Rrm4 shuttles along microtubules (64, 65). Although several
elements of these mRNA transport systems have been elucidated, they are far from
being fully understood (66).

Rbp3 likely does not function independently of other proteins. Mechanisms must exist
to sort and recognize the targeted membrane or cellular compartment and to prevent
premature translation. Therefore, we enriched proteins by co-IP to identify those
interacting directly with Rbp3 or forming larger complexes nearby. A clear result of this
approach was the enrichment of the entire set of ribosomal proteins (**Fig. 4**), indicating
that complete ribosomes, not just subunits, were near Rbp3. However, recent
fractionation of ribonucleoprotein complexes by Grad-Seq did not indicate co-
fractionation of Rbp3 and ribosomes (7, 8). Thus, while a direct association of Rbp3
with ribosomes cannot be definitively excluded, there is no direct evidence supporting
it, leading to the model shown in **Fig. 6A** in which Rbp3 is linked to ribosomes via a
bound mRNA molecule.

For mRNAs encoding PS core subunits, Rbp3 interacts with regions close to, or
overlapping the stop codons. These mRNAs are highly translated and associated with
polysomes, explaining the enrichment of ribosomal subunits in the Rbp3 co-IP. When
the first ribosomes reach the stop codon region, their helicase activity likely unwinds
the secondary structures where Rbp3 resides. As a result, it may be released and
initiate interaction with another mRNA. Nascent polypeptides probably begin inserting
into the membrane, keeping the mRNA-ribosome complexes associated with thylakoid
surfaces, even after Rbp3 is released from the 3′ region. However, somehow
membrane contact must be established and maintained. A conserved PDPRWA
sequence motif within the C-terminal region of Rbp3 orthologs (26) is located within a
large intrinsically disordered region of the protein (**Fig. 6B**). Prediction of putative lipid
binding sites using the DisoLipPred algorithm, which was also trained on bacterial

datasets (67), yielded a high score, suggesting its possible involvement in the
interaction of Rbp3 with the membrane surface.

An unexpected finding in our co-IP experiments was the enrichment of Slr1870 and
Slr0650, two structurally similar proteins with a LabA/NYN domain and an additional
OST-HTH domain in Slr1870. A third such protein, Slr0755, was not enriched. The
OST-HTH domain was predicted as RNA-binding and some eukaryotic proteins with
that domain nucleate or organize ribonucleoprotein complexes (68). The relationship
between Rbp3 and these proteins is unclear, but a similar structure is found in PIN-
related RNases, which may be involved in tRNA or rRNA maturation. These proteins
belong to the low-amplitude and bright (LabA) superfamily, including the
Synechococcus elongatus LabA, which is linked to circadian timing (69). A LabA
paralog, LalA, was described in *S. elongatus*, which is also involved in circadian gene
expression, as well as in cell growth regulation (70). Recently, the *S. elongatus* Rbp2
homolog was reported to interact with the circadian clock protein KaiC, and disruption
of its gene resulted in altered gene expression rhythms and KaiC localization (31). This
suggests a connection between the cyanobacterial Rbps and the circadian clock
system; however, Slr0755, the ortholog of *S. elongatus* LabA was not enriched in our
co-IP, and the KaiC-interacting protein was the *S. elongatus* Rpb2 homolog and not
Rbp3 addressed here.

Another interesting interacting protein was Slr0006, which binds tightly to membranes
(71) and colocalizes with 30S ribosomal proteins and rRNA (72). Slr0006 has a TsaC
fold, indicating its role in the N6-threonylcarbamoyladenosine modification in most
ANN-decoding tRNAs (46), aligning with the enrichment of ribosomal proteins and
other RNA maturation factors.

There is evidence of cross-talk between the three Rbps in *Synechocystis* 6803, as
indicated by the upregulation of *rbp1* in the Δ*rbp2*Δ*rbp3* double mutant (**Fig. 5C, E, F**)
and our observation that it is possible to generate double deletion mutants between
any combination of these *rbp* genes, but not triple deletions. How this cross-talk is
mediated is currently unknown, but we noticed the co-IP of possible regulatory proteins
with Rbp3 such as Sll0518 (**Fig. 4A**). The gene *sll0518* is located directly downstream
of *sll0517* encoding Rbp1, making it a candidate for possible cross-talk between
different Rbps, especially during cold stress acclimation.

Similar targeting mechanisms probably exist for plasma membrane proteins (73) and,

as our data indicate, potentially for cellular microcompartments, like carboxysomes.

Moreover, the complex mechanisms involved in mRNA targeting may have contributed

to the retention of genes in plant chloroplasts over evolutionary time scales (the 'mRNA

targeting hypothesis') (74). While some components must be distinct, others might be

shared. Clarification of the functions of the here identified non-ribosomal proteins and

of the transcript segments that distinguish them from other transcripts is a promising

topic for future research.

Materials and Methods

Culturing conditions and strain construction

Synechocystis 6803 GT-I (75) wild type and mutant strains were cultivated in copper-free BG11 supplemented with 20 mM TES buffer pH 7.5, under continuous white light of 50 µmol photons m^{-2} s^{-1} at 30°C (standard conditions). Mutant strains containing pVZ322 were cultivated in the presence of 50 µg/mL kanamycin and 2.5 µg/mL gentamycin, while Δ*rbp3* was kept at 50 µg/mL kanamycin. The Δ*rbp2*Δ*rbp3* double mutant strain was kept at 10 µg/mL chloramphenicol and 50 µg/mL kanamycin in the pre-cultures. Antibiotics were omitted during experimental assays. To test the acclimation responses of the Δ*rbp2*Δ*rbp3* double mutant and wild type to downshifts in temperature or to upshifts in light intensity, cultures were transferred at an OD_{750} of ~ 1.0 from 30°C to 20°C, or at an OD_{750} of ~0.6 from a light intensity of 50 to 250 µmol photons m^{-2} s^{-1}. Kanamycin was omitted during co-IP experiments. For mRNA-FISH and Rbp3-GFP localization microscopy, *Synechocystis* 6803 PCC-M (76) was used due its larger cell size, facilitating high-resolution microscopy and protein localization. For details of the mutant strain construction and spectrometric characterization, see ***SI Appendix.***

Spectroscopic analyses

Synechocystis 6803 cultures for spectroscopic analyses were grown as described above but at light intensities of 20 µmol photons m^{-2} s^{-1}. Room-temperature absorption spectra of cell suspensions were measured with a modernized Aminco DW-2000 ultraviolet-visible light spectrophotometer (Olis) and normalized to 750 nm. 77 K fluorescence emission spectra were recorded with a Perkin-Elmer LS55 luminescence spectrometer equipped with a liquid nitrogen housing. Cells were collected and prepared as previously reported (30). Fluorescence spectra were recorded with excitation at 435 nm or 600 nm and emission at 620–750 nm. Excitation and emission slit widths were 5 nm. Emission spectra were corrected for the instrument spectral response and normalized to the PSI or phycobilin fluorescence peaks after subtraction of the background signal.

Protein induction, cell harvest and lysis

The expression of *rbp3-3xFLAG* as well as *sfGFP-3xFLAG* was induced in
corresponding culture volumes at an OD_{750} of ~ 0.6 by adding Cu_2SO_4 to a final
concentration of 0.31 µM. After another 24 h of cultivation, the cells were harvested by
centrifugation (4,000 *x* g, 4°C, 20 min). Cell pellet is resuspended in the respective
lysis buffer. For cell lysis, mechanical disruption using a prechilled Precellys
homogenizer (Bertin Technologies) was used. To remove intact cells and glass beads,
the culture was centrifuged (800 *x* g, 5 min, 4°C), and the supernatant was collected
for further processing. To solubilize membrane proteins, the cell lysates were
incubated for 1 h in the presence of 2% n-dodecyl β-D-maltoside in the dark at 4°C.
Afterwards, cell debris was removed by centrifugation (21,000 *x* g, 4°C 1 h).
Supernatant was then used respectively.
623

Production of recombinant Rbp3

To obtain the intact recombinant Rbp3 protein, it was first expressed as a GST fusion
protein (GST-Rbp3) and then the GST tag was removed. The *rbp3* gene (*slr0193*) was
amplified and ligated into vector pGEX-6P-1 using the In-Fusion Cloning kit (TaKaRa,
Shiga, Japan) and the PCR primers pGEX-6P-1_inv-f/-r and Rbp3_iVEC-f/-r (**SI**
Appendix, **Table S5**). After verification of the sequences, the plasmids were
introduced into *E. coli* Rosetta competent cells (TaKaRa) and used for the expression
and purification of recombinant GST-Rbp3 protein as follows. The resulting *E. coli*
strain was grown in 2 L of LB medium at 37°C. When the culture reached an OD_{600} of
0.6, IPTG was added to the medium at a final concentration of 1 mM and the growth
temperature was shifted to 15°C. After 24 h, cells were harvested by centrifugation,
washed with purification buffer (10% w/v glycerol, 100 mM NaCl, 20 mM HEPES-
NaOH), and stored at −80°C until use. For protein purification, frozen cells were
suspended in 200 mL of purification buffer. The cells were disrupted by sonication and
centrifugated at 10,460 *x* *g* for 30 min at 4°C, and protein affinity purification was
performed using an ÄKTA start chromatography system with a GSTrap column (Cytiva,
MA, USA) according to the manufacturer's instructions. The protein was eluted with 50
mM reduced glutathione and dialyzed against the dialysis buffer 1 (100 mM HEPES-
KOH (pH 7.6), 100 mM NaCl, 1 mM 1,4-dithiothreitol (DTT), dialysis buffer 2 (50% w/v
dialysis buffer 1, 50% w/v glycerol). Following dialysis, GST-Rbp3 was cleaved by

PreScission Protease (Cytiva) to remove the GST-tag and dialyzed again. The purity

of the protein was confirmed by SDS–PAGE and used for the BLI analysis.

SDS-PAGE and Western blotting

Protein separation was done on by SDS-PAGE and protein detection with a

subsequent western blot as described before (77). In short, after co-IP beads in elution

buffer were mixed in a 5:1 ration with 5x protein loading dye (25 mM Tris-HCl pH 6.8,

25% glycerol, 10% SDS, 50 mM DTT, 0.05% bromophenol blue) and boiled for 5 min

at 95°C. Equal volumes of supernatant were separated on a 15% SDS-PAA gel (78).

After separation the proteins were transferred to a nitrocellulose membrane

(Hybond™-ECL, Cytiva) at 1.5 A cm^{-2} for 1 h. After transfer, the membrane was

blocked in 5% skimmed milk powder in TBS-T (50 mM Tris-HCl, 150 mM NaCl,

0.05% Tween20, pH 7.5) overnight at 4°C or for 1 h at room temperature. After

blocking, the membrane was washed three times with TBS-T before incubating with

mouse monoclonal anti-FLAG coupled to HRP (#A8592, Sigma) at a 1:5,000 titer in

TBS-T. After incubation for 1 h at room temperature, signals were visualize using

Western-Blot ECL spray (Advanta) and the FUSION SL Transilluminator (Vilber

Lourmat).

RNA preparation, Northern blotting and hybridization

Culture for RNA extraction was harvested by vacuum filtration on a hydrophilic

polyether sulfone filter (Pall Supor-800) and snap frozen in liquid nitrogen. For RNA

extraction, 1 mL PGTX (79) was given to the filter followed by an incubation of 15 min

at 65°C. RNA was washed and precipitated as described previously (80).

The RNA concentration was determined by a Nanodrop spectrophotometer (peqlab).

Equal amounts of RNA were diluted in water and 1:1 mixed with 2x RNA Gel Loading

Dye (Thermo). Afterwards the samples were denatured at 72°C for 5 min.

RNA was separated on a 1.5% formaldehyde-agarose gel and RNA transfer to a

positively charged nylon membrane (Amersham Hybond-N+, GE Healthcare) occurred

overnight using a capillary blot system. Synthesis of probes and detection of

radioactive signal has been described before (77).

Microarray analysis

The here used Agilent microarrays have been designed for the direct detection of
transcript after chemical labeling, without cDNA synthesis (81). The microarrays
contain oligonucleotide probes for all annotated mRNAs and the majority of non-coding
RNAs. Total RNA (5 µg) extracted from the wild type and deletion mutants grown at
standard temperature (30°C) and after 1 h of cold stress (20°C) or after 0.5 h at high
light was directly labeled using the ULS labeling kit for Agilent gene expression arrays
with Cy3 according to the manufacturer's protocol for Agilent one color microarrays
(Kreatech Diagnostics, B.V., Netherlands). Array analysis was performed using two
biological replicates and an internal technical replicate. Raw data were normexp
background corrected (82) and quantile normalized using the limma R package (83).
The transcriptome differences were calculated between wild type and mutant for the
two different temperatures and the two different light conditions. Significance criteria
for differential expression were $\log_2 FC \geq |1|$, adj. p -value ≤ 0.05. P-values were
adjusted for multiple testing with the Benjamini-Hochberg method. The array data has
been deposited in the GEO database under the accession number GSE269515.

Protein preparation and proteomic sample preparation

After harvesting, the cell pellet was resuspended in 1.5 mL FLAG-MS buffer (50 mM
HEPES/NaOH pH 7, 5 mM $MgCl_2$, 25 mM $CaCl_2$, 150 mM NaCl) containing protease
inhibitor (c0mplete, Roche). The cells were lysed as described above. Co-IP was
performed by incubating 50 µL packed volume of Anti-FLAG M2 magnetic beads
(Sigma) with the cleared cell lysate (1 h, 4°C, rotating). Afterwards the supernatant
was removed using the DynaMag™-2 Magnet rack (Thermo Fisher Scientific) and the
magnetic beads were washed 6x with FLAG-MS buffer. For additional details of the
proteomic sample preparation and analyses by MS, see ***SI Appendix***.

Crosslinking immunoprecipitation and sequencing (CLIP-Seq)

CLIP-Seq was carried out as described (10). Rbp3-3×FLAG and sfGFP-3×FLAG
expression was induced in 250 mL *Synechocystis* cultures with 0.31 µM Cu_2SO_4 at an
OD_{750} of 0.6-0.7. After 24 h, the cultures were transferred to 21   x14.5   x 5.5 cm^3
plastic trays and crosslinked in a UV Stratalinker 2400 (Stratagene) three times at 0.45
J/cm^{-2}. 25 mL culture aliquots were collected to prepare total RNA using the hot-phenol
extraction method (80). The remaining culture was collected as described above. Cell

pellets were resuspended in 1 mL lysis buffer (20 mM Tris/HCl pH 8.0, 1 mM $MgCl_2$,
150 mM KCl, 1 mM DTT) with 1 U of RNase inhibitor per sample (RiboLock, Thermo
Fisher Scientific) and lysed as described above. Supernatants were co-
immunoprecipitated by incubation with 50 µL packed bead volume of Anti-FLAG M2
magnetic beads (Sigma) for 1 h at 4°C.The DynaMag™-2 magnetic rack (Thermo
Fisher Scientific) was used for flow-through removal and washing. Beads were washed
twice with FLAG buffer, then twice with high salt FLAG buffer (50 mM HEPES/NaOH
pH 7, 5 mM $MgCl_2$, 25 mM $CaCl_2$, 1 M NaCl, 10% glycerol, 0.1% Tween20), and then
washed two more times with FLAG buffer. Beads were subjected to proteinase K
digestion (1% SDS, 10 mM Tris-HCl pH 7.5, 100 µg/mL proteinase K) for 30 min and
650 rpm at 50°C. Then, 200 µL PGTX (79) was added and the samples were boiled
for 15 min at 65°C to extract protein-bound RNA. After adding 140 µL
chloroform/isoamyl alcohol (24:1), the samples were incubated for 10 min at room
temperature with agitation. For phase separation, the samples were centrifuged in a
swing-out rotor at 3,260 x g for 3 min at room temperature. The upper aqueous phase
was transferred to an RNase-free tube and mixed in a 1:1 ratio with 100% EtOH and
processed via the clean and concentrator kit (Zymo Research). RNA was eluted in
15 µL RNase-free ddH_2O. Construction of cDNA libraries and sequence analysis were
performed by a commercial provider (Vertis Biotechnologie AG Freising, Germany).
The RNA samples were fragmented using ultrasound (4 pulses of 30 s each at 4°C).
An oligonucleotide adapter was ligated to the RNA 3' ends. M-MLV reverse
transcriptase and a 3′ primer complementary to the adapter sequence were used for
first-strand cDNA synthesis. The obtained cDNA was purified and the 5' Illumina
TruSeq sequencing adapter ligated to the cDNA 3' ends. The cDNA was amplified by
13 cycles of PCR (15 to 18 for the negative control), to about 10-20 ng/µL using a high-
fidelity DNA polymerase and introducing TruSeq barcode sequences as part of the 5'
and 3' TruSeq sequencing adapters. The cDNA was purified using the Agencourt
AMPure XP kit (Beckman Coulter Genomics) and analyzed by capillary
electrophoresis. For Illumina NextSeq sequencing, the samples were pooled in
approximately equimolar amounts. The cDNA pool was 200 – 600 bp size-fractionated
via a preparative agarose gel and sequenced on an Illumina NextSeq 500 system with
a read length of 75 bp.
Raw reads of the RNA-seq libraries were analyzed using the galaxy platform. The
galaxy workflow can be accessed and reproduced at the following link:

https://usegalaxy.eu/u/luisa_hemm/w/poi-3xflag-clip. In short, read quality was check
with FastQC in the beginning. After the quality check, reads were mapped to the
Synechocystis chromosome (GenBank accession NC_000911.1) and the 4 large
plasmids pSYSA, pSYSG, pSYSM and pSYSX (NC_005230.1, NC_005231.1,
NC_005229.1, NC_005232.1) using BWA-MEM (84) with default parameters.
Resulting data were filtered using the BAM filter (85) to remove unmapped reads as
well as reads smaller than 20 nucleotides. The annotation gff-file was based on the
reference GenBank annotations complemented by additional genes based on
transcriptomic analyses (86). htseq-count (87) (mode union, nonunique all, minaqual
0) was used to count the reads per transcript. Only transcripts with ≥ 10 reads in at
least three samples were subjected to differential expression analysis with DESeq2
(36). To visualize coverage of the reads, bam files were transformed into wiggle files
using RSeQC (88) and summarizing the reads from the triplicate analyses for each
position.

PNK assay

Procedures followed previously described methods (8, 35). Rbp3-3xFLAG and sfGFP-
3xFLAG expression was induced in 250 mL *Synechocystis* cultures with 0.31 µM
Cu_2SO_4 at an OD_{750} of 0.6–0.7. After 24 h, cultures were transferred to
21 x14.5 x 5.5 cm^3 plastic trays on ice and crosslinked three times at 0.75 J/cm^{-2} using
a UV Stratalinker 2400 (Stratagene) to induce protein-RNA covalent bonds. Negative
controls were not crosslinked. Cells were collected by centrifugation and the pellets
were resuspended in 800 µL of NT-P buffer (50 mM NaH_2PO_4, 300 mM NaCl, 0.05%
Tween, pH 8.0) containing c0mplete protease inhibitor (Roche). Cell lysis supernatants
were co-immunoprecipitated by incubation with 80 µL packed bead volume of Anti-
FLAG M2 magnetic beads (Sigma) for 1 h at 4°C with slight rotation. The DynaMag™-
2 magnetic rack (Thermo Fisher Scientific) was used for flow-through removal and
washing using 2 mL per wash step. Beads were washed twice with high salt NT-P
buffer (50 mM NaH_2PO_4, 1 M NaCl, 0.05% Tween, pH 8.0), once with NT-P buffer and
once with benzonase buffer (50 mM Tris-HCl, 1 mM $MgCl_2$, pH 8). Beads were
resuspended in 100 µL benzonase buffer containing 25 U benzonase (Sigma). After
incubation for 10 min at 37°C and 900 rpm, beads were washed once with high-salt
NT-P buffer and twice with FastAP buffer (10 mM Tris-HCl, 5 mM $MgCl_2$, 100 mM KCl,

pH 8). Beads were then resuspended in 150 μL FastAP buffer and incubated with 2 U

Fast-AP (Thermo) for 30 min at 37°C and 900 rpm. Beads were washed once with

high-salt NT-P buffer and twice with PNK buffer (50 mM Tris-HCl, 10 mM MgCl2, 0.1

mM spermidine, pH 8) and then resuspended in 100 μL PNK buffer containing 10 μCi

of [γ-^{32}P]-ATP and 1 μL PNK (Thermo). After incubation at 37°C for 30 min at 900 rpm

beads were washed once with high salt NT-P buffer and twice with NT-P buffer. For

elution, beads were resuspended in 35 μL TBS (50 mM Tris-HCl, 150 mM NaCl, pH

7.5) with 0.5 μg/μL FLAG peptide (Sigma) and incubated at room temperature for

15 min at 900 rpm. To the supernatant 5 μL protein loading dye (25 mM Tris-HCl pH

6.8, 25% glycerol, 10% SDS, 50 mM DTT, 0.05% bromophenol blue) was added and

samples were boiled for 5 min at 95°C. Proteins were separated by SDS-PAGE and

western blotting was done as described above. After transfer to the membrane, the

membrane was exposed for several hours to a phosphoimaging screen (Kodak) and

visualized using a Typhoon FLA 9500 (GE Healthcare), followed by western blot

incubation.

Bio-Layer Interferometry

Recombinant Rbp3 protein interactions with RNA oligonucleotide ligands were

examined using the BLI system, BLItz (Fortebio). Ligand RNA oligonucleotides were

synthesized with a biotinylated 5' end (Eurofins) and used for BLI analysis. The binding

buffer (4% glycerol, 10 mM Tris-HCl pH 7.5, 1 mM MgCl$_2$, 0.5 mM EDTA, 50 mM NaCl,

0.5 mM DTT, 0.05 mg/mL poly(dI-dC)·poly(dI-dC), 0.05% [w/v] BSA, 0.01% Tween 20)

was used to dilute both ligand RNA and analyte Rbp3 protein. Experiments were

conducted at room temperature. 0.5 pmol of the biotinylated ligand RNA

oligonucleotides were captured on a streptavidin biosensor (Sartorius) that had been

equilibrated with binding buffer. Rbp3 was bound to the ligand RNA-reacted biosensor

proteins at concentrations of 0.5, 1, 2 or 3 pmol/μL, and the association was analyzed.

The same binding buffer was used for the dissociation step.

mRNA fluorescent *in situ* hybridization

mRNA-FISH was performed as described (89), with adjustments for *Synechocystis*

6803 according to reference (30). Briefly, the target RNA was hybridized with 44–48

oligonucleotides, each 20 nt long, fused to a TAMRA fluorophore at the 3' end.

Oligonucleotide probes for mRNA-FISH were sourced from LGC Biosearch
Technologies (*SI Appendix,* **Table S6)**. For further procedural details, including cell
fixation, permeabilization, hybridization and buffer compositions, see reference (30).
For imaging hybridized *Synechocystis* 6803 PCC-M cells, fixed samples were mounted
on a high precision cover glass covered by a 1.5% agarose pad to minimize cell
movement. Imaging was conducted using an LSM 880 confocal microscope equipped
with an Airyscan detector (Carl Zeiss) and a Plan-Apochromat 63×/1.4NA oil objective
lens. Images were recorded in 16-bit format, at 488 × 488 pixels resolution, 8x line
averaging and a zoom value of 10. The 561 nm, 488 nm, and 633 nm lasers were used
to excite the TAMRA fluorophore, Rbp3-sfGFP fusion protein, and thylakoid membrane
photosynthetic pigments, respectively. To prevent signal bleed-through, a secondary
beam splitter and filter set were installed before the Airyscan detector. A short pass
615 nm beam splitter specified TAMRA emission, and a band pass filter from 495–550
nm isolated sfGFP emission. Image reconstruction used the Airyscan processing
software with automatic strength.

Prior to in-depth image analysis, autofluorescence from RNA-FISH and Rbp3-sfGFP
was removed by background subtraction as described (30). The ImageJ plugin
EzColocalization was used to determine the cell area, calculate mean fluorescence
intensity and perform colocalization analysis (90, 91). Binary images of cell outlines
were created from the thylakoid membrane channel using the auto threshold
command. The adjustable watershed with a tolerance of 5 was applied to distinguish
between adjacent cells. Cells not fully captured in the images were excluded from
analysis. Colocalization of hybridized target mRNA and Rbp3-sfGFP fluorescence was
assessed by calculating the Pearson correlation coefficient using Costes thresholding
(92).

Data availability

The CLIP-seq data have been deposited in the SRA database under the accession
number PRJNA1104562. The mass spectrometry proteomics data have been
deposited to the ProteomeXchange Consortium (93) via the PRIDE (94) partner
repository with the dataset identifier PXD045584. All mass spectrometry proteomics
datasets analyzed during this study are available online at the MassIVE repository

(http://massive.ucsd.edu/; dataset identifier: MSV000092937). The microarray data

have been deposited in the GEO database under the accession number GSE269515.

Funding

We appreciate the support by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) to LH, EL, OS, AW and WRH through the graduate school

MeInBio - 322977937/GRK2344 and grants HE 2544/22-1 to WRH and SCHI 871/11-

1 to OS. The Proteomic Platform – Core Facility was supported by the Medical Faculty

of the University of Freiburg to O.S. (2021/A3-Sch). CWM acknowledges support from

the Biotechnology and Biological Research Council (BB/W001012/1).

References

1. D. L. Kaplan, M. O'Donnell, Rho factor: transcription termination in four steps. *Curr. Biol. CB* **13**, R714-716 (2003).

2. H. Aiba, Mechanism of RNA silencing by Hfq-binding small RNAs. *Curr. Opin. Microbiol.* **10**, 134–139 (2007).

3. S. G. Chaulk, *et al.*, ProQ is an RNA chaperone that controls ProP levels in *Escherichia coli. Biochemistry* **50**, 3095–3106 (2011).

4. W. Bae, B. Xia, M. Inouye, K. Severinov, *Escherichia coli* CspA-family RNA chaperones are transcription antiterminators. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 7784–7789 (2000).

5. A. Bøggild, M. Overgaard, P. Valentin-Hansen, D. E. Brodersen, Cyanobacteria contain a structural homologue of the Hfq protein with altered RNA-binding properties. *FEBS J.* **276**, 3904–3915 (2009).

6. N. Schuergers, *et al.*, Binding of the RNA chaperone Hfq to the type IV pilus base is crucial for its function in *Synechocystis* sp. PCC 6803. *Mol. Microbiol.* **92**, 840–852 (2014).

7. M. Riediger, *et al.*, Analysis of a photosynthetic cyanobacterium rich in internal membrane systems via gradient profiling by sequencing (Grad-seq). *Plant Cell* **33**, 248–269 (2021).

8. M. Brenes-Álvarez, *et al.*, R-DeeP/TripepSVM identifies the RNA-binding OB-fold-like protein PatR as regulator of heterocyst patterning. *Nucleic Acids Res.* gkae1247 (2024). https://doi.org/10.1093/nar/gkae1247.

9. L. Hemm, *et al.*, RAPDOR: Using Jensen-Shannon Distance for the computational analysis of complex proteomics datasets. *bioRxiv* 2024–09 (2024).

10. L. Hemm, *et al.*, Interactors and effects of overexpressing YlxR/RnpM, a conserved RNA binding protein in cyanobacteria. *RNA Biol.* **21**, 1308–1326 (2024).

11. G. Dreyfuss, M. S. Swanson, S. Piñol-Roma, Heterogeneous nuclear ribonucleoprotein particles and the pathway of mRNA formation. *Trends Biochem. Sci.* **13**, 86–91 (1988).

12. C. Maris, C. Dominguez, F. H.-T. Allain, The RNA recognition motif, a plastic RNA-binding platform to regulate post-transcriptional gene expression. *FEBS J.* **272**, 2118–2131 (2005).

13. J. Roca-Martínez, H. Dhondge, M. Sattler, W. F. Vranken, Deciphering the RRM-RNA recognition code: A computational analysis. *PLOS Comput. Biol.* **19**, e1010859 (2023).

14. A. Zhang, W.-F. Liu, Y.-B. Yan, Role of the RRM domain in the activity, structure and stability of poly(A)-specific ribonuclease. *Arch. Biochem. Biophys.* **461**, 255–262 (2007).

15. A. B. Sachs, R. W. Davis, R. D. Kornberg, A single domain of yeast poly(A)-binding protein is necessary and sufficient for RNA binding and cell viability. *Mol. Cell. Biol.* **7**, 3268–3276 (1987).

16. L. Olgeiser, *et al.*, The key protein of endosomal mRNP transport Rrm4 binds translational landmark sites of cargo mRNAs. *EMBO Rep.* **20**, e46588 (2019).

17. M. Grskovic, M. W. Hentze, F. Gebauer, A co-repressor assembly nucleated by Sex-lethal in the 3′UTR mediates translational control of Drosophila msl-2 mRNA. *EMBO J.* **22**, 5571–5581 (2003).

18. G. Caponigro, R. Parker, Multiple functions for the poly(A)-binding protein in mRNA decapping and deadenylation in yeast. *Genes Dev.* **9**, 2421–2432 (1995).

19. K. Maruyama, N. Sato, N. Ohta, Conservation of structure and cold-regulation of
RNA-binding proteins in cyanobacteria: Probable convergent evolution with
eukaryotic glycine-rich RNA-binding proteins. *Nucleic Acids Res.* **27**, 2029–2036
(1999).
20. M. Sugita, M. Sugiura, The existence of eukaryotic ribonucleoprotein consensus
sequence-type RNA-binding proteins in a prokaryote, *Synechococcus* 6301.
Nucleic Acids Res. **22**, 25–31 (1994).
21. M. E. Mulligan, D. M. Jackman, S. T. Murphy, Heterocyst-forming filamentous
cyanobacteria encode proteins that resemble eukaryotic RNA-binding proteins of
the RNP family. *J. Mol. Biol.* **235**, 1162–1170 (1994).
22. N. Sato, A cold-regulated cyanobacterial gene cluster encodes RNA-binding
protein and ribosomal protein S21. *Plant Mol. Biol.* **24**, 819–823 (1994).
23. M. E. Mulligan, T. J. Belbin, Characterization of RNA-binding protein genes in
cyanobacteria. *Nucleic Acids Symp. Ser.* 140–2 (1995).
24. T. Kaneko, *et al.*, Sequence analysis of the genome of the unicellular
cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination
of the entire genome and assignment of potential protein-coding regions. *DNA*
Res. **3**, 109–136 (1996).
25. T. Hamano, *et al.*, Characterization of RNA-binding properties of three types of
RNA-binding proteins in *Anabaena* sp. PPC 7120. *Cell. Mol. Biol. Noisy--Gd. Fr.*
50, 613–624 (2004).
26. Y. Zhang, D. Wu, Y. Wang, X. Xu, Two types of C-terminal regions of RNA-binding
proteins play distinct roles in stress tolerance of *Synechocystis* sp. PCC 6803.
FEMS Microbiol. Lett. **369**, fnac021 (2022).
27. N. Sato, A family of cold-regulated RNA-binding protein genes in the
cyanobacterium *Anabaena variabilis* M3. *Nucleic Acids Res.* **23**, 2161–2167
(1995).
28. N. Sato, K. Maruyama, Differential regulation by low temperature of the gene for
an RNA-binding protein, rbpA3, in the cyanobacterium *Anabaena variabilis* strain
M3. *Plant Cell Physiol.* **38**, 81–86 (1997).
29. M. Mutsuda, M. Sugiura, M. Sugita, Physiological characterization of RNA-binding
protein-deficient cells from *Synechococcus* sp. strain PCC7942. *Plant Cell*
Physiol. **40**, 1203–1209 (1999).
30. M. Mahbub, *et al.*, mRNA localization, reaction centre biogenesis and thylakoid
membrane targeting in cyanobacteria. *Nat. Plants* **6**, 1179–1191 (2020).
31. B. M. McKnight, *et al.*, Roles for the *Synechococcus elongatus* RNA-binding
protein Rbp2 in regulating the circadian clock. *J. Biol. Rhythms* **38**, 447–460
(2023).
32. X. Tan, *et al.*, Role of Rbp1 in the acquired chill-light tolerance of cyanobacteria.
J. Bacteriol. **193**, 2675–2683 (2011).
33. Q. Tang, X. Tan, X. Xu, Effects of a type-II RNA-binding protein on fatty acid
composition in *Synechocystis* sp. PCC 6803. *Chin. Sci. Bull.* **55**, 2416–2421
(2010).
34. Y. Wang, X. Xu, Effects of Rbp3 on lipid peroxidation and salt tolerance in
Synechocystis sp. PCC 6803. *FEBS Lett.* **587**, 1446–1451 (2013).
35. T. S. Stenum, *et al.*, RNA interactome capture in *Escherichia coli* globally identifies
RNA-binding proteins. *Nucleic Acids Res.* **51**, 4572–4587 (2023).
36. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
37. S. B. Pereira, *et al.*, The role of the tyrosine kinase Wzc (Sll0923) and the
phosphatase Wzb (Slr0328) in the production of extracellular polymeric

substances (EPS) by *Synechocystis* PCC 6803. *MicrobiologyOpen* **8**, e00753
(2019).

38. V. Capuano, A. S. Braux, N. Tandeau de Marsac, J. Houmard, The "anchor polypeptide" of cyanobacterial phycobilisomes. Molecular characterization of the *Synechococcus* sp. PCC 6301 apcE gene. *J. Biol. Chem.* **266**, 7239–7247 (1991).

39. L. Zheng, *et al.*, Cryo-EM and femtosecond spectroscopic studies provide mechanistic insight into the energy transfer in CpcL-phycobilisomes. *Nat. Commun.* **14**, 3961 (2023).

40. M. Riediger, M. A. Hernández-Prieto, K. Song, W. R. Hess, M. E. Futschik, Genome-wide identification and characterization of Fur-binding sites in the cyanobacteria Synechocystis sp. PCC 6803 and PCC 6714. *DNA Res. Int. J. Rapid Publ. Rep. Genes Genomes* **28**, dsab023 (2021).

41. S. Watanabe, *et al.*, Regulation of RNase E during the UV stress response in the cyanobacterium *Synechocystis* sp. PCC 6803. *mLife* **2**, 43–57 (2023).

42. D. H. Mathews, *et al.*, Incorporating chemical modification constraints into a dynamic programming algorithm for prediction of RNA secondary structure. *Proc. Natl. Acad. Sci.* **101**, 7287–7292 (2004).

43. L. Shi, K. M. Bischoff, P. J. Kennelly, The icfG gene cluster of *Synechocystis* sp. strain PCC 6803 encodes an Rsb/Spo-like protein kinase, protein phosphatase, and two phosphoproteins. *J. Bacteriol.* **181**, 4761–4767 (1999).

44. L. Beuf, S. Bédu, M. C. Durand, F. Joset, A protein involved in co-ordinated regulation of inorganic carbon and glucose metabolism in the facultative photoautotrophic cyanobacterium *Synechocystis* PCC6803. *Plant Mol. Biol.* **25**, 855–864 (1994).

45. A. Santana-Sanchez, *et al.*, Flavodiiron proteins 1-to-4 function in versatile combinations in O2 photoreduction in cyanobacteria. *eLife* **8**, e45766 (2019).

46. L. L. Carvalho, T. A. Salminen, K. M. Dahlström, Slr0006-like proteins: A TsaC/TsaC2/YciO subfamily exclusive to cyanobacteria. *Mol. Phylogenet. Evol.* **109**, 1–10 (2017).

47. R. Narikawa, *et al.*, Novel photosensory two-component system (PixA-NixB-NixC) involved in the regulation of positive and negative phototaxis of cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* **52**, 2214–2224 (2011).

48. M. Mutsuda, M. Sugiura, Translation initiation of cyanobacterial rbcS mRNAs requires the 38-kDa ribosomal protein S1 but not the Shine-Dalgarno sequence: development of a cyanobacterial in vitro translation system. *J. Biol. Chem.* **281**, 38314–38321 (2006).

49. S. Nakagawa, Y. Niimura, K. Miura, T. Gojobori, Dynamic evolution of translation initiation mechanisms in prokaryotes. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 6382–6387 (2010).

50. S. Nagarajan, D. M. Sherman, I. Shaw, L. A. Sherman, Functions of the duplicated hik31 operons in central metabolism and responses to light, dark, and carbon sources in *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **194**, 448–459 (2012).

51. J. Giner-Lamia, L. López-Maury, J. C. Reyes, F. J. Florencio, The CopRS two-component system is responsible for resistance to copper in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.* **159**, 1806–1818 (2012).

52. J.-Y. Song, *et al.*, Near-UV cyanobacteriochrome signaling system elicits negative phototaxis in the cyanobacterium *Synechocystis* sp. PCC 6803. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 10780–10785 (2011).

53. X.-Y. Liu, *et al.*, Structures of cyanobacterial bicarbonate transporter SbtA and its complex with PII-like SbtB. *Cell Discov.* **7**, 63 (2021).

54. M. Marbouty, C. Saguez, C. Cassier-Chauvat, F. Chauvat, Characterization of the
FtsZ-interacting septal proteins SepF and Ftn6 in the spherical-celled
cyanobacterium *Synechocystis* strain PCC 6803. *J. Bacteriol.* **191**, 6178–6185
(2009).
55. V. Krauspe, *et al.*, Discovery of a small protein factor involved in the coordinated
degradation of phycobilisomes in cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.*
118, e2012277118 (2021).
56. A. Wilde, *et al.*, Inactivation of a *Synechocystis* sp strain PCC 6803 gene with
homology to conserved chloroplast open reading frame 184 increases the
photosystem II-to-photosystem I ratio. *Plant Cell* **7**, 649–658 (1995).
57. N. H. Mann, *et al.*, Involvement of an FtsH homologue in the assembly of
functional photosystem I in the cyanobacterium *Synechocystis* sp. PCC 6803.
FEBS Lett. **479**, 72–77 (2000).
58. K. Wang, M. Mahbub, G. Mastroianni, C. W. Mullineaux, mRNA localization and
thylakoid protein biogenesis in the filamentous heterocyst-forming
cyanobacterium Anabaena sp. PCC 7120. [Preprint] (2024). Available at:
https://www.biorxiv.org/content/10.1101/2024.04.17.589878v1 [Accessed 2 May
2024].
59. J. Ding, *et al.*, Crystal structure of the two-RRM domain of hnRNP A1 (UP1)
complexed with single-stranded telomeric DNA. *Genes Dev.* **13**, 1102–1115
(1999).
60. B. Soubise, Y. Jiang, N. Douet-Guilbert, M.-B. Troadec, RBM22, a key player of
pre-mRNA splicing and gene expression regulation, is altered in cancer. *Cancers*
14, 643 (2022).
61. C. Mayr, Regulation by 3′-untranslated regions. *Annu. Rev. Genet.* **51**, 171–194
(2017).
62. W. Ma, C. Mayr, A membraneless organelle associated with the endoplasmic
reticulum enables 3'UTR-mediated protein-protein interactions. *Cell* **175**, 1492-
1506.e19 (2018).
63. J. Béthune, R.-P. Jansen, M. Feldbrügge, K. Zarnack, Membrane-associated
RNA-binding proteins orchestrate organelle-coupled translation. *Trends Cell Biol.*
29, 178–188 (2019).
64. J. König, *et al.*, The fungal RNA-binding protein Rrm4 mediates long-distance
transport of ubi1 and rho3 mRNAs. *EMBO J.* **28**, 1855–1866 (2009).
65. P. Becht, J. König, M. Feldbrügge, The RNA-binding protein Rrm4 is essential for
polarity in *Ustilago maydis* and shuttles along microtubules. *J. Cell Sci.* **119**, 4964–
4973 (2006).
66. S.-K. Devan, *et al.*, A MademoiseLLE domain binding platform links the key RNA
transporter to endosomes. *PLOS Genet.* **18**, e1010269 (2022).
67. A. Katuwawala, B. Zhao, L. Kurgan, DisoLipPred: accurate prediction of
disordered lipid-binding residues in protein sequences with deep recurrent
networks and transfer learning. *Bioinforma. Oxf. Engl.* **38**, 115–124 (2021).
68. V. Anantharaman, D. Zhang, L. Aravind, OST-HTH: a novel predicted RNA-
binding domain. *Biol. Direct* **5**, 13 (2010).
69. Y. Taniguchi, *et al.*, labA: a novel gene required for negative feedback regulation
of the cyanobacterial circadian clock protein KaiC. *Genes Dev.* **21**, 60–70 (2007).
70. Y. Taniguchi, T. Nishikawa, T. Kondo, T. Oyama, Overexpression of lalA, a
paralog of labA, is capable of affecting both circadian gene expression and cell
growth in the cyanobacterium *Synechococcus elongatus* PCC 7942. *FEBS Lett.*
586, 753–759 (2012).

71. D. Carmel, P. Mulo, N. Battchikova, E.-M. Aro, Membrane attachment of Slr0006 in *Synechocystis* sp. PCC 6803 is determined by divalent ions. *Photosynth. Res.* **108**, 241–245 (2011).

72. D. Carmel, *et al.*, Structural model, physiology and regulation of Slr0006 in *Synechocystis* PCC 6803. *Arch. Microbiol.* **195**, 727–736 (2013).

73. M. Mahbub, C. W. Mullineaux, Locations of membrane protein production in a cyanobacterium. *J. Bacteriol.* **205**, e0020923 (2023).

74. W. R. Hess, A. Wilde, C. W. Mullineaux, Does mRNA targeting explain gene retention in chloroplasts? *Trends Plant Sci.* **30**, 147–155 (2025).

75. Y. Kanesaki, *et al.*, Identification of substrain-specific mutations by massively parallel whole-genome resequencing of *Synechocystis* sp. PCC 6803. *DNA Res.* **19**, 67–79 (2012).

76. D. Trautmann, B. Voß, A. Wilde, S. Al-Babili, W. R. Hess, Microevolution in cyanobacteria: Re-sequencing a motile substrain of *Synechocystis* sp. PCC 6803. *DNA Res.* **19**, 435–448 (2012).

77. A. Kaltenbrunner, *et al.*, Regulation of pSYSA defense plasmid copy number in *Synechocystis* through RNase E and a highly transcribed asRNA. *Front. Microbiol.* **14**, 1112307 (2023).

78. U. K. Laemmli, SDS-page Laemmli method. *Nature* **227**, 680–685 (1970).

79. F. L. Pinto, A. Thapper, W. Sontheim, P. Lindblad, Analysis of current and alternative phenol based RNA extraction methodologies for cyanobacteria. *BMC Mol. Biol.* **10**, 79 (2009).

80. S. Hein, I. Scholz, B. Voß, W. R. Hess, Adaptation and modification of three CRISPR loci in two closely related cyanobacteria. *RNA Biol.* **10**, 852–864 (2013).

81. B. Voß, W. R. Hess, The identification of bacterial non-coding RNAs through complementary approaches. *Handb. RNA Biochem. Second Complet. Revis. Enlarg. Ed.* 787–800 (2014).

82. M. E. Ritchie, *et al.*, A comparison of background correction methods for two-colour microarrays. *Bioinforma. Oxf. Engl.* **23**, 2700–2707 (2007).

83. M. E. Ritchie, *et al.*, limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

84. H. Li, R. Durbin, Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics* **26**, 589–595 (2010).

85. M. R. Breese, Y. Liu, NGSUtils: a software suite for analyzing and manipulating next-generation sequencing datasets. *Bioinforma. Oxf. Engl.* **29**, 494–496 (2013).

86. M. Kopf, *et al.*, Comparative analysis of the primary transcriptome of *Synechocystis* sp. PCC 6803. *DNA Res.* **21**, 527–539 (2014).

87. G. H. Putri, S. Anders, P. T. Pyl, J. E. Pimanda, F. Zanini, Analysing high-throughput sequencing data in Python with HTSeq 2.0. *Bioinformatics* **38**, 2943–2945 (2022).

88. L. Wang, S. Wang, W. Li, RSeQC: quality control of RNA-seq experiments. *Bioinformatics* **28**, 2184–2185 (2012).

89. S. O. Skinner, L. A. Sepúlveda, H. Xu, I. Golding, Measuring mRNA copy number in individual *Escherichia coli* cells using single-molecule fluorescent *in situ* hybridization. *Nat. Protoc.* **8**, 1100–1113 (2013).

90. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

91. W. Stauffer, H. Sheng, H. N. Lim, EzColocalization: An ImageJ plugin for visualizing and measuring colocalization in cells and organisms. *Sci. Rep.* **8**, 15764 (2018).

92. K. W. Dunn, M. M. Kamocka, J. H. McDonald, A practical guide to evaluating
colocalization in biological microscopy. *Am. J. Physiol. - Cell Physiol.* **300**, C723–
C742 (2011).
93. E. W. Deutsch, *et al.*, The ProteomeXchange consortium at 10 years: 2023
update. *Nucleic Acids Res.* **51**, D1539–D1548 (2023).
94. Y. Perez-Riverol, *et al.*, The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552
(2022).
95. K. Darty, A. Denise, Y. Ponty, VARNA: Interactive drawing and editing of the RNA
secondary structure. *Bioinformatics* **25**, 1974–1975 (2009).
96. S. Tyanova, *et al.*, The Perseus computational platform for comprehensive
analysis of (prote)omics data. *Nat. Methods* **13**, 731–740 (2016).
97. J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold.
Nature **596**, 583–589 (2021).
98. M. Mirdita, *et al.*, ColabFold: making protein folding accessible to all. *Nat. Methods*
19, 679–682 (2022).
99. G. Erdős, Z. Dosztányi, AIUPred: combining energy estimation with deep learning
for the enhanced prediction of protein disorder. *Nucleic Acids Res.* gkae385
(2024). https://doi.org/10.1093/nar/gkae385.

1122
1123

1124 **Figures**

1125
1126

Fig. 1. Phenotypical analysis of Rbp3 mutants. (A) Spot assays of the indicated
Synechocystis 6803 strains after 8 days of incubation under standard growth
conditions (40 µmol photons m^{-2} s^{-1}, continuous light 30°C). **(B)** Spot assays under
high light (300 µmol photons m^{-2} s^{-1}) and 5 days of incubation. In both conditions, the
growth of wild type *Synechocystis* GT-I strain was compared to the Δ*rbp3* deletion
mutant and the complementation strain Δ*rbp3*+Rbp3 that expressed *rbp3* from an
introduced pVZ322 plasmid under control of its native promoter. Culture aliquots were
inoculated in triplicates and diluted ten-fold by every step as indicated. Colony sizes
were measured and evaluated (right panels). Significances were calculated by the
Students t-test, for the numeric values see *SI Appendix,* **Table S7**. **(C)** Room-
temperature absorption spectra of cell suspensions for the GT-I wild type, Δ*rbp3* and
Δ*rbp3*+*rbp3*, normalized at 750 nm. **(D)** 77K fluorescence emission spectra with
chlorophyll excitation at 435 nm, normalized to the PSI peak. **(E)** 77K fluorescence
emission spectra with phycocyanin excitation at 600 nm, normalized to phycobilin
emission at 660 nm.

Fig. 2. Verification of RNA-binding of Rbp3. (A) Schematic overview of the PNK

assay (35). The protein of interest (POI) and RNA are crosslinked using UV-C. After

cell lysis, co-IP is performed using anti-FLAG M2 magnetic beads. Non-covalently

bound nucleic acids are removed by benzonase treatment. Nuclease-protected bound

RNA is labeled with ^{32}P using PNK. Purified proteins are separated by SDS-PAGE (78),

transferred to a nitrocellulose membrane and visualized by autoradiography for ^{32}P-

labeled RNA and by western blotting for proteins. Controls omit the crosslinking step.

(B) Rbp3 and sfGFP (negative control) PNK assays. Samples were cross-linked 24 h

after induction of protein expression. After separating complexes by SDS-PAGE and

western blotting, FLAG-tagged proteins were detected immunologically. Several bands

appeared in the protein lanes (left subpanels). The most prominent band indicates

monomeric Rbp3-3xFLAG at ~20 kDa. For sfGFP-3xFLAG, the strongest signal is at

~40 kDa, corresponding to monomeric sfGFP-3xFLAG. Bands below the signal of the

respective monomer indicate degraded protein fragments, as these increased in

intensity after UV treatment. Bands above the respective monomers belong to potential

multimers, stabilized by UV treatment. In the UV-treated Rbp3-3xFLAG sample, a

diffuse protein signal at ~30 kDa (marked with an arrow) overlaps the autoradiography

signal (left panel, + CL, RNA lane). The band at ~70 kDa might belong to a multimer

of Rbp3. **(C)** Volcano plot showing transcripts enriched in the Rbp3_3×FLAG CLIP

RNA samples compared to the sfGFP_3×FLAG CLIP RNA samples. Colored dots

indicate enriched transcripts encoding proteins involved in photosynthesis (green),

proteins with the GO term membrane (yellow), associated with the carboxysome

(purple), or others (dark gray). Light gray dots indicate non-enriched transcripts.

Functionalities or gene names are given for transcripts enriched with a $\log_2$ fold change

≥2, the transcript for Rbp3 is not shown. (**D**) Room-temperature absorption spectra of

cell suspensions for the GT-I wild type, Δ*rbp3* and Δ*rbp3*+*rbp3*-FLAG strains. **(E)** 77K

fluorescence emission spectra, with chlorophyll excitation at 435 nm and normalized

to PSI emission at 725 nm. **(F)** Confocal images showing the localization of *psaA*

mRNA (TAMRA, grey) relative to the thylakoid membrane (chlorophyll, blue) in PCC-

M (WT), *Δrbp3* and *Δrbp3* + Rbp3-sfGFP in the respective PCC-M WT background

strain; scale bar = 2 µm (see ***SI Appendix,* Fig. S2** for the spectral characterization of

the complementation strain expressing Rbp3-sfGFP and **Fig. S3** for the determination

of cell sizes). **(G)** Quantification of *psaA* mRNA fluorescence intensity comparing PCC-

M (WT), *Δrbp3* and *Δrbp3* + Rbp3-sfGFP strains; *n* = number of cells. *** significant

difference ($p < 0.001$). Significances were calculated by the students t-test assuming

equal variances, for the numeric values see ***SI Appendix,* Table S8**.

Fig. 3. Coverage peaks in enriched mRNA reveal Rbp3 binding sites. (A) Read coverage along the *psaAB* locus in the Rbp3-3xFLAG CLIP RNA compared to the sfGFP-3xFLAG CLIP RNA (negative control), total RNA from cultures expressing the Rbp3-3xFLAG or sfGFP-3xFLAG. All cDNA libraries were sequenced in biological

triplicates. **(B)** Secondary structure prediction around the *psaB* read coverage peak.

(**C**) Three *psaB* mRNA regions were selected to test interactions with Rbp3. These

regions encompass 50 nt of the *psaB* coverage peak (center), the preceding 50 nt of

coding sequence (left), or the following 50 nt encompassing the Rho-independent

terminator (right). (**D**) Recombinant Rbp3 was used in an *in vitro* Bio-Layer

Interferometry assay to determine its binding affinities towards the 50 nt segments from

panel C as synthetic RNA oligonucleotides. (**E**) Bio-Layer Interferometry assay with

the *psaB* coverage peak RNA oligonucleotide in which the predicted binding motif 2

was mutated as shown.

In panels B and C, the STOP codons are marked in red, and the putative RRM binding

sites predicted by the RRMScorer algorithm are circled in blue (13). The secondary

structures were calculated by RNAfold (42) and displayed using VARNA (95). For

further mRNA coverage plots, see ***SI Appendix,*** **Fig. S5**.

1197

Fig. 4. **Protein interaction partners of Rbp3 (A)** Volcano plot showing the protein

interaction partners of Rbp3-3xFLAG identified by mass spectrometry compared to

sfGFP-3xFLAG. Further details of the enriched proteins can be found in ***SI Appendix,***

Table S2. For the complete list of detected proteins, see *SI Appendix,* **Table S9**. **(B)** Heatmap showing the $\log_2$ (LFQ intensity) of all ribosomal proteins that could be detected. Hierarchical clustering was done using the Perseus software (96). Rbp3-3xFLAG and sfGFP-3xFLAG expression was induced 24 h prior to harvesting with 0.31 µM Cu_2SO_4. For the analysis, the Rbp3-3xFLAG as well as the sfGFP-3xFLAG bait protein were not included in the plot. Details of statistical analysis in *SI Appendix,* **Table S10**. **(C)** Left panel: Localization of hybridized 16S rRNA (TAMRA, magenta) and Rbp3-sfGFP (green) relative to the thylakoid membrane (blue) or the cellular outline (white) in *Δrbp3* + Rbp3-sfGFP (PCC-M background) cells. Right panel: Colocalization quantification of fluorescent pair (left panel, third image) by calculation of the Pearson's correlation coefficient. n = number of cells. **(D)** Left panel: Localization of hybridized *psaA* mRNA (magenta) and Rbp3-sfGFP (sfGFP, green) relative to the thylakoid membrane (blue) or the cellular outline (white) in *Δrbp3* + Rbp3-sfGFP cells. Right panel: Colocalization quantification of fluorescent pair (left panel, third image) by calculation of the Pearson's correlation coefficient. n = number of cells. Scale bar = 2 µm in panels C and D. For the numeric values in the calculation of statistical significances in panels C and D, see *SI Appendix,* **Table S11**.

1219

Fig. 5. **Transcriptomic analysis of Δ*rbp2*Δ*rbp3* double mutant. (A)** Time course of *psaAB* mRNA accumulation in *Synechocystis* 6803 wild type (GT-I) or the Δ*rbp2*Δ*rbp3* mutant (clone 1, clone 2) after transferring the cultures from 30°C to 20°C for the indicated times. **(B)** Time course of *psaAB* mRNA accumulation after transferring the cultures from 50 to 250 µmol photons m^{-2} s^{-1}. In both panels A and B, 5 µg of total RNA were separated on 1.5% formaldehyde-agarose gels and the RiboRuler High Range RNA ladder (Thermo Fisher) served as molecular mass standard. The single-stranded RNA probe encompassed the region from 100 to 252 nt after the *psaA* start codon. Hybridization against the 5S rRNA served as loading control. (**C, D, E and F)** Volcano plots showing transcripts with significantly increased or decreased levels in the comparison of the double mutant against wild type, 0.5 h after shift from 40 to 250 µmol photons m^{-2} s^{-1}, or 1 h after shifting the cultures from 30°C to 20°C. The respective comparison at T = 0 h is also shown for both conditions. Transcriptomic changes were captured using a custom-made microarray. Salmon dots display increased transcript levels; turquoise dots display decreased transcript levels. Both experiments were performed with two replicates each. Note, that values for *rbp2* and *rbp3* represent background. They were included for completeness and do not indicate detected transcripts. See also **Tables S3** and **S4** and **Datasets 1** and **2** for the complete set of transcriptomic differences.

1239

Fig. 6. Proposed model of transcript localization towards the membrane mediated by Rbp3. (A) Rbp3 binds distinct mRNA motifs such as a secondary structure overlapping the stop codon in case of the *psaAB* mRNA. The Rbp3 interaction with the membrane surface might involve a PDPRWA sequence motif conserved within

the C-terminal section of Rbp3 orthologs (26), identified by the DisoLipPred algorithm

(67) as a lipid binding site. These mRNAs are translated near the membrane surface,

explaining the high co-enrichment of ribosomal proteins with Rbp3. The helicase

activity of the ribosome may unwind the RNA secondary structure, releasing Rbp3. Our

data indicate that Rbp3 exists in both RNA-binding and non-binding states. The place

where RNA binding is initiated and the factors involved in intracellular transport are

unknown (question marks). The other RRM proteins, Rbp2 and Rbp1, likely also bind

mRNAs with specific, potentially overlapping targets with Rbp3. Evidence suggests

cross-talk among these Rbps, though its mechanisms are unclear. **(B)** Rbp3 structure

predicted by AlphaFold (97, 98). The RRM domain characterized by a four-stranded

antiparallel beta-sheet and two alpha helices, is boxed. The structure ends with a C-

terminal short beta fold followed by an alpha helix. Between the RRM domain and the

C-terminal structures, a long intrinsically disordered region is predicted, at the end of

which is the PDPRWA element. The reliability of the prediction is color-coded; the

intrinsically disordered region is confirmed by the AIUPred algorithm (99).

1259