Lacticaseibacillus rhamnosus* P118 enhances host tolerance to *Salmonella

infection by promoting microbe-derived indole metabolites

Baikui Wang[1,#], Xianqi Peng[2,3,#], Xiao Zhou[6], Abubakar Siddique[3], Jiayun Yao[2], Haiqi

Zhang[2], Weifen Li[5], Yan Li[3,4], Min Yue[1,3,4,*]

1 Key Laboratory of Systems Health Science of Zhejiang Province, School of Life

Science, Hangzhou Institute for Advanced Study, University of Chinese Academy of

Sciences, Hangzhou 310024, China

2 Zhejiang Institute of Freshwater Fisheries, Ministry of Agriculture and Rural Affairs

Key Laboratory of Healthy Freshwater Aquaculture, Key Laboratory of Fish Health

and Nutrition of Zhejiang Province, Key Laboratory of Fishery Environment and

Aquatic Product Quality and Safety of Huzhou City, Huzhou 313001, China

3 Department of Veterinary Medicine, Zhejiang University College of Animal

Sciences, Hangzhou 310058, China

4 Hainan Institute of Zhejiang University, Sanya 572025, China

5 Institute of Animal Nutrition and Feed Sciences, Zhejiang University College of

Animal Sciences, Hangzhou 310058, China

6. Ningbo Academy of Agricultural Sciences, Ningbo 315040, China

ORCID: Min Yue, https://orcid.org/0000-0002-6787-0794; Baikui Wang,

https://orcid.org/0000-0001-5283-7337; Xianqi Peng,

https://orcid.org/0000-0001-7143-8307

* Corresponding Author: Professor Min Yue (myue@zju.edu.cn)

[#] Baikui Wang and Xianqi Peng contributed equally to this work.

Abstract

Salmonella is one of the most common foodborne pathogens, resulting in inflammatory gastroenteritis and frequently accompanied by dysbiosis. Gut commensals, such as *Lactobacillus* species, have been proven to exhibit broad anti-bacterial activities and protect hosts against pathogenic infections. Here, *Lacticaseibacillus rhamnosus* strain P118, with great probiotic properties, was screened from 290 isolates recovered from fermented yoghurts and piglet intestines using traditional and *C. elegans*-infection screening strategies. Notably, P118 and its supernatant exhibited great antibacterial activities and attenuated *C. elegans* susceptibility to *Salmonella* infection. We found that P118 protected mice against *Salmonella* lethal infections by enhancing colonization resistance, reducing pathogen invasion, alleviating intestinal pro-inflammatory response, and improving microbial dysbiosis and metabolic disorders. Microbiota and fecal metabolome analyses suggested P118 administration significantly decreased the relative abundances of harmful microbes (e.g., *Salmonella*, *Anaeroplasma*, *Klebsiella*) and increased the fecal levels of tryptophan and its derivatives (indole, indole-3-acrylic acid, 5-hydroxytryptophan, 5-methoxyindoleacetate). Deterministic processes determined the gut microbial community assembly of P118-pretreated mice. Integrated omics further demonstrated that P118 probiotic activities in enhancing host tolerance to *Salmonella* infection were mediated by microbe-derived tryptophan/indole metabolites (e.g., indole-3-acrylic acid, indole, tryptophan, 5-methoxyindoleacetic acid, and 5-hydroxytryptophan). Collective results demonstrate that *L. rhamnosus*

P118 could enhance host tolerance to *Salmonella* infections via various pathways,

including direct antibacterial actions, inhibiting *Salmonella* colonization and invasion,

attenuating pro-inflammatory responses of intestinal macrophages, and modulating

gut microbiota mediated by microbe-derived indole metabolites.

Keywords: *Lacticaseibacillus rhamnosus*; *Salmonella* infection; intestinal

dysbacteriosis; tryptophan metabolism; indole-3 acrylic acid; pro-inflammatory

response

Introduction

Diarrhoeal diseases caused by infectious agents (e.g., pathogens and parasites) remain

a severe health burden worldwide. Approximately 1.7 billion childhood diarrhoea

cases are recorded annually, with diarrhea responsible for over 480 thousand deaths

for children aged <5 years and over 500 thousand deaths for adults aged >70 years

each year (Thiagarajah et al., 2015; UNICEF, 2022; WHO, 2017). As one of the most

common foodborne enteric pathogens, *Salmonella* infection leads to substantial

gastroenteritis incidents, imposing enormous economic burdens on global society and

posing disproportionate threats to animal and human health (Kim and Isaacson, 2017;

Stanaway et al., 2019). *Salmonella* has evolved strategies to subvert colonization

resistance conferred by intestinal commensals and evade host immune defense

responses, contributing to the increasing incidence of Salmonellosis (Caballero-Flores

et al., 2022; Fu et al., 2019). *Salmonella*-contaminated food (meat, eggs, dairy) is

considered the leading cause of Salmonellosis, estimated to cause a total global

economic loss of over $3.5 billion annually in the US (Kim and Isaacson, 2017).

Traditionally, antibiotic treatment has been the primary strategy to control *Salmonella*

infectious diseases (Eng et al., 2015). However, the global abuse of antibiotics has

raised significant concerns about antimicrobial resistance (AMR), foodborne

antibiotic residues, and compromises in treating antimicrobial resistant bacterial

infections (Fishbein et al., 2023; Olesen et al., 2018). These issues have inspired

interest in seeking alternative strategies, such as dietary interventions (e.g., probiotics,

prebiotics, natural products), to prevent *Salmonella* infectious disease outbreaks.

Microbial dysbiosis induced by antimicrobials, enteric infections, and irregular

dietary consumption increases the susceptibility to pathogen infection and other

diseases, such as inflammatory bowel disease (Gillis et al., 2018a; Levy et al., 2017).

Therefore, a balanced intestinal microbiota and metabolic functions are crucial for

maintaining gastrointestinal homeostasis. Gut commensal microbes and their

derivatives have been proven to exhibit broad antimicrobial activities and protect the

host against pathogen infections by providing colonization resistance against enteric

pathogens, limiting pathogen colonization, invasion and transmission, and

maintaining intestinal barrier function (Caballero-Flores et al., 2023; Jacobson et al.,

2018; Wan et al., 2019). As one of the intestinal commensal microbes, probiotic

species (without residues in raw food products), such as *Lactobacillus*,

Bifidobacterium, *Bacillus* and yeast, exert beneficial effects on the host by enhancing

colonization resistance and immune defense, inhibiting colonization and invasion

against pathogens, and maintaining gastrointestinal homeostasis (Gareau et al., 2010;

Xiao et al., 2021). Although many candidate probiotic isolates have been uncovered

from fermented foods and mammalian intestines (Sakandar and Zhang, 2021; Wang et

al., 2022), the traditional strategies for screening these candidates are both

time-consuming and labor-intensive, involving bacterial isolation, culturing,

phenotypic characterization, randomized controlled trials, and various *in vitro* and *in*

vivo tests to assess probiotic properties (Sun et al., 2022). While culture-dependent

methods are classic strategies, emerging evaluation strategies for candidates, such as

whole-genome sequencing-based approaches and small invertebrate models (e.g., *C.*

elegans, *Drosophila*), have gained significant attention due to their high-throughput,

replicable and standardized properties (Sun et al., 2022; Wang et al., 2022).

Intestinal macrophages play vital roles in maintaining gut homeostasis, regulating

inflammation, and particularly in promoting the resolution of inflammation (Lavelle

and Sokol, 2020; Na et al., 2019), which has been considered as a novel potential

target for controlling intestinal pro-inflammatory disorders. It has been reported that

probiotics-derived metabolites, such as bacteriocins, antimicrobial compounds,

short-chain fatty acids, and tryptophan/indole derivatives, can mediate the beneficial

effects of probiotics on the host health by interacting with the host gastrointestinal

immune cells and gut residents (Britton and Faith, 2021; Sanders et al., 2019).

Bacterial tryptophan metabolism produces indole derivatives (e.g., indole-3-acetic

acid, 3-indolepropionic acid, 3-indoleacrylic acid, indole-3-lactic acid,

indole-3-aldehyde, indole-3-acetaldehyde), which are potent bioactive ligands for the

aryl hydrocarbon receptor (AHR) (Agus et al., 2018; Koh and Backhed, 2020). These

indole derivatives play crucial roles in maintaining gut barrier integrity and

homeostasis by activating AHR signaling pathway (Agus et al., 2018). *L.*

reuteri-produced indole derivatives have been reported to exert anti-inflammatory

effects by activating AHR signaling pathway (Cervantes-Barragan et al., 2017).

Although the beneficial activities of probiotics in protecting the host against diverse

enteric pathogens have been extensively examined, the interactive roles of

pro-inflammatory macrophages in microbe-tryptophan/indole metabolites against

enteric pathogen infection have not been fully studied yet. Therefore, *L. rhamnosus*

P118 with potential probiotic properties was screened from 290 recovered isolates

using traditional and *C. elegans* infection screening strategies. P118 exhibited broad

antibacterial activities and reduced host susceptibility to enteric *Salmonella* infection

by improving intestinal dysbiosis and metabolic disorders, and inhibiting intestinal

pro-inflammatory responses. Integrated omics further demonstrated that P118

probiotic activities in enhancing host tolerance to *Salmonella* infection were mediated

by microbe-derived tryptophan/indole metabolites.

Results

Two screening approaches to convergent probiotic candidates

A total of 290 bacterial strains were isolated and identified from 32 samples of the

fermented yoghurt and piglet rectal contents collected across diverse regions within

China using MRS and BHI medium, which consists of 63 *Streptococcus* strains, 158

Lactobacillus/Lacticaseibacillus/*Limosilactobacillus* strains and 69 *Enterococcus*

strains (Figure 1A & B, Table 1). Two screening strategies were employed in the present study to further investigate the potential probiotic properties of the isolates: the traditional/classic approach and the *C. elegans* infection approach. In the traditional/classic approach, 27 isolated strains were screened by milk-clotting activity assay (Figure 1A, Table S1), among which two strains (P118 and P199) exhibited the highest tolerance to bile salt (0.3%~2.0%) (Figure 1C) and biofilm formation capabilities (Figure 1D). Compared with the P199 strain, P118 strain exhibited the highest susceptibility to multiple antimicrobials (Figure S1A). *C. elegans* mimics many critical aspects of animal intestinal physiology, indicating that *C. elegans* provides an invaluable model for investigating host-microbe interactions (Kumar et al., 2020). In the *C. elegans* infection approach, 8 out of 290 isolated strains significantly increased the survivals and prolonged the life span of *S.* Typhimurium-infected *C. elegans* ($P < 0.05$), and three isolates (P118, P119, P120) treated worms were more resistant to *S*. Typhimurium infection than the other isolates ($P < 0.05$) (Figure 1E, Table S2). Integrating the results of both screening approaches, *L. rhamnosus* P118 strain exhibits versatile capabilities as a probiotic candidate (Figure 1F).

To illustrate the probiotic properties of *L. rhamnosus* P118, antibacterial activity evaluation *in vitro* was further conducted, and the results showed that the fermented supernatants of P118 significantly inhibited the growth of multiple pathogens (e.g., *Salmonella*, *Yersinia*, *S. aureus*, *E. coli*, *C. rodentium*, *P. aeruginosa*, *R. anatipestifer*) (Figure S1B). Interestingly, the inhibitory effect of P118 against *S*. Typhimurium was

in dose- and oxygen-dependent manners and results showed that high-dose and

anaerobic cultures exhibited more antibacterial effects than low-dose and aerobic

cultures, respectively ($P < 0.05$) (Figure 1G). Active factors derived from P118 to

exert antibacterial effects were explored. The results showed that the antibacterial

activities of the fermented supernatant (pH = 3.4) were sensitive to alkalinity (pH >

5.0), trypsin, proteinase-K, pepsin and high temperature (> 70°C), but were resistant

to catalase treatment (Figure 1H-K). The fermented cultures and supernatants of P118

not only significantly inhibited the growth of *S*. Typhimurium *in vitro* (Figure 1L), but

also protected worms against *S*. Typhimurium infection (Figure 1M, Table S3).

Although they exhibited no antibacterial activities *in vitro* (Figure 1L), the protective

activities of heat treated P118 and P118 lysates against *S*. Typhimurium pathogenesis

were observed in *C. elegans* but weaker than that of the fermented cultures and

supernatants (Figure 1M).

P118 genomics analysis suggests genetic determinants for antibacterial actions

Subsequently, whole genome sequencing of P118 was performed to explore the

potential antibacterial factors based on genomic evidence. The results showed that

species-level identity for P118 isolate with 2.99 Mb genome size was confirmed at *P*

< 0.0001 (Figure S2A, Figure 1N, Table S4), and P118 draft genome was most closely

related to *L. rhamnosus* subspecies (NCTC13710 and BIO5326) (Figure 1O, Table S5,

Figure S2B). Genomic analysis using antiSMASH and BAGEL4 databases revealed

the presence of putative bacteriocin synthesis genes in P118 isolate (Figure 1P-Q).

Putative bacteriocins predicated in P118 isolate showed high homology with known

class II bacteriocins (IIa carnobacteriocin B2 produced by *C. maltaromaticum*, IIb

enterocin X produced by *Enterococcus faecium* KU-B5, IId/b lactococcin A/G, and

some novel bacteriocin-encoded hypothetical proteins). Collective data demonstrates

that *L. rhamnosus* P118 exhibits outstanding probiotic traits.

P118 protects lethal *S*. Typhimurium infections in a murine model

The protective activity of *L. rhamnosus* P118 against enteric pathogens was further

examined in the *Salmonella*-infected murine model (Figure 2A). Administration of

P118 significantly protected mice against *Salmonella*-induced deaths and body weight

losses (Figure 2B-C), accompanied by the alleviated splenomegaly, hepatomegaly,

shortened gastrointestinal tract and clinical and pathological score (Figure 2D-F,

Figure S3B). Meanwhile, P118 significantly limited *Salmonella* colonization in the

intestinal tissues (duodenum and colon) and invasion into the peripheral organs (liver

and spleen) of mice, thereby markedly reducing *Salmonella* fecal shedding (Figure

2G). Additionally, *Salmonella* infection-induced ileal mucosal damage was alleviated

by P118 administration, as illustrated by the increased ileal villus height (Figure 2H,

Figure S3C), ratio of villus height/crypt-depth (Figure 2H, Figure S3D), ileal

microvilli height and density (Figure 2H, Figure S3E-F), the upregulated ZO-1

mRNA expression (Figure S4A), and the downregulated *IL-1β* mRNA expression

(Figure S4E), accompanied by the reduced IL-1β level in serum (Figure S4D).

Intervention with P118 significantly increased the numbers of Ki67-positive cells

(proliferating cells), Muc2-positive cells (goblet cells), and decreased the numbers of

F4/80-positive cells (macrophages) and $F4/80^+$ $iNOS^+$ cells (pro-inflammatory

macrophages) in the ileum (Figure 2H) and also attenuated the pathological damages

of peripheral organs (liver and spleen) caused by *Salmonella* infection (Figure S3A &

B).

P118 improves *Salmonella* infection by modulating gut microbiota

Gut microbes play crucial roles in host physiological activities, and colonization

resistance against enteric pathogens (Jacobson et al., 2018), and dysbiosis is

supposedly associated with *Salmonella* infection (Gillis et al., 2018b). Significant

bacterial community structures among four groups were observed (Figure 3A, Table

S7), and most of the bacterial operational taxonomic units (OTUs) were classified as

group-specific OTUs (Figure 3B). Compared to the uninfected group, *Salmonella*

infection significantly depleted 167 OTUs and enriched 60 OTUs (Figure 3C),

whereas compared to the *Salmonella*-infected group, pretreatment with P118

significantly depleted 16 OTUs and enriched 52 OTUs (Figure 3D). Specifically,

Salmonella infection significantly increased the relative abundances of harmful

microbes (e.g., *Salmonella*, *Escherichia*-Shigella, *Klebsiella*, *Morganella*,

Akkermansia) and significantly reduced the relative abundances of SCFA-producing

and beneficial microbes (e.g., *Alloprevotella*, *Lactococcus*, *Faecalibacterium*,

Limosilactobacillus, *Parasutterella*, *Odoribacter*, *Dubosiella*, *Candidatus arthromitus*)

(Figure 3E, Figure S5B). Pretreatment with P118 significantly reduced the relative

abundances of pathogenic microbes (e.g., *Salmonella*, *Anaeroplasma*, *Klebsiella*,

Morganella) and significantly increased the relative abundances of beneficial or

commensal microbes (e.g., *Candidatus_Saccharimonas*, *Turicibacter*, *Enterorhabdus*)

(Figure 3F, Figure S5C).

Based on the changes in bacterial community structures, the internal driving forces of

gut microbial communities were investigated using ecological models. Both null

model and neutral community model analyses revealed that the stochastic processes

belonging to homogenizing dispersal exerted important roles in *Salmonella*-infected

mice gut microbiome assembly, whereas the deterministic processes (homogeneous

selection and variable selection) exerted more influences on P118-pretreated mice gut

microbiome assembly than *Salmonella*-infected mice (Figure S6A-B).

Microbe-derived tryptophan metabolites are associated with the protection

Significant differences in fecal metabolite structures among the three groups separated

into distinct clusters were also observed (Figure 4A, Table S8). Compared with the

uninfected group, *Salmonella* infection significantly upregulated 256 fecal

metabolites ($\log_2$(fold change) > 1, $P < 0.05$) and downregulated 402 fecal

metabolites ($\log_2$(fold change) < 1, $P < 0.05$) (Figure 4B), and pretreatment with P118

significantly enriched 100 metabolites ($\log_2$(fold change) > 1, $P < 0.05$) and depleted

81 metabolites ($\log_2$(fold change) < 1, $P < 0.05$) (Figure 4C). Among the significant

differential metabolites, 91 metabolites (VIP > 1) downregulated in

Salmonella-infected mice were upregulated explicitly in P118-pretreated mice (Figure

S7A&C, Table S9), and 43 metabolites (VIP > 1) upregulated in *Salmonella*-infected

mice were downregulated explicitly in P118-pretreated mice (Figure S7B&C, Table

S9). KEGG pathway analysis revealed that fecal metabolites were mainly enriched in

tryptophan metabolism, fatty acid biosynthesis, biosynthesis of amino acids and

neuroactive ligand-receptor interaction pathways in P118-pretreated mice (Figure 4D).

Interestingly, pretreatment with P118 significantly increased the fecal levels of

tryptophan and tryptophan derivatives (indole, indole-3-acrylic acid,

5-hydroxytryptophan, 5-methoxyindoleacetate) (Figure 4E). Pearson correlation

analysis showed that *S*. Typhimurium burdens (in the duodenum, colon, liver, spleen,

feces) and organ (spleen, liver) indices were negatively correlated with the levels of

fecal tryptophan and tryptophan derivatives (indole, indole-3-acrylic acid,

5-hydroxytryptophan, protocatechuic acid, trans-cinnamaldehyde,

5-methoxyindole-3-carbaldehyde) and positively correlated with the levels of

(serotonin, tryptophanol, 3-(2-hydroxyethyl) indole, tryptamine) (Figure 4F).

Conversely, the body weight was positively correlated with the levels of fecal

tryptophan and tryptophan derivatives (Figure 4F). Bio-Sankey network analysis

further showed that 26 genera were identified as potential bacteria that might

participate in the metabolic reaction R02340 (tryptophan synthase, indoleglycerol

phosphate and indole), of which five bacteria (*Bifidobacterium*, *Lacticaseibacillus*,

Enterococcus, *Staphylococcus*, and *Pseudomonas*, dark red if $P < 0.05$) were

positively associated with indole in the metabolic reaction R02340 (Figure 4G). These

results indicate that microbe-derived tryptophan metabolites are associated with

P118-mediated protection against *Salmonella* infection.

Based on the findings above that *Lacticaseibacillus* species were involved in

tryptophan biosynthesis and metabolism, *L. rhamnosus* P118 genome was reanalyzed

to determine whether P118 encode enzymes necessary to exert this function. The

genomic data further showed that P118 encoded various genes (e.g., *fldH*, *AraT*,

AspB-4, *AmiE*, *trpA*, *trpB*, *BCAT*, *IGPS*, *MT*) essential to biosynthesize and

metabolize tryptophan into tryptophan derivatives (Figure 4H, Table S10). The

metabolomic results of P118 cultures also validated the genomic data that P118 could

secrete a diverse profile of tryptophan-derived metabolites (e.g indole-3-acrylic acid,

indole, indole-3-lactic acid, DL-tryptophan, kynurenine, N-Acetyl-D-tryptophan,

5-methoxyindoleacetic acid, 5-hydroxytryptophan) (Table S11). Taken together, these

data suggest that P118-derived tryptophan metabolites might contribute to the

protection against *Salmonella* infection.

Indole-3-acrylic acid protects against *Salmonella* infection by inhibiting

macrophage pro-inflammatory responses

Based on the above results, exogenous indole-3-acrylic acid (IAA) was employed to

further explore its roles against *Salmonella* infection. The antibacterial assay results

showed that IAA (> 4.4 mM) significantly inhibited *Salmonella* growth (Figure S8).

Exogenous IAA administration significantly reduced mice susceptibility to

Salmonella infection，as evidenced by the increased body weight (Figure 5B), the

alleviated splenomegaly and hepatomegaly (Figure 5C, 5D, 5G), and the increased

colon length (Figure 5E). Meanwhile, IAA administration significantly inhibited

Salmonella colonization in the intestinal tissues (cecum and colon) and invasion into

the peripheral organs (liver and spleen) of mice, thereby markedly reducing

Salmonella fecal shedding (Figure 5F). Additionally, *Salmonella* infection-induced

ileal mucosa damage was alleviated by exogenous IAA, as illustrated by the increased

ileal villus height (Figure 5H, 5O), and the increased numbers of Muc2-positive

goblet cells in the ileum (Figure 5I, 5O). Exogenous IAA intervention significantly

reduced *Salmonella* infection-induced intestinal pro-inflammatory responses, as

evidenced by the decreased numbers of F4/80-positive macrophages (Figure 5O), the

downregulated mRNA (*IL-1β*, *IL-6*, *IL-18*, *TNF-α*, *iNOS*) expression (Figure 5J-N),

and the reduced protein (iNOS, IL-1β, IL-6, TNF-α, markers of pro-inflammatory

macrophages) expression (Figure 5O).

The *in vitro* experiments further showed that, although failed to increase the uptake of

Salmonella (Figure S9A, 0h), IAA treatment significantly reduced intracellular

Salmonella survival in RAW 264.7 macrophage cells (Figure S9A, 6h, 12h), which

could be reversed by pretreatment with AHR inhibitor CH-223191 (Figure S9A, 6h,

12h), indicating that IAA enhances the intracellular bactericidal capacity of RAW

264.7 cells. Furthermore, IAA treatment significantly inhibited *Salmonella*

infection-induced pro-inflammatory responses, as evidenced by inhibiting mRNA

(*IL-1β*, *IL-6*, *IL-18*, *TNF-α*, *iNOS*) expression and nitric oxide secretion of

Salmonella-infected RAW 264.7 macrophage cells, which blocked by the presence of

AHR inhibitor CH-223191 (Figure S9B-G).

Macrophages play important roles in initiating immune responses, maintaining gut

homeostasis and phagocytic clearance of pathogens (Na et al., 2019). To further

investigate the involved roles of macrophages in P118/IAA-mediated protective effect,

in vivo macrophage depletion using clodronate liposomes was conducted. As expected,

macrophage depletion significantly blocked the protection of P118 and exogenous

IAA against *Salmonella* infection (Figure 6A-O). P118 or exogenous IAA

administration failed to increase the body weight (Figure 6B) and colon length (Figure

6E), alleviate the splenomegaly and hepatomegaly (Figure 6C, 6D, 6F), inhibit the

colonization, invasion and shedding of *Salmonella* (Figure 6G) after the intestinal

macrophage depletion. Additionally, after depleting the intestinal macrophage, P118

or exogenous IAA administration also failed to attenuate *Salmonella*

infection-induced intestinal mucosa damage (Figure 6H, 6I, 6O) and

pro-inflammatory responses (Figure 6J-N, 6O). Taken together, these findings indicate

that P118 and IAA protect against *Salmonella* infection by inhibiting

pro-inflammatory responses of intestinal macrophage.

Discussion

Fermented foods and mammalian intestines are rich in lactic acid bacteria that can

produce organic acids and lactobacillin to inhibit overgrowth, adhesion, invasion and

toxin secretion of pathogens (Gareau et al., 2010; Wan et al., 2019). Traditionally, the

culturomics strategy, together with multiple evaluation assays, is the primary way to

search potential probiotic species from the complex microbial community, which is

both time- and labor-consuming (Wang et al., 2022). Although the whole genomic

sequencing-based approach is an emerging favorable way to screen potential probiotic

strains, it's hard to widely adopt based on numerous isolates due to the high

sequencing costs (Schwarze et al., 2020). Given that mimics many critical aspects of

animal intestinal physiology, *C. elegans* provides an invaluable model to investigate

host-microbe interactions and probiotic properties of beneficial microbes at low cost

(Kumar et al., 2020; Poupet et al., 2020). In the present study, we found *Salmonella*

infection significantly induced death and decreased the lifespan of *C. elegans*,

indicating *C. elegans* is very susceptible to *S*. Typhimurium infection, consistent with

a previous report (Rangan et al., 2016). Integrating the results of culturomics and *C.*

elegans infection strategies, *L. rhamnosus* P118 with excellent probiotic properties

was screened from 290 isolates and exhibited broad *in vitro* antibacterial activities.

Interestingly, high fermented dose and anaerobic cultures of P118 showed more

antibacterial effects than low dose and aerobic cultures, indicating that the active

factors were more released by P118 at the stationary phase under anaerobic conditions.

What's more, the antibacterial activities of the fermented supernatant were sensitive to

alkalinity, trypsin, proteinase-K, pepsin, and high temperature, indicating that the

active secreted factors in the fermented supernatant were protein- or peptide-based

active agents, consistent with a previous report that *L. rhamnosus* exerted antibacterial

effects by secreting bioactive peptides (Iram et al., 2022). Although limited studies

have reported bacteriocin-like activity in *L. rhamnosus*, it remains unclear whether

this species produces active bacteriocins (Kankainen et al., 2009). Nonetheless,

genomic data in this study revealed the presence of putative bacteriocin synthesis

genes in P118 genome. In agreement with a previous study (Kankainen et al., 2009),

the predicated bacteriocins showed high homology with known class II bacteriocins

(e.g., IIa carnobacteriocin B2 produced by *C. maltaromaticum*, IIb enterocin X

produced by *Enterococcus faecium* KU-B5), which needs to be further validated by

gene-editing technology and synthetic biology.

To further validate the antibacterial activities of *L. rhamnosus* P118 against enteric

pathogens, *Salmonella*-infected mice model was employed. We found that

administration of P118 significantly reduced mice susceptibility to *S*. Typhimurium

infection by improving intestinal health, dysbiosis and metabolic dysfunction.

Colonized *S*. Typhimurium in the intestine disrupts structural and functional intestinal

integrity, leading to substantial immune response and severe epithelial barrier damage,

and then migrates to internal organs (liver, spleen) via the blood circulatory system,

causing splenomegaly and hepatomegaly (Rogers et al., 2022; Wang et al., 2021). In

the present study, P118 significantly increased *Salmonella*-infected mice survival and

alleviated *Salmonella*-induced splenomegaly and hepatomegaly. The beneficial

attenuated effect was partly contributed by the reduced *Salmonella* colonization in

intestine and invasion into the peripheral organs, and eventually, the decreased fecal

shedding, consistent with previous reports (Liu et al., 2023; Wu et al., 2024). It's

reported that *S*. Typhimurium exploits and disrupts intestinal barrier integrity to

adhere, and disseminate deeper into intestinal mucosa, and evade immune clearance

(Bernal-Bayard and Ramos-Morales, 2018; Hiyoshi et al., 2022). In the current study,

P118 treatment also significantly improved the structural and functional intestinal

integrity, attenuated inflammatory responses, reduced the numbers of F4/80-positive

macrophages and proinflammatory macrophages (F4/80$^+$ iNOS$^+$ cells), and increased

the numbers of Ki67-positive and Muc2-positive goblet cell in the ileum,

demonstrating that P118 improved the intestinal health and reducing

pro-inflammatory responses of intestinal macrophages, which further confirms the

beneficial alleviated effect exerted by *L. rhamnosus* P118.

Intestinal microbes play essential roles in providing colonization resistance against

enteric pathogens and pathobionts, and limiting adhesion, invasion and transmission

of pathogens (Caballero-Flores et al., 2022; Jacobson et al., 2018). Microbial

dysbiosis and excessive immune responses induced by enteric pathogens can, in turn,

aggravate and exacerbate gut permeability and damage (Read et al., 2021; Rogers et

al., 2022). Many studies have reported that *Salmonella* infection was often

accompanied by microbial dysbiosis (Gillis et al., 2018a; Read et al., 2021; Wang et

al., 2021). The current results found that *L. rhamnosus* P118 treatment significantly

restored the bacterial community structures in *Salmonella*-infected mice, and

significantly reduced the relative abundances of harmful microbes (e.g., *Salmonella*,

Klebsiella, *Anaeroplasma*, *Morganella*), which belong to pathogens and pathobionts

(Dong et al., 2022; Galan, 2021; Laupland et al., 2022). Structure composition and

assembly of microbial communities are essential for ecosystem function (Shi et al.,

2023; Xun et al., 2019), and revealing the underlying mechanisms of microbial

community assembly is a major goal of microbial community ecology (Jiao et al.,

2020; Stegen et al., 2012). Generally, stochastic (e.g., dispersal events, ecological

drift, random birth, death) and deterministic (e.g., interspecies interactions (e.g.,

competition, facilitation, mutualisms and predation), species traits, environmental

factors) processes are two major ecological processes that drive microbiome assembly

(Stegen et al., 2012; Zhou and Ning, 2017). Deterministic processes mainly involve

non-random and niche-based mechanisms with abiotic and biotic factors that

influence microbial community assembly (Vellend, 2010), whereas stochastic

processes mainly reflect random changes in the relative abundance of species (Jiao et

al., 2020; Jiao et al., 2021). Although important in shaping the diversities of microbial

composition and functions, the relative contribution of microbiome assembly

processes varies with different habitats (Jiao et al., 2020; Jiao et al., 2021). The

present study revealed that the stochastic processes played dominant roles in intestinal

microbiome assembly in *Salmonella*-infected mice, indicating random changes

occurred in the relative abundance of species. Conversely, the deterministic processes

played more significant roles in driving intestinal microbial community assembly in

P118-pretreated mice than in *Salmonella*-infected mice, indicating abiotic factors (e.g.,

dietary intervention) or biotic interactions (e.g., microbial competition, facilitation,

mutualism, predation, host filtering) might influence the microbial community.

Accumulating evidence revealed that gut microbes-derived metabolites (e.g.,

short-chain fatty acids, tryptophan/indole-derived metabolites, bacteriocins, bile acid,

natural products) are crucial mediators in host physiological activities (Cani, 2019;

Wlodarska et al., 2017), and metabolic disorders are major risk factors for bacterial

infections (Olive and Sassetti, 2016). In this study, the results showed that *L.*

rhamnosus P118 improved *Salmonella* infection-induced fecal metabolic disorders.

Interestingly, P118 significantly increased the fecal levels of tryptophan and its

derivatives (indole, indole-3-acrylic acid, 5-hydroxytryptophan), and those

metabolites were negatively correlated with *S.* Typhimurium burdens (in the

duodenum, colon, liver, spleen, feces) and organ (spleen, liver) indices, and positively

correlated with the body weight of mice, indicating that microbiota-derived

tryptophan/indole metabolites play beneficial roles in P118 mediated probiotic

activities in enhancing host tolerance to *Salmonella* infection. It's reported that

through binding to the aryl hydrocarbon receptor (AHR) of host, bacterial tryptophan

derivatives act as triggers to stimulate immune responses and gut hormones to exert

anti-inflammatory activities, enhance intestinal epithelial barrier, and promote

gastrointestinal motility, which contributes to host gastrointestinal homeostasis and

health (Agus et al., 2018; Cani, 2019). Our genomic and metabolomic data further

revealed that P118 could produce a wide array of tryptophan-derived metabolites

owing to encoding a variety of enzyme genes necessary to metabolize tryptophan into

indole derivatives; consistently, some *Lactobacillus* species broadly conserved

enzyme encoding genes that are involved in tryptophan metabolism (Montgomery et

al., 2022). The above results indicate that *L. rhamnosus* P118- and microbe-derived

tryptophan/indole metabolites might play positive roles in P118-mediated probiotic

activities. Although significant increases in microbe-derived tryptophan/indole

metabolites were observed in P118-treated mice, and it was established that P118 can

metabolize tryptophan into indole derivatives, it remains to be investigated whether

the differential tryptophan/indole metabolites were directly derived from P118 or

from other microbes. To test the roles of differential tryptophan/indole metabolites in

P118-mediated beneficial effects, exogenous indole-3-acrylic acid, which

significantly and negatively correlated with *S.* Typhimurium burdens and

hepatosplenomegaly mentioned above, was employed to investigate its protective

effect against *Salmonella* infection. Interestingly, we found that indole-3-acrylic acid

administration significantly enhanced mice tolerance to *S.* Typhimurium infection by

improving intestinal health and attenuating pro-inflammatory responses of intestinal

macrophages. Furthermore, the *in vitro* results also provide that indole-3-acrylic acid

enhanced bactericidal capacity of RAW 264.7 cells and inhibited *S.*

Typhimurium-induced pro-inflammatory responses, consistent with a previous report

(Wlodarska et al., 2017). Taken together, these results demonstrate that *L. rhamnosus*

P118/microbe-derived indole-3-acrylic acid play beneficial roles in P118-mediated

probiotic activities.

Macrophages are important in initiating immune responses, maintaining intestinal

immune homeostasis, phagocytic clearance of pathogens, tissue repair, and host

defense (Na et al., 2019; Neurath, 2024). It's reported that macrophages with strong

plasticity could be polarized into pro-inflammatory (M1) phenotype or proresolving

(M2) phenotype (Na et al., 2019). M1 macrophage activation is crucial and necessary

for pathogen clearance (Na et al., 2019). However, exaggerated immune responses

triggered by M1 macrophage are detrimental to the host by inducing tissue damage

and cell deaths (Wang et al., 2020). The current study showed that both *L. rhamnosus*

P118   and   indole-3-acrylic   acid   significantly   attenuated   *S.*   Typhimurium

infection-induced pro-inflammatory responses, indicating that pro-inflammatory

macrophages might be key targets for *L. rhamnosus* P118 exerting probiotic effect.

This hypothesis was confirmed by macrophage depletion experiments that the

protective effects of *L. rhamnosus* P118 and indole-3-acrylic acid against *Salmonella*

infection were abrogated after intestinal macrophage depletion. Taken together, these

results indicate that *L. rhamnosus* P118/microbe-derived indole metabolites enhance

host tolerance to *Salmonella* infection by reducing intestinal pro-inflammatory

responses, which also provides a potential alternative strategy for using probiotics and

their derived metabolites to treat intestinal inflammatory disorders such as

inflammatory bowel disease (Lavelle and Sokol, 2020; Neurath, 2024).

Conclusion

In summary, two distinctive approaches were used to screen the probiotic candidate,

and the P118 strain was underscored for superficial effects on a murine infection

model. Further, bacterial genomic sequencing, gut microbiota and metabolomic

analysis pinpointed the microbe-derived tryptophan/indole could be the importance of

P118 probiotic properties. Nevertheless, the newly found P118 could enhance host

tolerance to *Salmonella* infections via various pathways, including direct antibacterial

actions, inhibiting *Salmonella* colonization and invasion, attenuating

pro-inflammatory responses of intestinal macrophages, and modulating gut

microbiota mediated by microbe-derived indole metabolites (Figure 7).

Materials and methods

Bacterial isolation and culture

Lactic acid bacteria (LAB) and *Enterococcus* strains were isolated from the fermented yoghurts collected from families in multiple cities of China and the intestinal contents from healthy piglets without pathogen infection and diarrhoea by our lab using de Man-Rogosa-Sharpe (MRS) with 2.0% $CaCO_3$ and Brain heart infusion (BHI) broths (Huankai Microbial, Guangzhou, China). Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry (MALDI-TOF MS) method was employed to identify of bacterial species (He et al., 2022). *Lacticaseibacillus rhamnosus* P118 (P118) and *Salmonella enterica* serovar Typhimurium SL1344 (STm) were cultured in MRS and Luria Broth (LB) medium at 37 °C overnight under aerobic conditions, separately. The final concentration of bacterial strains was constantly checked by the spreading plate method (Wang et al., 2023). After being centrifugated at 4000× g for 15 min at 4 °C, the fermented supernatant of *L. rhamnosus* P118 was collected by centrifugation at 4000× g for 15 min at 4 °C, and then filtered through a 0.22 μm membrane (Merck Millipore, Burlington, MA, USA) kept at 4 °C for further use.

Milk-clotting activity (MCA) analysis

For MCA, overnight fermented lactic acid bacteria strains under logarithmic growth were added to test tubes containing skim milk (10% skim milk with 10 mM $CaCl_2$) at 1:20 ratio (v/v) and then incubated at 37 °C. The coagulating time, final pH value and acidity of milk were measured during the curdling process.

Bile salt tolerance assay

507 The bile tolerance assay was conducted according to a previous report (Wang et al.,

508 2023). Briefly, the isolated strains under logarithmic growth were resuspended into

509 fresh MRS medium (1% v/v) containing different (wt/v, 0, 0.3%, 0.6%, 1.0%, 1.5%,

510 2.0%) porcine bile (MACKLIN®, Shanghai, China) and then plated into 96-well

511 plates, after incubation for 24h at 37 °C, the absorbance at 620 nm was measured by a

512 SpectraMax M5 reader (Molecular Devices, Sunnyvale, CA, USA).

513 **Biofilm forming assay**

514 Biofilm-forming assay of the isolated strains was performed as described previously

515 (Zhou et al., 2023). Briefly, 200 μL overnight cultures (adjusted to 2×10^6 CFU/mL)

516 were added into 96-well flat-bottom plates. After incubation for 36h at 37 °C, the

517 planktonic cells were discarded. After being gently washed three times with sterile

518 PBS, the adherent cells were stained with 200 μL of 0.4% crystal violet for 15 min at

519 room temperature and then washed with sterile PBS three times again. Finally, 200 μL

520 75% ethanol was added to dissolve the stained adherent cells for 15 min, and biofilm

521 was quantified by measuring the absorbance at 590 nm using a SpectraMax M5

522 reader.

523 **Antimicrobial susceptibility test**

524 Twelve antimicrobial agents (penicillins (PG), erythromycin (EM), lincomycin (LM),

525 gentamicin (GM), doxycycline (DOX), ceftriaxone (CTR), norfloxacin (NOR),

526 azithromycin (AZM), fleroxacin (FLE), vancomycin (VA), streptomycin (STR)) were

527 employed to evaluate the antimicrobial susceptibility of the candidate probiotics by

528 utilizing the Kirby-Bauer disk diffusion method (Hudzicki, 2009).

***Caenorhabditis elegans* infection**

C. elegans infection model was performed as previously described (Rangan et al.,

2016). *C. elegans* N2 (Bristol) were routinely raised on Nematode Growth Media

(NGM) feeding with *E. coli* OP50, and L4-stage worms were prepared according to

the description in WormBook (Girard et al., 2007). 200 μL of the isolated strains

(adjusted to 5×10^8 CFU/mL using M9 medium) and *L. rhamnosus* P118-related

substances (e.g., mixed cultures, fermented sterile supernatants, P118 lysate, dead

P118) were poured onto NGM in 24-well plates, separately, and dried at 22 °C for 4 h.

L4-stage worms were then transferred to the lawns of the poured NGM plates (30

worms/plate) with three plates per group for 24h. Then, the treated worms were

washed and transferred to lawns of STm (1×10^8 CFU/lawn) on NGM plates for

infection. Worm survival was monitored at 24-hour intervals for sixteen days.

Antibacterial assay

The agar diffusion method (Wang et al., 2023) was used to evaluate the antimicrobial

activity against multiple pathogens *in vitro*. Briefly, bacterial culture medium (LB,

BHI) containing 0.25% (v/v) different zoonotic pathogens under logarithmic growth

was poured into 10-cm plates (Corning, NY, USA), separately. Then, 200 μL of the

fermented sterile supernatant collected from the overnight cultured *L. rhamnosus*

P118 was added into 8 mm agar wells created by Oxford cup. The bacteriostatic effect

was evaluated by the growth inhibition zone around the 8-mm agar wells after

overnight incubation at 37 °C.

Whole genome sequencing and data analysis

Whole genome sequencing, quality control, assembly and annotation of *L. rhamnosus* P118 was conducted according to our previous study (Zhou et al., 2023) using PacBio Sequel platform and Illumina NovaSeq PE150 at Novogene (Novogene Co, Ltd., Beijing, China), and the P118 draft genome is available in the Sequence Read Archive under accession number PRJNA848987. *L. rhamnosus* P118 assembly was queried against the NCBI non-redundant prokaryotic genomes database using the Microbial Genomes Atlas (MIGA) webserver (database update to 10/10/2023) (Rodriguez et al., 2018). Taxonomic classification was inferred by the maximum average amino acid identity (AAI) against all genomes in the database, with p-values estimated from the distribution of all the reference genomes in NCBI's RefSeq at each taxonomic level as a readout of classification probability. Average nucleotide identity (ANI) and AAI tables of maximally the top-50 reference hits in the database were extracted and graphed as x-y scatter plots to determine the nearest phylogenetic neighbors. To generate phylogenic trees of isolates and MIGA-identified nearest phylogenetic neighbors, the interfered proteome from PROKKA isolate annotations, or as publicly available in NCBI for applicable reference genomes were uploaded to the ANI/AAI-Matrix calculator (Rodriguez-R and Konstantinidis, 2016). The resulting phylogenetic tree based on AAI was visualized using the interactive tree of life (iTOL) (Letunic and Bork, 2021).

L. rhamnosus P118 tryptophan-associated enzymes were identified as previously described (including TNA, TMO, TrpD, ArAT, ALD, IPDC, FldH, and AmiE) (Montgomery et al., 2022). Briefly, enzyme commission numbers, where available, or

alternatively enzyme names, were queried in PATRIC, PATRIC Global Family (PGF)

cross-genus identifiers were extracted and compiled for query within the *L.*

rhamnosus P118 genome. Protein sequences of *L. rhamnosus* P118 identified

enzymes were analyzed using InterProScan for additional functional prediction, and

sequence homology to the previously experimentally validated enzymes within other

bacterial species was determined using Blastp at 30% cross-genus identify and 90%

identity within each species.

***S.* Typhimurium murine infection model**

Animal assays, specifically eighty 6-week-old female C57BL/6 mice (Slac Animal

Inc., Shanghai, China), were conducted in the Laboratory Animal Center of Zhejiang

University under light-controlled (12h light/dark cycle), temperature-controlled (22 ±

2°C) and humidity-controlled (55 ± 5°C) conditions. Forty mice were randomly

selected for survival evaluation under *S.* Typhimurium infection, and the rest were

used for sample collection on day 3 post-infection (Figure 2A). The mice were

randomly divided into four groups (n = 10/group): Control group (C), *L. rhamnosus*

P118 group (P), *S.* Typhimurium-infected group (S), and *L. rhamnosus* P118

protective group (P+S). Mice in the C and S groups were drinking sterile water every

day, respectively. Mice in the P and P+ST groups were orally administered with *L.*

rhamnosus P118 (1×10^8 CFU/mouse) daily for seven days. After that, mice were

orally ingested with *S.* Typhimurium SL1344 (1×10^8 CFU/mouse) or sterile PBS. All

mice were allowed free access to water and food, and weighed every day. Mice were

euthanized on day 3 post-infection. Spleen and liver were weighted for organ index

calculation according to the formula: organ index = organ weight (g) / body weight (g)

*1000.

To evaluate the protective effect of indole-3-acrylic acid (IAA), eighteen mice were

randomly divided into three groups (n = 6/group): Control group (C), *S*.

Typhimurium-infected group (S), and indole-3-acrylic acid protective group (IAA+S).

Mice in the C and S groups were drinking sterile water every day, respectively. IAA

(Sigma, I3807) was dissolved in 1 M NaOH in PBS and adjusted pH to 7.4 with 1 M

HCl. Mice in the IAA+ST groups were orally administered with 75mg/kg IAA daily

for seven days. After that, mice were orally ingested with *S*. Typhimurium SL1344 or

sterile PBS. At 3 day-post-infection, mice were euthanized for sample collection.

***In vivo* macrophage depletion**

To evaluate the involved roles of macrophages, macrophage depletion assay was

conducted through intraperitoneal injection with 250 μL/mouse clodronate liposomes

(CLP, From Vrije Universiteit Amsterdam) twice (2 days prior and 1 day after

Salmonella infection, separately) according to the manufacturer's instructions.

Thirty-three mice were randomly divided into six groups (n = 3~6/group): Control

group (C), *S*. Typhimurium-infected group (S), Control group with macrophage

depletion (CLP+C), *S*. Typhimurium-infected group with macrophage depletion

(CLP+S), IAA protective group with macrophage depletion (CLP+IAA+S), and P118

protective group with macrophage depletion (CLP+P+S). At 3 day-post-infection,

mice were euthanized for sample collection.

Macrophage cell culture

Murine macrophage cell line RAW 264.7 was cultured in DMEM/F12 medium

(Gibco, Carlsbad, CA) supplemented with 10% FBS (Gibco, CA) and 1% (v/v)

antibiotic solution (100 μg/mL streptomycin + 100 U/mL penicillin, Sigma-Aldrich,

St. Louis, MO) at 37°C in a 5% humidified CO_2 incubator. If not mentioned, the

antibiotic solution (100 μg/mL streptomycin and 100 U/mL penicillin) was not added

into DMEM/F12 medium in the further experiment.

For *S*. Typhimurium killing assay, RAW 264.7 cells seeded into 12-well plates (2 ×

10^6 cells/well) were treated with 100 μM IAA for 6h, followed by infecting with *S*.

Typhimurium (MOI = 10) for 1h. After washing 3 times with sterile PBS, the infected

RAW 264.7 cells were incubated for 0h, 6h and 12h in DMEM/F12 medium

containing gentamicin (50 μg/mL). At each time point, the infected RAW 264.7 cells

were washed with sterile PBS for 4 times and lysed with 0.01% Triton X-100 diluted

in PBS. The serial 10-fold dilutions of cell lysates were immediately plated on LB

agar plates to determine bacterial viability.

For the immune response assay, RAW 264.7 cells seeded into 12-well plates (2 × 10^6

cells/well) were pretreated with CH-223191 (AHR inhibitor, 10 μM) for 6h,

subsequently treating cells with 100 μM IAA for another 6h. After 6h, RAW 264.7

cells were infected with *S*. Typhimurium (MOI = 10) for 2h. After washing 3 times

with sterile PBS, the infected RAW 264.7 cells were collected using RNAiso Plus

(TaKaRa, Dalian, China) for total RNA extracting and quantitative real-time PCR

(qPCR). The primer sets are listed in Table S6.

S. Typhimurium burden

S. Typhimurium loads in tissues (duodenum, cecum, colon, liver and spleen) and

shedding in feces were determined as previously described (Xu et al., 2018). Briefly,

tissues and fecal samples were collected under sterile conditions and homogenized in

sterile PBS containing 0.1% Triton X-100. Serial 10-fold dilutions of tissue

homogenates were spread onto SS agar plates in triplicate and then incubated at 37°C

overnight to quantify the bacterial colony-forming unit (CFU).

Histopathology and immunofluorescent analysis

Tissues (ileum, liver and spleen) fixed in 4% paraformaldehyde were embedded in

paraffin, sliced, dehydrated and then sectioned for hematoxylin and eosin (H&E)

staining. The tissue slices were imaged and analyzed by using Olympus Microscope

(Tokyo, Japan). For the immunofluorescent assay, the paraffin-embedded ileum

samples were incubated with primary antibodies against Ki67 (GB121141, Servicebio,

China), lysozyme (GB11345, Servicebio, China), Muc2 (GB11344, Servicebio,

China), F4/80 (GB113373, Servicebio, China), iNOS (GB11119, Servicebio, China),

IL-1β (GB11113, Servicebio, China), IL-6 (GB11117, Servicebio, China), TNF-α

(GB11188, Servicebio, China) for overnight at 4°C and then further incubated with

secondary antibodies (Alexa Fluor® 488-conjugated goat anti-rabbit/mouse IgG) or

HRP-conjugated goat anti-rabbit secondary antibody (GB23303, Servicebio, China)

for 60 min at room temperature. Finally, ileum slices were digitalized by Pannoramic

MIDI (3DHISTECH, Hungary).

Transmission electron microscope (TEM) observation

TEM observations of ileum tissues were prepared according to our previous study

(Peng et al., 2022). Briefly, after being fixed with 2.5% buffered glutaraldehyde,

ileum segments were washed three times by cold 100-mM phosphate buffers, and

then post-fixed in 0.1% osmium tetroxide for 2 h, rapidly dehydrated in ascending

grades of ethanol (30, 50, 70, 95, and 100%), and moved into a 1:1 mixture of

propylene oxide and epoxy araldite. Finally, the ileum samples were observed and

captured by the transmission electron microscope (JEOL, Tokyo, Japan).

ELISA assay

The contents of colonic interleukin (IL)-1β and IL-18 were determined by ELISA kits

(Solarbio, Beijing, China) according to the manufacturer's instructions.

qPCR

Total RNA extracted from intestinal tissues by RNAiso Plus kit (TAKARA, Dalian,

China) was reverse-transcripted using PrimeScript II 1st Strand cDNA Synthesis Kit

(TAKARA, Dalian, China) according to the manufacturer's instructions. The qPCR

was then performed on the StepOne real-time PCR system (Applied Biosystems)

using SYBR PremixEx TaqII (TAKARA). The primer sets are listed in Table S6. Fold

changes were calculated after normalizing to the housekeeping gene β-actin using the

$2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

Microbiota analysis

The TIANamp Stool DNA Kit (Tiangen, Beijing, China) was employed to extract

fecal bacterial genomic DNA, and fecal bacterial communities were investigated

using 16S rDNA sequencing by targeting the V3-V4 hypervariable region. The 16S

rDNA sequencing was then performed on an Illumina NovaSeq platform (Illumina

Inc., San Diego, CA, USA). The quality filter of the paired-end raw sequences and a

cluster of the filtered sequences into the operational taxonomic unit (OTU) at 97%

similarity was performed by QIIME software (version 1.9.1). Microbial OTU

representative sequences were assigned to a taxonomic lineage by the RDP classifier

based on the SILVA database (v 132 release).

To investigate the dissimilarities in bacterial communities and fecal metabolomes,

principal coordinates analysis (PCoA), analysis of similarities (ANOSIM),

permutational multivariate analysis of variance (PERMANOVA) and multi-response

permutation procedure (MRPP) were calculated using "vegan" package and

visualized using "ggplot2" package. Difference analysis of bacterial communities

based on OUT levels among groups was calculated using "DESeq2" package and was

envisioned by "UpSetR" and "ggplot2" packages. The significant differences in

bacterial taxonomies were analyzed and visualized by statistical analysis of taxonomic

and functional profiles (STAMP) software with a two-sided Welch's t-test (Parks et al.,

2014).

The sloan neutral model was employed to estimate the importance of neutral

processes in the assembly of bacterial communities using "MicEco" package (Sloan et

al., 2006). A null-model-based statistical framework was conducted to evaluate the

relative importance of determinism and stochasticity in bacterial community assembly

(Stegen et al., 2013; Xu et al., 2022). Two assembly processes were defined as

deterministic processes ($|\beta NTI|>2$) and stochastic processes ($|\beta NTI|\leq2$) based on the

absolute values of beta Nearest Taxon Index (βNTI) (Xu et al., 2020). Additionally,

five ecological assembly processes were then interpreted as homogeneous selection

(βNTI<-2), homogenizing dispersal (|βNTI|$\leq$2 and RC_{bray}<-0.95), undominated

(|βNTI|$\leq$2 and |RC_{bray}|<0.95), dispersal limitation (|βNTI|$\leq$2 and RC_{bray}>0.95) and

variable (heterogeneous) selection (βNTI>2) based on the threshold of the absolute

values of βNTI and Bray-Curtis-based Raup-Crick (RC_{bray}) (Jiao et al., 2020; Xu et al.,

2022).

Metabolomics analysis

Untargeted metabolomics was investigated to analyze the fecal metabolomes of mice.

After extracting metabolites from feces, UHPLC-MS/MS analyses of samples were

performed by Vanquish UHPLC systems (Thermo Fisher, Germany) coupled with

Orbitrap Q Exactive™ HF-X mass spectrometers (Thermo Fisher, Germany) in

Novogene Co., Ltd. (Beijing, China). Compound Discoverer 3.1 (CD3.1, Thermo

Fisher) was employed to conduct peak alignment, and quantify metabolites generated

by UHPLC-MS/MS. After normalizing to total spectral intensity and predicting

molecular formula, the peaks were queried against mzCloud, mzVault and MassList

databases to extract the quantitative results. Annotation of fecal metabolite signatures

was queried against the KEGG database, HMDB database and LIPIDMaps database.

Significant differences in fecal metabolites were analyzed using univariate analysis

(t-test) and partial least-squares discriminant analysis (PLS-DA), in which fecal

metabolites with variable importance in the projection (VIP) > 1 and $P < 0.05$ were

considered differential metabolites. Pearson correlation analysis between *S.*

Typhimurium burden (in tissues and feces) and fecal metabolites was analyzed and

visualized using "linkET" package (Huang, 2021). MetOrigin platform was employed

to analyze the sources of microbial metabolites and their metabolic functions in fecal

metabolomes (Yu et al., 2022).

Statistical Analysis

The significance of the results was analyzed by one-way analysis of variance

(ANOVA) with Tukey's multiple comparisons test using SPSS v24 (SPSS Inc.,

Chicago, IL, USA) and statistical graphs were visualized by GraphPad Prism v8.0

(GraphPad Software, CA) and package "ggplot2" of R software (v4.3.1). $* P < 0.05$,

$** P < 0.01$, $*** P < 0.001$.

Acknowledgements

This work was supported by the National Program on Key Research Project of China

(2022YFC2604201) as well as the European Union's Horizon 2020 Research and

Innovation Programme under Grant Agreement No. 861917-SAFFI, Zhejiang

Provincial Key R&D Program of China (2023C03045, 2022C02024), Key Research

and Development Program of Hangzhou (202203A08), Zhejiang Provincial Natural

Science Foundation of China (LQ23C180005).

Conflict of interest statement

No potential conflict of interest was reported by the author(s).

Data availability statement

16S rRNA sequencing data is deposited at the Genome Sequence Archive (GSA) of

the BIG Data Center (https://bigd.big.ac.cn/gsa/) under accession no. PRJCA021751.

L. rhamnosus P118 isolate draft genome assembly is publicly available in the NCBI

under accession no. <u>PRJNA848987</u>.

Ethics statement

752   Animal procedures followed the guidelines and were approved by the Institutional

753   Animal Care and Use Committee of Zhejiang University (Permission number:

754   ZJU20220295).

Author contributions

756   Baikui Wang and Xianqi Peng: Investigation, Validation, Methodology, Data

757   Analysis, Visualization, Formal analysis, Writing - original draft & review & editing;

758   Xiao Zhou, Jiayun Yao, Haiqi Zhang, Weifen Li, Yan Li: Investigation, Writing –

759   review; Min Yue: Conceptualization, Resources, Methodology, Supervision, Writing

760   – review & editing, Funding acquisition, Project administration.

References

Agus A, Planchais J, Sokol H. Gut microbiota regulation of tryptophan metabolism in
health and disease. Cell Host Microbe 2018;23:716-24.

Bernal-Bayard J, Ramos-Morales F. Molecular mechanisms used by salmonella to
evade the immune system. Curr Issues Mol Biol 2018;25:133-68.

Britton GJ, Faith JJ. Causative microbes in host-microbiome interactions. Annu Rev
Microbiol 2021;75:223-42.

Caballero-Flores G, Pickard JM, Nunez G. Microbiota-mediated colonization
resistance: Mechanisms and regulation. Nat Rev Microbiol 2022.

Caballero-Flores G, Pickard JM, Nunez G. Microbiota-mediated colonization
resistance: Mechanisms and regulation. Nat Rev Microbiol 2023;21:347-60.

Cani PD. Microbiota and metabolites in metabolic diseases. Nat Rev Endocrinol
2019;15:69-70.

Cervantes-Barragan L, Chai JN, Tianero MD, Di Luccia B, Ahern PP, Merriman J,
Cortez VS, Caparon MG, Donia MS, Gilfillan S, Cella M, Gordon JI, Hsieh
CS, Colonna M. Lactobacillus reuteri induces gut intraepithelial
cd4(+)cd8alphaalpha(+) t cells. Science 2017;357:806-10.

Dong N, Yang X, Chan EW, Zhang R, Chen S. Klebsiella species: Taxonomy,
hypervirulence and multidrug resistance. EBioMedicine 2022;79:103998.

Eng SK, Pusparajah P, Ab Mutalib NS, Ser HL, Chan KG, Lee LH. Salmonella: A
review on pathogenesis, epidemiology and antibiotic resistance. Frontiers in
Life Science 2015;8:284-93.

Fishbein SRS, Mahmud B, Dantas G. Antibiotic perturbations to the gut microbiome.
Nat Rev Microbiol 2023;21:772-88.

Fu A, Mo Q, Wu Y, Wang B, Liu R, Tang L, Zeng Z, Zhang X, Li W. Protective effect
of bacillus amyloliquefaciens against salmonella via polarizing macrophages
to m1 phenotype directly and to m2 depended on microbiota. Food & function
2019;10:7653-66.

Galan JE. Salmonella typhimurium and inflammation: A pathogen-centric affair. Nat
Rev Microbiol 2021;19:716-25.

Gareau MG, Sherman PM, Walker WA. Probiotics and the gut microbiota in intestinal
health and disease. Nat Rev Gastroenterol Hepatol 2010;7:503-14.

Gillis CC, Hughes ER, Spiga L, Winter MG, Zhu W, Furtado De Carvalho T, Chanin
RB, Behrendt CL, Hooper LV, Santos RL, Winter SE. Dysbiosis-associated
change in host metabolism generates lactate to support salmonella growth.
Cell Host Microbe 2018a;23:570.

Gillis CC, Hughes ER, Spiga L, Winter MG, Zhu W, Furtado De Carvalho T, Chanin
RB, Behrendt CL, Hooper LV, Santos RL, Winter SE. Dysbiosis-associated
change in host metabolism generates lactate to support salmonella growth.
Cell Host Microbe 2018b;23:54-64 e6.

Girard LR, Fiedler TJ, Harris TW, Carvalho F, Antoshechkin I, Han M, Sternberg PW,
Stein LD, Chalfie M. Wormbook: The online review of caenorhabditis elegans
biology. Nucleic Acids Res 2007;35:D472-5.

He D, Zeng W, Wang Y, Xing Y, Xiong K, Su N, Zhang C, Lu Y, Xing X. Isolation

and characterization of novel peptides from fermented products of

lactobacillus for ulcerative colitis prevention and treatment. Food Science and

Human Wellness 2022;11:1464-74.

Hiyoshi H, English BC, Diaz-Ochoa VE, Wangdi T, Zhang LF, Sakaguchi M, Haneda

T, Tsolis RM, Baumler AJ. Virulence factors perforate the pathogen-containing

vacuole to signal efferocytosis. Cell Host Microbe 2022;30:163-70 e6.

Huang H. Linket: Everything is linkable. R package version 0.0.3 2021.

Hudzicki J. Kirby-bauer disk diffusion susceptibility test protocol. American society

for microbiology 2009;15:55-63.

Iram D, Kindarle UA, Sansi MS, Meena S, Puniya AK, Vij S. Peptidomics-based

identification of an antimicrobial peptide derived from goat milk fermented by

lactobacillus rhamnosus (c25). J Food Biochem 2022;46:e14450.

Jacobson A, Lam L, Rajendram M, Tamburini F, Honeycutt J, Pham T, Van Treuren W,

Pruss K, Stabler SR, Lugo K, Bouley DM, Vilches-Moure JG, Smith M,

Sonnenburg JL, Bhatt AS, Huang KC, Monack D. A gut commensal-produced

metabolite mediates colonization resistance to salmonella infection. Cell Host

Microbe 2018;24:296-307 e7.

Jiao S, Yang YF, Xu YQ, Zhang J, Lu YH. Balance between community assembly

processes mediates species coexistence in agricultural soil microbiomes across

eastern china. Isme Journal 2020;14:202-16.

Jiao S, Zhang B, Zhang G, Chen W, Wei G. Stochastic community assembly

decreases soil fungal richness in arid ecosystems. Mol Ecol 2021;30:4338-48.

Kankainen M, Paulin L, Tynkkynen S, Von Ossowski I, Reunanen J, Partanen P,

Satokari R, Vesterlund S, Hendrickx AP, Lebeer S, De Keersmaecker SC,

Vanderleyden J, Hamalainen T, Laukkanen S, Salovuori N, Ritari J, Alatalo E,

Korpela R, Mattila-Sandholm T, Lassig A, Hatakka K, Kinnunen KT,

Karjalainen H, Saxelin M, Laakso K, Surakka A, Palva A, Salusjarvi T,

Auvinen P, De Vos WM. Comparative genomic analysis of lactobacillus

rhamnosus gg reveals pili containing a human- mucus binding protein. Proc

Natl Acad Sci U S A 2009;106:17193-8.

Kim HB, Isaacson RE. Salmonella in swine: Microbiota interactions. Annual Review

of Animal Biosciences, Vol 5 2017;5:43-63.

Koh A, Backhed F. From association to causality: The role of the gut microbiota and

its functional products on host metabolism. Mol Cell 2020;78:584-96.

Kumar A, Baruah A, Tomioka M, Iino Y, Kalita MC, Khan M. Caenorhabditis elegans:

A model to understand host-microbe interactions. Cell Mol Life Sci

2020;77:1229-49.

Laupland KB, Paterson DL, Edwards F, Stewart AG, Harris PNA. Morganella

morganii, an emerging cause of bloodstream infections. Microbiol Spectr

2022;10:e0056922.

Lavelle A, Sokol H. Gut microbiota-derived metabolites as key actors in

inflammatory bowel disease. Nature Reviews Gastroenterology & Hepatology

2020;17:223-37.

Letunic I, Bork P. Interactive tree of life (itol) v5: An online tool for phylogenetic tree

display and annotation. Nucleic acids research 2021;49:W293-W6.

Levy M, Kolodziejczyk AA, Thaiss CA, Elinav E. Dysbiosis and the immune system.

Nat Rev Immunol 2017;17:219-32.

Liu J, Liu H, Liu H, Teng Y, Qin N, Ren X, Xia X. Live and pasteurized akkermansia

muciniphila decrease susceptibility to salmonella typhimurium infection in

mice. J Adv Res 2023;52:89-102.

Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time

quantitative pcr and the 2(t)(-delta delta c) method. Methods 2001;25:402-8.

Montgomery TL, Eckstrom K, Lile KH, Caldwell S, Heney ER, Lahue KG,

D'alessandro A, Wargo MJ, Krementsov DN. Lactobacillus reuteri tryptophan

metabolism promotes host susceptibility to cns autoimmunity. Microbiome

2022;10:198.

Na YR, Stakenborg M, Seok SH, Matteoli G. Macrophages in intestinal inflammation

and resolution: A potential therapeutic target in ibd. Nat Rev Gastroenterol

Hepatol 2019;16:531-43.

Neurath MF. Strategies for targeting cytokines in inflammatory bowel disease. Nat Rev Immunol 2024.

Olesen SW, Barnett ML, Macfadden DR, Brownstein JS, Hernandez-Diaz S, Lipsitch M, Grad YH. The distribution of antibiotic use and its association with antibiotic resistance. Elife 2018;7.

Olive AJ, Sassetti CM. Metabolic crosstalk between host and pathogen: Sensing, adapting and competing. Nature Reviews Microbiology 2016;14:221-34.

Parks DH, Tyson GW, Hugenholtz P, Beiko RG. Stamp: Statistical analysis of taxonomic and functional profiles. Bioinformatics 2014;30:3123-4.

Peng X, Ed-Dra A, Song Y, Elbediwi M, Nambiar RB, Zhou X, Yue M. Lacticaseibacillus rhamnosus alleviates intestinal inflammation and promotes microbiota-mediated protection against salmonella fatal infections. Front Immunol 2022;13:973224.

Poupet C, Chassard C, Nivoliez A, Bornes S. Caenorhabditis elegans, a host to investigate the probiotic properties of beneficial microorganisms. Front Nutr 2020;7:135.

Rangan KJ, Pedicord VA, Wang YC, Kim B, Lu Y, Shaham S, Mucida D, Hang HC. A secreted bacterial peptidoglycan hydrolase enhances tolerance to enteric pathogens. Science 2016;353:1434-7.

Read E, Curtis MA, Neves JF. The role of oral bacteria in inflammatory bowel disease. Nat Rev Gastroenterol Hepatol 2021;18:731-42.

Rodriguez-R LM, Konstantinidis KT, The enveomics collection: A toolbox for specialized analyses of microbial genomes and metagenomes. PeerJ Preprints; 2016.p.

Rodriguez RL, Gunturu S, Harvey WT, Rossello-Mora R, Tiedje JM, Cole JR, Konstantinidis KT. The microbial genomes atlas (miga) webserver: Taxonomic and gene diversity analysis of archaea and bacteria at the whole genome level. Nucleic Acids Res 2018;46:W282-W8.

Rogers AP, Mileto SJ, Lyras D. Impact of enteric bacterial infections at and beyond the epithelial barrier. Nature Reviews Microbiology 2022.

894   Sakandar HA, Zhang HP. Trends in probiotic(s)-fermented milks and their in vivo

895   functionality: A review. Trends in Food Science & Technology

896   2021;110:55-65.

897   Sanders ME, Merenstein DJ, Reid G, Gibson GR, Rastall RA. Probiotics and

898   prebiotics in intestinal health and disease: From biology to the clinic. Nat Rev

899   Gastroenterol Hepatol 2019;16:605-16.

900   Schwarze K, Buchanan J, Fermont JM, Dreau H, Tilley MW, Taylor JM, Antoniou P,

901   Knight SJL, Camps C, Pentony MM, Kvikstad EM, Harris S, Popitsch N,

902   Pagnamenta AT, Schuh A, Taylor JC, Wordsworth S. The complete costs of

903   genome sequencing: A microcosting study in cancer and rare diseases from a

904   single center in the united kingdom. Genet Med 2020;22:85-94.

905   Shi J, Yang L, Liao Y, Li J, Jiao S, Shangguan Z, Deng L. Soil labile organic carbon

906   fractions mediate microbial community assembly processes during long-term

907   vegetation succession in a semiarid region. iMeta 2023:e142.

908   Sloan WT, Lunn M, Woodcock S, Head IM, Nee S, Curtis TP. Quantifying the roles of

909   immigration and chance in shaping prokaryote community structure. Environ

910   Microbiol 2006;8:732-40.

911   Stanaway JD, Parisi A, Sarkar K, Blacker BF, Reiner RC, Hay SI, Nixon MR,

912   Dolecek C, James SL, Mokdad AH, Abebe G, Ahmadian E, Alahdab F,

913   Alemnew BT, Alipour V, Bakeshei FA, Animut MD, Ansari F, Arabloo J,

914   Asfaw ET, Bagherzadeh M, Bassat Q, Belayneh YM, Carvalho F, Daryani A,

915   Demeke FM, Demis AB, Dubey M, Duken EE, Dunachie SJ, Eftekhari A,

916   Fernandes E, Fard RF, Gedefaw GA, Geta B, Gibney KB, Hasanzadeh A,

917   Hoang CL, Kasaeian A, Khater A, Kidanemariam ZT, Lakew AM,

918   Malekzadeh R, Melese A, Mengistu DT, Mestrovic T, Miazgowski B,

919   Mohammad KA, Mohammadian M, Mohammadian-Hafshejani A, Nguyen T,

920   Nguyen H, Nguyen H, Nirayo YL, Olagunju AT, Olagunju TO, Pourjafar H,

921   Qorbani M, Rabiee M, Rabiee N, Rafay A, Rezapour A, Samy AM, Sepanlou

922   SG, Shaikh MA, Sharif M, Shigematsu M, Tessema B, Tran BX, Ullah I,

923   Yimer EM, Zaidi Z, Murray CJL, Crump JA, Salm GN-T. The global burden

of non-typhoidal salmonella invasive disease: A systematic analysis for the global burden of disease study 2017. Lancet Infectious Diseases 2019;19:1312-24.

Stegen JC, Lin X, Konopka AE, Fredrickson JK. Stochastic and deterministic assembly processes in subsurface microbial communities. ISME J 2012;6:1653-64.

Stegen JC, Lin XJ, Fredrickson JK, Chen XY, Kennedy DW, Murray CJ, Rockhold ML, Konopka A. Quantifying community assembly processes and identifying features that impose them. Isme Journal 2013;7:2069-79.

Sun Y, Li HC, Zheng L, Li JZ, Hong Y, Liang PF, Kwok LY, Zuo YC, Zhang WY, Zhang HP. Iprobiotics: A machine learning platform for rapid identification of probiotic properties from whole-genome primary sequences. Briefings in Bioinformatics 2022;23.

Thiagarajah JR, Donowitz M, Verkman AS. Secretory diarrhoea: Mechanisms and emerging therapies. Nat Rev Gastroenterol Hepatol 2015;12:446-57.

Unicef, Diarrhoea remains a leading killer of young children, despite the availability of a simple treatment solution. U Data (Ed.). https://data.unicef.org/topic/child-health/diarrhoeal-disease/; 2022.p.

Vellend M. Conceptual synthesis in community ecology. Q Rev Biol 2010;85:183-206.

Wan MLY, Forsythe SJ, El-Nezami H. Probiotics interaction with foodborne pathogens: A potential alternative to antibiotics and future challenges. Critical reviews in food science and nutrition 2019;59:3320-33.

Wang B, Hussain A, Zhou Y, Zeng Z, Wang Q, Zou P, Gong L, Zhao P, Li W. Saccharomyces boulardii attenuates inflammatory response induced by clostridium perfringens via tlr4/tlr15-myd8 pathway in hd11 avian macrophages. Poult Sci 2020;99:5356-65.

Wang B, Ye X, Zhou Y, Zhao P, Mao Y. Glycyrrhizin attenuates salmonella typhimurium-induced tissue injury, inflammatory response, and intestinal dysbiosis in c57bl/6 mice. Front Vet Sci 2021;8:648698.

Wang B, Zhou Y, Wang Q, Xu S, Wang F, Yue M, Zeng Z, Li W. Lactiplantibacillus
plantarum lac16 attenuates enterohemorrhagic escherichia coli o157:H7
infection by inhibiting virulence traits and improving intestinal epithelial
barrier function. Cells 2023;12.

Wang L, Liu Q, Chen Y, Zheng X, Wang C, Qi Y, Dong Y, Xiao Y, Chen C, Chen T,
Huang Q, Zhai Z, Long C, Yang H, Li J, Wang L, Zhang G, Liao P, Liu YX,
Huang P, Huang J, Wang Q, Chu H, Yin J, Yin Y. Antioxidant potential of
pediococcus pentosaceus strains from the sow milk bacterial collection in
weaned piglets. Microbiome 2022;10:83.

Who, Diarrhoeal disease. 2017.p.

Wlodarska M, Luo C, Kolde R, D'hennezel E, Annand JW, Heim CE, Krastel P,
Schmitt EK, Omar AS, Creasey EA, Garner AL, Mohammadi S, O'connell DJ,
Abubucker S, Arthur TD, Franzosa EA, Huttenhower C, Murphy LO, Haiser
HJ, Vlamakis H, Porter JA, Xavier RJ. Indoleacrylic acid produced by
commensal peptostreptococcus species suppresses inflammation. Cell Host
Microbe 2017;22:25-37 e6.

Wu Y, Zhang X, Liu X, Li Y, Han D, Pi Y, Whitmore MA, Lu X, Zhang G, Zheng J,
Wang J. Strain specificity of lactobacilli with promoted colonization by
galactooligosaccharides administration in protecting intestinal barriers during
salmonella infection. J Adv Res 2024;56:1-14.

Xiao Y, Zhai Q, Zhang H, Chen W, Hill C. Gut colonization mechanisms of
lactobacillus and bifidobacterium: An argument for personalized designs.
Annu Rev Food Sci Technol 2021;12:213-33.

Xu Q, Ling N, Chen H, Duan Y, Wang S, Shen Q, Vandenkoornhuyse P. Long-term
chemical-only fertilization induces a diversity decline and deep selection on
the soil bacteria. mSystems 2020;5.

Xu Q, Vandenkoornhuyse P, Li L, Guo J, Zhu C, Guo S, Ling N, Shen Q. Microbial
generalists and specialists differently contribute to the community diversity in
farmland soils. J Adv Res 2022;40:17-27.

Xu XG, Gong L, Wang BK, Wu YP, Wang Y, Mei XQ, Xu H, Tang L, Liu RR, Zeng

ZH, Mao YL, Li WF. Glycyrrhizin attenuates salmonella enterica serovar
typhimurium infection: New insights into its protective mechanism. Frontiers
in Immunology 2018;9:2321.

Xun W, Li W, Xiong W, Ren Y, Liu Y, Miao Y, Xu Z, Zhang N, Shen Q, Zhang R.
Diversity-triggered deterministic bacterial assembly constrains community
functions. Nat Commun 2019;10:3833.

Yu G, Xu C, Zhang D, Ju F, Ni Y. Metorigin: Discriminating the origins of microbial
metabolites for integrative analysis of the gut microbiome and metabolome.
IMeta 2022;1:e10.

Zhou J, Ning D. Stochastic community assembly: Does it matter in microbial ecology?
Microbiology and Molecular Biology Reviews 2017;81:e00002-17.

Zhou X, Kang X, Chen J, Song Y, Jia C, Teng L, Tang Y, Jiang Z, Peng X, Tao X.
Genome degradation promotes salmonella pathoadaptation by remodeling
fimbriae-mediated proinflammatory response. National Science Review
2023:nwad228.

Figures

Figure 1. Isolation and antibacterial characterization of *L. rhamnosus* P118. (**A**) Screening flowchart of P118 *in vitro*. (**B**) Candidate probiotic strains. (**C**) Bile salt tolerance ability of strains. (**D**) Biofilm forming ability of strains. (**E**) Screening using

C. elegans infection model. (**F**) Interacted screening strategy. (**G**) Antibacterial ability

of P118 under aerobic or anaerobic culture conditions. (**H**) Broad-spectrum pH

tolerance of P118 supernatants (pH = 3.4) that adjusted to pH < 3.4 by 1M HCl or pH >

3.4 by 1M NaOH. Antibacterial effects of P118 supernatant under (**I**) 20 mg/mL

catalase (CAT), (**J**) 100μg/mL proteinases (trypsin, proteinase K, pepsin), and (**K**)

different temperatures (37, 50, 70, 90, 100, 120°C) boiled for 30 min treatments. (**L**)

Antibacterial effects of components of P118 (boiled at 120 °C for 30 min or was lysed

by ultrasonication at 240 W for 2 h). (**M**) Active ingredients of P118 protect *C.*

elegans against *S*. Typhimurium infection. (**N**) Taxonomic classification of P118 draft

genome. (**O**) The nearest subspecies phylogenetic neighbor of P118 draft genome was

determined by percentage shared genomic content graphed as ANI versus AAI. (**J, H**)

Prediction of secondary metabolites and bacteriocin protein-encoding gene clusters of

P118. (**G-M**) *S*. Typhimurium SL1344 was selected as an indicator pathogen.

Significant differences * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure 2. *L. rhamnosus* P118 enhances mice tolerance to *S*. Typhimurium infection.

(**A**) Experimental design. (**B**) Survival curve of mice infected with *S*. Typhimurium.

(**C**) Body weight. (**D**) Spleen index. (**E**) Representative images of spleen and intestine.

(**F**) Liver index. (**G**) *S*. Typhimurium burden in tissues and shedding in feces. (**H**)

Representative images of H&E staining, TEM, and immunostaining (DAPI, Ki67,

lysozyme, Muc2, F4/80, iNOS) in the ileum. Different lowercase letters indicate a

significant difference ($P < 0.05$). Significant differences * $P < 0.05$, ** $P < 0.01$.

1027

Figure 3. *L. rhamnosus* P118 improves *S*. Typhimurium infection-induced

dysbacteriosis. (**A**) Principal coordinates analysis (PCoA) based on Bray-Curtis

distance. (**B**) UpSetR plot based on bacterial absolute OTU abundances. (**C & D**) The

fold changes of bacterial absolute OTU abundances between two groups. (**E & F**)

Comparison of intestinal microbes by STAMP. The prefix "g_" represent the

annotated level of the genus.

1034

Figure 4. Microbe derived tryptophan metabolites are involved in mice tolerance to *S.*

Typhimurium. (**A**) Principal coordinates analysis (PCoA) based on Bray-Curtis

distance. (**B & C**) UpSetR plot based on fecal metabolites. (**D**) Metabolomics

pathway enrichment. (**E**) Comparison of tryptophan metabolism enriched pathway. (**F**)

Figure 5. Indole-3-acrylic acid enhances mice tolerance to *S*. Typhimurium infection.

(**A**) Experimental design. (**B**) Body weight change. (**C**) Spleen index. (**D**) Liver index.

(**E**) Colon length. (**F**) *S*. Typhimurium burden in tissues and shedding in feces. (**G**)

Representative images of spleen and liver. (**H**) Villus height of ileum. (**I**)

Muc2-positive cells in ileum. (**J-N**) mRNA expression levels in ileum. (**O**)

Representative images of H&E staining and immunostaining in the ileum. Different

1054

Figure 6. Macrophage depletion abrogates the protective effect of *L. rhamnosus* P118

and indole-3-acrylic acid against *S.* Typhimurium infection. (**A**) Experimental design.

(**B**) Body weight change. (**C**) Spleen index. (**D**) Liver index. (**E**) Colon length. (**F**)

Representative images of spleen and liver. (**G**) *S.* Typhimurium burden in tissues and

shedding in feces. (**H**) Villus height of ileum. (**I**) Muc2-positive cells in ileum. (**J-N**)

mRNA expression levels in ileum. (**O**) Representative images of H&E staining and

immunostaining in the ileum. Different lowercase letters indicate a significant

difference ($P < 0.05$). Significant differences ** $P < 0.01$.

1065

1066

1067

Figure 7. *Lacticaseibacillus rhamnosus* P118 strain with great probiotic properties was screened from 290 purified isolates through two distinctive screen approaches, and P118 strain was underscored for superficial protective effects on a murine infection model. Further, multi-omics analysis pinpointed the microbe-derived tryptophan/indole could be the importance of P118 probiotic properties. Nevertheless, the newly found P118 enhances host tolerance to *Salmonella* infections via various pathways, including direct antibacterial actions, inhibiting *Salmonella* colonization and invasion, attenuating pro-inflammatory responses of intestinal macrophages, and modulating gut microbiota mediated by microbe-derived indole metabolites.

1077

1078

1079 **Table 1** A list of probiotic strains recovered from examined samples

Source	Name	Size of strains
Yogurt	*Streptococcus salivarius ssp. thermophilus*	63
	Lacticaseibacillus paracasei	18
	Lactobacillus acidophilus	17
	Lactobacillus zeae	10
	Lacticaseibacillus rhamnosus	4
	Lacticaseibacillus casei	4
	Lactobacillus delbrueckii	6
piglet intestine	*Limosilactobacillus mucosae*	16
	Lactobacillus johnsonii	38
	Limosilactobacillus reuteri	16
	Limosilactobacillus vaginalis	11
	Lactobacillus crispatus	9
	Lactobacillus salivarius	6
	Lactobacillus amylovorus	3
	Enterococcus faecium	38
	Enterococcus faecalis	31
Total		290

1080 **Supplementary information**

1081

Figure S1 (**A**) Drug resistance and (**B**) broad-spectrum antibacterial activities of

candidate probiotic strains.

1084

Figure S2. (**A**) Circular genome map of *L. rhamnosus* P118. (**B**) phylogenetic trees of

L. rhamnosus P118 and nearest phylogenetic neighbors constructed by average amino

acid identify (AAI). Color gradients indicate the percentage of conserved average

nucleotide identity (ANI) between each isolate and respective nearest subspecies

phylogenetic neighbors.

Figure S3. Clinical and Histopathological observation. (**A**) Representative images of H&E staining tissues (liver, spleen). Scale bar with 100 μm. (**B**) Clinical symptom score. Histopathological quantification of H&E staining (**C & D**) and TEM (**E & F**) images observed in the ileum. Different lowercase letters indicate a significant difference ($P < 0.05$).

Figure S4. Gene and protein expression level of tight junction protein (ZO-1, Claudin-1) and inflammatory cytokines (IL-1β, IL-18) in ileum and serum of mice. Different lowercase letters indicate a significant difference ($P < 0.05$).

1100

Figure S5. Comparison of intestinal microbes by STAMP. The prefixes "g_" and "s_"

represent the annotated levels of genus and species.

1103

Figure S6. *L. rhamnosus* P118 alters gut microbial community assembly. (**A**)

Ecological processes of gut microbes. The inner circle indicates the percentage of

deterministic and stochastic assembly processes. The outer circle indicates the

contribution of detailed ecological processes. (**B**) The fit of neutral community model

(NCM) of community assembly for gut microbes. The best fit to NCM and 95% CIs

around the best fit to NCM were indicated by solid and dashed blue lines, respectively.

Light blue and dark red dots indicate OTUs that occur more (above prediction) or less

(below prediction) frequently than predicted by NCM, respectively, while black dots

indicate the OTUs that occur within neutral prediction (neutral distribution). Donut

charts shows the distinct fitting proportions of OTUs predicted by NCM.

1114

1115

Figure S7. Analysis of differential fecal metabolites. Venn diagram based on differential fecal metabolites (**A**) down-regulated in "S *vs* C" and up-regulated in "P+S *vs* S", (**B**) up-regulated in "S *vs* C" and down-regulated in "P+S *vs* S". (**C**)

1119　Significant differential fecal metabolites in "S *vs* C" and "P+S *vs* S".

1120

1121　**Figure S8.** Antibacterial activities of indole-3-acrylic acid.

1122

1123　**Figure S9.** Indole-3-acrylic acid enhances bactericidal capacity and exerts

1124   anti-inflammatory activity in RAW 264.7 cells. Different lowercase letters indicate a

1125   significant difference ($P < 0.05$).

1126

Table S1 Milk-clotting activity of lactic acid bacteria

Table S2 Significance between life span data was calculated using a Mantel-Cox

log-rank test

Table S3 Significance between life span data was calculated using a Mantel-Cox

log-rank test

Table S4 Species-level identity of *L. rhamnosus* P118. The draft genome of P118 was

queried using the NCBI non-redundant prokaryotic database available through MIGA

(http://www.microbial-genomes.org). P-values reflect confidence in the assignment as

each taxonomic rank.

Table S5 *L. rhamnosus* P118 draft genome nearest subspecies phylogenetic neighbors.

Average nucleotide identity (ANI) and average amino acid identity (AAI) using both

percent identity and a fraction of genome shared as determined through MIGA

(http://www.microbial-genomes.org) query of the NCBI non-redundant prokaryotic

genomes database were considered for ranking likelihood of nearest subspecies

phylogenetic neighbors for duplicate isolates of each *Lacticaseibacillus* species.

Table S6 Primers used for the quantitative PCR

Table S7 ANOSIM, PERMANOVA and MRPP analysis of microbial diversity among

treatments

Table S8 ANOSIM, PERMANOVA and MRPP analysis of fecal metabolites among

groups

Table S9 Significant differential fecal metabolites in "S vs C" and "P+S vs S"

Table S10 Prediction of aminotransferase class and tryptophan binding affinity in *L.*

rhamnosus P118

Table S11 Tryptophan metabolism and indole derivatives in *L. rhamnosus* P118

cultures