***Lacticaseibacillus rhamnosus* P118 enhances host tolerance to *Salmonella* infection**

by promoting microbe-derived indole metabolites

Baikui Wang[1,#], Xianqi Peng[2,3,#], Xiao Zhou[6], Xiuyan Jin[3], Abubakar Siddique[3], Jiayun

Yao[2], Haiqi Zhang[2], Weifen Li[5], Yan Li[3,4], Min Yue[1,3,4,*]

1 Key Laboratory of Systems Health Science of Zhejiang Province, School of Life

Science, Hangzhou Institute for Advanced Study, University of Chinese Academy of

Sciences, Hangzhou 310024, China

2 Zhejiang Institute of Freshwater Fisheries, Ministry of Agriculture and Rural Affairs

Key Laboratory of Healthy Freshwater Aquaculture, Key Laboratory of Fish Health

and Nutrition of Zhejiang Province, Key Laboratory of Fishery Environment and

Aquatic Product Quality and Safety of Huzhou City, Huzhou 313001, China

3 Department of Veterinary Medicine, Zhejiang University College of Animal Sciences,

Hangzhou 310058, China

4 Hainan Institute of Zhejiang University, Sanya 572025, China

5 Institute of Animal Nutrition and Feed Sciences, Zhejiang University College of

Animal Sciences, Hangzhou 310058, China

6. Ningbo Academy of Agricultural Sciences, Ningbo 315040, China

ORCID: Min Yue, https://orcid.org/0000-0002-6787-0794; Baikui Wang,

https://orcid.org/0000-0001-5283-7337; Xianqi Peng, https://orcid.org/0000-0001-

7143-8307

* Corresponding Author: Professor Min Yue (myue@zju.edu.cn)

[#] Baikui Wang and Xianqi Peng contributed equally to this work.

Abstract

Salmonella is one of the most common foodborne pathogens, resulting in inflammatory gastroenteritis and frequently accompanied by dysbiosis. Gut commensals, such as *Lactobacillus* species, have been proven to exhibit broad anti-bacterial activities and protect hosts against pathogenic infections. Here, *Lacticaseibacillus rhamnosus* strain P118, screened from 290 isolates recovered from fermented yoghurts and healthy piglet intestines using traditional and *Caenorhabditis elegans*-infection screening strategies, exerts great probiotic properties. Notably, P118 and its supernatant exhibited great antibacterial activities and attenuated *C. elegans* susceptibility to *Salmonella* infection. We found that P118 protected mice against *Salmonella* lethal infections by enhancing colonization resistance, reducing pathogen invasion, alleviating intestinal pro-inflammatory response, and improving microbial dysbiosis and fecal metabolite changes. Microbiota and fecal metabolome analyses suggested P118 administration significantly decreased the relative abundances of potentially harmful microbes (e.g., *Salmonella*, *Anaeroplasma*, *Klebsiella*) and increased the fecal levels of tryptophan and its derivatives (indole, indole-3-acrylic acid, 5-hydroxytryptophan, 5-methoxyindoleacetate). Deterministic processes determined the gut microbial community assembly of P118-pretreated mice. Integrated omics further demonstrated that P118 probiotic activities in enhancing host tolerance to *Salmonella* infection were mediated by microbe-derived tryptophan/indole metabolites (e.g., indole-3-acrylic acid, indole, tryptophan, 5-methoxyindoleacetic acid, and 5-hydroxytryptophan). Collective results demonstrate that *L. rhamnosus* P118 could enhance host tolerance to *Salmonella*

infections via various pathways, including direct antibacterial actions, inhibiting

Salmonella colonization and invasion, attenuating pro-inflammatory responses of

intestinal macrophages, and modulating gut microbiota mediated by microbe-derived

indole metabolites.

Keywords: *Lacticaseibacillus rhamnosus*; *Salmonella* infection; intestinal

dysbacteriosis; tryptophan metabolism; indole-3 acrylic acid; pro-inflammatory

response

Introduction

Diarrhoeal diseases caused by infectious agents (e.g., pathogens and parasites) remain

a severe health burden worldwide. Approximately 1.7 billion childhood diarrhoea cases

are recorded annually, with diarrhea responsible for over 480 thousand deaths for

children aged <5 years and over 500 thousand deaths for adults aged >70 years each

year (Thiagarajah et al., 2015; UNICEF, 2022; WHO, 2017). As one of the most

common foodborne enteric pathogens, *Salmonella* infection leads to substantial

gastroenteritis incidents, imposing enormous economic burdens on global society and

posing disproportionate threats to animal and human health (Kim and Isaacson, 2017;

Stanaway et al., 2019). *Salmonella* has evolved strategies to subvert colonization

resistance conferred by intestinal commensals and evade host immune defense

responses, contributing to the increasing incidence of Salmonellosis (Caballero-Flores

et al., 2022; Fu et al., 2019). *Salmonella*-contaminated food (meat, eggs, dairy) is

considered the leading cause of Salmonellosis, estimated to cause a total global

economic loss of over \$3.5 billion annually in the US (Kim and Isaacson, 2017).

Traditionally, antibiotic treatment has been the primary strategy to control *Salmonella*

infectious diseases (Eng et al., 2015). However, the global abuse of antibiotics has

raised significant concerns about antimicrobial resistance (AMR), foodborne antibiotic

residues, and compromises in treating antimicrobial resistant bacterial infections

(Fishbein et al., 2023; Olesen et al., 2018). These issues have inspired interest in seeking

alternative strategies, such as dietary interventions (e.g., probiotics, prebiotics, natural

products), to prevent *Salmonella* infectious disease outbreaks.

Microbial dysbiosis induced by antimicrobials, enteric infections, and irregular dietary

habits increases the susceptibility to pathogen infection and other diseases, such as

inflammatory bowel disease (Gillis et al., 2018b; Levy et al., 2017). Therefore, a

balanced intestinal microbiota and metabolic functions are crucial for maintaining

gastrointestinal homeostasis. Gut commensal microbes and their derivatives have been

proven to exhibit broad antimicrobial activities and protect the host against pathogen

infections by providing colonization resistance against enteric pathogens, limiting

pathogen colonization, invasion and transmission, and maintaining intestinal barrier

function (Caballero-Flores et al., 2023; Jacobson et al., 2018; Wan et al., 2019). As one

of the intestinal commensal microbes, probiotic species (without residues in raw food

products), such as *Lactobacillus*, *Bifidobacterium*, *Bacillus* and yeast (e.g.,

Saccharomyces boulardii, *S. cerevisiae*), exert beneficial effects on the host by

enhancing colonization resistance and immune defense, inhibiting colonization and

invasion against pathogens, and maintaining gastrointestinal homeostasis (Gareau et al.,

2010; Xiao et al., 2021). Although many candidate probiotic isolates have been

uncovered from fermented foods and mammalian intestines (Sakandar and Zhang, 2021;

Wang et al., 2022), the traditional strategies for screening these candidates are both

time-consuming and labor-intensive, involving bacterial isolation, culturing,

phenotypic characterization, randomized controlled trials, and various *in vitro* and *in*

vivo tests to assess probiotic properties (Sun et al., 2022). While culture-dependent

methods are classic strategies, emerging evaluation strategies for candidates, such as

whole-genome sequencing-based approaches and small invertebrate models (e.g., *C.*

elegans, *Drosophila*), have gained significant attention due to their high-throughput,

replicable and standardized properties (Sun et al., 2022; Wang et al., 2022).

Intestinal macrophages play vital roles in maintaining gut homeostasis, regulating

inflammation, and particularly in promoting the resolution of inflammation (Lavelle

and Sokol, 2020; Na et al., 2019), which has been considered as a novel potential target

for controlling intestinal pro-inflammatory disorders. It has been reported that

probiotics-derived metabolites, such as bacteriocins, antimicrobial compounds, short-

chain fatty acids, and tryptophan/indole derivatives, can mediate the beneficial effects

of probiotics on the host health by interacting with the host gastrointestinal immune

cells and gut residents (Britton and Faith, 2021; Sanders et al., 2019). Bacterial

tryptophan metabolism produces indole derivatives (e.g., indole-3-acetic acid, 3-

indolepropionic acid, 3-indoleacrylic acid, indole-3-lactic acid, indole-3-aldehyde,

indole-3-acetaldehyde), which are potent bioactive ligands for the aryl hydrocarbon

receptor (AHR) (Agus et al., 2018; Koh and Backhed, 2020). These indole derivatives

play crucial roles in maintaining gut barrier integrity and homeostasis by activating

AHR signaling pathway (Agus et al., 2018). *Limosilactobacillus reuteri*-produced

indole derivatives have been reported to exert anti-inflammatory effects by activating

AHR signaling pathway (Cervantes-Barragan et al., 2017). Although the beneficial

effects of probiotics in protecting the host against diverse enteric infections have been

extensively examined, the interactive roles between pro-inflammatory macrophages

and microbe-tryptophan/indole metabolites in combating enteric infections have not

been fully studied yet. Here, we found that *Lacticaseibacillus rhamnosus* P118 exerts

great probiotic properties after assessing by traditional and C. elegans-infection

screening strategies. P118 exhibited broad antibacterial activities and reduced host

susceptibility to enteric *Salmonella* infection by improving intestinal dysbiosis and

fecal metabolite changes, and inhibiting intestinal pro-inflammatory responses.

Integrated omics further demonstrated that P118 probiotic activities in enhancing host

tolerance to *Salmonella* infection were mediated by microbe-derived tryptophan/indole

metabolites.

Results

Two screening approaches to convergent probiotic candidates

After identified by MALDI-TOF MS, a total of 290 bacterial isolates were isolated and

identified from 33 fermented yoghurts samples and 6 healthy piglet rectal content

samples. Those isolates consist of 63 *Streptococcus* isolates, 158

Lactobacillus/Lacticaseibacillus/Limosilactobacillus isolates, and 69 *Enterococcus*

isolates (Figure 1A, Table 1). Two screening strategies were employed in the present

study to further investigate the potential probiotic properties of the isolates: the

traditional/classic approach and the *C. elegans* infection approach. In the

traditional/classic approach, 27 isolates were screened out by milk-clotting activity

assay (Figure 1A, Table S1), among which two isolates (P118 and P199) exhibited the

highest tolerance to bile salt (0.3%~2.0%) (Figure 1B) and biofilm formation

capabilities (Figure 1C). Compared with the P199 strain, P118 strain exhibited the

highest susceptibility to multiple antimicrobials (Figure S1A). *C. elegans* has been

widely used as an invaluable model for understanding the conserved mechanisms of

host-microbe interactions, due to the similarities of gut morphology and physiological

function with human and animal (Kumar et al., 2020). In the *C. elegans* infection

approach, 8 out of 290 isolates significantly increased the survivals and prolonged the

life span of *S*. Typhimurium-infected *C. elegans* ($P < 0.05$), and three isolates (P118,

P119, P120) treated worms were more resistant to *S*. Typhimurium infection than the

other isolates ($P < 0.05$) (Figure 1D, Table S2). Integrating the results of both screening

approaches, *L. rhamnosus* P118 strain exhibits versatile capabilities as a probiotic

candidate (Figure 1E).

To illustrate the probiotic properties of *L. rhamnosus* P118, antibacterial activity

evaluation *in vitro* was further conducted, and the results showed that the fermented

supernatants of P118 significantly inhibited the growth of multiple pathogens (e.g.,

Salmonella, *Yersinia*, *S. aureus*, *E. coli*, *C. rodentium*, *P. aeruginosa*, *R. anatipestifer*)

(Figure S1B). Interestingly, the inhibitory effect of P118 against *S*. Typhimurium was

in dose- and oxygen-dependent manners and results showed that high-dose and

anaerobic cultures exhibited more antibacterial effects than low-dose and aerobic

cultures, respectively ($P < 0.05$) (Figure 1F). Active factors derived from P118 to exert

antibacterial effects were explored. The results showed that the antibacterial activities

of the fermented supernatant (pH = 3.4) were sensitive to alkalinity (pH > 5.0), trypsin,

proteinase-K, pepsin and high temperature (> 70°C), but were resistant to catalase

treatment (Figure 1G-H). The fermented cultures and supernatants of P118 not only

significantly inhibited the growth of *S*. Typhimurium *in vitro* (Figure 1I), but also

protected worms against *S*. Typhimurium infection (Figure 1G, Table S3). Although

they exhibited no antibacterial activities *in vitro* (Figure 1I), the protective activities of

heat treated P118 and P118 lysates against *S*. Typhimurium pathogenesis were observed

in *C. elegans* but weaker than that of the fermented cultures and supernatants (Figure

1G).

P118 genomics analysis suggests genetic determinants for antibacterial actions

Subsequently, whole genome sequencing of P118 was performed to explore the

potential antibacterial factors based on genomic evidence. The results showed that

species-level identity for P118 isolate with 2.99 Mb genome size was confirmed at $P <$

0.0001 (Figure S2A, Figure 1K, Table S4), and P118 draft genome was most closely

related to *L. rhamnosus* subspecies (NCTC13710 and BIO5326) (Figure 1K, Table S5,

Figure S2B). Genomic analysis using antiSMASH and BAGEL4 databases revealed the

presence of putative bacteriocin synthesis genes in P118 isolate (Figure 1L). Putative

bacteriocins predicated in P118 isolate showed high homology with known class II

bacteriocins (IIa carnobacteriocin B2 produced by *C. maltaromaticum*, IIb enterocin X

produced by *Enterococcus faecium* KU-B5, IId/b lactococcin A/G, and some novel

bacteriocin-encoded hypothetical proteins). Collective data demonstrates that *L.*

rhamnosus P118 exhibits outstanding probiotic traits.

P118 protects lethal *S*. Typhimurium infections in a murine model

The protective activity of *L. rhamnosus* P118 against enteric pathogens was further

examined in the *Salmonella*-infected murine model (Figure 2A). Administration of

P118 significantly protected mice against *Salmonella*-induced deaths and body weight

losses (Figure 2B-C), accompanied by the alleviated splenomegaly, hepatomegaly,

shortened gastrointestinal tract and clinical and pathological score (Figure 2D-F, Figure

S3B). Meanwhile, P118 significantly limited *Salmonella* colonization in the intestinal

tissues (duodenum and colon) and invasion into the peripheral organs (liver and spleen)

of mice, thereby markedly reducing *Salmonella* fecal shedding (Figure 2G).

Additionally, *Salmonella* infection-induced ileal mucosal damage was alleviated by

P118 administration, as illustrated by the increased ileal villus height (Figure 2H, Figure

S3C), ratio of villus height/crypt-depth (Figure 2H, Figure S3D), ileal microvilli height

and density (Figure 2H, Figure S3E-F), the upregulated ZO-1 mRNA expression

(Figure S4A), and the downregulated *IL-1β* mRNA expression (Figure S4E),

accompanied by the reduced IL-1β level in serum (Figure S4D). Intervention with P118

significantly increased the numbers of Ki67-positive cells (proliferating cells), Muc2-

positive cells (goblet cells), and decreased the numbers of F4/80-positive cells

(macrophages) and F4/80$^+$ iNOS$^+$ cells (pro-inflammatory macrophages) in the ileum

(Figure 2H) and also attenuated the pathological damages of peripheral organs (liver

and spleen) caused by *Salmonella* infection (Figure S3A & B).

P118 improves *Salmonella* infection by modulating gut microbiota

Gut microbes play crucial roles in host physiological activities, and colonization

resistance against enteric pathogens (Jacobson et al., 2018), and dysbiosis is supposedly

associated with *Salmonella* infection (Gillis et al., 2018a). Significant bacterial

community structures among four groups were observed (Figure 3A, Table S8), and

most of the bacterial operational taxonomic units (OTUs) were classified as group-

specific OTUs (Figure 3B). Compared to the uninfected group, *Salmonella* infection

significantly depleted 167 OTUs and enriched 60 OTUs (Figure 3C), whereas

compared to the *Salmonella*-infected group, pretreatment with P118 significantly

depleted 16 OTUs and enriched 52 OTUs (Figure 3D). Specifically, *Salmonella*

infection significantly increased the relative abundances of potentially harmful

microbes (e.g., *Salmonella*, *Escherichia*-Shigella, *Klebsiella*, *Morganella*,

Akkermansia) and significantly reduced the relative abundances of SCFA-producing

and potentially beneficial microbes (e.g., *Alloprevotella*, *Lactococcus*,

Faecalibacterium, *Limosilactobacillus*, *Parasutterella*, *Odoribacter*, *Dubosiella*,

Candidatus arthromitus) (Figure 3E, Figure S5B). Pretreatment with P118 significantly

reduced the relative abundances of pathogenic microbes (e.g., *Salmonella*,

Anaeroplasma, *Klebsiella*, *Morganella*) and significantly increased the relative

abundances of potentially beneficial or commensal microbes (e.g.,

Candidatus_Saccharimonas, *Turicibacter*, *Enterorhabdus*) (Figure 3F, Figure S5C).

Based on the changes in bacterial community structures, the internal driving forces of

gut microbial communities were investigated using ecological models. Both null model

and neutral community model analyses revealed that the stochastic processes belonging

to homogenizing dispersal exerted important roles in *Salmonella*-infected mice gut

microbiome assembly, whereas the deterministic processes (homogeneous selection

and variable selection) exerted more influences on P118-pretreated mice gut

microbiome assembly than *Salmonella*-infected mice (Figure S6A-B).

Microbe-derived tryptophan metabolites are associated with the protection

Significant differences in fecal metabolite structures among the three groups separated

into distinct clusters were also observed (Figure 4A, Table S9). Compared with the

uninfected group, *Salmonella* infection significantly upregulated 256 fecal metabolites

($\log_2$(fold change) > 1, P < 0.05) and downregulated 402 fecal metabolites ($\log_2$(fold

change) < 1, P < 0.05) (Figure 4B), and pretreatment with P118 significantly enriched

100 metabolites ($\log_2$(fold change) > 1, P < 0.05) and depleted 81 metabolites ($\log_2$(fold

change) < 1, P < 0.05) (Figure 4C). Among the significant differential metabolites, 91

metabolites (VIP > 1) downregulated in *Salmonella*-infected mice were upregulated

explicitly in P118-pretreated mice (Figure S7A&C, Table S10), and 43 metabolites

(VIP > 1) upregulated in *Salmonella*-infected mice were downregulated explicitly in

P118-pretreated mice (Figure S7B&C, Table S10). KEGG pathway analysis revealed

that fecal metabolites were mainly enriched in tryptophan metabolism, fatty acid

biosynthesis, biosynthesis of amino acids and neuroactive ligand-receptor interaction

pathways in P118-pretreated mice (Figure 4D). Interestingly, pretreatment with P118

significantly increased the fecal levels of tryptophan and tryptophan derivatives (indole,

indole-3-acrylic acid, 5-hydroxytryptophan, 5-methoxyindoleacetate) (Figure 4E).

Pearson correlation analysis showed that *S.* Typhimurium burdens (in the duodenum,

colon, liver, spleen, feces) and organ (spleen, liver) indices were negatively correlated

with the levels of fecal tryptophan and tryptophan derivatives (indole, indole-3-acrylic

acid, 5-hydroxytryptophan, protocatechuic acid, trans-cinnamaldehyde, 5-

methoxyindole-3-carbaldehyde) and positively correlated with the levels of (serotonin,

tryptophanol, 3-(2-hydroxyethyl) indole, tryptamine) (Figure 4F). Conversely, the body

weight was positively correlated with the levels of fecal tryptophan and tryptophan

derivatives (Figure 4F). Bio-Sankey network analysis further showed that 26 genera

were identified as potential bacteria that might participate in the metabolic reaction

R02340 (tryptophan synthase, indoleglycerol phosphate and indole), of which five

bacteria (*Bifidobacterium*, *Lacticaseibacillus*, *Enterococcus*, *Staphylococcus*, and

Pseudomonas, dark red if $P < 0.05$) were positively associated with indole in the

metabolic reaction R02340 (Figure 4G). These results indicate that microbe-derived

tryptophan metabolites are associated with P118-mediated protection against

Salmonella infection.

Based on the findings above that *Lacticaseibacillus* species were involved in

tryptophan biosynthesis and metabolism, *L. rhamnosus* P118 genome was reanalyzed

to determine whether P118 encode enzymes necessary to exert this function. The

genomic data further showed that P118 encoded various genes (e.g., *fldH*, *AraT*, *AspB-*

4, *AmiE*, *trpA*, *trpB*, *BCAT*, *IGPS*, *MT*) essential to biosynthesize and metabolize

tryptophan into tryptophan derivatives (Figure 4H, Table S11). The metabolomic results

of P118 cultures also validated the genomic data that P118 could secrete a diverse

profile of tryptophan-derived metabolites (e.g indole-3-acrylic acid, indole, indole-3-

lactic acid, DL-tryptophan, kynurenine, N-Acetyl-D-tryptophan, 5-

methoxyindoleacetic acid, 5-hydroxytryptophan) (Table S12). Taken together, these

data suggest that P118-derived tryptophan metabolites might contribute to the

protection against *Salmonella* infection.

Indole-3-acrylic acid protects against *Salmonella* infection by inhibiting

macrophage pro-inflammatory responses

Based on the above results, exogenous indole-3-acrylic acid (IAA) was employed to

further explore its roles against *Salmonella* infection. The antibacterial assay results

showed that IAA (> 4.4 mM) significantly inhibited *Salmonella* growth (Figure S8).

Exogenous IAA administration significantly reduced mice susceptibility to *Salmonella*

infection，as evidenced by the increased body weight (Figure 5B), the alleviated

splenomegaly and hepatomegaly (Figure 5D, 5F), and the increased colon length

(Figure 5C). Meanwhile, IAA administration significantly inhibited *Salmonella*

colonization in the intestinal tissues (cecum and colon) and invasion into the peripheral

organs (liver and spleen) of mice, thereby markedly reducing *Salmonella* fecal shedding

(Figure 5E). Additionally, *Salmonella* infection-induced ileal mucosa damage was

alleviated by exogenous IAA, as illustrated by the increased ileal villus height (Figure

5G, 5I), and the increased numbers of Muc2-positive goblet cells in the ileum (Figure

5H, 5I). Exogenous IAA intervention significantly reduced *Salmonella* infection-

induced intestinal pro-inflammatory responses, as evidenced by the decreased numbers

of F4/80-positive macrophages (Figure 5I), the downregulated mRNA (*IL-1β*, *IL-6*, *IL-*

18, *TNF-α*, *iNOS*) expression (Figure 5H), and the reduced protein (iNOS, IL-1β, IL-6,

TNF-α, markers of pro-inflammatory macrophages) expression (Figure 5I).

The *in vitro* experiments further showed that, although failed to increase the uptake of

Salmonella (Figure S9A, 0h), IAA treatment significantly reduced intracellular

Salmonella survival in RAW 264.7 macrophage cells (Figure S9A, 6h, 12h), which

could be reversed by pretreatment with AHR inhibitor CH-223191 (Figure S9A, 6h,

12h), indicating that IAA enhances the intracellular bactericidal capacity of RAW 264.7

cells. Furthermore, IAA treatment significantly inhibited *Salmonella* infection-induced

pro-inflammatory responses, as evidenced by inhibiting mRNA (*IL-1β*, *IL-6*, *IL-18*,

TNF-α, *iNOS*) expression and nitric oxide secretion of *Salmonella*-infected RAW 264.7

macrophage cells, which blocked by the presence of AHR inhibitor CH-223191 (Figure

S9B-G).

Macrophages play important roles in initiating immune responses, maintaining gut

homeostasis and phagocytic clearance of pathogens (Na et al., 2019). To further

investigate the involved roles of macrophages in P118/IAA-mediated protective effect,

in vivo macrophage depletion using clodronate liposomes was conducted. As expected,

macrophage depletion significantly blocked the protection of P118 and exogenous IAA

against *Salmonella* infection (Figure 6A-I). P118 or exogenous IAA administration

failed to increase the body weight (Figure 6B) and colon length (Figure 6E), alleviate

the splenomegaly and hepatomegaly (Figure 6C, 6D), inhibit the colonization, invasion

and shedding of *Salmonella* (Figure 6F) after the intestinal macrophage depletion.

Additionally, after depleting the intestinal macrophage, P118 or exogenous IAA

administration also failed to attenuate *Salmonella* infection-induced intestinal mucosa

damage (Figure 6G, 6H, 6I) and pro-inflammatory responses (Figure 6H, 6I). Taken

together, these findings indicate that P118 and IAA protect against *Salmonella* infection

by inhibiting pro-inflammatory responses of intestinal macrophage.

Discussion

Fermented foods and mammalian intestines are rich in lactic acid bacteria that can

produce organic acids and lactobacillin to inhibit overgrowth, adhesion, invasion and

toxin secretion of pathogens (Gareau et al., 2010; Wan et al., 2019). Traditionally, the

culturomics strategy, together with multiple evaluation assays, is the primary way to

search potential probiotic species from the complex microbial community, which is

both time- and labor-consuming (Wang et al., 2022). Although the whole genomic

sequencing-based approach is an emerging favorable way to screen potential probiotic

strains, it's hard to widely adopt based on numerous isolates due to the high sequencing

costs (Schwarze et al., 2020). Given that the similarities of gut morphology and

physiological function with human and animal, *C. elegans* provides an invaluable

model to investigate host-microbe interactions and probiotic properties of beneficial

microbes at low cost (Kumar et al., 2020; Poupet et al., 2020). In the present study, we

found *Salmonella* infection significantly induced death and decreased the lifespan of *C.*

elegans, indicating *C. elegans* is very susceptible to *S.* Typhimurium infection,

consistent with a previous report (Rangan et al., 2016). Integrating the results of

culturomics and *C. elegans* infection strategies, *L. rhamnosus* P118 with excellent

probiotic properties was screened from 290 isolates and exhibited broad *in vitro*

antibacterial activities. Interestingly, high fermented dose and anaerobic cultures of

P118 showed more antibacterial effects than low dose and aerobic cultures, indicating

that the active factors were more released by P118 at the stationary phase under

anaerobic conditions. What's more, the antibacterial activities of the fermented

supernatant were sensitive to alkalinity, trypsin, proteinase-K, pepsin, and high

temperature, indicating that the active secreted factors in the fermented supernatant

were protein- or peptide-based active agents, consistent with a previous report that *L.*

rhamnosus exerted antibacterial effects by secreting bioactive peptides (Iram et al.,

2022). Although limited studies have reported bacteriocin-like activity in *L. rhamnosus*,

it remains unclear whether this species produces active bacteriocins (Kankainen et al.,

2009). Nonetheless, genomic data in this study revealed the presence of putative

bacteriocin synthesis genes in P118 genome. In agreement with a previous study

(Kankainen et al., 2009), the predicated bacteriocins showed high homology with

known class II bacteriocins (e.g., IIa carnobacteriocin B2 produced by *C.*

maltaromaticum, IIb enterocin X produced by *Enterococcus faecium* KU-B5), which

needs to be further validated by gene-editing technology and synthetic biology.

To further validate the antibacterial activities of *L. rhamnosus* P118 against enteric

pathogens, *Salmonella*-infected mice model was employed. We found that

administration of P118 significantly reduced mice susceptibility to *S*. Typhimurium

infection by improving intestinal health, dysbiosis and the changes of fecal metabolite

profiles. Colonized *S*. Typhimurium in the intestine disrupts structural and functional

intestinal integrity, leading to substantial immune response and severe epithelial barrier

damage, and then migrates to internal organs (liver, spleen) via the blood circulatory

system, causing splenomegaly and hepatomegaly (Rogers et al., 2022; Wang et al.,

2021a). In the present study, P118 significantly increased *Salmonella*-infected mice

survival and alleviated *Salmonella*-induced splenomegaly and hepatomegaly. The

beneficial attenuated effect was partly contributed by the reduced *Salmonella*

colonization in intestine and invasion into the peripheral organs, and eventually, the

decreased fecal shedding, consistent with previous reports (Liu et al., 2023; Wu et al.,

2024). It's reported that *S*. Typhimurium exploits and disrupts intestinal barrier integrity

to adhere, and disseminate deeper into intestinal mucosa, and evade immune clearance

(Bernal-Bayard and Ramos-Morales, 2018; Hiyoshi et al., 2022). In the current study,

P118 treatment also significantly improved the structural and functional intestinal

integrity, attenuated inflammatory responses, reduced the numbers of F4/80-positive

macrophages and proinflammatory macrophages ($F4/80^+$ $iNOS^+$ cells), and increased

the numbers of Ki67-positive and Muc2-positive goblet cell in the ileum, demonstrating

that P118 improved the intestinal health and reducing pro-inflammatory responses of

intestinal macrophages, which further confirms the beneficial alleviated effect exerted

by *L. rhamnosus* P118.

Intestinal microbes play essential roles in providing colonization resistance against

enteric pathogens and pathobionts, and limiting adhesion, invasion and transmission of

pathogens (Caballero-Flores et al., 2022; Jacobson et al., 2018). Microbial dysbiosis

and excessive immune responses induced by enteric pathogens can, in turn, aggravate

and exacerbate gut permeability and damage (Read et al., 2021; Rogers et al., 2022).

Many studies have reported that *Salmonella* infection was often accompanied by

microbial dysbiosis (Gillis et al., 2018b; Read et al., 2021; Wang et al., 2021a). The

current results found that *L. rhamnosus* P118 treatment significantly restored the

bacterial community structures in *Salmonella*-infected mice, and significantly reduced

the relative abundances of potentially harmful microbes (e.g., *Salmonella*, *Klebsiella*,

Anaeroplasma, *Morganella*), which belong to pathogens and pathobionts (Dong et al.,

2022; Galan, 2021; Laupland et al., 2022). Structure composition and assembly of

microbial communities are essential for ecosystem function (Shi et al., 2023; Xun et al.,

2019), and revealing the underlying mechanisms of microbial community assembly is

a major goal of microbial community ecology (Jiao et al., 2020; Stegen et al., 2012).

Generally, stochastic (e.g., dispersal events, ecological drift, random birth, death) and

deterministic (e.g., interspecies interactions (e.g., competition, facilitation, mutualisms

and predation), species traits, environmental factors) processes are two major

ecological processes that drive microbiome assembly (Stegen et al., 2012; Zhou and

Ning, 2017). Deterministic processes mainly involve non-random and niche-based

mechanisms with abiotic and biotic factors that influence microbial community

assembly (Vellend, 2010), whereas stochastic processes mainly reflect random changes

in the relative abundance of species (Jiao et al., 2020; Jiao et al., 2021). Although

important in shaping the diversities of microbial composition and functions, the relative

contribution of microbiome assembly processes varies with different habitats (Jiao et

al., 2020; Jiao et al., 2021). The present study revealed that the stochastic processes

played dominant roles in intestinal microbiome assembly in *Salmonella*-infected mice,

indicating random changes occurred in the relative abundance of species. Conversely,

the deterministic processes played more significant roles in driving intestinal microbial

community assembly in P118-pretreated mice than in *Salmonella*-infected mice,

indicating abiotic factors (e.g., dietary intervention) or biotic interactions (e.g.,

microbial competition, facilitation, mutualism, predation, host filtering) might

influence the microbial community.

Accumulating evidence revealed that gut microbes-derived metabolites (e.g., short-

chain fatty acids, tryptophan/indole-derived metabolites, bacteriocins, bile acid, natural

products) are crucial mediators in host physiological activities (Cani, 2019; Wlodarska

et al., 2017), and metabolic disorders are major risk factors for bacterial infections

(Olive and Sassetti, 2016). In this study, the results showed that *L. rhamnosus* P118

improved *Salmonella* infection-induced fecal metabolite changes. Interestingly, P118

significantly increased the fecal levels of tryptophan and its derivatives (indole, indole-

3-acrylic acid, 5-hydroxytryptophan), and those metabolites were negatively correlated

with *S.* Typhimurium burdens (in the duodenum, colon, liver, spleen, feces) and organ

(spleen, liver) indices, and positively correlated with the body weight of mice,

indicating that microbiota-derived tryptophan/indole metabolites play beneficial roles

in P118 mediated probiotic activities in enhancing host tolerance to *Salmonella*

infection. It's reported that through binding to the aryl hydrocarbon receptor (AHR) of

host, bacterial tryptophan derivatives act as triggers to stimulate immune responses and

gut hormones to exert anti-inflammatory activities, enhance intestinal epithelial barrier,

and promote gastrointestinal motility, which contributes to host gastrointestinal

homeostasis and health (Agus et al., 2018; Cani, 2019). Our genomic and metabolomic

data further revealed that P118 could produce a wide array of tryptophan-derived

metabolites owing to encoding a variety of enzyme genes necessary to metabolize

tryptophan into indole derivatives; consistently, some *Lactobacillus* species broadly

conserved enzyme encoding genes that are involved in tryptophan metabolism

(Montgomery et al., 2022). The above results indicate that *L. rhamnosus* P118- and

microbe-derived tryptophan/indole metabolites might play positive roles in P118-

mediated probiotic activities. Although significant increases in microbe-derived

tryptophan/indole metabolites were observed in P118-treated mice, and it was

established that P118 can metabolize tryptophan into indole derivatives, it remains to

be investigated whether the differential tryptophan/indole metabolites were directly

derived from P118 or from other microbes. To test the roles of differential

tryptophan/indole metabolites in P118-mediated beneficial effects, exogenous indole-

3-acrylic acid, which significantly and negatively correlated with *S.* Typhimurium

burdens and hepatosplenomegaly mentioned above, was employed to investigate its

protective effect against *Salmonella* infection. Interestingly, we found that indole-3-

acrylic acid administration significantly enhanced mice tolerance to *S*. Typhimurium

infection by improving intestinal health and attenuating pro-inflammatory responses of

intestinal macrophages. Furthermore, the *in vitro* results also provide that indole-3-

acrylic acid enhanced bactericidal capacity of RAW 264.7 cells and inhibited *S*.

Typhimurium-induced pro-inflammatory responses, consistent with a previous report

(Wlodarska et al., 2017). Taken together, these results demonstrate that *L. rhamnosus*

P118/microbe-derived indole-3-acrylic acid play beneficial roles in P118-mediated

probiotic activities.

Macrophages are important in initiating immune responses, maintaining intestinal

immune homeostasis, phagocytic clearance of pathogens, tissue repair, and host defense

(Na et al., 2019; Neurath, 2024). It's reported that macrophages with strong plasticity

could be polarized into pro-inflammatory (M1) phenotype or proresolving (M2)

phenotype (Na et al., 2019). M1 macrophage activation is crucial and necessary for

pathogen clearance (Na et al., 2019). However, exaggerated immune responses

triggered by M1 macrophage are detrimental to the host by inducing tissue damage and

cell deaths (Wang et al., 2020). The current study showed that both *L. rhamnosus* P118

and indole-3-acrylic acid significantly attenuated *S*. Typhimurium infection-induced

pro-inflammatory responses, indicating that pro-inflammatory macrophages might be

key targets for *L. rhamnosus* P118 exerting probiotic effect. This hypothesis was

confirmed by macrophage depletion experiments that the protective effects of *L.*

rhamnosus P118 and indole-3-acrylic acid against *Salmonella* infection were abrogated

after intestinal macrophage depletion. Taken together, these results indicate that *L.*

rhamnosus P118/microbe-derived indole metabolites enhance host tolerance to

Salmonella infection by reducing intestinal pro-inflammatory responses, which also

provides a potential alternative strategy for using probiotics and their derived

metabolites to treat intestinal inflammatory disorders such as inflammatory bowel

disease (Lavelle and Sokol, 2020; Neurath, 2024).

Conclusion

In summary, two distinctive approaches were used to screen the probiotic candidate,

and the P118 strain was underscored for beneficial effects on a murine infection model.

Further, bacterial genomic sequencing, gut microbiota and metabolomic analysis

pinpointed the microbe-derived tryptophan/indole could be the importance of P118

probiotic properties. Nevertheless, the newly found P118 could enhance host tolerance

to *Salmonella* infections via various pathways, including direct antibacterial actions,

inhibiting *Salmonella* colonization and invasion, attenuating pro-inflammatory

responses of intestinal macrophages, and modulating gut microbiota mediated by

microbe-derived indole metabolites (Figure 7). Further investigations are needed to

assess whether the mechanisms observed in P118 are strain-specific or broadly

applicable to other *L. rhamnosus* strains, or LAB species in general.

Materials and methods

Bacterial isolation and culture

Lactic acid bacteria (LAB) and *Enterococcus* strains were isolated from 39 samples: 33

fermented yoghurts samples (collected from families in multiple cities of China,

including Lanzhou, Urumqi, Guangzhou, Shenzhen, Shanghai, Hohhot, Nanjing,

Yangling, Dali, Zhengzhou, Shangqiu, Harbin, Kunming, Puer), and 6 healthy piglet

rectal content samples without pathogen infection and diarrhea in pig farm of Zhejiang

province (Table 1). Ten isolates were randomly selected from each sample. De Man-

Rogosa-Sharpe (MRS) with 2.0% $CaCO_3$ and Brain heart infusion (BHI) broths

(Huankai Microbial, Guangzhou, China) were used for bacteria isolation and

cultivation. Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass

Spectrometry (MALDI-TOF MS, Bruker Daltonik GmbH, Bremen, Germany) method

was employed to identify of bacterial species with a confidence level $\geq$ 90% (He et

al., 2022). *Lacticaseibacillus rhamnosus* P118 (P118) and *Salmonella enterica* serovar

Typhimurium SL1344 (STm) were cultured in MRS and Luria Broth (LB) medium at

37 °C overnight under aerobic conditions, separately. The final concentration of

bacterial isolates was constantly checked by the spreading plate method (Wang et al.,

2023). After being centrifugated at 4000× g for 15 min at 4 °C, the fermented

supernatant of *L. rhamnosus* P118 was collected by centrifugation at 4000× g for 15

min at 4 °C, and then filtered through a 0.22 μm membrane (Merck Millipore,

Burlington, MA, USA) kept at 4 °C for further use.

Probiotic property evaluation of candidate isolates

For milk-clotting activity analysis of probiotics (Zhang et al., 2023), overnight

fermented lactic acid bacteria isolates under logarithmic growth were added to test

tubes containing skim milk (10% skim milk with 10 mM $CaCl_2$) at 1:20 ratio (v/v) and

then incubated at 37 °C. The coagulating time, final pH value and acidity of milk were

measured during the curdling process.

The bile salt tolerance assay was conducted according to a previous report (Wang et al.,

2023). Briefly, the isolates under logarithmic growth were resuspended into fresh MRS

medium (1% v/v) containing different (wt/v, 0, 0.3%, 0.6%, 1.0%, 1.5%, 2.0%) porcine

bile (MACKLIN®, Shanghai, China) and then plated into 96-well plates, after

incubation for 24h at 37 °C, the absorbance at 620 nm was measured by a SpectraMax

M5 reader (Molecular Devices, Sunnyvale, CA, USA).

Biofilm-forming assay of the isolates was performed as described previously (Zhou et

al., 2023). Briefly, 200 µL overnight cultures (adjusted to 2×10^6 CFU/mL) were added

into 96-well flat-bottom plates. After incubation for 36h at 37 °C, the planktonic cells

were discarded. After being gently washed three times with sterile PBS, the adherent

cells were stained with 200 µL of 0.4% crystal violet for 15 min at room temperature

and then washed with sterile PBS three times again. Finally, 200 µL 75% ethanol was

added to dissolve the stained adherent cells for 15 min, and biofilm was quantified by

measuring the absorbance at 590 nm using a SpectraMax M5 reader.

For antimicrobial susceptibility test, twelve antimicrobial agents (penicillins (PG),

erythromycin (EM), lincomycin (LM), gentamicin (GM), doxycycline (DOX),

ceftriaxone (CTR), norfloxacin (NOR), azithromycin (AZM), fleroxacin (FLE),

vancomycin (VA), streptomycin (STR)) were employed to evaluate the antimicrobial

susceptibility of the candidate probiotics by utilizing the Kirby-Bauer disk diffusion

method (Hudzicki, 2009).

The agar diffusion method (Wang et al., 2023) was used to evaluate the antimicrobial

activity against multiple pathogens *in vitro*. Briefly, bacterial culture medium (LB, BHI)

containing 0.25% (v/v) different zoonotic pathogens under logarithmic growth was

poured into 10-cm plates (Corning, NY, USA), separately. Then, 200 µL of the

fermented sterile supernatant collected from the overnight cultured *L. rhamnosus* P118

was added into 8 mm agar wells created by Oxford cup. The bacteriostatic effect was

evaluated by the growth inhibition zone around the 8-mm agar wells after overnight

incubation at 37 °C.

***Caenorhabditis elegans* infection**

C. elegans infection model was performed as previously described (Rangan et al., 2016).

C. elegans N2 (Bristol) were routinely raised on Nematode Growth Media (NGM)

feeding with *E. coli* OP50, and L4-stage worms were prepared according to the

description in WormBook (Girard et al., 2007). 200 µL of the isolated isolates (adjusted

to 5×10^8 CFU/mL using M9 medium) and *L. rhamnosus* P118-related substances (e.g.,

mixed cultures, fermented sterile supernatants, P118 lysate, dead P118) were poured

onto NGM in 24-well plates, separately, and dried at 22 °C for 4 h. L4-stage worms

were then transferred to the lawns of the poured NGM plates (30 worms/plate) with

three plates per group for 24h. Then, the treated worms were washed and transferred to

lawns of STm (1×10^8 CFU/lawn) on NGM plates for infection. Worm survival was

monitored at 24-hour intervals for sixteen days.

Whole genome sequencing and data analysis

Whole genome sequencing, quality control, assembly and annotation of *L. rhamnosus*

P118 was conducted according to our previous study (Zhou et al., 2023) using PacBio

Sequel platform and Illumina NovaSeq PE150 at Novogene (Novogene Co, Ltd.,

Beijing, China), and the P118 draft genome is available in the Sequence Read Archive

under accession number PRJNA848987. *L. rhamnosus* P118 assembly was queried

against the NCBI non-redundant prokaryotic genomes database using the Microbial

Genomes Atlas (MIGA) webserver (database update to 10/10/2023) (Rodriguez et al.,

2018). Taxonomic classification was inferred by the maximum average amino acid

identity (AAI) against all genomes in the database, with p-values estimated from the

distribution of all the reference genomes in NCBI's RefSeq at each taxonomic level as

a readout of classification probability. Average nucleotide identity (ANI) and AAI

tables of maximally the top-50 reference hits in the database were extracted and graphed

as x-y scatter plots to determine the nearest phylogenetic neighbors. To generate

phylogenic trees of isolates and MIGA-identified nearest phylogenetic neighbors, the

interfered proteome from PROKKA isolate annotations, or as publicly available in

NCBI for applicable reference genomes were uploaded to the ANI/AAI-Matrix

calculator (Rodriguez-R and Konstantinidis, 2016). The resulting phylogenetic tree

based on AAI was visualized using the interactive tree of life (iTOL) (Letunic and Bork,

2021).

L. rhamnosus P118 tryptophan-associated enzymes were identified as previously

described (including TNA, TMO, TrpD, ArAT, ALD, IPDC, FldH, and AmiE)

(Montgomery et al., 2022). Briefly, enzyme commission numbers, where available, or

alternatively enzyme names, were queried in PATRIC, PATRIC Global Family (PGF)

cross-genus identifiers were extracted and compiled for query within the *L. rhamnosus*

P118 genome. Protein sequences of *L. rhamnosus* P118 identified enzymes were

analyzed using InterProScan for additional functional prediction, and sequence

homology to the previously experimentally validated enzymes within other bacterial

species was determined using Blastp at 30% cross-genus identify and 90% identity

within each species.

_S._ Typhimurium murine infection model

Animal assays, specifically eighty 6-week-old female C57BL/6 mice (Slac Animal Inc.,

Shanghai, China), were conducted in the Laboratory Animal Center of Zhejiang

University under light-controlled (12h light/dark cycle), temperature-controlled (22 ±

2°C) and humidity-controlled (55 ± 5°C) conditions. Forty mice were randomly

selected for survival evaluation under _S_. Typhimurium infection, and the rest were used

for sample collection on day 3 post-infection (Figure 2A). The mice were randomly

divided into four groups (n = 10/group): Control group (C), _L. rhamnosus_ P118 group

(P), _S_. Typhimurium-infected group (S), and _L. rhamnosus_ P118 protective group (P+S).

Mice in the C and S groups were drinking sterile water every day, respectively. Mice

in the P and P+ST groups were orally administered with _L. rhamnosus_ P118 (1×10^8

CFU/mouse) daily for seven days. After that, mice were orally ingested with _S_.

Typhimurium SL1344 (1×10^8 CFU/mouse) or sterile PBS. All mice were allowed free

access to water and food, and weighed every day. Mice were euthanized on day 3 post-

infection. Spleen and liver were weighted for organ index calculation according to the

formula: organ index = organ weight (g) / body weight (g) *1000.

To evaluate the protective effect of indole-3-acrylic acid (IAA), eighteen mice were

randomly divided into three groups (n = 6/group): Control group (C), _S_. Typhimurium-

infected group (S), and indole-3-acrylic acid protective group (IAA+S). Mice in the C

and S groups were drinking sterile water every day, respectively. IAA (Sigma, I3807)

was dissolved in 1 M NaOH in PBS and adjusted pH to 7.4 with 1 M HCl. Mice in the

IAA+ST groups were orally administered with 75mg/kg IAA daily for seven days. After

that, mice were orally ingested with *S.* Typhimurium SL1344 or sterile PBS. At 3 day-

post-infection, mice were euthanized for sample collection.

***In vivo* macrophage depletion**

To evaluate the involved roles of macrophages, macrophage depletion assay was

conducted through intraperitoneal injection with 250 μL/mouse clodronate liposomes

(CLP, From Vrije Universiteit Amsterdam) twice (2 days prior and 1 day after

Salmonella infection, separately) according to the manufacturer's instructions. Thirty-

three mice were randomly divided into six groups (n = 3~6/group): Control group (C),

S. Typhimurium-infected group (S), Control group with macrophage depletion

(CLP+C), *S.* Typhimurium-infected group with macrophage depletion (CLP+S), IAA

protective group with macrophage depletion (CLP+IAA+S), and P118 protective group

with macrophage depletion (CLP+P+S). At 3 day-post-infection, mice were euthanized

for sample collection.

Macrophage cell culture

Murine macrophage cell line RAW 264.7 was cultured in DMEM/F12 medium (Gibco,

Carlsbad, CA) supplemented with 10% FBS (Gibco, CA) and 1% (v/v) antibiotic

solution (100 μg/mL streptomycin + 100 U/mL penicillin, Sigma-Aldrich, St. Louis,

MO) at 37°C in a 5% humidified CO_2 incubator. If not mentioned, the antibiotic

solution (100 μg/mL streptomycin and 100 U/mL penicillin) was not added into

DMEM/F12 medium in the further experiment.

For *S*. Typhimurium killing assay, RAW 264.7 cells seeded into 12-well plates (2 ×

$10^{\wedge 6}$ cells/well) were treated with 100 μM IAA for 6h, followed by infecting with *S*.

Typhimurium (MOI = 10) for 1h. After washing 3 times with sterile PBS, the infected

RAW 264.7 cells were incubated for 0h, 6h and 12h in DMEM/F12 medium containing

gentamicin (50 μg/mL). At each time point, the infected RAW 264.7 cells were washed

with sterile PBS for 4 times and lysed with 0.01% Triton X-100 diluted in PBS. The

serial 10-fold dilutions of cell lysates were immediately plated on LB agar plates to

determine bacterial viability.

For the immune response assay, RAW 264.7 cells seeded into 12-well plates (2 × $10^{\wedge 6}$

cells/well) were pretreated with CH-223191 (AHR inhibitor, 10 μM) for 6h,

subsequently treating cells with 100 μM IAA for another 6h. After 6h, RAW 264.7 cells

were infected with *S*. Typhimurium (MOI = 10) for 2h. After washing 3 times with

sterile PBS, the infected RAW 264.7 cells were collected using RNAiso Plus (TaKaRa,

Dalian, China) for total RNA extracting and quantitative real-time PCR (qPCR). The

primer sets are listed in Table S6.

S. Typhimurium burden

S. Typhimurium loads in tissues (duodenum, cecum, colon, liver and spleen) and

shedding in feces were determined as previously described (Xu et al., 2018). Briefly,

tissues and fecal samples were collected under sterile conditions and homogenized in

sterile PBS containing 0.1% Triton X-100. Serial 10-fold dilutions of tissue

homogenates were spread onto SS agar plates in triplicate and then incubated at 37°C

overnight to quantify the bacterial colony-forming unit (CFU).

Histopathology, immunofluorescent analysis, transmission electron microscope

(TEM) observation and clinical symptom score

Tissues (ileum, liver and spleen) fixed in 4% paraformaldehyde were embedded in

paraffin, sliced, dehydrated and then sectioned for hematoxylin and eosin (H&E)

staining. The tissue slices were imaged and analyzed by using Olympus Microscope

(Tokyo, Japan). For the immunofluorescent assay, the paraffin-embedded ileum

samples were incubated with primary antibodies against Ki67 (GB121141, Servicebio,

China), lysozyme (GB11345, Servicebio, China), Muc2 (GB11344, Servicebio, China),

F4/80 (GB113373, Servicebio, China), iNOS (GB11119, Servicebio, China), IL-1β

(GB11113, Servicebio, China), IL-6 (GB11117, Servicebio, China), TNF-α (GB11188,

Servicebio, China) for overnight at 4°C and then further incubated with secondary

antibodies (Alexa Fluor® 488-conjugated goat anti-rabbit/mouse IgG) or HRP-

conjugated goat anti-rabbit secondary antibody (GB23303, Servicebio, China) for 60

min at room temperature. Finally, ileum slices were digitalized by Pannoramic MIDI

(3DHISTECH, Hungary).

TEM observations of ileum tissues were prepared according to our previous study

(Peng et al., 2022). Briefly, after being fixed with 2.5% buffered glutaraldehyde, ileum

segments were washed three times by cold 100-mM phosphate buffers, and then post-

fixed in 0.1% osmium tetroxide for 2 h, rapidly dehydrated in ascending grades of

ethanol (30, 50, 70, 95, and 100%), and moved into a 1:1 mixture of propylene oxide

and epoxy araldite. Finally, the ileum samples were observed and captured by the

transmission electron microscope (JEOL, Tokyo, Japan).

The clinical symptom scores were evaluated and graded by two blinded assessors

according to previous reported with modified (Burkholder et al., 2012), and the scoring

criteria are list in Table S7.

ELISA assay and qPCR

The serum contents of interleukin (IL)-1β and IL-18 were determined by ELISA kits

(Solarbio, Beijing, China) according to the manufacturer's instructions.

Total RNA extracted from intestinal tissues and RAW 264.7 cells by RNAiso Plus kit

(TAKARA, Dalian, China) was reverse-transcribed using PrimeScript II 1st Strand

cDNA Synthesis Kit (TAKARA, Dalian, China) according to the manufacturer's

instructions. The qPCR was then performed on the StepOne real-time PCR system

(Applied Biosystems) using SYBR PremixEx TaqII (TAKARA). The primer sets are

listed in Table S6. Fold changes were calculated after normalizing to the housekeeping

gene β-actin using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

Microbiota analysis

The TIANamp Stool DNA Kit (Tiangen, Beijing, China) was employed to extract fecal

bacterial genomic DNA, and fecal bacterial communities were investigated using 16S

rDNA sequencing by targeting the V3-V4 hypervariable region. The 16S rDNA

sequencing was then performed on an Illumina NovaSeq platform (Illumina Inc., San

Diego, CA, USA). The quality filter of the paired-end raw sequences and a cluster of

the filtered sequences into the operational taxonomic unit (OTU) at 97% similarity was

performed by QIIME software (version 1.9.1). Microbial OTU representative

sequences were assigned to a taxonomic lineage by the RDP classifier based on the

SILVA database (v 132 release).

To investigate the dissimilarities in bacterial communities and fecal metabolomes,

principal coordinates analysis (PCoA), analysis of similarities (ANOSIM),

permutational multivariate analysis of variance (PERMANOVA) and multi-response

permutation procedure (MRPP) were calculated using "vegan" package and visualized

using "ggplot2" package. Difference analysis of bacterial communities based on OUT

levels among groups was calculated using "DESeq2" package and was envisioned by

"UpSetR" and "ggplot2" packages. The significant differences in bacterial taxonomies

were analyzed and visualized by statistical analysis of taxonomic and functional

profiles (STAMP) software with a two-sided Welch's t-test (Parks et al., 2014).

The sloan neutral model was employed to estimate the importance of neutral processes

in the assembly of bacterial communities using "MicEco" package (Sloan et al., 2006).

A null-model-based statistical framework was conducted to evaluate the relative

importance of determinism and stochasticity in bacterial community assembly (Stegen

et al., 2013; Xu et al., 2022). Two assembly processes were defined as deterministic

processes ($|\beta NTI|>2$) and stochastic processes ($|\beta NTI|\leq2$) based on the absolute values

of beta Nearest Taxon Index (βNTI) (Xu et al., 2020). Additionally, five ecological

assembly processes were then interpreted as homogeneous selection ($\beta NTI<-2$),

homogenizing dispersal ($|\beta NTI|\leq2$ and $RC_{bray}<-0.95$), undominated ($|\beta NTI|\leq2$ and

$|RC_{bray}|<0.95$), dispersal limitation ($|\beta NTI|\leq2$ and $RC_{bray}>0.95$) and variable

(heterogeneous) selection (βNTI>2) based on the threshold of the absolute values of

βNTI and Bray-Curtis-based Raup-Crick (RC$_{bray}$) (Jiao et al., 2020; Xu et al., 2022).

Metabolomics analysis

Untargeted metabolomics was investigated to analyze the fecal metabolomes of mice.

The method for extracting metabolites from feces was conducted according to our

previous study described (Wang et al., 2021b). After extracting metabolites from feces,

UHPLC-MS/MS analyses of samples were performed by Vanquish UHPLC systems

(Thermo Fisher, Germany) coupled with Orbitrap Q Exactive^{TM} HF-X mass

spectrometers (Thermo Fisher, Germany) in Novogene Co., Ltd. (Beijing, China).

Compound Discoverer 3.1 (CD3.1, Thermo Fisher) was employed to conduct peak

alignment, and quantify metabolites generated by UHPLC-MS/MS. After normalizing

to total spectral intensity and predicting molecular formula, the peaks were queried

against mzCloud, mzVault and MassList databases to extract the quantitative results.

Annotation of fecal metabolite signatures was queried against the KEGG database,

HMDB database and LIPIDMaps database. Significant differences in fecal metabolites

were analyzed using univariate analysis (t-test) and partial least-squares discriminant

analysis (PLS-DA), in which fecal metabolites with variable importance in the

projection (VIP) > 1 and $P < 0.05$ were considered differential metabolites. Pearson

correlation analysis between *S.* Typhimurium burden (in tissues and feces) and fecal

metabolites was analyzed and visualized using "linkET" package (Huang, 2021).

MetOrigin platform was employed to analyze the sources of microbial metabolites and

their metabolic functions in fecal metabolomes (Yu et al., 2022).

Statistical Analysis

The significance of the results was analyzed by one-way analysis of variance (ANOVA) with Tukey's multiple comparisons test using SPSS v24 (SPSS Inc., Chicago, IL, USA) and statistical graphs were visualized by GraphPad Prism v8.0 (GraphPad Software, CA) and package "ggplot2" of R software (v4.3.1). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Acknowledgements

This work was supported by the National Program on Key Research Project of China (2022YFC2604201) as well as the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement No. 861917-SAFFI, Zhejiang Provincial Key R&D Program of China (2023C03045, 2022C02024), Key Research and Development Program of Hangzhou (202203A08), Zhejiang Provincial Natural Science Foundation of China (LQ23C180005).

Conflict of interest statement

No potential conflict of interest was reported by the author(s).

Data availability statement

16S rRNA sequencing data is deposited at the Genome Sequence Archive (GSA) of the BIG Data Center (https://bigd.big.ac.cn/gsa/) under accession no. PRJCA021751. *L. rhamnosus* P118 isolate draft genome assembly is publicly available in the NCBI under accession no. PRJNA848987.

Ethics statement

Animal procedures followed the guidelines and were approved by the Institutional

Animal Care and Use Committee of Zhejiang University (Permission number:

ZJU20220295).

Author contributions

Baikui Wang and Xianqi Peng: Investigation, Validation, Methodology, Data Analysis,

Visualization, Formal analysis, Writing - original draft & review & editing; Xiao Zhou,

Jiayun Yao, Haiqi Zhang, Weifen Li, Yan Li: Investigation, Writing – review; Min Yue:

Conceptualization, Resources, Methodology, Supervision, Writing – review & editing,

Funding acquisition, Project administration.

References

Agus A, Planchais J, Sokol H. Gut microbiota regulation of tryptophan metabolism in
health and disease. Cell Host Microbe 2018;23:716-24.

Bernal-Bayard J, Ramos-Morales F. Molecular mechanisms used by salmonella to
evade the immune system. Curr Issues Mol Biol 2018;25:133-68.

Britton GJ, Faith JJ. Causative microbes in host-microbiome interactions. Annu Rev
Microbiol 2021;75:223-42.

Burkholder T, Foltz C, Karlsson E, Linton CG, Smith JM. Health evaluation of
experimental laboratory mice. Curr Protoc Mouse Biol 2012;2:145-65.

Caballero-Flores G, Pickard JM, Nunez G. Microbiota-mediated colonization
resistance: Mechanisms and regulation. Nat Rev Microbiol 2022.

Caballero-Flores G, Pickard JM, Nunez G. Microbiota-mediated colonization
resistance: Mechanisms and regulation. Nat Rev Microbiol 2023;21:347-60.

Cani PD. Microbiota and metabolites in metabolic diseases. Nat Rev Endocrinol
2019;15:69-70.

Cervantes-Barragan L, Chai JN, Tianero MD, Di Luccia B, Ahern PP, Merriman J,
Cortez VS, Caparon MG, Donia MS, Gilfillan S, Cella M, Gordon JI, Hsieh CS,

Colonna M. Lactobacillus reuteri induces gut intraepithelial
cd4(+)cd8alphaalpha(+) t cells. Science 2017;357:806-10.

Dong N, Yang X, Chan EW, Zhang R, Chen S. Klebsiella species: Taxonomy,
hypervirulence and multidrug resistance. EBioMedicine 2022;79:103998.

Eng SK, Pusparajah P, Ab Mutalib NS, Ser HL, Chan KG, Lee LH. Salmonella: A
review on pathogenesis, epidemiology and antibiotic resistance. Frontiers in
Life Science 2015;8:284-93.

Fishbein SRS, Mahmud B, Dantas G. Antibiotic perturbations to the gut microbiome.
Nat Rev Microbiol 2023;21:772-88.

Fu A, Mo Q, Wu Y, Wang B, Liu R, Tang L, Zeng Z, Zhang X, Li W. Protective effect
of bacillus amyloliquefaciens against salmonella via polarizing macrophages to
m1 phenotype directly and to m2 depended on microbiota. Food & function
2019;10:7653-66.

Galan JE. Salmonella typhimurium and inflammation: A pathogen-centric affair. Nat
Rev Microbiol 2021;19:716-25.

Gareau MG, Sherman PM, Walker WA. Probiotics and the gut microbiota in intestinal
health and disease. Nat Rev Gastroenterol Hepatol 2010;7:503-14.

Gillis CC, Hughes ER, Spiga L, Winter MG, Zhu W, Furtado De Carvalho T, Chanin
RB, Behrendt CL, Hooper LV, Santos RL, Winter SE. Dysbiosis-associated
change in host metabolism generates lactate to support salmonella growth. Cell
Host Microbe 2018a;23:54-64 e6.

Gillis CC, Hughes ER, Spiga L, Winter MG, Zhu W, Furtado De Carvalho T, Chanin
RB, Behrendt CL, Hooper LV, Santos RL, Winter SE. Dysbiosis-associated
change in host metabolism generates lactate to support salmonella growth. Cell
Host Microbe 2018b;23:570.

Girard LR, Fiedler TJ, Harris TW, Carvalho F, Antoshechkin I, Han M, Sternberg PW,
Stein LD, Chalfie M. Wormbook: The online review of caenorhabditis elegans
biology. Nucleic Acids Res 2007;35:D472-5.

He D, Zeng W, Wang Y, Xing Y, Xiong K, Su N, Zhang C, Lu Y, Xing X. Isolation and
characterization of novel peptides from fermented products of lactobacillus for

ulcerative colitis prevention and treatment. Food Science and Human Wellness
2022;11:1464-74.

Hiyoshi H, English BC, Diaz-Ochoa VE, Wangdi T, Zhang LF, Sakaguchi M, Haneda
T, Tsolis RM, Baumler AJ. Virulence factors perforate the pathogen-containing
vacuole to signal efferocytosis. Cell Host Microbe 2022;30:163-70 e6.

Huang H. Linket: Everything is linkable. R package version 0.0.3 2021.

Hudzicki J. Kirby-bauer disk diffusion susceptibility test protocol. American society
for microbiology 2009;15:55-63.

Iram D, Kindarle UA, Sansi MS, Meena S, Puniya AK, Vij S. Peptidomics-based
identification of an antimicrobial peptide derived from goat milk fermented by
lactobacillus rhamnosus (c25). J Food Biochem 2022;46:e14450.

Jacobson A, Lam L, Rajendram M, Tamburini F, Honeycutt J, Pham T, Van Treuren W,
Pruss K, Stabler SR, Lugo K, Bouley DM, Vilches-Moure JG, Smith M,
Sonnenburg JL, Bhatt AS, Huang KC, Monack D. A gut commensal-produced
metabolite mediates colonization resistance to salmonella infection. Cell Host
Microbe 2018;24:296-307 e7.

Jiao S, Yang YF, Xu YQ, Zhang J, Lu YH. Balance between community assembly
processes mediates species coexistence in agricultural soil microbiomes across
eastern china. Isme Journal 2020;14:202-16.

Jiao S, Zhang B, Zhang G, Chen W, Wei G. Stochastic community assembly decreases
soil fungal richness in arid ecosystems. Mol Ecol 2021;30:4338-48.

Kankainen M, Paulin L, Tynkkynen S, Von Ossowski I, Reunanen J, Partanen P,
Satokari R, Vesterlund S, Hendrickx AP, Lebeer S, De Keersmaecker SC,
Vanderleyden J, Hamalainen T, Laukkanen S, Salovuori N, Ritari J, Alatalo E,
Korpela R, Mattila-Sandholm T, Lassig A, Hatakka K, Kinnunen KT,
Karjalainen H, Saxelin M, Laakso K, Surakka A, Palva A, Salusjarvi T, Auvinen
P, De Vos WM. Comparative genomic analysis of lactobacillus rhamnosus gg
reveals pili containing a human- mucus binding protein. Proc Natl Acad Sci U
S A 2009;106:17193-8.

Kim HB, Isaacson RE. Salmonella in swine: Microbiota interactions. Annual Review

of Animal Biosciences, Vol 5 2017;5:43-63.

Koh A, Backhed F. From association to causality: The role of the gut microbiota and its
functional products on host metabolism. Mol Cell 2020;78:584-96.

Kumar A, Baruah A, Tomioka M, Iino Y, Kalita MC, Khan M. Caenorhabditis elegans:
A model to understand host-microbe interactions. Cell Mol Life Sci
2020;77:1229-49.

Laupland KB, Paterson DL, Edwards F, Stewart AG, Harris PNA. Morganella morganii,
an emerging cause of bloodstream infections. Microbiol Spectr
2022;10:e0056922.

Lavelle A, Sokol H. Gut microbiota-derived metabolites as key actors in inflammatory
bowel disease. Nature Reviews Gastroenterology & Hepatology 2020;17:223-
37.

Letunic I, Bork P. Interactive tree of life (itol) v5: An online tool for phylogenetic tree
display and annotation. Nucleic acids research 2021;49:W293-W6.

Levy M, Kolodziejczyk AA, Thaiss CA, Elinav E. Dysbiosis and the immune system.
Nat Rev Immunol 2017;17:219-32.

Liu J, Liu H, Liu H, Teng Y, Qin N, Ren X, Xia X. Live and pasteurized akkermansia
muciniphila decrease susceptibility to salmonella typhimurium infection in
mice. J Adv Res 2023;52:89-102.

Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
quantitative pcr and the 2(t)(-delta delta c) method. Methods 2001;25:402-8.

Montgomery TL, Eckstrom K, Lile KH, Caldwell S, Heney ER, Lahue KG,
D'alessandro A, Wargo MJ, Krementsov DN. Lactobacillus reuteri tryptophan
metabolism promotes host susceptibility to cns autoimmunity. Microbiome
2022;10:198.

Na YR, Stakenborg M, Seok SH, Matteoli G. Macrophages in intestinal inflammation
and resolution: A potential therapeutic target in ibd. Nat Rev Gastroenterol
Hepatol 2019;16:531-43.

Neurath MF. Strategies for targeting cytokines in inflammatory bowel disease. Nat Rev
Immunol 2024.

Olesen SW, Barnett ML, Macfadden DR, Brownstein JS, Hernandez-Diaz S, Lipsitch
M, Grad YH. The distribution of antibiotic use and its association with antibiotic
resistance. Elife 2018;7.

Olive AJ, Sassetti CM. Metabolic crosstalk between host and pathogen: Sensing,
adapting and competing. Nature Reviews Microbiology 2016;14:221-34.

Parks DH, Tyson GW, Hugenholtz P, Beiko RG. Stamp: Statistical analysis of
taxonomic and functional profiles. Bioinformatics 2014;30:3123-4.

Peng X, Ed-Dra A, Song Y, Elbediwi M, Nambiar RB, Zhou X, Yue M.
Lacticaseibacillus rhamnosus alleviates intestinal inflammation and promotes
microbiota-mediated protection against salmonella fatal infections. Front
Immunol 2022;13:973224.

Poupet C, Chassard C, Nivoliez A, Bornes S. Caenorhabditis elegans, a host to
investigate the probiotic properties of beneficial microorganisms. Front Nutr
2020;7:135.

Rangan KJ, Pedicord VA, Wang YC, Kim B, Lu Y, Shaham S, Mucida D, Hang HC. A
secreted bacterial peptidoglycan hydrolase enhances tolerance to enteric
pathogens. Science 2016;353:1434-7.

Read E, Curtis MA, Neves JF. The role of oral bacteria in inflammatory bowel disease.
Nat Rev Gastroenterol Hepatol 2021;18:731-42.

Rodriguez-R LM, Konstantinidis KT, The enveomics collection: A toolbox for
specialized analyses of microbial genomes and metagenomes. PeerJ Preprints;
2016.p.

Rodriguez RL, Gunturu S, Harvey WT, Rossello-Mora R, Tiedje JM, Cole JR,
Konstantinidis KT. The microbial genomes atlas (miga) webserver: Taxonomic
and gene diversity analysis of archaea and bacteria at the whole genome level.
Nucleic Acids Res 2018;46:W282-W8.

Rogers AP, Mileto SJ, Lyras D. Impact of enteric bacterial infections at and beyond the
epithelial barrier. Nature Reviews Microbiology 2022.

Sakandar HA, Zhang HP. Trends in probiotic(s)-fermented milks and their in vivo
functionality: A review. Trends in Food Science & Technology 2021;110:55-65.

Sanders ME, Merenstein DJ, Reid G, Gibson GR, Rastall RA. Probiotics and prebiotics
in intestinal health and disease: From biology to the clinic. Nat Rev
Gastroenterol Hepatol 2019;16:605-16.

Schwarze K, Buchanan J, Fermont JM, Dreau H, Tilley MW, Taylor JM, Antoniou P,
Knight SJL, Camps C, Pentony MM, Kvikstad EM, Harris S, Popitsch N,
Pagnamenta AT, Schuh A, Taylor JC, Wordsworth S. The complete costs of
genome sequencing: A microcosting study in cancer and rare diseases from a
single center in the united kingdom. Genet Med 2020;22:85-94.

Shi J, Yang L, Liao Y, Li J, Jiao S, Shangguan Z, Deng L. Soil labile organic carbon
fractions mediate microbial community assembly processes during long-term
vegetation succession in a semiarid region. iMeta 2023:e142.

Sloan WT, Lunn M, Woodcock S, Head IM, Nee S, Curtis TP. Quantifying the roles of
immigration and chance in shaping prokaryote community structure. Environ
Microbiol 2006;8:732-40.

Stanaway JD, Parisi A, Sarkar K, Blacker BF, Reiner RC, Hay SI, Nixon MR, Dolecek
C, James SL, Mokdad AH, Abebe G, Ahmadian E, Alahdab F, Alemnew BT,
Alipour V, Bakeshei FA, Animut MD, Ansari F, Arabloo J, Asfaw ET,
Bagherzadeh M, Bassat Q, Belayneh YM, Carvalho F, Daryani A, Demeke FM,
Demis AB, Dubey M, Duken EE, Dunachie SJ, Eftekhari A, Fernandes E, Fard
RF, Gedefaw GA, Geta B, Gibney KB, Hasanzadeh A, Hoang CL, Kasaeian A,
Khater A, Kidanemariam ZT, Lakew AM, Malekzadeh R, Melese A, Mengistu
DT, Mestrovic T, Miazgowski B, Mohammad KA, Mohammadian M,
Mohammadian-Hafshejani A, Nguyen T, Nguyen H, Nguyen H, Nirayo YL,
Olagunju AT, Olagunju TO, Pourjafar H, Qorbani M, Rabiee M, Rabiee N,
Rafay A, Rezapour A, Samy AM, Sepanlou SG, Shaikh MA, Sharif M,
Shigematsu M, Tessema B, Tran BX, Ullah I, Yimer EM, Zaidi Z, Murray CJL,
Crump JA, Salm GN-T. The global burden of non-typhoidal salmonella invasive
disease: A systematic analysis for the global burden of disease study 2017.
Lancet Infectious Diseases 2019;19:1312-24.

Stegen JC, Lin X, Konopka AE, Fredrickson JK. Stochastic and deterministic assembly

925   processes in subsurface microbial communities. ISME J 2012;6:1653-64.

926   Stegen JC, Lin XJ, Fredrickson JK, Chen XY, Kennedy DW, Murray CJ, Rockhold ML,

927   Konopka A. Quantifying community assembly processes and identifying

928   features that impose them. Isme Journal 2013;7:2069-79.

929   Sun Y, Li HC, Zheng L, Li JZ, Hong Y, Liang PF, Kwok LY, Zuo YC, Zhang WY, Zhang

930   HP. Iprobiotics: A machine learning platform for rapid identification of

931   probiotic properties from whole-genome primary sequences. Briefings in

932   Bioinformatics 2022;23.

933   Thiagarajah JR, Donowitz M, Verkman AS. Secretory diarrhoea: Mechanisms and

934   emerging therapies. Nat Rev Gastroenterol Hepatol 2015;12:446-57.

935   Unicef, Diarrhoea remains a leading killer of young children, despite the availability of

936   a simple treatment solution. U Data (Ed.). https://data.unicef.org/topic/child-

937   health/diarrhoeal-disease/; 2022.p.

938   Vellend M. Conceptual synthesis in community ecology. Q Rev Biol 2010;85:183-206.

939   Wan MLY, Forsythe SJ, El-Nezami H. Probiotics interaction with foodborne pathogens:

940   A potential alternative to antibiotics and future challenges. Critical reviews in

941   food science and nutrition 2019;59:3320-33.

942   Wang B, Hussain A, Zhou Y, Zeng Z, Wang Q, Zou P, Gong L, Zhao P, Li W.

943   Saccharomyces boulardii attenuates inflammatory response induced by

944   clostridium perfringens via tlr4/tlr15-myd8 pathway in hd11 avian macrophages.

945   Poult Sci 2020;99:5356-65.

946   Wang B, Ye X, Zhou Y, Zhao P, Mao Y. Glycyrrhizin attenuates salmonella

947   typhimurium-induced tissue injury, inflammatory response, and intestinal

948   dysbiosis in c57bl/6 mice. Front Vet Sci 2021a;8:648698.

949   Wang B, Zhou Y, Mao Y, Gong L, Li X, Xu S, Wang F, Guo Q, Zhang H, Li W. Dietary

950   supplementation with lactobacillus plantarum ameliorates compromise of

951   growth performance by modulating short-chain fatty acids and intestinal

952   dysbiosis in broilers under clostridium perfringens challenge. Front Nutr

953   2021b;8:706148.

954   Wang B, Zhou Y, Wang Q, Xu S, Wang F, Yue M, Zeng Z, Li W. Lactiplantibacillus

plantarum lac16 attenuates enterohemorrhagic escherichia coli o157:H7 infection by inhibiting virulence traits and improving intestinal epithelial barrier function. Cells 2023;12.

Wang L, Liu Q, Chen Y, Zheng X, Wang C, Qi Y, Dong Y, Xiao Y, Chen C, Chen T, Huang Q, Zhai Z, Long C, Yang H, Li J, Wang L, Zhang G, Liao P, Liu YX, Huang P, Huang J, Wang Q, Chu H, Yin J, Yin Y. Antioxidant potential of pediococcus pentosaceus strains from the sow milk bacterial collection in weaned piglets. Microbiome 2022;10:83.

Who, Diarrhoeal disease. 2017.p.

Wlodarska M, Luo C, Kolde R, D'hennezel E, Annand JW, Heim CE, Krastel P, Schmitt EK, Omar AS, Creasey EA, Garner AL, Mohammadi S, O'connell DJ, Abubucker S, Arthur TD, Franzosa EA, Huttenhower C, Murphy LO, Haiser HJ, Vlamakis H, Porter JA, Xavier RJ. Indoleacrylic acid produced by commensal peptostreptococcus species suppresses inflammation. Cell Host Microbe 2017;22:25-37 e6.

Wu Y, Zhang X, Liu X, Li Y, Han D, Pi Y, Whitmore MA, Lu X, Zhang G, Zheng J, Wang J. Strain specificity of lactobacilli with promoted colonization by galactooligosaccharides administration in protecting intestinal barriers during salmonella infection. J Adv Res 2024;56:1-14.

Xiao Y, Zhai Q, Zhang H, Chen W, Hill C. Gut colonization mechanisms of lactobacillus and bifidobacterium: An argument for personalized designs. Annu Rev Food Sci Technol 2021;12:213-33.

Xu Q, Ling N, Chen H, Duan Y, Wang S, Shen Q, Vandenkoornhuyse P. Long-term chemical-only fertilization induces a diversity decline and deep selection on the soil bacteria. mSystems 2020;5.

Xu Q, Vandenkoornhuyse P, Li L, Guo J, Zhu C, Guo S, Ling N, Shen Q. Microbial generalists and specialists differently contribute to the community diversity in farmland soils. J Adv Res 2022;40:17-27.

Xu XG, Gong L, Wang BK, Wu YP, Wang Y, Mei XQ, Xu H, Tang L, Liu RR, Zeng ZH, Mao YL, Li WF. Glycyrrhizin attenuates salmonella enterica serovar

typhimurium infection: New insights into its protective mechanism. Frontiers
in Immunology 2018;9:2321.

Xun W, Li W, Xiong W, Ren Y, Liu Y, Miao Y, Xu Z, Zhang N, Shen Q, Zhang R.
Diversity-triggered deterministic bacterial assembly constrains community
functions. Nat Commun 2019;10:3833.

Yu G, Xu C, Zhang D, Ju F, Ni Y. Metorigin: Discriminating the origins of microbial
metabolites for integrative analysis of the gut microbiome and metabolome.
IMeta 2022;1:e10.

Zhang Y, Wang J, He J, Liu X, Sun J, Song X, Wu Y. Characteristics and application in
cheese making of newly isolated milk-clotting enzyme from bacillus
megaterium ly114. Food Res Int 2023;172:113202.

Zhou J, Ning D. Stochastic community assembly: Does it matter in microbial ecology?
Microbiology and Molecular Biology Reviews 2017;81:e00002-17.

Zhou X, Kang X, Chen J, Song Y, Jia C, Teng L, Tang Y, Jiang Z, Peng X, Tao X.
Genome degradation promotes salmonella pathoadaptation by remodeling
fimbriae-mediated proinflammatory response. National Science Review
2023:nwad228.

1002

Figures

Figure 1. Isolation and antibacterial characterization of *L. rhamnosus* P118. (**A**) Screening flowchart of P118 *in vitro* and candidate probiotic isolates identified by MALDI-TOF MS. (**B**) Bile salt tolerance ability of the isolates. (**C**) Biofilm forming

ability of isolates. (**D**) Screening using *C. elegans* infection model. (**E**) Interacted screening strategy. (**F**) Antibacterial ability of P118 under aerobic or anaerobic culture conditions. (**G**) Broad-spectrum pH tolerance of P118 supernatants (pH = 3.4) that adjusted to pH < 3.4 by 1M HCl or pH > 3.4 by 1M NaOH. (**H**) Antibacterial effects of P118 supernatant under 20 mg/mL catalase (CAT), 100μg/mL proteinases (trypsin, proteinase K, pepsin), and different temperatures (37, 50, 70, 90, 100, 120°C) boiled for 30 min treatments. (**I**) Antibacterial effects of components of P118 (boiled at 120 °C for 30 min or was lysed by ultrasonication at 240 W for 2 h). (**G**) Active ingredients of P118 protect *C. elegans* against *S*. Typhimurium infection. (**K**) Taxonomic classification of P118 draft genome queried against the NCBI non-redundant prokaryotic genomes database with p-values representing confidence of phylogenetic assignment, and the nearest subspecies phylogenetic neighbor of P118 draft genome was determined by percentage shared genomic content graphed as ANI versus AAI. (**L**) Prediction of secondary metabolites and bacteriocin protein-encoding gene clusters of P118 using antiSMASH and BAGEL4 databases. (**F-G**) *S*. Typhimurium SL1344 was selected as an indicator pathogen. Significant differences * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure 2. *L. rhamnosus* P118 enhances mice tolerance to *S*. Typhimurium infection. (**A**)

Experimental design. (**B**) Survival curve of mice infected with *S*. Typhimurium. (**C**)

Body weight. (**D**) Spleen index. (**E**) Representative images of spleen and intestine. (**F**)

Liver index. (**G**) *S.* Typhimurium burden in tissues and shedding in feces. (**H**)

Representative images of H&E staining, TEM, and immunostaining (DAPI, Ki67,

lysozyme, Muc2, F4/80, iNOS) in the ileum. Different lowercase letters indicate a

significant difference (*P* < 0.05). Significant differences * *P* < 0.05, ** *P* < 0.01. C:

PBS group; P: P118 administered group; S: *S.* Typhimurium-infected group; P+S: P118

protective group.

Figure 3. *L. rhamnosus* P118 improves *S.* Typhimurium infection-induced

dysbacteriosis. (**A**) Principal coordinates analysis (PCoA) based on Bray-Curtis

distance. (**B**) UpSetR plot based on bacterial absolute OTU abundances. (**C & D**) The

fold changes of bacterial absolute OTU abundances between two groups. (**E & F**)

Comparison of intestinal microbes by STAMP. The prefix "g_" represent the annotated

level of the genus. C: PBS group; P: P118 administered group; S: *S.* Typhimurium-

infected group; P+S: P118 protective group.

Figure 4. Microbe derived tryptophan metabolites are involved in mice tolerance to *S*.

Typhimurium. (**A**) Principal coordinates analysis (PCoA) based on Bray-Curtis

distance of fecal metabolites. (**B & C**) UpSetR plot based on fecal metabolites. (**D**)

Metabolomics pathway enrichment in "P+S *vs* S". (**E**) Comparison of fecal microbial

tryptophan metabolism enriched pathway among groups. (**F**) Pearson correlation

analysis among *S*. Typhimurium burden, organ indices, body weight and fecal

metabolites in mice. (**G**) Bio-Sankey network analysis between intestinal microbes and

fecal metabolites. (**H**) Pathway schematic of abbreviated mammalian and microbial

tryptophan metabolism. Different lowercase letters indicate a significant difference (*P*

< 0.05). Significant differences **P* < 0.05; ***P* < 0.01; ****P* < 0.001. C: PBS group; P:

P118 administered group; S: *S*. Typhimurium-infected group; P+S: P118 protective

group.

1056

1057 **Figure 5.** Indole-3-acrylic acid enhances mice tolerance to *S*. Typhimurium infection.

1058 (**A**) Experimental design. (**B**) Body weight change. (**C**) Colon length. (**D**) Spleen and

1059 liver indexes. (**E**) *S*. Typhimurium burden in tissues and shedding in feces. (**F**)

1060 Representative images of spleen and liver. (**G**) Villus height of ileum. (**H**) Muc2-

 positive cells and mRNA expression levels in ileum. (**I**) Representative images of H&E

 staining and immunostaining in the ileum. Different lowercase letters indicate a

 significant difference (*P* < 0.05). C: PBS group; S: *S*. Typhimurium-infected group;

 IAA+S: indole-3-acrylic acid protective group.

 Figure 6. Macrophage depletion abrogates the protective effect of *L. rhamnosus* P118

and indole-3-acrylic acid against *S*. Typhimurium infection. (**A**) Experimental design.

(**B**) Body weight change. (**C**) Spleen and liver indexes. (**D**) Representative images of

spleen and liver. (**E**) Colon length. (**F**) *S.* Typhimurium burden in tissues and shedding

in feces. (**G**) Villus height of ileum. (**H**) Muc2-positive cells and mRNA expression

levels in ileum. (**I**) Representative images of H&E staining and immunostaining in the

ileum. Different lowercase letters indicate a significant difference ($P < 0.05$). (**B-E**)

Significant differences ** $P < 0.01$ indicates "S *vs* C".CLP: macrophage depletion

reagent clodronate liposomes. C: PBS group; S: S. Typhimurium-infected group;

IAA+S: indole-3-acrylic acid protective group; P+S: P118 protective group.

1076

Figure 7. *Lacticaseibacillus rhamnosus* P118 strain with great probiotic properties was

screened from 290 purified isolates through two distinctive screen approaches, and

P118 strain was underscored for protective effects on a murine infection model. Further,

multi-omics analysis pinpointed the microbe-derived tryptophan/indole could be the

importance of P118 probiotic properties. Nevertheless, the newly found P118 enhances

host tolerance to *Salmonella* infections via various pathways, including direct

antibacterial actions, inhibiting *Salmonella* colonization and invasion, attenuating pro-

inflammatory responses of intestinal macrophages, and modulating gut microbiota

mediated by microbe-derived indole metabolites.

1086

1087

1088 **Table 1** A list of probiotic isolates recovered from examined samples

Source	Name	Number of isolates
Yogurt	*Streptococcus salivarius ssp. thermophilus*	63
	Lacticaseibacillus paracasei	18
	Lactobacillus acidophilus	17
	Lactobacillus zeae	10
	Lacticaseibacillus rhamnosus	4
	Lacticaseibacillus casei	4
	Lactobacillus delbrueckii	6
piglet intestine	*Limosilactobacillus mucosae*	16
	Lactobacillus johnsonii	38
	Limosilactobacillus reuteri	16
	Limosilactobacillus vaginalis	11
	Lactobacillus crispatus	9
	Lactobacillus salivarius	6
	Lactobacillus amylovorus	3
	Enterococcus faecium	38
	Enterococcus faecalis	31
Total		290

1089 **Supplementary information**

     Figure S1 (**A**) Drug resistance candidate probiotic isolates (P118 and P199). (**B, C**)

     Broad-spectrum antibacterial activities of P118 fermented supernatants and cultures.

     Figure S2. (**A**) Circular genome map of *L. rhamnosus* P118. (**B**) phylogenetic trees of

     L. rhamnosus P118 and nearest phylogenetic neighbors constructed by average amino

     acid identify (AAI). Color gradients indicate the percentage of conserved average

1097 nucleotide identity (ANI) between each isolate and respective nearest subspecies

1098 phylogenetic neighbors.

1099

Figure S3. Clinical and Histopathological observation. (**A**) Representative images of

H&E staining tissues (liver, spleen). Scale bar with 100 μm. (**B**) Clinical symptom score.

Histopathological quantification of H&E staining (**C & D**) and TEM (**E & F**) images

observed in the ileum. Different lowercase letters indicate a significant difference (*P* <

0.05). C: PBS group; P: P118 administered group; S: *S*. Typhimurium-infected group;

P+S: P118 protective group.

1106

1107 **Figure S4.** Gene and protein expression level of tight junction protein (ZO-1, Claudin-

1108 1) and inflammatory cytokines (IL-1β, IL-18) in ileum and serum of mice. Different

1109 lowercase letters indicate a significant difference ($P < 0.05$). C: PBS group; P: P118

1110 administered group; S: *S.* Typhimurium-infected group; P+S: P118 protective group.

Figure S5. Comparison of intestinal microbes by the analysis of statistical analysis of taxonomic and functional profiles (**STAMP**) software with a two-sided Welch's t-test. (**A**) C *vs* P, (**B**) C *vs* S, (**C**) S *vs* P+S. The prefixes "g_" and "s_" represent the annotated levels of genus and species. C: PBS group; P: P118 administered group; S: *S.* Typhimurium-infected group; P+S: P118 protective group.

Figure S6. *L. rhamnosus* P118 alters gut microbial community assembly. (**A**) Ecological processes of gut microbes. The inner circle indicates the percentage of deterministic and stochastic assembly processes. The outer circle indicates the contribution of detailed ecological processes. (**B**) The fit of neutral community model (NCM) of community assembly for gut microbes. The best fit to NCM and 95% CIs around the best fit to NCM were indicated by solid and dashed blue lines, respectively. Light blue and dark red dots indicate OTUs that occur more (above prediction) or less (below prediction) frequently than predicted by NCM, respectively, while black dots indicate the OTUs that occur within neutral prediction (neutral distribution). Donut charts shows the distinct fitting proportions of OTUs predicted by NCM. C: PBS group; P: P118 administered group; S: *S*. Typhimurium-infected group; P+S: P118 protective

1129   group.

1130

1131

Figure S7. Analysis of differential fecal metabolites. Venn diagram based on

1133 differential fecal metabolites (**A**) down-regulated in "S *vs* C" and up-regulated in "P+S

1134 *vs* S", (**B**) up-regulated in "S *vs* C" and down-regulated in "P+S *vs* S". (**C**) Significant

1135 differential fecal metabolites in "S *vs* C" and "P+S *vs* S". C: PBS group; P: P118

1136 administered group; S: *S.* Typhimurium-infected group; P+S: P118 protective group.

1137

1138 **Figure S8.** Antibacterial activities of indole-3-acrylic acid. GM: 100 μg/mL gentamicin.

Figure S9. Indole-3-acrylic acid enhances bactericidal capacity and exerts anti-inflammatory activity in RAW 264.7 cells. Different lowercase letters (a, b, c, d) indicate a significant difference ($P < 0.05$). C: PBS group; IAA: indole-3-acrylic acid group; S: *S*. Typhimurium-infected group; CH-223191: AHR inhibitor-treated group.

Table S1 Milk-clotting activity of lactic acid bacteria

Table S2 Significance between life span data was calculated using a Mantel-Cox log-rank test

Table S3 Significance between life span data was calculated using a Mantel-Cox log-rank test

Table S4 Species-level identity of *L. rhamnosus* P118. The draft genome of P118 was queried using the NCBI non-redundant prokaryotic database available through MIGA

(http://www.microbial-genomes.org). P-values reflect confidence in the assignment as

each taxonomic rank.

Table S5 *L. rhamnosus* P118 draft genome nearest subspecies phylogenetic neighbors.

Average nucleotide identity (ANI) and average amino acid identity (AAI) using both

percent identity and a fraction of genome shared as determined through MIGA

(http://www.microbial-genomes.org) query of the NCBI non-redundant prokaryotic

genomes database were considered for ranking likelihood of nearest subspecies

phylogenetic neighbors for duplicate isolates of each *Lacticaseibacillus* species.

Table S6 Primers used for the quantitative PCR

Table S7 The criteria of clinical symptom scores

Table S8 ANOSIM, PERMANOVA and MRPP analysis of microbial diversity among

treatments

Table S9 ANOSIM, PERMANOVA and MRPP analysis of fecal metabolites among

groups

Table S10 Significant differential fecal metabolites in "S *vs* C" and "P+S *vs* S"

Table S11 Prediction of aminotransferase class and tryptophan binding affinity in *L.*

rhamnosus P118

Table S12 Tryptophan metabolism and indole derivatives in *L. rhamnosus* P118

cultures